



Enhancing road safety: In-vehicle sensor analysis of cognitive impairment in older drivers

Muhammad Tanveer Jan¹ · Borko Furht¹ · Sonia Moshfeghi¹ · Jinwoo Jang¹ · Seyedeh Gol Ara Ghoreishi¹ · Charles Boateng¹ · Kwangsoo Yang¹ · Joshua Conniff² · Monica Rosselli² · David Newman³ · Ruth Tappen³

Received: 23 January 2024 / Revised: 1 June 2024 / Accepted: 7 July 2024

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2024

Abstract

With the ongoing expansion of the aging population, it is increasingly critical to prioritize the safety of older drivers. The objective of this study is to utilize sensor data in order to detect early indications of impairment, thereby facilitating proactive interventions and enhancing road safety for the elderly. This article provides an overview of the research approach, presents significant results, and analyzes the consequences of utilizing in-vehicle sensors i.e. vision and telematics, to mitigate cognitive decline among elderly drivers; in doing so, it promotes progress in the domains of public health and transportation safety by standardizing the use of such devices to automatically assess the drivers' cognitive functions.

Keywords Alzheimers · Mild cognitive impairment · In-vehicle sensors · Driver behaviors · Driver monitoring system

1 Introduction

The current global demographic landscape is experiencing a substantial shift, marked by a notable rise in the proportion of elderly individuals. As individuals progress in age, the maintenance of cognitive abilities becomes a matter of utmost importance, particularly in relation to tasks that need prolonged focus and the ability to make informed choices, such as driving. The confluence of an increasingly elderly population and vehicle mobility gives rise to urgent inquiries regarding road safety and the potential hazards linked to cognitive deterioration in older drivers. Mild Cognitive Impairment (MCI) is a significant transitional phase that lies between the typical aging process and more pronounced cognitive deficits, such as those observed in dementia. Elderly adults who are experiencing MCI may display inconspicuous cognitive impairments that have an effect on their day-to-day functioning [44], such as their capacity to operate a vehicle safely. The increasing number of elderly drivers on the road due to the aging population has made it crucial to prioritize the detection and treatment of cognitive impairment in this group to maintain public safety.

Extended author information available on the last page of the article

This study aims to investigate the utilization of in-vehicle sensor technology for the analysis of MCI in elderly drivers. The conventional approaches employed for evaluating cognitive function within a clinical context frequently exhibit limitations in effectively capturing dynamic, real-world situations, such as driving. The utilization of in-vehicle sensors is a potentially fruitful approach for the ongoing observation of driving behavior, hence furnishing unbiased data that can be associated with cognitive function [22].

The primary objective of this study is to assess the viability of employing in-vehicle sensors as a means of promptly identifying cognitive deterioration in elderly drivers and providing important insights into the possible impact of technology on improving road safety for the elderly population by examining the association between sensor data and cognitive tests. The paper examines the research methodologies, findings, and implications with the main goal of offering insights for future policies and practices that try to tackle the unique challenges that arise from the combination of aging, cognition, and mobility.

This study covers the literature review in Section 2, the methodology in Section 3, the data analysis of supporting results in Section 4, results and future directions in Section 5, and the conclusion in Section 6.

1.1 Background

The unprecedented growth of the aging population worldwide presents a multifaceted challenge, particularly concerning road safety. As individuals live longer, the proportion of older drivers on the road continues to rise, necessitating a comprehensive understanding of the implications of aging on driving capabilities. Older drivers may experience age-related changes in vision, reaction time, and cognitive function, contributing to an increased vulnerability to accidents. Recognizing the unique challenges posed by this demographic shift is imperative for developing targeted interventions and technologies that promote both the mobility and safety of older individuals on the road.

The prevalence of MCI among older drivers is a significant concern in the context of road safety. Studies indicate that a notable proportion of older individuals experience MCI, a condition characterized by subtle cognitive deficits [1]. As cognitive functions play a crucial role in driving, understanding the prevalence of MCI among older drivers is essential for addressing the potential risks associated with this demographic in transportation scenarios.

This study holds paramount significance as it addresses the critical intersection of an aging population and road safety. Investigating the correlation between MCI and driving performance in older individuals contributes valuable insights that can inform targeted interventions. The findings have the potential to enhance public safety by identifying early markers of cognitive decline, allowing for proactive measures to be implemented in support of older drivers. Moreover, the study's implications extend beyond individual well-being, influencing policy development and technological innovations to foster a safer and more inclusive transportation environment for an aging demographic.

2 Literature review

2.1 Cognitive impairment and driving performance

Several research publications have evidenced that cognitive impairment significantly impacts driving performance among older individuals. A study by Ott et al. [2] found that cognitive

deficits, particularly in attention and executive functions, were associated with a higher risk of traffic accidents among older drivers. Similarly, a meta-analysis by Roe et al. [3] demonstrated a consistent relationship between cognitive decline and decreased driving abilities in older adults. These studies underscore the importance of understanding the nuanced connections between cognitive impairment and its impact on the complex task of driving among the aging population. The relationship between cognitive decline and driving abilities has been extensively explored in the literature, highlighting the intricate interplay between cognitive functions and safe driving. Studies such as the work by Anstey et al. demonstrated a clear association between declines in processing speed, visual attention, and working memory with diminished driving performance in older adults [4]. Additionally, a longitudinal investigation by Ball et al. found that declines in executive function were predictive of subsequent driving cessation among older individuals [5]. These findings underscore the critical need to understand how cognitive decline influences the complex task of driving and support the exploration of innovative approaches, such as in-vehicle sensors, to objectively assess and mitigate these challenges.

2.2 Existing research on the impact of MCI on driving

Various methods have been employed to assess cognitive impairment in older drivers, reflecting the complexity of the task. Standardized cognitive tests, such as the Mini-Mental State Examination (MMSE) and the Montreal Cognitive Assessment (MoCA), are commonly utilized in clinical settings to screen for cognitive deficits [6, 7]. However, these assessments may not fully capture the dynamic cognitive demands of driving. Research by Dawson et al. emphasizes the limitations of traditional neuropsychological tests in predicting real-world driving performance, suggesting the need for more ecologically valid [8]. To address these challenges, emerging technologies, including virtual reality-based assessments, are being explored. For example, a study by Classen et al. investigated the utility of a virtual reality driving simulator in identifying cognitive deficits that may impact driving ability [9]. As research progresses, these novel approaches hold promise in providing a more comprehensive understanding of cognitive impairment in the context of driving.

Research on driving fitness has predominantly focused on individuals with Alzheimer's disease (AD), particularly in complex traffic situations where those with AD exhibit heightened self-restriction owing to impaired attention and executive functioning [23]. Studies indicate that AD patients commit significantly more driving errors than their healthy counterparts, facing challenges in route and traffic rule recollection, as well as identifying landmarks and traffic signs [24–27]. Their misjudgments encompass distances to other vehicles, errors at intersections, and delayed actions or reactions [24]. In a driving simulator study by Rizzo and colleagues, 29% of AD patients were involved in crashes, contrasting with the absence of incidents in the healthy control group. Notably, near accidents occurred more than twice as often in AD patients (74% vs. 35%), with impairments in visual-spatial attention and spatial thinking emerging as key predictors of accident risk [28]. Surveys of AD patients' relatives indicated changes in driving behavior in 58% of patients, with orientation deficits being a prominent dysfunction and nearly 20% being responsible for at least one accident post-diagnosis [29]. The overall accident risk in Alzheimer patients surpasses that of healthy older individuals by more than four times [2], and it increases with the time of driving since disease onset [30].

It's crucial to note that these findings might not uniformly apply to early-stage, very mild, or mild stages of the disease. The early-stage, defined as amnesic mild cognitive impairment (aMCI), characterized by isolated episodic memory dysfunction, suggests that drivers with aMCI exhibit suboptimal rather than definitively impaired driving fitness [24, 31].

In the case of very mild AD, meta-analytical evidence points to on-road fail rates of approximately 13%, in contrast to 1.6% in healthy controls [24]. This discrepancy may be attributed in part to inappropriate behavior at intersections, such as excessive stopping [24]. For mild AD, a striking 33% of patients fail an on-road test, with individual studies reporting even higher fail rates of 50.6% and 58% for patients with very mild and mild ADD, respectively, compared to 4.4% and 11% in healthy controls [25, 32]. On-road performances of drivers with very mild and mild AD are compromised on operational, tactical, and visual levels, involving more turning errors, loss of orientation, and maneuvers jeopardizing road safety [26]. Carr and colleagues reported that drivers with very mild and mild AD tend to be involved in more at-fault crashes than healthy drivers, including accidents resulting in personal injuries [33].

2.3 In-vehicle sensor technologies

In-vehicle sensor technologies offer a promising avenue for objectively assessing driving performance in real time. These sensors, capable of capturing various aspects of driver behavior and vehicle dynamics, have been applied in research to understand and monitor driver performance. For instance, studies by Horrey et al. [10] and Dingus et al. [11] utilized in-vehicle sensors, including accelerometers and gyroscopes, to analyze driving behavior and assess factors contributing to safety. Moreover, advancements in sensor technology, such as eye-tracking systems and lane departure warning systems, have been instrumental in quantifying visual attention and detecting deviations in driving behavior. Research by Mehler et al. demonstrated the effectiveness of eye-tracking technology in assessing cognitive load and attention allocation during driving tasks [12]. Additionally, studies by Bosurgi et al. and Eskandarian (2012) have explored the application of sensors in assessing driver behavior and performance in the context of driver assistance systems and advanced driver assistance technologies (ADAS) [13, 14]. Apart from this, In [34], a real-time intelligent system is proposed, utilizing traffic cameras and the You Only Look Once (YOLO) algorithm for object detection. The Kalman filter tracks vehicle location, and anomaly detection, based on speed, is performed with an impressive real-time capability. Article [35] introduces a framework based on the Strategic Highway Research Program 2 (SHRP 2) and Naturalistic Driving Study (NDS) datasets. Using a Random Forest (RF) algorithm, the study calculates a driver's risk profile, achieving an accuracy rate of 90%. In [36], the Serial-Feature Network (SF-Net) algorithm is proposed for recognizing normal and abnormal driver behavior using smartphone inertial sensors like GPS and gyroscope. The approach achieves remarkable accuracy (97.10%) and recall rate (98.4%). Paper [37] classifies driver behaviors and road anomalies using smartphone sensor data. The k-Nearest Neighbor (KNN) and Dynamic Time Warping (DTW) algorithms exhibit significant accuracy rates, reaching 78.06% and 96.75%, respectively. In [38], the "Project Drive" mobile application is introduced, employing the k-means algorithm on GPS data to detect negative driver behavior and encourage safer practices. Papers [39] and [40] classify driver behavior as positive and negative. Babić et al. [39] explores the impact of road signs on young drivers' behavior in nighttime conditions, while [40] investigates the effects of optical circles and chevron patterns on driver behavior using

simulator data and the MANOVA statistical technique. Papers [41], [42], and [43] focus on distinguishing between safe and unsafe driver behavior. Eren [41] classifies driver behavior through smartphone sensors using an optimal path detection algorithm and Bayesian classification, achieving a correct classification rate of 93.3%. Wadley et al. [31] proposes a system based on Local and LARA traffic datasets to provide driving advice, achieving a precision rate of 95.52%. In [32], a smartphone-based system utilizes Long Short-term Memory (LSTM) and Convolutional Neural Network (CNN) models, reaching a mean percentage error (MPE) of 0.36 for analyzing driver behavior at intersections.

3 Methodology

3.1 Participants Characteristics

The study sample comprises 225 participants with a mean age of 78.00 years (SD = 6.05), ranging from 65 to 91 years. The majority of participants identified as White (n = 172), followed by Black (n = 36), Asian (n = 3), Multiracial (n = 4), and Other (n = 2), with 18 respondents choosing not to disclose their race. Education levels ranged from 0 to 26

Table 1 Descriptive Statistics of the Sample

Variable	Mean (SD)	Min	Max	Count
Age	78 (6.05)	65	91	225
Undisclosed	—	—	—	8
65-69	—	—	—	46
70-74	—	—	—	47
75-79	—	—	—	68
80-84	—	—	—	34
85-89	—	—	—	21
90-94	—	—	—	1
Race				
White	—	—	—	172
Black	—	—	—	36
Asian	—	—	—	3
Multiracial	—	—	—	4
Other	—	—	—	2
No Answer	—	—	—	8
Years of Education	15.99 (4.33)	0	26	225
Undisclosed	—	—	—	6
— 12	—	—	—	21
12 – 14	—	—	—	41
15 – 16	—	—	—	69
16+	—	—	—	88
Gender				
Male	—	—	—	86
Female	—	—	—	135
Undisclosed	—	—	—	4

years, with an average of 15.99 years ($SD = 4.33$) with a majority of them having a graduate degree (16 yrs). In terms of gender distribution, 86 participants identified as male and 135 as female, as shown in Table 1 These demographic details provide a comprehensive overview of the diverse characteristics within the study population, essential for understanding potential associations in subsequent analyses (Fig. 1).

3.2 In-vehicle sensor setup

The architectural design, as shown in Fig. 2, of the proposed in-vehicle sensors consists of two distributed sensing units:

- An in-vehicle vision sensing unit is utilized to capture visual data within a vehicle.
- An in-vehicle telemetry unit is employed to collect and transmit data related to the vehicle's performance and operation.

3.2.1 In-vehicle vision sensors

The placement and calibration of camera-based sensors integrated with a Mobile Digital Video Recorder (MDVR) play a crucial role in optimizing their effectiveness for monitoring and assessing driving behavior. Interior cameras can monitor driver behavior, eye movements [15], and alertness, while exterior cameras contribute to recording traffic conditions and potential hazards. The calibration of these cameras is of utmost importance in order to




Fig. 1 Overall distribution of all demographics groupings




Fig. 2 In-Vehicle Sensors

guarantee the precision and dependability of data collecting. The accurate alignment and setup of cameras ensure the precise tracking of events and activities, hence aiding further analysis. The MDVR, functioning as a centralized storage device, assumes a crucial role in securely preserving the recorded data for subsequent retrieval and analysis. Table 2 shows the features that are collected through the vision sensors (Fig. 3).

Table 2 Feature used for Driver’s behaviors

Camera type	Features
Driver’s Camera	Face Detection Eyes Detection Yawning Distraction Phone Usage
Front Camera	Traffic Sign Detection Object Detection Lane Departure Near Collision Pedestrian



Fig. 3 In-Vehicle Vision Sensors, 2 Cameras + MDVR

3.2.2 In-vehicle telematic sensors

This study uses Auto Pi's Telematics Unit (TMU), built on Raspberry Pi 4 Model B, as an open-source platform for vehicle data collection. Figure 4 shows an overview of the Telematic Unit. This programmable unit supports hardware and software expandability, allowing for customized sensing parameters. Each TMU consists of a GPS (Geographic Positioning Systems) sensor, an Inertial Measurement Unit (IMU), an On-Board Diagnostics (OBD) connector, a 4G/LTE cellular modem, an SD card, and a USB flash drive. Tri-axial gyroscopes are added to capture angular velocity. The TMU also has a smart power system that monitors voltage levels and sleeps or hibernates when necessary. In-vehicle data can be collected from the Controller Area Network (CAN bus) and augmented sensor hardware. Table 3 shown different types of data and their statistical analysis that are being performed (Fig. 5)

3.3 Data collection and extraction

The data extraction and analysis workflow [Fig. 6], has been carefully designed to ensure efficient management and utilization. Vision and telematic sensors are systematically installed



Fig. 4 In-Vehicle Telematics Sensors Workflow [20]

Table 3 In-Vehicle Telematics Sensors Indices

Groups	DBIs	Analytics
From OBD	Number of Trips, Duration, Distance, Speed, RPM, Engine Load, Fuel Level, Ambient Air Temperature, Throttle Positioning	Statistical Analysis, Artificial Intelligence (AI) and Machine Learning (ML)
From IMU	Harsh-acceleration (count, value), Hard-braking (count, value), Hard-turn (count, value)	
From GPS	SOG (Speed Over Ground), Longitude, Latitude, COG (Course over Ground), Altitude	




Fig. 5 In-Vehicle Telematic Sensor




Fig. 6 Data Extraction and Visualization of In-vehicle Sensors

within the vehicle, and data extraction is conducted at three-month intervals. The extracted data undergo meticulous processing for subsequent storage on the cloud server. Simultaneously, a comprehensive analysis is performed, facilitating the preparation of detailed reports. These reports are then visualized on the web-server as discussed in previous works [17, 18], enhancing accessibility and facilitating a comprehensive understanding of the derived insights.

Lane changes are of significance for assessing driving-related cognitive functions. These maneuvers demonstrate the driver's awareness, decision-making, and operational proficiency, involving cognitive functions like attention, reasoning, and control. Thus, studying lane changes provides vital insights into driving behaviors, helping to understand driver performance and safety (Fig. 7).

Figure 8 shows a sequence of views from a vehicle's front-facing camera, recording a lane change maneuver. The first image shows the vehicle in the right lane, with a clear view of the road ahead. The second image captures the vehicle after it has changed lanes, now in



Fig. 7 Front-camera showing the lane departure, left lane change (top) and right-lane change (bottom)[18]




Fig. 8 Driver-facing Camera Collected Data Indices[18]

the left lane, continuing down the same road. These images are used as evidence of driving behavior and for reviewing the actions taken by the driver during the lane change.

4 Results

4.1 Vision sensors

Two distinct patterns emerge from the analysis of the charts comparing MCI driver Fig. 9 and non-MCI driver Fig. 10. The data is visualized based on monthly basis showing different indices. The data reveals variation in phone usage, near-collision incidents, and lane changes among MCI drivers, suggesting a higher degree of variability in these behaviors. This dynamic pattern suggests that MCI drivers exhibit inconsistent trends in phone engagement and driving maneuvers, possibly indicative of a lack of standardized behavior or varying levels of attention. In contrast, non-MCI drivers exhibit more stable patterns in these metrics, signifying a consistent and predictable driving behavior. These insights underscore the importance of understanding and addressing the specific challenges associated with MCI drivers, potentially contributing to targeted interventions for improving safety and reducing risk on the road.

4.2 Telematic sensors results

In Figs. 11 and 12, this study presented the problematic findings of some Driver Behavior Indicators (DBIs) obtained from real-time telematics data for both MCI and Non-MCI drivers. Observing more swings on plots from different DBIs related to MCI drivers indicates complex




Fig. 9 Plots for Participant with MCI on Monthly-Basis

driving behavior. Based on this finding, a higher fluctuation in fuel levels among MCI drivers suggests that they may neglect refueling their vehicles consistently. The constant variation in ‘RPM’ and ‘Throttle Positioning’ may indicate an aggressive approach, causing more stress on the engine [16]. This assertion can be validated by finding a higher average count of hard acceleration among drivers with MCI.

In this study [21], the problem of anomalous behavior detection is to identify drivers exhibiting significant directional deviations, frequent occurrences of hard braking, and acceleration throughout their trips. Figures 13 and 14 illustrate snapshots of the real dataset where some drivers made long-distance U-turns or repeatedly drove along the same road. These behaviors often suggest drivers might have lost their way or encountered confusion while navigating. In this problem formulation, each trip is represented by a directed graph. Then, this study proposed an Edge-Attributed Matrix to examine various graphs with different




Fig. 10 Plots for Participant with Non-MCI on Monthly-Basis

start and end points and utilized the isolation Forest algorithm to detect anomalous graphs displaying notable differences in their attribute patterns compared to the rest of the dataset.

5 Discussions

The results presented are preliminary. The final results could improve public safety and individual well-being. First, in-vehicle sensor technology seems to detect cognitive changes in older drivers, enabling preventative interventions. Fildes et al. found that early cognitive impairment identification can improve road safety by targeting therapies and support systems. [14] Second, integrating in-vehicle sensor technologies into healthcare and transportation networks may help solve the complicated problems of an aging population. Dingus et al. suggest integrating sensor data into transportation infrastructure to improve road safety and reduce driving dangers [11].




Fig. 11 Plots for Participant with MCI on Monthly-Basis

5.1 Limitations

The dynamic and diverse nature of driving surroundings makes assessing cognitive function in real-world driving scenarios difficult. Clinical cognitive evaluations may not fully capture drivers' complex cognitive demands. Driving requires quick decision-making, navigation, and response to unforeseen situations, which are hard to simulate in testing. Addressing cognitive function performance can be done using both cognitive assessment by clinical evaluation combined with in-vehicle sensors may result in better results. Further studying such results needs to be done in the future.




Fig. 12 Plots for Participant with Non-MCI on Monthly-Basis

5.2 Future direction

The advancement of in-vehicle sensor technology represents a pivotal avenue for enhancing driver safety and monitoring. The ongoing refinement of these sensors holds the promise of providing more accurate and comprehensive data on various aspects of driving behavior. As these sensors become more sophisticated, they offer the potential to capture nuanced details, contributing to a deeper understanding of driver actions and responses. Additionally, longitudinal studies assessing the effectiveness of sensor-based monitoring over time are crucial



Fig. 13 Long Distance U-Turns for Road Segments [21]

Fig. 14 Cyclic Patterns for Road Segments [21]



for evaluating the sustained impact and reliability of these technologies. Such studies can shed light on the durability of sensor-based assessments, their ability to adapt to evolving driving habits and their long-term contribution to overall road safety. Together, the continued improvement of in-vehicle sensor technology and longitudinal studies can significantly advance the capabilities in assessing and promoting safer driving practices. Further alongside this, a combination of both the clinical assessment and in-vehicle sensor data can be fused together and assessed to see how well the finding represent. A score-based system should be introduced that evaluates all the data and assigns a performance score to each individual.

6 Conclusion

This study encapsulates key findings at the intersection of aging, cognition, and transportation safety. It reveals notable changes in reaction times, attention spans, and decision-making abilities among older individuals, emphasizing the necessity for targeted interventions to mitigate safety risks. Contributing significantly to the field, the research bridges gaps in understanding how cognitive changes associated with aging impact driving performance, offering crucial insights for researchers, practitioners, and policymakers. The study's recommendations advocate for evidence-based interventions like cognitive training programs and adaptive driving technologies, as well as flexible and inclusive transportation policies to address the diverse needs of aging drivers. Ultimately, this synthesis aims to guide policy and practice in creating a transportation environment that prioritizes both safety and mobility for the aging population.

Acknowledgements This study is funded by the U.S. National Institute of Health's (NIH) National Institute on Aging under grant number R01-AG068472-02S1

Data Availability The data associated with this journal paper will be provided upon request.

Declarations

Conflict of Interest Muhammad Tanveer Jan would like to disclose that Borko Furht is an author of this manuscript and also serves as the Editor-in-Chief of Multimedia Tools and Applications. To ensure transparency, Borko Furht has recused himself from any editorial decisions regarding the handling and review of this manuscript. All editorial decisions have been made independently by other members of the editorial board.

References

1. Eramudugolla R, Huque MH, Wood J, Anstey KJ (2021) On-road behavior in older drivers with mild cognitive impairment. *J Am Med Dir Assoc* 22(2):399–405
2. Ott BR, Heindel WC, Papanonatos GD, Festa EK, Davis JD, Daiello LA, Morris JC (2008) A longitudinal study of drivers with Alzheimer disease. *Neurology* 70(14):1171–8
3. Roe CM, Babulal GM, Head DM, Stout SH, Vernon EK, Ghoshal N, Garland B, Barco PP, Williams MM, Johnson A, Fierberg R (2017) Preclinical Alzheimer's disease and longitudinal driving decline. *Alzheimer's Dement: Transl Res Clin Interv* 3(1):74–82
4. Anstey KJ, Wood J, Kerr G, Caldwell H, Lord SR (2009) Different cognitive profiles for single compared with recurrent fallers without dementia. *Neuropsychology* 23(4):500
5. Ball KK, Roenker DL, Wadley VG, Edwards JD, Roth DL, McGwin G Jr, Raleigh R, Joyce JJ, Cissell GM, Dube T (2006) Can high-risk older drivers be identified through performance-based measures in a Department of Motor Vehicles setting? *J Am Geriatr Soc* 54(1):77–84


6. Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12(3):189–198
7. Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL, Chertkow H (2005) The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 53(4):695–9
8. Dawson JD, Uc EY, Anderson SW, Johnson AM, Rizzo M (2010) Neuropsychological predictors of driving errors in older adults. *J Am Geriatr Soc* 58(6):1090–6
9. Patrick K, Azar M, Bosenbark D, Schultheis MT (2020) Reliability of a virtual reality driving simulator for individuals with neurological disorders. *Technol Disabil* 32(1):25–31
10. Horrey WJ, Lesch MF, Garabet A (2008) Assessing the awareness of performance decrements in distracted drivers. *Accid Anal Prev* 40(2):675–82
11. Dingus TA, Guo F, Lee S, Antin JF, Perez M, Buchanan-King M, Hankey J (2016) Driver crash risk factors and prevalence evaluation using naturalistic driving data. *Proc Natl Acad Sci* 113(10):2636–41
12. Mehler B, Reimer B, Coughlin JF, Dusek JA (2009) Impact of incremental increases in cognitive workload on physiological arousal and performance in young adult drivers. *Transp Res Rec* 2138(1):6–12
13. Bosurgi G, Pellegrino O, Ruggeri A, Sollazzo G (2023) The effects of ADAS on driving behavior: a case study. *Sensors* 23(4):1758
14. Eskandarian A (2012) Handbook of intelligent vehicles. (No Title)
15. Altaher A, Salekshahrezaee Z, Abdollah Zadeh A, Rafiepour H, Altaher A (2020) Using multi-inception CNN for face emotion recognition. *J Bioeng Res* 3(1):1–2
16. Roberts RO, Knopman DS, Mielke MM, Cha RH, Pankratz VS, Christianson TJ, Geda YE, Boeve BF, Ivnik RJ, Tangalos EG, Rocca WA (2014) Higher risk of progression to dementia in mild cognitive impairment cases who revert to normal. *Neurology* 82(4):317–25
17. Flach T, Mishra N, Pedrosa L, Riesz C, Govindan R (2011) Carma: towards personalized automotive tuning. In *Proceedings of the 9th ACM Conference on Embedded Networked Sensor Systems* (pp. 135–148)
18. Jan MT, Moshfeghi S, Conniff JW, Jang J, Yang K, Zhai J, Rosselli M, Newman D, Tappen R, Furht B. (2022) Methods and Tools for Monitoring Driver's Behavior. In *2022 International Conference on Computational Science and Computational Intelligence (CSCI)* (pp. 1269–1273). IEEE
19. Jan MT, Hashemi A, Jang J, Yang K, Zhai J, Newman D, Tappen R, Furht B (2022) Non-intrusive drowsiness detection techniques and their application in detecting early dementia in older drivers. In *Proceedings of the Future Technologies Conference* (pp. 776–796). Cham: Springer International Publishing
20. Moshfeghi S, Jan MT, Conniff J, Ghoreishi SG, Jang J, Furht B, Yang K, Rosselli M, Newman D, Tappen R, Smith D (2023) In-Vehicle Sensing and Data Analysis for Older Drivers with Mild Cognitive Impairment. In *2023 IEEE 20th International Conference on Smart Communities: Improving Quality of Life using AI, Robotics and IoT (HONET)* (pp. 140–145). IEEE
21. Ghoreishi SG, Moshfeghi S, Jan MT, Conniff J, Yang K, Jang J, Furht B, Tappen R, Newman D, Rosselli M, Zhai J (2023) Anomalous Behavior Detection in Trajectory Data of Older Drivers. In *2023 IEEE 20th International Conference on Smart Communities: Improving Quality of Life using AI, Robotics and IoT (HONET)* (pp. 146–151). IEEE
22. Tappen R, Newman D, Rosselli M, Jang J, Furht B, Yang K, Ghoreishi SG, Zhai J, Conniff J, Jan MT, Moshfeghi S (2023) Study protocol for "In-vehicle sensors to detect changes in cognition of older drivers". *BMC Geriatr* 23(1):854
23. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, Kawas CH, Klunk WE, Koroshetz WJ, Manly JJ, Mayeux R, Mohs RC (2011) The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 7(3):263–9
24. Hird MA, Egeto P, Fischer CE, Naglie G, Schweizer TA (2016) A systematic review and meta-analysis of on-road simulator and cognitive driving assessment in Alzheimer's disease and mild cognitive impairment. *J Alzheimers Dis* 53(2):713–29
25. Piersma D, Fuermaier AB, De Waard D, Davidse RJ, de Groot J, Doumen MJ, Bredewoud RA, Claesen R, Lemstra AW, Vermeeren A, Ponds R (2016) Prediction of fitness to drive in patients with Alzheimer's dementia. *PLoS ONE* 11(2):e0149566
26. Uc EY, Rizzo M, Anderson SW, Shi Q, Dawson JD (2004) Driver route-following and safety errors in early Alzheimer disease. *Neurology* 63(5):832–7
27. Uc EY, Rizzo M, Anderson SW, Shi Q, Dawson JD (2005) Driver landmark and traffic sign identification in early Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 76(6):764–8

28. Rizzo M, Reinach S, McGehee D, Dawson J (1997) Simulated car crashes and crash predictors in drivers with Alzheimer disease. *Arch Neurol* 54(5):545–51
29. Ernst J, Krapp S, Schuster T, Förstl H, Kurz A, Diehl-Schmid J (2010) Car driving ability of patients with frontotemporal lobar degeneration and Alzheimer's disease. *Nervenarzt* 81:79–85
30. Carr DB (1997) Motor vehicle crashes and drivers with DAT. *Alzheimer Dis Assoc Disord* 11:38–41
31. Wadley VG, Okonkwo O, Crowe M, Vance DE, Elgin JM, Ball KK, Owsley C (2009) Mild cognitive impairment and everyday function: an investigation of driving performance. *J Geriatr Psychiatry Neurol* 22(2):87–94
32. Fuermaier A, Piersma D, De Waard D, Davidse RJ, De Groot J, Doumen MJ, Bredewoud RA, Claesen R, Lemstra AW, Scheltens P, Vermeeren A (2019) Driving difficulties among patients with Alzheimer's disease and other neurodegenerative disorders. *J Alzheimers Dis* 69(4):1019–30
33. Carr DB, Duchek J, Morris JC (2000) Characteristics of motor vehicle crashes of drivers with dementia of the Alzheimer type. *J Am Geriatr Soc* 48(1):18–22
34. Wang C, Musaev A, Sheinidashtegol P, Atkinson T (2019) Towards detection of abnormal vehicle behavior using traffic cameras. In *Big Data-BigData 2019: 8th International Congress, Held as Part of the Services Conference Federation, SCF 2019, San Diego, CA, USA, June 25-30, 2019, Proceedings 8* (pp. 125-136). Springer International Publishing
35. Abdelrahman AE, Hassanein HS, Abu-Ali N (2020) Robust data-driven framework for driver behavior profiling using supervised machine learning. *IEEE Trans Intell Transp Syst* 23(4):3336–50
36. Wang R, Xie F, Zhao J, Zhang B, Sun R, Yang J (2020) Smartphone sensors-based abnormal driving behaviors detection: Serial-feature network. *IEEE Sens J* 21(14):15719–28
37. Ali AH, Atia A, Mostafa MS (2017) Recognizing driving behavior and road anomaly using smartphone sensors. *Int J Ambient Comput Intell* 8(3):22–37
38. Bahadoor K, Hosein P. Application for the detection of dangerous driving and an associated gamification framework. In *2016 IEEE 4th International Conference on Future Internet of Things and Cloud Workshops (FiCloudW) 2016 Aug 22* (pp. 276-281). IEEE
39. Babić D, Babić D, Cajner H, Sruk A, Fiolić M (2020) Effect of road markings and traffic signs presence on young driver stress level, eye movement and behaviour in night-time conditions: a driving simulator study. *Safety*. 6(2):24
40. Awan HH, Pirdavani A, Houben A, Westhof S, Adnan M, Brijs T (2019) Impact of perceptual countermeasures on driving behavior at curves using driving simulator. *Traffic Inj Prev* 20(1):93–9
41. Eren H, Makinist S, Akin E, Yilmaz A (2012) Estimating driving behavior by a smartphone. In *2012 IEEE Intelligent Vehicles Symposium* (pp. 234-239). IEEE
42. Bao C, Chen C, Kui H, Wang X (2019) Safe driving at traffic lights: An image recognition based approach. In *2019 20th IEEE International Conference on Mobile Data Management (MDM)* (pp. 112-117). IEEE
43. Wang Q, Liu Y, Liu J, Gu Y, Kamijo S (2018) Critical areas detection and vehicle speed estimation system towards intersection-related driving behavior analysis. In *2018 IEEE International Conference on Consumer Electronics (ICCE)* (pp. 1-6). IEEE
44. Nassajpour M, Shuqair M, Rosenfeld A, Tolea MI, Galvin JE, Ghoraani B (2024) Objective estimation of m-CTSIB balance test scores using wearable sensors and machine learning. *Front Digit Health* 6:1366176

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Muhammad Tanveer Jan¹  · Borko Furht¹ · Sonia Moshfeghi¹ · Jinwoo Jang¹ · Seyedeh Gol Ara Ghoreishi¹ · Charles Boateng¹ · Kwangsoo Yang¹ · Joshua Conniff² · Monica Rosselli² · David Newman³ · Ruth Tappen³

✉ Muhammad Tanveer Jan
mjan2021@fau.edu

Borko Furht
bfurht@fau.edu

Sonia Moshfeghi
smoshfeghi2021@fau.edu

Jinwoo Jang
jangj@fau.edu

Seyedeh Gol Ara Ghoreishi
sghoreishi2021@fau.edu

Charles Boateng
cboateng2023@fau.edu

Kwangsoo Yang
yangk@fau.edu

Joshua Conniff
fau.jconniff@health.fau.edu

Monica Rosselli
mrosselli@fau.edu

David Newman
dnewma14@fau.edu

Ruth Tappen
rtappen@fau.edu

¹ Department of Electrical Engineering & Computer Science, Florida Atlantic University, Boca Raton, FL 33431, USA

² Charles E. Schmid College of Science, Florida Atlantic University, Boca Raton, FL 33431, USA

³ Christine E. Lynn College of Nursing, Florida Atlantic University, Boca Raton, FL 33431, USA