



Bone Marrow Adipocyte Developmental Origin and Biology

Joanna Bukowska¹ · Trivia Frazier^{2,3} · Stanley Smith² · Theodore Brown⁴ · Robert Bender² · Michelle McCarthy⁴ · Xiyang Wu^{2,3} · Bruce A. Bunnell⁴ · Jeffrey M. Gimble^{2,3,4}

Published online: 18 April 2018

© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract

Purpose of Review This review explores how the relationships between bone marrow adipose tissue (BMAT) adipogenesis with advancing age, obesity, and/or bone diseases (osteopenia or osteoporosis) contribute to mechanisms underlying musculoskeletal pathophysiology.

Recent Findings Recent studies have re-defined adipose tissue as a dynamic, vital organ with functions extending beyond its historic identity restricted solely to that of an energy reservoir or sink. “State of the art” methodologies provide novel insights into the developmental origin, physiology, and function of different adipose tissue depots. These include genetic tracking of adipose progenitors, viral vectors application, and sophisticated non-invasive imaging modalities.

Summary While constricted within the rigid bone cavity, BMAT vigorously contributes to local and systemic metabolic processes including hematopoiesis, osteogenesis, and energy metabolism and undergoes dynamic changes as a function of age, diet, bone topography, or sex. These insights will impact future research and therapies relating to osteoporosis.

Keywords Beige cells · Bone marrow · Brown adipose tissue · Cytomegalovirus · White adipose tissue

Introduction to the Types of Adipocytes and Adipose Tissues

Adipose tissue is found throughout the body and is distinguished morphologically by the presence of adipocytic cells containing prominent lipid containing vacuoles. Currently, adipocytes can be categorized based on their metabolic/physiologic function, developmental origins, and characteristic protein biomarkers (Table 1). The most abundant human adult or white adipocyte, localized to white adipose tissue (WAT), serves multiple functions including energy storage,

adipokines secretion, immunomodulatory roles, and endocrine organ. Developmentally, white adipocytes are postulated to derive from resident adipose stromal/stem cells (ASC) or mesenchymal stromal/stem cells (MSC). These have been characterized based on flow cytometry as a hematopoietic lineage negative, CD31⁻ (non-endothelial cell), Sca1⁺, and CD29⁺ CD34⁺ CD24⁺ populations [1, 2]. Multiple studies of extramedullary adipose depots have documented the expression of pericytic and perivascular markers such as CD146 and 3G5 on ASC and MSC [3, 22, 23]. This has led to the hypothesis that capillary mural wall-associated pericytes are the original adipocyte progenitor cells; however, morphological and fate mapping studies now question this conclusion. Lineage tracing in transgenic mice using the pericytic lineage-specific Tbx18 transcription factor indicate that this cell population does not contribute directly to the formation of adipose and other tissues [24••]. Furthermore, there is in vitro and in vivo murine evidence associating adipogenic progenitors and adipose differentiation with the CD34 biomarker alone, exclusive of the CD146 pericytic antigen or a perivascular anatomical location [22, 25–27].

Brown adipocytes, localized to brown adipose tissues (BAT) located in the neck and around vital organs of newborn infants,

This article is part of the Topical Collection on *Bone Marrow and Adipose Tissue*

✉ Jeffrey M. Gimble
jgimble@tulane.edu

¹ Institute of Animal Reproduction and Food Research, Polish Academy of Sciences, Olsztyn, Poland

² LaCell LLC, New Orleans, LA, USA

³ Obatala Sciences, Inc., New Orleans, LA, USA

⁴ Center for Stem Cell Research and Regenerative Medicine, Tulane University School of Medicine, New Orleans, LA, USA

Table 1 Adipose tissue categories and their characteristic features

Tissue type	Tissue function	Developmental origin	Specific markers ^a	References
White adipose tissue (WAT)	Energy storage as triglycerides; Coordination of systemic metabolism (i.e., glucose metabolism); Insulation; Mechanical support for internal organs Endocrine organ (secretes adipokines: leptin, adiponectin, resistin, apelin, visfatin); Immunomodulation; Role in toxicokinetics of a persistent organic pollutant (POP);	Derived from a <i>Myf5</i> ⁻ precursors	Leptin; Resistin; Adiponectin	[1–11]
Brown adipose tissue (BAT)	Energy dissipation and heat production (transferring energy from nutrients to heat in the process involving UCP1 as a main agent that collapses the electron gradient to generate heat rather than ATP); Body weight regulation; Counteracting metabolic diseases (obesity, type 2 diabetes); Critical to newborn infants survival	Derived from <i>Myf5</i> ⁺ and <i>Pax7</i> ⁺ progenitor cells that also give rise to skeletal muscle	Uncoupling protein 1 (UCP1); PPAR γ coactivator-1 α (PGC-1 α); PR domain zinc finger protein 16 (PRDM16); Type 2 iodothyronine deiodinase (DIO2); V-like antigen 1 (EVA1); also known as myelin protein zero-like 2 (MPZL2)	[1–11]
Beige/brite adipose tissue	Thermogenic function; Counteracting metabolic diseases (obesity, type 2 diabetes)	Derived from a <i>Myf5</i> ⁻ precursors	Homeobox C9 (HOXC9); Transmembrane protein 26 (TMEM26); T-box 1 (<i>Tbx1</i>); Tumor necrosis factor receptor superfamily member 9 (TNFRSF9) also known as 4-1BB or CD137; Cytochrome c oxidase subunit 8B (COX8B); Cell death activator (CIDEA); Fatty acid elongase 3 (ELOVL3); Fibroblast growth factor 21 (FGF21)	[5–9, 11–17]
Bone marrow adipose tissue (BMAT; MAT)	Regulates hematopoietic stem cell biology; Contributes to hematopoiesis; participates in bone remodeling; bone lipid metabolism; Secretory tissue (releases leptin, adiponectin)	Some BMAT derived from a <i>Myf5</i> ⁻ precursors than can differentiate into osteoblasts and WAT; however, both BAT and beige are present in the bone marrow	Specific markers were not identified; however, it has been reported that high expression of BAT markers including PRDM16, PGC1 α , DIO2, and low levels of WAT markers (adiponectin and leptin), as well as BAT-like characteristics decrease with aging and diabetes.	[18, 19, 20, 21]

^a Adipose tissues share transcriptional cascade including peroxisome proliferator-activated receptor (PPAR)- and CCAAT/enhancer-binding proteins (C/EBPs) [6]

are less abundant than white adipocytes and display thermogenic properties. With the advent of non-invasive metabolic imaging methods monitoring radiolabeled glucose, BAT can be routinely visualized and its persistence in adult humans is now accepted [28, 29, 30]. Due to their expression of uncoupling protein 1 (UCP1), brown adipocytes convert energy directly into heat by acting to short circuit the mitochondrial membrane [31, 32]. Developmentally, brown adipocytes are more closely related to skeletal muscle progenitors than to white adipocytes and ASC. Additional brown adipocyte biomarkers include the transcription co-factors PR domain 16 (Prdm16) and PPAR γ co-activator 1 α (PGC1 α) [12, 33, 34].

Beige or brite (brown/white) adipocytes are a metabolically intermediate population of adipocytes. Like brown adipocytes, beige/brite adipocytes express UCP1 and are capable of thermogenic function; however, they are localized to subcutaneous and other white-associated adipose tissues under inductive conditions such as cold exposure and are developmentally linked to ASC and MSC rather than skeletal muscle [35]. Furthermore, beige/brite adipocytes express distinct biomarkers including *Tbx1*, TMEM16, *HoxC9*, CIDEA, CITED1, and CD137 [13–15]. There is substantial interest in developing pharmacological methods to induce beige/brite adipocytes as a mechanism to combat obesity [36].

Bone marrow adipose tissue (BMAT) (also known as marrow adipose tissue (MAT) or yellow adipose tissue (YAT)) is an adipose depot with unique features distinguishing it from the better characterized extramedullary sites. While the mechanical constraints impacting extramedullary subcutaneous and visceral adipose depots are due to soft tissue compressive forces, those faced by BMAT are more rigid due to the structural composition of trabecular and cortical bone. Furthermore, while extramedullary adipose depots contain cells capable of hematopoiesis, only BMAT supports this critical cell differentiation event on a daily high volume/turnover basis [37, 38]. There is a growing body of evidence indicating that the anatomical uniqueness of BMAT may be accompanied by a developmental origin distinct from both BAT and WAT [18, 39]. Bone marrow fat does not comprise a single, homogenous tissue. Indeed, BMAT may exist in two distinct populations: “regulated” (rMAT) that may influence hematopoiesis and “constitutive” BMAT (cMAT) important during early vertebrate development [19]. Although constricted within the rigid bone cavity, BMAT participates in both local and systemic metabolic processes [40]. Additionally, considering its contributions to hematopoiesis [19], osteogenesis [41], energy metabolism [20••], and as a functional endocrine tissue [21], there is a growing appreciation that BMAT is a functional, dynamic organ. This review evaluates recent advances in studies of the anatomy, development, and function of BMAT with particular attention directed towards cell tracking/fate mapping with genetic models, brown/beige mechanisms, and their contribution to marrow fat physiology. Additionally, the authors extrapolate from extramedullary adipose studies of viral infection to identify potential avenues for future investigations into the developmental origins and function of BMAT.

Age and Disease Dependent Time Course of Marrow Adipogenesis Using Non-invasive Methods

While routine clinical bone imaging methods, dual-energy X-ray-absorptiometry (DXA), and quantitative computed tomography (QCT) assess bone mineral density, they less effectively quantify BMAT and related non-mineralized bone components. To address this need, new imaging techniques have been developed to enable precise in situ characterization of BMAT. Positron emission tomography/computed tomography (PET/CT) scans determine tissue metabolic activity based on their uptake of radiolabeled pharmaceuticals (natural substrates including glucose and fatty acid, substrates analogs or drugs) that are injected intravenously or inhaled by the subject. By using PET and the glucose analog ^{18}F -fluoro-deoxy-glucose (^{18}F -FDG) as a tracer, it has been possible to extend studies concerning human BAT physiology in the context of

body-mass index (BMI) and ambient temperature [42]. The PET techniques have been used to evaluate bone marrow composition and function in patients with multiple myeloma [43], follicular lymphoma [44], and related disorders [45]. Likewise, high-resolution magnetic resonance imaging (MRI) and magnetic resonance spectroscopy (MRS) are alternative non-invasive techniques for quantification of BMAT [46, 47]. These techniques have provided new insights into bone pathophysiology in the context of metabolic diseases and cancers. MRI visualizes BM soft tissues based on their chemical composition and profile. The relative differences between fat, water, and protein ratios observed between red bone marrow and BMAT serve as quantitative and identifiable signatures relevant to clinical diagnostic and therapeutic purposes. Furthermore, MRI has considerable advantage due to its lack of ionizing radiation, excellent tissue contrast and ability to quantify low amount of fat [48]. Both MRI and MRS can be combined with bone mineral density (BMD) or DEXA to explore the bone-fat dynamic in laboratory animals. Scheller et al. [49] quantified BMAT volume accurately by staining bone with osmium tetroxide followed by μCT scanning or more precise nano-computed tomography (nanoCT). These studies revealed that single adipocytes that are spread with regions of active hematopoiesis (regulated MAT; rMAT) when exposed to cold temperature were reduced in size and number within the proximal tibia but remained unchanged in the distal tibia. This observation lends support to the hypothesis that region-specific differences exert regulatory control over adipogenesis within BMAT. Likewise, sophisticated proton magnetic resonance spectroscopy (^1H -MRS) evaluates not only total bone marrow fat content but also provides information about the fatty acid composition (assessment of saturated and unsaturated fatty acids) based on MR spectra [50, 51]. Using 3T full-body MRI scanner, Pansini et al. [47] compared normal spectroscopic bone marrow fat content (FC) between different regions of the hip in the context of subject age and sex. They further calculated the spectroscopic conversion index (SCI) based on the FC neck/FC greater trochanter ratio. While SCI increased with age in both genders, it remained lower in women compared to men, consistent with a role for sex steroids in controlling BM conversion. Together, these non-invasive clinical imaging tools will provide a comprehensive picture of BMAT adipogenesis as a function of age and disease status in humans as well as preclinical animal models.

Developmental Origin and Fate Map Tracking of Adipose Progenitor Cells

Bone marrow adipose tissue has long been recognized as the “yellow” bone marrow characterized by a metabolic profile and functions distinct from extramedullary fat depots. Current data reveal that BMAT plays vital role in bone remodeling;

this is highlighted by impaired bone alterations observed in a spectrum of diseases associated with increased marrow adipogenesis [51–53]. However, in healthy subjects, development of BMAT depends, among others, on age, bone topography, and sex. Furthermore, recent data in C57BL/6J (B6) male mice indicates that high-fat diet-induced obesity increased BMAT without altering bone mineral density (BMD) [54]. In humans, during intrauterine life and shortly after birth, bone cavities contain mainly hematopoietic red bone marrow. Deposition of BMAT starts during the childhood and lasts throughout adulthood. Adipose tissue expansion primarily occurs in the distal parts of the appendicular skeleton, then spreads towards more proximal portions whereas central bones (i.e., pelvis, ribs) retain red BM throughout life [47, 55]. By the age of 25 years, BMAT occupies approximately 70% of the BM volume and can reach up to 5% of total fat mass in adults [49]. Interestingly, a study by Griffith et al. shown that conversion on red BM into BMAT occurs with age and this process increases particularly sharply in women at menopause. Compared to age matched males, menopausal women displayed 10% higher levels of BMAT content [56].

Genetic Tracking of Adipose Progenitors

Recent studies have employed inducible genetic lineage tracing systems to monitor and quantify adipose tissue formation in vivo. In order to track adipogenesis in mice fed a high fat diet (HFD; 60% of calories from fat) or in response to cold (6 °C) or β -3 agonist exposure, Wang et al. developed the “AdipoChaser” mouse model [57]. This triple transgenic system uses doxycycline to permanently label mature adipocytes in white adipose tissue (WAT). During the early stage of HFD exposure (1 month), hypertrophy accounted for the development of both gonadal and subcutaneous fat. In contrast, prolonged HFD feeding (2 months) resulted in high capacity adipogenesis exclusively within the gonadal (epididymal) fat depots. Furthermore, exposure to cold or β -3 agonists induced gonadal WAT formation and the de novo appearance of beige/brite adipocytes within subcutaneous fat depots [57]. Similar genetic lineage tracing approaches examining cold-induced brown adipocytes (BA) have determined a distinct origin for newly appearing BA in the interscapular BAT (iBAT) and subcutaneous inguinal WAT. While iBAT derive from undifferentiated resident PDGFR α + progenitors, multilocular BA in the inguinal WAT derive exclusively from preexisting “white” adipocytes [58]. Focusing on committed osteoblast progenitors, Mizoguchi et al. used both *Lepr*/Tomato⁺ and P5-iOx/Tomato+ transgenic mice to evaluate marrow adipogenesis following irradiation or physical injury. Adipogenesis increased significantly following a pulsed 6 Gy radiation injury while P5-iOx/Tomato+ were detected in the newly formed chondrogenic tissue of the fracture area that also

expressed the chondrocyte marker Sox9. Together, this suggests that progenitor cells expressing *Lepr* or P5-iOx can be a source of osteo-, adipo-, and chondro-precursors and they contribute to granulation tissue formation and healing of the injured area [59]. In an independent study, Zhou et al. determined a role for leptin receptor *LepR*+ (*Lepr-cre; tdTomato*) mesenchymal stem cells (MSC) in adipogenesis and osteogenesis of adult mice; however, their role in prenatal bone formation was restricted [60] showing rare distribution in the bone marrow at embryonic day (E) 19.5. However, as early as postnatal day (P) 0.5, a sharp increase appeared in the number of Tomato+ cells that further increased with age.

While fate mapping showed that *Lep-cre* MSC cells were a rare population in adult murine bone marrow, representing ~ 0.3% of cells, their number increased considerably within the callus formed immediately following a bone fracture. Moreover, intrafemoral injection of *LepR*+ cells led to the appearance of Tomato signal positive osteoblasts, adipocytes, and chondrocytes. These findings support the hypothesis that *LepR*+ derived adipocytes and osteoblast contribute to bone marrow renewal and regeneration [60, 61]. However, studies conducted with mice expressing a conditional deletion of *LepR* in bone marrow MSC (*Prx1-Cre; LepR^{fl/fl}*) indicate that while *LepR* is required for increased bone marrow adipogenesis in response to an HFD, the receptor is not required for bone marrow adipose tissue formation following radiation injury [61]. Indeed, *LepR* deficiency in *Prx1-Cre; LepR^{fl/fl}* mice correlated with increased femoral trabecular bone volume relative to the controls. While irradiation increased adipocyte numbers in the marrow, the impact was comparable between *LepR*-deficient mice and their controls. While HFD feeding increased marrow adipogenesis at the expense of osteogenesis in control mice, bone marrow *LepR*-deficient mice displayed improved bone formation and reduced adipogenesis. This suggests that HFD exhibits leptin-dependent effects not only through the central hypothalamic system but also at the level of the MSC. Evidence supports a mechanistic role for leptin-dependent downstream activation of the *Jak2/Stat3* signaling pathway and *Cebpa* transcription resulting in *Wnt4* downregulation as a regulator of bone marrow adipogenesis [61]. Ambrosi et al. have examined BMAT adipogenesis in the context of aging and high fat diet using lineage tracing based on reporters driven by the promoters of identifying hematopoietic (*Vav1*), endothelial (*Cdh5/Tie2*), MSC (*PDGFR α*), and committed adipocytes (*Adipoq*) [62]. They noted that high fat diet with or without aging increased marrow adipogenesis and did so at the expense of hematopoiesis and osteogenesis [62]. The authors further validated their findings based on flow cytometric analysis and in vitro tri-lineage differentiation analyses [62]. Together, this body of work confirms the value of genetic lineage cell tracking in dissecting the developmental dynamics of bone marrow adipogenesis under physiological and pathological conditions. Further studies with these in vivo

genetic-based tools in the context of tumor formation and metastasis are warranted.

Role of Viral Vectors in Promoting or Inhibiting Adipogenic Development

There is a growing appreciation of the potential role of viral vectors as etiological agents responsible for adipogenesis and the development of obesity. The term “infectobesity,” first introduced by Dhurandhar [63], reflects the concept of close association between certain infections and obesity in human [64], non-human primate [65], and rodents [66]. Multiple viral agents have been reported to influence adipose tissue biology [67, 68]. To date, the most extensive studies have focused on human adenovirus type 36 (Ad-36) which increases adiposity and paradoxically reduces the concentrations of cholesterol and triglycerides in experimental infected animals [66]. Infection with Ad-36 had multiple effects on differentiation of 3T3-L1 preadipocytes cell line including increase in the cells numbers that differentiate into adipocytes, enhanced lipid accumulation, and upregulation of the adipocyte marker enzyme GPDH [68]. The *in vitro* induction of the adipogenic program of both rodent (3T3-L1 cell line) and human (hASC) adipocyte progenitors involved the Ad-36 viral E4 open reading frame (orf)-1 gene. Introduction of this gene was sufficient to upregulate cAMP levels, phosphatidylinositol 3-kinase (PKB) activity, and C/EBP- β expression [69]. In contrast, infection with human cytomegalovirus (HCMV) significantly reduced the adipogenic differentiation capacity of hASC [70]. Accordingly, HCMV-infected cells developed fewer lipid droplets when compared with mock-infected cells cultured under adipogenic differentiation conditions. Furthermore, the presence of HCMV altered lipid droplet formation and/or stability. While vacuoles in infected cells aggregated in the perinuclear region, these organelles were distributed regularly throughout the cytoplasm of uninfected control hASC. Parallel analyses by RT-PCR confirmed the downregulation of adipogenic-associated genes, e.g., peroxisome proliferator-activated receptor gamma (*PPAR* γ), adiponectin, fatty acid binding protein 4 (*FABP4*), lipoprotein lipase (*LPL*), and perilipin 1. Moreover, HCMV infection inhibited ASC expression of transcripts of anti-inflammatory mediators including *TSG-6* and *IL-6*. These studies emphasize the importance of determining the HCMV seropositivity of both ASC-donors and recipients since viral status can impact hASC immunomodulatory function and subsequent clinical outcome. Additionally, they raise questions about the role of adenoviral and cytomegaloviral infections as contributors to the regulation of bone marrow adipogenesis.

Conclusions and Future Directions

Studies during the past few years have substantially advanced the understanding of BMAT and its developmental origins; however, additional questions remain unanswered and merit further consideration. These include the following:

Are beige/brite mechanisms responsible for bone marrow adipogenesis seen with advancing age, osteopenia, and osteoporosis? There is a growing body of data indicating that bone marrow contains beige/brite adipocytes. Studies of heterotopic ossification in human and rodent models have detected UCP1, consistent with the presence of beige/brite or brown adipocytes [71, 72].

1. These findings indicate that beige/brite cells are associated and co-localized with peripheral nerves [73, 74]. These are intriguing observations and further work is necessary to determine if the same progenitor population contributes to age associated adipogenesis. Are beige/brite adipocytes an intermediate step in the pathway towards the acquisition of a yellow marrow? Can the endocrine/bone community re-purpose drugs developed for the treatment of obesity and metabolic syndrome targeting the beige/brown pathway to re-direct or commandeer the BMAT differentiation pathway, thereby promoting bone growth rather than age-associated osteopenia and osteoporosis?
2. Are viral agents, including but not limited to adenovirus and cytomegalovirus, contributors to bone marrow adipogenesis? The ability of viral vectors to alter adipogenesis presents an alternative view of extramedullary obesity as a possible infectious disease. This may have an equivalent impact on BMAT formation. Since Ad-36 and CMV induce adipogenic markers associated with a white or beige/brown phenotype, respectively, this raises questions concerning the role of viral infection or re-activation in the acquisition of marrow adiposity and bone loss. Additionally, it suggests studies examining the contribution of other common viral infections, such as hepatitis and HIV, to BMAT pathophysiology at the progenitor cell level.
3. Is extramedullary obesity linked directly to bone marrow adiposity? (Corollary: How does nutrition alter bone marrow adipogenesis?) This concept is by no means new as evidenced by the intriguing title of highly cited review article: “Mechanisms of Disease: Is Osteoporosis Obesity of the Bone Marrow?” [75].
4. There remains considerable interest in the links between high fat diets and marrow adipogenesis [4, 62]. Studies have found that male C57BL6 mice fed a high fat diet displayed increased bone mass relative to controls secondary to the mechanical strain of an increased body mass index; however, it was also associated with a reduced rate of bone formation and turnover [4].

5. High fat diet studies have been found to increase the number of lineage committed adipogenic progenitors in murine marrow, and this is enhanced with advancing age [62]. The production of dipeptidyl peptidase 4 (Dpp4) by the marrow adipocytes exerted a negative impact on bone repair and osteogenesis during fracture healing in this model [62]. These findings further confirm the importance of bone marrow adipocytes as a therapeutic target in the treatment of osteopenia, osteoporosis, and their comorbidities of fracture and sarcopenia.

Acknowledgements The authors thank Barbara Gawronska-Kozak Ph.D. for her critical review of the manuscript.

Compliance with Ethical Standards

Conflict of Interest Xiyang Wu and Smith report a patent on use of adipose cells in therapy in the submission process from LaCell LLC.

Jeff Gimble is a co-founder and co-owner of Talaria Antibodies, a polyclonal antibody production company; Obatala Sciences, a fat on a chip technology; and is a co-founder, co-owner, and employee at LaCell LLC. Xiyang Wu is a co-founder, co-owner, and R&D Director at LaCell LLC; a co-owner and co-founder of Obatala Sciences; and reports a patent on use of adipose cells in therapy in the submission process from LaCell LLC. Stanley Smith reports a patent on use of adipose cells in therapy in the submission process from LaCell LLC. Michelle McCarthy, Trivia Frazier, Joanna Bukowska, Theodore Brown, Robert Bender, and Bruce Bunnell declare no conflict of interest.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

1. Rodeheffer MS, Birsoy K, Friedman JM. Identification of white adipocyte progenitor cells in vivo. *Cell*. 2008;135:240–9.
2. Berry R, Rodeheffer MS. Characterization of the adipocyte cellular lineage in vivo. *Nat Cell Biol*. 2013;15:302–8.
3. Crisan M, Yap S, Casteilla L, Chen CW, Corselli M, Park TS, et al. A perivascular origin for mesenchymal stem cells in multiple human organs. *Cell Stem Cell*. 2008;3:301–13.
4. Lecka-Czernik B, Stechschulte LA, Czernik PJ, Dowling AR. High bone mass in adult mice with diet-induced obesity results from a combination of initial increase in bone mass followed by attenuation in bone formation; implications for high bone mass and decreased bone quality in obesity. *Mol Cell Endocrinol*. 2015;410:35–41.
5. Chu DT, Malinowska E, Gawronska-Kozak B, Kozak LP. Expression of adipocyte biomarkers in a primary cell culture models reflects preweaning adipobiology. *J Biol Chem*. 2014;289:18478–88.

6. Wang W, Seale P. Control of brown and beige fat development. *Nat Rev Mol Cell Biol*. 2016;17:691–702.
7. Gimble JM, Katz AJ, Bunnell BA. Adipose-derived stem cells for regenerative medicine. *Circ Res*. 2007;100:1249–60.
8. Coelho M, Oliveira T, Fernandes R. Biochemistry of adipose tissue: an endocrine organ. *Arch Med Sci*. 2013;9:191–200.
9. Berry DC, Stenesen D, Zeve D, Graff JM. The developmental origins of adipose tissue. *Development*. 2013;140:3939–49.
10. La Merrill M, Emond C, Kim MJ, Antignac JP, Le Bizec B, Clement K, et al. Toxicological function of adipose tissue: focus on persistent organic pollutants. *Environ Health Perspect*. 2013;121:162–9.
11. Giralt M, Villarroya F. White, brown, beige/brite: different adipose cells for different functions? *Endocrinology*. 2013;154:2992–3000.
12. Harms M, Seale P. Brown and beige fat: development, function and therapeutic potential. *Nat Med*. 2013;19:1252–63.
13. Carey AL, Vorlander C, Reddy-Luthmoodoo M, Natoli AK, Formosa MF, Bertovic DA, et al. Reduced UCP-1 content in in vitro differentiated beige/brite adipocytes derived from preadipocytes of human subcutaneous white adipose tissues in obesity. *PLoS One*. 2014;9:e91997.
14. Lecka-Czernik B, Stechschulte LA, Czernik PJ, Sherman SB, Huang S, Krings A. Marrow adipose tissue: skeletal location, sexual dimorphism, and response to sex steroid deficiency. *Front Endocrinol (Lausanne)*. 2017;8:188.
15. Garcia RA, Roemmich JN, Claycombe KJ. Evaluation of markers of beige adipocytes in white adipose tissue of the mouse. *Nutr Metab (Lond)*. 2016;13:24.
16. Wu J, Bostrom P, Sparks LM, Ye L, Choi JH, Giang AH, et al. Beige adipocytes are a distinct type of thermogenic fat cell in mouse and human. *Cell*. 2012;150:366–76.
17. Lee P, Werner CD, Kebebew E, Celi FS. Functional thermogenic beige adipogenesis is inducible in human neck fat. *Int J Obes*. 2014;38:170–6.
18. Krings A, Rahman S, Huang S, Lu Y, Czernik PJ, Lecka-Czernik B. Bone marrow fat has brown adipose tissue characteristics, which are attenuated with aging and diabetes. *Bone*. 2012;50:546–52.
19. Scheller EL, Doucette CR, Learman BS, Cawthorn WP, Khandaker S, Schell B, et al. Region-specific variation in the properties of skeletal adipocytes reveals regulated and constitutive marrow adipose tissues. *Nat Commun*. 2015;6:7808.
20. •• Rahman S, Lu Y, Czernik PJ, Rosen CJ, Enerback S, Lecka-Czernik B. Inducible brown adipose tissue, or beige fat, is anabolic for the skeleton. *Endocrinology*. 2013;154:2687–701. **Highly Important: First study to characterize the role of beige adipocytes in BMAT.**
21. Cawthorn WP, Scheller EL, Learman BS, Parlee SD, Simon BR, Mori H, et al. Bone marrow adipose tissue is an endocrine organ that contributes to increased circulating adiponectin during caloric restriction. *Cell Metab*. 2014;20:368–75.
22. Traktuev DO, Merfeld-Clauss S, Li J, Kolonin M, Arap W, Pasqualini R, et al. A population of multipotent CD34-positive adipose stromal cells share pericyte and mesenchymal surface markers, reside in a periendothelial location, and stabilize endothelial networks. *Circ Res*. 2008;102:77–85.
23. Zannettino AC, Paton S, Arthur A, Khor F, Itescu S, Gimble JM, et al. Multipotential human adipose-derived stromal stem cells exhibit a perivascular phenotype in vitro and in vivo. *J Cell Physiol*. 2008;214:413–21.
24. •• Guimaraes-Camboa N, Cattaneo P, Sun Y, Moore-Morris T, Gu Y, Dalton ND, et al. Pericytes of multiple organs do not behave as mesenchymal stem cells in vivo. *Cell Stem Cell*. 2017;20:345–59. **e5. Highly Important: This study challenges the concept that pericytes and MSC are equivalent.**

25. Gawronska-Kozak B, Staszkiwicz J, Gimble JM, Kirk-Ballard H. Recruitment of fat cell precursors during high fat diet in C57BL/6J mice is fat depot specific. *Obesity*. 2014;22:1091–102.
26. Frazier TPBA, Lee S, Abbott R, Tucker HA, Kaplan D, Wang M, et al. Serially transplanted non-pericytic CD146- adipose stromal/stem cells in silk bioscaffolds regenerate adipose tissue in vivo. *Stem Cells*. 2016;34:1097–111.
27. Philips BJ, Grahovac TL, Valentin JE, Chung CW, Bliley JM, Pfeifer ME, et al. Prevalence of endogenous CD34+ adipose stem cells predicts human fat graft retention in a xenograft model. *Plast Reconstr Surg*. 2013;132:845–58.
28. Cypess AM, Lehman S, Williams G, Tal I, Rodman D, Goldfine AB, et al. Identification and importance of brown adipose tissue in adult humans. *N Engl J Med*. 2009;360:1509–17.
29. Cypess AM, White AP, Vemochet C, Schulz TJ, Xue R, Sass CA, et al. Anatomical localization, gene expression profiling and functional characterization of adult human neck brown fat. *Nat Med*. 2013;19:635–9. **Important: This study is among the first to provide a detailed analysis of human brown fat.**
30. Virtanen KA, Lidell ME, Orava J, Heglind M, Westergren R, Niemi T, et al. Functional brown adipose tissue in healthy adults. *N Engl J Med*. 2009;360:1518–25.
31. Ricquier D. UCP1, the mitochondrial uncoupling protein of brown adipocyte: a personal contribution and a historical perspective. *Biochimie*. 2017;134:3–8.
32. Nicholls DG. The hunt for the molecular mechanism of brown fat thermogenesis. *Biochimie*. 2017;134:9–18.
33. Kajimura S, Seale P, Kubota K, Lunsford E, Frangioni JV, Gygi SP, et al. Initiation of myoblast to brown fat switch by a PRDM16-C/EBP-beta transcriptional complex. *Nature*. 2009;460:1154–8.
34. Seale P, Kajimura S, Yang W, Chin S, Rohas LM, Uldry M, et al. Transcriptional control of brown fat determination by PRDM16. *Cell Metab*. 2007;6:38–54.
35. Qian S, Huang H, Tang Q. Brown and beige fat: the metabolic function, induction, and therapeutic potential. *Front Med*. 2015;9:162–72.
36. Liu X, Cervantes C, Liu F. Common and distinct regulation of human and mouse brown and beige adipose tissues: a promising therapeutic target for obesity. *Protein Cell*. 2017;8:446–54.
37. Cousin B, Andre M, Arnaud E, Penicaud L, Casteilla L. Reconstitution of lethally irradiated mice by cells isolated from adipose tissue. *Biochem Biophys Res Commun*. 2003;301:1016–22.
38. Han J, Koh YJ, Moon HR, Ryoo HG, Cho CH, Kim I, et al. Adipose tissue is an extramedullary reservoir for functional hematopoietic stem and progenitor cells. *Blood*. 2010;115:957–64.
39. Chen J, Shi Y, Regan J, Karuppaiah K, Ornitz DM, Long F. Osx-Cre targets multiple cell types besides osteoblast lineage in postnatal mice. *PLoS One*. 2014;9:e85161.
40. Lecka-Czemik B. Marrow fat metabolism is linked to the systemic energy metabolism. *Bone*. 2012;50:534–9.
41. Botolin S, McCabe LR. Bone loss and increased bone adiposity in spontaneous and pharmacologically induced diabetic mice. *Endocrinology*. 2007;148:198–205.
42. Gifford A, Towse TF, Walker RC, Avison MJ, Welch EB. Human brown adipose tissue depots automatically segmented by positron emission tomography/computed tomography and registered magnetic resonance images. *J Vis Exp: JoVE*. 2015.
43. Derlin T, Peldschus K, Munster S, Bannas P, Herrmann J, Stubig T, et al. Comparative diagnostic performance of (1)(8)F-FDG PET/CT versus whole-body MRI for determination of remission status in multiple myeloma after stem cell transplantation. *Eur Radiol*. 2013;23:570–8.
44. Perry C, Lerman H, Joffe E, Sarid N, Amit O, Avivi I, et al. The value of PET/CT in detecting bone marrow involvement in patients with follicular lymphoma. *Medicine (Baltimore)*. 2016;95:e2910.
45. El Karak F, Bou-Orm IR, Ghosn M, Kattan J, Farhat F, Ibrahim T, et al. PET/CT scanner and bone marrow biopsy in detection of bone marrow involvement in diffuse large B-cell lymphoma. *PLoS One*. 2017;12:e0170299.
46. Li GW, Xu Z, Chen QW, Tian YN, Wang XY, Zhou L, et al. Quantitative evaluation of vertebral marrow adipose tissue in postmenopausal female using MRI chemical shift-based water-fat separation. *Clin Radiol*. 2014;69:254–62.
47. Pansini V, Monnet A, Salleron J, Hardouin P, Cortet B, Cotten A. 3 tesla (1) H MR spectroscopy of hip bone marrow in a healthy population, assessment of normal fat content values and influence of age and sex. *Journal of magnetic resonance imaging: JMIR*. 2014;39:369–76.
48. Kherlopian AR, Song T, Duan Q, Neimark MA, Po MJ, Gohagan JK, et al. A review of imaging techniques for systems biology. *BMC Syst Biol*. 2008;2:74.
49. Scheller EL, Troiano N, Vanhoutan JN, Boussein MA, Fretz JA, Xi Y, et al. Use of osmium tetroxide staining with microcomputerized tomography to visualize and quantify bone marrow adipose tissue in vivo. *Methods Enzymol*. 2014;537:123–39.
50. Ren J, Dimitrov I, Sherry AD, Malloy CR. Composition of adipose tissue and marrow fat in humans by 1H NMR at 7 Tesla. *J Lipid Res*. 2008;49:2055–62.
51. Yu EW, Greenblatt L, Eajazi A, Torriani M, Bredella MA. Marrow adipose tissue composition in adults with morbid obesity. *Bone*. 2017;97:38–42.
52. Schwartz AV, Sigurdsson S, Hue TF, Lang TF, Harris TB, Rosen CJ, et al. Vertebral bone marrow fat associated with lower trabecular BMD and prevalent vertebral fracture in older adults. *J Clin Endocrinol Metab*. 2013;98:2294–300.
53. Kuhn JP, Hernando D, Meffert PJ, Reeder S, Hosten N, Laqua R, et al. Proton-density fat fraction and simultaneous R2* estimation as an MRI tool for assessment of osteoporosis. *Eur Radiol*. 2013;23:3432–9.
54. Doucette CR, Horowitz MC, Berry R, MacDougald OA, Anunciado-Koza R, Koza RA, et al. High fat diet increases bone marrow adipose tissue (MAT) but does not alter trabecular or cortical bone mass in C57BL/6J mice. *J Cell Physiol*. 2015;230:2032–7.
55. Roldan-Valadez E, Pina-Jimenez C, Favila R, Rios C. Gender and age groups interactions in the quantification of bone marrow fat content in lumbar spine using 3T MR spectroscopy: a multivariate analysis of covariance (Mancova). *Eur J Radiol*. 2013;82:e697–702.
56. Griffith JF, Yeung DK, Ma HT, Leung JC, Kwok TC, Leung PC. Bone marrow fat content in the elderly: a reversal of sex difference seen in younger subjects. *Journal of Magnetic Resonance Imaging: JMIR*. 2012;36:225–30.
57. Wang QA, Tao C, Gupta RK, Scherer PE. Tracking adipogenesis during white adipose tissue development, expansion and regeneration. *Nat Med*. 2013;19:1338–44.
58. Lee YH, Petkova AP, Konkar AA, Granneman JG. Cellular origins of cold-induced brown adipocytes in adult mice. *FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology*. 2015;29:286–99.
59. Mizoguchi T, Pinho S, Ahmed J, Kunisaki Y, Hanoun M, Mendelson A, et al. Osterix marks distinct waves of primitive and definitive stromal progenitors during bone marrow development. *Dev Cell*. 2014;29:340–9.
60. Zhou BO, Yue R, Murphy MM, Peyer JG, Morrison SJ. Leptin-receptor-expressing mesenchymal stromal cells represent the main source of bone formed by adult bone marrow. *Cell Stem Cell*. 2014;15:154–68.
61. Yue R, Zhou BO, Shimada IS, Zhao Z, Morrison SJ. Leptin receptor promotes adipogenesis and reduces osteogenesis by regulating

- Mesenchymal stromal cells in adult bone marrow. *Cell Stem Cell*. 2016;18:782–96.
62. Ambrosi TH, Scialdone A, Graja A, Gohlke S, Jank AM, Bocian C, et al. Adipocyte accumulation in the bone marrow during obesity and aging impairs stem cell-based hematopoietic and bone regeneration. *Cell Stem Cell*. 2017;20:771–84. e6
 63. Dhurandhar NV. Infectobesity: obesity of infectious origin. *J Nutr*. 2001;131:2794S–7S.
 64. Atkinson RL, Dhurandhar NV, Allison DB, Bowen RL, Israel BA, Albu JB, et al. Human adenovirus-36 is associated with increased body weight and paradoxical reduction of serum lipids. *Int J Obes*. 2005;29:281–6.
 65. Dhurandhar NV, Whigham LD, Abbott DH, Schultz-Darken NJ, Israel BA, Bradley SM, et al. Human adenovirus Ad-36 promotes weight gain in male rhesus and marmoset monkeys. *J Nutr*. 2002;132:3155–60.
 66. Dhurandhar NV, Israel BA, Kolesar JM, Mayhew GF, Cook ME, Atkinson RL. Increased adiposity in animals due to a human virus. *International Journal of Obesity and Related Metabolic Disorders: Journal of the International Association for the Study of Obesity*. 2000;24:989–96.
 67. Atkinson RL. Viruses as an etiology of obesity. *Mayo Clin Proc*. 2007;82:1192–8.
 68. Vangipuram SD, Sheele J, Atkinson RL, Holland TC, Dhurandhar NV. A human adenovirus enhances preadipocyte differentiation. *Obes Res*. 2004;12:770–7.
 69. Rogers PM, Fusinski KA, Rathod MA, Loiler SA, Pasarica M, Shaw MK, et al. Human adenovirus Ad-36 induces adipogenesis via its E4 orf-1 gene. *Int J Obes*. 2008;32:397–406.
 70. Zvezdaryk KJ, Ferris MB, Strong AL, Morris CA, Bunnell BA, Dhurandhar NV, et al. Human cytomegalovirus infection of human adipose-derived stromal/stem cells restricts differentiation along the adipogenic lineage. *Adipocyte*. 2016;5:53–64.
 71. Olmsted-Davis E, Gannon FH, Ozen M, Ittmann MM, Gugala Z, Hipp JA, et al. Hypoxic adipocytes pattern early heterotopic bone formation. *Am J Pathol*. 2007;170:620–32.
 72. Salisbury EA, Dickerson AR, Davis TA, Forsberg JA, Davis AR, Olmsted-Davis EA. Characterization of brown adipose-like tissue in trauma-induced heterotopic ossification in humans. *Am J Pathol*. 2017;187:2071–9.
 73. Olmsted-Davis EA, Salisbury EA, Hoang D, Davis EL, Lazard Z, Sonnet C, et al. Progenitors in peripheral nerves launch heterotopic ossification. *Stem Cells Transl Med*. 2017;6:1109–19.
 74. Salisbury EA, Lazard ZW, Ubogu EE, Davis AR, Olmsted-Davis EA. Transient brown adipocyte-like cells derive from peripheral nerve progenitors in response to bone morphogenetic protein 2. *Stem Cells Transl Med*. 2012;1:874–85.
 75. Rosen CJ, Bouxsein ML. Mechanisms of disease: is osteoporosis the obesity of bone? *Nat Clin Pract Rheumatol*. 2006;2:35–43.