

Early-Life Adversity, Systemic Inflammation and Comorbid Physical and Psychiatric Illnesses of Adult Life



Maria Antonietta Nettis, Carmine M. Pariante, and Valeria Mondelli

Contents

1	Introduction	208
2	Association Between Childhood Trauma and Psychosis	209
3	Childhood Trauma and Immune Activation	210
3.1	Preclinical Evidence of the Link Between Early Stress and Increased Inflammation	211
3.2	The Emerging Evidence of Neuroinflammation	212
3.3	From Animal to Human Models	212
4	An Inflammatory Pathway Linking Childhood Trauma and Psychosis?	213
4.1	The Allostatic Systems: The Role of the Nervous System and of the Hypothalamic-Pituitary-Adrenal Axis	214
4.2	The Role of Neuroinflammation	215
4.3	Possible Role of the Interaction of Childhood Trauma with Other Environmental Factors	216
5	Inflammation and Physical Health	217
5.1	Inflammation and Metabolic Complications in Psychosis	217
5.2	Childhood Trauma and Metabolic Abnormalities	218
5.3	Childhood Trauma and Metabolic Abnormalities in Patients with Psychosis	219
6	Conclusion	220
	References	221

M. A. Nettis and C. M. Pariante

King's College London, Institute of Psychiatry, Psychology and Neuroscience, Department of Psychological Medicine, London, UK

National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London, London, UK

V. Mondelli (✉)

King's College London, Institute of Psychiatry, Psychology and Neuroscience, Department of Psychological Medicine, London, UK

National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London, London, UK

Maurice Wohl Clinical Neuroscience Institute, London, UK

e-mail: valeria.mondelli@kcl.ac.uk

Abstract Recently, the evidence of increased immune activation in patients with schizophrenia has suggested a role for the immune system in the development of psychosis. However, what is causing this increased immune activation and how this leads to the development of psychopathology remain still unclear. In this chapter we discuss the evidence about the role of childhood trauma as possible underlying cause of the increased immune activation in patients with schizophrenia. According to preclinical and clinical models, early adverse events can disrupt the homeostatic control of immune responses and lead to enduring inflammatory dysregulation at a peripheral and central level. In fact, persisting systemic inflammation may facilitate peripheral tissues damage and breach the blood-brain barrier, leading to microglia activation and to neuroinflammation.

Such chronic immune dysregulation also appear to partially explain the frequent comorbidity between psychosis and metabolic abnormalities, which have previously mainly considered as side effect of antipsychotic treatment.

Overall, this evidence suggests that early stress may contribute to development of schizophrenia spectrum disorders through a modulation of the peripheral and central immune system and support the immune pathways as possible future therapeutic approach for psychosis.

Keywords Childhood maltreatment · Childhood trauma · Comorbidities · Immune activation · Inflammation · Metabolic abnormalities · Psychosis · Schizophrenia

1 Introduction

In recent decades, research on schizophrenia has increasingly focused on the role of the immune system in the development of psychosis. In particular, the evidence of comorbidity between psychosis and metabolic abnormalities, together contributing to a reduced life expectancy, suggests that specific biological factors, such as an immune dysregulation, may link the aetiology of psychosis and physical health problems. A chronic low-grade immune activation has been consistently shown by studies in patients with schizophrenia spectrum disorders. These immune abnormalities have been detected not only in peripheral tissues (such as blood) but also in the brain, and they have been suggested to contribute to the development of other medical comorbidities, including metabolic and cardiovascular disorders (Leonard et al. 2012). However, what is causing this increased immune activation and how this leads to the development of psychopathology remain still unclear. Different causes have been considered, including perinatal infection, disruption of microbiota and psychosocial stress. In this chapter we are going to first discuss the evidence about the role of psychosocial stress, and more specifically childhood trauma, as possible underlying cause of the increased immune activation in patients with schizophrenia. In the second part of this chapter, we will analyse the evidence supporting the underlying role of childhood trauma and related immune dysregulation in the link between psychosis and metabolic disorders.

Childhood trauma is defined as harm, potential of harm or threat of a harm resulting from commission or omission by child's caregiver (Sideli et al. 2012). This definition includes a number of adverse experiences, such as physical, sexual and emotional abuse, neglect, parental death and bullying, but the most common forms of trauma, reported by both men and women, are physical abuse, physical neglect and emotional abuse, all of which are likely to co-occur (Scherrer et al. 2004). Neglect, the failure to provide for all aspects of the child's well-being, is considered the most frequent form with 78.5% of children exposed in the general population; physical abuse is reported in 17.6%, and it is defined as the use of physical force that harms the child's health, survival, development or dignity (Dubowitz et al. 2004). Approximately 8% of males and 20% of females universally experienced childhood sexual abuse (the involvement in sexual activity that a child is unable to give consent to or is not developmentally prepared for), with the highest prevalence in Africa (34.4%), followed by Asia (in particular China and India), America and Europe, with 23.9%, 10.1% and 9.2%, respectively (Verdolini et al. 2015). Finally, emotional abuse is the failure to provide children with a supportive environment. Actually, it seems to have higher prevalence than sexual and physical abuse, but it is more difficult to measure and quantify (Holmes and Slap 1998).

2 Association Between Childhood Trauma and Psychosis

Childhood exposure to adverse experiences has been associated with an increased risk of psychosis, mood disorders and other medical disorders in general, like cardiovascular disorders or diabetes (Varese et al. 2012; Coelho et al. 2014). Several studies have outlined the high prevalence of a history of childhood adversities among patients with psychosis. For example, in a comprehensive review in 2005, Read and colleagues reviewed 39 studies on female in- and outpatients with schizophrenia and demonstrated that the majority of female and male patients reported either childhood sexual abuse (CSA) or childhood physical abuse (CPA) (Read et al. 2005). A more recent study by Varese et al. (2012) found that childhood trauma increases the risk of psychosis with an OR of 2.8. Moreover, they showed that if the adversities considered as risk factors were entirely removed from the population, the number of people with psychosis would be reduced by 33%. It has also been showed that, with the exception of parental death, all types of adversities were related to an increased risk of psychosis. This suggests that psychosis risk is increased by the exposure to adverse experiences in general, rather than to a specific type of trauma.

When looking at studies investigating subjects at "ultrahigh risk" of developing psychosis, trauma has been repeatedly found to predict transition to psychosis in this population (Mayo et al. 2017). Sexual abuse has been shown to be the most common form of childhood trauma associated with later psychosis conversion, followed by

physical abuse (Bechdolf et al. 2005; Conus 2010). Finally, similarly to findings on sexual abuse history, increased severity and duration of individuals' bullying history has also been linked to the emergence of psychotic symptoms (Arseneault et al. 2011).

The most consistent finding about the association between childhood maltreatment and psychosis is that a history of childhood trauma increases specifically positive symptoms. Already in 1994, Ross and colleagues found that in-patients with schizophrenia with a history of sexual or physical abuse in their childhood had significantly more positive schizophrenia symptoms, especially hallucinations, and slightly fewer negative symptoms than those not abused (Ross et al. 1994). This evidence has been replicated in more recent studies (Ajnakina et al. 2016). Similarly, Bendall et al. (2013) reported that first-episode psychosis (FEP) patients exposed to childhood sexual abuse had more severe hallucinations and delusions (Bendall et al. 2013). According to their findings, childhood sexual abuse, especially rape, was associated with auditory verbal hallucinations, whereas victimization (physical abuse and bullying) predicted paranoia as well as auditory verbal hallucinations. Separation experiences (placements in foster care or institutions) were also associated with paranoia.

Trauma has also been suggested to influence the content of hallucinations, with patients experiencing hallucinations with themes similar to their trauma (Hardy et al. 2005). Furthermore, childhood trauma has a negative effect on cognitive functions in healthy individuals as well as in patients with psychosis and their high-risk offspring, in particular in relation to general cognitive abilities, memory and executive functions (Aas et al. 2011; Berthelot et al. 2015; Bucker et al. 2012). By contrast, several studies (Read and Ross 2003; Resnick et al. 2003; Lysaker et al. 2001) have found no differences in negative symptoms prevalence between abused and non-abused in-patients, whereas two adult in-patient studies found slightly fewer negative symptoms in abused subjects (McCormick and Goff 1991; Ross et al. 1994; Ajnakina et al. 2016).

3 Childhood Trauma and Immune Activation

Childhood adversities can be considered as environmental insults happening in a critical developmental phase of life. Interestingly, they have been suggested to alter the immune system function and to finally result in a chronic immune activation. Supporting this hypothesis, childhood adversities are strongly associated with acute and persistent inflammatory dysregulation, namely, increased blood levels of C reactive protein (CRP), interleukin-6 (IL-6) and tumour necrosis factor alpha (TNF- α), and such inflammation persists throughout adulthood (Coelho et al. 2014) offering a potential molecular pathway by which early trauma may increase susceptibility to psychiatric disorders (Baumeister et al. 2016). The most supported hypothesis is that early stress causes a pro-inflammatory status that may influence the response to inflammatory challenges during adult life, further contributing to

behavioural and cognitive alterations. As a result, childhood adversities may become an important preventable cause of psychiatric disorders, including psychosis.

As far as evidence of peripheral inflammation is concerned, we have recently conducted a meta-analysis of all studies investigating history of childhood maltreatment and inflammatory markers in adulthood (Baumeister et al. 2016). The meta-analysis included studies on both general and clinical populations (with psychiatric and physical disorders) and focused on CRP, IL-6 and TNF- α as these are the most examined inflammatory markers in psychiatric research. Results showed a significant association between childhood trauma and inflammatory markers in adulthood, with greatest effect size for TNF- α ($z = 0.20$, 95% CI = 0.10–0.29) followed by IL-6 ($z = 0.09$, 95% CI = 0.04–0.15) and then CRP ($z = 0.08$, 95% CI = 0.04–0.11). Another important finding from this study was that different types of trauma exposure impacted differentially on the inflammatory markers: physical and sexual abuse was associated with significant increased TNF- α and IL-6, but not CRP. Conversely, CRP was primarily related to parental absence during early development. How different types of trauma may impact different aspects of immune activity remains unknown; however, it has been suggested that factors such as context, time and duration of stress exposure may interact with individual trauma type in modulating immune response.

3.1 Preclinical Evidence of the Link Between Early Stress and Increased Inflammation

The first evidence of an association between early stress and immune system dysregulation was derived by preclinical studies. After Ader and colleagues found that rats handled before weaning showed slower development of a transplanted tumour (Ader and Friedman 1965), a number of authors started investigating the association between early-life stress and later immune functioning in rodents and non-human primates (Hennessy et al. 2010; Shanks and Lightman 2001). Stress exposure in early life (i.e., across postnatal days 1–20 in rodents) has been re-created in experimental models including neonatal handling, maternal separation, maternal deprivation, nursery rearing, early weaning and dexamethasone treatment. For example, maternal separation has been associated with an increased stress reactivity. Measures of immune functioning have included pro-inflammatory cytokines in the plasma, antigen-induced immunoproliferation in the spleen, expression levels of pro-inflammatory genes in the brain and intestinal microflora. Maternal separation in rats has been associated with elevated pro-inflammatory cytokines in the plasma (Reus et al. 2013; Wieck et al. 2013).

In studies with non-human primates, maternal separation led to an increase in macrophage activity (Coe et al. 1988) and long-term up-regulation in pro-inflammatory gene transcription in monocytes (Cole et al. 2012).

3.2 *The Emerging Evidence of Neuroinflammation*

The second interesting line of evidence coming from preclinical models is the link between early-life stress and markers of immune function in the central nervous system (CNS) (Reus et al. 2013) in adulthood. For example, in adult rats, early maternal separation is associated with greater synaptic levels of the receptor for the pro-inflammatory cytokine interleukin-1 (IL-1) (Viviani et al. 2014), greater number and motility of cortical microglial processes (Takatsuru et al. 2015) and greater microglia activation (Ganguly and Brenhouse 2015; Mondelli et al. 2017). Microglia are myeloid cells which provide the main form of adaptive immune response in the central nervous system (CNS) and play a crucial role in neuroinflammation. Indeed, neuroinflammation may lead to cognitive and behavioural dysfunction and psychopathology. A model of stress-induced central inflammation has already been created in animals by using stressful physical stimuli, such as restraint and foot shock, which could be applied to parent animals to induce maternal or perinatal stress (Mondelli et al. 2017).

The immediate consequences of early-life stress on animals' brain were investigated by Roque and colleagues in 2016 (Roque et al. 2016). The authors analysed the effect of maternal separation on astrocyte and microglial cell morphology in the hippocampus and hypothalamus of male rat pups and found that maternal separation (MS) caused microglia activation and decrease in astrocyte density in both areas. Moreover, compared with controls, MS rats showed increased IL-1 β in the hippocampus and increased TNF- α and IL-6 levels in the hypothalamus. Similarly, Diz-Chaves et al. (2012) showed that prenatal stress induces a basal pro-inflammatory status in the hippocampal formation during adulthood that results in an enhanced activation of microglia and astrocytes in response to pro-inflammatory insults. These findings suggest that early stress may contribute to decreased hippocampal neurogenesis and alter the neuroendocrine axis, leading to both psychopathological manifestation and metabolic abnormalities.

3.3 *From Animal to Human Models*

Preclinical evidence has led to a new insight about the link between stress and inflammation in humans. As far as peripheral inflammation is concerned, it is clear that the immune activation resulting from stress is able to protect the body from possible external challenges. Once released, inflammatory mediators are able to coordinate a variety of cell functions that stimulate and enhance immune activation. In particular, IL-1, IL-6 and TNF- α promote the differentiation of lymphocytes called *cytotoxic T cells*, which kill pathogens that are introduced into the body during physical wounding. These cytokines also promote increased vascular permeability and cellular adhesion, which allows immune cells to leave the blood vessels and migrate to tissues (Dhabhar et al. 2012). However, when such immune

activation persists chronically, it may damage vessels and peripheral tissues and lead to cardiovascular diseases.

Moreover, the hypothesis that stress may also reach the brain has progressively been investigated in humans. On one hand, stress-induced peripheral inflammation has been showed to affect the brain and to alter neural activity through different pathways: by active transport of cytokines, by involving macrophage-like cells residing in circumventricular organs or by the release of second messengers, which in turn stimulate local cytokines. These, in turn, can cross the blood-brain barrier and determine microglia activation (Cattaneo et al. 2015). They influence cell proliferation and survival depending on their inflammatory state (Mondelli et al. 2017). In response to harmful stimuli, microglial cells undergo a number of changes (Walker et al. 2014) including production of pro-inflammatory cytokines and the expression of several cell surface antigens that promote oxidative stress in the brain.

In humans, central inflammation (in terms of microglial activation) can be investigated with positron emission tomography (PET) using radio ligands for the 18-kDa translocator protein (TSPO), that is, a five-membrane domain protein localized mainly in the outer mitochondrial membrane of steroid-synthesizing cells, including those in the central nervous system. TSPO is involved in the transport of cholesterol into mitochondria. Peripheral lipopolysaccharide injection, used as immune challenge in primates, has been shown to increase TSPO expression in a uniform manner across the brain. This, in turn, seems to lead to microglial activation and to the expression of histological markers of brain activation in human post-mortem tissue (Mondelli et al. 2017).

4 An Inflammatory Pathway Linking Childhood Trauma and Psychosis?

As a result of the association of early adversities with both elevated inflammation and with psychosis risk, it is not surprising that patients with schizophrenia who have history of childhood trauma tend to show increased levels of some pro-inflammatory markers, including IL-6 and TNF- α (Dennison et al. 2012). In agreement with these findings, we have also previously shown that FEP patients with history of childhood sexual abuse have higher body mass index (BMI) and increased CRP levels in comparison with controls and patients without a history of sexual abuse (Hepgul et al. 2012). In addition, in another study we have also reported that FEP patients with childhood trauma have significantly higher serum levels of TNF- α and monocyte chemo-attractant protein-1 compared with patients without childhood trauma (di Nicola et al. 2013). The biological pathways linking stress to psychosis remain partly unclear, and although peripheral immune activation has been identified as an important link between the two, other biological systems and mechanisms appear to play a role. In particular, the hypothalamic-pituitary-adrenal (HPA) axis and the effects of neuroinflammation have been both explored in the link between stress and

psychosis and offer further understanding behind the presence of an increased peripheral inflammation in psychotic patients with experience of childhood trauma.

4.1 *The Allostatic Systems: The Role of the Nervous System and of the Hypothalamic-Pituitary-Adrenal Axis*

A suggested mechanism involved in childhood trauma embedding and then leading to chronic inflammation is the association between childhood adversities and enduring changes in the nervous, the endocrine and the immune systems that are highly integrated. Thus, the activation of one of these systems commonly triggers responses in the others (Danese and McEwen 2012).

4.1.1 The Nervous System

A neurobiological network including cerebral areas such as the thalamus, the sensory cortex and the amygdala is involved in detecting environmental threats. In response to psychosocial stressors, such as early adversities, the amygdala triggers firing in the locus coeruleus, which increases alertness and attention to the environment and induces a bodily response through the activation of the sympathetic nervous system (SNS) (the “fight or flight response”).

The SNS, in turn, affects the immune system and regulates pro-inflammatory cytokine production by releasing the neurotransmitter norepinephrine into peripheral tissues. In fact, by binding to β -adrenergic and α -adrenergic receptors, norepinephrine regulates the transcription of pro-inflammatory cytokine genes interleukin (IL)-1 and tumor necrosis factor (TNF)- α , leading to systemic immune activation (Danese and McEwen 2012). Then, a chronic or repeated stimulation of the sympathetic nervous system leads to chronically increased levels of inflammation. In the same time, key anti-inflammatory pathways, such as the *hypothalamic-pituitary-adrenal (HPA)* axis, are progressively downregulated under stressful conditions.

4.1.2 The Endocrine System: The Hypothalamic-Pituitary-Adrenal Axis

The principal endocrine effectors of the stress response are localized in the paraventricular nucleus (PVN) of the hypothalamus, the anterior lobe of the pituitary gland and the adrenal gland. Together, these structures are known as the hypothalamic-pituitary-adrenal axis.

Neurons localized in the medial parvocellular subdivision of the PVN secrete corticotropin-releasing factor (CRF), the principle regulator of the HPA axis. Stress (in this case, early adversities) triggers the release of CRF into hypophysial portal vessels that access the anterior pituitary gland. In turn, CRF induces the release of

adrenocorticotrophic hormone (ACTH) from corticotrophic cells into the systemic circulation. Then circulating ACTH targets the adrenal cortex, where it stimulates glucocorticoid synthesis and secretion from the zona fasciculata (Smith and Vale 2006).

Glucocorticoids, such as cortisol, exert their activity by binding to glucocorticoid receptors (GR). However, chronic stress can cause persistent elevated basal glucocorticoids levels and impair glucocorticoids ability to exert the negative feedback on the HPA axis itself. Moreover, glucocorticoids exert a natural anti-inflammatory action, binding GR in the body, but chronic stress leads to an impaired GR sensitivity (glucocorticoid resistance) and then to a prolonged/increased immune activation (Miller et al. 1999). Both HPA axis hyperactivity and increased immune activation are present in maltreated children (Danese and McEwen 2012) and persist in adulthood with several detrimental effects. As a consequence, subsequent responses to stressful situations may be abnormal. Second, the elevation of glucocorticoids levels due to glucocorticoid resistance may endanger the hippocampus and make it vulnerable to potential injury as well as stimulate striatal dopamine release that make patients more prone to psychosis (Read et al. 2001). Finally, this chronic alteration of the endocrine and immune system may progressively promote metabolic abnormalities and explain their common comorbidity with psychotic disorders. In fact, high cortisol levels can increase glucose levels and fat deposition (Dinan et al. 2004); clinical evidence has also shown that stress can lead to an increased intake of high-calories food (Dallman et al. 2005) and precipitate binge eating (Freeman and Gil 2004).

4.2 *The Role of Neuroinflammation*

Another hypothesized mechanism linking early trauma to psychosis is neuroinflammation. Preclinical studies suggest that microglia could be an important mediator of the association between psychosocial stress, in particular early stress, and psychiatric disorders, including psychosis. In fact, microglia activation may in turn be associated with impairment of neurogenesis and lead to structural and functional changes in the brain that predispose individuals to mental illnesses. In humans, ten studies have been published that investigated microglial activity using PET brain imaging in patients with psychosis, but results are inconsistent, probably because of the use of different radiotracers and different ways of analysing the data. Indeed, four of these studies found an increase in TSPO binding, one found an increase in medicated patients but not in drug-naive patients, and the other five did not find significant differences between patients and matched controls (Mondelli et al. 2017).

Findings suggesting the presence of neuroinflammation have already led researchers to support the development of new treatments. In particular, drugs that inhibit or reduce microglial activation are being tested in a number of studies (Mondelli et al. 2017). A preclinical study from Giovanoli et al. (2016)

have explored whether an early anti-inflammatory intervention with minocycline during peripubertal stress exposure might prevent the subsequent emergence of adult behavioural pathology (Giovanoli et al. 2016). They used an environmental two-hit model in mice, in which prenatal maternal administration of the viral mimetic poly(I:C) served as the first hit and exposure to subchronic unpredictable stress during peripubertal maturation as the second hit. Using this model, they examined the effectiveness of the tetracycline antibiotic minocycline, a broad-spectrum tetracyclic antibiotic displaying neuroprotective properties, given during stress exposure, to block stress-induced inflammatory responses and to prevent subsequent behavioural abnormalities. They found that combined exposure to prenatal immune activation and peripubertal stress caused significant behavioural dysfunctions, which minocycline treatment during stress exposure was able to prevent. In addition, the same pharmacological intervention blocked hippocampal and prefrontal microglia activation and interleukin-1 β expression in offspring exposed to prenatal infection and peripubertal stress. Minocycline has been suggested as a new potential therapy for negative symptoms in clinical studies of patients with schizophrenia. In two clinical trials comparing minocycline versus placebo, both added to the standard care, patients receiving minocycline showed a greater reduction in negative symptoms (Levkovitz et al. 2010; Chaudhry et al. 2012).

4.3 Possible Role of the Interaction of Childhood Trauma with Other Environmental Factors

A number of environmental insults have been associated with a risk of psychosis, including socioeconomic status, cannabis use and urbanicity. Epidemiological studies have focused their interest on the interaction between childhood adversities and other environmental risk factors in order to provide broader insights into inflammatory trajectories leading to psychosis onset. In particular, there are studies showing that childhood traumatic events are associated with increased cannabis use in adulthood (Harley et al. 2010; Houston et al. 2011; Konings et al. 2012). Some preclinical data demonstrate that Δ^9 -THC in adolescent mice triggers immune dysfunctions that last long after the end of abuse, switching the murine immune system to pro-inflammatory status in adulthood.

Such a pro-inflammatory state they may play a key role in the pathogenesis of neuropsychiatric disorders by modulating neurotransmitter and neuropeptide systems (Kronfol and Remick 2000; Muller and Ackenheil 1998) such as central monoamine activity (DeLisi 1992; Zalcman et al. 1994). This may partly explain why the onset of cannabis consumption at an earlier age has been identified by several authors as a factor contributing to poor prognosis in schizophrenia (Veen et al. 2004; Busse et al. 2012). A modulation of microglia function by cannabinoids,

both endogenous and synthetic ones, has also been suggested. This dysregulation might cause an alteration in the neuronal architecture or the neurotransmitter flow (Busse et al. 2012; Leweke and Koethe 2008; Bernstein et al. 2009) meaning that cannabinoid action may not be limited to direct changes at the neurotransmitter level; it may alter the “immune atmosphere” of the brain (Skaper et al. 2013).

5 Inflammation and Physical Health

Another interesting aspect of high levels of inflammation is that they are generally associated with physical health comorbidities, in particular cardiovascular diseases, in both the general population and in patients with psychosis (Russell et al. 2015).

This is because the functions of the metabolic and immune system are highly interdependent. An example is the overlapping function of macrophages and adipocytes. In obesity, these two kinds of cells contribute together to the production of inflammatory mediators, which, together with fatty acids, are able to inhibit the downstream signalling of insulin receptors, eventually leading to insulin resistance. Consistently, metabolic-related conditions as obesity, type 2 diabetes and insulin resistance are usually associated with chronic inflammation (Wellen and Hotamisligil 2005), in particular with high levels of IL-6 and CRP (Capuzzi and Freeman 2007). Inflammation is also involved in all stages of atherothrombosis, the underlying cause of approximately 80% of all sudden cardiac death (SCD). In fact, inflammatory cells such as macrophages and T lymphocytes eventually contribute to the formation of the atheromatous lesion, which consists of a lipid pool protected by a fibrous cap. Moreover, the activation of these cells leads to the release of additional mediators, including adhesion molecules, cytokines, chemokines and growth factors, all of which play important roles in atherogenesis and in the development of cardiovascular events (Willerson and Ridker 2004). In patients with psychosis, cardiovascular issues are even more present than in the general population. For example, as early as 1960 a survey in the USA showed that 40% of hospitalized patients with schizophrenia were overweight, in contrast to 20% of the general population (Gordon et al. 1960), and in 2004, a cross-sectional study by Kato and colleagues found that the prevalence of metabolic syndrome in patients with schizophrenia was 63% and that they had threefold greater risk to develop this syndrome than the general population (Kato et al. 2004).

5.1 *Inflammation and Metabolic Complications in Psychosis*

Metabolic abnormalities and weight gain in patients with psychosis are widely associated to antipsychotic treatment (Allison and Casey 2001; Mondelli et al. 2013). However, studies have shown impaired glucose tolerance, increased visceral fat and

increased obesity and hypertension also in drug-naïve patients (Correll et al. 2014), suggesting an impaired metabolism regardless of medication. For instance, some authors found that prevalence of diabetes in schizophrenia exceeds that in the general population well before the widespread use of the new (atypical) antipsychotic drugs (Dixon et al. 2000). Therefore, impaired glucose metabolism may be associated with schizophrenia rather than be only a side effect of antipsychotic treatment. To this regard, Perry and colleagues have conducted an interesting meta-analysis on pre-diabetic markers in antipsychotic naïve FEP. They included 12 studies (a total of 1,137 participants), and they found first-episode psychosis to be related to insulin resistance and impaired glucose tolerance (but not fasting plasma glucose) in the absence of medication (Perry et al. 2016).

On the other hand, in diabetic patients, untreated hyperglycaemia has been associated with alterations of mood state and acute psychosis manifestations (Sahoo et al. 2016).

Again, stress and inflammation are possible common pathways explaining the link between psychosis and metabolic abnormalities. In particular, childhood trauma may have a key role in mediating both.

Interestingly, inflammation has already been suggested to link cardiovascular abnormalities and depression. CRP, IL-1 and IL-6 have been associated with atherosclerosis and depression alike, both in healthy subjects and in cardiac patients. Previous studies have suggested both that inflammation increases risk of depression and that depression causes inflammation in patients with cardiovascular disorders (Elderon and Whooley 2013). Thus, although it remains difficult to disentangle whether such inflammatory mediators serve as triggers of both depression and CVD, act on the causal pathway between them or result from both conditions, at least it is now clear that inflammatory pathways may be the shared biological mechanisms between mental and physical diseases.

5.2 *Childhood Trauma and Metabolic Abnormalities*

Again, animal models, like those involving non-human primates, have been a useful source of evidence suggesting that adverse childhood experiences may influence physical health, especially obesity risk.

Kaufman et al. (2007) showed that compared with normally reared monkeys, those exposed to early stress exhibit greater weight, BMI, abdominal circumference, glucagon-like peptide-1 and decreased glucose disposal rates during hyperinsulinaemic-euglycaemic clamps. In their model of early-life stress (variable foraging demand [VFD]), food insecurity is imposed on monkey mothers for 16 weeks beginning when their nursing offspring are 3–5 months of age. VFD resulted in a range of neurobiological abnormalities, including dysregulation of the HPA axis, manifested in abnormal cerebrospinal fluid cortisol and corticotropin-releasing factor levels. These data suggest that early-life stress during a critical

period of neurodevelopment can result in the prepubertal emergence of obesity and insulin resistance.

Evidence of the link between early trauma and metabolic abnormalities is also quite clear from more recent human studies. A meta-analysis conducted by Danese and Tan (2014) on 41 studies in humans suggests that childhood maltreatment predicts obesity, independently from the measures and the definitions used and other potential confounding variables. Stressful psychosocial experiences in childhood might thus be conceptualized as potentially modifiable risk factors for obesity. Thus, prevention or effective treatment of severe cases of childhood maltreatment could avoid development of obesity in adulthood. However, it is still unclear if and how the effect of maltreatment on obesity could be modified through intervention.

Li et al. (2017) showed that childhood maltreatment is an independent risk factor for developing prediabetes. They recruited 121 participants from the general population, either with ($n = 69$) or without history of childhood maltreatment ($n = 52$). The authors found a 15% higher glucose area under the OGTT curve in the maltreated group, together with impaired insulin sensitivity. This group also showed higher CRP and TNF- α levels, both positively correlating with severity of childhood trauma ($r = 0.21, 0.23$, respectively, both $p < 0.05$). These data suggest an important relationship between childhood maltreatment and increased risk for prediabetic state due to glucose intolerance and impaired insulin sensitivity and beta cells function. Finally, Rich-Edwards et al. (2012) also showed that severe child abuse is a risk factor for early adult cardiovascular disorders, while Suglia et al. (2014) found that women who experienced sexual abuse in early childhood had a higher prevalence of hypertension (prevalence ratio (PR) 1.43 95% CI 1.00, 2.05) compared with women who did not experience such maltreatment.

5.3 Childhood Trauma and Metabolic Abnormalities in Patients with Psychosis

When looking at schizophrenia spectrum patients, only three studies have addressed the contribution of childhood trauma to cardio-metabolic risk.

In our study by Hepgul et al. (2012), we found that FEP patients with a history of childhood sexual abuse had higher BMI and CRP levels when compared with healthy controls and patients without childhood sexual abuse. CRP has been recently considered a marker of increased risk of diabetes and other metabolic dysfunction (Bassuk et al. 2004) and has also been similarly linked to chronic psychosocial stress (Miller 2008). Another study (Misiak et al. 2015) revealed that a history of childhood adversities, especially sexual and emotional abuse, is associated with higher systolic and diastolic blood pressure as well as greater levels of low-density lipoproteins (LDL) in FEP patients. Moreover, a study by Rajkumar (2015) found an association between childhood trauma and higher BMI and systolic blood pressure in patients with schizophrenia. In particular, physical abuse was linked to elevated

systolic blood pressure, while emotional abuse and neglect in women were linked to being overweight.

Finally, whether psychotropic drugs may play a role in leading to inflammation, by interacting with childhood adversities, is still unclear. The association between antipsychotic medication and an impaired metabolism is well known (Martin Otano et al. 2013), so that treated patients often show an overall altered metabolic-inflammatory status. Given the already mentioned association between sexual abuse and severity of hallucinations and delusions, it could be hypothesized that higher doses of antipsychotics are prescribed to patients with a history of childhood maltreatment. As a consequence, side effects of medication would be another mechanism linking childhood maltreatment and adulthood inflammation. However, studies investigating this hypothesis are currently lacking, so this should be the aim of future works.

6 Conclusion

In conclusion, susceptibility to psychosis is likely to be influenced by personal history of exposure to early traumatic events. In this model, priming adverse events can disrupt the homeostatic control of immune responses leading to enduring inflammatory dysregulation at a peripheral and central level. In particular, persisting systemic low-grade chronic inflammation may facilitate peripheral tissues damage and breach the blood-brain barrier with activation of microglia and development of neuroinflammation. The immune dysregulation also explains the development of multi-axial comorbidity including metabolic abnormalities and cardiovascular disorders, which have previously mainly considered as side effect of antipsychotic treatment (Fig. 1).




Fig. 1 Proposed model through which childhood adversities contribute to development of psychosis and comorbid metabolic abnormalities via activation of peripheral and central immune system

Two important considerations should be drawn from this work: First, research in the field of immunopsychiatry may lead to the development of new treatments targeting immune pathways in future therapeutic approaches to psychosis. Second, it is clear that early stress might be an important factor to early detect, prevent and better manage the development and treatment of schizophrenia spectrum disorders and any related physical comorbidities.

Acknowledgements This work has been supported by the South London and Maudsley (SLaM) NHS Foundation Trust & Institute of Psychiatry NIHR Biomedical Research Centre (BRC) for Mental Health; a strategic award from the Wellcome Trust (Consortium of Neuroimmunology of Mood Disorders and Alzheimer's Disease); and the grants 'Persistent Fatigue Induced by Interferon-alpha: A New Immunological Model for Chronic Fatigue Syndrome' (MR/J002739/1), and 'Immuno-psychiatry: a consortium to test the opportunity for immunotherapeutics in psychiatry' (MR/L014815/1), from the Medical Research Council (UK).

References

- Aas M, Dazzan P, Fisher HL, Morgan C, Morgan K, Reichenberg A, Zanelli J, Fearon P, Jones PB, Murray RM, Pariante CM (2011) Childhood trauma and cognitive function in first-episode affective and non-affective psychosis. *Schizophr Res* 129(1):12–19
- Ader R, Friedman SB (1965) Differential early experiences and susceptibility to transplanted tumor in the rat. *J Comp Physiol Psychol* 59:361–364
- Ajnakina O, Trotta A, Oakley-Hannibal E, di Forti M, Stilo SA, Kolliakou A, Gardner-Sood P, Gaughran F, David AS, Dazzan P, Pariante C, Mondelli V, Morgan C, Vassos E, Murray RM, Fisher HL (2016) Impact of childhood adversities on specific symptom dimensions in first-episode psychosis. *Psychol Med* 46(2):317–326
- Allison DB, Casey DE (2001) Antipsychotic-induced weight gain: a review of the literature. *J Clin Psychiatry* 62(Suppl 7):22–31
- Arseneault L, Cannon M, Fisher HL, Polanczyk G, Moffitt TE, Caspi A (2011) Childhood trauma and children's emerging psychotic symptoms: a genetically sensitive longitudinal cohort study. *Am J Psychiatry* 168(1):65–72
- Bassuk SS, Rifai N, Ridker PM (2004) High-sensitivity C-reactive protein: clinical importance. *Curr Probl Cardiol* 29(8):439–493
- Baumeister D, Akhtar R, Ciufolini S, Pariante CM, Mondelli V (2016) Childhood trauma and adulthood inflammation: a meta-analysis of peripheral C-reactive protein, interleukin-6 and tumour necrosis factor-alpha. *Mol Psychiatry* 21(5):642–649
- Bechdolf A, Pohlmann B, Geyer C, Ferber C, Klosterkötter J, Gouzoulis-Mayfrank E (2005) Motivational interviewing for patients with comorbid schizophrenia and substance abuse disorders: a review. *Fortschr Neurol Psychiatr* 73(12):728–735
- Bendall S, Hulbert CA, Alvarez-Jimenez M, Allott K, McGorry PD, Jackson HJ (2013) Testing a model of the relationship between childhood sexual abuse and psychosis in a first-episode psychosis group: the role of hallucinations and delusions, posttraumatic intrusions, and selective attention. *J Nerv Ment Dis* 201(11):941–947
- Bernstein HG, Steiner J, Bogerts B (2009) Glial cells in schizophrenia: pathophysiological significance and possible consequences for therapy. *Expert Rev Neurother* 9(7):1059–1071
- Berthelot N, Paccalet T, Gilbert E, Moreau I, Merette C, Gingras N, Rouleau N, Maziade M (2015) Childhood abuse and neglect may induce deficits in cognitive precursors of psychosis in high-risk children. *J Psychiatry Neurosci* 40(5):336–343

- Bucker J, Kapczinski F, Post R, Cereser KM, Szobot C, Yatham LN, Kapczinski NS, Kauer-Sant'Anna M (2012) Cognitive impairment in school-aged children with early trauma. *Compr Psychiatry* 53(6):758–764
- Busse S, Busse M, Schiltz K, Bielau H, Gos T, Brisch R, Mawrin C, Schmitt A, Jordan W, Muller UJ, Bernstein HG, Bogerts B, Steiner J (2012) Different distribution patterns of lymphocytes and microglia in the hippocampus of patients with residual versus paranoid schizophrenia: further evidence for disease course-related immune alterations? *Brain Behav Immun* 26(8):1273–1279
- Capuzzi D, Freeman J (2007) C-reactive protein and cardiovascular risk in the metabolic syndrome and type 2 diabetes: controversy and challenge. *Clin Diabetes* 25(1):16–22
- Cattaneo A, Macchi F, Plazzotta G, Veronica B, Bocchio-Chiavetto L, Riva MA, Pariante CM (2015) Inflammation and neuronal plasticity: a link between childhood trauma and depression pathogenesis. *Front Cell Neurosci* 9:40
- Chaudhry IB, Hallak J, Husain N, Minhas F, Stirling J, Richardson P, Dursun S, Dunn G, Deakin B (2012) Minocycline benefits negative symptoms in early schizophrenia: a randomised double-blind placebo-controlled clinical trial in patients on standard treatment. *J Psychopharmacol* 26(9):1185–1193
- Coe CL, Rosenberg LT, Levine S (1988) Effect of maternal separation on the complement system and antibody responses in infant primates. *Int J Neurosci* 40(3–4):289–302
- Coelho R, Viola TW, Walss-Bass C, Brietzke E, Grassi-Oliveira R (2014) Childhood maltreatment and inflammatory markers: a systematic review. *Acta Psychiatr Scand* 129(3):180–192
- Cole SW, Conti G, Arevalo JM, Ruggiero AM, Heckman JJ, Suomi SJ (2012) Transcriptional modulation of the developing immune system by early life social adversity. *Proc Natl Acad Sci U S A* 109(50):20578–20583
- Conus P (2010) First episode of mood disorders: an opportunity for early intervention in bipolar disorders. *Encéphale* 36(Suppl 3):S71–S76
- Correll CU, Robinson DG, Schooler NR, Brunette MF, Mueser KT, Rosenheck RA, Marcy P, Addington J, Estroff SE, Robinson J, Penn DL, Azrin S, Goldstein A, Severe J, Heinsen R, Kane JM (2014) Cardiometabolic risk in patients with first-episode schizophrenia spectrum disorders: baseline results from the RAISE-ETP study. *JAMA Psychiat* 71(12):1350–1363
- Dallman MF, Pecoraro NC, la Fleur SE (2005) Chronic stress and comfort foods: self-medication and abdominal obesity. *Brain Behav Immun* 19(4):275–280
- Danese A, McEwen BS (2012) Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiol Behav* 106(1):29–39
- Danese A, Tan M (2014) Childhood maltreatment and obesity: systematic review and meta-analysis. *Mol Psychiatry* 19(5):544–554
- DeLisi LE (1992) The significance of age of onset for schizophrenia. *Schizophr Bull* 18(2):209–215
- Dennison U, McKernan D, Cryan J, Dinan T (2012) Schizophrenia patients with a history of childhood trauma have a pro-inflammatory phenotype. *Psychol Med* 42(9):1865–1871
- Dhabhar FS, Malarkey WB, Neri E, McEwen BS (2012) Stress-induced redistribution of immune cells – from barracks to boulevards to battlefields: a tale of three hormones – Curt Richter Award winner. *Psychoneuroendocrinology* 37(9):1345–1368
- di Nicola M, Cattaneo A, Hepgul N, Di Forti M, Aitchison KJ, Janiri L, Murray RM, Dazzan P, Pariante CM, Mondelli V (2013) Serum and gene expression profile of cytokines in first-episode psychosis. *Brain Behav Immun* 31:90–95
- Dinan T, Peveler R, Holt R (2004) Understanding schizophrenia and diabetes. *Hosp Med* 65(8):485–488
- Dixon L, Weiden P, Delahanty J, Goldberg R, Postrado L, Lucksted A, Lehman A (2000) Prevalence and correlates of diabetes in national schizophrenia samples. *Schizophr Bull* 26(4):903–912
- Diz-Chaves Y, Pernia O, Carrero P, Garcia-Segura LM (2012) Prenatal stress causes alterations in the morphology of microglia and the inflammatory response of the hippocampus of adult female mice. *J Neuroinflammation* 9:71

- Dubowitz H, Pitts SC, Black MM (2004) Measurement of three major subtypes of child neglect. *Child Maltreat* 9(4):344–356
- Elderon L, Whooley MA (2013) Depression and cardiovascular disease. *Prog Cardiovasc Dis* 55(6):511–523
- Freeman LM, Gil KM (2004) Daily stress, coping, and dietary restraint in binge eating. *Int J Eat Disord* 36(2):204–212
- Ganguly P, Brenhouse HC (2015) Broken or maladaptive? Altered trajectories in neuroinflammation and behavior after early life adversity. *Dev Cogn Neurosci* 11:18–30
- Giovanoli S, Engler H, Engler A, Richetto J, Feldon J, Riva MA, Schedlowski M, Meyer U (2016) Preventive effects of minocycline in a neurodevelopmental two-hit model with relevance to schizophrenia. *Transl Psychiatry* 5(6):38
- Gordon HL, Law A, Hohman KE, Groth C (1960) The problem of overweight in hospitalized psychotic patients. *Psychiatry Q* 34:69–82
- Hardy A, Fowler D, Freeman D, Smith B, Steel C, Evans J, Garety P, Kuipers E, Bebbington P, Dunn G (2005) Trauma and hallucinatory experience in psychosis. *J Nerv Ment Dis* 193(8):501–507
- Harley M, Kelleher I, Clarke M, Lynch F, Arseneault L, Connor D, Fitzpatrick C, Cannon M (2010) Cannabis use and childhood trauma interact additively to increase the risk of psychotic symptoms in adolescence. *Psychol Med* 40(10):1627–1634
- Hennessy MB, Deak T, Schiml-Webb PA (2010) Early attachment-figure separation and increased risk for later depression: potential mediation by proinflammatory processes. *Neurosci Biobehav Rev* 34(6):782–790
- Hepgul N, Pariante CM, Dipasquale S, DiForti M, Taylor H, Marques TR, Morgan C, Dazzan P, Murray RM, Mondelli V (2012) Childhood maltreatment is associated with increased body mass index and increased C-reactive protein levels in first-episode psychosis patients. *Psychol Med* 42(9):1893–1901
- Holmes WC, Slap GB (1998) Sexual abuse of boys: definition, prevalence, correlates, sequelae, and management. *JAMA* 280(21):1855–1862
- Houston JE, Murphy J, Shevlin M, Adamson G (2011) Cannabis use and psychosis: re-visiting the role of childhood trauma. *Psychol Med* 41(11):2339–2348
- Kato MM, Currier MB, Gomez CM, Hall L, Gonzalez-Blanco M (2004) Prevalence of metabolic syndrome in Hispanic and non-Hispanic patients with schizophrenia. *Prim Care Companion J Clin Psychiatry* 6(2):74–77
- Kaufman D, Banerji MA, Shorman I, Smith EL, Coplan JD, Rosenblum LA, Kral JG (2007) Early-life stress and the development of obesity and insulin resistance in juvenile bonnet macaques. *Diabetes* 56(5):1382–1386
- Konings M, Stefanis N, Kuepper R, de Graaf R, ten Have M, van Os J, Bakoula C, Henquet C (2012) Replication in two independent population-based samples that childhood maltreatment and cannabis use synergistically impact on psychosis risk. *Psychol Med* 42(1):149–159
- Kronfol Z, Remick DG (2000) Cytokines and the brain: implications for clinical psychiatry. *Am J Psychiatry* 157(5):683–694
- Leonard BE, Schwarz M, Myint AM (2012) The metabolic syndrome in schizophrenia: is inflammation a contributing cause? *J Psychopharmacol* 26(5 Suppl):33–41
- Levkovitz Y, Mendlovich S, Riwkes S, Braw Y, Levkovitch-Verbin H, Gal G, Fennig S, Treves I, Kron S (2010) A double-blind, randomized study of minocycline for the treatment of negative and cognitive symptoms in early-phase schizophrenia. *J Clin Psychiatry* 71(2):138–149
- Leweke FM, Koethe D (2008) Cannabis and psychiatric disorders: it is not only addiction. *Addict Biol* 13(2):264–275
- Li L, Garvey WT, Gower BA (2017) Childhood maltreatment is an independent risk factor for prediabetic disturbances in glucose regulation. *Front Endocrinol (Lausanne)* 8:151
- Lysaker PH, Meyer PS, Evans JD, Clements CA, Marks KA (2001) Childhood sexual trauma and psychosocial functioning in adults with schizophrenia. *Psychiatr Serv* 52(11):1485–1488

- Martin Otano L, Barbadillo Izquierdo L, Galdeano Mondragon A, Alonso Pinedo M, Querejeta Ayerdi I (2013) After six months of anti-psychotic treatment: is the improvement in mental health at the expense of physical health? *Rev Psiquiatr Salud Ment* 6(1):26–32
- Mayo D, Corey S, Kelly LH, Yohannes S, Youngquist AL, Stuart BK, Niendam TA, Loewy RL (2017) The role of trauma and stressful life events among individuals at clinical high risk for psychosis: a review. *Front Psych* 8:55
- McCormick S, Goff DC (1991) Apparent decrease in schizophrenia. *Br J Psychiatry* 159:725
- Miller AH (2008) Inflammation versus glucocorticoids as purveyors of pathology during stress: have we reached the tipping point? *Biol Psychiatry* 64(4):263–265
- Miller AH, Pariante CM, Pearce BD (1999) Effects of cytokines on glucocorticoid receptor expression and function. Glucocorticoid resistance and relevance to depression. *Adv Exp Med Biol* 461:107–116
- Misiak B, Kiejna A, Frydecka D (2015) The history of childhood trauma is associated with lipid disturbances and blood pressure in adult first-episode schizophrenia patients. *Gen Hosp Psychiatry* 37(4):365–367
- Mondelli V, Anacker C, Vernon AC, Cattaneo A, Natesan S, Modo M, Dazzan P, Kapur S, Pariante CM (2013) Haloperidol and olanzapine mediate metabolic abnormalities through different molecular pathways. *Transl Psychiatry* 3:e208
- Mondelli V, Vernon AC, Turkheimer F, Dazzan P, Pariante CM (2017) Brain microglia in psychiatric disorders. *Lancet Psychiatry* 4(7):563–572
- Muller N, Ackenheil M (1998) Psychoneuroimmunology and the cytokine action in the CNS: implications for psychiatric disorders. *Prog Neuro-Psychopharmacol Biol Psychiatry* 22(1):1–33
- Perry BI, McIntosh G, Weich S, Singh S, Rees K (2016) The association between first-episode psychosis and abnormal glycaemic control: systematic review and meta-analysis. *Lancet Psychiatry* 3(11):1049–1058
- Rajkumar RP (2015) The impact of childhood adversity on the clinical features of schizophrenia. *Schizophr Res Treat* 2015:532082
- Read J, Ross CA (2003) Psychological trauma and psychosis: another reason why people diagnosed schizophrenic must be offered psychological therapies. *J Am Acad Psychoanal Dyn Psychiatry* 31(1):247–268
- Read J, Perry BD, Moskowitz A, Connolly J (2001) The contribution of early traumatic events to schizophrenia in some patients: a traumagenic neurodevelopmental model. *Psychiatry* 64(4):319–345
- Read J, van Os J, Morrison AP, Ross CA (2005) Childhood trauma, psychosis and schizophrenia: a literature review with theoretical and clinical implications. *Acta Psychiatr Scand* 112:330–350
- Resnick SG, Bond GR, Mueser KT (2003) Trauma and posttraumatic stress disorder in people with schizophrenia. *J Abnorm Psychol* 112(3):415–423
- Reus GZ, Dos Santos MA, Abelaira HM, Ribeiro KF, Petronilho F, Vuolo F, Colpo GD, Pfaffenseller B, Kapczinski F, Dal-Pizzol F, Quevedo J (2013) Imipramine reverses alterations in cytokines and BDNF levels induced by maternal deprivation in adult rats. *Behav Brain Res* 242:40–46
- Rich-Edwards JW, Mason S, Rexrode K, Spiegelman D, Hibert E, Kawachi I, Jun HJ, Wright RJ (2012) Physical and sexual abuse in childhood as predictors of early-onset cardiovascular events in women. *Circulation* 126(8):920–927
- Roque A, Ochoa-Zarzosa A, Torner L (2016) Maternal separation activates microglial cells and induces an inflammatory response in the hippocampus of male rat pups, independently of hypothalamic and peripheral cytokine levels. *Brain Behav Immun* 55:39–48
- Ross CA, Anderson G, Clark P (1994) Childhood abuse and the positive symptoms of schizophrenia. *Hosp Community Psychiatry* 45(5):489–491
- Russell A, Ciufolini S, Gardner-Sood P, Bonaccorso S, Gaughran F, Dazzan P, Pariante CM, Mondelli V (2015) Inflammation and metabolic changes in first episode psychosis: preliminary results from a longitudinal study. *Brain Behav Immun* 49:25–29

- Sahoo S, Mehra A, Grover S (2016) Acute hyperglycemia associated with psychotic symptoms in a patient with type 1 diabetes mellitus: a case report. *Innov Clin Neurosci* 13(11–12):25–27
- Scherrer JF, Waterman BM, Heath AC, Bucholz KK, True WR, Jacob T (2004) Are substance use, abuse and dependence associated with study participation? Predictors of offspring nonparticipation in a twin-family study. *J Stud Alcohol* 65(1):140–144
- Shanks N, Lightman SL (2001) The maternal-neonatal neuro-immune interface: are there long-term implications for inflammatory or stress-related disease? *J Clin Invest* 108(11):1567–1573
- Sideli L, Mule A, La Barbera D, Murray RM (2012) Do child abuse and maltreatment increase risk of schizophrenia? *Psychiatry Investig* 9(2):87–99
- Skaper SD, Facci L, Giusti P (2013) Glia and mast cells as targets for palmitoylethanolamide, an anti-inflammatory and neuroprotective lipid mediator. *Mol Neurobiol* 48(2):340–352
- Smith SM, Vale WW (2006) The role of the hypothalamic-pituitary-adrenal axis in neuroendocrine responses to stress. *Dialogues Clin Neurosci* 8(4):383–395
- Suglia SF, Clark CJ, Boynton-Jarrett R, Kressin NR, Koenen KC (2014) Child maltreatment and hypertension in young adulthood. *BMC Public Health* 14:1149
- Takatsuru Y, Nabekura J, Ishikawa T, Kohsaka S, Koibuchi N (2015) Early-life stress increases the motility of microglia in adulthood. *J Physiol Sci* 65(2):187–194
- Varesse F, Smeets F, Drukker M, Lieverse R, Lataster T, Viechtbauer W, Read J, van Os J, Bentall RP (2012) Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophr Bull* 38(4):661–671
- Veen ND, Selten JP, van der Tweel I, Feller WG, Hoek HW, Kahn RS (2004) Cannabis use and age at onset of schizophrenia. *Am J Psychiatry* 161(3):501–506
- Verdolini N, Attademo L, Agius M, Ferranti L, Moretti P, Quartesan R (2015) Traumatic events in childhood and their association with psychiatric illness in the adult. *Psychiatr Danub* 27(Suppl 1): S60–S70
- Viviani B, Boraso M, Valero M, Gardoni F, Marco EM, Llorente R, Corsini E, Galli CL, di Luca M, Marinovich M, Lopez-Gallardo M, Viveros MP (2014) Early maternal deprivation immunologically primes hippocampal synapses by redistributing interleukin-1 receptor type I in a sex dependent manner. *Brain Behav Immun* 35:135–143
- Walker FR, Beynon SB, Jones KA, Zhao Z, Kongsui R, Cairns M, Nilsson M (2014) Dynamic structural remodelling of microglia in health and disease: a review of the models, the signals and the mechanisms. *Brain Behav Immun* 37:1–14
- Wellen KE, Hotamisligil GS (2005) Inflammation, stress, and diabetes. *J Clin Invest* 115(5):1111–1119
- Wieck A, Andersen SL, Brenhouse HC (2013) Evidence for a neuroinflammatory mechanism in delayed effects of early life adversity in rats: relationship to cortical NMDA receptor expression. *Brain Behav Immun* 28:218–226
- Willerson JT, Ridker PM (2004) Inflammation as a cardiovascular risk factor. *Circulation* 109:II2–II10
- Zalcman S, Green-Johnson JM, Murray L, Nance DM, Dyck D, Anisman H, Greenberg AH (1994) Cytokine-specific central monoamine alterations induced by interleukin-1, -2 and -6. *Brain Res* 643(1–2):40–49