

Erratum to: LC-MS/MS Analysis of Cerebrospinal Fluid Metabolites in the Pterin Biosynthetic Pathway

Erland Arning · Teodoro Bottiglieri

Received: 21 May 2014 / Revised: 18 June 2014 / Accepted: 01 July 2014 / Published online: 08 June 2016
© SSIEM and Springer-Verlag Berlin Heidelberg 2014

Erratum to: JIMD Reports
DOI: 10.1007/8904_2014_336

The last row in Table 2 was incorrectly typeset in the original publication; the correct values are provided in the table below:

Table 2 Assay precision, linearity, and limit of detection (LOD)

Analyte (nmol/L)	Internal standard	Intra-assay (<i>n</i> = 10)		Inter-assay (<i>n</i> = 20)		Linearity	LOD
		Level 1 (CV%)	Level 2 (CV%)	Level 1 (CV%)	Level 2 (CV%)		
BH4	¹⁵ N-BH ₄	55.1 ± 1.7 (3.0)	156.1 ± 7.2 (4.6)	50.2 ± 2.9 (5.9)	149.0 ± 6.2 (4.2)	3–200	1
BH2	¹⁵ N-BH ₂	41.0 ± 1.8 (4.4)	170.9 ± 5.9 (3.5)	38.6 ± 3.2 (8.2)	152.7 ± 12.8 (8.4)	3–200	1
Neopterin	¹⁵ N-Neopterin	35.6 ± 2.0 (5.6)	160.2 ± 7.6 (4.7)	29.1 ± 2.8 (9.5)	141.3 ± 10.3 (7.3)	3–200	1
Sepiapterin	¹⁵ N-BH ₂	17.5 ± 0.9 (5.3)	101.3 ± 4.4 (4.3)	23.1 ± 3.3 (14.4)	119.7 ± 15.0 (12.5)	3–200	1

Data expressed as nmol/L, mean ± standard deviation (coefficient of variation, %)

The online versions of the original chapter can be found under
DOI 10.1007/8904_2014_336

E. Arning (✉) · T. Bottiglieri
Baylor Research Institute, Institute of Metabolic Disease, Dallas,
TX 75226, USA
e-mail: erlanda@baylorhealth.edu