

# **Erratum to: Treatment with Mefolinate (5-Methyltetrahydrofolate), but Not Folic Acid or Folinic Acid, Leads to Measurable 5-Methyltetrahydrofolate in Cerebrospinal Fluid in Methylenetetrahydrofolate Reductase Deficiency**

**L. Knowles · A.A.M. Morris · J.H. Walter**

Received: 18 October 2015 / Revised: 07 December 2015 / Accepted: 10 December 2015 / Published online: 21 June 2016  
© SSIEM and Springer-Verlag Berlin Heidelberg 2016

Erratum to: JIMD Reports  
DOI: 10.1007/8904\_2016\_529

The final sentence of the abstract should read ‘Only treatment with oral 5-MTHF given as calcium mefolinate at doses of 15–60 mg/day resulted in an increase in CSF 5-MTHF’.

---

The online version of the original chapter can be found under  
DOI 10.1007/8904\_2016\_529

---

---

L. Knowles · A.A.M. Morris · J.H. Walter  
Willink Biochemical Genetics Unit, Genetic Medicine, Manchester  
Academic Health Sciences Centre, Central Manchester University  
Hospitals NHS Foundation Trust, Manchester, UK

J.H. Walter (✉)  
Willink Biochemical Genetics Unit, Manchester Centre for Genomic  
Medicine, University of Manchester, Central Manchester University  
Hospitals NHS Foundation Trust, St Mary’s Hospital, Oxford Road,  
Manchester M13 9WL, UK  
e-mail: john.walter@cmft.nhs.uk