

Henryk A. Domanski, Xiaohua Qian,  
Måns Åkerman, and Donald E. Stanley

## 13.1 Introduction

### 13.1.1 Evaluation of Soft Tissue Lesions by FNA

Over the years, with advances in our understanding of soft tissue tumors, availability of an increasing number of diagnostic ancillary tests, and matured biopsy techniques with or without image guidance, needle biopsy investigation of a soft tissue lesion has largely replaced open excisional biopsy as the primary diagnostic tool to establish malignancy and assess histologic type and grade [1]. Fine needle aspiration cytology (FNAC) has been used as the first-line approach for several decades in many medical centers in Europe and as a complementary tool to the needle core biopsy in most centers in northern America [2–8].

According to the guidelines of the Scandinavian Sarcoma Group, patients present with a suspicion of soft tissue sarcoma (any deep intra- or intermuscular tumors and superficial tumors >5 cm) should be referred to the centers with

histopathologic and cytologic expertise and the capability for a clinical–radiologic correlation.

Multidisciplinary management of soft tissue neoplasms during the past few decades favorably influenced outcomes for patients suffering from sarcoma. The best approach to a successful diagnostic evaluation of soft tissue lesions is to combine clinical data and radiographic findings with the morphology interpretation that should be accomplished with the aid of supportive ancillary techniques [9].

Accumulated experience from Sweden and other groups with expertise in FNAC of soft tissue and bone lesions has indicated that FNAC is suitable to establish a metastatic carcinoma (*see* Fig. 13.1), melanoma, or lymphoma, to document a recurrence or metastasis of a previously treated sarcoma, and to diagnose a benign soft tissue lesion, as well as a primary soft tissue sarcoma [6, 10–13].

### 13.1.2 Sampling Technique


Technically, FNA biopsy of a soft tissue lesion is not significantly different from sampling a lesion from other parts of the body. Certain parameters that are particularly important in FNA of a soft tissue lesion include anatomic site, size of the target, and tumor consistency and its mobility. A syringe holder, 10 ml syringe, and needles of varying lengths are recommended. Needles wider than 22 gauge are seldom necessary. For deep-seated lesions, a needle with a stylet may be used to avoid sampling of the adjacent normal tissue. If a sarcoma is suspected, aspirating through the vertex and through a single tract is preferred. The direction of the needle should be changed with each pass to cover different parts of the tumor. The preferred areas of needle entrance are often marked by the orthopaedic surgeon as part of surgery planning so as to ensure that the needle tracts are completely removed during the subsequent surgery (*see* Fig. 13.2) [14].

---


H.A. Domanski, MD, PhD (✉) • M. Åkerman, MD, PhD  
Department of Pathology, Skåne University Hospital,  
Sölvegatan 25, 22185 Lund, Sweden  
e-mail: henryk.domanski@med.lu.se; birgitta.berglof@telia.com

X. Qian, MD, PhD  
Department of Pathology, Brigham & Women's Hospital,  
Harvard Medical School, 75 Francis St., Boston, MA 02115, USA  
e-mail: xqian@partners.org

D.E. Stanley, MD  
Department of Pathology, Maine Medical Center,  
22 Bramhall St., Portland, ME 04101, USA  
e-mail: stanld@mmc.org



**Fig. 13.1** Bone and soft tissue metastases. (a) Metastasis of colonic adenocarcinoma: clusters of columnar cells arranged in solid nests. Note mucinous background (H&E). (b) Tumor tissue architecture appears in cell-block sections (H&E; Cell Block)



**Fig. 13.2** Sampling technique. (a and b) Areas of the skin where the needle should be inserted were marked by the orthopaedic surgeons

### 13.1.3 Ancillary Techniques

Ancillary techniques used in the work-up of FNA samplings have been described in Sect. 1.4. The most commonly used ancillary test is immunocytochemistry (IC), which is preferably performed on cell-block preparations or liquid-base preparations (Thin Prep [TP]). Useful antibodies in the FNA diagnosis of soft tissue tumors are listed in Table 13.1. Cytogenetic and molecular techniques play an increasingly important role in the evaluation of soft tissue neoplasms. FNA preparations are especially suitable for molecular genetic techniques such as reverse transcriptase–polymerase

chain reaction (RT-PCR) and fluorescence in situ hybridization (FISH) (Table 13.2) [15]. The use of electron microscopic (EM) examination has decreased considerably because of the highly effective IC and molecular genetic techniques to determine the line of differentiation of a soft tissue tumor.

### 13.1.4 Reporting the Diagnosis

The goal of a biopsy is to establish malignancy (sarcoma versus benign soft tissue lesion/neoplasm), to determine the

**Table 13.1** Useful antibodies in the diagnosis of soft tissue neoplasm

Antibodies	Neoplasm
Muscle-specific actin	Leiomyoma; leiomyosarcoma; myofibroblastic tumors; rhabdomyosarcoma; pseudosarcomas
Smooth muscle actin (SMA)	Leiomyoma; leiomyosarcoma; myofibroblastic tumors; pseudosarcomas; PEComa
Desmin	Leiomyoma; leiomyosarcoma; rhabdomyosarcoma; PEComa desmoplastic small round cell tumor; extrarenal malignant rhabdoid tumor (some); angiomatoid fibrous histiocytoma (some)
Caldesmon	Leiomyoma; leiomyosarcoma
Myoglobin	Rhabdomyosarcoma
MyoD1	Rhabdomyosarcoma; alveolar soft past sarcoma (cytoplasmic)
S-100 protein	Benign nerve sheath tumors; malignant peripheral nerve sheath tumor; granular cell tumor; clear cell sarcoma; round cell liposarcoma; ossifying fibromyxoid tumor; synovial sarcoma ~30 %
CD34	Dermatofibrosarcoma protuberans; hemangiopericytoma/solitary fibrous tumor; epithelioid sarcoma; hemangiomas; angiosarcoma; gastrointestinal stromal tumor; spindle-cell lipoma/pleomorphic lipoma
CD31	Hemangiomas; angiosarcoma; hemangioendothelioma
Fli-1/ERG	Ewing sarcoma; angiosarcoma
CD99	Ewing sarcoma (diffuse membranous); synovial sarcoma; alveolar rhabdomyosarcoma (some); mesenchymal chondrosarcoma; solitary fibrous tumor
Neuron-specific enolase (NSE)	Neuroblastoma; Ewing sarcoma (some); desmoplastic small round cell tumor
Chromogranin	Neuroblastoma (often—in undifferentiated tumors); Ewing sarcoma (some)
Synaptophysin	Neuroblastoma (often—in undifferentiated tumors); Ewing sarcoma (some)
Cytokeratin	Synovial sarcoma; epithelioid sarcoma; angiosarcoma (epithelioid angiosarcoma); epithelioid hemangioendothelioma; desmoplastic small round cell tumor; leiomyosarcoma ~30 %; extrarenal rhabdoid tumor
Epithelial membrane antigen (EMA)	Synovial sarcoma; epithelioid sarcoma; epithelioid angiosarcoma; epithelioid hemangioendothelioma; low-grade fibromyxoid sarcoma (some); perineurioma; angiomatoid fibrous histiocytoma
HMB45; Melan-A; tyrosinase	Melanotic schwannoma; clear cell sarcoma; PEComa
MDM2/CDK4	Atypical lipomatous tumor; dedifferentiated liposarcoma
Claudin-1	Synovial sarcoma; epithelioid sarcoma; perineurioma; Ewing sarcoma (some)
Bcl-2	Synovial sarcoma; solitary fibrous tumor; spindle-cell lipoma/pleomorphic lipoma
CD117 (c-kit); DOG 1	Gastrointestinal stromal tumor
MUC-4	Low-grade fibromyxoid sarcoma; sclerosing epithelioid fibrosarcoma
ALK	Inflammatory myofibroblastic tumor
β-catenin (aberrant nuclear expression)	Desmoid fibromatosis ( normal cells with cytoplasmic staining)
INI-1 (loss of expression)	Proximal-type epithelioid sarcoma, extrarenal rhabdoid tumor, epithelioid malignant peripheral nerve sheath tumor (some)

*PEComa* perivascular epithelioid cell tumor

histologic type (specific sarcoma versus metastasis or lymphoma), and to assess the histologic grade (dedifferentiated liposarcoma versus well-differentiated liposarcoma). However, it is not always necessary to reach a specific histologic type.

The necessary diagnostic level of an FNA depends on treatment choice. For many sarcomas seated in the extremities or chest wall, primary radical resection is envisaged; in these instances, therefore, the most important information for treatment planning is the size and site of the mass and its proximity to vessels, nerve bundles, and bone. A correct diagnosis of sarcoma is a sufficient diagnostic level to plan surgical treatment. When neoadjuvant

therapy (chemo- or radiotherapy) is the treatment option before surgery (*e.g.*, pediatric sarcomas such as small blue round cell tumors and other high-grade sarcomas), the FNA diagnosis must include the correct histologic type and grade.

FNA diagnosis of a benign soft tissue neoplasm should also include histologic type or at least the basic histogenesis (*e.g.*, lipomatous tumor, nerve sheath tumor, pseudosarcomas), as different benign tumors are managed differently. Most nonsymptomatic benign tumors (lipoma, nodular fasciitis) could be put for “wait and watch,” and some are treated with limited excision (schwannoma, myxoma), whereas a few such tumors need a wider excision to prevent

**Table 13.2** Selected soft tissue tumors with cytogenetic alterations and commercially available FISH probes

Tumor	Cytogenetic aberration(s)	FISH probe(s)
Angiomatoid fibrous histiocytoma	t(12;16)(q13;p11)	<i>FUS</i>
	t(12;22)(q13;q12)	<i>EWSR1</i>
	t(2;22)(q34;q12)	<i>EWSR1</i>
Alveolar soft part sarcoma	der(X)t(X;17)(p11;q25)	<i>TFE3</i>
Clear cell sarcoma	t(12;22)(q13;q12)	<i>EWSR1</i>
	t(2;22)(q33;q12)	<i>EWSR1</i>
Dermatofibrosarcoma protuberans/giant-cell fibroblastoma	r(17;22)(q21;q13)/t(17;22)(q21;q13)	<i>PDGFB</i>
Desmoid fibromatosis	Trisomy 8 and 20	<i>CEP8/CEP20</i>
Desmoplastic small round cell tumor	t(11;22)(p13;q12)	<i>EWSR1</i>
Ewing sarcoma/primitive neuroectodermal tumor	t(11;22)(q24;q12)	<i>EWSR1</i>
	t(21;22)(q22.2;q12)	<i>EWSR1</i>
	t(2;22)(q33;q12)	<i>EWSR1</i>
	t(7;22)(p22;q12)	<i>EWSR1</i>
	t(17;22)(q12;q12)	<i>EWSR1</i>
	t(20;22)(q13;q12)	<i>EWSR1</i>
Extraskeletal myxoid chondrosarcoma	t(9;22)(q22;q12)	<i>EWSR1</i>
	t(9;17)(q22;q11)	<i>NR4A3</i>
	t(9;17)(q22;q11)	<i>NR4A3</i>
	t(3;9)(q11;q22)	<i>NR4A3</i>
Inflammatory myofibroblastic tumor	2p23 rearrangement	<i>ALK</i>
Infantile fibrosarcoma	t(12;15)(p13;q26)	<i>ETV6</i>
Lipoma, ordinary	12q14.3 rearrangement	<i>HMGA2</i>
Liposarcoma, well differentiated/dedifferentiated	Ring/giant marker chromosomes from chromosome 12q	<i>HMGA2/MDM2</i>
Liposarcoma, myxoid/round cell	t(12;16)(q13;p11)	<i>DDIT3/FUS</i>
	t(12;22)(q13;q12)	<i>DDIT3/EWSR1</i>
Low-grade fibromyxoid sarcoma	t(7;16)(q34;p11)	<i>FUS</i>
	t(1;16)(p11;p11)	<i>FUS</i>
Myoepithelioma, soft tissue	t(19;22)(q13;q12)	<i>EWSR1</i>
	t(1;22)(q23;q12)	<i>EWSR1</i>
	t(6;22)(p21;q12)	<i>EWSR1</i>
Rhabdomyosarcoma, alveolar	t(2;13)(q35;q14)	<i>FOXO1</i>
	t(1;13)(p36;q14), double minutes	<i>FOXO1</i>
Synovial sarcoma	t(X;18)(p11;q11)	<i>SS18</i>

*FISH* fluorescence in situ hybridization

recurrence. At present, criteria for cytologic diagnosis have been thoroughly defined in a number of sarcomas and benign soft tissue neoplasms. To facilitate evaluation and treatment based on FNAC, a standardized reporting of FNA results is necessary. At Lund Sarcoma Centre, the following diagnostic categories are used:

- Benign (histologic type)
- Sarcoma (histologic type and grade)
- Malignancy other than sarcoma
- Inconclusive
- Insufficient

The main diagnosis of either *benign* or *sarcoma* should be given, whenever possible, supplemented with histologic type of the neoplasm, and, in case of sarcoma malignancy, grade (low grade or high grade).

*Inconclusive* indicates inability to reliably determine whether a lesion is benign or malignant; whether a true sarcoma or other malignancy. *Insufficient* indicates technically failed biopsy.

### 13.1.5 Grading of Soft Tissue Sarcomas in FNA


The most widely used sarcoma grading system on histology is the FNCLCC system base on tumor differentiation, mitotic rate, and tumor necrosis [16]. Recently, vascular invasion and pattern of peripheral growth (pushing/infiltrative) of sarcomas are also shown to be important prognostic parameters [17].

However, assessment of these parameters is not feasible in FNA samples. Nevertheless, in technically satisfactory specimens, it is often possible to categorize a sarcoma into morphologically low-grade or high-grade malignancy. In a subset of sarcomas, a specific histologic diagnosis rendered may determine the grade (*see* Fig. 13.3) [10, 18–22].

### 13.1.6 Diagnostic Accuracy

If FNA biopsy is performed in experienced hands and with adequate samples, the diagnostic accuracy is high. In a retrospective 20-year study of 517 soft tissue tumors (315 benign





**Fig. 13.3** Sarcoma grading in FNA. Grade of malignancy may be determined by histologic diagnosis. (a) High-grade undifferentiated pleomorphic sarcoma: high grade (grade 3/3) by definition. (b) Myxoid liposarcoma: low grade (grade 2/3) by definition (H&E)

tumors and 202 sarcomas) from the Musculoskeletal Tumor Centre, Lund University Hospital, 28 false diagnoses (5 %) were rendered including 14 false-positive and 14 false-negative diagnoses, an inconclusive diagnosis in 13 cases (3 %), and insufficient for diagnosis in 29 tumors (6 %) [23]. In published reports during the past 10 years, a correct classification of soft tissue tumors as benign or malignant has accuracy around 90 %. In some reports, diagnostic accuracy of FNA can reach up to 97 % [6–8, 24].

### 13.1.7 Limitations of FNA in the Diagnosis of Soft Tissue Tumors

Despite of the reported high degree of accuracy in distinguishing malignant soft tissue tumors from benign ones, FNAC has been less successful in establishing their histologic types and grades, compared to an excisional biopsy. There are three major limitations in FNA of soft tissue lesions: (1) sampling error (the tumor or the diagnostic area was missed); (2) insufficient or suboptimal material

aspirated (poor yield, technically inferior smearing, necrosis or hemorrhage, or insufficient material for diagnostic ancillary studies); and (3) misinterpretation of the material (rare or new entities).

### 13.1.8 Complications of FNA of Soft Tissue and Bone Tumors

Complications of a correctly performed FNA in examination of soft tissue and bone lesions are minor and include small hematomas and localized tenderness. The risk of tumor cell spread in the needle tract must be addressed. This complication is strongly related to the size of the needles and the number of passes with the needles; using regular 22–25 gauge needles, for example, the incidence of this event is exceedingly low [25]. To prevent this complication, the same needle-insertion point is recommended for several punctures when sampling a suspected sarcoma. Tattooing of the needle-insertion point helps to remove the needle track during surgery (*see* Figs. 1.52 and 1.53) [14].

## 13.2 Normal Elements

### 13.2.1 Fibroblasts/Myofibroblasts


Fibroblasts are spindle-shaped cells with slender contours, containing ovoid or rounded nuclei with fine, evenly distributed chromatin and inconspicuous nucleoli. Well-preserved fibroblasts show unipolar or bipolar cytoplasmic processes (*see* Fig. 13.4a). Bare nuclei are also common findings in smears. In reactive connective tissue, fibroblasts/myofibroblasts present as fusiform, rounded, or triangular cells of variable sizes, with enlarged, irregular nuclei, and occasionally coarse chromatin and prominent macro nucleoli. In addition, reactive fibroblasts/myofibroblasts contain frequently abundant cytoplasm with one to several processes or angulated cytoplasmic extensions (*see* Fig. 13.4b–d). Binucleated cells are also common.

### 13.2.2 Fat Tissue

Normal fat tissue in FNA smears occurs as clusters of fat cells with abundant univacuolated cytoplasm and peripherally situated small nuclei. Slender capillaries are frequent findings in the clusters (*see* Fig. 13.5a). Smears from reactive fat tissue may show a myxoid background, increasing numbers of fibroblasts and endothelial cells, and occasionally adipocytes with a multivacuolated cytoplasm resembling lipoblasts (*see* Fig. 13.5b–e). Histiocytes with vacuolated or foamy cytoplasm seen in reactive fat may be occasionally difficult to distinguish from hibernoma cells.

### 13.2.3 Striated Muscle

Muscle fibers are pink or amphophilic on Papanicolaou staining, eosinophilic with hematoxylin and eosin (H&E),




**Fig. 13.4** Fibroblasts/myofibroblasts. (a) Loose cluster of well-preserved fibroblasts with uniform spindle-shaped nuclei and cytoplasmic processes (MGG). (b–d) Reactive fibroblasts/myofibroblasts

with enlarged nuclei, macronucleoli, and abundant cytoplasm with one to several processes or angulated cytoplasmic extensions (MGG; H&E)




and deep blue in May–Grünwald–Giemsa (MGG)-stained preparations. Peripherally placed small nuclei and cross-striations may be observed occasionally (*see* Fig. 13.6). Damaged and regenerating striated muscle fibers appear in

smears as multinucleated cells with varying shapes of “muscle giant cells,” which have nuclei arranged in rows or eccentrically placed. Nucleoli may be large and prominent (*see* Fig. 13.7).




**Fig. 13.5** Fat tissue. (a) Normal fat tissue. Clusters of adipocytes with abundant univacuolated cytoplasm and peripherally situated small nuclei. (b) Smears from reactive adipose tissue: an increasing number

of fibroblasts and endothelial cells (H&E). Occasional myxoid background (c) (MGG) and adipocytes (d and e) with a multivacuolated cytoplasm resembling lipoblasts (H&E)



**Fig. 13.6** Striated muscle. (a, b) Muscle fibers are deep blue in MGG and eosinophilic in H&E stainings. Note peripherally placed small nuclei and cross-striations (MGG; H&E)



**Fig. 13.7** Damaged and regenerating striated muscle. (a–c) “Muscle giant cells” with nuclei arranged in rows or eccentrically placed and prominent nucleoli (MGG; H&E)




### 13.3 Non-neoplastic Soft Tissue Tumors

#### 13.3.1 Pseudosarcomatous Proliferations

Relatively common soft tissue lesions constitute several entities, which are mimics of malignant spindle-cell and pleomorphic neoplasms. These benign proliferations can cause considerable diagnostic difficulties in FNAC, as smears are usually cellular containing proliferating fibroblasts and myofibroblasts with a wide variation in size and shape of the nuclei. Spindle-shaped cells with cytoplasmic processes and fusiform nuclei are the most common cell type, but plump cells with ovoid, rounded, or irregular nuclei are also present. Despite hypercellularity and nuclear pleomorphism throughout the smears, the chromatin in all cells is finely granular. Another typical finding is the presence of polyhedral or triangular cells with abundant cytoplasm and one or two rounded nuclei at the periphery near the cytoplasmic membrane, resembling ganglion cells.

#### 13.3.2 Nodular Fasciitis


Nodular fasciitis (NF) occurs in any age group but most commonly in young adults and children. Most NF occurs subcutaneously as a rapidly growing, painful, or tender mass, and the common sites are the upper extremities, trunk, and head and neck region. The lesion is composed of fibroblasts and myofibroblasts showing variable degrees of anisocytosis and anisokaryosis (*see* Fig. 13.8). Because of the above mentioned clinical presentation, most nodular fasciitis is needed at its early phase. In this phase, the matrix is often myxoid, occasionally giving the impression of a low-grade myxoid sarcoma such as myxofibrosarcoma (*see* Fig. 13.9). Persistent NF often yields smears containing relatively uniform, haphazardly arranged myofibroblasts producing a so-called cell culture appearance. Collagenous matrix is often present (*see* Fig. 13.10). The vast majority of NFs regress




**Fig. 13.8** Nodular fasciitis. (a) Cellular clusters of spindle-shaped myofibroblasts with cytoplasmic processes and fusiform nuclei showing variable degrees of anisocytosis and anisokaryosis. (b) Occasional

osteoclast-like giant cells (H&E). (c, d) Dispersed spindle and plump cells with ovoid, irregular nuclei, abundant cytoplasm, and occasional mitoses (MGG)





**Fig. 13.9** Nodular fasciitis. (a, b) Clusters of spindle cells or dispersed spindle cells in a prominent myxoid background. Note the admixed inflammatory cell (MGG)



**Fig. 13.10** Nodular fasciitis. (a) Cluster of myofibroblasts embedded in collagenous matrix. (b, c) Loosely cohesive and dispersed relatively uniform myofibroblasts resembling a so-called cell culture appearance (H&E)

spontaneously, and a “wait and see” treatment option is indicated in cases with a typical clinical presentation and when characteristic cytomorphology is seen in FNA smears [26–28].


*Cytologic features:*

- Cellular aspirates
- Myxoid background matrix
- Dispersed spindle cells mixed with clusters and sheets of spindle cells
- Cell culture-like appearance
- Variable anisocytosis and anisokaryosis
- Ganglion-like cells
- Admixture of inflammatory cells and histiocytes
- Occasional multinucleated giant cells
- Mitoses

*Differential diagnosis and problems in diagnosis:*


- Pleomorphic, spindle-cell, and myxoid sarcomas
- Desmoid fibromatosis

The related pseudosarcomatous lesions include proliferative fasciitis and proliferative myositis, which develop predominantly in adults. Proliferative myositis commonly arises in the trunk, whereas proliferative fasciitis is more common in the extremities. Both proliferative fasciitis and myositis share many cytologic features with nodular fasciitis. However, the myxoid background matrix is less prominent, and the ganglion-like cells are usually numerous and often exhibit large nucleoli in proliferative fasciitis (*see Fig. 13.11*). In proliferative myositis, smears contain multinucleated regenerating muscle fibers in addition to the fibroblastic–myofibroblastic cells (*see Fig. 13.12a, b*). Mitotic figures can be found in both smears (*see Fig. 13.12c*).



**Fig. 13.11** Proliferative fasciitis. (a–c) Loosely cohesive clusters or dispersed relatively pleomorphic cells with round to oval and spindle-shaped nuclei. Note admixture of numerous ganglion cell-like cells with binucleation and abundant cytoplasm (H&E; MGG)





**Fig. 13.12** Proliferative myositis. (a, b) Dispersed fibroblastic–myofibroblastic cells with admixture of muscle fragments, multinucleated giant cells, and regenerating muscle fibers. (c) Mitotic figures can be occasionally found in the smears (MGG)

### 13.3.3 Myositis Ossificans

Myositis ossificans (MO) is a rapidly growing soft tissue lesion, which is often mistaken clinically as sarcoma especially osteosarcoma. It most often arises in the subcutaneous tissue or musculature of the extremities in young adults. The mass is often tender and presents ossification in a zonal pattern. The characteristic cytologic finding is the mixture of proliferating fibroblasts/myofibroblasts, osteoblasts, osteoclast giant cells, occasionally regenerating muscle fibers (“muscle giant cells”) (see Fig. 13.13), and rarely bone fragments. MO may resolve spontaneously some weeks/months after FNA examination [29].

#### *Cytologic features:*

- Proliferating spindle cells—fibroblasts/myofibroblasts with variable anisocytosis and anisokaryosis
- Osteoblasts, often with reactive changes
- Osteoclasts
- Muscle “giant cells”

- Occasionally small calcifications
- Rare mitoses

#### *Differential diagnosis and problems in diagnosis:*

- Pleomorphic soft tissue sarcoma
- Extraskelletal osteosarcoma

### 13.3.4 Ischemic Fasciitis (Atypical Decubital Fibroplasia)


Ischemic fasciitis is a reactive process arising as a poorly circumscribed, non-ulcerated mass in the deep subcutaneous tissue, and is usually associated with chronic pressure of areas over the bone. Ischemic fasciitis is composed of large ganglion-like cells, fibroblasts, and myofibroblasts set in a myxoid stroma. In addition, smears may contain inflammatory cells, necrotic debris, and cells with large, irregular nuclei and dense and smudged chromatin (see Fig. 1.50).




### 13.3.5 Amyloidoma (Tumoral Amyloidosis)

Amyloidoma is an uncommon lesion caused by amyloid deposition and not necessarily associated with systemic

amyloidosis. Soft tissue and bone amyloidomas are very rare [30, 31]. Smears are hypocellular and contain fragments of amorphous material, with admixed bland fibroblasts and occasional calcifications (*see Fig. 13.14*).




**Fig. 13.13** Myositis ossificans. (a–c) Mixture of proliferating myofibroblasts, osteoblasts, and osteoclast-like giant cells (MGG) without cytologic atypia. (d) Rarely bone fragments (H&E)



**Fig. 13.14** Tumoral amyloidosis. (a, b) Deposition of amorphous material with bland fibroblasts and calcifications (H&E)





**Fig. 13.15** Gout. (a) In low-power view, brown-stained sheath of crystals with adjacent multinucleated giant cells (MGG). (b–d) Monosodium urate crystals are easily identified in high power with (d) or without polarized light (MGG; H&E)

### 13.3.6 Gout

Disturbed uric acid metabolism results in gout tophi, which is defined by monosodium urate crystal deposition in the joint spaces and/or in the soft tissue adjacent to the joints. Laboratory tests showing an elevated serum

uric acid level and microscopic examination of joint effusions help to establish the diagnosis. FNA cytologic diagnosis of soft tissue gout tophi can be easily rendered because monosodium urate crystals are readily identifiable on smears with or without polarized light (*see* Fig. 13.15) [32].

## 13.4 Benign Adipocytic Tumors

### 13.4.1 Lipoma

Lipomas, the most common soft tissue tumor of adults, usually present as slowly growing, solitary or multiple, subcutaneous masses and can be occasionally deeply seated (intramuscular or intermuscular lipoma). They are composed of mature adipose tissue but may include other connective tissue elements and regressive/degenerative areas, which may cause difficulties in FNA examination.

#### *Cytologic features:*

- Fatty tissue fragments composed of cells containing a single fat vacuole and a small, dark, peripheral nucleus
- Variable number of capillaries within the fatty tissue fragments
- Few dissociated adipocytes
- Fragments of striated muscle or regenerating muscle in inter/intramuscular lipoma
- Occasional myxoid matrix
- Rarely metaplastic cartilage/bone

Aspirates from lipoma are identical to those from normal adipose tissue. It is important to ensure that the needle is placed within the mass to avoid contamination with subcutaneous fat during the biopsy of a deeply seated mass.

### 13.4.2 Angiolipoma

Angiolipomas are subcutaneous lipomatous tumors. They are often multiple and tender at palpation. Most angiolipomas are smaller than 2 cm. An angiolipoma should be suspected when the tumor is small and tender at palpation, in conjunction with the FNA findings of numerous branching capillary vessels within the fat tissue fragments.

### 13.4.3 Lipoblastoma/Lipoblastomatosis


This tumor of infancy most commonly involves the extremities and presents with either a well-circumscribed subcutaneous tumor (lipoblastoma) or a deep-seated, diffusely infiltrative, ill-defined mass (lipoblastomatosis). Histologically composed of immature fat with mesenchymal, myxoid, and fibrotic areas, the tumor is thought to be capable of maturation to a common lipoma. The typical features include small uniform adipocytes with vacuolated cytoplasm and round uniform nuclei in a myxoid background matrix, admixed with branching strands of capillaries (see Fig. 13.16). Smears may also be dominated by large ordinary univacuolated adipocytes. Although lipoblastoma mimics myxoid liposarcoma cytologically, two tumors occur in different age groups: lipoblastoma in boys younger than 3 years old and myxoid liposarcoma in young adults [33].

#### *Cytologic features:*

- Fatty tissue fragments with variable amounts of myxoid matrix
- Branching strands of thin capillaries
- Few dissociated adipocytes
- Occasionally uni- or multivacuolated lipoblast-like cells
- Occasionally hibernoma-like cells


### 13.4.4 Spindle Cell/Pleomorphic Lipoma

These two benign lipomatous tumors share very similar clinical, morphologic, and cytogenetic features and represent a morphologic continuum. For both tumors, the typical clinical setting is a subcutaneous tumor in the posterior neck region, upper back, and over the shoulders in middle-aged men. Head and mouth areas, including tongue and orbit, are less common sites.



**Fig. 13.16** Lipoblastoma. (a, b) FNA smears contain large fragments of fatty tissue with variable amounts of myxoid matrix and branching strands of thin capillaries





**Fig. 13.17** Spindle-cell lipoma. (a, b) Mixture of loosely cohesive clusters and dispersed bland spindle cells, mature adipose tissue, and fragments of brightly eosinophilic collagen–hyaline fibers (H&E). (c, d) Myxoid background and mast cells are common (MGG)

The cytomorphology of spindle-cell lipoma is often distinct, although proportions of different elements vary from case to case. The smears contain fat fragments mixed with fascicles of uniform spindle cells with elongated, uniform nuclei and collagen fibers, often in a myxoid background matrix (see Fig. 13.17) [34]. The characteristic finding of pleomorphic lipoma is the presence of floret cells: multinucleated giant cells with hyperchromatic nuclei and a moderate amount of cytoplasm (see Fig. 13.18).

*Cytologic features of spindle-cell lipoma:*

- Mixture of mature adipose tissue and dispersed or clustered, bland spindle cells
- Fragments of brightly eosinophilic (H&E) collagen–hyaline fibers
- Myxoid background common
- Mast cells (particularly in cases with the myxoid background)

*Differential diagnosis and problems in diagnosis:*

- Schwannoma
- Dermatofibrosarcoma protuberans

- Low-grade myxofibrosarcoma
- Myxoid liposarcoma

*Cytologic features of pleomorphic lipoma:*


- Fragments of mature fat
- Variable number of floret cells
- Variable number of spindle cells
- Occasionally myxoid background matrix
- Occasionally mast cells

*Differential diagnosis and problems in diagnosis:*


- Well-differentiated liposarcoma (atypical lipomatous tumor)

A diagnostically important but variable sign in smears is the presence of fragments of eosinophilic (H&E) collagen–hyaline fibers. Depending on the areas sampled from this very heterogenous tumor, aspirates from spindle-cell lipoma may show a predominance of adipose tissue fragments or a predominance of spindle-cell fascicles or an abundant myxoid matrix. In addition to floret cells, smears from pleomorphic lipomas may contain collagen fibers and areas of myxoid stroma similar to spindle-cell lipoma. Hybrid forms with both spindle and pleomorphic





**Fig. 13.18** Pleomorphic lipoma. (a) In scanning power fragments of mature fat, variable number of floret cells and spindle cells (H&E). (b, c) Floret cells and occasionally prominent myxoid background matrix (H&E; MGG)



**Fig. 13.19** Spindle-cell lipoma. Positive CD34 staining in the spindle cells component (ThinPrep)


lipoma features are not uncommon. The spindle cells are positive for CD34 (see Fig. 13.19) and negative for S-100 protein.

### 13.4.5 Hibernoma

Hibernoma is a rare lipomatous tumor of brown fat derivation. It is commonly situated in the interscapular region, on the back or chest wall (sites of normal deposits of brown fat), but can also occur in the extremities. Hibernomas are usually subcutaneous but may also be deep seated (intramuscular). In smears, clusters or fragments of ordinary large adipocytes are intermingled with round to oval and polygonal cells of variable sizes, with vacuolated or granular cytoplasm and centrally placed small uniform nuclei. The tissue fragments often contain numerous capillary vessels (see Fig. 13.20). Ordinary lipoma-like fat cells may dominate the smears and the typical hibernoma cells may be in minority, thus posing challenges to diagnose [35].

#### Cytologic features:

- Fragments of adipocytes and hibernoma cells
- Hibernoma cells: abundant microvacuolated to granular cytoplasm
- Small, bland nuclei



**Fig. 13.20** Hibernoma. (a–c) Clusters, sheets of round to oval and polygonal cells with abundant microvacuolated or granular cytoplasm, and centrally placed small uniform nuclei, sometimes mimicking a

lipoblast (b) (MGG; H&E). (d) Overall hibernoma architecture is preserved in cell-block material (H&E; Cell Block)

- Numerous capillaries
- Occasionally lipoblast-like cells

*Differential diagnosis and problems in diagnosis:*

- Well-differentiated liposarcoma (atypical lipomatous tumor)
- Lipoma with fat necrosis
- Granular cell tumor
- Adult rhabdomyoma

### 13.4.6 Chondroid Lipoma

Chondroid lipoma is an infrequent deep-seated benign lipomatous tumor occurring in the proximal extremities, limb girdles, trunk, and head and neck region. FNA smears show clusters of mature adipocytes with admixture of lipoblasts and small chondroid cells in a background of abundant myxochondroid matrix (see Fig. 13.21) [36].

Despite variation in size and shape, the nuclei of small chondroid cells are bland appearing. One helpful feature in distinguishing it from myxoid liposarcoma is the lack of a plexiform capillary network in the cellular clusters.


*Cytologic features:*

- Variable but commonly abundant myxochondroid background matrix
- Sheets and clusters of ordinary adipocytes
- Sheets and clusters of uni- or multivacuolated, lipoblast-like cells
- Small chondroid cells with irregular, bland, occasionally coffee bean-shaped nuclei and granular cytoplasm
- Lack of a plexiform capillary network seen in myxoid liposarcoma

*Differential diagnosis and problems in diagnosis:*

- Myxoid liposarcoma
- Extraskeletal myxoid chondrosarcoma





**Fig. 13.21** Chondroid lipoma. (a–e) Clusters and small fragments of mature adipocytes with admixture of lipoblasts, small chondroid cells, and abundant myxochondroid background matrix (MGG). Note the lipoblast-like cells in (c)

### 13.4.7 Diagnostic Pitfalls in Benign Adipose Tumors

Distinguishing various benign lipomatous neoplasms from liposarcomas is important for proper clinical management.

The main clue to a benign diagnosis is the absence of atypical lipoblasts. Clinical details such as patient age, tumor location, and size are also valuable in the differential diagnosis.

The cytologic features of variants of benign lipomatous tumors and their mimics are summarized in Tables 13.3 and 13.4.

**Table 13.3** Cytologic features of benign lipomatous tumors and their mimics

Tumor	Cytologic features	Mimics
Common lipoma	Fragments of adipose tissue with large, univacuolated, mature fat cells with a small dark peripheral nucleus. Few dissociated adipocytes. Variable capillary vessels within the fatty tissue fragments. Occasional myxoid matrix, metaplastic cartilage/bone	Atypical lipomatous tumor
Inter- and intramuscular lipoma	As common lipoma + fragments of striated muscle or regenerating muscle, occasional muscle giant cells	Intramuscular angioma with prominent fat
Angiolipoma	As common lipoma + numerous branching capillary vessel fragments are in the fat tissue fragments	
Lipoblastoma/lipoblastomatosis	Clusters of small uniform adipocytes with vacuolated cytoplasm and round uniform nuclei. Myxoid background matrix. Branching strands of capillaries. Commonly admixture of normal looking fatty cells. Occasionally hibernoma-like cells and uni- or multivacuolated lipoblast-like cells	Myxoid liposarcoma
Spindle-cell lipoma	Mixture of mature adipose tissue and dispersed or clustered, bland spindle cells in variable proportions. Fragments of brightly eosinophilic (hematoxylin and eosin) collagen–hyaline fibers. Myxoid background and mast cells common	Schwannoma Dermatofibrosarcoma protuberans Low-grade myxofibrosarcoma Myxoid liposarcoma
Pleomorphic lipoma	Fragments of mature fat. Variable number of spindle cells and floret cells. Occasionally myxoid background matrix and mast cells	Well-differentiated liposarcoma (atypical lipomatous tumor)
Hibernoma	Clusters and fragments of large univacuolated, mature fat cells mixed with variable proportions of “hibernoma cells” (multivacuolated or granulated cytoplasm). Capillary network. Occasionally dissociated or small groups of hibernoma cells. Smears may be dominated by large mature fat cells	Well-differentiated liposarcoma (atypical lipomatous tumor) Lipoma with fat necrosis Granular cell tumor Adult rhabdomyoma
Chondroid lipoma	Clusters of mature fat cells mixed with uni- or multivacuolated, lipoblast-like cells. Variable but commonly abundant myxochondroid background matrix. Small chondroid cells with irregular, bland, occasionally coffee bean-shaped nuclei and granular cytoplasm	Myxoid liposarcoma Extraskeletal myxoid chondrosarcoma

**Table 13.4** Cytologic features in mimics of benign lipomatous tumors

Tumor	Cytologic features
Well-differentiated liposarcoma/atypical lipomatous tumor	Fragments/clusters of mature fat cells. Large atypical stromal cells with irregular, hyperchromatic, within and outside fragments/clusters. Rare atypical lipoblasts. Occasional floret cells
Myxoid liposarcoma	Tissue fragments with myxoid matrix and branching capillary network. Uniform, round to oval or spindle-shaped primitive cells. Uni- or multivacuolated lipoblasts, some with signet-ring features. No mitoses. Few dissociated tumor cells
Low-grade myxofibrosarcoma	Dispersed cells and cell clusters. Abundant myxoid background. Slight to moderate atypical spindle cells. Fragments of curved vessels embedded in the myxoid matrix. Occasional pseudolipoblasts
Extraskeletal myxoid chondrosarcoma	Dispersed cells, branching cords, strands, and cell balls. Spindle-shaped, fusiform, or round cells with ovoid or rounded nuclei. Bland nuclear chromatin and small nucleoli. Myxoid background matrix. Chondroblasts-like lacunar structures. Lack of significant vascularity
Schwannoma	Cohesive fragments of different sized, often with irregular borders (“of a jigsaw puzzle pieces”). Rare dispersed cells. Cells with indistinct cytoplasm and elongated, slender nuclei with pointed ends, boomerang-shaped or comma-like nuclei. Fibrillary background. Occasional nuclear palisading and Verocay bodies. Occasionally small lymphocyte-like nuclei or slight to moderate nuclear pleomorphism
Adult rhabdomyoma	Large, rounded, polygonal, or elongated cells, dispersed or arranged in cohesive, syncytial clusters. Abundant eosinophilic (hematoxylin and eosin) granular cytoplasm. Small nuclei with prominent nucleoli. Stripped nuclei occasionally. Rarely visible cross-striation
Granular cells tumor	Mixture of poorly cohesive sheets and single cells or bare nuclei. Cells with fragile granular cytoplasm, often stripped nuclei and cytoplasmic granular background. Preserved cells with abundant granular cytoplasm and ill-defined cytoplasmic borders. Small, rounded to oval nuclei with finely granular chromatin and small nucleoli. Occasionally moderately pleomorphic cells



## 13.5 Malignant Adipocytic Tumors

It has been estimated that about 20 % of soft tissue sarcomas in adults are liposarcomas. Most liposarcomas are deep seated, intra- or intermuscular, and the most common sites are the lower extremities, trunk, and retroperitoneum. There are three clinically, morphologically, and cytogenetically distinct entities: well-differentiated/dedifferentiated liposarcoma, myxoid/round cell liposarcoma, and pleomorphic liposarcoma [1].


### 13.5.1 Atypical Lipomatous Tumor/Well-Differentiated Liposarcoma

Well-differentiated liposarcoma is a locally aggressive, non-metastasizing adipocytic neoplasm occurring predominantly in the retroperitoneum or mediastinum. When the same tumor arises at surgically curable sites such as the lower extremities, the term “atypical lipomatous tumor” (ALT) is preferred [1]. Cytologic findings in these tumors include the mixture of

ordinary lipoma-like fragments and atypical stromal cells with enlarged, irregular hyperchromatic nuclei (*see Fig. 13.22*). Atypical lipoblasts with cytoplasmic vacuoles and scalloped nuclei may be present but are not necessary for the diagnosis. Dedifferentiated liposarcoma is diagnosed when a well-differentiated liposarcoma also displays non-lipogenic sarcomatous areas (*see Fig. 13.22e*). Well-differentiated liposarcoma/ALT is characterized by supernumerary ring and giant marker chromosomes, resulting in MDM2 and CDK4 amplification. These genetic abnormalities can be detected by FISH showing MDM2 amplification and/or by immunohistochemistry showing MDM2 and/or CDK4 nuclear reactivities (*see Fig. 13.22f* and Table 13.2) [37].


*Cytologic features:*

- Clusters or sheets of lipogenic cells
- Large atypical stromal cells with irregular, hyperchromatic nuclei
- Rare, multivacuolated atypical lipoblasts
- Occasional floret cells
- Non-lipogenic elements suggestive of dedifferentiation



**Fig. 13.22** Atypical lipomatous tumor/well-differentiated liposarcoma. (a, b) Mixture of ordinary lipoma-like fragments and atypical stromal cells with enlarged, irregular hyperchromatic nuclei (H&E). (c, d) Occasional myxoid background and floret cells in smears (MGG).

(e) Dedifferentiated liposarcoma with non-lipogenic pleomorphic/spindle cells (Pap stain). Immunoreactivity with MDM2 (f) and CDK4, indicative of MDM2/CDK4 amplification, is characteristic for ALT/well-differentiated liposarcoma and dedifferentiated liposarcoma



**Fig. 13.22** (continued)

*Differential diagnosis of well-differentiated liposarcoma/ALT:*

Variants of benign lipomatous tumor:

- Hibernoma, pleomorphic lipoma, and chondroid lipoma
- Fat necrosis with lipophages
- Any other mesenchymal tumor with nuclear atypia and a fat component

*Differential diagnosis of dedifferentiated liposarcoma:*

- Undifferentiated pleomorphic sarcoma
- Myxofibrosarcoma
- Myogenic sarcoma
- Osteosarcoma
- Other fibroblastic/myofibroblastic tumor

### 13.5.2 Myxoid/Round Cell Liposarcoma

Myxoid/round cell liposarcoma accounts for 15–20 % of liposarcomas. It occurs in young adults and the most common site of involvement is in the deep soft tissue of the thigh.

#### 13.5.2.1 Myxoid Liposarcoma

*Cytologic features:*

- Tissue fragments with myxoid matrix and a branching capillary network
- Monotonous appearance with minimal pleomorphism
- Uni- or multivacuolated lipoblasts, some with signet-ring features
- Round to ovoid non-lipogenic primitive mesenchymal cells
- No mitoses
- Few dispersed tumor cells

*Differential diagnosis and problem in diagnosis:*

- Spindle-cell lipoma with abundant myxoid matrix
- Intramuscular myxoma, cellular type
- Low-grade myxofibrosarcoma

The diagnostic features of myxoid liposarcoma are the triad of abundant myxoid background matrix, fragments of tumor tissue with a branching network of thin capillaries, and slightly atypical lipoblasts as well as round to ovoid primitive cells, often associated with the capillaries in tissue fragments (see Figs. 13.23 and 13.24).

#### 13.5.2.2 Round Cell Liposarcoma (Hypercellular Myxoid Liposarcoma)

A subset of myxoid liposarcomas have hypercellular areas containing numerous round to ovoid cells with scant cytoplasm, mixed with few to none lipoblasts. The myxoid matrix is scanty and the capillary network less prominent, often obscured by clusters of tumor cells (see Fig. 13.25) [38, 39].

*Cytologic features:*


- Rich yield of clusters and clumps of round or ovoid tumor cells embedded in matrix
- Tumor cells with a high nuclear/cytoplasmic (N/C) ratio and round nuclei with vesicular chromatin
- Less conspicuous myxoid matrix and capillaries compared to myxoid liposarcoma
- Few to none lipoblasts

*Differential diagnosis and problems in diagnosis:*

- Other types of round cell sarcoma infiltrating adipose tissue
- Soft tissue metastasis of renal cell carcinoma

Most myxoid/round cell liposarcoma is characterized by the recurrent translocation t(12;16)(q13;p11) with fusion of the *DDIT3* gene on chromosome 12 and the *FUS* gene on





**Fig. 13.23** Myxoid liposarcoma. Tissue fragments with round to ovoid primitive mesenchymal cells, myxoid matrix, and branching capillary network (a–c) (H&E); (d) (MGG)

chromosome 16. In a small subset case, an *EWSRI-DDIT3* fusion resulting from the translocation  $t(12;22)(q13;q12)$  is present. The genetic abnormality can be detected by FISH for *DDIT3* rearrangement, which is very helpful in difficult cases, especially the round cell variant. Immunocytochemistry is usually neither necessary nor helpful to establish the diagnosis (Table 13.2) [40].

### 13.5.3 Pleomorphic Liposarcoma

Pleomorphic liposarcoma is a rare, high-grade liposarcoma containing markedly atypical, bizarre, and giant multinucleated tumor cells with adipocytic differentiation (see Fig. 13.26). Most cases occur in the extremities in elderly

patients. The most important clue to the diagnosis is the presence of highly atypical uni- or multinucleated lipoblasts. Mitoses and necrosis are common [37].


#### *Cytologic features:*

- Dispersed cells and cell clusters
- Pleomorphic tumor cells including multinucleated giant tumor cells
- Variable presence of highly atypical, uni- or multinucleated lipoblasts
- Cells with hyaline cytoplasmic droplets occasionally present
- Necrosis


#### *Differential diagnosis and problem in diagnosis:*

- Pleomorphic high-grade sarcoma of another lineage
- Poorly differentiated carcinoma






**Fig. 13.24** Lipoblasts in myxoid liposarcoma. (a–d) Uni- or multivacuolated lipoblasts, some with signet-ring features, often associated with the capillary network (MGG)




**Fig. 13.25** Round cell liposarcoma (hypercellular myxoid liposarcoma). (a–d) Capillary network less prominent, often obscured by clusters of tumor cells (H&E; MGG)





**Fig. 13.25** (continued)



**Fig. 13.26** Pleomorphic liposarcoma. (a–c) Smears are similar to non-lipogenic high-grade sarcomas but with admixed atypical lipoblasts (MGG). (d) Markedly atypical, bizarre, and giant multinucleated tumor cells (H&E)



## 13.6 Fibroblastic/Myofibroblastic Tumors


### 13.6.1 Desmoid Fibromatosis

Fibromatoses represent a broad spectrum of infiltrative and non-metastasizing myofibroblastic neoplasms. Extra-abdominal deep fibromatosis arises usually in the proximal parts of extremities, in the pelvic girdle and shoulder region, commonly in young adults and teenagers. Fibromatosis is composed of fascicles of fibroblasts/myofibroblasts embedded in a collagenous stroma. Because of abundant collagenous stroma, it is difficult to obtain sufficient material from the lesion, and therefore, vigorous aspiration using larger needles and/or core needle biopsy is often necessary [41, 42]. The mixture of uniform, occasional slightly atypical spindle cells and fragments of paucicellular collagenous stroma in aspirates from a deep-seated, firm, soft tissue tumor is the hallmark of desmoid fibromatosis. Bland spindle cells

arranged in long fascicles is an additional important cytologic feature suggestive of desmoid (*see* Fig. 13.27) [41]. Smears from desmoids may contain myxoid matrix, evoking the differential diagnosis of low-grade myxoid spindle-cell lesions, such as nodular fasciitis or low-grade fibromyxoid sarcoma. Demonstration of aberrant nuclear expression of  $\beta$ -catenin is helpful to confirm the diagnosis (*see* Fig. 13.27g).

#### Cytologic features:


- Variable yield
- Clusters of or dispersed fibroblasts
- Bland or slightly atypical spindle cells in collagenous stroma arranged in long fascicles
- Fibroblasts with elongated fusiform nuclei, insignificant nucleoli, and slight anisokaryosis
- Cytoplasmic processes are seen in preserved cells but stripped nuclei are a common finding
- Fragments of paucicellular collagenous stroma



**Fig. 13.27** Desmoid fibromatosis. (a–d) The mixture of cohesive or loosely cohesive clusters and dispersed uniform or slightly atypical spindle cells and fragments of paucicellular collagenous stroma (MGG; H&E). (e, f) Bland spindle cells in long fascicles, a useful clue

suggestive of fibromatosis (H&E; MGG). (g) Aberrant nuclear expression of  $\beta$ -catenin is characteristic for desmoid fibromatosis. Note the negative nuclear staining in adjacent endothelial cells





**Fig. 13.27** (continued)

- Occasional regenerating striated muscle fibers (muscle giant cells)

*Differential diagnosis and problems in diagnosis:*

- Nodular fasciitis
- Low-grade fibromyxoid sarcoma
- Deep-seated leiomyoma
- Low-grade malignant peripheral nerve sheath tumor (MPNST)
- Monophasic synovial sarcoma
- Fibrosarcoma

### 13.6.2 Elastofibroma Dorsi

Elastofibroma dorsi (EFD) is a relatively rare, soft tissue mass, probably of reactive nature. The lesion is typically slowly growing, located near the inferior margin of the scapula or between the inferior part of scapula and the chest wall in elderly women. Elastofibroma is composed of (myo)fibroblasts, abundant hypocellular collagenous stroma, and fat.


The main diagnostic feature is the presence of elastic fibers with serrated borders, corresponding to faulty elastin fibrillogenesis [43]. FNA smears show small sheets or clusters of uniform spindle cells, mature adipocytes, fragments of acellular collagen bundles and fibers, and the degenerated elastic fibers presented as linear (“braid-like”), globular, and stellate structures with serrated edges (*see Fig. 13.28*). Diagnostic difficulties arise when the material is poor, fat cells are dominant, and the typical serrated elastic fibers are missing.

*Cytologic features:*

- Variably cellular smears
- Fat tissue intermingled with collagen fragments/collagen fibers and/or uniform spindle cells
- Spindle cells in loosely cohesive groups or dispersed
- Elastic fibers present as linear (“braid-like”), globular bodies with shell-like and stellate appearances, and the characteristic serrated “moth-eaten” borders

*Differential diagnosis and problems in diagnosis:*

- Extra-abdominal desmoid



**Fig. 13.28** Elastofibroma dorsi. (a) Smears show small clusters of and dispersed uniform spindle cells, mature adipocytes and fragments of acellular collagen with admixture of degenerated elastic fibers. (b–d) Elastic

fibers presented as linear (“braid-like”), distinct globular and stellate structures with serrated edges in high-magnification image (ThinPrep, Pap stain; Cell Block; Elastin stain; Hologic; Bedford, MA, USA)

### 13.6.3 Benign Fibrous Histiocytoma

Benign fibrous histiocytoma, a common neoplasm, typically arises in the skin or subcutaneous tissue, occasionally extends to deep soft tissue. The cytologic features are described in Sect. 14.10.


### 13.6.4 Extrapleural Solitary Fibrous Tumor

Solitary fibrous tumor (SFT) is a mostly benign fibroblastic tumor, which can occur practically at any location in the body besides the pleura. The common sites of involvement include deep soft tissue of the lower extremities, head and neck region, mediastinum, and retroperitoneum. Many SFTs

were previously termed hemangiopericytomas, and the rare, fat-forming variant was called lipomatous hemangiopericytoma [1]. Both malignant and dedifferentiated forms have been described [44, 45].

The aspirate yields variably cellular smears containing bland spindle cells, either dispersed or in tight clusters. The tumor cells are fibroblast like with spindle or ovoid nuclei and scanty cytoplasm. Naked nuclei, ropy collagen fibers, and mast cells are common (*see* Fig. 13.29). The prominent “staghorn” vascular pattern seen in histology is rarely appreciated in FNA smears but can be seen in cell-block sections (*see* Fig. 13.29d). The smears from the fat-forming variant SFT show three-dimensional clusters of uniform spindle cells admixed with mature adipose tissue. A correct diagnosis of SFT is difficult based on cytomorphology alone





**Fig. 13.29** Solitary fibrous tumor. (a, b) A mixture of dispersed cells and fascicle-like clusters of uniform population of bland spindle cells with inconspicuous nucleoli (MGG; H&E). (c) Clusters of spindle

cells mixed with mature fat in the fat-forming variant (MGG). (d) Classic architecture of SFT with HPC-like vessels better appreciated in the cell-block sections (H&E)

[46, 47]. Immunocytochemistry is often diagnostically helpful as the spindle cells of SFT display characteristic reactivities for CD34, CD99, and bcl-2 and limited reactivities for EMA and keratins; the profile distinguishes it from synovial sarcoma.

*Cytologic features:*

- Variable cellularity
- A mixture of dispersed cells and tight, fascicle-like clusters
- Uniform population of bland spindle cells with inconspicuous nucleoli
- Ropy collagen fibers
- Stripped nuclei
- Mast cells
- Three-dimensional clusters of uniform spindle cells mixed with mature fat in the fat-forming variant


*Differential diagnosis and problems in diagnosis:*

- Monophasic synovial sarcoma
- Fibromatosis
- Low-grade fibromyxoid sarcoma

- Low-grade malignant peripheral nerve sheath tumor (MPNST)
- Benign or malignant lipomatous tumors versus fat-forming variant SFT

### 13.6.5 Fibromatosis Colli (Torticollis)

Fibromatosis colli is a very firm tumor-like lesion in the sternocleidomastoid muscle in newborn infants. A diffuse proliferation of fibroblasts is present within the muscle tissue. The involved muscle fibers are atrophic and contain regenerating muscle giant cells. The FNA smears of fibromatosis colli include a double population of bland spindle cells and muscle giant cells with nuclei showing prominent nucleoli (*see Fig. 13.30*). There is a risk to misinterpret muscle giant cells as malignant cells. It is important to remember that many tumors regress spontaneously; the primary therapy is expectancy and clinical symptom control.



**Fig. 13.30** Fibromatosis colli (torticollis). (a–c) Hypocoellular smears with bland spindle cells and muscle giant cell with prominent nuclei (MGG)

*Cytologic features:*

- Bland fibroblasts and myofibroblasts dispersed or clustered
- Muscle giant cells and tadpole cells
- Occasionally small tufts of myxoid matrix
- Stripped nuclei

*Differential diagnosis and problems in diagnosis:*

- Pleomorphic sarcoma

### 13.6.6 Fibrous Hamartoma of Infancy

Fibrous hamartoma of infancy is a rare subcutaneous mass composed of a mixture of myxoid areas with small round primitive cells, septa or bands of fibrous tissue, and mature adipose tissue. Typical sites are the upper arms, shoulders, and axillary regions. In our limited experience of FNA of this entity, the spindle cells in fibrous hamartoma of infancy (see Fig. 13.31) resemble those in infantile fibrosarcoma. Most

infantile fibrosarcomas arise in the extremities and yield hypercellular smears compared to poor cellularity of spindle-cell component seen in fibrous hamartoma of infancy.

*Cytologic features:*

- A mixture of normal fatty tissue and bland spindle cells in sheets and clusters
- Variable amounts of myxoid matrix


*Differential diagnosis and problems of diagnosis:*

- Infantile fibrosarcoma

### 13.6.7 Low-Grade Fibromyxoid Sarcoma

Low-grade fibromyxoid sarcoma (LGFMS) is an uncommon, slowly growing, malignant fibroblastic neoplasm. It arises in the deep soft tissues of the extremities and trunk in young adults but may also occur in other age groups and locations. In histologic sections, LGFMS displays bland spindle cells arranged in a whorled pattern in a variable myxoid and





**Fig. 13.31** Fibrous hamartoma of infancy. (a–c) Mixture of normal fatty tissue and bland spindle cells in sheets and clusters (MGG); (H&E). (d) Variable amounts of dispersed spindle cells or bare nuclei (MGG)


collagenous stroma. In the myxoid areas, curvilinear vessels are a typical finding. The bland appearance may lead to the misperception of a benign spindle-cell or a myxoid lesion. Some tumors display rosette-like structures with a central hyalinized core surrounded by fibroblast-like cells. It is very difficult to render a diagnosis of LGFMS based on FNAC alone due to the morphologic overlapping with other spindle-cell and myxoid lesions. FNA smears usually show a mixture of uniform spindle to polygonal/rounded cells, embedded in a collagenous and myxoid matrix, and scattered single cells or bare nuclei with a mild anisokaryosis (see Fig. 13.32). The cytologic features of LGFMS have been described in case reports and in one series of eight cases [48–50]. LGFMS is characterized by a *FUS-CREB3L2* or a *FUS-CREB3L1* gene fusion, which can be detected by FISH, a very helpful ancillary test for difficult myxoid tumors [15, 40].

*Cytologic features:*

- Dispersed cells and cellular fascicles/clusters
- Fibroblast-like spindle cells with slightly atypical nuclei
- Monotonous appearance with mild anisokaryosis
- Stripped nuclei common
- Often abundant myxoid matrix
- Fragments of collagen tissue in the background
- Occasional fragments of curvilinear vessel in the myxoid areas

*Differential diagnosis and problems in diagnosis:*

- Intramuscular myxoma, cellular type
- Schwannoma
- Desmoid fibromatosis
- Low-grade myxofibrosarcoma
- Dermatofibrosarcoma protuberans (myxoid variant)



**Fig. 13.32** Low-grade fibromyxoid sarcoma. (a–c) Mixture of uniform or slightly pleomorphic spindle and polygonal/rounded cells, embedded in a collagenous and myxoid matrix and scattered single

cells or bare nuclei with mild to moderate anisokaryosis (MGG). (d) Occasionally large cohesive clusters of bland spindle cells embedded in collagenous/myxoid matrix (H&E)

### 13.6.8 Myxofibrosarcoma

Myxofibrosarcoma (MFS) is defined as a fibroblastic malignancy with variably myxoid stroma, cellular pleomorphism, and a prominent curvilinear vascularity. It was previously called “myxoid MFH.” MFS commonly arises in the subcutaneous tissue in the extremities of elderly patients but may also arise in deep soft tissue. Low-grade MFS is characterized by slightly and moderately atypical spindle cells, whereas high-grade MFS presents marked cellular and nuclear pleomorphism including multinucleated bizarre tumor cells. From the clinical management point of view, it is most important to distinguish low-grade MFS from benign myxoid lesions such as intramuscular myxoma and nodular fasciitis. The presence of moderate nuclear pleomorphism and fragments of curved capillaries in the myxoid matrix are the most important clues to the diagnosis of MFS (see Fig. 13.33) [51–53].


#### *Cytologic features:*

- Dispersed cells and cell clusters
- Abundant myxoid background
- Fragments of curved vessels embedded in the myxoid matrix
- Slight to moderate atypical spindle cells in low-grade neoplasm
- Marked cellular and nuclear pleomorphism in high-grade neoplasm
- Occasional pseudolipoblasts

#### *Differential diagnosis and problems in diagnosis:*

- Intramuscular myxoma, cellular type
- Myxoid liposarcoma
- Nodular fasciitis
- Low-grade fibromyxoid sarcoma
- Spindle-cell lipoma






**Fig. 13.33** Myxofibrosarcoma. (a–c) Loosely cohesive cell clusters of pleomorphic spindle cells and dispersed atypical single cells in a myxoid background. Note fragments of curved vessels (MGG; H&E) (d)

Pseudolipoblasts are common (Pap stain). (e) Low-grade myxofibrosarcoma is sometimes indistinguishable from benign cellular myxoid neoplasms, such as cellular intramuscular myxoma (H&E)





**Fig. 13.34** Dermatofibrosarcoma protuberans (DFSP). (a, b) Compact, often three-dimensional clusters of spindle cells with poorly defined cytoplasmic borders (MGG; H&E). (c, d) Cell-block sections show a

storiform pattern and positive CD34 staining by immunohistochemistry (Cell Block; Shandon kit; H&E; CD34+) (Image courtesy of Elwira Bakula-Zalewska, MD, PhD, Institute of Oncology, Warsaw, Poland)

### 13.6.9 Dermatofibrosarcoma Protuberans

Dermatofibrosarcoma protuberans (DFSP) is a low-grade, locally aggressive fibroblastic neoplasm involving both the dermis and subcutis. Occasionally, higher-grade fibrosarcomatous progression occurs. FNA smears from DFSP are characterized by compact, often three-dimensional clusters of spindle cells. These spindle cells have poorly defined cytoplasmic borders and relatively uniform, spindle-shaped or oval nuclei with finely dispersed chromatin (see Fig. 13.34). Collagen matrix with embedded spindle cells may show myxoid changes. Numerous dissociated spindle cells or naked nuclei are also commonly seen [54–56]. The FNA cytomorphology alone is not sufficiently characteristic to permit a confident diagnosis of DFSP. A definitive diagnosis requires the confirmatory CD34 positivity by immunocytochemistry [56].


#### *Cytologic features:*

- Variable yield
- Tight clusters or fascicles of spindle cells embedded in a collagen matrix
- Dispersed spindle cells and stripped nuclei
- Slight to moderate cellular and nuclear atypia but bland nuclear chromatin and inconspicuous nucleoli
- Pale and poorly defined cytoplasm, better-preserved cells with bipolar cytoplasmic extensions
- Occasionally fat fragments with admixture of spindle cells

#### *Differential diagnosis and problems in diagnosis:*

- Cellular benign fibrous histiocytoma
- Neurofibroma
- Schwannoma
- Nodular fasciitis
- Solitary fibrous tumor
- Other spindle-cell sarcoma arising in cutis–subcutis





**Fig. 13.35** Infantile fibrosarcoma. (a–d) Cellular aspirate with tightly clusters and fascicles of mildly to moderately pleomorphic spindle cells (MGG; H&E)

### 13.6.10 Adult Fibrosarcoma

Adult fibrosarcoma was once considered the most common sarcoma in adults and is now a diagnosis of exclusion. It arises commonly in the deep soft tissue of the extremities, trunk, and head and neck region. Before the era of ancillary techniques, many cases of monophasic synovial sarcoma and malignant peripheral nerve sheath tumor were labeled as fibrosarcoma. The neoplastic cells resemble fibroblasts with no more than a moderate degree of pleomorphism, often arranged in fascicles or a herringbone-like pattern on histology. Dedifferentiated sarcoma and fibrosarcomatous progression from DFSP should be excluded. Adult fibrosarcoma is often diagnosed as a low- or high-grade spindle-cell sarcoma not otherwise specified (NOS) in FNA.

#### *Cytologic features:*

- Spindle cells arranged in thigh clusters, fascicles, or as dispersed cells
- Variable cellular and nuclear atypia
- Stripped nuclei are common findings

#### *Differential diagnosis and problems in diagnosis:*

- Desmoid fibromatosis
- Solitary fibrous tumor
- Synovial sarcoma
- Malignant peripheral nerve sheath tumor
- Nodular fasciitis
- Dedifferentiated liposarcoma

### 13.6.11 Infantile Fibrosarcoma

Infantile fibrosarcoma is commonly seen before the age of 2 years; it occurs as a congenital neoplasm as well. Most tumors arise in the extremities as a large, fast-growing, non-tender mass. The main histologic features are fibroblast-like cells with moderate atypia and often numerous mitoses arranged in tight fascicles. Foci of myxoid stroma, areas of round cells, and a hemangiopericytoma-like vascular pattern may be seen. FNA smears show tight clusters and fascicles of slight to moderate pleomorphic spindle cells (*see* Fig. 13.35).

Spindle-cell population in smears from infantile fibrosarcoma and from fibrous hamartoma of infancy may show similar features. In addition, the mixture of spindle and round cells in aspirate may give a false impression of embryonal rhabdomyosarcoma. It has a distinctive *ETV6-NTRK3* gene fusion and is related to cellular congenital mesoblastic nephroma [57].

*Cytologic features:*

- Cellular aspirate
- Tight clusters and fascicles of slight to moderate pleomorphic spindle cells
- Spindle cells with bland nuclei
- Numerous mitoses

*Differential diagnosis and problems in diagnosis:*


- Fibrous hamartoma of infancy
- Childhood fibromatoses
- Embryonal rhabdomyosarcoma

### 13.6.12 Inflammatory Myofibroblastic Tumor

Inflammatory myofibroblastic tumor (IMT) was once regarded as a reactive process, which was fallen in the spectrum of lesions termed “inflammatory pseudotumor.” Over the past two decades, IMT has been recognized as a rare low-grade neoplasm with distinctive clinical, pathological, and molecular features. IMT shows a predilection for the visceral soft tissue of children and adolescents. The FNA cytologic features of IMT are somewhat nonspecific, variable, and usually impose a benign, reactive impression. Typically, the smears show a mixture of plump myofibroblasts, inflammatory cells (mainly plasma cells and lymphocytes), and occasional large ganglion-like cells (*see* Fig. 13.36a) [58, 59]. Approximately 50 % of IMTs show chromosomal translocations involving *ALK* gene rearrangements resulting in *ALK* protein overexpression (*see* Fig. 13.36b), especially in young patients [60].

*Cytologic features:*

- Hypercellular smears
- Plump myofibroblasts arranged in thigh clusters or dispersed cells
- Inflammatory infiltrate including plasma cells, lymphocytes, and histiocytes
- Occasional ganglion-like cells



**Fig. 13.36** Inflammatory myofibroblastic tumor. (a) Nodular fasciitis-like smears containing clusters of plump myofibroblasts and admixed inflammatory infiltrate rich in plasma cells (Pap stain). *ALK* protein overexpression (b) is seen in about 50 % of cases

*Differential diagnosis and problems in diagnosis:*

- Reactive myofibroblastic proliferation
- Angiomatoid fibrous histiocytoma
- Follicular dendritic cell sarcoma
- Langerhans cell histiocytosis




## 13.7 So-Called Fibrohistiocytic Tumors

### 13.7.1 Tenosynovial Giant Cell Tumor (Localized and Diffuse Types)


Tenosynovial giant-cell tumors are among the most-targeted benign neoplasms by FNAC. The localized and diffuse types of tenosynovial giant-cell tumor share a common pathogenesis and a similar morphology but differ in their growth patterns, clinical features, and biological behaviors. Both can be intra- or extra-articular. The localized type (giant-cell tumor of tendon sheath) is a relatively common

neoplasm of the digits with a female predominance. It is a painless, slowly growing, well-circumscribed tumor, usually smaller than 4 cm in size. The diffuse type (pigmented villonodular tenosynovitis) is a locally aggressive neoplasm affecting mainly the knee and hip in younger patients. It presents as an ill-defined, periarticular mass, often larger than 5 cm. Both neoplasms are composed of mononuclear cells with round to oval nuclei, admixed with osteoclast-like giant cells, xanthoma cells, and siderophages, embedded in a collagenous, occasionally hyalinized matrix. All these components can be seen in the FNA smears (*see* Fig. 13.37) [61–63].



**Fig. 13.37** Tenosynovial giant-cell tumor (localized and diffuse types). (a–c) Dispersed and small clusters of mononuclear cells with round to oval nuclei and mild to moderate cellular and nuclear

pleomorphism with admixture of osteoclast-like giant cells, xanthomatous histiocytes and siderophages (H&E; MGG). (d) Similar morphology in cell-block sections (H&E)



**Fig. 13.38** Undifferentiated sarcoma. (a, b) Polygonal and epithelioid cells with marked cellular and nuclear pleomorphism (MGG; H&E)

*Cytologic features:*

- Variable yield, hypocellular smears in tumors with hyalinized stroma
- Dispersed and clusters of mononuclear cells with round to oval nuclei (sometimes plasma cell-like)
- Mild to moderate cellular and nuclear pleomorphism
- Osteoclast-like giant cells
- Xanthomatous histiocytes and siderophages
- Mitoses may be found

*Differential diagnosis and problems in diagnosis:*

- Deep benign fibrous histiocytoma
- Reactive synovial hyperplasia
- Giant cell-rich sarcoma

### 13.7.2 Undifferentiated Sarcoma

Undifferentiated sarcomas are defined as a heterogeneous group of sarcomas without evidence of line of differentiation by presently available technology in the latest 2013 World Health Organization (WHO) classification [1]. Morphologically, they can be either spindle cell, epithelioid, round cell, or pleomorphic (formerly often known as pleomorphic malignant fibrous histiocytoma [MFH]) (see Fig. 13.38). MFH was for many years considered a distinct entity. Several subtypes had been described and the most common one was the pleomorphic type. Retrospective examinations of series of pleomorphic MFH have identified lipogenic, myogenic,

and Schwann cell differentiation, as well as non-mesenchymal histogenesis in the majority of neoplasms diagnosed initially as MFH. Now, these groups of sarcomas are the diagnoses of exclusions. Because of limited sampling, undifferentiated sarcoma should only be made on FNAC if immunocytochemical studies are possible.

*Cytologic features:*

- Hypercellular smears
- Necrosis and hemorrhage
- Mixture of dispersed cells, cell clusters, and small tissue fragments in variable proportions
- Variable cellular morphology, commonly spindle-, polygonal-, and epithelioid cells with marked cellular and nuclear pleomorphism, coarse chromatin, and large nucleoli
- Uni- and multinucleated giant cells
- Atypical mitoses

*Differential diagnosis and problems in diagnosis:*

- Other pleomorphic sarcomas with a specific line of differentiation (pleomorphic liposarcoma, pleomorphic leiomyosarcoma, pleomorphic MPNST, and pleomorphic rhabdomyosarcoma)
- Dedifferentiated sarcomas
- Radiation-associated pleomorphic sarcoma
- Non-mesenchymal tumors (anaplastic large cell lymphoma, soft tissue metastases of anaplastic carcinoma, and sarcoma-like malignant melanoma)
- Sarcomatoid mesothelioma



## 13.8 Smooth Muscle Tumors

Benign smooth muscle neoplasms such as cutaneous leiomyoma, angioleiomyoma, and uncommon deep leiomyoma of soft tissue are rare targets for FNA. Leiomyosarcoma of soft tissue, one of the most common soft tissue sarcomas, is relatively frequently examined by FNA cytology.

### 13.8.1 Leiomyoma of Deep Soft Tissue

Leiomyoma of deep soft tissue is a rare, slowly growing, well-circumscribed benign neoplasm, commonly arising in the retroperitoneum and different parts of the abdominal cavity in women. The cells have the morphology reminiscence of normal smooth muscle cells (*see* Fig. 13.39).

Low-grade leiomyosarcoma, an important diagnostic pitfall, displays more nuclear pleomorphism and coarser chromatin than leiomyoma [64].

*Cytologic features:*

- A mixture of clusters and dispersed, bland spindle cells
- Elongated, blunt-ended, cigar-shaped, occasionally truncated nuclei with insignificant nucleoli
- Collagenous matrix stains bluish-red (MGG) and often in the cell clusters
- Occasional mild to moderate cellular pleomorphism
- Some cells display grey or blue-grey cytoplasm with cytoplasmic extensions
- No mitotic figures

*Differential diagnosis and problems in diagnosis:*

- Extra-abdominal desmoid fibromatosis
- Low-grade leiomyosarcoma
- Gastrointestinal stromal tumor, spindle-cell type
- Schwannoma


- Low-grade malignant peripheral nerve sheath tumor (MPNST)
- Monophasic synovial sarcoma

### 13.8.2 Angioleiomyoma


Angioleiomyoma (ALM) is a benign neoplasm that usually presents as a tender or painful nodule in the skin or subcutaneous tissue in adults, commonly located in the lower extremities, but it can be widely distributed over the body. Clinically, ALM can be easily confused with glomus tumor, cutaneous cylindroma, and schwannoma. Morphologically, it may overlap with myopericytoma. One series of FNA of ALM has been published to date [65]. The most common findings are variable proportions of benign smooth muscle cells and benign uniform spindle cells, either dissociated or arranged in fascicles. Small fragments of collagenous matrix are also present (*see* Fig. 13.40).

### 13.8.3 Leiomyosarcoma


Leiomyosarcoma (LMS) accounts for 10–15 % of all primary soft tissue sarcomas [8]. Soft tissue LMS occurs most often in the retroperitoneum and limbs but also in other locations such as visceral organs and bones. Superficial LMS most often shows an indolent clinical course, whereas deeply seated soft tissue or retroperitoneal LMS shows an aggressive behavior with a high risk of metastasis. The morphology of LMS is similar in both settings. Typical tumor cells have elongated, cigar-shaped nuclei, sometimes segmented “in tandem” position (*see* Figs. 13.41 and 13.42). In most cases, a characteristic fascicular/pleomorphic pattern is recognizable at low-power examination. A pure spindle-cell or epithelioid pattern is less common (*see* Fig. 13.43). In high-grade



**Fig. 13.39** Leiomyoma of deep soft tissue. (a, b) Mixture of clusters and dispersed bland spindle cells with elongated, blunt-ended, cigar-shaped nuclei (MGG)




**Fig. 13.40** Angioleiomyoma. (a, b) Fascicles of smooth muscle cells and uniform spindle cells in a collagenous matrix (MGG)




**Fig. 13.41** Leiomyosarcoma. (a–d) Small clusters and dissociated tumor cells with elongated, cigar-shaped, occasionally truncated or “in tandem” position nuclei (H&E; MGG)





**Fig. 13.42** Leiomyosarcoma. (a, b) Loosely cohesive clusters with a mixed fascicular/pleomorphic pattern (H&E; MGG)



**Fig. 13.43** Leiomyosarcoma. (a, b) Cohesive fascicles of atypical spindle cells (MGG; H&E). (c) Similar pattern is obvious in the cell-block section. (d) Strong desmin positivity supports diagnosis (H&E; desmin; cell block)

LMS, marked cellular atypia, pleomorphism, necrosis, and multinucleated giant cells (some osteoclast-like giant cells) are common [66, 67].

*Cytologic features:*

- Hypercellular smears (hypocellular in tumors with hyalinized matrix)
- Three major cellular patterns in low power: mixed fascicular/pleomorphic, predominantly fascicular, and predominantly pleomorphic
- Minor component of dispersed, well-preserved cells with dense cytoplasm
- Stripped atypical, degenerate nuclei, with blue- or magenta-colored background in fascicles or clusters (MGG)
- Nuclei with elongated, blunted ends, cigar-shaped, occasionally segmented “in tandem” position
- Epithelioid tumor cells in epithelioid variant
- Pleomorphic, occasionally bizarre, multinucleated cells
- Mitoses and necrosis
- Osteoclasts

*Differential diagnosis and problems in diagnosis:*

- Soft tissue leiomyoma
- Nodular fasciitis
- Schwannoma
- Malignant peripheral nerve sheath tumor (MPNST)
- Myofibroblastic sarcomas
- Undifferentiated pleomorphic/spindle-cell sarcoma
- Dedifferentiated sarcoma
- Metastatic sarcomatoid carcinoma
- Metastatic melanoma

Epithelioid variant leiomyosarcoma may be misdiagnosed as undifferentiated carcinoma. In LMS, at least few cells with cigar-shaped or blunt-ended nuclei are present. Immunocytochemical examinations are helpful in doubtful cases (*see* Fig. 13.43). Leiomyosarcoma stains for smooth muscle actin (SMA), desmin, and caldesmon, although sometimes only focally. Focal cytokeratin positivity has been reported in leiomyosarcoma. The cytomorphology of leiomyosarcoma has been thoroughly evaluated in a few large series [66, 67].



### 13.9 Striated Muscle Tumors

Benign and malignant striated muscle tumors are rare in adults in general. Embryonal and alveolar subtypes of rhabdomyosarcoma (RMS) are among the most common sarcomas of childhood, whereas pleomorphic RMS occurs exclusively in adults.

#### 13.9.1 Adult Rhabdomyoma

Adult rhabdomyoma is a rare benign neoplasm showing mature skeletal muscle differentiation. Most arise in the head and neck region, especially the tongue and floor of mouth. Smears display dispersed cells or cohesive clusters of large, round to polygonal or elongated cells with abundant eosinophilic granular cytoplasm, often peripherally located, uniform round nuclei with large nucleoli (*see* Fig. 13.44). Cytoplasm may be vacuolated due to dissolved glycogen.


#### *Cytologic features:*

- Large rounded, polygonal or elongated cells, dispersed or arranged in cohesive clusters
- Abundant eosinophilic (H&E) granular cytoplasm
- Small, peripherally located nuclei with prominent nucleoli
- Occasional stripped nuclei
- Rarely visible cross-striation


#### *Differential diagnosis and problems in diagnosis:*

- Granular cell tumor
- Hibernoma

FNAC evaluation of rhabdomyoma may be difficult due to the similarity of the tumor cells to normal striated muscle and to other tumors with cells containing abundant granular cytoplasm [68, 69]. Hibernoma cells have also granular and/or vacuolated cytoplasm in smears, but the cells are generally smaller than those in rhabdomyoma. Granular cell tumors have no cytoplasmic vacuoles and diffusely stain with S-100 protein, whereas desmin and myoglobin expression is typical for rhabdomyoma.



**Fig. 13.44** Adult rhabdomyoma. (a, b) Large, rounded, polygonal, or elongated cells with small round to oval nuclei, fine chromatin, and abundant granular cytoplasm. (c) Rarely visible cross-striation (MGG)



**Fig. 13.45** Embryonal rhabdomyosarcoma. (a–d) Hypercellular smears with loosely cohesive clusters and dispersed single cells of various morphology (triangular, rounded, tadpole- or strap-shaped) (H&E; MGG). Mitoses and multinucleation are common

### 13.9.2 Embryonal Rhabdomyosarcoma

Embryonal rhabdomyosarcoma (RMS) is the most common soft tissue sarcoma in children and accounts for approximately 50 % of diagnosed rhabdomyosarcomas. Common locations include the head and neck region, genitourinary tract, and retroperitoneum. Typically, the smears are variably cellular, with marked cellular pleomorphism. Primitive spindle cells and rhabdomyoblast-like cells of various morphology (triangular, rounded, and tadpole- or strap-shaped) are commonly seen (see Fig. 13.45) [70, 71]. Tumor cells express desmin, myogenin, and MyoD1. Embryonal RMS has a better prognosis than alveolar RMS.

#### *Cytologic features:*

- Mixture of loosely cohesive clusters and dispersed cells
- Spindle cells or round cells are most common
- Cellular and nuclear pleomorphism
- Nucleoli may be prominent
- Rhabdomyoblast-like tumor cells (triangular, rounded, and tadpole- or strap-shaped) variably present

- Rhabdomyoblasts with eosinophilic (H&E) or grey-blue (MGG) dense cytoplasm
- Areas of myxoid matrix variably present

#### *Differential diagnosis and problems in diagnosis:*


- Alveolar rhabdomyosarcoma
- Infantile fibrosarcoma
- Leiomyosarcoma (LMS)

LMS is extremely rare in children and does not express myogenin or MyoD1. Spindle-cell neoplasm such as infantile fibrosarcoma is excluded when tumor cells are positive for desmin, myogenin, or MyoD1.

### 13.9.3 Alveolar Rhabdomyosarcoma

Alveolar RMS (ARMS) accounts for about 20 % of all pediatric RMSs and occurs in older children and adolescents. In contrast to embryonal RMS, the cellular composition of smears is more uniform with predominantly small- to medium-sized round or ovoid cells with round or pear-shaped hyperchromatic





**Fig. 13.46** Alveolar rhabdomyosarcoma. (a, b) Loosely cohesive clusters and dispersed, single, small- and medium-sized cells with hyperchromatic nuclei and prominent nucleoli (H&E). (c) Rhabdomyoblastic

morphology (MGG). (d, e) Diffuse and strong desmin and myogenin positivities support with diagnosis (desmin; myogenin)

nuclei, often with large nucleoli (*see* Fig. 13.46). Multinucleated tumor giant cells with small nuclei are common [71]. Similar to the embryonal subtype, tumor cells are immunoreactive with desmin, myogenin, and MyoD1. Aberrant expression of keratins, CD99, S-100 protein, lymphoid markers, and NSE poses

significant diagnostic challenges and confusion with other small round blue cell tumors. Most ARMS have either *PAX3-FOXO1* or *PAX7-FOXO1* gene fusion, which can be detected by FISH [15]. Identification of ARMS is clinically important as it has unfavorable prognosis and is treated more aggressively.

*Cytologic features:*

- Hypercellular smears
- Cell clusters mixed with dispersed cells
- Stripped nuclei commonly mixed with grey-blue blebs of cytoplasm
- Small- or medium-sized, round to polygonal, and ovoid or pear-shaped cells
- Scanty cytoplasm, coarse nuclear chromatin, and often large nucleoli
- Cytoplasmic vacuolization
- Mitoses
- Binucleated and multinucleated tumor cells

*Differential diagnosis and problems in diagnosis:*

- Ewing sarcoma
- Poorly differentiated synovial sarcoma
- Precursor lymphoblastic lymphoma
- Poorly differentiated neuroendocrine carcinoma

Alveolar RMS belongs to a group of round cell malignant neoplasms and distinguishing ARMS from other small round cell tumors can be difficult in routinely stained smears. The final diagnosis should be confirmed with ancillary tests such as immunocytochemical and molecular genetic studies (Table 13.1 and 13.2).

**13.9.4 Pleomorphic Rhabdomyosarcoma**

Pleomorphic RMS is an uncommon sarcoma of adults older than 50 years of age. Common sites are the limbs. This is a high-grade sarcoma with marked cellular and nuclear pleomorphism. The FNA smears are hypercellular, containing highly atypical spindle or polygonal cells, and large rhabdoid cells with abundant eosinophilic cytoplasm (H&E) and eccentrically located nuclei [72, 73]. These rhabdoid cells are similar to those seen as a heterologous component in other sarcomas such as MPNST. Tumor cells stain for desmin and myogenin.

*Cytologic features:*

- Cell clusters and dispersed cells
- Marked cellular and nuclear pleomorphism
- Mixture of atypical spindle cells, rhabdoid cells, and multinucleated tumor cells

*Differential diagnosis and problems in diagnosis:*

- Pleomorphic variant of other sarcomas
- Malignant peripheral nerve sheath tumor with heterologous rhabdoid differentiation
- Dedifferentiated liposarcoma with heterologous rhabdoid differentiation



## 13.10 Tumors of Peripheral Nerves

### 13.10.1 Benign Nerve Sheath Tumors

#### 13.10.1.1 Schwannoma (Neurilemmoma)

Schwannoma occurs in all age groups but most frequently in adults between 40 and 60 years old. Most schwannomas are deep seated, arising in the limbs, head and neck region, retroperitoneum, and posterior mediastinum.

Schwannoma is a neoplasm commonly examined by FNAC. Needling the lesion itself may produce a sharp pain radiating along the nerve, a valuable clue to the diagnosis; however, it is not entirely specific and can occur in any soft tissue lesion seated close to a nerve. Smears are variably but often hypercellular and contain irregular tissue fragments of spindle cells embedded in a fibrillary stroma. It is often possible to identify both the highly cellular Antoni A and the poorly cellular myxoid Antoni B components (*see* Fig. 13.47). Most spindle cells exhibit neurogenic differentiation including elongated, wavy nuclei that are often pointed or folded and sometimes show a fishhook-like appearance. Many schwannomas show cystic changes resulting in poor cellularity or nondiagnostic yield [74, 75].

*Cytologic features of Antoni A-type schwannoma:*

- Cohesive tissue fragments of different sizes, often with irregular borders (“of a jigsaw puzzle pieces”)
- Rare dispersed cells
- In the fragments, cells have indistinct cytoplasm and elongated, slender nuclei with pointed ends, boomerang-shaped, or comma-like nuclei embedded in a fibrillary stroma
- Occasional nuclear palisading and Verocay bodies
- Occasionally small round nuclei
- Occasionally slight to moderate nuclear pleomorphism

*Cytologic features of Antoni B-type schwannoma:*

- Poor cellular smears with variable areas of myxoid matrix
- Dispersed cells more common than in smears from Antoni A areas
- Cystic degeneration with presence of histiocytes and inflammatory cells

*Differential diagnosis and problems in diagnosis:*

- Spindle-cell lipoma
- Solitary fibrous tumor
- Low-grade MPNST
- Low-grade LMS
- Meningioma

- Myxoid soft tissue tumors (Antoni B-type neurilemmomas with large foci of myxoid matrix)
- Pleomorphic sarcomas (smears from ancient schwannomas)

A common problem is to distinguish sarcomas from ancient schwannomas, which often exhibit nuclear pleomorphism with marked anisokaryosis and hyperchromasia. Many of those large nuclei, however, show evenly distributed chromatin and typical large intranuclear vacuoles (“Kern-loche”) (*see* Fig. 13.48). It is usually possible to find other components of a schwannoma in addition to the scattered atypical cells (*see* Fig. 13.42) [76]. Another diagnostic pitfall is misinterpreting cellular schwannoma as a low-grade spindle-cell sarcoma [77]. Demonstration of diffuse and strong positivity for S-100 protein by immunocytochemistry is very helpful to confirm the diagnosis of schwannoma.

#### 13.10.1.2 Neurofibroma


Neurofibromas, especially large neurofibromas in patients with von Recklinghausen’s disease, are occasionally referred for FNA when there is a question of malignant transformation. Neurofibroma is composed of a mixture of Schwann cells, perineural-like cells, fibroblasts, mast cells, and scattered axons all embedded in a stroma. Because the stroma is typically fibromyxoid, at times collagenous or hyalinized, FNA of these stroma-rich areas can result in poor or hypocellular smears. The aspirated cells are similar to those in smears from schwannoma: cohesive spindle cells with slender and often comma-like, bent, or wavy nuclei. Different cell types in neurofibroma are indistinguishable in smears. A specific diagnosis of neurofibroma based on FNA alone can be difficult. Immunocytochemical studies are often necessary.

*Cytologic features:*

- Variable cellularity
- Occasionally myxoid matrix
- Small, poorly cohesive cell clusters and dispersed cells
- Bland fibroblast-like spindle cells and spindle cells reminiscent of those in schwannoma
- Stripped nuclei

*Differential diagnosis and problems in diagnosis:*


- Antoni B-type schwannoma
- Intramuscular myxoma
- Solitary fibrous tumor
- Spindle-cell lipoma
- Low-grade malignant peripheral nerve sheath tumor



**Fig. 13.47** Schwannoma. (a–d) Cohesive tissue fragments of different sizes with spindle cells showing pointed ends embedded in a fibrillary stroma. Occasional nuclear palisading (c) and round cell component

(d) (H&E; MGG). Verocay body better appreciated in the cell-block section (e) (H&E)





**Fig. 13.48** Ancient schwannoma. (a–d) Irregular clusters and some dispersed single cells with marked anisokaryosis and hyperchromasia. Note typical intranuclear inclusions (“Kern-loche”) (H&E; MGG)

### 13.10.1.3 Granular Cell Tumor (Sect. 14.9)

Granular cell tumor is an uncommon benign neoplasm commonly arising in the tongue, gastrointestinal tract, mediastinum, skin and subcutaneous tissue, breast, and other sites in the middle-aged adults.

The tumor cells have round to oval nuclei, inconspicuous nucleoli, and abundant granular cytoplasm. Because of the fragility of tumor cells, smears often contain stripped nuclei in a background of finely granular material (see Figs. 14.32 and 14.33). Nuclei are small and dark but occasionally with mild to moderate atypia. The tumor cells stain for S-100 protein (see Fig. 14.33), NSE, CD68, and inhibin contain PAS-positive diastase-resistant cytoplasmic granules representing the phagolysosomes. Malignant granular cell tumors are


very rare and smears show morphologic criteria of malignancy such as predominance of pleomorphic spindle cells with large nucleoli, mitoses, and necrosis [78].

#### *Cytologic features:*

- Dispersed cells and poorly cohesive clusters
- Intact polygonal cells with abundant granular cytoplasm
- Stripped nuclei in a granular background
- Rounded, small nuclei with insignificant nucleoli
- Cells with large nuclei with coarse chromatin and macronucleoli may be present

#### *Differential diagnosis and problems in diagnosis:*

- Hibernoma
- Adult rhabdomyoma
- Alveolar soft part sarcoma



**Fig. 13.49** Perineurioma. (a–c) Poorly preserved, elongated spindle cells with bland nuclei in the myxoid background (H&E; MGG)

#### 13.10.1.4 Perineurioma

Soft tissue perineurioma is a rare benign peripheral nerve sheath neoplasm composed of cells resembling the normal perineural cells. A specific diagnosis is most likely not feasible based on cytomorphology alone because the FNA smears share a similar appearance with other low-grade spindle-cell and myxoid neoplasms, such as cellular intramuscular myxoma. Nevertheless, the presence of elongated spindle cells with bland oval nuclei and characteristic long, thin, bipolar cytoplasmic processes in a myxoid stroma is a typical finding (see Fig. 13.49) [79, 80]. The perineurioma cells are positive for CD34, epithelial membrane antigen (EMA), and claudin-1 (as normal perineural cells) and negative for S-100 protein.

*Cytologic features:*

- Variable cellularity
- Often myxoid background
- Elongated cells with oval nuclei and long, thin, bipolar cytoplasmic processes
- Few vessel fragments


*Differential diagnosis and problems in diagnosis:*

- Cellular intramuscular myxoma
- Schwannoma with prominent Antoni B area
- Neurofibroma
- Myxofibrosarcoma, low grade
- Low-grade fibromyxoid sarcoma

#### 13.10.2 Malignant Peripheral Nerve Sheath Tumor

Malignant peripheral nerve sheath tumor (MPNST) is a sarcoma arising from a peripheral nerve or as a malignant transformation of neurofibromas in patients with von Recklinghausen's disease (NF1). Common sites of involvement include large- and medium-sized nerves of the upper arm, buttock, brachial plexus, and paraspinal nerves. The cytomorphologic features, reflecting very diverse histologic appearances, are extremely variable and nonspecific. The most commonly described tumor cells are elongated with





**Fig. 13.50** Malignant peripheral nerve sheath tumor. (a–c) Clusters of spindle cells with hyperchromatic nuclei in fascicles (a) and in loosely clusters (b, c) (H&E; MGG). Without knowledge of clinical history and

ancillary immunostains with S-100 protein and GFAP, correct diagnosis of MPNST is difficult to render based on cytomorphology alone

fusiform, often comma-shaped, wavy, or buckled nuclei, but other forms of neoplastic cells are also present. Pleomorphic and multinucleated tumor cells are found in high-grade tumors. Polygonal or rounded cells may predominate in epithelioid MPNST. Common helpful histologic features such as whorled perivascular arrangement of cells, fascicles, or a storiform pattern (*see* Fig. 13.50), and alternative cellular and hypocellular areas are not perceptible in FNA smears. Heterologous components (cartilage, bone, epithelial glands, or rhabdoid cells) present focally in approximately 10–15 % of MPNST are rarely sampled by FNA [81–83]. Although focal S-100 protein and/or GFAP positivity is helpful; a panel of antibodies (*e.g.*, CD34, cytokeratins, EMA, desmin, MDM2) should be used to exclude its many morphologic mimics.

*Cytologic features:*

- Hypercellular smears
- Mixture of cell clusters, fascicles, and dispersed cells
- A fibrillar background may be found in clusters or fascicles
- Predominantly spindle-shaped cells often with bipolar cytoplasmic processes and fusiform, wavy, or comma-like nuclei
- Often hyperchromatic nuclei with large nucleoli
- Variable presence of pleomorphic and/or multinucleated tumor cells
- Infrequent heterologous components
- Rounded or polygonal cells predominate smears of epithelioid MPNST
- Mitotic figures

*Differential diagnosis and problems in diagnosis:*

- Ancient schwannoma
- Cellular schwannoma
- Leiomyosarcoma
- Synovial sarcoma
- Sarcomatoid carcinoma
- Spindle-cell melanoma
- Malignant solitary fibrous tumor
- Dedifferentiated liposarcoma
- Myoepithelial carcinoma (epithelioid MPNST)

## 13.11 Vascular Neoplasm

Vascular neoplasms comprise a wide spectrum of entities, including benign lesions such as hemangioma; low-grade malignancy such as epithelioid hemangioendothelioma (EHE) and high-grade sarcomas such as epithelioid angiosarcoma. Benign entities such as subcutaneous and intramuscular hemangioma are those most often referred for FNA. Cytologic features are usually nonspecific; a diagnosis of benign vascular tumor must be supplemented by clinical and radiographic findings. The cytologic features of angiosarcoma and EHE have been reported in some series and case reports [84–86].

### 13.11.1 Hemangioma (Sect. 14.7)

*Cytologic features* (Figs. 14.25, 14.26, and 14.27):

- Often, when a needle reaches the tumor, the syringe is filled with blood before aspiration
- Cell poor, bloody smears
- Small sheets and clusters of uniform spindle cells with thin cytoplasmic processes
- Fusiform nuclei with pointed ends
- Variable presence of fragments of small capillaries, occasionally with metachromatic matrix
- Histiocytes in variable numbers, some with hemosiderin pigment
- Occasionally adipose tissue fragments, at times with fragments of vessels

*Differential diagnosis and problems in diagnosis:*

- Benign spindle-cell neoplasms
- Angiomatoid fibrous histiocytoma
- Angiosarcoma
- Benign lipomatous tumors

A specific diagnosis of hemangioma may be made with confirmatory endothelial markers such as CD31 (*see* Fig. 14.26), factor VIII, and Fli-1.

### 13.11.2 Angiosarcoma (Sect. 14.8)

The vast majority of angiosarcomas are cutaneous tumors, and approximately 20 % arise in the soft tissue. Smears of angiosarcomas are most often diagnosed as malignant. A correct diagnosis of angiosarcoma is difficult to reach in routinely stained smears. Immunocytochemical studies with endothelial markers are necessary to establish the diagnosis (*see* Fig. 14.30). An exception is for those cases in which vasoformative structures such as small acinar-like formations with central erythrocytes or vacuolated cells with single erythrocytes are present.

*Cytologic features:*

- Variable yield
  - Often hemorrhagic aspirates
  - Mixture of dispersed cells, cells in clusters or groups
  - Occasionally acinar structures and intracytoplasmic erythrocytes
  - Variable cellular and nuclear pleomorphism
  - Variable presence pleomorphic/polygonal cells and spindle cells
  - Signet ring-like cells often with single erythrocytes within cytoplasmic vacuoles
  - Epithelial-like cells in epithelioid angiosarcoma
- Differential diagnosis and problems in diagnosis:*
- Spindle-cell sarcoma of various lines of differentiation
  - Pleomorphic sarcoma of various lines of differentiation
  - Epithelioid sarcoma of various lines of differentiation
  - Carcinoma metastasis
  - Malignant melanoma

Tumor cells in epithelioid angiosarcoma express cytokeratins in approximately 50 % of cases, which can lead to the misinterpretation of metastatic carcinoma.



## 13.12 Neuroectodermal Tumors

### 13.12.1 Neuroblastoma

Neuroblastoma is the most common extracranial solid malignant tumor in children. Ninety percent of neuroblastomas are diagnosed before the age of 5 years. As neuroblastoma and ganglioneuroblastoma originate in sympathetic ganglia, neuroblastomas may arise in the adrenal medulla or from any sympathetic ganglia within the retroperitoneum, posterior mediastinum, neck, and sacral region.

Neuroblastomas are composed of clusters of small tumor cells with rounded or irregular nuclei showing finely granular chromatin and small nucleoli. Variable amount of intercellular fibrillary material from neuritic cell processes (neuropil) is present. A typical microscopic finding is the rosette-like structures with a central tangle of fibrillary material (Homer-Wright


rosettes). In more mature tumors, large ganglion-like cells are present and in undifferentiated neuroblastoma there is no fibrillary matrix. Criteria for cytologic diagnosis are described in detail in Sect. 3.1 on pediatric tumors.

### 13.12.2 Ganglioneuroblastoma

Typical features in smears are large ganglion cell-like cells mixed with the small cell population.

### 13.12.3 Ganglioneuroma

In ganglioneuroma smears, both large ganglion cells and schwannoma-like tissue fragments are present (*see* Fig. 13.51) [87].




**Fig. 13.51** Ganglioneuroma. (a) Double cell population of large ganglion cells and schwannoma-like tissue fragments (b) a cluster rich in ganglion cells (MGG)

### 13.12.4 Ewing Sarcoma

Extrasosseous Ewing sarcoma (ES) is a primitive round cell sarcoma showing varying degrees of neuroectodermal differentiation. It mainly occurs in adolescents and young adults but affects elderly patients as well. It predominantly arises at

extraskelatal sites such as deep soft tissue of chest wall (the so-called Askin tumor), subcutis, and extremities, but it can also occur in lung, kidney, and genital organs. In our files, there is a patient with a tumor in the myocardium.

Cytologic features are similar to conventional ES in bone (*see* Chaps. 15 and 16) (Fig. 13.52).



**Fig. 13.52** Ewing sarcoma. (a) Dual appearances: small dark and large light cells and tigroid background (b) obvious in air-dried smears (MGG). Irregular cluster of loosely cohesive cells (c) and rosette formation (d) are less common (H&E)



### 13.13 Tumors of Unknown, Uncertain, or Debated Histogenesis

#### 13.13.1 Intramuscular Myxoma


Intramuscular myxoma is a benign tumor typically found in middle-aged or elderly patients. Predilection sites are thigh, buttock, and the shoulder region. As they are deep seated and rather firm at palpation, they might be perceived as malignant clinically. The smears are extremely hypocellular and contain abundant myxoid background and few bland, spindle to stellate cells with long cytoplasmic processes (*see* Fig. 13.53). Fragments of capillaries are rare or absent. Scattered macrophages with cytoplasmic vacuolation are often seen.

#### Cytologic features:

- Droplets of stringy, glue-like colorless fluid at aspiration
- Abundant myxoid matrix
- Dispersed cells more common than small tissue fragments
- Slender tumor cells with small, bland nuclei and (very) long, thin, uni- or bipolar cytoplasmic processes
- Scattered macrophage-like cells with vacuolated cytoplasm
- Infrequent small vessel fragments.


#### Differential diagnosis and problems in diagnosis:

- Schwannoma with foci of myxoid stroma
- Ganglion
- Myxoid liposarcoma
- Low-grade myxofibrosarcoma
- Low-grade fibromyxoid sarcoma
- Extraskeletal myxoid chondrosarcoma



**Fig. 13.53** Intramuscular myxoma. (a–c) Hypocellular smears with abundant myxoid background and small dyscohesive clusters of bland, spindle to stellate cells with long cytoplasmic processes (MGG;

H&E) (d) Cell-block section showing similar morphology of bland spindle cells in a myxoid matrix (H&E)



**Fig. 13.54** Angiomatoid fibrous histiocytoma (AFH). (a–d) Cellular clusters of oval to spindled histiocytoid cells with ill-defined cell borders, ovoid to folded nuclei, finely dispersed chromatin, and inconspicuous

nucleoli. In large cellular clusters, capillary structures with spindled endothelial cells are seen (a). Cytologic atypia and pleomorphism (b, d) may be present, without correlation of malignancy (H&E; MGG)

### 13.13.2 Angiomatoid Fibrous Histiocytoma

Angiomatoid fibrous histiocytoma (AFH) is an indolent neoplasm commonly arising in the subcutaneous tissue of the extremities in children and young adults. AFH may present as a partly cystic mass with hemorrhage. FNA smears show variable cellularity and contain ovoid to spindled histiocytoid cells that may be isolated or in clusters. Some of these cells are atypical and others contain hemosiderin. Large cellular clusters with a capillary structure and a whorled arrangement of tumor cells can be appreciated in some cases (see Fig. 13.54) [88]. FNA cytomorphology is nonspecific. Clinical correlation and ancillary studies are necessary to render the diagnosis. Tumor cells are positive for desmin and EMA in approximately 50 % cases. Most of these tumors show an *EWSR1–CREB1* fusion gene, which can be detected by FISH studies (Table 13.2).


#### *Cytologic features:*

- Variable cellularity
- Dispersed cells and cell clusters
- Ovoid to spindled histiocytoid cells
- Bloody background and hemosiderin
- Infrequent lymphoplasmacytic infiltrate
- Occasional whorled arrangement of tumor cells associated with vessels

#### *Differential diagnosis and problems in diagnosis:*

- Hemangioma
- Benign fibrous histiocytoma
- Nodular fasciitis
- Inflammatory myofibroblastic tumor
- Follicular dendritic sarcoma
- Metastatic carcinoma





**Fig. 13.55** Ossifying fibromyxoid tumor. (a–d) Dispersed, bland, round (epithelioid-like), or ovoid cells with mostly poorly preserved cytoplasm, associated with a variable myxoid matrix (a–c, MGG; d, H&E)

### 13.13.3 Ossifying Fibromyxoid Tumor

Ossifying fibromyxoid tumor (OFMT) is a rare subcutaneous tumor, predominantly in adults. Extremities are the most common sites but OFMT in the trunk and head and neck region has been described. It is a lobulated, well-circumscribed lesion with fibrous capsule often containing a more or less complete shell of mature metaplastic bone. The uniform, oval to spindled neoplastic cells form cords or trabeculae that are situated in fibromyxoid stroma on histology and present as dispersed single cells, clusters of cells, and/or acinar-like structures on FNA smears (see Fig. 13.55) [89, 90]. OFMT was considered as a benign neoplasm until 1995 when Kilpatrick et al. [91] reported six malignant OFMT.

#### *Cytologic features:*


- Variable cellularity (tumors with an extensive shell of bone difficult to aspirate)
- Dispersed cells, cell clusters, and acinar-like structures in a variable myxoid matrix

- Rounded (epithelioid-like) or ovoid nuclei with central nucleoli and slightly anisokaryosis
- Rather abundant cytoplasm

#### *Differential diagnosis and problems in diagnosis:*

- Epithelioid peripheral nerve sheath tumors
- Mixed tumor of soft tissue
- Epithelioid smooth muscle tumors
- Thyroid neoplasms (head and neck tumors)
- Extraskeletal myxoid chondrosarcoma
- Myxoid liposarcoma

The cytology of OFMT in FNA has not been sufficiently investigated, except for a few case reports [89, 90, 92]. When the FNA material is sufficient, IHC is of diagnostic value; positivity for S-100 protein (>90 % cases) and desmin (40–50 % cases) would be supportive for OFMT. Recently, *PHF1* gene rearrangement has been found in typical or atypical OFMT but only rarely in malignant ones [93].



**Fig. 13.56** Mixed tumor/myoepithelioma of soft tissue. (a, b) Epithelial- and myoepithelial-like cells in a chondromyxoid stroma resembling pleomorphic adenoma of salivary gland (MGG)


### 13.13.4 Mixed Tumor/Myoepithelioma of Soft Tissue

Myoepithelial tumors of soft tissue are morphologically and immunophenotypically similar to their counterparts in the salivary gland. They arise commonly in the subcutaneous tissue in adults. These neoplasms are composed of epithelial- and myoepithelial-like cells in a chondromyxoid or hyalinized stroma. In FNA smears, their cytomorphology resembles that of pleomorphic adenoma of salivary gland and chondroid syringoma (see Fig. 13.56). Myoepithelial carcinomas show undifferentiated, large, round cell morphology with a high mitotic rate and necrosis. Interestingly, *EWSR1* gene rearrangement has been detected in these tumors [94]. Most importantly, differential diagnoses are chondroid syringoma and extraskeletal myxoid chondrosarcoma.

### 13.13.5 Synovial Sarcoma

Synovial sarcoma (SS) accounts for 5–10 % of soft tissue sarcomas. It may occur at any age, but more than half of the patients are between the ages of 10 and 35 years. Most SS are deep seated and arise in the extremities, trunk, and head and neck region, but they may arise elsewhere in the body. SS is monophasic (see Fig. 13.57), biphasic (see Fig. 13.58), or poorly differentiated. Three morphologic variants of the poorly differentiated SS have been described; the small cell variant resembling Ewing sarcoma (see Fig. 13.58), a spindle-cell variant resembling MPNST and epithelioid variant with rhabdoid features [95–98]. The majority of SSs stain positively for EMA and cytokeratins 7 and 19. More than 50 % stain positively for CD99 and Bcl-2. These stainings may appear focally, especially in the poorly differentiated





**Fig. 13.57** Synovial sarcoma. (a–c) Tissue fragments mixed with dispersed uniform spindle cells in almost equal proportions. Note nuclear hyperchromasia and mitosis (H&E). (d) Cell-block section with hyperchromatic spindle cells in a collagenous stroma (H&E)

areas. A translocation  $t(X;18)(p11;q11)$  is present in most SSs, resulting in *SS18-SSX* gene fusion, which can be detected by FISH (Table 13.2).

*Cytologic features:*

- Often hypercellular yield
- Tissue fragments mixed with dispersed cells in almost equal proportions
- Stripped nuclei
- Branching capillary strands bordered by tumor cells
- Spindle cells with bland fusiform or ovoid nuclei and inconspicuous nucleoli
- Mitotic figures
- Small acinar-like and alveolar structures in biphasic tumors


- Often scattered mast cells

*Differential diagnosis and problems in diagnosis:*

- Solitary fibrous tumor (monophasic SS)
- Thymoma (biphasic SS)
- Carcinosarcoma (biphasic SS)
- MPNST (poorly differentiated or monophasic SS)
- Ewing sarcoma (poorly differentiated SS)


A cellular aspirate containing bland spindle cells with mitotic figures and scattered mast cells from a deep-seated tumor in an adolescent or middle-aged adult is suggestive of monophasic fibrous synovial sarcoma. However, a definitive diagnosis of a synovial sarcoma should be confirmed by ancillary techniques such as immunocytochemical and molecular genetics examinations.





**Fig. 13.58** Synovial sarcoma. (a, b) Biphasic subtype showing small acinar-like and alveolar structures (MGG). (c, d) Poorly differentiated synovial sarcoma resembling Ewing sarcoma (H&E)





**Fig. 13.59** Alveolar soft part sarcoma. (a, b) Large, dyshesive tumor cells with round nuclei and abundant granular, fragile cytoplasm. (c, d) Presence of stripped nuclei with prominent, centrally situated nucleoli

in a cytoplasmic granular background is a useful clue to the diagnosis (MGG; H&E)

### 13.13.6 Alveolar Soft Part Sarcoma

Alveolar soft part sarcoma (ASPS) is a rare soft tissue sarcoma accounting for less than 1 % of all sarcomas and mainly seen in adolescents and young adults with a slight female predominance. Common sites are deep soft tissue of the lower limbs or limb girdles, and head and neck region. The cytomorphic features of ASPS include large, dyshesive tumor cells with round nuclei and prominent centrally situated nucleoli. Tumor cells have abundant granular, fragile cytoplasm. Presence of stripped nuclei in a cytoplasmic granular background is a useful clue to the diagnosis. The typical alveolar architecture on histology is imperceptible on FNA smears [99, 100]. Intracytoplasmic glycogen may be seen in cell-block sections. The most characteristic rhomboid crystals containing actin


filaments may be evident by ultrastructural examination (*see* Fig. 13.59). ASPS is characterized by an *ASPSCR1-TFE3* gene fusion and nuclear overexpression of TFE3 protein.

#### *Cytologic features:*

- Variable cellularity
- Large, dyshesive epithelioid cells with abundant granular cytoplasm
- Round nuclei with very prominent nucleoli
- Bare nuclei and granular cytoplasmic background

#### *Differential diagnosis and problems in diagnosis:*

- Granular cell tumor
- Metastatic renal cell carcinoma
- Adult rhabdomyoma
- Paraganglioma
- PEComa



**Fig. 13.60** Epithelioid sarcoma. (a, b) Dispersed, round, or polygonal cells with an eccentrically placed nucleus and dense cytoplasm (a, MGG; b, Pap stain). (c) Loss of nuclear expression of INI1 is seen in both conventional and proximal-type epithelioid sarcoma

### 13.13.7 Epithelioid Sarcoma

Epithelioid sarcoma (ES) is a rare malignant mesenchymal neoplasm showing epithelioid morphology and phenotype. It arises in the skin and subcutaneous tissue or tendon aponeurosis of distal extremities, mainly hands and wrists of young adults and adolescents. There are two subtypes: the conventional/distal form and proximal type, which mainly occurs in proximal/truncal regions. FNA smears show a mixture of spindle cells and epithelioid cells, with eccentrically placed nuclei and eosinophilic to dense cytoplasm in the H&E-stained smears (see Fig. 13.60a). It may be difficult to aspirate cellular material likely due to necrosis and/or hyalinization. ES tumor cells express cytokeratins, especially keratins 8 and 19 and/or EMA, and more than 50 % of cases express CD34. Loss of nuclear expression of

INI1 is characteristic for both types of ES (see Fig. 13.60b) [101, 102].


#### *Cytologic features:*

- Variable cellularity
- Dyshesive and small clusters of spindled to epithelioid cells
- Round to oval nuclei with vesicular chromatin and prominent nuclei
- Dense to eosinophilic cytoplasm
- Necrotic background

#### *Differential diagnosis and problems in diagnosis:*

- Metastatic carcinoma
- Metastatic melanoma
- Epithelioid hemangioendothelioma
- Clear cell sarcoma
- Granulomatous inflammation





**Fig. 13.61** Clear cell sarcoma (malignant melanoma of soft parts). (a, b) Polygonal or spindly shaped cells with rather abundant, clear cytoplasm and large, round, or ovoid nuclei. (c) Binucleation is common (MGG)

### 13.13.8 Clear Cell Sarcoma (Malignant Melanoma of Soft Parts)

Clear cell sarcoma (CCS), first described by Enzinger in 1965, is a rare, distinct entity with melanocytic differentiation. It predominantly arises in the deep soft tissue of the extremities in young adults, and it is often found in close proximity to aponeurotic structures and tendons. FNA smears are at least moderately cellular, containing mostly dispersed cells. Small clusters of loosely cohesive cells can also be seen. Tumor cells are polygonal or spindly shaped with rather abundant clear or pale cytoplasm and large round or ovoid nuclei and prominent nucleoli (see Fig. 13.61). Intranuclear cytoplasmic pseudoinclusions, melanin, multinucleation, and tigroid background can also present [103, 104]. Similarly to malignant melanoma, CCS tumor


cells are diffusely and strongly positive for S-100 protein and HMB-45. Metastatic malignant melanoma is the most important differential diagnosis. CCS harbors the characteristic translocation  $t(12;22)(q13;q12)$  with *EWSRI-ATF1* fusion, which is not present in malignant melanoma.

#### *Cytologic features:*

- Moderately cellular smears with tigroid background
- Dispersed cells and occasional loose clusters
- Polygonal or spindle cells with abundant clear to pale cytoplasm
- Round to ovoid nuclei with vesicular chromatin and macronucleoli

#### *Differential diagnosis and problems in diagnosis:*

- Metastasis melanoma
- Metastatic carcinoma with clear cell features
- Alveolar soft part sarcoma



**Fig. 13.62** Extraskeletal myxoid chondrosarcoma. (a–c) Dispersed cells, branching cords, strands, and cell balls of spindly shaped, fusiform, or round cells with ovoid or rounded nuclei in the myxoid

background matrix (H&E; MGG). (d) The typical histologic appearance of branching cords and strands of bland spindle cells is best appreciated in cell-block sections (H&E; Shandon kit)

### 13.13.9 Extraskeletal Myxoid Chondrosarcoma

Extraskeletal myxoid chondrosarcoma (EMC) is a malignant mesenchymal tumor of unknown differentiation in adults, despite bearing “chondrosarcoma” in the name. Most EMCs arise in the deep soft tissue of the trunk and the limbs, with the thigh being the most common site. FNA smears show uniform, fusiform to spindle or round epithelioid tumor cells arranged as cords, strands, or balls in a myxoid and fibrillary matrix. Chondroblasts-like lacunar structures mimicking real chondrosarcoma are common findings (see Fig. 13.62) [105]. The immunophenotype is not specific, up to 20 % stain for S-100 protein and about 30 % for CD117 (KIT) and occasionally for neuroendocrine markers. A characteristic translocation  $t(9;22)(q22/q3;q12)$ , resulting in *EWSR1-NR4A3* fusion gene, is present in most cases. In about 25 %

cases, a translocation  $t(9;17)(q22;q11)$  is identified, which is probably connected with neuroendocrine differentiation [106].


#### *Cytologic features:*

- Myxoid background matrix
- Dispersed cells, branching cords, strands, and cell balls
- Spindly shaped, fusiform or round cells with ovoid or rounded nuclei
- Bland nuclear chromatin and small nucleoli
- Chondroblasts-like lacunar structures
- Lack of significant vascularity

#### *Differential diagnosis and problems in diagnosis:*

- Low-grade myxofibrosarcoma
- Low-grade fibromyxoid sarcoma
- Mixed tumor/myoepithelioma of soft tissue
- Myxoid/round cell liposarcoma
- Chondroid lipoma





**Fig. 13.63** Desmoplastic, small round cell tumor. (a) Loosely cohesive small clusters of ovoid undifferentiated cells with slightly irregular nuclear contours, finely granular chromatin, and scant cytoplasm (Pap stain). (b) Small nests of tumor cells with associated desmoplastic

stroma is characteristic (Cell Block; H&E). Demonstration of polyphenotypic differentiation with positive immunoreactivity for cytokeratins, EMA, desmin (c), and neuroendocrine markers is diagnostically helpful. Note the perinuclear dot-like staining pattern for desmin


### 13.13.10 Desmoplastic Small Round Cell Tumor

Desmoplastic small round cell tumor (DSRCT) is an aggressive malignant mesenchymal tumor of unknown histogenesis. It is predominantly found in the abdominal cavity in adolescents and young adults with a striking male predominance. The cytomorphology of DSRCT, though similar to other small round cell malignancies, displays some distinct features: relatively low cellularity, some cohesiveness retained, slightly angulated nuclei, and acinar-like structures (see Fig. 13.63). The characteristic finding of small nests of tumor cells associated with desmoplastic stroma is better appreciated in cell-block sections. DSRCT exhibits polyphenotypic differentiation, with immunoreactivity for cytokeratins, desmin, and neuroendocrine markers. DSRCT is

characterized by a unique translocation  $t(11;22)(p13;q12)$  with *EWSR1WT1* gene fusion, with can be detected by FISH studies (Table 13.2).

#### *Cytologic features:*

- Moderately cellular smears with loosely cohesive small clusters
  - Desmoplastic stroma fragments
  - Uniform undifferentiated cells with slightly angulated nuclei and scant cytoplasm
  - Nuclear molding and small acinar-like structures
- #### *Differential diagnosis and problems in diagnosis:*
- Metastatic carcinoma (basaloid squamous cell carcinoma)
  - Rhabdomyosarcoma
  - Poorly differentiated synovial sarcoma
  - Ewing sarcoma
  - Neuroendocrine tumor



**Fig. 13.64** Extrarenal rhabdoid tumor of soft tissue. (a) Smears containing both dispersed cells and cohesive cellular clusters composed of typical large rhabdoid cells and smaller round cells. (b, c) Rhabdoid

cells with eccentrically placed large nuclei and abundant cytoplasm with perinuclear densities (MGG)

### 13.13.11 Extrarenal Rhabdoid Tumor of Soft Tissue

This rare, highly malignant tumor of soft tissue occurs in deep soft tissue of the neck and paraspinal region in infants and children. Congenital cases have been reported. This tumor also affects skin and visceral organs such as liver and retroperitoneum. The main cytologic feature is a mixture of dispersed cells and cell clusters of large rhabdoid cells and smaller round or polygonal or spindled cells. The rhabdoid cells have eccentrically placed large nuclei with macronucleoli and perinuclear cytoplasmic densities (*see* Fig. 13.64). Although a diagnosis of a malignancy is relatively easy to render from FNA smears, a correct specific diagnosis needs

the demonstration of *SMARCB1* gene loss by cytogenetic techniques and/or INI1 protein loss by immunohistochemistry [107, 108].

#### *Cytologic features:*

- Variable cellularity with dispersed cells and clusters
- Large rhabdoid cells with eccentric nuclei, prominent nucleoli, and perinuclear density
- Smaller round, polygonal, and spindled cells
- Occasional binucleation and multinucleation
- Brisk mitoses and necrosis

#### *Differential diagnosis and problems in diagnosis:*

- Rhabdomyosarcoma
- Epithelioid sarcoma
- Other sarcomas with rhabdoid differentiation



## References

1. Fletcher CDM, Bridge J, Hogendoorn P, Mertens F, editors. World Health Organization classification of tumours. Pathology and genetics of tumours of soft tissue and bone. 4th ed. Geneva: IARC Press; 2013.
2. Akerman M, Rydholm A, Persson BM. Aspiration cytology of soft-tissue tumors. The 10-year experience at an orthopedic oncology center. *Acta Orthop Scand*. 1985;56:407–12.
3. Akerman M, Willen H. Critical review on the role of fine needle aspiration in soft tissue tumors. *Pathol Case Rev*. 1998;3:111–7.
4. Kilpatrick SE, Cappellari JO, Bos GD, Gold SH, Ward WG. Is fine-needle aspiration biopsy a practical alternative to open biopsy for the primary diagnosis of sarcoma? Experience with 140 patients. *Am J Clin Pathol*. 2001;115:59–68.
5. Domanski HA, Akerman M, Carlen B, Engellau J, Gustafson P, Jonsson K, et al. Core-needle biopsy performed by the cytopathologist: a technique to complement fine-needle aspiration of soft tissue and bone lesions. *Cancer*. 2005;105:229–39.
6. Ng VY, Thomas K, Crist M, Wakely Jr PE, Mayerson J. Fine needle aspiration for clinical triage of extremity soft tissue masses. *Clin Orthop Relat Res*. 2010;468:1120–8.
7. Yang J, Frassica FJ, Fayad L, Clark DP, Weber KL. Analysis of nondiagnostic results after image-guided needle biopsies of musculoskeletal lesions. *Clin Orthop Relat Res*. 2010;468:3103–11.
8. Huang AJ, Kattapuram SV. Musculoskeletal neoplasms: biopsy and intervention. *Radiol Clin North Am*. 2011;49:1287–305. vii.
9. Domanski HA. Fine-needle aspiration cytology of soft tissue lesions: diagnostic challenges. *Diagn Cytopathol*. 2007;35:768–73.
10. Kilpatrick SE, Ward WG, Cappellari JO, Bos GD. Fine-needle aspiration biopsy of soft tissue sarcomas. A cytomorphologic analysis with emphasis on histologic subtyping, grading, and therapeutic significance. *Am J Clin Pathol*. 1999;112:179–88.
11. Layfield LJ. *Cytopathology of bone and soft tissue tumors*. New York: Oxford University Press; 2002.
12. Akerman M, Domanski HA. The cytology of soft tissue tumours. *Monogr Clin Cytol*. 2003;16:IX–X, 1–112.
13. Klijanienko J, Lagace R. *Soft tissue tumor: a multidisciplinary, decisional diagnostic approach*. Hoboken: Wiley-Blackwell; 2011.
14. Domanski HA, Klijanienko J, Lagace R. Sampling procedure, fine needle aspiration and core needle biopsy. In: *Soft tissue tumors*. Hoboken: Wiley-Blackwell; 2011.
15. Dal Cin P, Qian X, Cibas ES. The marriage of cytology and cytogenetics. *Cancer Cytopathol*. 2013;121:279–90. [Epub ahead of print].
16. Trojani M, Contesso G, Coindre JM, Rouesse J, Bui NB, de Mascarel A, et al. Soft-tissue sarcomas of adults; study of pathological prognostic variables and definition of a histopathological grading system. *Int J Cancer*. 1984;33:37–42.
17. Caneiro A, Bendahl PO, Engellau J, Domanski HA, Fletcher CD, Rissler P, et al. A prognostic model for soft tissue sarcoma of the extremities and trunk wall based on size, vascular invasion, necrosis, and growth pattern. *Cancer*. 2011;117:1279–87.
18. Weir MM, Rosenberg AE, Bell DA. Grading of spindle cell sarcomas in fine-needle aspiration biopsy specimens. *Am J Clin Pathol*. 1999;112:784–90.
19. Palmer HE, Mukunyadzi P, Culbreth W, Thomas JR. Subgrouping and grading of soft-tissue sarcomas by fine-needle aspiration cytology: a histopathologic correlation study. *Diagn Cytopathol*. 2001;24:307–16.
20. Mallik MK, Dey P, Gupta SK, Vasishta RK. Grading of soft tissue sarcomas on fine-needle aspiration cytology smear. *Diagn Cytopathol*. 2010;38:109–12.
21. Deyrup AT, Weiss SW. Grading of soft tissue sarcomas: the challenge of providing precise information in an imprecise world. *Histopathology*. 2006;48:42–50.
22. Jones C, Liu K, Hirschowitz S, Klipfel N, Layfield LJ. Concordance of histopathologic and cytologic grading in musculoskeletal sarcomas: can grades obtained from analysis of the fine-needle aspirates serve as the basis for therapeutic decisions? *Cancer*. 2002;96:83–91.
23. Willen H, Akerman M, Carlen B. Fine needle aspiration (FNA) in the diagnosis of soft tissue tumours: a review of 22 years experience. *Cytopathology*. 1995;6:236–47.
24. Dey P, Mallik MK, Gupta SK, Vasishta RK. Role of fine needle aspiration cytology in the diagnosis of soft tissue tumours and tumour-like lesions. *Cytopathology*. 2004;15:32–7.
25. Robertson EG, Baxter G. Tumour seeding following percutaneous needle biopsy: the real story! *Clin Radiol*. 2011;66:1007–14.
26. Stanley MW, Skoog L, Tani EM, Horwitz CA. Nodular fasciitis: spontaneous resolution following diagnosis by fine-needle aspiration. *Diagn Cytopathol*. 1993;9:322–4.
27. Saad RS, Takei H, Lipscomb J, Ruiz B. Nodular fasciitis of parotid region: a pitfall in the diagnosis of pleomorphic adenomas on fine-needle aspiration cytology. *Diagn Cytopathol*. 2005;33:191–4.
28. Plaza JA, Mayerson J, Wakely Jr PE. Nodular fasciitis of the hand: a potential diagnostic pitfall in fine-needle aspiration cytopathology. *Am J Clin Pathol*. 2005;123:388–93.
29. Klapsinou E, Despoina P, Dimitra D. Cytologic findings and potential pitfalls in proliferative myositis and myositis ossificans diagnosed by fine needle aspiration cytology: report of four cases and review of the literature. *Diagn Cytopathol*. 2012;40:239–44.
30. Factor RE, Layfield LJ, Grossmann AH, Crim JR, Price SL, Randall RL. Fine-needle aspiration diagnosis of an intraosseous amyloidoma. *Diagn Cytopathol*. 2012;40 Suppl 2:E114–7.
31. Elkins CT, Scharschmidt TJ, Wakely Jr PE. Amyloidomas of soft parts: diagnosis by fine-needle aspiration. *Diagn Cytopathol*. 2012;40 Suppl 2:E126–30.
32. Bhadani PP, Sah SP, Sen R, Singh RK. Diagnostic value of fine needle aspiration cytology in gouty tophi: a report of 7 cases. *Acta Cytol*. 2006;50:101–4.
33. Kloboves-Prevodnik VV, Us-Krasovec M, Gale N, Lamovec J. Cytological features of lipoblastoma: a report of three cases. *Diagn Cytopathol*. 2005;33:195–200.
34. Domanski HA, Carlen B, Jonsson K, Mertens F, Akerman M. Distinct cytologic features of spindle cell lipoma. A cytologic-histologic study with clinical, radiologic, electron microscopic, and cytogenetic correlations. *Cancer*. 2001;93:381–9.
35. Lemos MM, Kindblom LG, Meis-Kindblom JM, Remotti F, Ryd W, Gunterberg B, et al. Fine-needle aspiration characteristics of hibernoma. *Cancer*. 2001;93:206–10.
36. Gisselsson D, Domanski HA, Hoglund M, Carlen B, Mertens F, Willen H, et al. Unique cytological features and chromosome aberrations in chondroid lipoma: a case report based on fine-needle aspiration cytology, histopathology, electron microscopy, chromosome banding, and molecular cytogenetics. *Am J Surg Pathol*. 1999;23:1300–4.
37. Marino-Enriquez A, Hornick JL, Cibas ES, Qian X. Dedifferentiated liposarcoma and pleomorphic liposarcoma: a comparative study on FNA cytomorphology and MDM2/CDK4 expression. *Mod Pathol*. 2013;26:97A.
38. Szadowska A, Lasota J. Fine needle aspiration cytology of myxoid liposarcoma; a study of 18 tumours. *Cytopathology*. 1993;4:99–106.
39. Klijanienko J, Caillaud JM, Lagace R. Fine-needle aspiration in liposarcoma: cytohistologic correlative study including well-differentiated, myxoid, and pleomorphic variants. *Diagn Cytopathol*. 2004;30:307–12.
40. Downs-Kelly E, Goldblum JR, Patel RM, Weiss SW, Folpe AL, Mertens F, et al. The utility of fluorescence in situ hybridization (FISH) in the diagnosis of myxoid soft tissue neoplasms. *Am J Surg Pathol*. 2008;32:8–13.

41. Rege TA, Madan R, Qian X. Long fascicular tissue fragments in desmoid fibromatosis by fine needle aspiration: a new cytologic feature. *Diagn Cytopathol.* 2012;40:45–7.
42. Owens CL, Sharma R, Ali SZ. Deep fibromatosis (desmoid tumor): cytopathologic characteristics, clinicoradiologic features, and immunohistochemical findings on fine-needle aspiration. *Cancer.* 2007;111:166–72.
43. Domanski HA, Carlen B, Sloth M, Rydholm A. Elastofibroma dorsi has distinct cytomorphologic features, making diagnostic surgical biopsy unnecessary: cytomorphologic study with clinical, radiologic, and electron microscopic correlations. *Diagn Cytopathol.* 2003;29:327–33.
44. Bishop JA, Rekhtman N, Chun J, Wakely Jr PE, Ali SZ. Malignant solitary fibrous tumor: cytopathologic findings and differential diagnosis. *Cancer Cytopathol.* 2012;118:83–9.
45. Lee JC, Fletcher CD. Malignant fat-forming solitary fibrous tumor (so-called “lipomatous hemangiopericytoma”): clinicopathologic analysis of 14 cases. *Am J Surg Pathol.* 2011;35:1177–85.
46. Domanski HA. Fine-needle aspiration smears from lipomatous hemangiopericytoma need not be confused with myxoid liposarcoma. *Diagn Cytopathol.* 2003;29:287–91.
47. Gupta N, Barwad A, Katamuthu K, Rajwanshi A, Radotra BD, Nijhawan R, et al. Solitary fibrous tumour: a diagnostic challenge for the cytopathologist. *Cytopathology.* 2012;23:250–5.
48. Domanski HA, Mertens F, Panagopoulos I, Akerman M. Low-grade fibromyxoid sarcoma is difficult to diagnose by fine needle aspiration cytology: a cytomorphological study of eight cases. *Cytopathology.* 2009;20:304–14.
49. Silverman JF, Nathan G, Olson PR, Prichard J, Cohen JK. Fine-needle aspiration cytology of low-grade fibromyxoid sarcoma of the renal capsule (capsuloma). *Diagn Cytopathol.* 2000;23:279–83.
50. Lindberg GM, Maitra A, Gokaslan ST, Saboorian MH, Albores-Saavedra J. Low grade fibromyxoid sarcoma: fine-needle aspiration cytology with histologic, cytogenetic, immunohistochemical, and ultrastructural correlation. *Cancer.* 1999;87:75–82.
51. Olson MT, Ali SZ. Myxofibrosarcoma: cytomorphologic findings and differential diagnosis on fine needle aspiration. *Acta Cytol.* 2012;56:15–24.
52. Merck C, Hagmar B. Myxofibrosarcoma: a correlative cytologic and histologic study of 13 cases examined by fine needle aspiration cytology. *Acta Cytol.* 1980;24:137–44.
53. Kilpatrick SE, Ward WG. Myxofibrosarcoma of soft tissues: cytomorphologic analysis of a series. *Diagn Cytopathol.* 1999;20:6–9.
54. Klijanienko J, Caillaud JM, Lagace R. Fine-needle aspiration of primary and recurrent dermatofibrosarcoma protuberans. *Diagn Cytopathol.* 2004;30:261–5.
55. Domanski HA. FNA diagnosis of dermatofibrosarcoma protuberans. *Diagn Cytopathol.* 2005;32:299–302.
56. Domanski HA, Gustafson P. Cytologic features of primary, recurrent, and metastatic dermatofibrosarcoma protuberans. *Cancer.* 2002;96:351–61.
57. Rubin BP, Chen CJ, Morgan TW, Xiao S, Grier HE, Kozakewich HP, et al. Congenital mesoblastic nephroma t(12;15) is associated with ETV6-NTRK3 gene fusion: cytogenetic and molecular relationship to congenital (infantile) fibrosarcoma. *Am J Pathol.* 1998;153:1451–8.
58. Hannah CD, Oliver DH, Liu J. Fine needle aspiration biopsy and immunostaining findings in an aggressive inflammatory myofibroblastic tumor of the lung: a case report. *Acta Cytol.* 2007;51:239–43.
59. Stoll LM, Li QK. Cytology of fine-needle aspiration of inflammatory myofibroblastic tumor. *Diagn Cytopathol.* 2011;39:663–72.
60. Gleason BC, Hornick JL. Inflammatory myofibroblastic tumours: where are we now? *J Clin Pathol.* 2008;61:428–37.
61. Wakely Jr PE, Frable WJ. Fine-needle aspiration biopsy cytology of giant-cell tumor of tendon sheath. *Am J Clin Pathol.* 1994;102:87–90.
62. Iyer VK, Kapila K, Verma K. Fine-needle aspiration cytology of giant cell tumor of tendon sheath. *Diagn Cytopathol.* 2003;29:105–10.
63. Ho CY, Maleki Z. Giant cell tumor of tendon sheath: cytomorphologic and radiologic findings in 41 patients. *Diagn Cytopathol.* 2012;40:E94–8.
64. Tao LC, Davidson DD. Aspiration biopsy cytology of smooth muscle tumors. A cytologic approach to the differentiation between leiomyosarcoma and leiomyoma. *Acta Cytol.* 1993;37:300–8.
65. Domanski HA. Cytologic features of angioleiomyoma: cytologic-histologic study of 10 cases. *Diagn Cytopathol.* 2002;27:161–6.
66. Domanski HA, Akerman M, Rissler P, Gustafson P. Fine-needle aspiration of soft tissue leiomyosarcoma: an analysis of the most common cytologic findings and the value of ancillary techniques. *Diagn Cytopathol.* 2006;34:597–604.
67. Klijanienko J, Caillaud JM, Lagace R, Vielh P. Fine-needle aspiration of leiomyosarcoma: a correlative cytohistopathological study of 96 tumors in 68 patients. *Diagn Cytopathol.* 2003;28:119–25.
68. Domanski HA, Dawiskiba S. Adult rhabdomyoma in fine needle aspirates. A report of two cases. *Acta Cytol.* 2000;44:223–6.
69. Gupta N, Rajwanshi A, Mohindra S, Vasishta RK, Batra C, Gupta AK. Diagnosis of adult rhabdomyoma by fine needle aspiration cytology: a report of 2 cases. *Acta Cytol.* 2010;54:968–72.
70. Seidal T, Walaas L, Kindblom LG, Angervall L. Cytology of embryonal rhabdomyosarcoma: a cytologic, light microscopic, electron microscopic, and immunohistochemical study of seven cases. *Diagn Cytopathol.* 1988;4:292–9.
71. Klijanienko J, Caillaud JM, Orbach D, Brisse H, Lagace R, Vielh P, et al. Cyto-histological correlations in primary, recurrent and metastatic rhabdomyosarcoma: the Institut Curie’s experience. *Diagn Cytopathol.* 2007;35:482–7.
72. Kishore B, Khare P, Gupta RJ, Gupta C, Khare V. A rare case of paratesticular pleomorphic rhabdomyosarcoma diagnosed by fine needle aspiration: a case report. *Diagn Cytopathol.* 2010;38:121–6.
73. Ali SZ, Smilari TF, Teichberg S, Hajdu SI. Pleomorphic rhabdomyosarcoma of the heart metastatic to bone. Report of a case with fine needle aspiration biopsy findings. *Acta Cytol.* 1995;39:555–8.
74. Domanski HA, Akerman M, Engellau J, Gustafson P, Mertens F, Rydholm A. Fine-needle aspiration of neurilemmoma (schwannoma). A clinicocytopathologic study of 116 patients. *Diagn Cytopathol.* 2006;34:403–12.
75. Klijanienko J, Caillaud JM, Lagace R. Cytohistologic correlations in schwannomas (neurilemmomas), including “ancient,” cellular, and epithelioid variants. *Diagn Cytopathol.* 2006;34:517–22.
76. Dodd LG, Marom EM, Dash RC, Matthews MR, McLendon RE. Fine-needle aspiration cytology of “ancient” schwannoma. *Diagn Cytopathol.* 1999;20:307–11.
77. Henke AC, Salomao DR, Hughes JH. Cellular schwannoma mimics a sarcoma: an example of a potential pitfall in aspiration cytodiagnosis. *Diagn Cytopathol.* 1999;20:312–6.
78. Wiczorek TJ, Krane JF, Domanski HA, Akerman M, Carlen B, Misdraji J, et al. Cytologic findings in granular cell tumors, with emphasis on the diagnosis of malignant granular cell tumor by fine-needle aspiration biopsy. *Cancer.* 2001;93:398–408.
79. Lee LH, Bos GD, Marsh Jr WL, Wakely Jr PE. Fine-needle aspiration cytology of sclerosing perineurioma. *Ann Diagn Pathol.* 2004;8:80–6.
80. Housini I, Dabbs DJ. Fine needle aspiration cytology of perineurioma. Report of a case with histologic, immunohistochemical and ultrastructural studies. *Acta Cytol.* 1990;34:420–4.



81. McGee Jr RS, Ward WG, Kilpatrick SE. Malignant peripheral nerve sheath tumor: a fine-needle aspiration biopsy study. *Diagn Cytopathol.* 1997;17:298–305.
82. Kljanienko J, Caillaud JM, Lagace R, Vielh P. Cytohistologic correlations of 24 malignant peripheral nerve sheath tumor (MPNST) in 17 patients: the Institut Curie experience. *Diagn Cytopathol.* 2002;27:103–8.
83. Wakely Jr PE, Ali SZ, Bishop JA. The cytopathology of malignant peripheral nerve sheath tumor: a report of 55 fine-needle aspiration cases. *Cancer Cytopathol.* 2012;120:334–41.
84. Boucher LD, Swanson PE, Stanley MW, Silverman JF, Raab SS, Geisinger KR. Cytology of angiosarcoma. Findings in fourteen fine-needle aspiration biopsy specimens and one pleural fluid specimen. *Am J Clin Pathol.* 2000;114:210–9.
85. Wakely Jr PE, Frable WJ, Kneisl JS. Aspiration cytopathology of epithelioid angiosarcoma. *Cancer.* 2000;90:245–51.
86. Kilpatrick SE, Kopyly PD, Ward WG, Richards 2nd F. Epithelioid hemangioendothelioma of bone and soft tissue: a fine-needle aspiration biopsy study with histologic and immunohistochemical confirmation. *Diagn Cytopathol.* 1998;19:38–43.
87. Domanski HA. Fine-needle aspiration of ganglioneuroma. *Diagn Cytopathol.* 2005;32:363–6.
88. Qian X, Hornick JL, Cibas ES, Dal Cin P, Domanski HA. Angiomatoid fibrous histiocytoma a series of five cytologic cases with literature review and emphasis on diagnostic pitfalls. *Diagn Cytopathol.* 2012;40 Suppl 2:E86–93.
89. Mohanty SK, Srinivasan R, Rajwanshi A, Vasishta RK, Vignesh PS. Cytologic diagnosis of ossifying fibromyxoid tumor of soft tissue: a case report. *Diagn Cytopathol.* 2004;30:41–5.
90. Minami R, Yamamoto T, Tsukamoto R, Maeda S. Fine needle aspiration cytology of the malignant variant of ossifying fibromyxoid tumor of soft parts: a case report. *Acta Cytol.* 2001;45:745–55.
91. Kilpatrick SE, Ward WG, Mozes M, Miettinen M, Fukunaga M, Fletcher CD. Atypical and malignant variants of ossifying fibromyxoid tumor. Clinicopathologic analysis of six cases. *Am J Surg Pathol.* 1995;19:1039–46.
92. Gupta S, Gupta R, Singh S, Pant L. Ossifying fibromyxoid tumour of soft parts: report of a case diagnosed on fine needle aspiration cytology. *Cytopathology.* 2012;23:126–8.
93. Gebre-Medhin S, Nord KH, Moller E, Mandahl N, Magnusson L, Nilsson J, et al. Recurrent rearrangement of the PHF1 gene in ossifying fibromyxoid tumors. *Am J Pathol.* 2012;181:1069–77.
94. Antonescu CR, Zhang L, Chang NE, Pawel BR, Travis W, Katabi N, et al. EWSR1-POU5F1 fusion in soft tissue myoepithelial tumors. A molecular analysis of sixty-six cases, including soft tissue, bone, and visceral lesions, showing common involvement of the EWSR1 gene. *Genes Chromosomes Cancer.* 2010;49:1114–24.
95. Akerman M, Domanski HA. The complex cytological features of synovial sarcoma in fine needle aspirates, an analysis of four illustrative cases. *Cytopathology.* 2007;18:234–40.
96. Akerman M, Ryd W, Skytting B, Scandinavian Sarcoma Group. Fine-needle aspiration of synovial sarcoma: criteria for diagnosis: retrospective reexamination of 37 cases, including ancillary diagnostics. A Scandinavian Sarcoma Group study. *Diagn Cytopathol.* 2003;28:232–8.
97. Kljanienko J, Caillaud JM, Lagace R, Vielh P. Cytohistologic correlations in 56 synovial sarcomas in 36 patients: the Institut Curie experience. *Diagn Cytopathol.* 2002;27:96–102.
98. Silverman JF, Landreneau RJ, Sturgis CD, Raab SS, Fox KR, Jasnosh KM, et al. Small-cell variant of synovial sarcoma: fine-needle aspiration with ancillary features and potential diagnostic pitfalls. *Diagn Cytopathol.* 2000;23:118–23.
99. Wakely Jr PE, McDermott JE, Ali SZ. Cytopathology of alveolar soft part sarcoma: a report of 10 cases. *Cancer.* 2009;117:500–7.
100. Lopez-Ferrer P, Jimenez-Heffernan JA, Vicandi B, Gonzalez-Peramato P, Viguer JM. Cytologic features of alveolar soft part sarcoma: report of three cases. *Diagn Cytopathol.* 2002;27:115–9.
101. Hollmann TJ, Hornick JL. INI1-deficient tumors: diagnostic features and molecular genetics. *Am J Surg Pathol.* 2011;35:e47–63.
102. Cardillo M, Zakowski MF, Lin O. Fine-needle aspiration of epithelioid sarcoma: cytology findings in nine cases. *Cancer.* 2001;93:246–51.
103. Creager AJ, Pitman MB, Geisinger KR. Cytologic features of clear cell sarcoma (malignant melanoma) of soft parts: a study of fine-needle aspirates and exfoliative specimens. *Am J Clin Pathol.* 2002;117:217–24.
104. Rau AR, Kini H, Verghese R. Tigroid background in fine-needle aspiration cytology of clear cell sarcoma. *Diagn Cytopathol.* 2006;34:355–7.
105. Jakowski JD, Wakely Jr PE. Cytopathology of extraskeletal myxoid chondrosarcoma: report of 8 cases. *Cancer.* 2007;111:298–305.
106. Domanski HA, Carlen B, Mertens F, Akerman M. Extraskeletal myxoid chondrosarcoma with neuroendocrine differentiation: a case report with fine-needle aspiration biopsy, histopathology, electron microscopy, and cytogenetics. *Ultrastruct Pathol.* 2003;27:363–8.
107. Drut R, Drut RM. Renal and extrarenal congenital rhabdoid tumor: diagnosis by fine-needle aspiration biopsy and FISH. *Diagn Cytopathol.* 2002;27:32–4.
108. Thomson TA, Kljanienko J, Couturier J, Brisse H, Pierron G, Freneaux P, et al. Fine-needle aspiration of renal and extrarenal rhabdoid tumors: the experience of the Institut Curie regarding 20 tumors in 13 patients. *Cancer Cytopathol.* 2011;119:49–57.