

Ravish Shah and Anil K. Agarwal

13.1 Introduction

The superiority of the native arteriovenous fistula (AVF) over other types of accesses including arteriovenous graft (AVG) and tunneled dialysis catheters (TDC) for chronic hemodialysis is a well-recognized fact. It has been shown to have superior patency rates and lower complication rate including a low risk of infection and a lower intervention rate to maintain its patency [1, 2]. This is the core reason underlying the development of guidelines and the Fistula First project that have led to AVF creation in a majority of patients with end-stage renal disease (ESRD). Unfortunately, a considerable number of fistulas (28–53 %) fail to mature sufficiently to support dialysis therapy [3–6]. Failure to mature (FTM) often commits these patients to a tunneled dialysis catheter for a variable length of time until they have a well-functioning arteriovenous (AV) access [3]. In addition to the risk of infection and central venous stenosis, the catheters also contribute to inadequate dialysis and poor patient outcomes [3]. Therefore, early recognition and timely intervention in cases of an AVF with FTM are critically important [3].

13.1.1 Failure to Mature: Definition

Fistula failure can be classified as early and late. Early failure is a true FTM that refers to the cases in which the AV fistula never develops to the point that it can be used or fails within the first 3 months of usage [1]. Late failure refers to those cases where the AVF fails after 3 months of successful usage [7, 8]. Although there might be considerable overlap in the causes of both early and late failure, early failure has gained significant attention as recent data have demonstrated that a great majority of the failed fistulas can be salvaged

using percutaneous interventions [9–12]. While it not infrequent to abandon these AVFs with early failure, aggressive evaluation and treatment have been shown to result in the salvage of vast majority of these accesses [10].

13.1.2 Risk Factors for Failure of Maturation

FTM is a common problem occurring in 28–53 % of native AVF [3–6, 13]. Several studies have looked at factors that might predict fistula maturation. Preoperative vascular mapping has been shown to improve the rate of fistula placement and overall surgical success rate [14–16]. Creation of AVF using very small arteries (e.g., < 1.6 mm in diameter) and veins is likely to fail, although the precise cutoff hinges on the available surgical experience and expertise [14]. Perhaps the most critical determinant of fistula maturation is the functional ability of the artery and vein to dilate and achieve a rapid increase in blood flow after surgery [14]. Several studies have shown that postoperative flow rate measured by Doppler ultrasound in a forearm fistula is a moderately good predictor of fistula maturation [17, 18]. In addition, these studies have reported using a cutoff between 400 and 500 ml/min at 2–8 weeks as a predictor of fistula maturation. Clinical examination of the fistula may be as accurate as Doppler flow measurements [17–19]. Other predictors of AVF failure include age >65 year, diabetes, female gender, and high body mass index (>27), although in the majority of the patients with early FTM, anatomic abnormalities as detected by angiography are present [1].

13.1.3 Causes of Early Fistula Failure

Fistulas can fail to mature for a number of reasons including:

1. Inflow problems: poor arterial inflow and juxta-anastomotic venous stenosis (JAVS).
2. Outflow problems: failure for the vein to “arterialize” and presence of large and/or multiple accessory veins.

R. Shah, MD • A.K. Agarwal, MD, FACP, FASN, FNKF (✉)
Division of Nephrology, The Ohio State University,
395 W 12th Avenue, Ground Floor, Columbus, OH 43210, USA
e-mail: anil.agarwal@osumc.edu

- Other technical factors at the time of surgery: a deep fistula, although mature, might not be easily accessible for cannulation and may require transposition in order to support dialysis adequately. Majority of these causes can and must be identified early in order to salvage the AVF, frequently using percutaneous interventions.

Inflow Problems

A good inflow is critical for fistula maturation and for attaining adequate flow rates to deliver dialysis. After AVF creation, the arterial flow is expected to increase, with increasing arterial diameter and changes in flow pattern [14]. Vascular remodeling and dilation are typically attained by longitudinal shear stress and circumferential deformation, in the milieu of vasoactive factors [14, 20]. This process may continue over a long period of time and contributes to maturation. Rarely, a small-size artery or presence of arterial disease such as atherosclerosis can result in early fistula failure. However, this can be identified and prevented by a comprehensive patient evaluation prior to access placement.

JAVS is one of the most common causes of maturation failure in angiographically evaluated AVF [15] (Fig. 13.1). Although precise etiology is not clear, it is postulated that the JAVS occurs in the swing segment of the vein, where the vein is mobilized to connect with the artery and suffers stretching, torsion, and spasm [21]. It is unclear as to what extent these factors contribute to JAVS; however, the net effect of JAVS is to reduce fistula inflow. JVAS often occurs early in the process and often results in early access failure. In one single-center retrospective study, the authors reported their 12-year experience of radiological management of stenosis and thrombosis in both AVF and AVGs [22]. Of the total 283 patients with AVF, majority of the patients had forearm AVF (74 %) and 74 patients had upper arm AVF. In patients with forearm AVF, JVAS was present in almost half of these patients leading to an inflow problem (Fig. 13.2a). However, of the 74 patients with the upper arm AVF, outflow venous stenosis was predominantly reported in 55 % of cases

($n=41$) (Fig. 13.2b). The vast majority of the stenoses (86 %) were less than two cm long [22]. Fortunately, JVAS is amenable to treatment. It can generally be successfully treated by percutaneous angioplasty surgically [1, 10, 23]. A retrospective analysis of prospectively collected data compared outcomes and cost of surgery ($n=21$) and percutaneous transluminal angioplasty (PTA) ($n=43$) for JAVS in a total of 64 patients [24]. Although the results showed similar cost and success rate, adjusted relative risk was 2.77 for restenosis within the PTA group. The primary 1-year unassisted patency rate for surgery was 91 ± 6 % as compared with 54 ± 8 % with PTA, although adjusted-assisted primary patency rates were similar in the two groups. The surgical approach that had the advantage of less restenosis however was more invasive, involved small but significant risk of loss of venous capital, and was associated with a higher median cost, primarily because of the procedure-related hospitalization. It is important to note that the study was not randomized and only included patients with mature AVF based on the choice made on the basis of available expertise and technical facilities as suggested by the authors. It is worth reemphasizing that JAVS can be easily diagnosed by physical examination [25, 26].

Outflow Problems

Venous dilatation ensues after AVF creation, initially as a result of increased venous pressure and later because of the increase in flow-mediated shear stress [14, 15]. For an AVF to develop and provide satisfactory hemodialysis, there must also be sufficient blood flow through the outflow vein. The absence of good outflow will result in failure of the access. Anomalies that lead to outflow problems include veins that are too small for fistula development, veins that are fibrotic or stenotic, or presence of side branches, referred to as accessory veins. Failure of the dilation of outflow vein has been suggested to be a common cause of maturation failure [27]. Venous stenosis causes failure of the majority of AVF, and endovascular techniques have become popular in the treat-




Fig.13.1 Fistulogram of radiocephalic AVF showing arteriovenous anastomotic stenosis and juxta-anastomotic stenosis of cephalic vein and accessory veins in the forearm




Fig. 13.2 (a, b) Common sites of stenosis in AVF. (a) In wrist AVF. (b) In upper arm AVF (Adapted from Turmel-Rodrigues et al. [22])




Fig. 13.3 (a) Cephalic vein in forearm with severe stenosis that can be angioplastied for maturation of AVF. (b) Cephalic vein in forearm after angioplasty leading to maturation of AVF

ment of most venous stenoses (Fig. 13.3a, b). However, recurrent lesions remain problematic, especially with long segment of severely narrow lesions [28]. Close surveillance and repeated interventions are generally required to maintain patency, although the restenosis at 6 month is significantly less with AVF, compared with AVG [29].

Although, a single cephalic vein stretching from the wrist to the antecubital space is preferred, in many cases, it may be accompanied by one or more accessory veins [25]. These accessory veins are normal anatomy. All of the veins receiving the drainage from the newly created anastomosis enlarge after creation of AVF, and a small accessory vein may

become enlarged with time. The accessory veins must be distinguished from the collateral veins which are pathological and are associated with a downstream (antegrade) stenosis. Ideally the presence of an accessory vein may be viewed as an advantage since it might provide an additional venous channel suitable for cannulation. However, when large (>25 % of the diameter of main AVF), these can steal enough blood flow so that the main fistula channel does not dilate, often resulting in early AVF failure [25, 30] (Fig. 13.4). The accessory veins can often be diagnosed by physical examination [31, 32]. Frequently they are visible or can be detected by palpating the fistula. Also, the thrill that is palpable over




Fig. 13.4 Fistulogram of forearm radiocephalic AVF showing multiple large accessory veins




Fig. 13.5 Physical examination of accessory vein. When the fistula is occluded at *point A*, the thrill will disappear at the anastomosis. As the point of occlusion is moved upward past the accessory vein to *point B*, the thrill will continue when the fistula is occluded (Adapted from Beathard [25])

the arterial anastomosis usually disappears when the downstream (antegrade) fistula is manually occluded, but it does not disappear if an outflow channel (accessory vein) is present below the point of occlusion [25] (Fig. 13.5). Ligation of these accessory veins will redirect the flow to the main channel and promote the development of a usable AVF [1, 10]. Accessory veins together with the JAVS represent the two most common causes of early AVF failure [33, 34]. These two lesions are often present simultaneously [22, 35].

13.1.4 Identification and Management of Early AVF Failure

Identification of patients who are at risk of early AVF failure is critical in order to perform timely intervention to salvage the AVF [1, 15]. Physical examination of the AV access is not only easy to perform and inexpensive, it provides a high level of accuracy [25, 36]. Both accessory veins along with JAVS can be easily identified by a thorough physical examination of the AV access [7]. Please refer to the chapter on “Approach to an AV Access” for details regarding access examination. While ultrasonography can identify these lesions successfully, it is not readily available in all centers and is not free of added cost.

Given the fact that there is very little change in the fistula blood flow or diameter after the first month along with the finding that AVF maturation can be judged with high accuracy via physical examination, it is recommended that all

newly created AVFs should be evaluated by an experienced examiner at 4 weeks [1, 10, 37, 38]. An angiographic study must be performed for non-maturing or poorly mature AVF [3]. In patients who have not initiated dialysis, there is often a concern with use of radiocontrast. However, a small amount of contrast use has been shown to be safe in the evaluation of AVF.

An early identification and intervention approach is critical to the successful maturation of AVF for two reasons. First, a majority of fistulas with early failure demonstrate stenotic lesions within the access circuit [1, 10, 11]. In addition, vascular stenosis is a progressive process eventually culminating in access thrombosis, with the risk of permanently losing the access. Failure to act promptly in these AVFs will result in a loss of the opportunity to salvage an AVF. Second, those patients with early fistula failure are often committed to a tunneled dialysis catheter exposing them to all the dreaded complications of catheter use. Hence, early intervention to identify and salvage early AVF failure becomes an important part of preventing AVF loss and minimizing complications related to tunneled dialysis catheters. Such an approach also supports the “catheter last” approach that the experts advocate.

13.1.5 Specific Interventions

Once a patient with early AVF failure has been identified, appropriate action to salvage the AVF should be taken in a timely manner. As previously mentioned, studies have demonstrated that the two most common problems observed in early AVF failure are the presence of stenosis and accessory veins [9–11]. Fortunately, a great majority of these failed fistulas can be salvaged using percutaneous techniques [1, 11, 39].

13.2 Angioplasty

Endovascular intervention to salvage an immature or failing AVF has become routine. Using radiocontrast, an angiogram of the AVF (commonly termed as “fistulogram”) is done to diagnose the presence of anatomic abnormalities, which can commonly be treated with percutaneous transluminal angioplasty (PTA). PTA is typically indicated when there is >50% stenosis of AVF or AVG [15, 32]. In addition, an inflow lesion, if identified, may be amenable to PTA via a retrograde fistula approach. In a prospective observational study, 100 patients with early failure underwent evaluation and treatment at six freestanding outpatient vascular access centers [1]. Vascular stenosis and the presence of a significant accessory vein alone or in combination were found to be the most common offenders. Venous stenosis was present in 78% of

the cases. A majority (48 %) of these lesions were found to be close to the anastomosis (JAVS). A significant accessory vein was present in 46 % of the cases. Percutaneous balloon angioplasty and accessory vein obliteration using one of the three techniques (percutaneous ligation using 3/0 nylon, venous cut down, or coil insertion) were used to salvage the failed AVF. Angioplasty was performed with a 98 % success rate, and there was 100 % success rate for accessory vein ligation. These interventions resulted in dialysis initiation using the AVF in 92 % of the cases [1]. Upon further analysis, 84 % of the AVF were functional at 3 months, 72 % at 6 months, and 68 % at 12 months [1]. The overall complication rate in this series was 4 %, exclusively seen in patient who underwent angioplasty. Of these, only one patient (1 %) had a major complication consisted of a vein rupture with an expanding hematoma resulting in loss of the access. The three minor complications included low-grade hematomas requiring no treatment and no sequelae [1].

13.3 Accessory/Branching Vein Ligation

Ligation of the accessory veins can be performed surgically or percutaneously with suture ligation and/or embolization. Suture ligation is useful in patients with superficial accessory veins given minimal distance for subcutaneous dissection [40]. Coils within superficial veins can be irritating to patients and possibly erode through the skin. However, coil embolization is preferred in those with deep accessory veins as cutdown suture ligation is more difficult with potential risks of nerve/muscle and tendon injury [40]. Using a percutaneous ligation technique, a separate report also described accessory vein ligation of fistulas that failed to achieve adequate blood flow or size for successful cannulation. Authors reported that of the 17 AV fistulas, 15 (88 %) successfully matured at 1.7 months (± 1 month) after the procedure and were functioning at 44.5 (± 12 weeks) after the first use [9]. In another series of 119 patients with AVF complicated by maturation failure, 29.4 % had a significant accessory vein but that was the sole cause of AVF dysfunction in only 3.4 % [39]. The AVF salvage rate for all lesions was 83 % in this series. These reports suggest that early intervention for maturation failure can salvage a majority of AVF using endovascular techniques [1, 11, 15].

13.4 Sequential Dilatation

Occasionally early fistula failure is found due to a long segment of the vein which is diffusely small or stenosed. Recent reports have highlighted a newer technique (sequential dilatation or balloon-assisted maturation) to salvage an AVF that fails to develop because of diffuse stenosis [5, 41]. In this

technique, the AVF is gradually dilated with a progressively increasing size of angioplasty balloon at 2- to 4-weeks intervals until a size that is optimal for dialysis cannulation is achieved. The goal is to progressively dilate the outflow vein to a point that it is usable for repetitive cannulation and will also deliver adequate blood flow. Dilatation time is typically <20 s mainly to reduce the chance of thrombosis [40]. In addition, shutting down or occluding flow to the AVF by compressing the anastomosis during vein dilation is recommended to prevent venous tears resulting in blood leaking out subsequently causing ecchymosis [40]. In addition, balloon dilatation is usually performed starting from the central to the peripheral vein to reduce the likelihood of blood extravasation as it is easier to pull back a balloon than push it forward [40].

13.5 Surgical Techniques

Surgical interventions include patch angioplasty, creation of a combination of fistula and graft (“graftula”), creation of a new anastomosis for a juxta-anastomotic lesion, and superficialization procedures [3, 40]. However, large-scale randomized prospective studies examining the role of surgical approach in the salvage of AVF with early failure are lacking. Inability to navigate the wire across a stenotic lesion during percutaneous approach and deep location of an AVF are some of the indications for surgical intervention [3].

13.6 Stents in AV Access

Stents have a very limited role in salvage of immature AVF. When dealing with the stenosis, patients with >30 % residual stenosis after PTA of venous stenosis or those with recurrence of the stenosis within 3 months and requiring repeated intervention should be considered for a stent placement [15]. Stents can also be useful in the case of vessel rupture during angioplasty that does not respond to conservative measures. However, the latter is generally associated with poor primary patency [42]. Stents can also be used when PTA has failed and surgery is not feasible due to a variety of reasons. Although stents have been used in coronary and peripheral arterial circulation with decent success, dialysis access demonstrates unique pathologies being in a venous circulation. Self-expanding rather than balloon-expanded stents are commonly used for VA [43]. These include bare metal stainless steel stents or nitinol shape memory alloy recoverable technology (SMART) stents that are made of nickel titanium alloy [15]. These have physical characteristics that allow more deformability as compared with bare metal stents. Stent grafts are composed of nitinol skeleton covered by graft material on both sides. Stents available until recently

have been used off-label to improve patency in patients with VA stenosis, primarily in AVG, with variable results. Stent placement has several disadvantages including migration, fracture, and in-stent restenosis [15]. Infectious complications are usually not evident until many days after the procedure [44]. In addition, due to the stent placed in the venous segment, loss of vein length may jeopardize future AVF creation [15]. Despite the recent advances in knowledge, both technical and theoretical, the role of stent placement in the management of hemodialysis access dysfunction remains controversial. It will remain so until large, multicenter, prospective, randomized, controlled trials are conducted [44]. Stent placement should be utilized only after considering the type, location, and frequency of recurrence of the lesion. Possibility of a secondary AVF must be considered to avoid the loss of available venous length from stent placement.

13.7 Thrombectomy

If the immature AVF is thrombosed, then one can perform a thrombectomy with simple PTA maceration of the clot in most cases [5]. There is typically a minimal amount of thrombus usually located at the juxta-anastomotic region. Use of heparin is typically indicated. The treatment should also include prompt detection and treatment of the underlying anatomic abnormality and evaluation and management of outflow, including central veins. Percutaneous declotting of AVF is more difficult than declotting of AVG, with success rates that vary between 73 and 96 % in published literature [45]. With the advent of new technology and growing expertise in the field on interventional nephrology, the results of percutaneous techniques have improved significantly and are now comparable to surgical thrombectomy with restoring AVF patency in >90 % of cases [46]. However, the results seem to vary with operator experience and available resources.

13.8 Prevention of Early FTM

Appropriate preoperative evaluation of the patients prior to AVF creation will not only increase chances of AVF creation but also of AVF maturation. Use of physical examination, ultrasonography, and occasional venography are recommended based on individual case. Use of certain pharmacologic agents, especially the antiplatelet agents, has been noted to be associated with improved survival of AVF but has not been proved conclusively to improve the use of AVF in randomized controlled trials despite reduction in AVF thrombosis [47–50]. Many novel therapies are being evaluated to improve maturation of AVF. Local delivery of endothelial cells as a wrap can reduce development of neointimal

hyperplasia at the arteriovenous anastomosis [51]. Perivascular wraps of antiproliferative agents (paclitaxel) and gene therapy with adenoviral vectors have been tried [52]. Use of venous and arterial allografts as well as decellularized xenografts has been tried in those with unsuitable veins. Better hemodynamics by way of using a premade arteriovenous anastomosis is also being tried in clinical studies.

Conclusion

It is crucial to evaluate a newly created AVF at 4–6 weeks after placement to identify candidates with early AVF failure. Physical examination is a simple but efficient modality of identifying such candidates. Once identified, these patients then should be referred to an interventionalist for evaluation and appropriate intervention. Delays in such intervention may result in the delivery of dialysis with a tunneled dialysis catheter, rendering the patient susceptible to higher complications as well as to a risk of eventual thrombosis leading to permanent loss of access. By use of the percutaneous endovascular techniques such as balloon angioplasty and vein obliteration, majority of early fistula failures can be rescued.

References

1. Beathard GA, Arnold P, Jackson J, Litchfield T, Physician Operators Forum of RMS Lifeline, Inc. Aggressive treatment of early fistula failure. *Kidney Int.* 2003;64:1487–94.
2. Allon M, Robbin ML. Increasing arteriovenous fistulas in hemodialysis patients: problems and solutions. *Kidney Int.* 2002;62:1109–24.
3. Asif A, Roy-Chaudhury P, Beathard GA. Early arteriovenous fistula failure: a logical proposal for when and how to intervene. *Clin J Am Soc Nephrol.* 2006;1:332–9.
4. Allon M, Lockhart ME, Lilly RZ, et al. Effect of preoperative sonographic mapping on vascular access outcomes in hemodialysis patients. *Kidney Int.* 2001;60:2013–20.
5. Asif A, Cherla G, Merrill D, Cipleu CD, Briones P, Pennell P. Conversion of tunneled hemodialysis catheter-consigned patients to arteriovenous fistula. *Kidney Int.* 2005;67:2399–406.
6. Won T, Jang JW, Lee S, Han JJ, Park YS, Ahn JH. Effects of intraoperative blood flow on the early patency of radiocephalic fistulas. *Ann Vasc Surg.* 2000;14:468–72.
7. Beathard GA. Strategy for maximizing the use of arteriovenous fistulae. *Semin Dial.* 2000;13:291–6.
8. Beathard GA. Angioplasty for arteriovenous grafts and fistulae. *Semin Nephrol.* 2002;22:202–10.
9. Faiyaz R, Abreo K, Zaman F, et al. Salvage of poorly developed arteriovenous fistulae with percutaneous ligation of accessory veins. *Am J Kidney Dis.* 2002;39:824–7.
10. Beathard GA, Settle SM, Shields MW. Salvage of the nonfunctioning arteriovenous fistula. *Am J Kidney Dis.* 1999;33:910–6.
11. Turmel-Rodrigues L, Mouton A, Birmele B, et al. Salvage of immature forearm fistulas for hemodialysis by interventional radiology. *Nephrol Dial Transplant.* 2001;16:2365–71.
12. Corpataux JM, Haesler E, Silacci P, et al. Low-pressure environment and remodelling of the forearm vein in Brescia-Cimino haemodialysis access. *Nephrol Dial Transplant.* 2002;17:1057–62.

13. Miller PE, Tolwani A, Luscly CP, Deierhoi MH, Bailey R, Redden DT, Allon M. Predictors of adequacy of arteriovenous fistulas in hemodialysis patients. *Kidney Int.* 1999;56:275–80.
14. Dixon BS. Why don't fistulas mature? *Kidney Int.* 2006;70:1413–22.
15. Agarwal A, Asif A. Maintenance and salvage of vascular access: current concepts. *Intervent Nephrol.* 2009;8(5):353–9.
16. Silva Jr MB, Hobson II RW, Pappas PJ, et al. A strategy for increasing use of autogenous hemodialysis access procedures: impact of preoperative noninvasive evaluation. *J Vasc Surg.* 1998;27:302–7.
17. Lin SL, Chen HS, Huang CH, et al. Predicting the outcome of hemodialysis arteriovenous fistulae using duplex ultrasonography. *J Formos Med Assoc.* 1997;96:864–8.
18. Robbin ML, Oser RF, Allon M, et al. Hemodialysis access graft stenosis: US detection. *Radiology.* 1998;208:655–61.
19. Wong V, Ward R, Taylor J, et al. Factors associated with early failure of arteriovenous fistulae for haemodialysis access. *Eur J Vasc Endovasc Surg.* 1996;12:207–13.
20. Kim YO, Choi YJ, Kim JI, et al. The impact of intima-media thickness of radial artery on early failure of radiocephalic arteriovenous fistula in hemodialysis patients. *J Korean Med Sci.* 2006;21:284–9.
21. Konner K, Nonnast-Daniel B, Ritz E. The arteriovenous fistula. *J Am Soc Nephrol.* 2003;14:1669–80.
22. Turmel-Rodrigues L, Pengloan J, Baudin S, et al. Treatment of stenosis and thrombosis in haemodialysis fistulas and grafts by interventional radiology. *Nephrol Dial Transplant.* 2000;15:2032–6.
23. Romero A, Polo JR, Morato EG, et al. Salvage of angioaccess after late thrombosis of radiocephalic fistulas for hemodialysis. *Int Surg.* 1999;71:122–4.
24. Tessitore N, Mansueto G, Lipari G, et al. Endovascular versus surgical preemptive repair of forearm arteriovenous fistula juxta-anastomotic stenosis: analysis of data collected prospectively from 1999 to 2004. *Clin J Am Soc Nephrol.* 2006;1:448–54.
25. Beathard GA. Physical examination of the dialysis vascular access. *Semin Dial.* 1998;11:231–6.
26. Beathard GA. Physical examination: the forgotten tool. In: Gray R, Sands J, editors. *A multidisciplinary approach for hemodialysis access.* New York: Lippincott Williams & Wilkins; 2002. p. 111–8.
27. Tordoir JH, Rooyens P, Dammers R, Van der Sande FM, De Haan M, Yo TI. Prospective evaluation of failure modes in autogenous radiocephalic wrist access for hemodialysis. *Nephrol Dial Transplant.* 2003;18:378–83.
28. Clark TW, Hirsch DA, Jindal KJ, et al. Outcome and prognostic factors of restenosis after percutaneous treatment of native hemodialysis fistulas. *J Vasc Interv Radiol.* 2002;13:51–9.
29. Turmel-Rodrigues L, Pengloan J, Blanchier D, et al. Insufficient dialysis shunts: improved long term patency rates with close hemodynamic monitoring, repeat percutaneous balloon angioplasty, and stent placement. *Radiology.* 1993;187:273–8.
30. Miles AM. Upper limb ischemia after vascular access surgery: differential diagnosis and management. *Semin Dial.* 2000;13:312–6.
31. Leon C, Orozco-Vargas LC, et al. Accuracy of Physical Examination in the Detection of Arteriovenous Graft Stenosis. *Seminars in Dialysis.* 2008;21(1):85–8.
32. National Kidney Foundation. *K/DOQI clinical practice guidelines for vascular access, guideline 19: treatment of stenosis without thrombosis in dialysis AV grafts and primary AV fistulae.* *Am J Kidney Dis.* 2002;37(1):S163–4.
33. Beathard GA. Percutaneous transvenous angioplasty in the treatment of vascular access stenosis. *Kidney Int.* 1992;42:1390–7.
34. Vesely TM, Siegel JB. Use of the peripheral cutting balloon to treat hemodialysis-related stenoses. *J Vasc Interv Radiol.* 2005;16:1593–603.
35. Tessitore N, Mansueto G, Bedogna V, et al. A prospective controlled trial on effect of percutaneous transluminal angioplasty on functioning arteriovenous fistulae survival. *J Am Soc Nephrol.* 2003;14:1623–7.
36. Migliacci R, Selli ML, Falcinelli F, et al. Assessment of occlusion of the vascular access in patients on chronic hemodialysis: comparison of physical examination with continuous-wave Doppler ultrasound. *Nephron.* 1999;82(1):7.
37. Robbin ML, Chamberlain NE, Lockhart ME, et al. Hemodialysis arteriovenous fistula maturity: US evaluation. *Radiology.* 2002;225:59–64.
38. Robbin ML, Gallichio MH, Deierhoi MH, et al. US vascular mapping before hemodialysis access placement. *Radiology.* 2000;217:83–8.
39. Nassar GM, Nguyen B, Rhee E, Achkar K. Endovascular treatment of the “failing to mature” arteriovenous fistula. *Clin J Am Soc Nephrol.* 2006;1:275–80.
40. Falk A, Rajan D. The immature or failure to mature fistula. In: *Essentials of percutaneous dialysis interventions.* New York: Springer; 2011. p. 323–39.
41. Achkar K, Nassar GM. Salvage of a severely dysfunctional arteriovenous fistula with a strictured and occluded outflow tract. *Semin Dial.* 2005;18:336–42.
42. Rajan D, Clark TW. Patency of wallstents placed at the venous anastomosis of dialysis grafts for salvage of angioplasty-induced rupture. *Cardiovasc Intervent Radiol.* 2003;26:242–5.
43. Urbanes AQ. Proper use and indications for stents in vascular access salvage. *Quest.* 2008;15:15–9.
44. Yevzlin A, Asif A. Stent placement in hemodialysis access: historical lessons, the state of the art and future directions. *Clin J Am Soc Nephrol.* 2009;4:996–1008.
45. Moossavi S, Reagan J, Pierson ED, Kasey JM, et al. Non-surgical salvage of thrombosed arteriovenous fistulae: a case series and review of the literature. *Semin Dial.* 2008;20:459–64.
46. Schon D, Mishler R. Salvage of occluded arteriovenous fistulae. *Am J Kidney Dis.* 2000;36:804–10.
47. Dember LM, Beck GJ, Allon M, et al. Effect of clopidogrel on early failure of arteriovenous fistulas for hemodialysis: a randomized controlled trial. *JAMA.* 2008;299:2164.
48. Saran R, Dykstra DM, Wolfe RA, Gillespie B, Held PJ, Young EW, Dialysis Outcomes and Practice Patterns Study. Association between vascular access failure and the use of specific drugs: the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Am J Kidney Dis.* 2002;40:1255–63.
49. Yevzlin AS, Conley EL, Sanchez RJ, Young HN, Becker BN. Vascular access outcomes and medication use: a USRDS study. *Semin Dial.* 2006;19:535–9.
50. Hasegawa T, et al. Consistent aspirin use associated with improved arteriovenous fistula survival among incident hemodialysis patients in the dialysis outcomes and practice patterns study. *Clin J Am Soc Nephrol.* 2008;3:1373–8.
51. Nuget HM, Groothuis A, Seifert P, Guerraro JL, Nedelman M, Mohankumar T, Edelman ER. Perivascular endothelial implants inhibit intimal hyperplasia in a model of arteriovenous fistulae: a safety and efficacy study in the pig. *J Vasc Res.* 2002;39:524–33.
52. Roy-Chaudhury P, Kelly BS, Zhang J, Narayana A, Desai P, Melham D, Duncan H, Heffelfinger SC. Hemodialysis vascular access dysfunction: from pathophysiology to novel therapies. *Blood Purif.* 2003;21:99–110.