

Chapter 2

Key Players in the Regulation of Fungal Secondary Metabolism

Benjamin P. Knox and Nancy P. Keller

Introduction

Beyond their environmental ubiquity and critical roles in nearly every ecological niche as primary decomposers, fungi are well known for producing a wealth of low molecular weight molecules called secondary metabolites, which are also known as natural products. Although the true ecological function of most secondary metabolites (SMs) is still unknown, their roles as biotic and abiotic protectants or defensive metabolites is emerging [1–4]. Furthermore, their significant impact on human well-being, both positive and negative, makes them attractive study targets.

With a broad spectrum of biological activity, SMs can have major influences on human health. For example, subsets of SMs known as mycotoxins are responsible for millions of dollars in crop loss annually just in the USA alone [5]. When mycotoxin contamination of consumables goes undetected, the resulting mycotoxicoses have additional health and economic consequences, which has been documented throughout recorded history [6]. Crop losses and health consequences are especially devastating in developing countries where testing for toxin contamination is either not well established or is nonexistent [7]. Conversely, the diverse pharmacodynamics of medically relevant SMs such as the β (beta)-lactam antibiotics penicillin and cephalosporin, and the popular cholesterol-lowering agent lovastatin, are examples of how fungal SMs have had positive impacts on human well-being.

B. P. Knox (✉)

Department of Medical Microbiology and Immunology, University of Wisconsin-Madison,
1550 Linden Drive, 53706 Madison, Wisconsin, USA
e-mail: bpknox@wisc.edu

N. P. Keller

Department of Medical Microbiology and Immunology, Department of Bacteriology,
University of Wisconsin-Madison, 1550 Linden Drive, 53706 Madison, Wisconsin, USA
e-mail: npkeller@wisc.edu

At the genomic level, one of the defining hallmarks of SM is the grouping of biosynthetic genes into discreet clusters [8] and general localization to subtelomeric regions of the chromosome [9]. Many, but not all, clusters contain cluster-specific transcription factors that regulate expression of the biosynthetic genes for their respective metabolites. This clustering, coupled with unique chromosomal location, allows for multiple regulatory layers giving the producing fungus precise spatial and temporal control over metabolite expression and likely contributes to intra-, and possibly inter-kingdom, horizontal cluster transfer [10–12]. Additionally, SM production is often tightly correlated with growth and development [13], and often-times disruption of one process will have a significant effect on the other.

The study of fungal SMs has established several model fungal systems and particular metabolites with facile laboratory characterization allowing clear study of their expression and regulation. Yet, as an increasing number of fungal genomes become available, it has become clear that the number of metabolites observed under routine laboratory conditions is a paltry representation of the theoretic yield based on predicted *in silico* genomic regions containing signature SM biosynthetic cluster motifs [14]. Resulting from these observations, endeavors to unlock these “cryptic” gene clusters have shed light on SM regulators working at the level of individual metabolites or at a larger, global scale.

In this chapter, we review the general mechanisms of key players in the regulation of fungal SM. As will become evident in the following paragraphs, fungal SM regulation does not follow a strict hierarchical regime and is composed of overlapping and interconnected pathways making regulatory classifications of regulators less amenable to simple progressional delineations. As such, we will group and present key players based on the largely cluster-specific $\text{Zn(II)}_2\text{Cys}_6$ family of transcription factors, then investigate additional families that possess more pleiotropic regulatory characteristics, and finish with global regulators and multiprotein complexes that respond to environmental cues.

$\text{Zn(II)}_2\text{Cys}_6$

The grouping of SM biosynthetic genes into discreet and contiguous clusters is a distinguishing feature of SM in fungi [8]. In-cluster transcription factors are commonly found that exhibit regulatory control over their respective cluster’s transcription of biosynthetic genes. As the production and regulation of SMs in fungi are replete with diversity, it is not surprising that multiple families of transcription factors regulate these processes. Therefore, we will begin with the largest cluster-specific family of fungal transcription factors (Table 2.1) [15–35].

Unique to fungi are the $\text{Zn(II)}_2\text{Cys}_6$ family of transcription factors. Some distinguishing features of these proteins are the binding of two zinc atoms to six cysteine residues and their ability to bind DNA as monomers, homodimers, or heterodimers [36]. Common nomenclatures for $\text{Zn(II)}_2\text{Cys}_6$ transcription factors include: C_6 , Zn_2C_6 , zinc binuclear cluster, as well as zinc cluster. Compared to other families of transcription factors, $\text{Zn(II)}_2\text{Cys}_6$ proteins are the most common family of dedicated cluster regulators for fungal SM.

Table 2.1 Examples of Zn(II)₂Cys₆ cluster-specific transcription factors (adapted from Yin and Keller 2011 [15])

Transcription factor	Target cluster	Species	Source
AflR	Aflatoxin/ sterigmatocystin	<i>Aspergillus nidulans</i> <i>Aspergillus flavus</i> <i>Aspergillus parasiticus</i>	Brown et al. 1996 [16] Woloshuk et al. 1994 [17] Chang et al. 1995 [18]
AfoA	Asperfuranone	<i>Aspergillus nidulans</i>	Chiang et al. 2009 [19]
ApdR	Aspyridone	<i>Aspergillus nidulans</i>	Bergmann et al. 2007 [20]
Bik5	Bikaverin	<i>Fusarium fujikuroi</i>	Wiemann et al. 2009 [21]
CTB8	Cercosporin	<i>Cercospora nicotianae</i>	Chen et al. 2007 [22]
CtnA	Citrinin	<i>Monascus purpureus</i>	Shimizu et al. 2007 [23]
CtnR	Asperfuranone	<i>Aspergillus nidulans</i>	Chiang et al. 2009 [19]
DEP6	Depudecin	<i>Alternaria brassicicola</i>	Wight et al. 2009 [24]
FapR	Fumagillin/pseurotin	<i>Aspergillus fumigatus</i>	Wiemann et al. 2013 [25]
FUM21	Fumonisin	<i>Fusarium verticillioides</i>	Brown et al. 2007 [26]
GIP2	Aurofusarin	<i>Gibberella zeae</i>	Kim et al. 2006 [27]
GliZ	Gliotoxin	<i>Aspergillus fumigatus</i>	Bok et al. 2006b [28]
LovE	Lovastatin	<i>Aspergillus terreus</i>	Kennedy et al. 1999 [29]
MdpE	Monodictyphenone	<i>Aspergillus nidulans</i>	Chiang et al. 2010 [30]
MlcR	Compactin	<i>Penicillium citrinum</i>	Abe et al. 2002b [31]
MokH	Monacolin K	<i>Monascus pilosus</i>	Chen et al. 2010 [32]
ORFR	AK-toxin	<i>Alternaria alternata</i>	Tanaka and Tsuge 2000 [33]
SirZ	Sirodesmin PL	<i>Leptosphaeria maculans</i>	Fox et al. 2008 [34]
ZFR1	Fumonisin	<i>Fusarium verticillioides</i>	Flaherty and Woloshuk 2004 [35]

To date, the most well-characterized cluster-specific transcription factor is the aflatoxin (AF)/sterigmatocystin (ST) regulator AflR. As the paradigm for Zn(II)₂Cys₆-mediated cluster regulation, AflR has unequivocally established the importance of studying transcriptional regulators in the context of SM expression and regulation. Deletion of *aflR* in *Aspergillus nidulans* produced a condition in which ST production was suppressed at the transcriptional level, even under ST-stimulating conditions [37] and, conversely, increasing expression of *aflR* in normally non-AF producing conditions resulted in concomitant expression of AF biosynthetic genes [18].

Many of the SM clusters characterized in *A. nidulans* contain Zn(II)₂Cys₆ proteins including AfoA and CtnR, ApdR, and MdpE required for asperfuranone, aspyridone, and monodictyphenone gene expression, respectively [19, 20, 30].

Overexpression of $\text{Zn(II)}_2\text{Cys}_6$ proteins can be sufficient to activate silent SM clusters as was first reported for ApdR, thus representing a critical proof-of-principle for fungal genome mining in the postgenomic era [20]. This technique, however, does not always work, resulting in the development of alternative methods to bypass $\text{Zn(II)}_2\text{Cys}_6$ regulation altogether and target individual keystone SM genes [38]. Cluster-specific $\text{Zn(II)}_2\text{Cys}_6$ transcription factors regulating SM production in plant pathogenic fungi include CTB8 for cercosporin, DEP6 for depudecin, FUM21 and ZFR1 for fumonisin, GIP2 for aurofusarin, ORFR for AK-toxin, and SirZ for sirodesmin PL in *Cercospora nicotianae*, *Alternaria brassicicola*, *Fusarium verticillioides*, *Gibberella zeae*, *Alternaria alternata*, and *Leptosphaeria maculans*, respectively [22, 24, 26, 27, 33–35]. Additionally, Bik5 controls synthesis of the mycelial pigment bikaverin in *Fusarium fujikuroi* [21]. Understanding $\text{Zn(II)}_2\text{Cys}_6$ cluster regulation is also critical in biotechnologically relevant fungi given that this family of transcription factors can also control production of the important antihypercholesterolemic agents lovastatin and monacolin K, as well as the pravastatin sodium precursor compactin, regulated by LovE, MokH, and MlcR in *Aspergillus terreus*, *Monascus pilosus*, and *Penicillium citrinum*, respectively [29, 31, 32, 39]. Recently, an interesting new twist to the canonical view of a $\text{Zn(II)}_2\text{Cys}_6$ regulating a single SM came with the finding of the transcription factor FapR that simultaneously regulates biosynthetic genes of the intertwined fumagillin and pseurotin cluster in *Aspergillus fumigatus* [25]. Table 2.1 provides a few additional examples of these types of transcription factors [15–35].

CYS₂HIS₂

Common to fungi (Table 2.2) and all other eukaryotes are the Cys₂His₂ family of transcription factors [21, 40–69]. Defining features of this family include two or more of the conserved, repeating amino acid zinc finger units that bind a single zinc atom [36]. Cys₂His₂ proteins bind DNA as monomers and can also be referred to as C₂H₂ and classical zinc finger transcription factors.

For surviving a multitude of environmental challenges, fungi often require structural pigments for withstanding biotic and abiotic stresses [70]. A unique group of orthologous Cys₂His₂ transcription factors has been found in several plant pathogenic fungi, all positively regulating biosynthesis of the structural pigment melanin. The proteins Cmr1p, Pig1p, Cmr1, and BMR1 are found in *Colletrichum lagenarium*, *Magnaporthe grisea*, *Cochliobolus heterostrophus*, and *Bipolaris oryzae*, respectively [43, 45, 46]. In addition to two Cys₂His₂ motifs, these four proteins also possess a $\text{Zn(II)}_2\text{Cys}_6$ sequence. The earliest recorded examples of SM regulation by Cys₂His₂ proteins are MRTRI6 and Tri6 regulating trichothecene mycotoxin gene clusters in the plant pathogenic fungi *Myrothecium roridum* and *Fusarium sporotrichioides*, respectively [50, 58]. In one of the few cases of an SM-specific regulator gene lying outside of the cluster it controls, the Cys₂His₂ transcription factor ScpR on chromosome II was shown to activate the asperfuranone cluster in

Table 2.2 Transcription factor families and their regulated metabolite(s)

Regulator	Metabolite(s)	Species	Source
Cys ₂ His ₂			
AreA	AF gibberellin fumonisin	<i>Aspergillus parasiticus</i> <i>Gibberella fujikuroi</i> <i>Fusarium verticillioides</i>	Chang et al. 2000 [40] Mihlan et al. 2003 [41] Kim and Woloshuk 2008 [42]
BMR1	Melanin	<i>Bipolaris oryzae</i>	Kihara et al. 2008 [43]
BcYOH1	Botrydial/botcinic acid	<i>Botrytis cinerea</i>	Simon et al. 2013 [44]
Cmr1	Melanin	<i>Cochliobolus heterostrophus</i>	Eliahu et al. 2007 [45]
Cmr1p	Melanin	<i>Colletrichum lagenarium</i>	Tsuji et al. 2000 [46]
CreA	Flavipucine cephalosporin	<i>Aspergillus terreus</i> <i>Acremonium chrysogenum</i>	Gressler et al. 2011 [47] Jekosch and Kück 2000 [48, 49]
MRTRI6	Trichothecene	<i>Myrothecium roridum</i>	Trapp et al. 1998 [50]
PacC	Penicillin ST/AF cephalosporin bikaverin fumonisin gluconic acid	<i>Aspergillus nidulans</i> <i>A. nidulans/A. parasiticus</i> <i>Acremonium chrysogenum</i> <i>Fusarium fujikuroi</i> <i>Fusarium verticillioides</i> <i>Penicillium expansum</i>	Bergh and Brakhage 1998 [51] Keller et al. 1997 [52] Schmitt et al. 2001 [53] Wiemann et al. 2009 [21] Flaherty et al. 2003 [54] Barad et al. 2013 [55]
Pig1p	Melanin	<i>Magnaporthe grisea</i>	Tsuji et al. 2000 [46]
Sda1	Fumonisin B ₁	<i>Fusarium verticillioides</i>	Malapi-Wight et al. 2013 [56]
ScpR	Asperfuranone	<i>Aspergillus nidulans</i>	Bergmann et al. 2010 [57]
Tri6	Trichothecene	<i>Fusarium sporotrichioides</i>	Hohn et al. 1999 [58]
bZip			
Aoyap1	Ochratoxin	<i>Aspergillus ochraceus</i>	Reverberi et al. 2012 [59]
AtfB	AF	<i>Aspergillus parasiticus</i>	Roze et al. 2011 [60]
HapX	Ferricrocin	<i>Aspergillus nidulans</i>	Eisendle et al. 2006 [61]
MeaB	Bikaverin AF	<i>Fusarium fujikuroi</i> <i>Aspergillus flavus</i>	Wagner et al. 2010 [62] Amaike et al. 2013 [63]
RsmA	ST/asperthecin gliotoxin	<i>Aspergillus nidulans</i> <i>Aspergillus fumigatus</i>	Yin et al. 2012 [64] Sekonyela et al. 2013 [65]
ToxE	HC-toxin	<i>Cochliobolus carbonum</i>	Bussink et al. 2001 [66]; Pedley and Walton 2001 [67]
Winged helix			
AcFKH1	Cephalosporin C	<i>Acremonium chrysogenum</i>	Schmitt et al. 2004 [68]
CPCR1	Cephalosporin C	<i>Acremonium chrysogenum</i>	Schmitt and Kuck 2000 [69]

A. nidulans, likely through binding the promoter region of *afmA*, embedded in the asperuranone cluster located on chromosome VIII [57].

Recently, the characterization of two new Cys₂His₂ transcription factors has shown regulation of other physiological processes beyond SM. Exhibiting positive regulation over the toxins botrydial and botcinic acid in *Botrytis cinerea*, BcYOH1 exhibited a more global regulatory role as it also affects mechanisms in detoxification, virulence, and carbohydrate metabolism [44], whereas *sda1* knockout strains of *F. verticillioides* had excessive fumonisin B₁ biosynthesis, reduced capacity to form conidia, and an inability to grow on selected carbon sources [56].

bZIP

Found in all eukaryotic organisms, basic leucine zipper (bZIP) transcription factors are characterized by basic and leucine zipper regions. The basic region dictates sequence-specific DNA-binding whereas the leucine zipper region mediates dimerization of the protein. As dimers, bZIPs target palindromic DNA sequences. Many fungal bZIPs (Table 2.2 [21, 40–69]) have been characterized as stress response transcription factors, responding to a variety of environmental stresses that appears to link them to SM production [60, 71, 72]. bZIPs associated with SM regulation and stress include RsmA regulating sterigmatocystin and asperthecin in *A. nidulans* and gliotoxin in *A. fumigatus* [64, 65], AtfB regulating aflatoxin in *Aspergillus parasiticus* [60], and Aoyap1 regulating ochratoxin in *Aspergillus ochraceus* [59]. The bZIP protein MeaB, involved in nitrogen regulation, has also been associated with regulation of SMs including bikaverin in *F. fujikuroi* and AF in *A. flavus* [62, 63].

In the plant pathogen *Cochliobolus carbonum*, HC-toxin biosynthetic genes are regulated by the hybrid bZIP/ankyrin repeat transcription factor ToxE [73]. Despite having the basic region characteristic of bZIPs, ToxE lacks the leucine zipper sequence but possesses four ankyrin repeats. Both the basic region and ankyrin repeat region mediate DNA binding to promoter regions of all HC-toxin biosynthesis genes [67]. ToxE, together with the putative transcription factor Bap1 from the tomato pathogen *Cladosporium fulvum*, represent a potentially novel class of fungal-specific hybrid transcription factors possessing bZIP and ankyrin repeat characteristics [66].

Winged Helix

Winged helix proteins are found in all organisms and belong within the general helix-turn-helix structural group of proteins. Broadly speaking, the structure of a winged helix protein consists of two wings, three α (alpha) helices, and three β (beta) strands [74]. The industrially relevant fungus *Acremonium chrysogenum* is well

known for production of the antibiotic cephalosporin C. Cluster-specific regulation of cephalosporin C production in *A. chrysogenum* was first shown to be controlled by the transcription factor CPC1, which belongs to the winged helix subfamily of regulatory factor X (RFX) proteins [69]. Another regulator of cephalosporin C, AcFKH1 belongs to a subfamily of winged helix proteins possessing a forkhead associated domain (FHA) and a forkhead DNA-binding domain (FKH) [68]. Consistent with the observation that SM production is often inextricably linked to morphological development, it was shown that CPC1 is not only required for cephalosporin C production, but also for the formation of arthrospores and whereas *AcFKH1* deletion strains still retained the ability to form arthrospores despite possessing swollen and highly septate hyphae [75]. To the best of our knowledge, cephalosporin C is the only fungal SM currently known to be regulated by winged helix proteins (Table 2.2 [21, 40–69]).

Global Regulators

AreA—Nitrogen

So far we have reviewed transcription factors largely characterized as cluster-specific and will now move into global regulators that translate environmental cues into SM and concomitant physiological responses. In *A. nidulans* an increase in ST, and an increase in the rate of sexual development, is observed when grown on nitrate, whereas an opposite response for both phenotypes is observed on ammonium media [76]. Conversely, ammonium stimulates AF biosynthesis in *A. parasiticus* and nitrate inhibits its production [77]. Comparing the SM response to varying nitrogen sources between these two *Aspergillus* species suggests a highly dynamic interplay between environment, SM adaptation, and developmental regime.

Among fungi, the highly conserved global transcription factor AreA is responsible for repression of nitrogen metabolism in the presence of glutamine or ammonium, and is a member of the GATA family of transcription factors, which are conserved among eukaryotes and are characterized by their Cys₂Hys₂ zinc finger DNA binding domains [78]. Beyond its regulatory role in primary metabolism, AreA modulates morphological development and SM regulation in filamentous fungi and is likely responsible for the aforementioned species-specific mycotoxin responses based on the observation of multiple GATA sequences in the aflatoxin and sterigmatocystin regulatory genes *aflR* and *aflJ*, and subsequent AreA binding to these regions in *A. parasiticus* [40]. The rice pathogen *Gibberella fujikuroi* exhibits decreased transcript levels for nearly all structural genes for the SM gibberellin in an *areA* deletion mutant, showing direct regulation of this cluster by AreA [41]. Additionally, AreA in the maize pathogen *F. verticillioides* is a positive regulator of the toxic SM fumonisin [42].

PacC—pH

Beyond nutritional requirements, another environmental parameter essential for growth and development is pH. The global regulator PacC is a conserved Cis_2His_2 zinc finger transcription factor among fungi [79], capable of regulating a suite of physiological processes, including SM production, in response to environmental pH [80]. Studies investigating PacC regulation of SM have shown that this alkali-activated transcription factor dynamically controls metabolites to be expressed in a pH environment most suitable for compound bioactivity and maximum niche exploitation [81]. In *A. nidulans*, penicillin production was shown to be increased in an alkaline environment [51, 82], possibly as an ecological adaptation by the fungus to outcompete increased bacterial competition in high pH environments [81]. Also in *A. nidulans*, and in contrast to penicillin production, ST synthesis was repressed by alkaline pH as was AF synthesis in *A. parasiticus* [52]. Expression of another important β (beta)-lactam, cephalosporin, by *A. chrysogenum* was shown to be regulated by PacC through binding to promoter regions of structural genes [53]. Expression of the mycelial pigment bikaverin from the rice pathogen *F. fujikuroi* also exhibits PacC regulation through inhibition of its synthesis in a high pH environment [21]. Production of the mycotoxin fumonisin is also downregulated under elevated pH conditions by the maize pathogen *F. verticillioides* [54]. Interestingly, a recent study provides evidence for a dynamic role of PacC in tree fruit spoilage by the phytopathogen *Penicillium expansum* such that mechanisms of acidification after initial colonization, via secretion of organic acids such as gluconic acid, are mediated by PacC to control subsequent expression of the mycotoxin patulin [55]. Like other strains deficient in global regulators of SM, PacC deletion strains exhibit developmental phenotypes such as reduced conidiation [79] or growth inhibitions [21, 54], illustrating a pivotal role for pH sensing and homeostasis in proper growth, development, and SM expression.

CreA—Carbon

When grown in media rich in glucose, filamentous fungi downregulate genes required for metabolizing other carbon sources via a phenomenon called carbon catabolite repression [83], mediated largely by the Cys_2His_2 zinc finger transcription factor CreA [84]. Some examples of CreA-mediated regulation of SM include an overproducing cephalosporin strain of *A. chrysogenum*, obtained by classical mutagenesis techniques, which appeared to have deregulation of the CreA homologue Cre1 compared to wild-type strains [48]. Additionally, putative Cre1 binding sites found within the promoter regions of two cephalosporin biosynthetic genes suggest carbon catabolite repression of this metabolite [49]. In *A. nidulans*, glucose represses antibiotic production independent of a mutated CreA binding site in the penicillin biosynthetic gene *ipnA* promoter region [82] and was still unaffected by a CreA mutant background [85]. Although the role of CreA in regulating SM is less

clear-cut compared to AreA and PacC, it is important to acknowledge the role of this global regulator in environmental sensing and the concomitant physiological response and consider how SMs might be affected. For example, the stringency of glucose repression, likely mediated by CreA, on a cryptic SM cluster in *A. terreus* was recently shown to be insurmountable even by overexpression of a putative in-cluster transcription factor [47].

Velvet Complex—Light

Like most organisms, filamentous fungi sense and respond to light. Conserved throughout the filamentous and dimorphic ascomycetes and possibly basidiomycetes [86] is the light-sensing heterotrimeric velvet complex consisting of LaeA, and the velvet proteins VeA and VelB [87]. The velvet complex links sexual development with SM production in response to light and accomplishes this through tightly regulated spatial compartmentalization of velvet complex components. VelB has nuclear and cytoplasmic localization regardless of illumination status [87], whereas VeA is cytoplasmic under light conditions and migrates to the nucleus in the absence of light [88] as a heterodimer with VelB, and LaeA is constitutively nuclear [89]. Consequently, the three velvet complex proteins are only colocalized to the nucleus under dark conditions, allowing formation of the fully functional heterotrimeric velvet complex. The mechanism of velvet complex regulation of SMs is best studied in *A. nidulans* but increasingly well known in other fungi [90–97]. Once assembled in the nucleus of *A. nidulans*, the velvet complex drives sexual development and production of SMs, whereas these processes are repressed under illuminating conditions resulting from a dissociated velvet complex [81]. Consequently, when localization of members of this complex is compromised, such as with VeA via interacting with the newly described LaeA-like methyltransferase LlmF, neither sexual development nor SM production can proceed normally [98]. The LlmF mechanism of VeA control appears conserved in other ascomycetes as well [99].

The finding that LaeA exhibited some similarity to methyltransferases involved in histone modification, coupled with the observation that SM clusters were most frequently found in subtelomeric regions of the genome [9], led to the hypothesis that SM clusters could be under epigenetic regulation [100]. This indeed has proved to be the case and for detailed coverage of histone modifications and their influence on SM production, we direct the reader to Chap. 3 for chromatin-based regulation of secondary metabolism.

CBC—Iron

Beyond the velvet complex, which is required for coordinating sexual development and SM production in response to light, another well-characterized global regulatory complex in filamentous fungi is the CCAAT-binding complex (CBC). Unlike velvet

components, which are only conserved among filamentous fungi, CBC complexes are found in all eukaryotic organisms [101]. AnCF, the CBC in *A. nidulans* formerly known as PENR1 [101] is composed of the proteins HapB, HapC, and HapE. The AnCF is a positive regulator of genes required for penicillin biosynthesis including *ipnA* and *aata* [102], is a negative autoregulator of *hapB* [103], and coordinates physiological processes in response to cellular redox state [104] and environmental iron [105]. With respect to iron depletion, CBC-mediated upregulation of *sidC* in *A. nidulans* is dependent upon physical interaction with the bZIP protein HapX [105] with SidC being a core enzyme for synthesis of ferricrocin, an SM siderophore that is essential for intracellular iron homeostasis and morphological development [61].

Conclusion

The study of SM in fungi originated primarily for two reasons: (1) to understand the regulation of mycotoxin gene clusters with the goal of using this knowledge to ameliorate the deleterious costs of crop contamination with mycotoxins and (2) to identify compounds with novel bioactivities applicable for pharmaceutical, medicinal, and agricultural uses. Since then, affecting all aspects of fungal biology, the integrated and critical roles of SMs have emerged as being greater than anticipated. SM regulators originally characterized as cluster-specific transcription factors have later been shown to have regulatory functions beyond their native cluster. For example, the canonical AF/ST regulator and $Zn(II)_2Cys_6$ transcription factor *aflR* was shown in *A. parasiticus* to regulate several genes beyond the defined boundaries of the cluster [64, 106]. Additionally, the Cys_2Hys_2 transcription factor Tri6 from *F. graminearum* was shown to regulate 192 genes, ranging from central metabolism to virulence, beyond the known targets within the trichothecene gene cluster [107], leading the authors to propose characterizing Tri6 as a global regulator much like the other known global Cys_2Hys_2 regulators AreA, PacC, and CreA.

Beyond expanding the roles of well-established regulators, research aimed toward elucidating new biosynthetic pathways and compounds has also benefited from exploiting the tight interplay of primary and SM. In one particular study, carbon catabolite repression was so stringent over a cryptic cluster, the authors utilized a suite of growth media containing various carbon sources to define nutrient conditions favorable for cluster activation, eventually identifying the compounds hydroisoflavipucin and dihydroisoflavipucin [47].

One of the greatest challenges in studying fungal SM will be to make sense of how key regulators fit into an ever-expanding network of global interactions linking primary and secondary metabolism with growth and development and how this modulates in response to nutrient availability, biotic and abiotic stresses, and niche exploitation. Taken together, it is apparent that ongoing and future studies will necessarily approach fungal SMs not from the standpoint that they are “accessory” molecules, but rather integrated members of a complex metabolic network affecting every facet of fungal biology.

References

1. Cho Y, Srivastava A, Ohm R, Lawrence CB, Wang K-H, Grigoriev IV et al (2012) Transcription factor Amr1 induces melanin biosynthesis and suppresses virulence in *Alternaria brassicicola*. *PLoS Pathog* 8:e1002974
2. Ortiz SC, Trienens M, Rohlf M (2013) Induced fungal resistance to insect grazing: reciprocal fitness consequences and fungal gene expression in the *Drosophila-Aspergillus* model system. *PLoS ONE* 8:e74951
3. Mousa WK, Raizada MN (2013) The diversity of anti-microbial secondary metabolites produced by fungal endophytes: an interdisciplinary perspective. *Front Microbiol* 4:1–18
4. Subramani R, Kumar R, Prasad P, Aalbersberg W (2013) Cytotoxic and antibacterial substances against multi-drug resistant pathogens from marine sponge symbiont: citrinin, a secondary metabolite of *Penicillium* sp. *Asian Pac J Trop Biomed* 3:291–296
5. Robens J, Cardwell K (2003) The costs of mycotoxin management to the USA: management of aflatoxins in the United States. *Toxin reviews. J Toxicol-Toxin Rev* 22:139–152
6. Bryden WL (2012) Mycotoxin contamination of the feed supply chain: implications for animal productivity and feed security. *Anim Feed Sci Technol* 173:134–158
7. Gieseke KE and Centers for Disease Control and Prevention (2004) Outbreak of aflatoxin poisoning—eastern and central provinces, Kenya, January–July 2004. *Public Health Fac Publ* 53:790–792
8. Hoffmeister D, Keller NP (2006) Natural products of filamentous fungi: enzymes, genes, and their regulation. *Nat Prod Rep* 24:393–416
9. Palmer JM, Keller NP (2010) Secondary metabolism in fungi: does chromosomal location matter? *Curr Opin Microbiol* 13:431–436
10. Khaldi N, Collemare J, Lebrun M-H, Wolfe KH (2008) Evidence for horizontal transfer of a secondary metabolite gene cluster between fungi. *Genome Biol* 9(1):1–10
11. Schmitt I, Lumbsch HT (2009) Ancient horizontal gene transfer from bacteria enhances biosynthetic capabilities of fungi. *PLoS ONE* 4:e4437
12. Slot JC, Rokas A (2011) Horizontal transfer of a large and highly toxic secondary metabolic gene cluster between fungi. *Curr Biol* 21:134–139
13. Calvo AM, Wilson RA, Bok JW, Keller NP (2002) Relationship between secondary metabolism and fungal development. *Microbiol Mol Biol Rev* 66:447–459
14. Andersen MR, Nielsen JB, Klitgaard A, Petersen LM, Zachariassen M, Hansen TJ, Blicher LH, Gotfredsen CH, Larsen TO, Nielsen KF, Mortensen UH (2013) Accurate prediction of secondary metabolite gene clusters in filamentous fungi. *Proc Natl Acad Sci USA* 110:E99–E107
15. Yin W, Keller NP (2011) Transcriptional regulatory elements in fungal secondary metabolism. *J Microbiol* 49:329–339
16. Brown DW, Yu JH, Kelkar HS, Fernandes M, Nesbitt TC, Keller NP, Adams TH, Leonard TJ (1996) Twenty-five coregulated transcripts define a sterigmatocystin gene cluster in *Aspergillus nidulans*. *Proc Natl Acad Sci USA* 93:1418–1422
17. Woloshuk CP, Foutz KR, Brewer JF, Bhatnagar D, Cleveland TE, Payne GA (1994) Molecular characterization of aflR, a regulatory locus for aflatoxin biosynthesis. *Appl Environ Microbiol* 60:2408–2414
18. Chang P-K, Ehrlich KC, Yu J, Bhatnagar D, Cleveland TE (1995) Increased expression of *Aspergillus parasiticus* aflR, encoding a sequence-specific DNA-binding protein, relieves nitrate inhibition of aflatoxin biosynthesis. *Appl Environ Microb* 61:2372–2377
19. Chiang Y-M, Szweczyk E, Davidson AD, Keller N, Oakley BR, Wang CCC (2009) A gene cluster containing two fungal polyketide synthases encodes the biosynthetic pathway for a polyketide, asperfuranone, in *Aspergillus nidulans*. *J Am Chem Soc* 131:2965–2970
20. Bergmann S, Schümman J, Scherlach K, Lange C, Brakhage AA, Hertweck C (2007) Genomics-driven discovery of PKS-NRPS hybrid metabolites from *Aspergillus nidulans*. *Nat Chem Biol* 3:213–217

21. Wiemann P, Willmann A, Straeten M, Kleigrew K, Beyer M, Humpf H-U, Tudzynski B (2009) Biosynthesis of the red pigment bikaverin in *Fusarium fujikuroi*: genes, their function and regulation. *Mol Microbiol* 72:931–946
22. Chen H, Lee M-H, Daub ME, Chung K-R (2007) Molecular analysis of the cercosporin biosynthetic gene cluster in *Cercospora nicotianae*. *Mol Microbiol* 64:755–770
23. Shimizu T, Kinoshita H, Nihira T (2007) Identification and in vivo functional analysis by gene disruption of *ctnA*, an activator gene involved in citrinin biosynthesis in *Monascus purpureus*. *Appl Environ Microbiol* 73(16):5097–5103
24. Wight WD, Kim K-H, Lawrence CB, Walton JD (2009) Biosynthesis and role in virulence of the histone deacetylase inhibitor depudecin from *Alternaria brassicicola*. *Mol Plant Microbe Interact* 22:1258–1267
25. Wiemann P, Guo CJ, Palmer JM, Sekonyela R, Wang CCC, Keller NP (2013) Prototype of an intertwined secondary metabolite supercluster. *Proc Natl Acad Sci USA* 110(42):17065–17070
26. Brown DW, Butchko R, Busman M, Proctor R (2007) The *Fusarium verticillioides* FUM gene cluster encodes a Zn(II)2Cys6 protein that affects FUM gene expression and fumonisin production. *Eukaryot Cell* 6:1210–1218
27. Kim J-E, Jin J, Kim H, Kim J-C, Yun S-H, Lee Y-W (2006) GIP2, a putative transcription factor that regulates the aurofusarin biosynthetic gene cluster in *Gibberella zeae*. *Appl Environ Microbiol* 72:1645–1652
28. Bok JW, Chung D, Balajee SA, Marr K, Andes D, Nielsen KF, Frisvad JC, Kirby KA, Keller NP (2006b) GliZ, a transcriptional regulator of gliotoxin biosynthesis, contributes to *Aspergillus fumigatus* virulence. *Infect Immun* 74:6761–6768
29. Kennedy J, Auclair K, Kendrew SG, Park C, Vederas JC, Hutchinson CR (1999) Modulation of polyketide synthase activity by accessory proteins during lovastatin biosynthesis. *Science* 284:1368–1372
30. Chiang Y-M, Szewczyk E, Davidson AD, Entwistle R, Keller NP, Wang CCC, Oakley BR (2010) Characterization of the *Aspergillus nidulans* monodictyphenone gene cluster. *Appl Environ Microbiol* 76:2067–2074
31. Abe Y, Ono C, Hosobuchi M, Yoshikawa H (2002b) Functional analysis of *mlcR*, a regulatory gene for ML-236B (compactin) biosynthesis in *Penicillium citrinum*. *Mol Genet Genomics* 268:352–361
32. Chen Y-P, Yuan G-F, Hsieh S-Y, Lin Y-S, Wang W-Y, Liaw L-L et al (2010) Identification of the *mokH* gene encoding transcription factor for the upregulation of monacolin K biosynthesis in *Monascus pilosus*. *J Agric Food Chem* 58:287–293
33. Tanaka A, Tsuge T (2000) Structural and functional complexity of the genomic region controlling AK-toxin biosynthesis and pathogenicity in the Japanese pear pathotype of *Alternaria alternata*. *Mol Plant Microbe Interact* 13:975–986
34. Fox EM, Gardiner DM, Keller NP, Howlett BJ (2008) A Zn(II)2Cys6 DNA binding protein regulates the sirodesmin PL biosynthetic gene cluster in *Leptosphaeria maculans*. *Fungal Genet Biol* 45:671–682
35. Flaherty JE, Woloshuk CP (2004) Regulation of fumonisin biosynthesis in *Fusarium verticillioides* by a zinc binuclear cluster-type gene, ZFRI. *Appl Environ Microbiol* 70:2653–2659
36. MacPherson S, Laroche M, Turcotte B (2006) A fungal family of transcriptional regulators: the zinc cluster proteins. *Microbiol Mol Biol Rev* 70:583–604
37. Yu J-H, Butchko RAE, Fernandes M, Keller NP, Leonard TJ, Adams TH (1996) Conservation of structure and function of the aflatoxin regulatory gene *aflR* from *Aspergillus nidulans* and *A. flavus*. *Curr Genet* 29:549–555
38. Ahuja M, Chiang Y-M, Chang S-L, Praseuth MB, Entwistle R, Sanchez JF, Lo H-C, Yeh H-H, Oakley BR, Wang CCC (2012) Illuminating the diversity of aromatic polyketide synthases in *Aspergillus nidulans*. *J Am Chem Soc* 134:8212–8221
39. Abe Y, Suzuki T, Ono C, Iwamoto K, Hosobuchi M, Yoshikawa H (2002a) Molecular cloning and characterization of an ML-236B (compactin) biosynthetic gene cluster in *Penicillium citrinum*. *Mol Genet Genomics* 267:636–646

40. Chang P-K, Yu J, Bhatnagar D, Cleveland TE (2000) Characterization of the *Aspergillus parasiticus* major nitrogen regulatory gene, *areA*. *Biochim Biophys Acta* 1491:263–266
41. Mihan M, Homann V, Liu T-WD, Tudzynski B (2003) *AREA* directly mediates nitrogen regulation of gibberellin biosynthesis in *Gibberella fujikuroi*, but its activity is not affected by NMR. *Mol Microbiol* 47:975–991
42. Kim H, Woloshuk CP (2008) Role of *AREA*, a regulator of nitrogen metabolism, during colonization of maize kernels and fumonisin biosynthesis in *Fusarium verticillioides*. *Fungal Genet Biol* 45:947–953
43. Kihara J, Moriwaki A, Tanaka N, Tanaka C, Ueno M, Arase S (2008) Characterization of the *BMR1* gene encoding a transcription factor for melanin biosynthesis genes in the phytopathogenic fungus *Bipolaris oryzae*. *FEMS Microbiol Lett* 281:221–227
44. Simon A, Dalmais B, Morgant G, Viaud M (2013) Screening of a *Botrytis cinerea* one-hybrid library reveals a *Cys2His2* transcription factor involved in the regulation of secondary metabolism gene clusters. *Fungal Genet Biol* 52:9–19
45. Eliahu N, Igbaria A, Rose MS, Horwitz BA, Lev S (2007) Melanin biosynthesis in the maize pathogen *Cochliobolus heterostrophus* depends on two mitogen-activated protein kinases, *Chk1* and *Mps1*, and the transcription factor *Cmr1*. *Eukaryot Cell* 6:421–429
46. Tsuji G, Kenmochi Y, Takano Y, Sweigard J, Farrall L, Furusawa I, Horino O, Kubo Y (2000) Novel fungal transcriptional activators, *Cmr1p* of *Colletotrichum lagenarium* and *pig1p* of *Magnaporthe grisea*, contain *Cys2His2* zinc finger and *Zn(II)2Cys6* binuclear cluster DNA-binding motifs and regulate transcription of melanin biosynthesis genes in a developmentally specific manner. *Mol Microbiol* 38:940–954
47. Gressler M, Zaehle C, Scherlach K, Hertweck C, Brock M (2011) Multifactorial induction of an orphan PKS-NRPS gene cluster in *Aspergillus terreus*. *Chem* 18:198–209
48. Jekosch K, Kück U (2000a) Glucose dependent transcriptional expression of the *cre1* gene in *Acremonium chrysogenum* strains showing different levels of cephalosporin C production. *Curr Genet* 37:388–395
49. Jekosch K, Kück U (2000b) Loss of glucose repression in an *Acremonium chrysogenum* beta-lactam producer strain and its restoration by multiple copies of the *cre1* gene. *Appl Microbiol Biotechnol* 54:556–563
50. Trapp SC, Hohn TM, McCormick S, Jarvis BB (1998) Characterization of the gene cluster for biosynthesis of macrocyclic trichothecenes in *Myrothecium roridum*. *Mol Gen Genet* 257:421–432
51. Bergh KT, Brakhage AA (1998) Regulation of the *Aspergillus nidulans* penicillin biosynthesis gene *acvA* (*pcbAB*) by amino acids: implication for involvement of transcription factor *PACC*. *Appl Environ Microbiol* 64:843–849
52. Keller NP, Nesbitt C, Sarr B, Phillips TD, Burow GB (1997) pH regulation of sterigmatocystin and aflatoxin biosynthesis in *Aspergillus* spp. *Phytopathology* 87:643–648
53. Schmitt EK, Kempken R, Kuck U (2001) Functional analysis of promoter sequences of cephalosporin C biosynthesis genes from *Acremonium chrysogenum*: specific DNA-protein interactions and characterization of the transcription factor *PACC*. *Mol Genet Genomics* 265:508–518
54. Flaherty JE, Pirttilä AM, Bluhm BH, Woloshuk CP (2003) *PAC1*, a pH-regulatory gene from *Fusarium verticillioides*. *Appl Environ Microbiol* 69:5222–5227
55. Barad S, Horowitz S, Kobiler I, Sherman A, Prusky DB (2013) Accumulation of the mycotoxin patulin in the presence of gluconic acid contributes to pathogenicity of *Penicillium expansum*. *Mol Plant-Microbe Interact* 27:66–77
56. Malapi-Wight M, Smith J, Campbell J, Bluhm BH, Shim W-B (2013) *Sda1*, a *Cys2-His2* zinc finger transcription factor, is involved in polyol metabolism and fumonisin B1 production in *Fusarium verticillioides*. *PLoS ONE* 8:e67656
57. Bergmann S, Funk AN, Scherlach K, Schroeckh V, Shelest E, Horn U, Hertweck C, Brakhage AA (2010) Activation of a silent fungal polyketide biosynthesis pathway through regulatory cross talk with a cryptic nonribosomal peptide synthetase gene cluster. *Appl Environ Microbiol* 76(24):8143–8149

58. Hohn TM, Krishna R, Proctor RH (1999) Characterization of a transcriptional activator controlling trichothecene toxin biosynthesis. *Fungal Genet Biol* 26:224–235
59. Reverberi M, Gazzetti K, Punelli F, Scarpari M, Zjalic S, Ricelli A, Fabbri AA, Fanelli C (2012) Aoyap1 regulates OTA synthesis by controlling cell redox balance in *Aspergillus ochraceus*. *Appl Microbiol Biotechnol* 95:1293–1304
60. Roze LV, Chanda A, Wee J, Awad D, Linz JE (2011) Stress-related transcription factor AtfB integrates secondary metabolism with oxidative stress response in aspergilli. *J Biol Chem* 286:35137–35148
61. Eisendle M, Schrettl M, Kragl C, Müller D, Illmer P, Haas H (2006) The intracellular siderophore ferricrocin is involved in iron storage, oxidative-stress resistance, germination, and sexual development in *Aspergillus nidulans*. *Eukaryot Cell* 5:1596–1603
62. Wagner D, Schmeinck A, Mos M, Morozov IY, Caddick MX, Tudzynski B (2010) The bZIP transcription factor MeaB mediates nitrogen metabolite repression at specific loci. *Eukaryot Cell* 9:1588–1601
63. Amaike S, Affeldt KJ, Yin W-B, Franke S, Choithani A, Keller NP (2013) The bZIP protein MeaB mediates virulence attributes in *Aspergillus flavus*. *PLoS ONE* 8:e74030
64. Yin W-B, Amaike S, Wohlbach DJ, Gasch AP, Chiang Y-M, Wang CCC, Bok JW, Rohlf M, Keller NP (2012) An *Aspergillus nidulans* bZIP response pathway hardwired for defensive secondary metabolism operates through aflR. *Mol Microbiol* 83:1024–1034
65. Sekonyela R, Palmer JM, Bok J-W, Jain S, Berthier E, Forseth R, Schroeder F, Keller NP (2013) RsmA regulates *Aspergillus fumigatus* gliotoxin cluster metabolites including cyclo(L-Phe-L-Ser), a potential new diagnostic marker for invasive aspergillosis. *PLoS ONE* 8:e62591
66. Bussink HJ, Clark A, Oliver R (2001) The *Cladosporium fulvum* Bap1 gene: evidence for a novel class of Yap-related transcription factors with ankyrin repeats in phytopathogenic fungi. *Eur J plant Pathol* 107:655–659
67. Pedley KF, Walton JD (2001) Regulation of cyclic peptide biosynthesis in a plant pathogenic fungus by a novel transcription factor. *Proc Natl Acad Sci USA* 98:14174–14179
68. Schmitt EK, Hoff B, Kück U (2004) AcFKH1, a novel member of the forkhead family, associates with the RFX transcription factor CPC1 in the cephalosporin C-producing fungus *Acremonium chrysogenum*. *Gene* 342:269–281
69. Schmitt EK, Kück U (2000) The fungal CPC1 protein, which binds specifically to β -lactam biosynthesis genes, is related to human regulatory factor X transcription factors. *J Biol Chem* 275:9348–9357
70. Atanasova L, Knox BP, Kubicek CP, Druzhinina IS, Baker SE (2013) The polyketide synthase gene pks4 of *Trichoderma reesei* provides pigmentation and stress resistance. *Eukaryot Cell* 12:1499–1508
71. Hong S-Y, Roze LV, Linz JE (2013) Oxidative stress-related transcription factors in the regulation of secondary metabolism. *Toxins* 5:683–702
72. Temme N, Oeser B, Massaroli M, Heller J, Simon A, Collado IG, Viaud M, Tudzynski P (2012) BcAtf1, a global regulator, controls various differentiation processes and phytotoxin production in *Botrytis cinerea*. *Mol Plant Pathol* 13:704–718
73. Ahn JH, Walton JD (1998) Regulation of cyclic peptide biosynthesis and pathogenicity in *Cochliobolus carbonum* by TOXE_p, a novel protein with a bZIP basic DNA-binding motif and four ankyrin repeats. *Mol Gen Genet* 260:462–469
74. Gajiwala KS, Burley SK (2000) Winged helix proteins. *Curr Opin Struct Biol* 10:110–116
75. Hoff B, Schmitt EK, Kück U (2005) CPC1, but not its interacting transcription factor AcFKH1, controls fungal arthrospore formation in *Acremonium chrysogenum*. *Mol Microbiol* 56:1220–1233
76. Bayram O, Braus GH (2012) Coordination of secondary metabolism and development in fungi: the velvet family of regulatory proteins. *FEMS Microbiol Rev* 36(1):1–24
77. Feng GH, Leonard TJ (1998) Culture conditions control expression of the genes for aflatoxin and sterigmatocystin biosynthesis in *Aspergillus parasiticus* and *A. nidulans*. *Appl Environ Microbiol* 64:2275–2277

78. Wilson R, Arst H (1998) Mutational analysis of AREA, a transcriptional activator mediating nitrogen metabolite repression in *Aspergillus nidulans* and a member of the “streetwise” GATA family. *Microbiol Mol Biol Rev* 62:586–596
79. Tilburn J, Sarkar S, Widdick D, Espeso E, Orejas M, Mungroo J, Peñalva MA, Arst HN Jr (1995) The *Aspergillus* PacC zinc finger transcription factor mediates regulation of both acid- and alkaline-expressed genes by ambient pH. *EMBO J* 14:779–790
80. Trushina N, Levin M, Mukherjee PK, Horwitz BA (2013 Jan) PacC and pH-dependent transcriptome of the mycotrophic fungus *Trichoderma virens*. *BMC Genomics* 14(138):1–21
81. Brakhage AA (2013) Regulation of fungal secondary metabolism. *Nat Rev Microbiol* 11:21–32
82. Espeso EA, Tilburn J, Arst HN, Peñalva MA (1993) pH regulation is a major determinant in expression of a fungal penicillin biosynthetic gene. *EMBO J* 12:3947–3956
83. Ronne H (1995) Glucose repression in fungi. *Trends Genet* 11:12–17
84. Dowzer CEA, Kelly JM (1991) Analysis of the *creA* gene, a regulator of carbon catabolite repression in *Aspergillus nidulans*. *Mol Cell Biol* 11:5701–5709
85. Brakhage AA, Browne P, Turner G (1992) Regulation of *Aspergillus nidulans* penicillin biosynthesis and penicillin biosynthesis genes *acvA* and *ipnA* by glucose. *J Bacteriol* 174:3789–3799
86. Ni M, Yu J-H (2007) A novel regulator couples sporogenesis and trehalose biogenesis in *Aspergillus nidulans*. *PLoS ONE* 2:e970
87. Bayram O, Krappmann S, Ni M, Bok JW, Helmstaedt K, Valerius O, Braus-Stronmeyer S, Kwon N-J, Keller NP, Yu J-H, Braus GH (2008) VelB/VeA/LaeA complex coordinates light signal with fungal development and secondary metabolism. *Science* 320:1504–1506
88. Stinnett SM, Espeso E, Cobeño L, Araújo-Bazán L, Calvo AM (2007) *Aspergillus nidulans* VeA subcellular localization is dependent on the importin alpha carrier and on light. *Mol Microbiol* 63:242–255
89. Bok JW, Keller NP (2004) *LaeA*, a regulator of secondary metabolism in *Aspergillus* spp. *Eukaryot Cell* 3:527–535
90. Chettri P, Calvo AM, Cary JW, Dhingra S, Guo Y, McDougal RL, Bradshaw RE (2012) The *veA* gene of the pine needle pathogen *Dothistroma septosporum* regulates sporulation and secondary metabolism. *Fungal Genet Biol* 49:141–151
91. Hoff B, Kamerewerd J, Sigl C, Mitterbauer R, Zadra I, Kürnsteiner H, Kück U (2010) Two components of a velvet-like complex control hyphal morphogenesis, conidiophore development, and penicillin biosynthesis in *Penicillium chrysogenum*. *Eukaryot Cell* 9:1236–1250
92. Kopke K, Hoff B, Bloemendal S, Katschorowski A, Kamerewerd J, Kück U (2013) Members of the *Penicillium chrysogenum* velvet complex play functionally opposing roles in the regulation of penicillin biosynthesis and conidiation. *Eukaryot Cell* 12:299–310
93. López-Berges MS, Hera C, Sulyok M, Schäfer K, Capilla J, Guarro J, Di Pietro A (2013) The velvet complex governs mycotoxin production and virulence of *Fusarium oxysporum* on plant and mammalian hosts. *Mol Microbiol* 87:49–65
94. Veiga T, Nijland JG, Driessen AJM, Bovenberg RAL, Touw H, van den Berg MA, Pronk JT, Daran J-M (2012) Impact of velvet complex on transcriptome and penicillin G production in glucose-limited chemostat cultures of a β -lactam high-producing *Penicillium chrysogenum* strain. *OMICS* 16:320–333
95. Wiemann P, Brown DW, Kleigrewe K, Bok JW, Keller NP, Humpf H-U, Tudzynski B (2011) FfVel1 and FfLae1, components of a velvet-like complex in *Fusarium fujikuroi*, affect differentiation, secondary metabolism and virulence. *Mol Microbiol* 77:972–994
96. Wu D, Oide S, Zhang N, Choi MY, Turgeon BG (2012) ChLae1 and ChVel1 regulate T-toxin production, virulence, oxidative stress response, and development of the maize pathogen *Cochliobolus heterostrophus*. *PLoS Pathog* 8(2):e1002542
97. Yang Q, Chen Y, Ma Z (2013) Involvement of BcVeA and BcVelB in regulating conidiation, pigmentation and virulence in *Botrytis cinerea*. *Fungal Genet Biol* 50:63–71
98. Palmer JM, Theisen JM, Duran RM, Grayburn WS, Calvo AM, Keller NP (2013) Secondary metabolism and development is mediated by LlmF control of VeA subcellular localization in *Aspergillus nidulans*. *PLoS Genet* 9:e1003193

99. Bi Q, Wu D, Zhu X, Gillian Turgeon B (2013) Cochliobolus heterostrophus Llm1— a Lae1-like methyltransferase regulates T-toxin production, virulence, and development. *Fungal Genet Biol* 51:21–33
100. Bok JW, Noordermeer D, Kale SP, Keller NP (2006a) Secondary metabolic gene cluster silencing in *Aspergillus nidulans*. *Mol Microbiol* 61:1636–1645
101. Brakhage AA, Andrianopoulos A, Kato M, Steidl S, Davis MA, Tsukagoshi N, Hynes MJ (1999) HAP-like CCAAT-binding complexes in filamentous fungi: implications for biotechnology. *Fungal Genet Biol* 27:243–252
102. Brakhage AA, Thön M, Spröte P, Scharf DH, Al-Abdallah Q, Wolke SM, Hortschansky P (2009) Aspects on evolution of fungal β -lactam biosynthesis gene clusters and recruitment of trans-acting factors. *Phytochemistry* 70:1801–1811
103. Steidl S, Hynes MJ, Brakhage AA (2001) The *Aspergillus nidulans* multimeric CCAAT binding complex AnCF is negatively autoregulated via its hapB subunit gene. *J Mol Biol* 306:643–653
104. Thön M, Abdallah Q A, Hortschansky P, Scharf DH, Eisendle M, Haas H, Brakhage AA (2010) The CCAAT-binding complex coordinates the oxidative stress response in eukaryotes. *Nucleic Acids Res* 38:1098–1113
105. Hortschansky P, Eisendle M, Al-Abdallah Q, Schmidt AD, Bergmann S, Thön M, Kniemeyer O, Abt B, Seeber B, Werner ER, Kato M, Brakhage AA, Haas H (2007) Interaction of HapX with the CCAAT-binding complex—a novel mechanism of gene regulation by iron. *EMBO J* 26:3157–3168
106. Price MS, Yu J, Nierman WC, Kim HS, Pritchard B, Jacobus CA, Bhatnagar D, Cleveland TE, Payne GA (2006) The aflatoxin pathway regulator AflR induces gene transcription inside and outside of the aflatoxin biosynthetic cluster. *FEMS Microbiol Lett* 255:275–279
107. Nasmith CG, Walkowiak S, Wang L, Leung WWY, Gong Y, Johnston A, Harris LJ, Guttman DS, Subramaniam R (2011) Tri6 is a global transcription regulator in the phytopathogen *Fusarium graminearum*. *PLoS Pathog* 7:e1002266