

Chapter 1

Overview of Basic Immunology and Translational Relevance for Clinical Investigators



Betty Stephen and Joud Hajjar

Abstract Tumor exists as a complex network of structures with an ability to evolve and evade the host immune surveillance mechanism. The immune milieu which includes macrophages, dendritic cells, natural killer cells, neutrophils, mast cells, B cells, and T cells are found in the core, the invasive margin, or the adjacent stromal or lymphoid component of the tumor. The immune infiltrate is heterogeneous and varies within a patient and between patients of the same tumor histology. The location, density, functionality, and the crosstalk between the immune cells in the tumor microenvironment influence the nature of immune response, prognosis, and treatment outcomes in cancer patients. Therefore, an understanding of the characteristics of the immune cells and their role in tumor immune surveillance is of paramount importance to identify immune targets and to develop novel immune therapeutics in the war against cancer. In this chapter we provide an overview of the individual components of the human immune system and the translational relevance of predictive biomarkers.

Keywords Adaptive · Biomarkers · CTLA-4 · Immune checkpoints · Immunology · Immunotherapy · Innate · PD-1 · PD-L1 · Resistance · Response · T cells · Translational

The human immune system is an elaborate and dynamic network of cells that work together to defend the human body against attacks by foreign agents including malignant cells. There are two levels of immunity, the innate immunity and the adaptive immunity. The innate immunity constitutes the first line of defense against pathogens, which includes the anatomic and physiologic barriers, phagocytic leukocytes, dendritic cells (DC), natural killer (NK) cells, and the circulating plasma proteins [1]. Elie Metchnikoff, a pathologist and Father of natural immunity, was

B. Stephen

The University of Texas MD Anderson Cancer Center, Houston, TX, USA

e-mail: BAStephen@mdanderson.org

J. Hajjar (✉)

Baylor College of Medicine and Texas Children's Hospital, Houston, TX, USA

e-mail: joud.hajjar@bcm.edu

© Springer Nature Switzerland AG 2018

A. Naing, J. Hajjar (eds.), *Immunotherapy*, Advances in Experimental Medicine and Biology 995, https://doi.org/10.1007/978-3-030-02505-2_1

the first to describe the concept of leukocyte recruitment and phagocytosis of microorganisms [2]. The adaptive immune system is a more versatile mechanism of defense provided by the B lymphocytes and the T lymphocytes, which has been attributed to Paul Ehrlich, the physicist who described the side-chain theory of antibody formation [3]. The innate and adaptive immune systems are distinct but interactive components of the human immune system that collectively contribute to the defense operations against foreign proteins [4]. In this chapter we will discuss the fundamental components of the immune system and their development, how innate immunity interfaces with adaptive immune responses to eliminate tumor cells, and the development of immunotherapeutic strategies to combat cancer.

Innate Immune System

An association between inflammation and tumorigenesis has long been described, but has been established with turn of the century [5]. The human body is constantly exposed to a highly diverse world of foreign proteins every day, which are rapidly eliminated in a normal healthy individual by the components of the innate immune system. Speed is the essence of innate immune response; however, they are non-specific in nature, of limited duration, and lack immunologic memory [6]. Traditionally, the cellular components of the innate immune system, which includes the macrophages, neutrophils, eosinophils, basophils, mast cells, NK cells, and DCs, are associated with elimination of microbial agents and activation of the more efficient, antigen-specific adaptive immune response in the event of failure [4, 6]. And, the humoral elements of the innate immune system that includes the complement proteins and C-reactive protein are considered as a regulator of inflammatory process [4]. However, accumulating evidence suggests that the innate and adaptive immune system, triggered by the tumor antigens, play a significant role in the recognition and elimination of malignant cells as well [7]. In the process, several noxious reactive chemicals, cytokines, and chemokines are released, which damages the surrounding healthy tissue [8]. The inflammatory microenvironment also induces genomic instability and enhances rate of molecular alterations [9]. The resultant process of repeated cell renewal and proliferation sets the stage for chronic inflammation that produces a microenvironment conducive for malignant transformation of cells [10]. For this reason tumors are sometimes described as “wounds that do not heal” [11].

Cellular Components of the Innate Immune System

All the cells of the immune system originate from the pluripotent hematopoietic stem cells (HSCs) in the bone marrow. The HSCs divide to produce the common lymphoid progenitor (CLP) and the common myeloid progenitor (CMP) cells. The

CLP give rise to the T and B lymphocytes that are responsible for adaptive immunity, and the NK cells; while, the CMP give rise to the cells of the innate immune system, leukocytes (neutrophils, monocytes, basophils, and eosinophils), mast cells, DCs, erythrocytes, and the megakaryocytes.

Leukocytes

The primary function of the leukocytes is to protect the body against invading microorganisms. However, microenvironmental factors at the site of inflammation produces substantial changes in the phenotype and functional status of individual cells that favor initiation and progression of tumor [12, 13].

Neutrophils

They account for 50–70% of circulating leukocytes [14] and form the indispensable first line of defense against pathogenic microorganisms. They originate from the CMP cells in the bone marrow in response to several cytokines including granulocyte colony stimulating factor (G-CSF) and granulocyte macrophage colony stimulating factor (GM-CSF) [14, 15]. They circulate in the blood as dormant cells and are recruited to sites of infection by specific chemokines, cytokines, and cell adhesion molecules [16]. The microbes are then taken up by the process of phagocytosis and destroyed by high concentrations of microbicidal granules or by respiratory burst associated with production of highly toxic reactive oxygen species in the pathogen-containing vacuole [14]. In addition, the activated neutrophils upregulates the production of cytokines [including tumor necrosis factor- α , interleukin (IL)-1 β , IL-1R α , IL-12, and vascular endothelial growth factor (VEGF)] and chemokines (including IL-8) critical for chemotaxis and recruitment of additional neutrophils, macrophages, and T cells [17, 18].

Beyond the classical role of professional phagocytes, neutrophils play a significant role in tumor biology [1, 19]. Neutrophils are recruited to the tumor microenvironment (TME) through local production of chemokines such as IL-8, macrophage inflammatory protein-1 α (MIP-1 α /CCL3), and human granulocyte chemotactic protein-2 (huGCP-2/CXCL6) [20]. Tumor-associated neutrophils (TANs) are markedly different from naïve neutrophils. TANs exhibit dual conflicting roles at the molecular level [20]. They either take up an anti-tumorigenic (N1) or a pro-tumorigenic (N2) phenotype [14, 21]. In untreated tumors, the regulatory cytokine transforming growth factor-beta (TGF- β) in the tumor cells drives the differentiation of TANs towards N2 phenotype [13]. These neutrophils locally produce neutrophil elastase (ELA2) [22], oncostatin M [23], and alarmins S100A8/9 [24] that promotes proliferation, survival, metastasis, and resistance of tumor cells to chemotherapy. In addition, N2 TANs promote immunosuppression and tumor progression by releasing growth-stimulating signals, angiogenic factors, and matrix-degrading enzymes [13, 20, 25]. Neutrophils thus assume multiple roles in development and progression

of tumor cells [26]. However, under certain conditions such as TGF- β blockade, TANs assume a N1 phenotype, which are more cytotoxic due to enhanced expression of immune activating cytokines and chemokines, and lower levels of arginase [13]. N1 TANs also communicate with DCs to trigger an adaptive immune response [27]. In addition, they facilitate intratumoral CD8+ T cell infiltration and activation through production of chemokines (like CCL3, CXCL9, and CXCL10) and pro-inflammatory cytokines (i.e., IL-12, TNF- α , GM-CSF, and VEGF) [28]. This phenotype has the potential to inhibit progression of the tumor, indicating the possibility of immunostimulation through TGF- β blockade [13].

Monocytes and Macrophages

Monocytes are derived from the CMP cells. They are large, mononuclear cells that account for 5–7% of circulating leukocytes. These monocytes migrate into the tissues, where they differentiate rapidly and mature into distinct macrophages depending on tissue of activation, the Langerhans cells in the epidermis, Kupffer cells in the liver, and microglial cells in the central nervous system [29]. Macrophages perform many functions. Primarily, they engulf and destroy the invading microorganisms. They also release cytokines and chemokines to recruit other cells of the immune system to the site of inflammation. Macrophages also induce expression of co-stimulatory molecules on the antigen presenting cells (APCs) to initiate adaptive immune response and help in the disposal of pathogens destroyed by adaptive immune response [2].

Similar to TANs, monocytes are attracted to the TME by tumor-derived chemokines such as CCL2, CCL5, CCL7, CCL8 or cytokines such as VEGF, platelet derived growth factor (PDGF), TGF- β , GM-CSF, and M-CSF [30–33], where they differentiate into tissue-resident macrophages [34]. The tumor-associated macrophages (TAMs) assume either anti-tumorigenic M1 phenotype (classically activated) or pro-tumorigenic M2 phenotype (alternatively activated) reflecting the functional plastic nature of these cells [35]. The cytokine profile of the TME plays a central role in the phenotype orientation of the differentiating macrophages [36]. In general, M-CSF, TGF- β , and IL-10, the principal cytokines present in the TME strongly inhibits IL-12 production and NF- κ B activation in TAMs [37]. This skews the differentiation of monocytes to macrophages M2 phenotype, characterized by IL-12^{low} IL-10^{high} [30, 38]. These macrophages migrate to hypoxic areas within the tumor and promote tumor progression by inducing angiogenesis through expression of factors such as VEGF, angiopoietins, pro-angiogenic cytokines, and IL-1; remodeling of stromal matrix by producing a variety of matrix metalloproteinases (MMP) such as MMP1 and MMP9; and by suppressing adaptive immunity through production of prostaglandins, IL-4, IL-6, IL-10, TGF- β and indoleamine 2,3-dioxygenase (IDO) metabolites, and induction of T regulatory (Treg) cells [33, 38]. This enables the tumor cells to escape into surrounding stroma and ultimately metastasize to distant sites. However, classical macrophage activation occurs under certain conditions, for

example in the presence of GM-CSF, microbial products, lipopolysaccharides, or interferon- γ (IFN- γ), where TAMs are educated to assume the more cytotoxic, antigen presenting, IL-12^{high} IL-10^{low} M1 phenotype [33]. They kill microbes and tumor cells by producing copious amounts of pro-inflammatory cytokines such as IL-12 and IL-23, toxic intermediates—nitric oxide, reactive oxygen intermediates (ROI), and TNF [30, 33]. The cytokines also initiate T-helper 1 (Th1) adaptive immunity. Though high macrophage content is often correlated with poor patient prognosis in breast [39, 40], bladder [41], endometrial [42], and cervical cancers [43], TAMs in tumor tissue confer survival advantage to patients with prostate cancer [44] and colon cancer [45]. Pharmacological skewing of macrophage polarization from M2 to M1 phenotype is likely to provide therapeutic benefit to cancer patients.

Eosinophils

Eosinophils are derived from the CMP cells and they constitute less than 5% of circulating leukocytes [2, 46]. Traditionally, eosinophils are associated with host defense against large, multicellular parasitic helminths and fungi with allergic conditions [47]. Eosinophils express a number of receptors such as chemokine receptors, cytokine receptors, immunoglobulin (Ig) receptors, Toll-like pattern recognition receptors, and histamine receptors [48]. Engagement of these receptors causes the release of highly cytotoxic proteins, such as major basic protein, eosinophil-derived neurotoxin or eosinophil peroxidase, pro-inflammatory cytokines and growth factors (IL-2, -3, -4, -5, -6, -10, -12, and -13, IFN- γ , TNF- α , GM-CSF, TGF- α/β), chemokines, including RANTES(CCL5), eotaxin-1 (CCL11), CXCL5, and lipid mediators (platelet-activating factor and leukotriene C4) from the large, highly cytotoxic, secretory cytoplasmic granules at the sites of allergic inflammation [48, 49].

In addition, eosinophils are found in the tumor infiltrating area [1]. Tumor-associated tissue eosinophilia has been associated with improved patient outcomes in a variety of solid tumors including colorectal cancer [50], oral squamous cell carcinoma (SCC) [51], laryngeal and bladder carcinoma [52]. Though an understanding of the function of eosinophils in cancer has remained elusive, it has become apparent that eosinophils express major histocompatibility complex (MHC) class II and co-stimulatory molecules [CD40, CD28/86, cytotoxic T lymphocyte associated protein 4 (CTLA-4)] [53, 54], whereby they function as APCs and initiate antigen-specific immune responses by the T cells [55]. Kinetic studies have demonstrated that chemotactic factors such as eotaxins and damage-associated molecular patterns (DAMPs), high mobility group box 1 (HMGB1) released by necrotic tumor cells, preferentially induce eosinophilic migration to tumors [56, 57] prior to infiltration by CD8+ T cells [58]. Tumor-associated tissue eosinophils in its active form release chemokines such as CCL5, CXCL9, and CXCL10 that attracts CD8+ T cells to the tumor [59]. Tumor-associated tissue eosinophilia in the presence of tumor-specific CD8+ T cells produces significant changes in the TME such as polarization of TAM to M1 phenotype and vascular normalization of the tumor, resulting in increased T cell infiltration, enhanced tumor rejection, and improved patient survival [58].

Basophils

They originate from the CMP cell in the bone marrow and are released into circulation as mature cells [2]. They account for less than 1% of circulating leukocytes and were therefore considered redundant to mast cells functionally till about 15 years ago [60]. Basophils travel to the sites of allergic inflammation and microbial assault in response to cytokines and chemokines released locally [60]. IgE-mediated activation of basophils induces proliferation and rapid release of several inflammatory mediators such as histamine, leukotriene C4, prostaglandins, and significant amount of IL-4 and IL-13 [61]. IL-4 and IL-13, released within an hour of stimulation, serve as chemoattractants for other immune cells and direct the differentiation of naïve T cells towards Th2 phenotype resulting in Th2-(allergic)-type immune responses in an IgE-dependent and IgE-independent manner [62, 63]. Further, basophils express CD40 ligand, which on binding with CD40 on B cell, induces transformation of B cells to plasma cells and promotes production of IgE antibodies [63].

Though the role of basophils in tumorigenesis has not been clearly understood, it is believed that basophils promote neoplastic angiogenesis [64]. Basophils express angiopoietin-1 and angiopoietin-2 messenger RNAs in the cytoplasmic vacuoles, and VEGFR-2 and Tie1 receptors on the cell surface. And, activation of basophils releases pro-angiogenic factors VEGF-A and VEGF-B through a crosstalk between the basophils and the mast cells, contributing to neoplastic angiogenesis. Further, the correlation between basophils in the tumor draining lymph node with Th2 inflammation in patients with pancreatic ductal adenocarcinomas and the emergence of basophils as an independent prognostic factor of poor survival after surgery suggests a role for basophils in tumor development and disease recurrence [65].

Mast Cells

Mast cells are tissue-based inflammatory cells of hematopoietic origin [66]. The origin of mast cell has long been debated. Recently Qi et al. identified pre-basophil and mast cell progenitors (pre-BMP), a population of granulocyte-macrophage progenitors (GMPs) with a capacity to differentiate into basophils and mast cells while still retaining a limited capacity to differentiate into myeloid cells [67]. The pre-BMPs circulate in the blood and reach the peripheral tissue, where they get differentiated into basophils and mast cells in the presence of mutually exclusive transcription factors, C/EBP α and MITF, respectively [67]. Basophils and mast cells share many characteristics such as expression of IgE receptors, presence of same granules, and secretion of similar mediators of immune response and cytokines when stimulated. Both offer protection against parasites and are key players in the Th2-(allergic)-type immune responses [68, 69]. However, mast cells show marked differences in their histochemical, biochemical, and functional characteristics based on their phenotype and the cytokine milieu, a phenomenon called “mast cell heterogeneity” [70]. Mast cells express several surface receptors including KIT IgG

receptor, and Toll-like receptors (TLRs) [70]. The characteristic feature of mast cells is the presence of dense metachromatic granules in the cytoplasm containing histamine and heparin which are explosively released on contact with allergens [71]. Tissue mast cells besides being the largest storehouse of histamine, with the exception of gastrointestinal tract and central nervous system, also contain several preformed mediators such as heparin, serotonin, tryptases, and chymases; lipid mediators; cytokines such as TNF- α/β , IFN- α/β , IL-1 α/β , IL-5, -6, -13, -16, and -18; chemokines such as IL-8 (CXCL8), I-309 (CCL1), MCP-1 (CCL2), MIP-1 α S (CCL3), MIP1 β (CCL4), MCP-3 (CCL7), RANTES (CCL5), eotaxin (CCL11), MCAF (MCP-1); and growth factors such as SCF, M-CSF, GM-CSF, bFGF, VEGF, NGF, and PDGF [71], which are synthesized and rapidly released on activation by IgE- or IgG-dependent mechanisms. Strategic location of the mast cells at the interface between mucosal and environmental surfaces, for example, near blood vessels, nerves, glands, and beneath epithelial surfaces [68, 70], and their ability to store TNF- α in a preformed state allows mast cells to orchestrate the first response to invading pathogens [66]. Different stimuli activate different pathways resulting in different cocktail of molecules released by mast cells, which significantly influences T cell differentiation and the subsequent adaptive immune response [66].

Increased numbers of mast cells found in many tumors may have a double-edged function in tumor development. Infiltration of tumor by mast cells has been associated with poor prognosis in some cancers such as prostate cancer [72], lip cancer [73], and diffuse large B cell lymphoma [74]. This may be because intratumoral mast cells, which are a rich source of pro-angiogenic and tumor growth stimulatory mediators, stimulate or modulate angiogenesis and peritumoral mast cells, which are rich sources of tryptase and chymase, promote extracellular matrix degradation and tumor invasion, resulting in tumor progression [73, 75, 76]. On the contrary, mast cell infiltration has been associated with good prognosis in breast [77], ovarian [78], lung [79], and colorectal cancers [80]. This is due to release of several antitumoral factors by stromal mast cells including cytotoxic endogenous peroxidase, cytokines like IL-1, IL-4, IL-6, and TNF- α that induces apoptosis of endothelial cells, chymase, which inhibits angiogenesis, and tryptase leading to tumor fibrosis [78, 81, 82]. It is therefore evident that the density and location of mast cells within the tumor samples and the crosstalk between mast cells and stromal cells are better predictors of patient survival as they modulate the immune response [1].

Dendritic Cells

DCs are professional APCs that are resident in most tissues of the body and concentrated in the secondary lymphoid tissues [83]. In the steady state, they originate from the monocyte and dendritic cell progenitor (MDP) derived from the CMP cells in the bone marrow [84]. The MDPs give rise to monocytes and common DC progenitors (CDPs) in the bone marrow [85]. The CDPs give rise to pre-DCs, which migrate from the bone marrow through the blood to lymphoid and non-lymphoid tissues, where they differentiate to produce conventional DCs (cDCs). The pre-DCs

lack the form and function of DCs, but with microbial or inflammatory stimuli they develop into DCs [86]. Plasmacytoid DCs is an example of pre-DCs found in blood, thymus, bone marrow, and secondary lymphoid tissue, which produce type I IFN- α in response to viral exposure. The cDCs are broadly classified into migratory DCs and lymphoid tissue-resident DCs. The migratory DCs (Langerhans cells and dermal DCs) are immature DCs present in the peripheral tissue, which are very effective in capturing antigens. They sample the environment using several receptors including the TLRs and (NOD)-like receptors (NLRs). On encountering a pathogen, endocytosis is upregulated transiently to facilitate accumulation of large quantities of antigens by the immature DCs that are phagocytic and macropinocytic in the peripheral tissue [3]. Immature DCs are relatively inefficient in presenting the peptide-MHC complexes at the surface due to reduced formation of antigenic peptides [3], ubiquitination of MHC class II molecules in the lysosomes, and poor expression of co-stimulatory ligands (CD80, CD86) [3, 87]. Shortly thereafter, functional maturation of DCs ensues triggering the antigen presenting machinery, which is the critical link between innate and adaptive immunity [88]. Endocytosis by the DCs decreases and expression of MHC-I, MHC-II, and co-stimulatory molecules increases at the surface possibly due to cessation of ubiquitination of MHC class II molecules [87]. As a result, the mature DCs degrade the pathogen and present the antigenic peptides on MHC Class I or II molecules on the cell surface to naïve T cells, express co-stimulatory ligands (CD80, CD86) simultaneously, and migrate to the T cell zones of the lymphoid tissue [3]. Binding of the ligands to the co-stimulatory molecules on T cells leads to activation of T cells [87]. Based on the type of pathogen and other maturation signals received, the activated T cells are educated to proliferate and differentiate to become potent effector cytotoxic T cells or helper T cells [3]. DCs can also directly present the intact antigen to and activate the antigen-specific B cells [3]. The lymphoid tissue-resident DCs (CD8+ and CD8- splenic cDCs and thymic cDCs) are immature DCs uniquely located in regions where naïve T cells are activated [87]. They present the antigens in the lymphoid organ to the T cells [86]. They are likely responsible for maintaining peripheral tolerance in the steady state. Under inflammatory conditions, some DCs may arise from the CLP cells and from the monocytes [2]. An example of inflammatory DC is the tumor necrosis factor- and inducible nitric-oxide synthase-producing DCs (Tip DCs) [86].

Under normal conditions, DCs are responsible for maintaining immune tolerance to host cells [3] DCs are generally phenotypically and functionally immature in the steady state. Immature state is characterized by ubiquitination and intracellular accumulation of MHC class II molecules and low levels of co-stimulatory molecules [83]. Therefore in the absence of infections, though DCs continuously present self-antigens and nonpathogenic environmental antigens to T cells, this induces the production of Tregs instead of effector T cells. In the development of cancer, where the tumor cells are more similar to normal cells, DCs are therefore more likely to induce peripheral tolerance in the absence of inflammation. Further, other mechanisms of immune suppression such as expression of PD-L1 and PD-L2, TGF- β , and IDO inhibit DC and T cell function and facilitate escape of tumor cells

from immune recognition. This may explain why vaccines did not succeed as an effective treatment modality in cancer patients [3]. DCs are aptly called the gatekeepers of the immune system because of their ability to inspect the microenvironment, interpret the cues in the environment, and instruct the immune cells to respond quickly and appropriately between tolerogenic and immunogenic function [83].

Natural Killer Cells

NK cells are the most powerful lymphocytes of the innate immune system with robust cytotoxic activity. They originate from the CLP cells in the bone marrow and account for 15% of all the circulating lymphocytes [1]. Besides, they are located in many peripheral tissues. Though NK cells do not express antigen-specific surface receptors such as the classical membrane-bound Igs of B cells or the T cell receptor (TCR) of the T cell, they express a wide range of activating and inhibitory cell surface receptors. As the primary function of NK cells is to identify and eliminate cells that fail to produce self MHC class I molecules, NK cells during the process of maturation are educated to identify “missing self” through the expression of several cell surface inhibitory receptors such as killer cell inhibitory receptor-L (KIR-L), which specifically binds with MHC class I ligands [89]. Engagement of these receptors by cognate MHC class I ligands constitutively expressed in normal cells in steady state conditions ensures self-tolerance by transducing inhibitory signals [90]. It is the absence of these MHC class I ligands on tumor cells and cells in distress as in viral infection that marks them for destruction by NK cells [89].

The effector function of NK cells is triggered by the engagement of cell surface activating receptors including the potent NKG2D receptor, killer-cell Ig-like receptors (KIR-S), TLR, and NLR that identifies non-self-infected cells and self-cells under stress by recognizing pathogen-associated molecular patterns (PAMPs) [91]. However, activation of the NK cells is dependent on cellular crosstalk with accessory cells such as DCs, neutrophils, macrophages, and mast cells, and/or a cytokine microenvironment that includes IL-2, IFN- α/β , IL-12, IL-15, IL-18, or IL-21 [92, 93]. The DCs, which are key partners to NK cells, lie in close proximity to the NK cells and prime the NK cells either directly by contact or by secretion of the cytokines, IFN- α , IL-2, IL-12, IL-15, or IL-18 [94]. Activated NK cells induce cytotoxicity and/or promote cytokine production [94]. NK cells kill tumor cells by releasing cytoplasmic granules containing perforin and granzymes or by expressing Fas ligand (CD95) or TNF- α -related apoptosis-inducing ligand (TRAIL) that binds with death receptors on the tumor cells triggering apoptosis [95]. Tumor cells however evolve and evade destruction by NK cells [95]. A common escape mechanism used by tumor cells is the proteolytic shedding of NKG2D ligands [96]. Further, chronic stimulation of NKG2D pathway by tumor-associated expression of TGF- β and NKG2D ligands (including MHC class I homologues MICA and MICB) on the surface of tumor cells can functionally impair NKG2D pathway by inducing endocytosis and destruction of the potent activating NKG2D receptors on NK cells [97, 98]. This result in markedly reduced expression of NKG2D on NK cells, which

promotes T cell silencing and evasion of immune surveillance by tumor cells. Nevertheless, NK cells prosecute tumor cells through other mechanisms such as antibody-dependent cell cytotoxicity [99]. NK cells express other activating receptors such as CD16, Fc- γ receptor IIIa (FCGR3A), which binds to the Fc region of Ig [100]. This enables the NK cells to identify antibody-coated tumor cells and destroys them by releasing perforins.

At least two functional subsets of NK cells have been described based on the expression of CD56 and CD16 [101]. The CD56^{dim} CD16⁺ NK cells account for 90% of circulatory NK cells. These cells are attracted to peripheral tissues by several chemokines. They express perforin, natural cytotoxicity receptors (NCR), and KIRs. On activation, the CD56^{dim} CD16⁺ NK cells are more cytotoxic and secrete low levels of cytokines. On the other hand, CD56^{bright} CD16⁻ NK cells are primarily located in the secondary lymphoid tissue and account for less than 10% of circulatory NK cells. They lack perforin, NCR, and KIRs. On activation by IL-2, the CD56^{bright} CD16⁻ NK cells produce cytokines, mainly IFN- γ , GM-CSF, and TNF- α . However, on prolonged stimulation by IL-2, they express perforin, NCR, and KIRs, and acquire cytotoxic function.

Though NK cells are traditionally characterized as cells of innate immunity, they also exhibit T cell characteristics and are capable of mounting rapid and robust immune response on secondary exposure [102]. The immune memory function of NK cells lasts for several months after the initial exposure, is antigen-specific, and transferable to naïve animals [102]. Though NK cells are potent killers with immune memory, only modest success in clinical setting has been achieved as their effectiveness has been hampered by their limited ability to infiltrate tumor cells [103].

Adaptive Immune System

The hallmark of adaptive immunity, mediated by the T lymphocytes (T cells) and B lymphocytes (B cells), is the specificity of the immune response to antigenic stimuli. Another unique feature of adaptive immunity is its ability to confer lasting immunological memory that results in more rapid and robust immune response with subsequent exposure to the same antigen [2]. Contrary to innate immune response, which is immediate in onset due to the presence of germline-encoded cell surface receptors, the adaptive immune response is a slower process, as the lymphocytes on activation undergo clonal expansion to attain sufficient numbers before the effector cells mount an immune response [29]. There are two classes of adaptive immune response, the humoral and cell-mediated. The humoral immune response is mediated by the B lymphocytes against antigens present outside the cells, in the blood and body fluids. On the other hand, the cell-mediated immune response is mediated by the T lymphocytes against intracellular pathogens presented as small antigenic determinants on MHC molecules.

Cellular Components of the Adaptive Immune System

The T and B lymphocytes originate from the CLP, a specialized type of stem cell originating from the pluripotent HSCs [2].

T Lymphocytes

The lymphoid progenitor cells migrate from the bone marrow to the thymus, where they undergo four stages of differentiation and proliferation, including developmental checkpoints to ensure that cells which fail to recognize antigen-MHC complexes or distinguish self-antigens do not mature [104]. As the lymphoid progenitor cells migrate through the cortex, they undergo an education program based on the constant interaction with the thymic epithelial cells [105]. The lymphoid progenitor cells that enter the thymus at the cortico-medullary junction do not express TCR, or CD4 or CD8 co-receptors and are therefore called CD4/CD8 double-negative (DN) lymphocytes (DN1) [106]. As they move through the cortex from the cortico-medullary junction to the capsule, the lymphoid progenitor cells lose their ability to form B cells or NK cells and become committed T cell precursors (DN2) [107]. Following T lineage commitment and expression of recombination-activating gene 1 (RAG1), the TCR- β chain is rearranged and paired with the pre-T α chain, resulting in expression of pre-TCRs (DN3) [104]. Subsequently, intense proliferation results in generation of multiple thymocytes (DN4). With appropriate cytokine stimulation, they express CD8 co-receptors first and then CD4 co-receptors to become double-positive (DP) thymocytes. This is accompanied by rearrangements in the TCR α chain, which results in generation of complete $\alpha\beta$ TCRs. Then, DP thymocytes interact with TECs and further development into naïve T cells is dependent on their ability to bind with MHC class I or class II molecules associated with self-peptides (positive selection) [104, 108]. Approximately 90% of DP thymocytes express TCRs that fail to bind with MHC molecules, resulting in delayed apoptosis of these cells (death by neglect). Based on their interaction with MHC molecules, the DP thymocytes differentiate into single positive T cell by silencing of the transcription of one co-receptor locus [105, 109].

In the medulla, T cells are screened for reactivity against wide range of tissue-specific proteins including self-peptides expressed by the thymic medullary epithelial cells [29]. The T cells that express TCRs with high affinity for self-peptides undergo rapid apoptosis and are later cleared by thymic macrophages (negative selection). T cells that express intermediate level of TCR signaling enter into a maturation phase by the process of positive selection. The T cells that express TCRs that bind with MHC Class I molecule mature into a single positive CD8 mature T cell (CD8+ T cell), while those that express TCRs that bind with MHC Class II molecule mature into a single positive CD4 mature T cell (CD4+ T cell). These naïve T cells then sample the environment in the medulla for antigen presenting DCs. On exposure to antigenic determinants presented by the APCs, the T cells are

activated in the presence of co-stimulation of CD28 by B7 molecules (CD80 and CD86) on the APCs, to form effector T cells that either destroy the pathogenic agent or attract other immune cells to the site. In the absence of antigenic stimuli in the medulla, the naïve T cells enter the blood stream and travel to the peripheral lymphoid tissue and enter the paracortical region of the LN. In the tumor draining LNs, naïve T cells are activated on encountering tumor antigen in the context of MHC molecule and co-stimulation of the constitutively expressed CD28 on the surface of T cells by B7 proteins (CD80 or CD86) expressed on the same APC [110]. This results in clonal expansion and differentiation of naïve T cells in the lymph nodes into effector T cells (CD4+ helper T cells or CD8+ cytotoxic T cells). Depending on the cytokine milieu and the transcription factors in the TME, the CD4+ helper T cells differentiate into several subtypes that includes Th1 [111], T-helper 2 (Th2) [112], T-helper 17 (Th17) [113], induced Tregs (iTregs) [114], follicular helper T cell (Tfh) [115], and T-helper 9 (Th9) [116]. These helper T cells secrete cytokines and chemokines that regulate the immune response. Th1 cells favor cell-mediated immunity by activation of CD8 T cells to mount an immune response against intracellular pathogens, while Th2 cells favor humoral immunity by activation of B cells against extracellular parasites. On the other hand, CD8+ effector T cells activated by antigen presentation on the MHC class I molecule or through CD4 helper T cells are directly cytotoxic. Hence, they migrate to the tumor and destroy the tumor cells. In addition, some of the activated T cells and B cells differentiate into memory cells that are responsible for the long-lasting immunological memory [117]. Subsequent exposure to the same antigen results in more rapid and robust immune response.

Regulation of T cell response is a delicate balance between co-stimulatory and inhibitory signals that serve as immune checkpoints. Co-stimulatory receptors include CD28, inducible T cell co-stimulator (ICOS), 4-1BB (CD-137), OX40 (CD-134), and glucocorticoid-induced TNFR-related protein (GITR), while CTLA-4, programmed cell death 1 (PD-1), lymphocyte activation gene-3 (Lag-3), T cell immunoglobulin-3 (Tim-3), and T cell immunoreceptor with immunoglobulin and ITIM domain (TIGIT) are co-inhibitory [118]. CD28 is the primary co-stimulatory molecule constitutively expressed on the surface of naïve T cells. On ligand binding with B7-1 and B7-2 on APCs, they provide the essential co-stimulatory signal for T cell activation and downstream signaling [119]. Besides CD28, there are other co-signaling receptors of the TNF receptor superfamily including 4-1BB [120], OX40 [121], and GITR [122] that synergize with TCR signaling to promote cytokine production and T cell survival. The stimulatory effect of T cells is counterbalanced by a suppressive mechanism in order to maintain immune homeostasis. Activated T cells simultaneously express CTLA-4 and PD-1 on their surface as immune checkpoints [123–125]. CTLA-4, a CD28 homologue with a higher affinity to bind with B7 molecules, is an early co-inhibitory signal that regulates T cell activity during the priming phase. On engagement with B7, CTLA-4 blocks CD28 co-stimulation and abrogates T cell activity and cytokine production. On the other hand, PD-1, a CD28 family member, is a late co-inhibitory signal that regulates T cell activity during the effector phase in the peripheral tissue. PD-1 interacts with two ligands, PD-L1 and PD-L2. PD-L1 is expressed on many cells including the tumor cells,

activated B and T cells in response to IFN- γ produced by the activated T cells, while PD-L2 is expressed exclusively on macrophages and DCs [126]. Unlike CTLA-4, the PD-1 to PD-L1 ligand binding does not interfere with co-stimulation, but down-regulates B and T cell proliferation and cytokine production by interfering with signaling pathways downstream of TCRs and B cell receptors (BCRs) [127]. Besides CTLA-4 and PD-1, there are other next generation co-inhibitory receptors such as Lag-3, Tim-3, and TIGIT, which are expressed on distinct lymphocyte subsets that are responsible for differential suppression of immune response [128]. For example, Tim-3 pathway may regulate immune responses in the gut, while TIGIT may regulate in the lungs and Lag-3 in the pancreas. Similarly, they exhibit functional specification in that TIGIT may selectively suppress pro-inflammatory response of Th1 and Th17 cells, while promoting Th2 cell response [129]. Besides immune checkpoints, a chief contributor to this immunosuppressive effect is the Tregs, which are specialized T cells that suppress the cytotoxic function of other T cells [130]. They are classified as thymus-derived natural Tregs (nTregs) and peripherally derived inducible Tregs (iTregs). nTregs characterized by surface expression of the CD4 and CD25 antigens and by the nuclear expression of forkhead box P3 (FOXP3) are positively selected thymocytes with relatively high affinity for self-antigens presented on MHC class II molecules. On the contrary, iTregs differentiate from naïve CD4 T cells in the periphery in the presence of TGF- β . They exert their immunosuppressive action by the expression of immunosuppressive cytokines such as IL10 and TGF- β [114]. Decreasing the activity of Treg cells enhances both innate and adaptive immune response, which can be utilized to treat cancer [131]. Thus, under normal conditions, coordinated regulation of immune activation and suppressive pathways play an important role in the maintenance of peripheral tolerance and regulation of the amplitude and duration of T cell responses [132].

B Lymphocytes

The B cells develop from the HSCs in the liver during fetal life and continue in the bone marrow in adult life [2]. The four subsets of B cell precursors that develop from the lymphoid progenitor cells, pre-pro-B cells, early pro-B cells, late pro-B cells, and pre-B cells are devoid of surface Ig [133]. In the presence of RAG 1 and 2, these cells constantly interact with the bone marrow stromal cells that provide critical growth factors, chemokines, and cytokines for B cell development. The B cell precursors undergo sequential rearrangement of the genes encoding for the heavy chain (H) [134]. The DJ rearrangement occurs in the early pro-B cells followed by VDJ rearrangements in the late pro-B cells resulting in the formation of a large pre-B cell with a complete Ig μ heavy chain in the cytoplasm [2]. The μ heavy chain combines with the surrogate light chain (L) and two invariant accessory chains Ig α and Ig β to form the pre-BCR, which is transiently expressed on the surface of pre-B cells, positively selecting these cells for further development. This initiates a negative feedback loop by which it shuts down RAG expression, halts the H gene

rearrangement in the pre-B cell, prevents the rearrangement of the second H (allelic exclusion), and signals the proliferation of pre-B cells. The RAG genes are re-expressed, which induces rearrangement of the genes encoding the L in positively selected pre-B cells that leads to formation of an immature B cell with the expression of a complete IgM BCR on the surface of the cell. This triggers the cessation of L gene rearrangement. As a vast repertoire of BCRs capable of recognizing a huge diversity of antigens including self-antigens are developed, the immature B cells are tested for reactivity to autoantigens before leaving the bone marrow. When immature B cells express a non-auto-reactive BCR with optimal downstream signaling, RAG expression is downregulated, which allows for positive selection of these cells to enter the spleen as transitional B cells. Whereas, immature B cells that express a non-auto-reactive BCR with low basal BCR signaling insufficient to downregulate RAG expression and immature B cells that are strongly self-reactive are negatively selected for elimination by apoptosis (clonal deletion). Alternatively, these cells may be inactivated (anergy) or may undergo receptor editing, a process by which secondary rearrangement of L leads to formation of new BCRs that are not self-reactive, which allows for subsequent positive selection of these cells for further development [135].

The immature B cells enter the spleen as transitional cells. Very few cells progress from T1 to T2 stage as most of the T1 cells undergo clonal deletion or anergy due to strong reactivity to self-antigens that are expressed only in the peripheral tissue [136]. And, the transition from T1 to T2 cell is dependent on basal tonic BCR signaling. The T2 cells receive pro-survival signals through B cell-activating factor (BAFF)-R and differentiate into naïve B cell expressing both IgM and IgG surface receptors. Guided by the strength of BCR signal, naïve B cell differentiates into either follicular (FO) B cells with intermediate BCR signals and expression of bruton tyrosine kinase, or marginal zone (MZ) B cell with weak BCR signal and expression of NOTCH2 [136, 137]. The MZ B cells located within the splenic white pulp are resting mature B cells that do not circulate. They have limited antigen specificity and are activated by non-protein antigens such as common blood-borne pathogens independent of T cells. On activation, they rapidly develop into short-lived plasma cells secreting low affinity IgM antibodies and do not produce memory cells. The FO B cells that circulate between the blood and the spleen are located adjacent to T cell-rich areas in secondary lymphoid organs and are activated by foreign proteins in a T cell-dependent manner [138]. The antigens bound to membrane-bound Ig are internalized by FO B cells and presented on MHC class II molecules to the CD4 helper T cells. The activated T cells express CD40L, a co-stimulatory molecule, and other cytokines required for B cell activation [2]. The activated B cells undergo clonal expansion to differentiate into plasma cells that produce large amounts of high affinity secreted antibody. Some of the activated B cells migrate into the lymphoid follicle to form a germinal center, where they undergo extensive proliferation, Ig class switching, and somatic hypermutation to generate long-lived plasma cells or memory B cells. These plasma cells leave the germinal center and migrate to the bone marrow, where they continue to produce antibodies even after elimination of the antigens. On reinfection, these circulating

antibodies provide immediate protection and activate the memory cells located in the peripheral lymphoid tissue.

Immunoglobulins

Immunoglobulins are Y-shaped heterodimers composed of two identical L chains and two identical H chains [139]. The two H chains are attached to each other by multiple disulfide bonds and each L chain is attached to an H chain by a disulfide bond. Each L and H chain is divided into a variable and constant region. The variable region in each L and H chain has three complementarity determining regions (CDRs). The three CDRs in one L chain pairs with the three CDRs in the H chain in each arm of the Y to form a paratope, the antigen binding site. Each paratope is specific for an epitope of the antigen, which determines the specificity of the Ig. The constant region of the H chain is identical for all the Igs of the same class, but different between classes. So also, all the Igs in a class have either λ or κ L chains. Proteolytic digestion with papain divides the Ig into three functional units, two antigen binding fragments (Fab) and the crystallizable fragment (Fc). Each Fab fragment contains a complete L chain and one variable and one constant domain of H chain, which includes the antigen binding site. The Fc fragment contains two constant domains of the H chain. This is the effector domain of the Ig which activates the NK cells, classical complement pathway, and phagocytosis [140].

Based on the amino acid sequences in the constant region of the H chains, human antibodies are classified as IgM, IgD, IgG, IgE, and IgA [139]. Accordingly, they have diverse biologic functions. IgM is the earliest antibody expressed on the surface during B cell development and it is the major class of Ig that is secreted on first exposure to the antigen. IgG is the major antibody in the blood that is produced in large quantities during secondary immune response and is responsible for clearance of opsonized pathogens and neutralization of toxins and viruses. IgA is the principal antibody in body secretions and contributes to nearly 50% of protein content in colostrum and protects mucosal surfaces from toxins, virus, and bacteria. Membrane-bound IgD is expressed in small amounts when the immature B cells leave the bone marrow and they regulate the cell's activation. IgE is found in trace amounts in the blood, but it is a very potent Ig expressed during hypersensitivity or allergic reactions and parasitic infestations.

Each B cell in the body produces only one kind of antibody [140]. When a naïve B cell is activated, it proliferates and differentiates into a clone of plasma cells, which produces large amount of secreted antibodies that have the same antigen binding site as the BCR that was activated and is specific for a single epitope. Hence they are called monoclonal antibodies (mAb). Polyclonal antibodies are secreted by different B cell clones that bind with different epitopes on the same antigen.

Monoclonal antibodies have revolutionized the use of Igs as a therapeutic agent. However, engineering mAb is not without challenge. The first mAb engineered for human use was a murine antibody [141]. They were highly immunogenic with limited biological efficacy and very short half-life. This limitation was overcome by

genetically engineering human protein formats of mAb. Chimeric mAbs that are 70% human, created by fusing murine variable region with human constant region [142]. Later, humanized mAbs that are 85–90% human, where only the CDRs are murine, were developed [143]. Currently, fully human mAbs produced by phage display are available [144]. The process of humanization has made the mAbs less immunogenic than murine mAbs. As a result, several mAbs that target growth factor receptor [such as epidermal growth factor (cetuximab), human epidermal growth factor receptor 2 (trastuzumab)], TME, and tumor antigens have been approved for treatment of colorectal, breast, and lung cancer [145]. The humanness of mAbs is indicated by the nomenclature. For example, -xi- indicates chimeric mAbs (rituximab), -zu- indicates humanized (bevacizumab), and -u- indicates fully human mAb (ipilimumab).

The Immune System in Action!

Summary of the Immune Responses Against Tumor Cells

In the fight against cancer, greater understanding of the immunoregulatory processes of TME is critical for development of immunotherapy. The TME is composed of a variety of cells such as macrophages, DCs, NK cells, mast cells, naïve lymphocytes, B cells, cytotoxic T cells, helper T cells, memory cells, Tregs, myeloid-derived suppressor cells (MDSCs), and stromal cells [146]. Despite the dynamic interaction between these elements in the TME and the tumor, the cancer cells develop cellular processes to subvert the immune attack and become resilient. Thus a comprehensive understanding of the interactions between the tumor and the elements in the TME will help to identify novel targets and therapeutic strategies to combat resistance to therapy.

The human immune system exhibit a dual role in cancer. Though the primary function of the immune system is to eliminate tumor cells, they also shape immunogenicity and promote tumor progression through a dynamic process called cancer immunoediting [147]. This process includes three distinct phases: elimination, equilibrium, and escape. During the elimination phase (cancer immunosurveillance), the challenge lies in the ability of the immune system to recognize the subtle differences between self and transformed self of the malignant cells [148]. The tumor cells express several danger signals such as NKG2D ligands and surface calreticulin, and produce minor disruptions in the surrounding tissue, resulting in the release of inflammatory signals such as IFN- γ , IFN α/β , TNF, and IL-12, which recruit NK cells, DCs, and macrophages to the tumor site. This results in apoptosis and death of tumor cells. The liberated tumor antigens are then presented by the APCs on MHC molecules to T cells. This initiates tumor-specific adaptive immune response. The cytotoxic T cells interact with the Fas and TRAIL receptors on tumor cells, or secrete granzymes and perforins to induce tumor cell apoptosis. Thus,

innate and adaptive immune cells have the capacity to completely eliminate the tumor cells and halt the immunoediting process.

During the equilibrium phase, there is continuous interaction between the immune cells and tumor cells that have escaped elimination phase. The tumor and the immune cells exist in a state of equilibrium that prevents expansion of the tumor cells. However, this continuous immune pressure selects or promotes the formation of new variants of tumor cells with reduced immunogenicity that escapes recognition by immune system [148]. This is the longest phase in the immunoediting process, when the tumor cell variants reside in a latent form before escaping eventually [149].

During the escape phase, tumor cells adopt several mechanisms to evade immunosurveillance [150]. Tumor cells downregulate expression of tumor antigens or MHC class I molecules to reduce immune recognition and antigen presentation to tumor-specific T cells, preventing activation of T cells. Tumor cells may also upregulate expression of pro-survival growth factors such as EGFR and HER2. In addition, the tumor cells frequently develop a host of immunosuppressive defense mechanisms to escape immune surveillance through a process called immune tolerance [7]. For example, tumor cells may express suppressive surface ligands, PD-L1 or PD-L2, that engage with PD-1 receptors on activated T cells resulting in T cell exhaustion; or release immunosuppressive molecules such as IDO [151]. Under hypoxic conditions, the TME may release VEGF, which suppresses T cell adhesion to tumor endothelium and impedes T cell infiltration of the tumor. Similarly, TAMs in the presence of IL-4, IL-10, and TGF- β may polarize to assume M2 phenotype and express high levels of IL-10 and low levels of IL-12. These macrophages suppress T cell activity and promote angiogenesis and tumor growth [152]. In addition, MDSCs, which are immature innate immune cells in the TME, utilize various mechanisms such as expression of IL-10, TGF- β , and Tregs to produce immune suppression, resulting in tumor progression [153, 154]. As a result, immunologically sculpted tumor cells with increased resistance emerge, resulting in uncontrolled growth of the tumor with overt clinical disease. It is therefore critical to overcome these barriers to elicit clinical response to therapeutic agents.

Cancer Immunotherapy

Immunotherapy has revolutionized cancer treatment due to its ability to produce durable responses in patients with certain types of advanced cancer. Though several immunotherapeutics including IL-2, IFN- α , and Sipuleucel-T vaccine were investigated, only small improvements in efficacy were observed. Several mAbs have also been used in the treatment of cancer [155] based on their ability to inhibit ligand binding and downstream signaling (cetuximab), target the TME (bevacizumab), and target immunosuppressive cytokines (GC-1008, an anti-TGF- β antibody) [156].

But it is the discovery of immune checkpoint CTLA-4 and a deeper understanding of the immune regulatory pathways that led to a major breakthrough in cancer immunotherapy [157]. Subsequent to the discovery that activated T cells express

CTLA-4, which on binding with B7 molecules on the APC blocks co-stimulation of T cells resulting in immune suppression, a series of experiments were performed to unleash the immune harnessing power of T cells to combat cancer. This led to the development of the concept of immune checkpoint blockade and discovery of ipilimumab, a CTLA-4 inhibitor, which produced durable responses in about 20% of patients and considerable improvement in the overall survival (OS) of patients with metastatic melanoma, resulting in FDA approval of the drug in 2011 [158]. The dramatic response with ipilimumab laid the foundation for exploration of other T cell inhibitory pathways. Based on strong preclinical evidence, several clinical trials were conducted to evaluate the efficacy of PD-1/PD-L1 pathway blockade by mAbs [159–163]. As a result of durable responses and survival benefits produced in several tumor types, FDA granted accelerated approval of several immune checkpoint inhibitors (ICPis) as listed in Table 1.1 [164]. This offers proof of concept that checkpoint inhibition provides durable and meaningful response in a subset of patients with responsive tumors.

Besides CTLA-4 and PD-1/PD-L1 signaling pathways, other immune regulatory pathways are being investigated as potential therapeutic targets. IDO is one such immunosuppressive pathway exploited by tumor cells to evade immune surveillance [165]. Several IDO inhibitors are under clinical development including INCB024360 [166, 167], indoximod [168], IDO peptide vaccine [169], BMS-986205 [170], and NLG919 [171]. A robust therapeutic immune response is produced not only by releasing the “brakes” on T cells, but also by stepping on the “gas.” T cell co-stimulation through receptors, like OX40 or 4-1BB, provides a potent “go” signal that actively promotes the optimal “killer” CD8 T cell responses [172]. Several ongoing clinical trials are investigating immune checkpoint agonist therapies as single-agent or in combination with other immunotherapies, chemotherapy, targeted therapy, or radiotherapy.

Despite the success with ICPis (CTLA-4, PD-1/PD-L1 blockade) in various tumor types, many patients are primarily resistant or develop resistance to treatment after an initial period of response [173]. Among several mechanistic approaches being investigated in the clinic to overcome primary and secondary resistance to the ICPis, there is growing evidence that combination therapies are far more effective than monotherapies to combat resistance mechanisms as tumors use multiple pathways to evade immune elimination [174]. Further, as these co-inhibitory receptors have non-redundant signaling pathways, a combined blockade of these mechanistically different pathways may be synergistic in restoring T cell-mediated immune response [128]. Recently, FDA approved nivolumab in combination with ipilimumab for the treatment of patients with BRAF V600 wild-type, unresectable or metastatic melanoma and advanced renal cell carcinoma [164]. There is intense research to identify optimal combinations that would increase the response rate and the duration of response. Targeted therapies are known to produce rapid onset of tumor regression [175]. However, the response is short-lived. On the contrary, immunotherapies produce more durable response; but, it takes longer to initiate tumor regression. Due to their complimentary outcomes, combinations of targeted and immunotherapy are being investigated in several clinical trials and emerging

Table 1.1 FDA-approved immune checkpoint inhibitors and indications^a

Drug	Immune checkpoint(s)	FDA-approved tumor type ^b
Ipilimumab	CTLA-4	Unresectable or metastatic melanoma
Nivolumab	PD-1	Unresectable or metastatic melanoma
		Metastatic non-small cell lung cancer
		Advanced renal cell carcinoma
		Classical Hodgkin lymphoma
		Recurrent or metastatic squamous cell carcinoma of the head and neck
		Locally advanced or metastatic urothelial carcinoma
		Mismatch repair deficient and microsatellite instability high metastatic colorectal cancer
Hepatocellular carcinoma		
Pembrolizumab	PD-1	Unresectable or metastatic melanoma
		PD-L1-positive non-small cell lung cancer
		Recurrent or metastatic squamous cell carcinoma of the head and neck
		Classical Hodgkin lymphoma
		Locally advanced or metastatic urothelial carcinoma
		Unresectable or metastatic microsatellite instability-high or mismatch repair deficient solid tumors
		Recurrent locally advanced or metastatic PD-L1-positive gastric or gastroesophageal junction adenocarcinoma
Atezolizumab	PD-L1	Metastatic urothelial carcinoma
		Metastatic non-small cell lung cancer
Durvalumab	PD-L1	Locally advanced or metastatic urothelial carcinoma
		Unresectable stage III non-small cell lung cancer
Avelumab	PD-L1	Metastatic Merkel cell carcinoma
		Locally advanced or metastatic urothelial carcinoma
Nivolumab in combination with ipilimumab	PD-1 and CTLA-4	Unresectable or metastatic melanoma Advanced renal cell carcinoma

^aList of FDA-approved immune checkpoint inhibitors as of May 15, 2018, adapted from: <https://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm279174.htm>

^bTumor type must meet the criteria listed in the abovementioned website

data suggests that such combinations may potentially be synergistic [176]. Similarly, radiation-induced immunomodulatory changes provide local control and prolong survival, but are insufficient to shift the balance of the immunosuppressive TME to

achieve tumor rejection [177]. To overcome this limitation, clinical studies evaluating the combination of radiotherapy and ICPis are currently underway [178, 179].

Emerging data suggest that activation of innate immune system could break the immunosuppressive dynamics of TME to evoke an effective antitumor immune response. Importantly, this process leads to initiation of adaptive immune response by enhancement of the T cell priming process. TLRs, the most important receptors in innate immunity exhibit dual role in cancer [180]. While some TLRs on cancer cells favor tumor progression [181, 182] and promote resistance to chemotherapy, most TLRs on immune cells serve as sensors [180]. Activation of these TLRs by foreign antigens triggers a cascade of pro-inflammatory reactions that ultimately initiates an adaptive immune response. Thus TLRs have been identified as potential targets and several TLR agonists (TLR3, TLR4, TLR5, and TLR7 agonists) are being investigated for clinical application [183, 184]. Similarly, an endoplasmic-reticulum-membrane protein STING (Stimulator of Interferon Genes) that is highly expressed in the APCs mediates potent antitumor activity by induction of innate immunity and initiation of adaptive immunity [184]. Typically, self DNA is located in the nucleus or mitochondrion, while microbial/tumor-derived DNA is located in the cytoplasm. By virtue of their location, the tumor-derived DNA is identified by several cytosolic DNA sensors triggering activation of STING signaling in the APCs [185]. The resultant downstream signaling through STING pathway results in phosphorylation of interferon regulatory factor 3 (IRF3) and nuclear factor- κ B and subsequent induction of pro-inflammatory molecules, IFN β and cytokines such as TNF, IL-1 β , and IL-6. In the process, IFNs also promote cross-priming of T cells by the DCs resulting in initiation of adaptive immune response [186]. As activation of STING pathway promotes T cell priming and induction of adaptive immune mechanism, several STING agonists as vaccine adjuvants and in combination with other immunomodulators are being investigated [187–189]. Thus strategies that bridge the innate and adaptive immune response may have therapeutic utility.

Translational Relevance

Immunotherapeutic agents have revolutionized the treatment paradigm of patients with advanced cancer. However, significant survival benefit has been observed only in a subset of patients. Biomarker-driven drug development is therefore critical, as it may help physicians to preselect patients who are most likely to derive benefit, and more importantly, allow patients who are less likely to benefit to look for alternate therapies and spare them from avoidable immune-related toxicities and cost of treatment [190]. Some of the important biomarkers of response are:

PD-L1 Expression

Early phase I trials suggest that cell surface expression of PD-L1 on tumor cells in pretreatment tissue samples could serve as biomarker of response to treatment with anti-PD-1/PD-L1 therapies. In a phase I study of MDX-1106, an anti-PD-1 inhibitor, in 39 patients with advanced cancers, tumor biopsies from 9 patients were analyzed for PD-L1 expression by immunohistochemistry (IHC) [159]. Objective response was observed in 3 of 4 patients (75%) with PD-L1-positive tumors, while none of the 5 patients with PD-L1-negative tumors had a response. Similar results were observed in another phase I study of BMS-936558 (nivolumab), an anti-PD1 therapy, in which pretreatment tumor tissue from 42 patients with advanced cancer was analyzed for PD-L1 expression by IHC [191]. Nine of 25 patients (36%) with PD-L1-positive tumors had objective response, while none of the 17 patients with PD-L1-negative tumors had a response indicating the possibility of an association between PD-L1 expression on pretreatment samples and objective response. Recently, FDA approved expression of PD-L1 by IHC using 22C3 pharmDx as a diagnostic test for selecting NSCLC patients for treatment with pembrolizumab [192]. However, PD-L1 expression in pretreatment tumor tissue as an absolute biomarker to predict response to PD-1/PD-L1 pathway inhibitors has been questioned for various reasons. In a phase I study conducted to evaluate the safety and efficacy of MPDL3280A, an anti-PD-L1 inhibitor, objective response rate (ORR) of 46% was reported in patients with high PD-L1 expression on pretreatment immune cells, 17% in patients with moderate PD-L1 expression, 21% in patients with minimal PD-L1 expression, and 13% in patients with absent PD-L1-expression in tumor immune cells [193]. Surprisingly, response to treatment was observed even in patients with PD-L1-negative disease. In addition, the association between response to therapy and PD-L1 status was discordant depending on PD-L1 expression on tumor cells or tumor immune cells. PD-L1 expression on tumor-infiltrating immune cells was significantly associated with response to MPDL3280A ($P = 0.007$), whereas, PD-L1 expression on tumor cells was not significantly associated with response ($P = 0.079$). In addition, in a phase III study, survival benefits were seen in NSCLC patients treated with atezolizumab compared to docetaxel regardless of PD-L1 expression in the tumor or immune cells [194]. There is also marked heterogeneity in PD-L1 expression between samples from the primary and metastatic sites in the same individual [195]. Further, the predictive potential of PD-L1 expression is challenged due to technical issues such as lack of standardized PD-L1 diagnostic assay, use of different PD-L1 antibody clones by multiple immune assays, different staining procedures for IHC staining, different cut-off values and scoring patterns [196]. As a result, there is lack of defined criteria to determine PD-L1-status of the patient. The above findings suggest that though PD-L1 expression in tumor tissue may indicate an increased likelihood of response to treatment with PD-1/PD-L1 inhibitors, it may not be a definitive biomarker to exclude PD-L1-negative patients from therapy [193, 197].

Tumor Infiltrating Lymphocytes (TILs)

There is a broad literature of evidence that infiltration of tumor tissue by T cells, specifically CD8+ T cell density at the invasive tumor edge, is associated with improved survival in patients with melanoma, breast, ovarian, lung, esophageal, gastric, renal cell, colorectal and bladder carcinoma among other solid tumors [198–200]. On the contrary, infiltration of the tumor tissue by Tregs is associated with poor survival in ovarian, breast cancer, hepatocellular carcinoma [201–203]. Interestingly, strong intratumoral infiltration by CD8+ T cells and Th1 cells did not favor immune elimination of tumors in patients with mismatch repair-deficient colorectal cancer [204]. Despite a hostile TME, the tumors survived due to strong co-expression of several immune checkpoints such as PD-1, PD-L1, CTLA-4, Lag-3, and IDO in the invasive margin, stroma, and TILs. This finding suggests that the tumors may be responsive to checkpoint blockade. As a result, MMR status may be predictive of response to checkpoint inhibition.

Further, the type, density, and location of immune cells within the tumor (collectively known as immune contexture) have prognostic value. Multiple immune markers including total T lymphocytes (CD3), T cell effectors (CD8), their associated cytotoxic molecule (GZMB), and memory T cells (CD45RO) in the center of tumor (CT) and the invasive margin (IM) were quantified using IHC in tumors from 415 colorectal cancer patients [205]. The immune cell densities in each tumor region were higher in patients without recurrence than in patients with recurrence and were predictive of disease free survival (DFS) and OS. These results were independent of the staging of the tumor indicating the role of adaptive immune response in preventing tumor recurrence. In addition, presence of markers for Th1 polarization, cytotoxic and memory cells were predictive of low recurrence rate.

Baseline expression of TILs may not always suggest response to immune checkpoint blockade. TILs may not always predict response to ICPis. For example, CD8+ T cells at the IM were positively associated to response with pembrolizumab in patients with metastatic melanoma [206], but not in patients with unresectable stage III/IV melanoma treated with ipilimumab [207]. However, on treatment increase in the levels of tumor infiltrating T cells at the CT and IM were predictive of response to treatment with ICPi in several studies [206–208]. The antitumor activity was largely dependent on pre-existing adaptive immune mechanism as evidenced by the presence of higher numbers of CD8-, PD-1-, and PD-L1-expressing cells in the baseline samples [206].

Immunoscore

Immunoscore is a methodology by which in situ immune infiltrate is quantified. This supersedes the TNM classification of tumors used for estimation of the degree of progression of the tumor to make informed treatment decisions [205]. Marked

variations in clinical outcomes among patients with the same stage of disease were observed with TNM classification, partly due to failure to include the immune cells in the TME in TNM classification of tumors. As the interaction between the tumor cells and the immune cells play an important role in immune escape and progression of the tumor, immune contexture discussed above is a better prognostic indicator than TNM classification [209]. Therefore, a new scoring system was derived from immune contexture called the immunoscore, which is a ratio of the densities of two lymphocyte populations, CD3/CD45RO, CD3/CD8 or CD8/CD45RO, in the CT and IM. Due to difficulty in staining methods, a combination of two markers (CD3+ and CD8+) in CT and IM has been used by the worldwide immunoscore consortium in the development and validation of immunoscore as prognostic markers in different patient populations. The score ranges from immunoscore 0 (I0), when the densities of both the lymphocyte populations are low in both the regions to immunoscore 4 (I4), when the densities of both the lymphocyte populations are high in both the regions. This score is the strongest prognostic indicator of DFS and OS in patients with local and metastatic disease [210]. Recently, the consensus immunoscore was validated in a study conducted by an international consortium of centers in 13 countries [211]. In the analysis that included tissue samples from 2681 colorectal cancer patients, patients with a high immunoscore had the lowest risk of recurrence in 5 years, prolonged DFS and OS, a finding that has been confirmed in both the internal and external validation set. This scoring system will help to stratify patients based on the risk of recurrence. However, the universal application of immunoscore across tumor types has to be determined.

T Cell Receptor Diversity

As T cells play an important role in recognition and eradication of cancer cells, a diverse TCR repertoire will allow for detection of wide range of foreign antigens. On activation, TCRs undergo clonal expansion. Thus characterization and estimation of TCR repertoire diversity by next generation sequencing of CDR3 may provide insight into antitumor activity of ICPis. In a melanoma patient with metastatic lesion to the brain that progressed on ipilimumab, a durable complete clinical response was achieved with sequential whole brain radiation therapy and pembrolizumab [212]. A high-throughput CDR3 sequencing of the intratumoral T cells in the brain metastasis obtained before treatment and the circulating peripheral T cells obtained sequentially during treatment showed that the dominant CD8+ T cell clone in the brain metastasis (pretreatment) had clonally expanded on treatment with pembrolizumab and was detected as the most frequently occurring clone in the blood. This indicates presence of pre-existing but inadequate adaptive immune response that was bolstered by treatment with pembrolizumab. Similar on-treatment clonal expansion of a CD8+ T cell clone present in the metastatic site prior to treatment was seen in a NSCLC patient who experienced pathological complete response with nivolumab [213]. In 10 patients with metastatic melanoma treated with

nivolumab [214], oligoclonal expansion of certain TCR- β clonotypes was observed in post-treatment tumor tissues of responders. Similar results were also observed in 25 patients with metastatic melanoma treated with pembrolizumab [206]. TCR sequencing of pre- and post-treatment samples showed the number of clones that had expanded was ten times more in the responders than in non-responders. Further, clinical response was associated with a more restricted TCR beta chain usage in pre-dosing samples. Thus, a diverse TCR repertoire at baseline and on-treatment tumor antigen-specific clonal expansion may be predictive of response to treatment with ICPis.

Mutation Load and Molecular Alterations

Tumors with high mutational load such as melanoma, NSCLC, head and neck squamous cell carcinoma (HNSCC) are more likely to respond to treatment with ICPis as neoepitomes generated by somatic mutations function as neoantigens and elicit a brisk immune response [215]. In several clinical trials, higher clinical benefit rate and longer progression-free survival have been reported in patients with high mutation burden treated with ICPis [215–217]. It is for the same reason that improved treatment outcomes with ICPis have been reported in patients with solid tumors, colorectal cancer patients in particular, with defects in the mismatch repair (MMR) mechanism [218, 219]. However, Snyder and colleagues described that while high mutational load correlated to sustained response to CTLA-4 blockade, not all melanoma patients with high mutational load responded to therapy [216]. But, the presence of tetrapeptide neoepitope signature in these patients with high mutation load correlated strongly with long-term clinical benefit and OS. On the contrary, tumors with low mutational loads (e.g., pancreatic and prostate cancer) were not responsive to ICPi. Also, molecular alterations in the PI3K pathway may promote tumor immune evasion through constitutive expression of PD-L1 [220]. Assessment of PD-L1 expression in such conditions may predict response with PD-1/PD-L1 inhibitors. Similarly, increased expression of VEGF promotes angiogenesis and is associated with poor prognosis [199].

Immune Gene Signature

Differential expression of genes may help to identify phenotypes responsive to treatment with ICPis. For example, loss-of-function *BRCA2* mutations with specific mutational signatures were identified in responding melanoma tumors sampled from patients on treatment with anti-PD-1 agents [217]. Likewise, in melanoma patients treated with pembrolizumab, an IFN- γ 10-gene and an expanded immune 28-gene signatures in pretreatment samples were significantly associated with ORR and PFS

[221]. On further evaluation, more refined immune signatures were found to produce similar results in patients with HNSCC and gastric cancer [222]. High pretreatment levels of IFN- γ mRNA and PD-L1 protein expression were associated with increased ORR and longer OS in NSCLC patients treated with durvalumab [223]. A similar association between high expression of T-effector-associated, IFN- γ -associated and PD-L1 genes in tumor tissue and improved OS was seen in NSCLC patients treated with atezolizumab [224]. The T-effector-associated and IFN- γ -associated gene expression was associated with PD-L1 expression on immune cells and not on tumor cells suggesting the role of pre-existing adaptive immune response. On the contrary, a group of 26 innate anti-PD-1 resistance (IPRES) signatures characterized by higher expression of mesenchymal transition, angiogenesis, hypoxia, and wound healing genes were identified in pretreatment melanoma tumors resistant to anti-PD-1 therapy [217]. The IPRES signature was also found in non-responsive pretreatment tumor samples from patients with other solid tumors such as adenocarcinoma of the lung, colon, and pancreas and clear cell carcinoma of kidney. Thus immune-related gene expression signatures may be associated with treatment outcomes.

Cancer Immunogram

The cancer immunogram model was developed to overcome the limitation that no single biomarker can truly reflect the dynamic interaction between the immune cells and tumor. Based on the assumption that T cells are the ultimate effectors of antitumor activity, seven parameters were included in the model to understand the interaction between the tumor and the immune cells in the TME of the patient [225]. The seven parameters and their potential biomarkers in parenthesis are: (1) tumor foreignness (mutation load), (2) general immune status (lymphocyte count), (3) immune cell infiltration (intratumoral T cells), (4) absence of checkpoints (PD-L1), (5) absence of soluble inhibitors (IL-6 and C-reactive protein [CRP]), (6) absence of inhibitory tumor metabolism (lactate dehydrogenase [LDH], glucose utilization), (7) and tumor sensitivity to immune effectors (MHC expression, IFN- γ sensitivity). The data points for each of the seven parameters are plotted in a radar plot and the line joining the individual data points provides a personalized framework reflecting the interaction in the TME. The gaps in the radar plot indicate potential therapeutic strategies that may evoke an effective immune response in the patient.

A modified immunogram has been developed based on the seven steps in the cancer immunity cycle for use in NSCLC patients [226]. The eight axes of the immunogram score (IGS) are: IGS₁, existence of T cell immunity in the tumor; IGS₂, tumor antigenicity (existence of neoantigens and cancer germline antigens); IGS₃, priming and activation (presence of activated DCs); IGS₄, trafficking and T cell infiltration; IGS₅, recognition of tumor antigens; IGS₆, absence of inhibitory cells (Tregs and MDSCs); IGS₇, absence of checkpoint expression (PD-1, PD-L1, etc.); and IGS₈, absence of inhibitory molecules (IDO 1, arginase 1, etc.). High scores for IGS₁₋₅ indicate a favorable environment for development of T cell immunity. On the

contrary, high scores for IGS₆₋₈ indicate immune suppression. Based on the radar plot, three groups of patients have been identified. Patients with high IGS₁₋₅ and low IGS₆₋₈ represent T cell-rich phenotype where antitumor activity is dampened by an immunosuppressive TME, patients with low IGS₁, IGS₃₋₅ represent T cell-poor phenotype with defects in the T cell priming process, and patients in whom IGS₂, IGS₆₋₈ are maintained represent an intermediate phenotype. Thus, the immunogram helps to identify areas of therapeutic focus to elicit an effective antitumor response. Cancer immunograms are promising for personalized approach to immunotherapy.

Serum Biomarkers

Several routinely available peripheral blood parameters have been evaluated as a biomarker of response to treatment with ICPis [208, 227–234]. Most common among them are absolute lymphocyte count (ALC), absolute eosinophil count (AEC), LDH, and CRP. In a compassionate use trial with ipilimumab in patients with advanced refractory melanoma, ALC $\geq 1000/\mu\text{L}$ after two treatments with ipilimumab was significantly associated with clinical benefit and OS [230, 231]. Though ALC at baseline and after one dose of ipilimumab showed only a trend for improved treatment outcomes, they may be prognostic because a threshold ALC of 1000 cells/ μL may be required for adequate activation of the immune system for patients to derive meaningful antitumor response with therapy. Similar results were seen in several clinical trials in patients with melanoma treated with ipilimumab [230–234], where an increase in ALC levels from baseline was associated with improved OS and disease control compared to patients with stable or decreasing levels. Likewise increase in AEC levels after two courses of ipilimumab was associated with OS [230] and was an independent predictor of response in patients with melanoma [235]. On the other hand, elevated level of LDH at baseline was an independent predictor of poor survival [230, 236]. Despite the association between these peripheral blood parameters and treatment outcomes, there is no validated biomarker available for use in the clinic.

Circulating Biomarkers

Serial assessment of circulating tumor cells (CTCs) and circulating tumor DNA (ctDNA), which is a measure of tumor burden, may predict response to treatment with checkpoint inhibitors. The association between ctDNA and treatment outcomes was evaluated in three groups of patients treated with PD-1 inhibitors as single agents or in combination with ipilimumab [237]. Group A included patients with undetectable ctDNA at baseline and during treatment, Group B had patients with detectable ctDNA at baseline but undetectable early during therapy, and Group C included patients with detectable ctDNA at baseline and during therapy. Compared

to baseline ctDNA, persistent on treatment levels of ctDNA was associated with decreased ORR and poor survival. On the other hand, increase in circulating levels of immune cells, Ki-67+ T cells, was associated with clinical benefit in NSCLC patients on treatment with PD-1 inhibitors [238]. If these findings are validated in large prospective cohorts, in the context of intratumoral heterogeneity, minimally invasive and easily accessible liquid biopsies may serve as a more comprehensive alternate technique for biomarker assessment.

Microbiome Assessment

Emerging data indicate that gut microbiome may be associated with response to treatment with PD-1 inhibitors. Alpha diversity of gut microbiomes in fecal samples was significantly higher in patients with metastatic melanoma responding (CR/PR/SD \geq 6 months) to treatment with PD-1 inhibitors [239]. And, patients with higher alpha diversity had longer PFS compared to patients with low or intermediate diversity. Further, the gut microbiome was enriched for Clostridiales in responders and Bacteroidales in non-responders. And patients with abundance of Faecalibacterium genus in Clostridiales order had significantly longer PFS compared to patients with abundance of Bacteroidales. Thus favorable gut microbiome may enhance antitumor response in patients treated with checkpoint inhibitors.

Due to the dynamic nature of immune response, development of immune oncology biomarkers is challenging. To this end, immune monitoring assays have been developed to perform genomic, proteomic, and functional studies on paired tumor and blood samples obtained before and after treatment with immunotherapeutic agents [197]. It is expected that correlation of changes in these biomarkers to treatment outcomes would provide mechanistic insight into pathways of response or resistance to immunotherapeutic agents that could guide the development of biomarker-driven, synergistic, immunotherapy-based treatment combinations. In addition, biomarkers may vary depending on the mechanism of action of the immunotherapeutic agent [159, 191]. Therefore, identification of a single immunologic biomarker may not be predictive of response [197]. This indicates a need to identify multi-factorial biomarker panels that would help to determine the immunogenic nature of the tumor and predict response or resistance to treatment. For example, presence of intratumoral CD8+ T cells, expression of PD-L1 on tumor cells, and increased mutational load have been associated with greater likelihood of response to PD-1/PD-L1 checkpoint inhibition [240].

Conclusion

Seminal studies have described the different components of the innate and adaptive immune system. Though they are two distinct arms of the human immune system, they are intricately organized in time and space and are critically dependent upon

one another. While the blockade of immune checkpoints by mAbs to unleash the antitumor immune response by T cells has now emerged as a powerful therapeutic tool in the treatment of advanced cancer, components of the innate immune system contribute to the activation and development of adaptive immunity. Improved understanding of the interaction between the tumor cells and the immune cells in the complex TME through rigorous immune profiling will guide the future development of new immunotherapeutic strategies as well as the identification of potential biomarkers of clinical response.

References

1. Benito-Martin A, Di Giannatale A, Ceder S, Peinado H. The new deal: a potential role for secreted vesicles in innate immunity and tumor progression. *Front Immunol.* 2015;6:66.
2. Murphy K, Weaver C. *Janeway's immunobiology.* 9th ed. New York: Garland Science, Taylor & Francis; 2016.
3. Mellman I. Dendritic cells: master regulators of the immune response. *Cancer Immunol Res.* 2013;1:145–9.
4. Turvey SE, Broide DH. Innate immunity. *J Allergy Clin Immunol.* 2010;125:S24–32.
5. Balkwill F, Mantovani A. Inflammation and cancer: back to Virchow? *Lancet.* 2001;357:539–45.
6. Janeway CA Jr, Medzhitov R. Innate immune recognition. *Annu Rev Immunol.* 2002;20:197–216.
7. Finn OJ. Immuno-oncology: understanding the function and dysfunction of the immune system in cancer. *Ann Oncol.* 2012;23(Suppl 8):viii6–9.
8. Lin WW, Karin M. A cytokine-mediated link between innate immunity, inflammation, and cancer. *J Clin Invest.* 2007;117:1175–83.
9. Grivennikov SI, Greten FR, Karin M. Immunity, inflammation, and cancer. *Cell.* 2010;140:883–99.
10. Fedeles BI, Freudenthal BD, Yau E, Singh V, Chang SC, Li DY, Delaney JC, Wilson SH, Essigmann JM. Intrinsic mutagenic properties of 5-chlorocytosine: a mechanistic connection between chronic inflammation and cancer. *Proc Natl Acad Sci U S A.* 2015;112:E4571–E80.
11. Dvorak HF, Flier J, Frank H. Tumors - wounds that do not heal - similarities between tumor stroma generation and wound-healing. *N Engl J Med.* 1986;315:1650–9.
12. Galli SJ, Borregaard N, Wynn TA. Phenotypic and functional plasticity of cells of innate immunity: macrophages, mast cells and neutrophils. *Nat Immunol.* 2011;12:1035–44.
13. Fridlender ZG, Sun J, Kim S, Kapoor V, Cheng G, Ling L, Worthen GS, Albelda SM. Polarization of tumor-associated neutrophil phenotype by TGF-beta: "N1" versus "N2" TAN. *Cancer Cell.* 2009;16:183–94.
14. Mayadas TN, Cullere X, Lowell CA. The multifaceted functions of neutrophils. *Annu Rev Pathol.* 2014;9:181–218.
15. Borregaard N. Neutrophils, from marrow to microbes. *Immunity.* 2010;33:657–70.
16. Kobayashi Y. Neutrophil infiltration and chemokines. *Crit Rev Immunol.* 2006;26:307–15.
17. Scapini P, Carletto A, Nardelli B, Calzetti F, Roschke V, Merigo F, Tamassia N, Pieropan S, Biasi D, Sbarbati A, Sozzani S, Bambara L, Cassatella MA. Proinflammatory mediators elicit secretion of the intracellular B-lymphocyte stimulator pool (BLyS) that is stored in activated neutrophils: implications for inflammatory diseases. *Blood.* 2005;105:830–7.
18. Theilgaard-Monch K, Knudsen S, Follin P, Borregaard N. The transcriptional activation program of human neutrophils in skin lesions supports their important role in wound healing. *J Immunol.* 2004;172:7684–93.

19. Fridlender ZG, Albelda SM. Tumor-associated neutrophils: friend or foe? *Carcinogenesis*. 2012;33:949–55.
20. Piccard H, Muschel RJ, Opdenakker G. On the dual roles and polarized phenotypes of neutrophils in tumor development and progression. *Crit Rev Oncol Hematol*. 2012;82:296–309.
21. Gregory AD, Houghton AM. Tumor-associated neutrophils: new targets for cancer therapy. *Cancer Res*. 2011;71:2411–6.
22. Houghton AM, Rzymkiewicz DM, Ji H, Gregory AD, Egea EE, Metz HE, Stolz DB, Land SR, Marconcini LA, Kliment CR, Jenkins KM, Beaulieu KA, Mouded M, Frank SJ, Wong KK, Shapiro SD. Neutrophil elastase-mediated degradation of IRS-1 accelerates lung tumor growth. *Nat Med*. 2010;16:219–23.
23. Queen MM, Ryan RE, Holzer RG, Keller-Peck CR, Jorcyk CL. Breast cancer cells stimulate neutrophils to produce oncostatin M: potential implications for tumor progression. *Cancer Res*. 2005;65:8896–904.
24. Acharyya S, Oskarsson T, Vanharanta S, Malladi S, Kim J, Morris PG, Manova-Todorova K, Leversha M, Hogg N, Seshan VE, Norton L, Brogi E, Massague J. A CXCL1 paracrine network links cancer chemoresistance and metastasis. *Cell*. 2012;150:165–78.
25. Shojaei F, Singh M, Thompson JD, Ferrara N. Role of Bv8 in neutrophil-dependent angiogenesis in a transgenic model of cancer progression. *Proc Natl Acad Sci U S A*. 2008;105:2640–5.
26. Liang W, Ferrara N. The complex role of neutrophils in tumor angiogenesis and metastasis. *Cancer Immunol Res*. 2016;4:83–91.
27. van Gisbergen KPJM, Geijtenbeek TBH, van Kooyk Y. Close encounters of neutrophils and DCs. *Trends Immunol*. 2005;26:626–31.
28. Scapini P, Lapinet-Vera JA, Gasperini S, Calzetti F, Bazzoni F, Cassatella MA. The neutrophil as a cellular source of chemokines. *Immunol Rev*. 2000;177:195–203.
29. Chaplin DD. Overview of the immune response. *J Allergy Clin Immunol*. 2010;125:S3–23.
30. Mantovani A, Schioppa T, Porta C, Allavena P, Sica A. Role of tumor-associated macrophages in tumor progression and invasion. *Cancer Metastasis Rev*. 2006;25:315–22.
31. Lin EY, Nguyen AV, Russell RG, Pollard JW. Colony-stimulating factor 1 promotes progression of mammary tumors to malignancy. *J Exp Med*. 2001;193:727–40.
32. Duyndam MC, Hillhorst MC, Schluper HM, Verheul HM, van Diest PJ, Kraal G, Pinedo HM, Boven E. Vascular endothelial growth factor-165 overexpression stimulates angiogenesis and induces cyst formation and macrophage infiltration in human ovarian cancer xenografts. *Am J Pathol*. 2002;160:537–48.
33. Sica A, Schioppa T, Mantovani A, Allavena P. Tumour-associated macrophages are a distinct M2 polarised population promoting tumour progression: potential targets of anti-cancer therapy. *Eur J Cancer*. 2006;42:717–27.
34. Sica A, Allavena P, Mantovani A. Cancer related inflammation: the macrophage connection. *Cancer Lett*. 2008;267:204–15.
35. Solinas G, Germano G, Mantovani A, Allavena P. Tumor-associated macrophages (TAM) as major players of the cancer-related inflammation. *J Leukoc Biol*. 2009;86:1065–73.
36. Pollard JW. Tumour-educated macrophages promote tumour progression and metastasis. *Nat Rev Cancer*. 2004;4:71–8.
37. Sica A, Saccani A, Bottazzi B, Polentarutti N, Vecchi A, van Damme J, Mantovani A. Autocrine production of IL-10 mediates defective IL-12 production and NF-kappa B activation in tumor-associated macrophages. *J Immunol*. 2000;164:762–7.
38. Mantovani A, Allavena P, Sica A. Tumour-associated macrophages as a prototypic type II polarised phagocyte population: role in tumour progression. *Eur J Cancer*. 2004;40:1660–7.
39. Tsutsui S, Yasuda K, Suzuki K, Tahara K, Higashi H, Era S. Macrophage infiltration and its prognostic implications in breast cancer: the relationship with VEGF expression and microvessel density. *Oncol Rep*. 2005;14:425–31.
40. Zhang J, Yan Y, Yang Y, Wang L, Li M, Wang J, Liu X, Duan X, Wang J. High infiltration of tumor-associated macrophages influences poor prognosis in human gastric cancer patients, associates with the phenomenon of EMT. *Medicine (Baltimore)*. 2016;95:e2636.

41. Hanada T, Nakagawa M, Emoto A, Nomura T, Nasu N, Nomura Y. Prognostic value of tumor-associated macrophage count in human bladder cancer. *Int J Urol.* 2000;7:263–9.
42. Salvesen HB, Akslen LA. Significance of tumour-associated macrophages, vascular endothelial growth factor and thrombospondin-1 expression for tumour angiogenesis and prognosis in endometrial carcinomas. *Int J Cancer.* 1999;84:538–43.
43. Fujimoto J, Sakaguchi H, Aoki I, Tamaya T. Clinical implications of expression of interleukin 8 related to angiogenesis in uterine cervical cancers. *Cancer Res.* 2000;60:2632–5.
44. Shimura S, Yang G, Ebara S, Wheeler TM, Frolov A, Thompson TC. Reduced infiltration of tumor-associated macrophages in human prostate cancer: association with cancer progression. *Cancer Res.* 2000;60:5857–61.
45. Forssell J, Oberg A, Henriksson ML, Stenling R, Jung A, Palmqvist R. High macrophage infiltration along the tumor front correlates with improved survival in colon cancer. *Clin Cancer Res.* 2007;13:1472–9.
46. Fulkerson PC, Rothenberg ME. Targeting eosinophils in allergy, inflammation and beyond. *Nat Rev Drug Discov.* 2013;12:117–29.
47. Rothenberg ME, Hogan SP. The eosinophil. *Annu Rev Immunol.* 2006;24:147–74.
48. Kita H. Eosinophils: multifaceted biological properties and roles in health and disease. *Immunol Rev.* 2011;242:161–77.
49. Muniz VS, Weller PF, Neves JS. Eosinophil crystalloid granules: structure, function, and beyond. *J Leukoc Biol.* 2012;92:281–8.
50. Fernandez-Acenero MJ, Galindo-Gallego M, Sanz J, Aljama A. Prognostic influence of tumor-associated eosinophilic infiltrate in colorectal carcinoma. *Cancer.* 2000;88:1544–8.
51. Dorta RG, Landman G, Kowalski LP, Lauris JRP, Latorre MRDO, Oliveira DT. Tumour-associated tissue eosinophilia as a prognostic factor in oral squamous cell carcinomas. *Histopathology.* 2002;41:152–7.
52. Costello R, O’Callaghan T, Sebahoun G. [Eosinophils and antitumour response]. *Rev Med Interne.* 2005;26:479–84.
53. Ohkawara Y, Lim KG, Xing Z, Glibetic M, Nakano K, Dolovich J, Croitoru K, Weller PF, Jordana M. CD40 expression by human peripheral blood eosinophils. *J Clin Invest.* 1996;97:1761–6.
54. Woerly G, Roger N, Loiseau S, Dombrowicz D, Capron A, Capron M. Expression of CD28 and CD86 by human eosinophils and role in the secretion of type 1 cytokines (interleukin 2 and interferon gamma): inhibition by immunoglobulin a complexes. *J Exp Med.* 1999;190:487–95.
55. Shi HZ, Humbles A, Gerard C, Jin Z, Weller PF. Lymph node trafficking and antigen presentation by endobronchial eosinophils. *J Clin Invest.* 2000;105:945–53.
56. Lotfi R, Herzog GI, DeMarco RA, Beer-Stolz D, Lee JJ, Rubartelli A, Schrezenmeier H, Lotze MT. Eosinophils oxidize damage-associated molecular pattern molecules derived from stressed cells. *J Immunol.* 2009;183:5023–31.
57. Cormier SA, Taranova AG, Bedient C, Nguyen T, Protheroe C, Pero R, Dimina D, Ochkur SI, O’Neill K, Colbert D, Lombardi TR, Constant S, McGarry MP, Lee JJ, Lee NA. Pivotal advance: eosinophil infiltration of solid tumors is an early and persistent inflammatory host response. *J Leukoc Biol.* 2006;79:1131–9.
58. Minton K. Granulocytes: eosinophils enable the antitumour T cell response. *Nat Rev Immunol.* 2015;15:333.
59. Carretero R, Sektiglu IM, Garbi N, Salgado OC, Beckhove P, Hammerling GJ. Eosinophils orchestrate cancer rejection by normalizing tumor vessels and enhancing infiltration of CD8(+) T cells. *Nat Immunol.* 2015;16:609–17.
60. Falcone FH, Zillikens D, Gibbs BF. The 21st century renaissance of the basophil? Current insights into its role in allergic responses and innate immunity. *Exp Dermatol.* 2006;15:855–64.
61. Schroeder JT, DW MG Jr, Lichtenstein LM. Human basophils: mediator release and cytokine production. *Adv Immunol.* 2001;77:93–122.

62. Haas H, Falcone FH, Holland MJ, Schramm G, Haisch K, Gibbs BF, Bufe A, Schlaak M. Early interleukin-4: its role in the switch towards a Th2 response and IgE-mediated allergy. *Int Arch Allergy Immunol.* 1999;119:86–94.
63. Schroeder JT. Basophils beyond effector cells of allergic inflammation. *Adv Immunol.* 2009;101:123–61.
64. Prevette N, Staiano RI, Granata F, Detoraki A, Necchi V, Ricci V, Triggiani M, De Paulis A, Marone G, Genovese A. Expression and function of angiopoietins and their tie receptors in human basophils and mast cells. *J Biol Regul Homeost Agents.* 2013;27:827–39.
65. De Monte L, Wormann S, Brunetto E, Heltai S, Magliacane G, Reni M, Paganoni AM, Recalde H, Mondino A, Falconi M, Aleotti F, Balzano G, Ul HA, Doglioni C, Protti MP. Basophil recruitment into tumor-draining lymph nodes correlates with Th2 inflammation and reduced survival in pancreatic cancer patients. *Cancer Res.* 2016;76:1792–803.
66. Frossi B, De Carli M, Pucillo C. The mast cell: an antenna of the microenvironment that directs the immune response. *J Leukoc Biol.* 2004;75:579–85.
67. Qi X, Hong J, Chaves L, Zhuang Y, Chen Y, Wang D, Chabon J, Graham B, Ohmori K, Li Y, Huang H. Antagonistic regulation by the transcription factors C/EBPalpha and MITF specifies basophil and mast cell fates. *Immunity.* 2013;39:97–110.
68. Marone G, Galli SJ, Kitamura Y. Probing the roles of mast cells and basophils in natural and acquired immunity, physiology and disease. *Trends Immunol.* 2002;23:425–7.
69. Galli SJ, Franco CB. Basophils are back! *Immunity.* 2008;28:495–7.
70. Stone KD, Prussin C, Metcalfe DD. IgE, mast cells, basophils, and eosinophils. *J Allergy Clin Immunol.* 2010;125:S73–80.
71. Metcalfe DD. Mast cells and mastocytosis. *Blood.* 2008;112:946–56.
72. Nonomura N, Takayama H, Nishimura K, Oka D, Nakai Y, Shiba M, Tsujimura A, Nakayama M, Aozasa K, Okuyama A. Decreased number of mast cells infiltrating into needle biopsy specimens leads to a better prognosis of prostate cancer. *Br J Cancer.* 2007;97:952–6.
73. Rojas IG, Spencer ML, Martinez A, Maurelia MA, Rudolph MI. Characterization of mast cell subpopulations in lip cancer. *J Oral Pathol Med.* 2005;34:268–73.
74. Fukushima H, Ohsawa M, Ikura Y, Naruko T, Sugama Y, Suekane T, Kitabayashi C, Inoue T, Hino M, Ueda M. Mast cells in diffuse large B-cell lymphoma; their role in fibrosis. *Histopathology.* 2006;49:498–505.
75. Kormelink TG, Abudukelimu A, Redegeld FA. Mast cells as target in cancer therapy. *Curr Pharm Des.* 2009;15:1868–78.
76. Ribatti D, Vacca A, Nico B, Crivellato E, Roncali L, Dammacco F. The role of mast cells in tumour angiogenesis. *Br J Haematol.* 2001;115:514–21.
77. Rajput AB, Turbin DA, Cheang MC, Voduc DK, Leung S, Gelmon KA, Gilks CB, Huntsman DG. Stromal mast cells in invasive breast cancer are a marker of favourable prognosis: a study of 4,444 cases. *Breast Cancer Res Treat.* 2008;107:249–57.
78. Chan JK, Magistris A, Loizzi V, Lin F, Rutgers J, Osann K, DiSaia PJ, Samoszuk M. Mast cell density, angiogenesis, blood clotting, and prognosis in women with advanced ovarian cancer. *Gynecol Oncol.* 2005;99:20–5.
79. Welsh TJ, Green RH, Richardson D, Waller DA, O’Byrne KJ, Bradding P. Macrophage and mast-cell invasion of tumor cell islets confers a marked survival advantage in non-small-cell lung cancer. *J Clin Oncol.* 2005;23:8959–67.
80. Tan SY, Fan Y, Luo HS, Shen ZX, Guo Y, Zhao LJ. Prognostic significance of cell infiltrations of immunosurveillance in colorectal cancer. *World J Gastroenterol.* 2005;11:1210–4.
81. Latti S, Leskinen M, Shiota N, Wang YF, Kovanen PT, Lindstedt KA. Mast cell-mediated apoptosis of endothelial cells in vitro: a paracrine mechanism involving TNF-alpha-mediated down-regulation of bcl-2 expression. *J Cell Physiol.* 2003;195:130–8.
82. Leskinen MJ, Lindstedt KA, Wang YF, Kovanen PT. Mast cell chymase induces smooth muscle cell apoptosis by a mechanism involving fibronectin degradation and disruption of focal adhesions. *Arterioscler Thromb Vasc Biol.* 2003;23:238–43.

83. Hammer GE, Ma A. Molecular control of steady-state dendritic cell maturation and immune homeostasis. *Annu Rev Immunol.* 2013;31:743–91.
84. Liu K, Nussenzweig MC. Origin and development of dendritic cells. *Immunol Rev.* 2010;234:45–54.
85. Liu K, Victora GD, Schwickert TA, Guermonprez P, Meredith MM, Yao K, Chu FF, Randolph GJ, Rudensky AY, Nussenzweig M. In vivo analysis of dendritic cell development and homeostasis. *Science.* 2009;324:392–7.
86. Shortman K, Naik SH. Steady-state and inflammatory dendritic-cell development. *Nat Rev Immunol.* 2007;7:19–30.
87. Trombetta ES, Mellman I. Cell biology of antigen processing in vitro and in vivo. *Annu Rev Immunol.* 2005;23:975–1028.
88. Steinman RM. Decisions about dendritic cells: past, present, and future. *Annu Rev Immunol.* 2012;30(30):1–22.
89. Ljunggren HG, Karre K. In search of the missing self - Mhc molecules and Nk cell recognition. *Immunol Today.* 1990;11:237–44.
90. Vivier E, Nunes JA, Vely F. Natural killer cell signaling pathways. *Science.* 2004;306:1517–9.
91. Tomasello E, Blery M, Vely F, Vivier E. Signaling pathways engaged by NK cell receptors: double concerto for activating receptors, inhibitory receptors and NK cells. *Semin Immunol.* 2000;12:139–47.
92. Strengell M, Matikainen S, Siren J, Lehtonen A, Foster D, Julkunen I, Sareneva T. IL-21 in synergy with IL-15 or IL-18 enhances IFN-gamma production in human NK and T cells. *J Immunol.* 2003;170:5464–9.
93. Brady J, Carotta S, Thong RP, Chan CJ, Hayakawa Y, Smyth MJ, Nutt SL. The interactions of multiple cytokines control NK cell maturation. *J Immunol.* 2010;185:6679–88.
94. Lunemann A, Lunemann JD, Munz C. Regulatory NK-cell functions in inflammation and autoimmunity. *Mol Med.* 2009;15:352–8.
95. Becknell B, Caligiuri MA. Natural killer cells in innate immunity and cancer. *J Immunother.* 2008;31:685–92.
96. Kaiser BK, Yim D, Chow IT, Gonzalez S, Dai Z, Mann HH, Strong RK, Groh V, Spies T. Disulphide-isomerase-enabled shedding of tumour-associated NKG2D ligands. *Nature.* 2007;447:482–6.
97. Groh V, Wu J, Yee C, Spies T. Tumour-derived soluble MIC ligands impair expression of NKG2D and T-cell activation. *Nature.* 2002;419:734–8.
98. Castriconi R, Cantoni C, Della Chiesa M, Vitale M, Marcenaro E, Conte R, Biassoni R, Bottino C, Moretta L, Moretta A. Transforming growth factor beta 1 inhibits expression of NKp30 and NKG2D receptors: consequences for the NK-mediated killing of dendritic cells. *Proc Natl Acad Sci U S A.* 2003;100:4120–5.
99. Sconocchia G, Titus JA, Segal DM. Signaling pathways regulating CD44-dependent cytotoxicity in natural killer cells. *Blood.* 1997;90:716–25.
100. Wang W, Erbe AK, Hank JA, Morris ZS, Sondel PM. NK cell-mediated antibody-dependent cellular cytotoxicity in cancer immunotherapy. *Front Immunol.* 2015;6:368.
101. Ferlazzo G, Thomas D, Lin SL, Goodman K, Morandi B, Muller WA, Moretta A, Munz C. The abundant NK cells in human secondary lymphoid tissues require activation to express killer cell Ig-like receptors and become cytolytic. *J Immunol.* 2004;172:1455–62.
102. Sun JC, Beilke JN, Lanier LL. Adaptive immune features of natural killer cells. *Nature.* 2009;457:557–61.
103. Albertsson PA, Basse PH, Hokland M, Goldfarb RH, Nagelkerke JF, Nanmark U, Kuppen PJ. NK cells and the tumour microenvironment: implications for NK-cell function and anti-tumour activity. *Trends Immunol.* 2003;24:603–9.
104. Robey E, Fowlkes BJ. Selective events in T cell development. *Annu Rev Immunol.* 1994;12:675–705.
105. Germain RN. T-cell development and the CD4-CD8 lineage decision. *Nat Rev Immunol.* 2002;2:309–22.

106. Scollay R, Wilson A, D'Amico A, Kelly K, Egerton M, Pearse M, Wu L, Shortman K. Developmental status and reconstitution potential of subpopulations of murine thymocytes. *Immunol Rev.* 1988;104:81–120.
107. Blackburn CC, Manley NR. Developing a new paradigm for thymus organogenesis. *Nat Rev Immunol.* 2004;4:278–89.
108. Vonboehmer H, Teh HS, Kisielow P. The thymus selects the useful, neglects the useless and destroys the harmful. *Immunol Today.* 1989;10:57–61.
109. Leung RK, Thomson K, Gallimore A, Jones E, Van den Broek M, Siervo S, Alsheikhly AR, McMichael A, Rahemtulla A. Deletion of the CD4 silencer element supports a stochastic mechanism of thymocyte lineage commitment. *Nat Immunol.* 2001;2:1167–73.
110. Sharma P, Wagner K, Wolchok JD, Allison JP. Novel cancer immunotherapy agents with survival benefit: recent successes and next steps. *Nat Rev Cancer.* 2011;11:805–12.
111. Lugo-Villarino G, Maldonado-Lopez R, Possemato R, Penaranda C, Glimcher LH. T-bet is required for optimal production of IFN-gamma and antigen-specific T cell activation by dendritic cells. *Proc Natl Acad Sci U S A.* 2003;100:7749–54.
112. Zhu JF, Guo LY, Watson CJ, Hu-Li J, Paul WE. Stat6 is necessary and sufficient for IL-4's role in Th2 differentiation and cell expansion. *J Immunol.* 2001;166:7276–81.
113. Zhou L, Lopes JE, Chong MM, Ivanov II, Min R, Victora GD, Shen Y, Du J, Rubtsov YP, Rudensky AY, Ziegler SF, Littman DR. TGF-beta-induced Foxp3 inhibits T(H)17 cell differentiation by antagonizing RORgamma function. *Nature.* 2008;453:236–40.
114. Chen WJ, Jin WW, Hardegen N, Lei KJ, Li L, Marinos N, McGrady G, Wahl SM. Conversion of peripheral CD4(+)CD25(-) naive T cells to CD4(+)CD25(+) regulatory T cells by TGF-beta induction of transcription factor Foxp3. *J Exp Med.* 2003;198:1875–86.
115. Nurieva RI, Chung Y, Hwang D, Yang XO, Kang HS, Ma L, Wang YH, Watowich SS, Jetten AM, Tian Q, Dong C. Generation of T follicular helper cells is mediated by interleukin-21 but independent of T helper 1, 2, or 17 cell lineages. *Immunity.* 2008;29:138–49.
116. Staudt V, Bothur E, Klein M, Lingnau K, Reuter S, Grebe N, Gerlitzki B, Hoffmann M, Ulges A, Taube C, Dehzad N, Becker M, Stassen M, Steinborn A, Lohoff M, Schild H, Schmitt E, Bopp T. Interferon-regulatory factor 4 is essential for the developmental program of T helper 9 cells. *Immunity.* 2010;33:192–202.
117. Saule P, Trauet J, Dutriez V, Lekeux W, Dessaint JP, Labalette M. Accumulation of memory T cells from childhood to old age: central and effector memory cells in CD4(+) versus effector memory and terminally differentiated memory cells in CD8(+) compartment. *Mech Ageing Dev.* 2006;127:274–81.
118. Chen L, Flies DB. Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat Rev Immunol.* 2013;13:227–42.
119. Linsley PS, Clark EA, Ledbetter JA. T-cell antigen CD28 mediates adhesion with B cells by interacting with activation antigen B7/BB-1. *Proc Natl Acad Sci U S A.* 1990;87:5031–5.
120. Nam KO, Kang H, Shin SM, Cho KH, Kwon B, Kwon BS, Kim SJ, Lee HW. Cross-linking of 4-1BB activates TCR-signaling pathways in CD8+ T lymphocytes. *J Immunol.* 2005;174:1898–905.
121. Godfrey WR, Fagnoni FF, Harara MA, Buck D, Engleman EG. Identification of a human OX-40 ligand, a costimulator of CD4+ T cells with homology to tumor necrosis factor. *J Exp Med.* 1994;180:757–62.
122. Nocentini G, Riccardi C. GITR: a modulator of immune response and inflammation. *Adv Exp Med Biol.* 2009;647:156–73.
123. Linsley PS, Brady W, Urnes M, Grosmaire LS, Damle NK, Ledbetter JA. CTLA-4 is a second receptor for the B cell activation antigen B7. *J Exp Med.* 1991;174:561–9.
124. Freeman GJ, Long AJ, Iwai Y, Bourque K, Chernova T, Nishimura H, Fitz LJ, Malenkovich N, Okazaki T, Byrne MC, Horton HF, Fouser L, Carter L, Ling V, Bowman MR, Carreno BM, Collins M, Wood CR, Honjo T. Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med.* 2000;192:1027–34.

125. Buchbinder EI, Desai A. CTLA-4 and PD-1 pathways similarities, differences, and implications of their inhibition. *Am J Clin Oncol.* 2016;39:98–106.
126. Liang SC, Latchman YE, Buhlmann JE, Tomczak MF, Horwitz BH, Freeman GJ, Sharpe AH. Regulation of PD-1, PD-L1, and PD-L2 expression during normal and autoimmune responses. *Eur J Immunol.* 2003;33:2706–16.
127. Okazaki T, Maeda A, Nishimura H, Kurosaki T, Honjo T. PD-1 immunoreceptor inhibits B cell receptor-mediated signaling by recruiting src homology 2-domain-containing tyrosine phosphatase 2 to phosphotyrosine. *Proc Natl Acad Sci U S A.* 2001;98:13866–71.
128. Anderson AC, Joller N, Kuchroo VK. Lag-3, Tim-3, and TIGIT: co-inhibitory receptors with specialized functions in immune regulation. *Immunity.* 2016;44:989–1004.
129. Joller N, Lozano E, Burkett PR, Patel B, Xiao S, Zhu C, Xia J, Tan TG, Sefik E, Yajnik V, Sharpe AH, Quintana FJ, Mathis D, Benoist C, Hafler DA, Kuchroo VK. Treg cells expressing the coinhibitory molecule TIGIT selectively inhibit proinflammatory Th1 and Th17 cell responses. *Immunity.* 2014;40:569–81.
130. Jordan MS, Boesteanu A, Reed AJ, Petrone AL, Holenbeck AE, Lerman MA, Najj A, Caton AJ. Thymic selection of CD4(+)CD25(+) regulatory T cells induced by an agonist self-peptide. *Nat Immunol.* 2001;2:301–6.
131. Croft M. The role of TNF superfamily members in T-cell function and diseases. *Nat Rev Immunol.* 2009;9:271–85.
132. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer.* 2012;12:252–64.
133. Nagasawa T. Microenvironmental niches in the bone marrow required for B-cell development. *Nat Rev Immunol.* 2006;6:107–16.
134. Hardy RR, Carmack CE, Shinton SA, Kemp JD, Hayakawa K. Resolution and characterization of pro-B and pre-pro-B cell stages in normal mouse bone marrow. *J Exp Med.* 1991;173:1213–25.
135. Tiegs SL, Russell DM, Nemazee D. Receptor editing in self-reactive bone marrow B cells. *J Exp Med.* 1993;177:1009–20.
136. Carsetti R, Kohler G, Lamers MC. Transitional B cells are the target of negative selection in the B cell compartment. *J Exp Med.* 1995;181:2129–40.
137. Pieper K, Grimbacher B, Eibel H. B-cell biology and development. *J Allergy Clin Immunol.* 2013;131:959–71.
138. Shlomchik MJ, Weisel F. Germinal center selection and the development of memory B and plasma cells. *Immunol Rev.* 2012;247:52–63.
139. Schroeder HW Jr, Cavacini L. Structure and function of immunoglobulins. *J Allergy Clin Immunol.* 2010;125:S41–52.
140. Lipman NS, Jackson LR, Trudel LJ, Weis-Garcia F. Monoclonal versus polyclonal antibodies: distinguishing characteristics, applications, and information resources. *ILAR J.* 2005;46:258–68.
141. Kung P, Goldstein G, Reinherz EL, Schlossman SF. Monoclonal antibodies defining distinctive human T cell surface antigens. *Science.* 1979;206:347–9.
142. Morrison SL, Johnson MJ, Herzenberg LA, Oi VT. Chimeric human antibody molecules: mouse antigen-binding domains with human constant region domains. *Proc Natl Acad Sci U S A.* 1984;81:6851–5.
143. Riechmann L, Clark M, Waldmann H, Winter G. Reshaping human antibodies for therapy. *Nature.* 1988;332:323–7.
144. Harding FA, Stickler MM, Razo J, DuBridgde RB. The immunogenicity of humanized and fully human antibodies: residual immunogenicity resides in the CDR regions. *MAbs.* 2010;2:256–65.
145. Scott AM, Wolchok JD, Old LJ. Antibody therapy of cancer. *Nat Rev Cancer.* 2012;12:278–87.
146. Fridman WH, Pages F, Sautes-Fridman C, Galon J. The immune contexture in human tumours: impact on clinical outcome. *Nat Rev Cancer.* 2012;12:298–306.

147. Dunn GP, Old LJ, Schreiber RD. The three Es of cancer immunoediting. *Annu Rev Immunol.* 2004;22:329–60.
148. Dunn GP, Bruce AT, Ikeda H, Old LJ, Schreiber RD. Cancer immunoediting: from immunosurveillance to tumor escape. *Nat Immunol.* 2002;3:991–8.
149. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell.* 2000;100:57–70.
150. Teng MW, Galon J, Fridman WH, Smyth MJ. From mice to humans: developments in cancer immunoediting. *J Clin Invest.* 2015;125:3338–46.
151. Mellman I, Coukos G, Dranoff G. Cancer immunotherapy comes of age. *Nature.* 2011;480:480–9.
152. Gabrilovich DI, Ostrand-Rosenberg S, Bronte V. Coordinated regulation of myeloid cells by tumours. *Nat Rev Immunol.* 2012;12:253–68.
153. Huang B, Pan PY, Li QS, Sato AI, Levy DE, Bromberg J, Divino CM, Chen SH. Gr-1(+) CD115(+) immature myeloid suppressor cells mediate the development of tumor-induced T regulatory cells and T-cell anergy in tumor-bearing host. *Cancer Res.* 2006;66:1123–31.
154. Lindau D, Gielen P, Kroesen M, Wesseling P, Adema GJ. The immunosuppressive tumour network: myeloid-derived suppressor cells, regulatory T cells and natural killer T cells. *Immunology.* 2013;138:105–15.
155. Tangri S, LiCalsi C, Sidney J, Sette A. Rationally engineered proteins or antibodies with absent or reduced immunogenicity. *Curr Med Chem.* 2002;9:2191–9.
156. Weiner LM, Surana R, Wang SZ. Monoclonal antibodies: versatile platforms for cancer immunotherapy. *Nat Rev Immunol.* 2010;10:317–27.
157. Krummel MF, Allison JP. Cd28 and Ctl4 have opposing effects on the response of T-cells to stimulation. *J Exp Med.* 1995;182:459–65.
158. Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, Haanen JB, Gonzalez R, Robert C, Schadendorf D, Hassel JC, Akerley W, van den Eertwegh AJM, Lutzky J, Lorigan P, Vaubel JM, Linette GP, Hogg D, Ottensmeier CH, Lebke C, Peschel C, Quirt I, Clark JI, Wolchok JD, Weber JS, Tian J, Yellin MJ, Nichol GM, Hoos A, Urba WJ. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med.* 2010;363:711–23.
159. Brahmer JR, Drake CG, Wollner I, Powderly JD, Picus J, Sharfman WH, Stankevich E, Pons A, Salay TM, McMiller TL, Gilson MM, Wang C, Selby M, Taube JM, Anders R, Chen L, Korman AJ, Pardoll DM, Lowy I, Topalian SL. Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: safety, clinical activity, pharmacodynamics, and immunologic correlates. *J Clin Oncol.* 2010;28:3167–75.
160. Borghaei H, Paz-Ares L, Horn L, Spigel DR, Steins M, Ready NE, Chow LQ, Vokes EE, Felip E, Holgado E, Barlesi F, Kohlhaufl M, Arrieta O, Burgio MA, Fayette J, Lena H, Poddubskaya E, Gerber DE, Gettinger SN, Rudin CM, Rizvi N, Crino L, Blumenschein GR Jr, Antonia SJ, Dorange C, Harbison CT, Graf Finckenstein F, Brahmer JR. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med.* 2015;373:1627–39.
161. Brahmer J, Reckamp KL, Baas P, Crino L, Eberhardt WE, Poddubskaya E, Antonia S, Pluzanski A, Vokes EE, Holgado E, Waterhouse D, Ready N, Gainor J, Aren Frontera O, Havel L, Steins M, Garassino MC, Aerts JG, Domine M, Paz-Ares L, Reck M, Baudalet C, Harbison CT, Lestini B, Spigel DR. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med.* 2015;373:123–35.
162. Herbst RS, Baas P, Kim DW, Felip E, Perez-Gracia JL, Han JY, Molina J, Kim JH, Arvis CD, Ahn MJ, Majem M, Fidler MJ, de Castro G Jr, Garrido M, Lubiniecki GM, Shentu Y, Im E, Dolled-Filhart M, Garon EB. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet.* 2016;387:1540–50.
163. Rosenberg JE, Hoffman-Censits J, Powles T, van der Heijden MS, Balar AV, Necchi A, Dawson N, O'Donnell PH, Balmanoukian A, Loriot Y, Srinivas S, Retz MM, Grivas P, Joseph RW, Galsky MD, Fleming MT, Petrylak DP, Perez-Gracia JL, Burris HA, Castellano D, Canil C, Bellmunt J, Bajorin D, Nickles D, Bourgon R, Frampton GM, Cui N, Mariathasan S, Abidoye O, Fine GD, Dreicer R. Atezolizumab in patients with locally advanced and

- metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet*. 2016;387:1909–20.
164. U.S. Food and Drug Administration. Hematology/oncology (cancer) approvals & safety notifications. 2018.
 165. Moon YW, Hajjar J, Hwu P, Naing A. Targeting the indoleamine 2,3-dioxygenase pathway in cancer. *J Immunother Cancer*. 2015;3:51.
 166. Koblisch HK, Hansbury MJ, Bowman KJ, Yang G, Neilan CL, Haley PJ, Burn TC, Waeltz P, Sparks RB, Yue EW, Combs AP, Scherle PA, Vaddi K, Fridman JS. [INCB preclin] Hydroxyamidine inhibitors of indoleamine-2,3-dioxygenase potently suppress systemic tryptophan catabolism and the growth of IDO-expressing tumors. *Mol Cancer Ther*. 2010;9:489–98.
 167. Liu X, Shin N, Koblisch HK, Yang G, Wang Q, Wang K, Leffet L, Hansbury MJ, Thomas B, Rupar M, Waeltz P, Bowman KJ, Polam P, Sparks RB, Yue EW, Li Y, Wynn R, Fridman JS, Burn TC, Combs AP, Newton RC, Scherle PA. [INCB preclin] selective inhibition of IDO1 effectively regulates mediators of antitumor immunity. *Blood*. 2010;115:3520–30.
 168. Metz R, Rust S, Duhadaway JB, Mautino MR, Munn DH, Vahanian NN, Link CJ, Prendergast GC. [Indoximod preclin] IDO inhibits a tryptophan sufficiency signal that stimulates mTOR: a novel IDO effector pathway targeted by D-1-methyl-tryptophan. *Oncoimmunology*. 2012;1:1460–8.
 169. Iversen TZ, Engell-Noerregaard L, Ellebaek E, Andersen R, Larsen SK, Bjoern J, Zeyher C, Gouttefangeas C, Thomsen BM, Holm B, Straten PT, Mellemegaard A, Andersen MH, Svane IM. [IDO pep vac] Long-lasting disease stabilization in the absence of toxicity in metastatic lung cancer patients vaccinated with an epitope derived from indoleamine 2,3 dioxygenase. *Clin Cancer Res*. 2014;20:221–32.
 170. Siu LL, Gelmon K, Chu Q, Pachynski R, Alese O, Basciano P, Walker J, Mitra P, Zhu L, Phillips P, Hunt J, Desai J. Abstract CT116: BMS-986205, an optimized indoleamine 2,3-dioxygenase 1 (IDO1) inhibitor, is well tolerated with potent pharmacodynamic (PD) activity, alone and in combination with nivolumab (nivo) in advanced cancers in a phase 1/2a trial. *Cancer Res*. 2017;77:CT116-CT.
 171. Mautino MR, Jaipuri FA, Waldo J, Kumar S, Adams J, Allen CV, Marciniowicz-Flick A, Munn D, Vahanian N, Link CJJ. NLG919, a novel indoleamine-2,3-dioxygenase (IDO)-pathway inhibitor drug candidate for cancer therapy. *AACR*; 2013. p. 491.
 172. Linch SN, McNamara MJ, Redmond WL. OX40 agonists and combination immunotherapy: putting the pedal to the metal. *Front Oncol*. 2015;5:34.
 173. Aspeslagh S, Postel-Vinay S, Rusakiewicz S, Soria JC, Zitvogel L, Marabelle A. Rationale for anti-OX40 cancer immunotherapy. *Eur J Cancer*. 2016;52:50–66.
 174. Topalian SL, Weiner GJ, Pardoll DM. Cancer immunotherapy comes of age. *J Clin Oncol*. 2011;29:4828–36.
 175. Wargo JA, Cooper ZA, Flaherty KT. Universes collide: combining immunotherapy with targeted therapy for cancer. *Cancer Discov*. 2014;4:1377–86.
 176. Sullivan RJ, Gonzalez R, Lewis KD, Hamid O, Infante JR, Patel MR, Hodi FS, Wallin J, Pitcher B, Cha E, Roberts L, Ballinger M, Hwu P. Atezolizumab (a) + cobimetinib (C) + vemurafenib (V) in BRAFV600-mutant metastatic melanoma (mel): updated safety and clinical activity. *J Clin Oncol*. 2017;35:3063.
 177. Formenti SC, Demaria S. Combining radiotherapy and cancer immunotherapy: a paradigm shift. *J Natl Cancer Inst*. 2013;105:256–65.
 178. Golden EB, Chachoua A, Fenton-Kerimian MB, Demaria S, Formenti SC. Abscopal responses in metastatic non-small cell lung cancer (NSCLC) patients treated on a phase 2 study of combined radiation therapy and ipilimumab: evidence for the in situ vaccination hypothesis of radiation. *Int J Radiat Oncol*. 2015;93:S66–S7.
 179. Fiorica F, Belluomini L, Stefanelli A, Santini A, Urbini B, Giorgi C, Frassoldati A. Immune checkpoint inhibitor nivolumab and radiotherapy in pretreated lung cancer patients: efficacy and safety of combination. *Am J Clin Oncol*. 2018;41:1101–05.

180. Cai Z, Sanchez A, Shi Z, Zhang T, Liu M, Zhang D. Activation of toll-like receptor 5 on breast cancer cells by flagellin suppresses cell proliferation and tumor growth. *Cancer Res.* 2011;71:2466–75.
181. Wolska A, Lech-Maranda E, Robak T. Toll-like receptors and their role in carcinogenesis and anti-tumor treatment. *Cell Mol Biol Lett.* 2009;14:248–72.
182. Liu Y, Yan W, Tohme S, Chen M, Fu Y, Tian D, Lotze M, Tang DL, Tsung A. Hypoxia induced HMGB1 and mitochondrial DNA interactions mediate tumor growth in hepatocellular carcinoma through toll-like receptor 9. *J Hepatol.* 2015;63:114–21.
183. Shi M, Chen X, Ye K, Yao Y, Li Y. Application potential of toll-like receptors in cancer immunotherapy: systematic review. *Medicine (Baltimore).* 2016;95:e3951.
184. Li K, Qu S, Chen X, Wu Q, Shi M: Promising targets for cancer immunotherapy: TLRs, RLRs, and STING-mediated innate immune pathways. *Int J Mol Sci* 2017;18. pii: E404.
185. Ishikawa H, Ma Z, Barber GN. STING regulates intracellular DNA-mediated, type I interferon-dependent innate immunity. *Nature.* 2009;461:788–92.
186. Fuertes MB, Kacha AK, Kline J, Woo SR, Kranz DM, Murphy KM, Gajewski TF. Host type I IFN signals are required for antitumor CD8+ T cell responses through CD8 α + dendritic cells. *J Exp Med.* 2011;208:2005–16.
187. Corrales L, Glickman LH, McWhirter SM, Kanne DB, Sivick KE, Katibah GE, Woo SR, Lemmens E, Banda T, Leong JJ, Metchette K, Dubensky TW Jr, Gajewski TF. Direct activation of STING in the tumor microenvironment leads to potent and systemic tumor regression and immunity. *Cell Rep.* 2015;11:1018–30.
188. Fu J, Kanne DB, Leong M, Glickman LH, McWhirter SM, Lemmens E, Mechette K, Leong JJ, Lauer P, Liu W, Sivick KE, Zeng Q, Soares KC, Zheng L, Portnoy DA, Woodward JJ, Pardoll DM, Dubensky TW, Kim Y. STING agonist formulated cancer vaccines can cure established tumors resistant to PD-1 blockade. *Sci Transl Med.* 2015;7:283ra52.
189. Deng LF, Liang H, Xu M, Yang XM, Burnette B, Arina A, Li XD, Mauceri H, Beckett M, Darga T, Huang XN, Gajewski TF, Chen ZJJ, Fu YX, Weichselbaum RR. STING-dependent cytosolic DNA sensing promotes radiation-induced type I interferon-dependent antitumor immunity in immunogenic tumors. *Immunity.* 2014;41:843–52.
190. Topalian SL, Taube JM, Anders RA, Pardoll DM. Mechanism-driven biomarkers to guide immune checkpoint blockade in cancer therapy. *Nat Rev Cancer.* 2016;16:275–87.
191. Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF, Powderly JD, Carvajal RD, Sosman JA, Atkins MB, Leming PD, Spigel DR, Antonia SJ, Horn L, Drake CG, Pardoll DM, Chen L, Sharfman WH, Anders RA, Taube JM, McMiller TL, Xu H, Korman AJ, Jure-Kunkel M, Agrawal S, McDonald D, Kollia GD, Gupta A, Wigginton JM, Sznol M. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med.* 2012;366:2443–54.
192. U.S. Food and Drug Administration. FDA approves Keytruda for advanced non-small cell lung cancer. 2015. <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm465444.htm>.
193. Herbst RS, Soria JC, Kowanetz M, Fine GD, Hamid O, Gordon MS, Sosman JA, McDermott DF, Powderly JD, Gettinger SN, Kohrt HE, Horn L, Lawrence DP, Rost S, Leabman M, Xiao Y, Mokatrinn A, Koeppen H, Hegde PS, Mellman I, Chen DS, Hodi FS. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature.* 2014;515:563–7.
194. Rittmeyer A, Barlesi F, Waterkamp D, Park K, Ciardiello F, von Pawel J, Gadgeel SM, Hida T, Kowalski DM, Dols MC, Cortinovis DL, Leach J, Polikoff J, Barrios C, Kabbinar F, Frontera OA, De Marinis F, Turna H, Lee JS, Ballinger M, Kowanetz M, He P, Chen DS, Sandler A, Gandara DR, Grp OS. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet.* 2017;389:255–65.
195. Madore J, Vilain RE, Menzies AM, Kakavand H, Wilmott JS, Hyman J, Yearley JH, Kefford RF, Thompson JF, Long GV, Hersey P, Scolyer RA. PD-L1 expression in melanoma shows

- marked heterogeneity within and between patients: implications for anti-PD-1/PD-L1 clinical trials. *Pigment Cell Melanoma Res.* 2015;28:245–53.
196. Rosell R, Palmero R. PD-L1 expression associated with better response to EGFR tyrosine kinase inhibitors. *Cancer Biol Med.* 2015;12:71–3.
 197. Sharma P, Allison JP. The future of immune checkpoint therapy. *Science.* 2015;348:56–61.
 198. Hadrup S, Donia M, Thor Straten P. Effector CD4 and CD8 T cells and their role in the tumor microenvironment. *Cancer Microenviron.* 2013;6:123–33.
 199. Zhang L, Conejo-Garcia JR, Katsaros D, Gimotty PA, Massobrio M, Regnani G, Makrigiannakis A, Gray H, Schlienger K, Liebman MN, Rubin SC, Coukos G. Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med.* 2003;348:203–13.
 200. Ruffini E, Asioli S, Filosso PL, Lyberis P, Bruna MC, Macri L, Daniele L, Oliaro A. Clinical significance of tumor-infiltrating lymphocytes in lung neoplasms. *Ann Thorac Surg.* 2009;87:365–71; discussion 371–2
 201. Curiel TJ, Coukos G, Zou L, Alvarez X, Cheng P, Mottram P, Evdemon-Hogan M, Conejo-Garcia JR, Zhang L, Burow M, Zhu Y, Wei S, Kryczek I, Daniel B, Gordon A, Myers L, Lackner A, Disis ML, Knutson KL, Chen L, Zou W. Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival. *Nat Med.* 2004;10:942–9.
 202. Gobert M, Treilleux I, Bendriss-Vermare N, Bachelot T, Goddard-Leon S, Arfi V, Biota C, Doffin AC, Durand I, Olive D, Perez S, Pasqual N, Faure C, Coquard IR, Puisieux A, Caux C, Blay JY, Menetrier-Caux C. Regulatory T cells recruited through CCL22/CCR4 are selectively activated in lymphoid infiltrates surrounding primary breast tumors and lead to an adverse clinical outcome. *Cancer Res.* 2009;69:2000–9.
 203. Fu JL, Xu DP, Liu ZW, Shi M, Zhao P, Fu BY, Zhang Z, Yang HY, Zhang H, Zhou CB, Ya JX, Jin L, Wang HF, Yang YP, Fu YX, Wang FS. Increased regulatory T cells correlate with CD8 T-cell impairment and poor survival in hepatocellular carcinoma patients. *Gastroenterology.* 2007;132:2328–39.
 204. Llosa NJ, Cruise M, Tam A, Wicks EC, Hechenbleikner EM, Taube JM, Blosser RL, Fan HN, Wang H, Lubber BS, Zhang M, Papadopoulos N, Kinzler KW, Vogelstein B, Sears CL, Anders RA, Pardoll DM, Housseau F. The vigorous immune microenvironment of microsatellite instable colon cancer is balanced by multiple counter-inhibitory checkpoints. *Cancer Discov.* 2015;5:43–51.
 205. Galon J, Costes A, Sanchez-Cabo F, Kirilovsky A, Mlecnik B, Lagorce-Pages C, Tosolini M, Camus M, Berger A, Wind P, Zinzindohoue F, Bruneval P, Cugnenc PH, Trajanoski Z, Fridman WH, Pages F. Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science.* 2006;313:1960–4.
 206. Tumeh PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJ, Robert L, Chmielowski B, Spasic M, Henry G, Ciobanu V, West AN, Carmona M, Kivork C, Seja E, Cherry G, Gutierrez AJ, Grogan TR, Mateus C, Tomasic G, Glaspy JA, Emerson RO, Robins H, Pierce RH, Elashoff DA, Robert C, Ribas A. PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature.* 2014;515:568–71.
 207. Hamid O, Schmidt H, Nissan A, Ridolfi L, Aamdal S, Hansson J, Guida M, Hyams DM, Gomez H, Bastholt L, Chasalow SD, Berman D. A prospective phase II trial exploring the association between tumor microenvironment biomarkers and clinical activity of ipilimumab in advanced melanoma. *J Transl Med.* 2011;9:204.
 208. Martens A, Wistuba-Hamprecht K, Yuan J, Postow MA, Wong P, Capone M, Madonna G, Khammari A, Schilling B, Sucker A, Schadendorf D, Martus P, Dreno B, Ascierto PA, Wolchok JD, Pawelec G, Garbe C, Weide B. Increases in absolute lymphocytes and circulating CD4+ and CD8+ T cells are associated with positive clinical outcome of melanoma patients treated with ipilimumab. *Clin Cancer Res.* 2016;22:4848–58.
 209. Galon J, Mlecnik B, Bindea G, Angell HK, Berger A, Lagorce C, Lugli A, Zlobec I, Hartmann A, Bifulco C, Nagtegaal ID, Palmqvist R, Masucci GV, Botti G, Tatangelo F, Delrio P, Maio M, Laghi L, Grizzi F, Asslaber M, D'Arrigo C, Vidal-Vanaclocha F, Zavadova E, Chouchane L,

- Ohashi PS, Hafezi-Bakhtiari S, Wouters BG, Roehrl M, Nguyen L, Kawakami Y, Hazama S, Okuno K, Ogino S, Gibbs P, Waring P, Sato N, Torigoe T, Itoh K, Patel PS, Shukla SN, Wang YL, Kopetz S, Sinicrope FA, Scripcariu V, Ascierto PA, Marincola FM, Fox BA, Pages F. Towards the introduction of the 'Immunoscore' in the classification of malignant tumours. *J Pathol.* 2014;232:199–209.
210. Mlecnik B, Van den Eynde M, Bindea G, Church SE, Vasaturo A, Fredriksen T, Lafontaine L, Haicheur N, Marliot F, Debetancourt D, Pairet G, Jouret-Mourin A, Gigot JF, Hubert C, Danse E, Dragean C, Carrasco J, Humblet Y, Valge-Archer V, Berger A, Pages F, Machiels JP, Galon J. Comprehensive intrametastatic immune quantification and major impact of immunoscore on survival. *J Natl Cancer Inst.* 2018;110:97–108.
211. Pagès F, Mlecnik B, Marliot F, Bindea G, Ou F-S, Bifulco C, Lugli A, Zlobec I, Rau TT, Berger MD, Nagtegaal ID, Vink-Börger E, Hartmann A, Geppert C, Kolwelter J, Merkel S, Grützmann R, Van den Eynde M, Jouret-Mourin A, Kartheuser A, Léonard D, Remue C, Wang JY, Bavi P, MHA R, Ohashi PS, Nguyen LT, Han S, HL MG, Hafezi-Bakhtiari S, Wouters BG, Masucci GV, Andersson EK, Zavadova E, Vocka M, Spacek J, Petruzella L, Konopasek B, Dunder P, Skalova H, Nemejcova K, Botti G, Tatangelo F, Delrio P, Ciliberto G, Maio M, Laghi L, Grizzi F, Fredriksen T, Buttard B, Angelova M, Vasaturo A, Maby P, Church SE, Angell HK, Lafontaine L, Bruni D, El Sissy C, Haicheur N, Kirilovsky A, Berger A, Lagorce C, Meyers JP, Paustian C, Feng Z, Ballesteros-Merino C, Dijkstra J, van de Water C, van Lent-van Vliet S, Knijn N, Muşină A-M, Scripcariu D-V, Popivanova B, Xu M, Fujita T, Hazama S, Suzuki N, Nagano H, Okuno K, Torigoe T, Sato N, Furuhashi T, Takemasa I, Itoh K, Patel PS, Vora HH, Shah B, Patel JB, Rajvik KN, Pandya SJ, Shukla SN, Wang Y, Zhang G, Kawakami Y, Marincola FM, Ascierto PA, Sargent DJ, Fox BA, Galon J. International validation of the consensus immunoscore for the classification of colon cancer: a prognostic and accuracy study. *Lancet.* 2018;391:2128–39.
212. Haymaker CL, Kim D, Uemura M, Vence LM, Phillip A, McQuail N, Brown PD, Fernandez I, Hudgens CW, Creasy C, Hwu WJ, Sharma P, Tetzlaff MT, Allison JP, Hwu P, Bernatchez C, Diab A. Metastatic melanoma patient had a complete response with clonal expansion after whole brain radiation and PD-1 blockade. *Cancer Immunol Res.* 2017;5:100–5.
213. Olugbile S, Park J-H, Hoffman P, Szeto L, Patel J, Vigneswaran WT, Vokes E, Nakamura Y, Klyotani K. Sustained oligoclonal T cell expansion correlates with durable response to immune checkpoint blockade in lung cancer. *J Cancer Sci Ther.* 2017;9:717–22.
214. Inoue H, Park JH, Kiyotani K, Zewde M, Miyashita A, Jinnin M, Kuniwa Y, Okuyama R, Tanaka R, Fujisawa Y, Kato H, Morita A, Asai J, Katoh N, Yokota K, Akiyama M, Ihn H, Fukushima S, Nakamura Y. Intratumoral expression levels of PD-L1, GZMA, and HLA-A along with oligoclonal T cell expansion associate with response to nivolumab in metastatic melanoma. *Oncoimmunology.* 2016;5:e1204507.
215. Rizvi NA, Hellmann MD, Snyder A, Kvistborg P, Makarov V, Havel JJ, Lee W, Yuan JD, Wong P, Ho TS, Miller ML, Rekhtman N, Moreira AL, Ibrahim F, Bruggeman C, Gasmfi B, Zappasodi R, Maeda Y, Sander C, Garon EB, Merghoub T, Wolchok JD, Schumacher TN, Chan TA. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science.* 2015;348:124–8.
216. Snyder A, Makarov V, Merghoub T, Yuan J, Zaretsky JM, Desrichard A, Walsh LA, Postow MA, Wong P, Ho TS, Hollmann TJ, Bruggeman C, Kannan K, Li Y, Elipenahli C, Liu C, Harbison CT, Wang L, Ribas A, Wolchok JD, Chan TA. Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N Engl J Med.* 2014;371:2189–99.
217. Hugo W, Zaretsky JM, Sun L, Song C, Moreno BH, Hu-Lieskovan S, Berent-Maoz B, Pang J, Chmielowski B, Cherry G, Seja E, Lomeli S, Kong X, Kelley MC, Sosman JA, Johnson DB, Ribas A, Lo RS. Genomic and transcriptomic features of response to anti-PD-1 therapy in metastatic melanoma. *Cell.* 2016;165:35–44.
218. Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, Skora AD, Luber BS, Azad NS, Laheru D, Biedrzycki B, Donehower RC, Zaheer A, Fisher GA, Crocenzi TS, Lee JJ, Duffy SM, Goldberg RM, de la Chapelle A, Koshiji M, Bhaijee F, Huebner T, Hruban RH, Wood LD, Cuka N, Pardoll DM, Papadopoulos N, Kinzler KW, Zhou S, Cornish TC, Taube

- JM, Anders RA, Eshleman JR, Vogelstein B, Diaz LA Jr. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med*. 2015;372:2509–20.
219. Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, Bignell GR, Bolli N, Borg A, Borresen-Dale AL, Boyault S, Burkhardt B, Butler AP, Caldas C, Davies HR, Desmedt C, Eils R, Eyfjord JE, Foekens JA, Greaves M, Hosoda F, Hutter B, Ilcic T, Imbeaud S, Imielinski M, Jager N, Jones DT, Jones D, Knappskog S, Kool M, Lakhani SR, Lopez-Otin C, Martin S, Munshi NC, Nakamura H, Northcott PA, Pajic M, Papaemmanuil E, Paradiso A, Pearson JV, Puente XS, Raine K, Ramakrishna M, Richardson AL, Richter J, Rosenstiel P, Schlesner M, Schumacher TN, Span PN, Teague JW, Totoki Y, Tutt AN, Valdes-Mas R, van Buuren MM, van't Veer L, Vincent-Salomon A, Waddell N, Yates LR, Zucman-Rossi J, Futreal PA, McDermott U, Lichter P, Meyerson M, Grimmond SM, Siebert R, Campo E, Shibata T, Pfister SM, Campbell PJ, Stratton MR. Signatures of mutational processes in human cancer. *Nature*. 2013;500:415–21.
 220. Lastwika KJ, Wilson W, Li QK, Norris J, Xu HY, Ghazarian SR, Kitagawa H, Kawabata S, Taube JM, Yao S, Liu LN, Gills JJ, Dennis PA. Control of PD-L1 expression by oncogenic activation of the AKT-mTOR pathway in non-small cell lung cancer. *Cancer Res*. 2016;76:227–38.
 221. Ribas A, Robert C, Hodi FS, Wolchok JD, Joshua AM, Hwu WJ, Weber JS, Zarour HM, Kefford R, Loboda A, Albright A, Kang SP, Ebbinghaus S, Yearley J, Murphy E, Nebozhyn M, Lunceford JK, McClanahan T, Ayers M, Daud A. Association of response to programmed death receptor 1 (PD-1) blockade with pembrolizumab (MK-3475) with an interferon-inflammatory immune gene signature. *J Clin Oncol*. 2015;33.
 222. Ayers M, Lunceford J, Nebozhyn M, Murphy E, Loboda A, Albright A, Cheng J, Kang SP, Ebbinghaus S, Yearley J, Shankaran V, Seiwert T, Ribas A, McClanahan T. Relationship between immune gene signatures and clinical response to PD-1 blockade with pembrolizumab (MK-3475) in patients with advanced solid tumors. *J Immunother*. 2015;3:P80.
 223. Higgs BW, Morehouse C, Streicher K, Rebelatto MC, Steele K, Jin X, Pilataxi F, Brohawn PZ, Blake-Haskins JA, Gupta AK, Ranade K. Relationship of baseline tumoral IFN γ mRNA and PD-L1 protein expression to overall survival in durvalumab-treated NSCLC patients. *J Clin Oncol*. 2016;34:3036.
 224. Fehrenbacher L, Spira A, Ballinger M, Kowanzet M, Vansteenkiste J, Mazieres J, Park K, Smith D, Artal-Cortes A, Lewanski C, Braiteh F, Waterkamp D, He P, Zou W, Chen DS, Yi J, Sandler A, Rittmeyer A, Group PS. Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial. *Lancet*. 2016;387:1837–46.
 225. Blank CU, Haanen JB, Ribas A, Schumacher TN. Cancer immunology. The “cancer immunogram”. *Science*. 2016;352:658–60.
 226. Karasaki T, Nagayama K, Kuwano H, Nitadori JI, Sato M, Anraku M, Hosoi A, Matsushita H, Morishita Y, Kashiwabara K, Takazawa M, Ohara O, Kakimi K, Nakajima J. An immunogram for the cancer-immunity cycle: towards personalized immunotherapy of lung cancer. *J Thorac Oncol*. 2017;12:791–803.
 227. Martens A, Wistuba-Hamprecht K, Geukes Foppen M, Yuan J, Postow MA, Wong P, Romano E, Khammari A, Dreno B, Capone M, Ascierto PA, Di Giacomo AM, Maio M, Schilling B, Sucker A, Schadendorf D, Hassel JC, Eigentler TK, Martus P, Wolchok JD, Blank C, Pawelec G, Garbe C, Weide B. Baseline peripheral blood biomarkers associated with clinical outcome of advanced melanoma patients treated with ipilimumab. *Clin Cancer Res*. 2016;22:2908–18.
 228. Hopkins AM, Rowland A, Kichenadasse G, Wiese MD, Gurney H, McKinnon RA, Karapetis CS, Sorich MJ. Predicting response and toxicity to immune checkpoint inhibitors using routinely available blood and clinical markers. *Br J Cancer*. 2017;117:913–20.
 229. Manson G, Norwood J, Marabelle A, Kohrt H, Houot R. Biomarkers associated with checkpoint inhibitors. *Ann Oncol*. 2016;27:1199–206.
 230. Delyon J, Mateus C, Lefeuvre D, Lanoy E, Zitvogel L, Chaput N, Roy S, Eggermont AMM, Routier E, Robert C. Experience in daily practice with ipilimumab for the treatment of patients with metastatic melanoma: an early increase in lymphocyte and eosinophil counts is associated with improved survival. *Ann Oncol*. 2013;24:1697–703.

231. Ku GY, Yuan JD, Page DB, Schroeder SEA, Panageas KS, Carvajal RD, Chapman PB, Schwartz GK, Allison JP, Wolchok JD. Single-institution experience with ipilimumab in advanced melanoma patients in the compassionate use setting lymphocyte count after 2 doses correlates with survival. *Cancer*. 2010;116:1767–75.
232. Wilgenhof S, Du Four S, Vandenbroucke F, Everaert H, Salmon I, Lienard D, Marmol VD, Neyns B. Single-center experience with ipilimumab in an expanded access program for patients with pretreated advanced melanoma. *J Immunother*. 2013;36:215–22.
233. Di Giacomo AM, Danielli R, Calabro L, Bertocci E, Nannicini C, Giannarelli D, Balestrazzi A, Vigni F, Riversi V, Miracco C, Biagioli M, Altomonte M, Maio M. Ipilimumab experience in heavily pretreated patients with melanoma in an expanded access program at the University Hospital of Siena (Italy). *Cancer Immunol Immunother*. 2011;60:467–77.
234. Simeone E, Gentilcore G, Giannarelli D, Grimaldi AM, Caraco C, Curvietto M, Esposito A, Paone M, Palla M, Cavalcanti E, Sandomenico F, Petrillo A, Botti G, Fulcinitti F, Palmieri G, Queirolo P, Marchetti P, Ferraresi V, Rinaldi G, Pistillo MP, Ciliberto G, Mozzillo N, Ascierto PA. Immunological and biological changes during ipilimumab treatment and their potential correlation with clinical response and survival in patients with advanced melanoma. *Cancer Immunol Immunother*. 2014;63:675–83.
235. Gebhardt C, Sevko A, Jiang HH, Lichtenberger R, Reith M, Tarnanidis K, Holland-Letz T, Umansky L, Beckhove P, Sucker A, Schadendorf D, Utikal J, Umansky V. Myeloid cells and related chronic inflammatory factors as novel predictive markers in melanoma treatment with ipilimumab. *Clin Cancer Res*. 2015;21:5453–9.
236. Kelderman S, Heemskerk B, van Tinteren H, van den Brom RR, Hospers GA, van den Eertwegh AJ, Kapiteijn EW, de Groot JW, Soetekouw P, Jansen RL, Fiets E, Furness AJ, Renn A, Krzystanek M, Szallasi Z, Lorigan P, Gore ME, Schumacher TN, Haanen JB, Larkin JM, Blank CU. Lactate dehydrogenase as a selection criterion for ipilimumab treatment in metastatic melanoma. *Cancer Immunol Immunother*. 2014;63:449–58.
237. Lee JH, Long GV, Boyd S, Lo S, Menzies AM, Tembe V, Guminski A, Jakrot V, Scolyer RA, Mann GJ, Kefford RF, Carlino MS, Rizos H. Circulating tumour DNA predicts response to anti-PD1 antibodies in metastatic melanoma. *Ann Oncol*. 2017;28:1130–6.
238. Kamphorst AO, Pillai RN, Yang S, Nasti TH, Akondy RS, Wieland A, Sica GL, Yu K, Koenig L, Patel NT, Behera M, Wu H, McCausland M, Chen ZJ, Zhang C, Khuri FR, Owonikoko TK, Ahmed R, Ramalingam SS. Proliferation of PD-1+CD8 T cells in peripheral blood after PD-1-targeted therapy in lung cancer patients. *Proc Natl Acad Sci U S A*. 2017;114:4993–8.
239. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpnits TV, Prieto PA, Vicente D, Hoffman K, Wei SC, Cogdill AP, Zhao L, Hudgens CW, Hutchinson DS, Manzo T, de Macedo MP, Cotechini T, Kumar T, Chen WS, Reddy SM, Sloane RS, Galloway-Pena J, Jiang H, Chen PL, Shpall EJ, Rezvani K, Alousi AM, Chemaly RF, Shelburne S, Vence LM, Okhuysen PC, Jensen VB, Swennes AG, McAllister F, Sanchez EMR, Zhang Y, Le Chatelier E, Zitvogel L, Pons N, Austin-Breneman JL, Haydu LE, Burton EM, Gardner JM, Sirmans E, Hu J, Lazar AJ, Tsujikawa T, Diab A, Tawbi H, Glitza IC, Hwu WJ, Patel SP, Woodman SE, Amaria RN, Davies MA, Gershenwald JE, Hwu P, Lee JE, Zhang J, Coussens LM, Cooper ZA, Futreal PA, Daniel CR, Ajami NJ, Petrosino JF, Tetzlaff MT, Sharma P, Allison JP, Jenq RR, Wargo JA. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science*. 2018;359:97–103.
240. Topalian SL, Taube JM, Anders RA, Pardoll DM. Mechanism-driven biomarkers to guide immune checkpoint blockade in cancer therapy. *Nat Rev Cancer*. 2016;16:275–87.