

Interpolation Between Hilbert Spaces



Yacin Ameur

Abstract This note comprises a synthesis of certain results in the theory of exact interpolation between Hilbert spaces. In particular, we examine various characterizations of interpolation spaces and their relations to a number of results in operator theory and in function theory.

Keywords Interpolation · Hilbert space · Calderón pair · Pick function · Matrix monotonicity

Mathematics Subject Classification (2010) 46B70, 47A57, 47A63

1 Interpolation Theoretic Notions

1.1 Interpolation Norms

When X, Y are normed spaces, we use the symbol $\mathcal{L}(X; Y)$ to denote the totality of bounded linear maps $T : X \rightarrow Y$ with the operator norm

$$\|T\|_{\mathcal{L}(X;Y)} = \sup \{ \|Tx\|_Y ; \|x\|_X \leq 1 \}.$$

When $X = Y$ we simply write $\mathcal{L}(X)$.

Consider a pair of Hilbert spaces $\overline{\mathcal{H}} = (\mathcal{H}_0, \mathcal{H}_1)$ which is *regular* in the sense that $\mathcal{H}_0 \cap \mathcal{H}_1$ is dense in \mathcal{H}_0 as well as in \mathcal{H}_1 . We assume that the pair is *compatible*, i.e., both \mathcal{H}_i are embedded in some common Hausdorff topological vector space \mathcal{M} .

Y. Ameur (✉)

Department of Mathematics, Faculty of Science, Lund University, Lund, Sweden

e-mail: Yacin.Ameur@math.lu.se; yacin@maths.lth.se

We define the K -functional¹ for the couple $\overline{\mathcal{H}}$ by

$$K(t, x) = K(t, x; \overline{\mathcal{H}}) = \inf_{x=x_0+x_1} \{ \|x_0\|_0^2 + t \|x_1\|_1^2 \}, \quad t > 0, x \in \mathcal{M}.$$

The *sum* of the spaces \mathcal{H}_0 and \mathcal{H}_1 is defined to be the space consisting of all $x \in \mathcal{M}$ such that the quantity $\|x\|_\Sigma^2 := K(1, x)$ is finite; we denote this space by the symbols

$$\Sigma = \Sigma(\overline{\mathcal{H}}) = \mathcal{H}_0 + \mathcal{H}_1.$$

We shall soon see that Σ is a Hilbert space (see Lemma 1.1). The *intersection*

$$\Delta = \Delta(\overline{\mathcal{H}}) = \mathcal{H}_0 \cap \mathcal{H}_1$$

is a Hilbert space under the norm $\|x\|_\Delta^2 := \|x\|_0^2 + \|x\|_1^2$.

A map $T : \Sigma(\overline{\mathcal{H}}) \rightarrow \Sigma(\overline{\mathcal{K}})$ is called a *couple map* from $\overline{\mathcal{H}}$ to $\overline{\mathcal{K}}$ if the restriction of T to \mathcal{H}_i maps \mathcal{H}_i boundedly into \mathcal{K}_i for $i = 0, 1$. We use the notations $T \in \mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{K}})$ or $T : \overline{\mathcal{H}} \rightarrow \overline{\mathcal{K}}$ to denote that T is a couple map. It is easy to check that $\mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{K}})$ as a Banach space, when equipped with the norm

$$\|T\|_{\mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{K}})} := \max_{j=0,1} \{ \|T\|_{\mathcal{L}(\mathcal{H}_j; \mathcal{K}_j)} \}. \quad (1.1)$$

If $\|T\|_{\mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{K}})} \leq 1$ we speak of a *contraction* from $\overline{\mathcal{H}}$ to $\overline{\mathcal{K}}$.

A Banach space X such that $\Delta \subset X \subset \Sigma$ (continuous inclusions) is called *intermediate* with respect to the pair $\overline{\mathcal{H}}$.

Let X, Y be intermediate spaces with respect to couples $\overline{\mathcal{H}}, \overline{\mathcal{K}}$, respectively. We say that X, Y are (relative) *interpolation spaces* if there is a constant C such that $T : \overline{\mathcal{H}} \rightarrow \overline{\mathcal{K}}$ implies that $T : X \rightarrow Y$ and

$$\|T\|_{\mathcal{L}(X; Y)} \leq C \|T\|_{\mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{K}})}. \quad (1.2)$$

In the case when $C = 1$ we speak about *exact interpolation*. When $\overline{\mathcal{H}} = \overline{\mathcal{K}}$ and $X = Y$ we simply say that X is an (exact) interpolation space with respect to $\overline{\mathcal{H}}$.

Let H be a suitable function of two positive variables and X, Y spaces intermediate to the couples $\overline{\mathcal{H}}, \overline{\mathcal{K}}$, respectively. We say that the spaces X, Y are of *type H* (relative to $\overline{\mathcal{H}}, \overline{\mathcal{K}}$) if for any positive numbers M_0, M_1 we have

$$\|T\|_{\mathcal{L}(\mathcal{H}_i; \mathcal{K}_i)} \leq M_i, \quad i = 0, 1 \quad \text{implies} \quad \|T\|_{\mathcal{L}(X; Y)} \leq H(M_0, M_1). \quad (1.3)$$

¹More precisely, this is the *quadratic version* of the classical Peetre K -functional.

The case $H(x, y) = \max\{x, y\}$ corresponds to exact interpolation, while $H(x, y) = x^{1-\theta}y^\theta$ corresponds to the convexity estimate

$$\|T\|_{\mathcal{L}(X;Y)} \leq \|T\|_{\mathcal{L}(\mathcal{H}_0;\mathcal{K}_0)}^{1-\theta} \|T\|_{\mathcal{L}(\mathcal{H}_1;\mathcal{K}_1)}^\theta. \tag{1.4}$$

In the situation of (1.4), one says that the interpolation spaces X, Y are of *exponent* θ with respect to the pairs $\overline{\mathcal{H}}, \overline{\mathcal{K}}$.

1.2 *K*-Spaces

Given a regular Hilbert couple $\overline{\mathcal{H}}$ and a positive Radon measure ϱ on the compactified half-line $[0, \infty]$ we define an intermediate quadratic norm by

$$\|x\|_*^2 = \|x\|_\varrho^2 = \int_{[0,\infty]} (1+t^{-1}) K(t, x; \overline{\mathcal{H}}) d\varrho(t). \tag{1.5}$$

Here the integrand $k(t) = (1+t^{-1}) K(t, x)$ is defined at the points 0 and ∞ by $k(0) = \|x\|_1^2$ and $k(\infty) = \|x\|_0^2$; we shall write \mathcal{H}_* or \mathcal{H}_ϱ for the Hilbert space defined by the norm (1.5).

Let $T \in \mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{K}})$ and suppose that $\|T\|_{\mathcal{L}(\mathcal{H}_i;\mathcal{K}_i)} \leq M_i$, then

$$K(t, Tx; \overline{\mathcal{K}}) \leq M_0^2 K(M_1^2 t / M_0^2, x; \overline{\mathcal{H}}), \quad x \in \Sigma. \tag{1.6}$$

In particular, $M_i \leq 1$ for $i = 0, 1$ implies $\|Tx\|_{\mathcal{K}_\varrho} \leq \|x\|_{\mathcal{H}_\varrho}$ for all $x \in \mathcal{H}_\varrho$. It follows that the spaces $\mathcal{H}_\varrho, \mathcal{K}_\varrho$ are exact interpolation spaces with respect to $\overline{\mathcal{H}}, \overline{\mathcal{K}}$.

1.2.1 Geometric Interpolation

When the measure ϱ is given by

$$d\varrho(t) = c_\theta \frac{t^{-\theta}}{1+t} dt, \quad c_\theta = \frac{\pi}{\sin \theta \pi}, \quad 0 < \theta < 1,$$

we denote the norm (1.5) by

$$\|x\|_\theta^2 := c_\theta \int_0^\infty t^{-\theta} K(t, x) \frac{dt}{t}. \tag{1.7}$$

The corresponding space \mathcal{H}_θ is easily seen to be of exponent θ with respect to $\overline{\mathcal{H}}$. In Sect. 3.1, we will recognize \mathcal{H}_θ as the geometric interpolation space which has been studied independently by several authors, see [25, 27, 40].

1.3 Pick Functions

Let $\overline{\mathcal{H}}$ be a regular Hilbert couple. The squared norm $\|x\|_1^2$ is a densely defined quadratic form in \mathcal{H}_0 , which we represent as

$$\|x\|_1^2 = \langle Ax, x \rangle_0 = \|A^{1/2}x\|_0^2$$

where A is a densely defined, positive, injective (perhaps unbounded) operator in \mathcal{H}_0 . The domain of the positive square-root $A^{1/2}$ is Δ .

Lemma 1.1 *We have in terms of the functional calculus in \mathcal{H}_0*

$$K(t, x) = \left\langle \frac{tA}{1+tA}x, x \right\rangle_0, \quad t > 0. \quad (1.8)$$

In the formula (1.8), we have identified the bounded operator $\frac{tA}{1+tA}$ with its extension to \mathcal{H}_0 .

Proof Fix $x \in \Delta$. By a straightforward convexity argument, there is a unique decomposition $x = x_{0,t} + x_{1,t}$ which is *optimal* in the sense that

$$K(t, x) = \|x_{0,t}\|_0^2 + t \|x_{1,t}\|_1^2. \quad (1.9)$$

It follows that $x_{i,t} \in \Delta$ for $i = 0, 1$. Moreover, for all $y \in \Delta$ we have

$$\frac{d}{d\epsilon} \{ \|x_{0,t} + \epsilon y\|_0^2 + t \|x_{1,t} - \epsilon y\|_1^2 \}_{\epsilon=0} = 0,$$

i.e.,

$$\langle A^{-1/2}x_{0,t} - tA^{1/2}x_{1,t}, A^{1/2}y \rangle_0 = 0, \quad y \in \Delta.$$

By regularity, we conclude that $A^{-1/2}x_{0,t} = tA^{1/2}x_{1,t}$, whence

$$x_{0,t} = \frac{tA}{1+tA}x \quad \text{and} \quad x_{1,t} = \frac{1}{1+tA}x. \quad (1.10)$$

(Note that the operators in (1.10) extend to bounded operators on \mathcal{H}_0 .) Inserting the relations (1.10) into (1.9), one finishes the proof of the lemma. \square

Now fix a positive Radon measure ϱ on $[0, \infty]$. The norm in the space \mathcal{H}_ϱ (see (1.5)) can be written

$$\|x\|_\varrho^2 = \langle h(A)x, x \rangle_0, \quad (1.11)$$

where

$$h(\lambda) = \int_{[0, \infty]} \frac{(1+t)\lambda}{1+t\lambda} d\varrho(t). \tag{1.12}$$

The class of functions representable in this form for some positive Radon measure ϱ is the class P' of *Pick functions, positive and regular on \mathbf{R}_+* .

Notice that for the definition (1.11) to make sense, we just need h to be defined on $\sigma(A) \setminus \{0\}$, where $\sigma(A)$ is the spectrum of A . (The value $h(0)$ is irrelevant since A is injective).

A calculus exercise shows that for the space \mathcal{H}_θ (see (1.7)) we have

$$\|x\|_\theta^2 = \langle A^\theta x, x \rangle_0. \tag{1.13}$$

1.4 Quadratic Interpolation Norms

Let \mathcal{H}_* be any *quadratic* intermediate space relative to $\overline{\mathcal{H}}$. We write

$$\|x\|_*^2 = \langle Bx, x \rangle_0$$

where B is a positive injective operator in \mathcal{H}_0 (the domain of $B^{1/2}$ is Δ).

For a map $T \in \mathcal{L}(\overline{\mathcal{H}})$ we shall often use the simplified notations

$$\|T\| = \|T\|_{\mathcal{L}(\mathcal{H}_0)} \quad , \quad \|T\|_A = \|T\|_{\mathcal{L}(\mathcal{H}_1)} \quad , \quad \|T\|_B = \|T\|_{\mathcal{L}(\mathcal{H}_*)}.$$

The reader can check the identities

$$\|T\|_A = \|A^{1/2}TA^{-1/2}\| \quad \text{and} \quad \|T\|_B = \|B^{1/2}TB^{-1/2}\|.$$

We shall refer to the following lemma as *Donoghue's lemma*, cf. [13, Lemma 1].

Lemma 1.2 *If \mathcal{H}_* is exact interpolation with respect to $\overline{\mathcal{H}}$, then B commutes with every projection which commutes with A and $B = h(A)$ where h is some positive Borel function on $\sigma(A)$.*

Proof For an orthogonal projection E on \mathcal{H}_0 , the condition $\|E\|_A \leq 1$ is equivalent to that $EAE \leq A$, i.e., that E commutes with A . The hypothesis that \mathcal{H}_* be exact interpolation thus implies that every spectral projection of A commutes with B . It now follows from von Neumann's bicommutator theorem that $B = h(A)$ for some positive Borel function h on $\sigma(A)$. □

In view of the lemma, the characterization of the exact quadratic interpolation norms of a given type H reduces to the characterization of functions $h : \sigma(A) \rightarrow \mathbf{R}_+$ such that for all $T \in \mathcal{L}(\overline{\mathcal{H}})$

$$\|T\| \leq M_0 \quad \text{and} \quad \|T\|_A \leq M_1 \quad \Rightarrow \quad \|T\|_{h(A)} \leq H(M_0, M_1), \tag{1.14}$$

or alternatively,

$$T^*T \leq M_0^2 \quad \text{and} \quad T^*AT \leq M_1^2 A \quad \Rightarrow \quad T^*h(A)T \leq H(M_0, M_1)^2 h(A). \quad (1.15)$$

The set of functions $h : \sigma(A) \rightarrow \mathbf{R}_+$ satisfying these equivalent conditions forms a convex cone $C_{H,A}$; its elements are called *interpolation functions of type H relative to A* . In the case when $H(x, y) = \max\{x, y\}$ we simply write C_A for $C_{H,A}$ and speak of *exact interpolation functions relative to A* .

1.5 Exact Calderón Pairs and the K -Property

Given two intermediate normed spaces Y, X relative to $\overline{\mathcal{H}}, \overline{\mathcal{K}}$, we say that they are (relatively) *exact K -monotonic* if the conditions

$$x^0 \in X \quad \text{and} \quad K(t, y^0; \overline{\mathcal{H}}) \leq K(t, x^0; \overline{\mathcal{K}}), \quad t > 0$$

imply that

$$y^0 \in Y \quad \text{and} \quad \|y^0\|_Y \leq \|x^0\|_X.$$

It is easy to see that *exact K -monotonicity implies exact interpolation*.

Proof of this. If $\|T\|_{\mathcal{L}(\overline{\mathcal{K}}; \overline{\mathcal{H}})} \leq 1$, then $\forall x, t: K(t, Tx; \overline{\mathcal{H}}) \leq K(t, x; \overline{\mathcal{K}})$ whence $\|Tx\|_Y \leq \|x\|_X$, by exact K -monotonicity. Hence $\|T\|_{\mathcal{L}(X; Y)} \leq 1$. \square

Two pairs $\overline{\mathcal{H}}, \overline{\mathcal{K}}$ are called *exact relative Calderón pairs* if any two exact interpolation (Banach-) spaces Y, X are exact K -monotonic. Thus, with respect to exact Calderón pairs, exact interpolation is equivalent to exact K -monotonicity. The term ‘‘Calderón pair’’ was coined after thorough investigation of Calderón’s study of the pair (L_1, L_∞) , see [10, 11].

In our present discussion, it is not convenient to work directly with the definition of exact Calderón pairs. Instead, we shall use the following, closely related notion.

We say that a pair of couples $\overline{\mathcal{H}}, \overline{\mathcal{K}}$ has the *relative (exact) K -property* if for all $x^0 \in \Sigma(\overline{\mathcal{K}})$ and $y^0 \in \Sigma(\overline{\mathcal{H}})$ such that

$$K(t, y^0; \overline{\mathcal{H}}) \leq K(t, x^0; \overline{\mathcal{K}}), \quad t > 0, \quad (1.16)$$

there exists a map $T \in \mathcal{L}(\overline{\mathcal{K}}; \overline{\mathcal{H}})$ such that $Tx^0 = y^0$ and $\|T\|_{\mathcal{L}(\overline{\mathcal{K}}; \overline{\mathcal{H}})} \leq 1$.

Lemma 1.3 *If $\overline{\mathcal{H}}, \overline{\mathcal{K}}$ have the relative K -property, then they are exact relative Calderón pairs.*

Proof Let Y, X be exact interpolation spaces relative to $\overline{\mathcal{H}}, \overline{\mathcal{K}}$ and take $x^0 \in X$ and $y^0 \in \Sigma(\overline{\mathcal{H}})$ such that (1.16) holds. By the K -property there is $T : \overline{\mathcal{K}} \rightarrow \overline{\mathcal{H}}$ such that $Tx^0 = y^0$ and $\|T\| \leq 1$. Then $\|T\|_{\mathcal{L}(X;Y)} \leq 1$, and so $\|y^0\|_Y = \|Tx^0\|_Y \leq \|x^0\|_X$. We have shown that Y, X are exact K -monotonic. \square

In the diagonal case $\overline{\mathcal{H}} = \overline{\mathcal{K}}$, we simply say that $\overline{\mathcal{H}}$ is an *exact Calderón couple* if for intermediate spaces Y, X , the property of being exact interpolation is equivalent to being exact K -monotonic. Likewise, we say that $\overline{\mathcal{H}}$ has the *K -property* if the pair of couples $\overline{\mathcal{H}}, \overline{\mathcal{H}}$ has that property.

Remark 1.4 For an operator $T : \overline{\mathcal{K}} \rightarrow \overline{\mathcal{H}}$ to be a contraction, it is necessary and sufficient that

$$K(t, Tx; \overline{\mathcal{H}}) \leq K(t, x; \overline{\mathcal{K}}), \quad x \in \Sigma(\overline{\mathcal{K}}), t > 0. \tag{1.17}$$

Indeed, the necessity is immediate. To prove the sufficiency it suffices to observe that letting $t \rightarrow \infty$ in (1.17) gives $\|Tx\|_0 \leq \|x\|_0$, and dividing (1.17) by t , and then letting $t \rightarrow 0$, gives that $\|Tx\|_1 \leq \|x\|_1$.

2 Mapping Properties of Hilbert Couples

2.1 Main Results

We shall elaborate on the following main result from [2].

Theorem I *Any pair of regular Hilbert couples $\overline{\mathcal{H}}, \overline{\mathcal{K}}$ has the relative K -property.*

Before we come to the proof of Theorem I, we note some consequences of it. We first have the following corollary, which shows that a strong form of the K -property is true.

Corollary 2.1 *Let $\overline{\mathcal{H}}$ be a regular Hilbert couple and $x^0, y^0 \in \Sigma$ elements such that*

$$K(t, y^0) \leq M_0^2 K(M_1^2 t / M_0^2, x^0), \quad t > 0. \tag{2.1}$$

Then

- (i) *There exists a map $T \in \mathcal{L}(\overline{\mathcal{H}})$ such that $Tx^0 = y^0$ and $\|T\|_{\mathcal{L}(\mathcal{H}_i)} \leq M_i$, $i = 0, 1$.*
- (ii) *If $x^0 \in X$ where X is an interpolation space of type H , then*

$$\|y^0\|_X \leq H(M_0, M_1) \|x^0\|_X.$$

Proof (i) Introduce a new couple $\overline{\mathcal{K}}$ by letting $\|x\|_{\mathcal{K}_i} = M_i \|x\|_{\mathcal{H}_i}$. The relation (2.1) then says that

$$K(t, y^0; \overline{\mathcal{H}}) \leq K(t, x^0; \overline{\mathcal{K}}), \quad t > 0.$$

By Theorem I there is a contraction $T : \overline{\mathcal{K}} \rightarrow \overline{\mathcal{H}}$ such that $Tx^0 = y^0$. It now suffices to note that $\|T\|_{\mathcal{L}(\mathcal{H}_i)} = M_i \|T\|_{\mathcal{L}(\mathcal{K}_i; \mathcal{H}_i)}$; (ii) then follows from Lemma 1.3. \square

We next mention some equivalent versions of Theorem I, which uses the families of functionals K_p and E_p defined (for $p \geq 1$ and $t, s > 0$) via

$$\begin{aligned} K_p(t) &= K_p(t, x) = K_p(t, x; \overline{\mathcal{H}}) = \inf_{x=x_0+x_1} \{ \|x_0\|_0^p + t \|x_1\|_1^p \} \\ E_p(s) &= E_p(s, x) = E_p(s, x; \overline{\mathcal{H}}) = \inf_{\|x_0\|_0^p \leq s} \{ \|x - x_0\|_1^p \}. \end{aligned} \quad (2.2)$$

Note that $K = K_2$ and that $E_p(s) = E_1(s^{1/p})^p$; the E -functionals are used in approximation theory. One has that E_p is decreasing and convex on \mathbf{R}_+ and that

$$K_p(t) = \inf_{s>0} \{ s + tE_p(s) \},$$

which means that K_p is a kind of *Legendre transform* of E_p . The inverse Legendre transformation takes the form

$$E_p(s) = \sup_{t>0} \left\{ \frac{K_p(t)}{t} - \frac{s}{t} \right\}.$$

It is now immediate that, for all $x \in \Sigma(\overline{\mathcal{K}})$ and $y \in \Sigma(\overline{\mathcal{H}})$, we have

$$K_p(t, y) \leq K_p(t, x), \quad t > 0 \quad \Leftrightarrow \quad E_p(s, y) \leq E_p(s, x), \quad s > 0. \quad (2.3)$$

Since moreover $E_p(s) = E_2(s^{2/p})^{p/2}$, the conditions in (2.3) are equivalent to that $K(t, y) \leq K(t, x)$ for all $t > 0$. We have shown the following result.

Corollary 2.2 *In Theorem I, one can substitute the K -functional for any of the functionals K_p or E_p .*

Define an exact interpolation norm $\|\cdot\|_{\varrho, p}$ relative to $\overline{\mathcal{H}}$ by

$$\|x\|_{\varrho, p}^p = \int_{[0, \infty]} (1 + t^{-1}) K_p(t, x) d\varrho(t)$$

where ϱ is a positive Radon measure on $[0, \infty]$. This norm is non-quadratic when $p \neq 2$, but is of course equivalent to the quadratic norm corresponding to $p = 2$.

2.2 Reduction to the Diagonal Case

It is not hard to reduce the discussion of Theorem I to a diagonal situation.

Lemma 2.3 *If the K -property holds for regular Hilbert couples in the diagonal case $\overline{\mathcal{H}} = \overline{\mathcal{K}}$, then it holds in general.*

Proof Fix elements $y^0 \in \Sigma(\overline{\mathcal{H}})$ and $x^0 \in \Sigma(\overline{\mathcal{K}})$ such that the inequality (1.16) holds. We must construct a map $T : \overline{\mathcal{K}} \rightarrow \overline{\mathcal{H}}$ such that $Tx^0 = y^0$ and $\|T\| \leq 1$.

To do this, we form the direct sum $\overline{\mathcal{S}} = (\mathcal{H}_0 \oplus \mathcal{K}_0, \mathcal{H}_1 \oplus \mathcal{K}_1)$. It is clear that $\mathcal{S}_0 + \mathcal{S}_1 = (\mathcal{H}_0 + \mathcal{H}_1) \oplus (\mathcal{K}_0 + \mathcal{K}_1)$, and that

$$K(t, x \oplus y; \overline{\mathcal{S}}) = K(t, x; \overline{\mathcal{H}}) + K(t, y; \overline{\mathcal{K}}).$$

Then

$$K(t, 0 \oplus y^0; \overline{\mathcal{S}}) \leq K(t, x^0 \oplus 0; \overline{\mathcal{S}}).$$

Hence assuming that the couple $\overline{\mathcal{S}}$ has the K -property, we can assert the existence of a map $S \in \mathcal{L}(\overline{\mathcal{S}})$ such that $S(x^0 \oplus 0) = 0 \oplus y^0$ and $\|S\| \leq 1$. Letting $P : \mathcal{S}_0 + \mathcal{S}_1 \rightarrow \mathcal{K}_0 + \mathcal{K}_1$ be the orthogonal projection, the assignment $Tx = PS(x \oplus 0)$ now defines a map such that $Tx^0 = y^0$ and $\|T\|_{\mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{K}})} \leq 1$. \square

2.3 The Principal Case

The core content of Theorem I is contained in the following statement.

Theorem 2.4 *Suppose that a regular Hilbert couple $\overline{\mathcal{H}}$ is finite dimensional and that all eigenvalues of the corresponding operator A are of unit multiplicity. Then $\overline{\mathcal{H}}$ has the K -property.*

We shall settle for proving Lemma 2.4 in this section, postponing to Sect. 5 the general case of Theorem I.

To prepare for the proof, we write the eigenvalues λ_i of A in the increasing order,

$$\sigma(A) = \{\lambda_i\}_1^n \quad \text{where} \quad 0 < \lambda_1 < \cdots < \lambda_n.$$

Let e_i be corresponding eigenvectors of unit length for the norm of \mathcal{H}_0 . Then for a vector $x = \sum x_i e_i$ we have

$$\|x\|_0^2 = \sum_1^n |x_i|^2, \quad \|x\|_1^2 = \sum_1^n \lambda_i |x_i|^2.$$

Working in the coordinate system (e_i) , the couple $\overline{\mathcal{H}}$ becomes identified with the

n -dimensional weighted ℓ_2 couple

$$\overline{\ell}_2^n(\lambda) := (\ell_2^n, \ell_2^n(\lambda)),$$

where we write λ for the sequence $(\lambda_i)_1^n$.

We will henceforth identify a vector $x = \sum x_i e_i$ with the point $x = (x_i)_1^n$ in \mathbf{C}^n ; accordingly, the space $\mathcal{L}(\ell_2^n)$ is identified with the C^* -algebra $M_n(\mathbf{C})$ of complex $n \times n$ matrices.

It will be convenient to reparametrize the K -functional for the couple $\overline{\ell}_2^n(\lambda)$ and write

$$k_\lambda(t, x) := K\left(1/t, x; \overline{\ell}_2^n(\lambda)\right). \quad (2.4)$$

By Lemma 1.1 we have

$$k_\lambda(t, x) = \sum_{i=1}^n \frac{\lambda_i}{t + \lambda_i} |x_i|^2, \quad x \in \mathbf{C}^n. \quad (2.5)$$

2.4 Basic Reductions

To prove that the couple $\overline{\ell}_2^n(\lambda)$ has the K -property, we introduce an auxiliary parameter $\rho > 1$. The exact value of ρ will change meaning during the course of the argument, the main point being that it can be chosen arbitrarily close to 1.

Initially, we pick any $\rho > 1$ such that $\rho\lambda_i < \lambda_{i+1}$ for all i ; we assume also that we are given two elements $x^0, y^0 \in \mathbf{C}^n$ such that

$$k_\lambda(t, y^0) < \frac{1}{\rho} k_\lambda(t, x^0), \quad t \geq 0. \quad (2.6)$$

We must construct a matrix $T \in M_n(\mathbf{C})$ such that

$$Tx^0 = y^0 \quad \text{and} \quad k_\lambda(t, Tx) \leq k_\lambda(t, x), \quad x \in \mathbf{C}^n, t > 0. \quad (2.7)$$

Define $\tilde{x}^0 = (|x_i^0|)_1^n$ and $\tilde{y}^0 = (|y_i^0|)_1^n$ and suppose that

$$k_\lambda(t, \tilde{y}^0) < \frac{1}{\rho} k_\lambda(t, \tilde{x}^0), \quad t \geq 0.$$

Suppose that we can find an operator $T_0 \in M_n(\mathbf{C})$ such that $T_0\tilde{x}^0 = \tilde{y}^0$ and $k_\lambda(t, T_0x) < k_\lambda(t, x)$ for all $x \in \mathbf{C}^n$ and $t > 0$. Writing $x_k^0 = e^{i\theta_k}\tilde{x}_k^0$ and $y_k^0 = e^{i\varphi_k}\tilde{y}_k^0$ where $\theta_k, \varphi_k \in \mathbf{R}$, we then have $Tx^0 = y^0$ and $k_\lambda(t, Tx) < k_\lambda(t, x)$ where

$$T = \text{diag}(e^{i\varphi_k})T_0\text{diag}(e^{-i\theta_k}).$$

Replacing x^0, y^0 by \tilde{x}^0, \tilde{y}^0 we can thus assume that the coordinates x_i^0 and y_i^0 are non-negative; replacing them by small perturbations if necessary, we can assume that they are strictly positive, at the expense of slightly diminishing the number ρ .

Now put $\beta_i = \lambda_i$ and $\alpha_i = \rho\lambda_i$. Our assumption on ρ means that

$$0 < \beta_1 < \alpha_1 < \cdots < \beta_n < \alpha_n.$$

Using the explicit expression for the K -functional, it is plain to check that

$$k_\beta(t, x) \leq k_\alpha(t, x) \leq \rho k_\beta(t, x), \quad x \in \mathbf{C}^n, t \geq 0.$$

Our assumption (2.6) therefore implies that

$$k_\alpha(t, y^0) < k_\beta(t, x^0), \quad t \geq 0. \tag{2.8}$$

We shall verify the existence of a matrix $T = T_\rho = T_{\rho, x^0, y^0}$ such that

$$Tx^0 = y^0 \quad \text{and} \quad k_\alpha(t, Tx) \leq k_\beta(t, x), \quad x \in \mathbf{C}^n, t > 0. \tag{2.9}$$

It is clear by compactness that, as $\rho \downarrow 1$, the corresponding matrices T_ρ will cluster at some point T satisfying $Tx^0 = y^0$ and $\|T\|_{\mathcal{L}(\overline{\mathcal{H}})} \leq 1$. (See Remark 1.4.)

In conclusion, the proof of Theorem 2.4 will be complete when we can construct a matrix T satisfying (2.9) with ρ arbitrarily close to 1.

2.5 Construction of T

Let \mathcal{P}_k denote the linear space of complex polynomials of degree at most k . We shall use the polynomials

$$L_\alpha(t) = \prod_1^n (t + \alpha_i) \quad , \quad L_\beta(t) = \prod_1^n (t + \beta_i),$$

and the product $L = L_\alpha L_\beta$. Notice that

$$L'(-\alpha_i) < 0 \quad , \quad L'(-\beta_i) > 0. \tag{2.10}$$

Recalling the formula (2.5), it is clear that we can define a real polynomial $P \in \mathcal{P}_{2n-1}$ by

$$\frac{P(t)}{L(t)} = k_\beta(t, x^0) - k_\alpha(t, y^0). \tag{2.11}$$

Clearly $P(t) > 0$ when $t \geq 0$. Moreover, a consideration of the residues at the poles

of the right-hand member shows that P is uniquely defined by the values

$$P(-\beta_i) = (x_i^0)^2 \beta_i L'(-\beta_i) \quad , \quad P(-\alpha_i) = -(y_i^0)^2 \alpha_i L'(-\alpha_i). \quad (2.12)$$

Combining with (2.10), we conclude that

$$P(-\alpha_i) > 0 \quad \text{and} \quad P(-\beta_i) > 0. \quad (2.13)$$

Perturbing the problem slightly, it is clear that we can assume that P has exact degree $2n - 1$, and that all zeros of P have multiplicity 1. (We here diminish the value of $\rho > 1$ somewhat, if necessary.)

Now, P has $2n - 1$ simple zeros, which we split according to

$$P^{-1}(\{0\}) = \{-r_i\}_{i=1}^{2m-1} \cup \{-c_i, -\bar{c}_i\}_{i=1}^{n-m},$$

where the r_i are positive and the c_i are non-real, and chosen to have positive imaginary parts. The following is the key observation.

Lemma 2.5 *We have that*

$$L'(-\beta_i) P(-\beta_i) > 0 \quad , \quad L'(-\alpha_i) P(-\alpha_i) < 0 \quad (2.14)$$

and there is a splitting $\{r_i\}_{i=1}^{2m-1} = \{\delta_i\}_{i=1}^m \cup \{\gamma_i\}_{i=1}^{m-1}$ such that

$$L(-\delta_j) P'(-\delta_j) > 0 \quad , \quad L(-\gamma_k) P'(-\gamma_k) < 0. \quad (2.15)$$

Proof The inequalities (2.14) follow immediately from (2.13) and (2.10). It remains to prove (2.15).

Let $-h$ denote the leftmost real zero of the polynomial LP (of degree $4n - 1$). We claim that $P(-h) = 0$. If this were not the case, we would have $h = \alpha_n$. Since the degree of P is odd, $P(-t)$ is negative for large values of t , and so $P(-\alpha_n) < 0$ contradicting (2.13). We have shown that $P(-h) = 0$. Since all zeros of LP have multiplicity 1, we have $(LP)'(-h) \neq 0$, whence

$$L(-h)P'(-h) = (LP)'(-h) > 0.$$

We write $\delta_m = h$ and put $P_*(t) = P(t)/(t + \delta_m)$. Since $t + \delta_m > 0$ for $t \in \{-\alpha_i, -\beta_i\}_1^n$, we have by (2.13) that for all i

$$P_*(-\alpha_i) > 0 \quad \text{and} \quad P_*(-\beta_i) > 0.$$

Denote by $\{-r_j^*\}_{j=1}^{2m-2}$ the real zeros of P_* . Since the degree of LP_* is even and the polynomial $(LP_*)'$ has alternating signs in the set $\{-\alpha_i, -\beta_i\}_{i=1}^n \cup \{-r_i^*\}_{i=1}^{2m-2}$, we can split the zeros of P_* as $\{-\delta_i, -\gamma_i\}_{i=1}^{m-1}$, where

$$L(-\delta_i)P_*'(-\delta_i) > 0 \quad , \quad L(-\gamma_i)P_*'(-\gamma_i) < 0. \quad (2.16)$$

Since $P'(-r_j^*) = (\delta_m - r_j^*)P'_*(-r_j^*)$ and $\delta_m > r_j^*$, the signs of $P'(-r_j^*)$ and $P'_*(-r_j^*)$ are equal, proving (2.15). \square

Recall that $\{-c_i\}_1^{n-m}$ denote the zeros of P such that $\text{Im } c_i > 0$. We put (with the convention that an empty product equals 1)

$$L_\delta(t) = \prod_{i=1}^m (t + \delta_i) \quad , \quad L_\gamma(t) = \prod_{i=1}^{m-1} (t + \gamma_i) \quad , \quad L_c(t) = \prod_{i=1}^{n-m} (t + c_i).$$

We define a linear map $F : \mathbf{C}^{n+m} \rightarrow \mathbf{C}^{n+m-1}$ in the following way. First define a subspace $U \subset \mathcal{P}_{2n-1}$ by

$$U = \{ L_c q ; q \in \mathcal{P}_{n+m-1} \}.$$

Notice that U has dimension $n+m-1$ and that $P \in U$; in fact $P = a L_c L_c^* L_\delta L_\gamma$ where a is the leading coefficient and the $*$ -operation is defined by $L^*(z) = \overline{L(\bar{z})}$.

For a polynomial $Q \in U$ we have

$$\begin{aligned} \frac{|Q(t)|^2}{L(t)P(t)} &= \sum_{i=1}^n |x_i|^2 \frac{\beta_i}{t + \beta_i} + \sum_{i=1}^n |x'_i|^2 \frac{\delta_i}{t + \delta_i} \\ &\quad - \sum_{i=1}^n |y_i|^2 \frac{\alpha_i}{t + \alpha_i} - \sum_{i=1}^{m-1} |y'_i|^2 \frac{\gamma_i}{t + \gamma_i}, \end{aligned} \tag{2.17}$$

where, for definiteness,

$$x_i = \frac{Q(-\beta_i)}{\sqrt{\beta_i L'(-\beta_i) P(-\beta_i)}} \quad ; \quad x'_j = \frac{Q(-\delta_j)}{\sqrt{\delta_j L'(-\delta_j) P(-\delta_j)}} \tag{2.18}$$

$$y_i = \frac{Q(-\alpha_i)}{\sqrt{-\alpha_i L'(-\alpha_i) P(-\alpha_i)}} \quad ; \quad y'_j = \frac{Q(-\gamma_j)}{\sqrt{-\gamma_j L'(-\gamma_j) P(-\gamma_j)}}. \tag{2.19}$$

The identities in (2.18) give rise to a linear map

$$M : \mathbf{C}^n \oplus \mathbf{C}^m \rightarrow U \quad ; \quad [x; x'] \mapsto Q. \tag{2.20}$$

We can similarly regard (2.19) as a linear map

$$N : U \rightarrow \mathbf{C}^n \oplus \mathbf{C}^{m-1} \quad ; \quad Q \mapsto [y; y']. \tag{2.21}$$

Our desired map F is defined as the composite

$$F = NM : \mathbf{C}^n \oplus \mathbf{C}^m \rightarrow \mathbf{C}^n \oplus \mathbf{C}^{m-1} \quad ; \quad [x; x'] \mapsto [y; y'].$$

Notice that if $Q = M [x; x']$ and $[y; y'] = F [x; x']$ then (2.17) means that

$$k_{\beta \oplus \delta} (t, [x; x']) - k_{\alpha \oplus \gamma} (t, F [x; x']) = \frac{|Q(t)|^2}{L(t)P(t)} \geq 0, \quad t \geq 0.$$

This implies that F is a contraction from $\overline{\ell_2^{n+m}}(\beta \oplus \delta)$ to $\overline{\ell_2^{n+m-1}}(\alpha \oplus \gamma)$.

We now define T as a ‘‘compression’’ of F . Namely, let $E : \mathbf{C}^n \oplus \mathbf{C}^{m-1} \rightarrow \mathbf{C}^n$ be the projection onto the first n coordinates, and define an operator T on \mathbf{C}^n by

$$Tx = EF [x; 0], \quad x \in \mathbf{C}^n.$$

Taking $Q = P$ in (2.17) we see that $Tx^0 = y^0$. Moreover,

$$\begin{aligned} k_{\beta} (t, x) - k_{\alpha} (t, Tx) &= \sum_{i=1}^n |x_i|^2 \frac{\beta_i}{t + \beta_i} - \sum_{i=1}^n |y_i|^2 \frac{\alpha_i}{t + \alpha_i} \\ &\geq \sum_{i=1}^n |x_i|^2 \frac{\beta_i}{t + \beta_i} - \sum_{i=1}^n |y_i|^2 \frac{\alpha_i}{t + \alpha_i} - \sum_{j=1}^{m-1} |y'_j|^2 \frac{\gamma_j}{t + \gamma_j} \\ &= k_{\beta \oplus \delta} (t, [x; 0]) - k_{\alpha \oplus \gamma} (t, F [x; 0]) = \frac{|Q(t)|^2}{L(t)P(t)}. \end{aligned}$$

Since the right-hand side is non-negative, we have shown that

$$k_{\alpha} (t, Tx) \leq k_{\beta} (t, x), \quad t > 0, x \in \mathbf{C}^n,$$

as desired. The proof of Theorem 2.4 is finished. \square

2.6 Real Scalars

Theorem 2.4 holds also in the case of Euclidean spaces over the real scalar field. To see this, assume without loss of generality that the vectors $x^0, y^0 \in \mathbf{C}^n$ have *real entries* (still satisfying $k_{\lambda} (t, y^0) \leq k_{\lambda} (t, x^0)$ for all $t > 0$).

By Theorem 2.4 we can find a (complex) contraction T of $\overline{\ell_2^n}(\lambda)$ such that $Tx^0 = y^0$. It is clear that the operator T^* defined by $T^*x = \overline{T(\bar{x})}$ satisfies those same conditions. Replacing T by $\frac{1}{2}(T + T^*)$ we obtain a real matrix $T \in M_n(\mathbf{R})$, which is a contraction of $\overline{\ell_2^n}(\lambda)$ and maps x^0 to y^0 . \square

2.7 Explicit Representations

We here deduce an explicit representation for the operator T constructed above.

Let x^0 and y^0 be two non-negative vectors such that

$$k_\lambda(t, y^0) \leq k_\lambda(t, x^0), \quad t > 0.$$

For small $\rho > 0$ we perturb x^0, y^0 slightly to vectors \tilde{x}^0, \tilde{y}^0 which satisfy the conditions imposed on the previous subsections. We can then construct a matrix $T = T_\rho$ such that

$$T\tilde{x}^0 = \tilde{y}^0 \quad \text{and} \quad k_\alpha(t, Tx) \leq k_\beta(t, x), \quad t > 0, x \in \mathbf{C}^n, \quad (2.22)$$

where $\beta = \lambda$ and $\alpha = \rho\lambda$. As $\rho, \tilde{x}^0, \tilde{y}^0$ approaches 1, x^0 , respectively y^0 , it is clear that any cluster point T of the set of contractions T_ρ will satisfy

$$Tx^0 = y^0 \quad \text{and} \quad k_\lambda(t, Tx) \leq k_\lambda(t, x), \quad t > 0, x \in \mathbf{C}^n.$$

Theorem 2.6 *The matrix $T = T_\rho = (\tau_{ik})_{i,k=1}^n$ where*

$$\tau_{ik} = \operatorname{Re} \left[\frac{1}{\alpha_i - \beta_k} \frac{\tilde{x}_k^0 \beta_k L_\delta(-\alpha_i) L_c(-\alpha_i) L_\alpha(-\beta_k)}{\tilde{y}_i^0 \alpha_i L_\delta(-\beta_k) L_c(-\beta_k) L'_\alpha(-\alpha_i)} \right] \quad (2.23)$$

satisfies (2.22).

Proof The range of the map $\mathbf{C}^n \rightarrow U, x \mapsto M[x; 0]$ (see 2.20) is precisely the n -dimensional subspace

$$V := L_\delta L_c \cdot \mathcal{P}_{n-1} = \{L_\delta L_c R; R \in \mathcal{P}_{n-1}\} \subset U. \quad (2.24)$$

We introduce a basis $(Q_k)_{k=1}^n$ for V by

$$Q_k(t) = \frac{L_\delta(t) L_c(t) L_\beta(t)}{t + \beta_k} \frac{\sqrt{\beta_k L'(-\beta_k) P(-\beta_k)}}{L_\delta(-\beta_k) L_c(-\beta_k) L'_\beta(-\beta_k)}.$$

Then

$$\frac{Q_k(-\beta_i)}{\sqrt{\beta_i L'(-\beta_i) P(-\beta_i)}} = \begin{cases} 1 & i = k, \\ 0 & i \neq k. \end{cases}$$

Denoting by (e_i) the canonical basis in \mathbf{C}^n and using (2.18), (2.19) we get

$$\begin{aligned} \tau_{ik} &= (Te_k)_i = \frac{Q_k(-\alpha_i)}{\sqrt{\alpha_i L'(-\alpha_i) P(-\alpha_i)}} \\ &= \frac{1}{\beta_k - \alpha_i} \frac{L_\delta(-\alpha_i) L_c(-\alpha_i) L_\beta(-\alpha_i)}{L_\delta(-\beta_k) L_c(-\beta_k) L'_\beta(-\beta_k)} \left(\frac{\beta_k L'(-\beta_k) P(-\beta_k)}{-\alpha_i L'(-\alpha_i) P(-\alpha_i)} \right)^{1/2}. \end{aligned}$$

Inserting the expressions (2.12) for $P(-\alpha_i)$ and $P(-\beta_k)$ and taking real parts (see the remarks in Sect. 2.6), we obtain the formula (2.23). \square

Remark 2.7 It is easy to see that, if we pick all matrix-elements real, some elements τ_{ik} of the matrix T in (2.23) will be negative, even while the numbers x_i^0 and y_k^0 are positive. It was proved in [2], Theorem 2.3, that this is necessarily so. Indeed, one there constructs an example of a five-dimensional couple $\overline{\ell}_2^5(\lambda)$ and two vectors $x^0, y^0 \in \mathbf{R}^5$ having non-negative entries such that *no* contraction $T = (\tau_{ik})_{i,k=1}^5$ on $\overline{\ell}_2^5(\lambda)$ having all matrix entries $\tau_{ik} \geq 0$ can satisfy $Tx^0 = y^0$. On the other hand, if one settles for using a matrix with $\|T\| \leq \sqrt{2}$, then it is possible to find one with only non-negative matrix entries. Indeed, such a matrix was used by Sedaev [35], see also [38].

2.8 On Sharpness of the Norm-Bounds

We shall show that if $m < n$ (i.e., if the polynomial P has at least one non-real zero), then the norm $\|T\|_{\mathcal{L}(\mathcal{H}_i)}$ of the contraction T constructed above is very close to 1 for $i = 0, 1$.

We first claim that $\|T\|_{\mathcal{L}(\mathcal{H}_0)} = 1$. To see this, we notice that if $m < n$, then there is a non-trivial polynomial $Q^{(1)}$ in the space V (see (2.24)) which vanishes at the points $0, \gamma_1, \dots, \gamma_{m-1}$. If $x_i^{(1)}$ and $y_i^{(1)}$ are defined by the formulas (2.18) and (2.19) (while $(x_j^{(1)})' = (y_k^{(1)})' = 0$), we then have $Tx^{(1)} = y^{(1)}$ and

$$k_\beta(t, x^{(1)}) - k_\alpha(t, y^{(1)}) = \frac{|Q^{(1)}(t)|^2}{L(t)P(t)}, \quad t > 0.$$

Choosing $t = 0$ we conclude that $\|x^{(1)}\|_{\ell_2^n}^2 - \|Tx^{(1)}\|_{\ell_2^n}^2 = 0$, whence $\|T\|_{\mathcal{L}(\mathcal{H}_0)} \geq 1$, proving our claim.

Similarly, the condition $m < n$ implies the existence of a polynomial $Q^{(2)} \in V$ of degree at most $n + m - 2$ vanishing at the points $\gamma_1, \dots, \gamma_{m-1}$. Constructing vectors $x^{(2)}, y^{(2)}$ via (2.18) and (2.19) we will have $Tx^{(2)} = y^{(2)}$ and

$$k_\beta(t, x^{(2)}) - k_\alpha(t, y^{(2)}) = \frac{|Q^{(2)}(t)|^2}{L(t)P(t)}, \quad t > 0.$$

Multiplying this relation by t and then sending $t \rightarrow \infty$, we find that $\|x^{(2)}\|_{\ell_2(\beta)}^2 - \|Tx^{(2)}\|_{\ell_2(\alpha)}^2 = 0$, which implies $\|T\|_{\mathcal{L}(\mathcal{H}_1)} \geq \rho^{-1/2}$.

2.9 A Remark on Weighted ℓ_p -Couples

As far as we are aware, if $1 < p < \infty$ and $p \neq 2$, it is still an open question whether the couple $\overline{\ell}_p^n(\lambda) = (\ell_p^n, \ell_p^n(\lambda))$ is an exact Calderón couple or not. (When $p = 1$ or $p = \infty$ it is exact Calderón; see [36] for the case $p = 1$; the case $p = \infty$ is essentially just the Hahn–Banach theorem.)

It is well known, and easy to prove, that the K_p -functional (see (2.2)) corresponding to the couple $\overline{\ell}_p^n(\lambda)$ is given by the explicit formula

$$K_p(t, x; \overline{\ell}_p^n(\lambda)) = \sum_{i=1}^n |x_i|^p \frac{t\lambda_i}{(1 + (t\lambda_i)^{\frac{1}{p-1}})^{p-1}}.$$

It was proved by Sedaev [35] (cf. [38]) that if $K_p(t, y^0; \overline{\ell}_p^n(\lambda)) \leq K_p(t, x^0; \overline{\ell}_p^n(\lambda))$ for all $t > 0$ then there is $T : \overline{\ell}_p^n(\lambda) \rightarrow \overline{\ell}_p^n(\lambda)$ of norm at most $2^{1/p'}$ such that $Tx^0 = y^0$. (Here p' is the exponent conjugate to p .)

Although our present estimates are particular for the case $p = 2$, our construction still shows that, if we re-define $P(t)$ to be the polynomial

$$\frac{P(t)}{L(t)} = \sum_1^n (\tilde{x}_i^0)^p \frac{\beta_i}{t + \beta_i} - \sum_1^n (\tilde{y}_i^0)^p \frac{\alpha_i}{t + \alpha_i}, \tag{2.25}$$

then the matrix T defined by

$$\tau_{ik} = \operatorname{Re} \left[\frac{1}{\alpha_i - \beta_k} \frac{(\tilde{x}_k^0)^{p-1}}{(\tilde{y}_i^0)^{p-1}} \frac{\beta_k L_\delta(-\alpha_i) L_c(-\alpha_i) L_\alpha(-\beta_k)}{\alpha_i L_\delta(-\beta_k) L_c(-\beta_k) L'_\alpha(-\alpha_i)} \right] \tag{2.26}$$

will satisfy $T\tilde{x}^0 = \tilde{y}^0$, at least, provided that $P(t) > 0$ when $t \geq 0$. (Here L_δ and L_c are constructed from the zeros of P as in the case $p = 2$.)

The matrix (2.26) differs from those used by Sedaev [35] and Sparr [38]. Indeed the matrices from [35, 38] have *non-negative entries*, while this is not so for the matrices (2.26). It seems to be an interesting problem to estimate the norm $\|T\|_{\mathcal{L}(\overline{\ell}_p^n(\lambda))}$ for the matrix (2.26), when $p \neq 2$. The motivation for this type of question is somewhat elaborated in Sect. 6.7, but we shall not discuss it further here.

2.10 A Comparison with Löwner's Matrix

In this subsection, we briefly explain how our matrix T is related to the matrix used by Löwner [26] in his original work on monotone matrix functions.²

We shall presently display four kinds of partial isometries; Löwner's matrix will be recognized as one of them. In all cases, operators with the required properties can alternatively be found using the more general construction in Theorem 2.4.

The following discussion was inspired by the earlier work of Sparr [39], who seems to have been the first to note that Löwner's matrix could be constructed in a similar way.

In this subsection, scalars are assumed to be real. In particular, when we write " ℓ_2^n " we mean the (real) Euclidean n -dimensional space.

Suppose that two vectors $x^0, y^0 \in \mathbf{R}^n$ satisfy

$$k_\lambda(t, y^0) \leq k_\lambda(t, x^0), \quad t > 0.$$

Let

$$L_\lambda(t) = \prod_1^n (t + \lambda_i),$$

and let $P \in \mathcal{P}_{n-1}$ be the polynomial fulfilling

$$\frac{P(t)}{L_\lambda(t)} = k_\lambda(t, x^0) - k_\lambda(t, y^0) = \sum_{i=1}^n \frac{\lambda_i}{t + \lambda_i} \left[(x_i^0)^2 - (y_i^0)^2 \right].$$

By assumption, $P(t) \geq 0$ for $t \geq 0$. Moreover, P is uniquely determined by the n conditions

$$P(-\lambda_i) = \frac{(x_i^0)^2 - (y_i^0)^2}{\lambda_i L'_\lambda(-\lambda_i)}.$$

Let $u_1, v_1, u_2, v_2, \dots$ denote the canonical basis of ℓ_2^n and let

$$\ell_2^n = O \oplus E$$

be the corresponding splitting, i.e.,

$$O = \text{span} \{u_i\} \quad , \quad E = \text{span} \{v_i\}.$$

²By "Löwner's matrix," we mean the unitary matrix denoted "V" in Donoghue's book [12], on p. 71. A more explicit construction of this matrix is found in [26], where it is called "T."

Notice that

$$\dim O = \lfloor (n-1)/2 \rfloor + 1 \quad , \quad \dim E = \lfloor (n-2)/2 \rfloor + 1,$$

where $\lfloor x \rfloor$ is the integer part of a real number x .

We shall construct matrices $T \in M_n(\mathbf{R})$ such that

$$Tx^0 = y^0 \quad \text{and} \quad k_\lambda(t, Tx) \leq k_\lambda(t, x), \quad t > 0, x \in \mathbf{R}^n, \quad (2.27)$$

in the following special cases:

- (1) $P(t) = q(t)^2$ where $q \in \mathcal{P}_{(n-1)/2}(\mathbf{R})$, $x^0 \in O$, and $y^0 \in E$,
- (2) $P(t) = tq(t)^2$ where $q \in \mathcal{P}_{(n-2)/2}(\mathbf{R})$, $x^0 \in E$, and $y^0 \in O$.

Here \mathcal{P}_x should be interpreted as $\mathcal{P}_{\lfloor x \rfloor}$.

Remark 2.8 In this connection, it is interesting to recall the well-known fact that any polynomial P which is non-negative on \mathbf{R}_+ can be written $P(t) = q_0(t)^2 + tq_1(t)^2$ for some real polynomials q_0 and q_1 .

To proceed with the solution, we rename the λ_i as $\lambda_i = \xi_i$ when i is odd and $\lambda_i = \eta_i$ when i is even. We also write

$$L_\xi(t) = \prod_{i \text{ odd}} (t + \xi_i) \quad , \quad L_\eta(t) = \prod_{i \text{ even}} (t + \eta_i),$$

and write $L = L_\xi L_\eta$. Notice that $L'_\lambda(-\xi_i) > 0$ and $L'_\lambda(-\eta_i) < 0$.

2.10.1 Case 1

Suppose that $P(t) = q(t)^2$, $q \in \mathcal{P}_{(n-1)/2}(\mathbf{R})$, $x^0 \in O$, and $y^0 \in E$. Then

$$\frac{q(t)^2}{L_\lambda(t)} = \sum_{k \text{ odd}} \frac{\xi_k}{t + \xi_k} (x_k^0)^2 - \sum_{i \text{ even}} \frac{\eta_i}{t + \eta_i} (y_i^0)^2,$$

where

$$x_k^0 = \frac{\varepsilon_k q(-\xi_k)}{\sqrt{\xi_k L'_\lambda(-\xi_k)}} \quad , \quad y_i^0 = \frac{\zeta_i q(-\eta_i)}{\sqrt{-\eta_i L'_\lambda(-\eta_i)}} \quad (2.28)$$

for some choice of signs $\varepsilon_k, \zeta_i \in \{\pm 1\}$.

By (2.28) are defined linear maps

$$O \rightarrow \mathcal{P}_{(n-1)/2}(\mathbf{R}) \quad : \quad x \mapsto Q \quad ; \quad \mathcal{P}_{(n-1)/2}(\mathbf{R}) \rightarrow E \quad : \quad Q \mapsto y.$$

The composition is a linear map

$$T_0 : O \rightarrow E \quad : \quad x \mapsto y.$$

We now define $T \in M_n(\mathbf{R})$ by

$$T : O \oplus E \rightarrow O \oplus E \quad : \quad [x; v] \mapsto [0; T_0x].$$

Then clearly $Tx^0 = y^0$ and

$$\begin{aligned} k_\lambda(t, [x; v]) - k_\lambda(t, T[x; v]) \\ &\geq k_\xi(t, x) - k_\eta(t, T_0x) \\ &= \frac{Q(t)^2}{L_\lambda(t)} \geq 0, \quad t > 0, x \in O, v \in E. \end{aligned} \tag{2.29}$$

We have verified (2.27) in case 1. A computation similar to the one in the proof of Theorem 2.6 shows that, with respect to the bases u_k and v_i ,

$$(T_0)_{ik} = \frac{\varepsilon_k \zeta_i}{\xi_k - \eta_i} \frac{L_\xi(-\eta_i)}{L'_\xi(-\xi_k)} \left(\frac{\xi_k L'_\xi(-\xi_k) L_\eta(-\xi_k)}{-\eta_i L_\xi(-\eta_i) L'_\eta(-\eta_i)} \right)^{1/2}.$$

Notice that, multiplying (2.29) by t , then letting $t \rightarrow \infty$ implies that

$$\sum_{k \text{ odd}} x_k^2 \xi_k - \sum_{i \text{ even}} (T_0x)_i^2 \eta_i = 0.$$

This means that T is a partial isometry from O to E with respect to the norm of $\ell_2^n(\lambda)$.

2.10.2 Case 2

Now assume that $P(t) = tq(t)^2$, $q \in \mathcal{P}_{(n-2)/2}(\mathbf{R})$, $x^0 \in E$, and $y^0 \in O$. Then

$$\frac{tq(t)^2}{L_\lambda(t)} = - \sum_{i \text{ odd}} (y_i^0)^2 \frac{\xi_i}{t + \xi_i} + \sum_{k \text{ even}} \frac{\eta_k}{t + \eta_k} (x_k^0)^2,$$

where

$$y_i^0 = \frac{\varepsilon'_i q(-\xi_i)}{\sqrt{L'_\lambda(-\xi_i)}}, \quad x_k^0 = \frac{-\zeta'_k q(-\eta_k)}{\sqrt{-L'_\lambda(-\eta_k)}} \tag{2.30}$$

for some $\varepsilon'_i, \zeta'_k \in \{\pm 1\}$.

By (2.30) are defined linear maps

$$E \rightarrow \mathcal{P}_{(n-2)/2}(\mathbf{R}) \quad : \quad x \mapsto Q \quad ; \quad \mathcal{P}_{(n-2)/2}(\mathbf{R}) \rightarrow O \quad : \quad Q \mapsto y.$$

We denote their composite by

$$T_1 : E \rightarrow O \quad : \quad x \mapsto y.$$

Define $T \in M_n(\mathbf{R})$ by

$$T : O \oplus E \rightarrow O \oplus E \quad : \quad [u; x] \mapsto [T_1x; 0].$$

We then have

$$\begin{aligned} -k_\lambda(t, T[u; x]) + k_\lambda(t, [u; x]) & \\ & \geq -k_\xi(t, T_1x) + k_\eta(t, x) \\ & = \frac{tQ(t)^2}{L_\lambda(t)} \geq 0, \quad t > 0, u \in O, x \in E, \end{aligned} \tag{2.31}$$

and (2.27) is verified also in case 2.

A computation shows that, with respect to the bases v_k and u_i ,

$$(T_1)_{ik} = \frac{\varepsilon'_i \zeta'_k}{\eta_k - \xi_i} \frac{L_\eta(-\xi_i)}{L'_\eta(-\eta_k)} \left(\frac{-L_\xi(-\eta_k)L'_\eta(-\eta_k)}{L'_\xi(-\xi_i)L_\eta(-\xi_i)} \right)^{1/2}.$$

Inserting $t = 0$ in (2.31) we find that

$$-\sum_{i \text{ odd}} (T_1x)_i^2 + \sum_{k \text{ even}} (x_k)^2 = 0,$$

i.e., T is a partial isometry form E to O with respect to the norm of ℓ_2^n .

In the case of even n , the matrix T_1 coincides with Löwner's matrix.

3 Quadratic Interpolation Spaces

3.1 A Classification of Quadratic Interpolation Spaces

Recall that an intermediate space X with respect to $\overline{\mathcal{H}}$ is said to be of *type H* if $\|T\|_{\mathcal{L}(\mathcal{H}_i)} \leq M_i$ for $i = 0, 1$ implies that $\|T\|_{\mathcal{L}(X)} \leq H(M_0, M_1)$. We shall henceforth make a mild restriction, and assume that H be homogeneous of degree

one. This means that we can write

$$H(s, t)^2 = s^2 \mathbf{H}(t^2/s^2) \tag{3.1}$$

for some function \mathbf{H} of one positive variable. In this situation, we will say that X is of type \mathbf{H} . The definition is chosen so that the estimates $\|T\|_{\mathcal{L}(\mathcal{H}_i)}^2 \leq M_i$ for $i = 0, 1$ imply $\|T\|_{\mathcal{L}(X)}^2 \leq M_0 \mathbf{H}(M_1/M_0)$.

In the following we will make the *standing assumptions*: \mathbf{H} is an increasing, continuous, and positive function on \mathbf{R}_+ with $\mathbf{H}(1) = 1$ and $\mathbf{H}(t) \leq \max\{1, t\}$.

Notice that our assumptions imply that all spaces of type \mathbf{H} are exact interpolation. Note also that $\mathbf{H}(t) = t^\theta$ corresponds to geometric interpolation of exponent θ .

Suppose now that $\overline{\mathcal{H}}$ is a regular Hilbert couple and that \mathcal{H}_* is an exact interpolation space with corresponding operator B . By Donoghue's lemma, we have that $B = h(A)$ for some positive Borel function h on $\sigma(A)$.

The statement that \mathcal{H}_* is intermediate relative to $\overline{\mathcal{H}}$ is equivalent to that

$$c_1 \frac{A}{1+A} \leq B \leq c_2(1+A) \tag{3.2}$$

for some positive numbers c_1 and c_2 .

Let us momentarily assume that \mathcal{H}_0 be *separable*. (This restriction is removed in Remark 3.1.) We can then define the *scalar-valued spectral measure* of A ,

$$\nu_A(\omega) = \sum 2^{-k} \langle E(\omega)e_k, e_k \rangle_0$$

where E is the spectral measure of A , $\{e_k; k = 1, 2, \dots\}$ is an orthonormal basis for \mathcal{H}_0 , and ω is a Borel set. Then, for Borel functions h_0, h_1 on $\sigma(A)$, one has that $h_1 = h_2$ almost everywhere with respect to ν_A if and only if $h_1(A) = h_2(A)$.

Note that the regularity of $\overline{\mathcal{H}}$ means that $\nu_A(\{0\}) = 0$.

Theorem II *If \mathcal{H}_* is of type \mathbf{H} with respect to $\overline{\mathcal{H}}$, then $B = h(A)$ where the function h can be modified on a null-set with respect to ν_A so that*

$$h(\lambda)/h(\mu) \leq \mathbf{H}(\lambda/\mu), \quad \lambda, \mu \in \sigma(A) \setminus \{0\}. \tag{3.3}$$

Proof Fix a (large) compact subset $K \subset \sigma(A) \cap \mathbf{R}_+$ and put $\mathcal{H}'_0 = \mathcal{H}'_1 = E_K(\mathcal{H}_0)$ where E is the spectral measure of A , and the norms are defined by restriction,

$$\|x\|_{\mathcal{H}'_i} = \|x\|_{\mathcal{H}_i} \quad , \quad \|x\|_{\mathcal{H}'_*} = \|x\|_{\mathcal{H}_*} \quad , \quad x \in E_K(\mathcal{H}_0).$$

It is clear that the operator A' corresponding to $\overline{\mathcal{H}'}$ is the compression of A to \mathcal{H}'_0 and likewise the operator B' corresponding to \mathcal{H}'_* is the compression of B to \mathcal{H}'_0 . Moreover, \mathcal{H}'_* is of interpolation type \mathbf{H} with respect to $\overline{\mathcal{H}'}$ and the operator $B' = (h|_K)(A')$. For this reason, and since the compact set K is arbitrary, it clearly

suffices to prove the statement with $\overline{\mathcal{H}}$ replaced by $\overline{\mathcal{H}'}$. Then A is bounded above and below. Moreover, by (3.2), also B is bounded above and below.

Let $c < 1$ be a positive number such that $\sigma(A) \subset (c, c^{-1})$. For a fixed $\varepsilon > 0$ with $\varepsilon < c/2$ we set

$$E_\lambda = \sigma(A) \cap (\lambda - \varepsilon, \lambda + \varepsilon)$$

and consider the functions

$$m_\varepsilon(\lambda) = \text{ess inf}_{E_\lambda} h, \quad M_\varepsilon(\lambda) = \text{ess sup}_{E_\lambda} h,$$

the essential inf and sup being taken with respect to ν_A .

Now fix a small positive number ε' and two unit vectors e_λ, e_μ supported by E_λ, E_μ , respectively, such that

$$\|e_\lambda\|_*^2 \geq M_\varepsilon(\lambda) - \varepsilon', \quad \|e_\mu\|_*^2 \leq m_\varepsilon(\mu) + \varepsilon'.$$

Now fix $\lambda, \mu \in \sigma(A)$ and let $Tx = \langle x, e_\mu \rangle_0 e_\lambda$. Then

$$\begin{aligned} \|Tx\|_1^2 &= |\langle x, e_\mu \rangle_0|^2 \|e_\lambda\|_1^2 \leq \frac{1}{(\mu - \varepsilon)^2} |\langle x, e_\mu \rangle_1|^2 (\lambda + \varepsilon) \\ &\leq \frac{(\mu + \varepsilon)(\lambda + \varepsilon)}{(\mu - \varepsilon)^2} \|x\|_1^2. \end{aligned}$$

Likewise,

$$\|Tx\|_0^2 \leq |\langle x, e_\mu \rangle_0|^2 \leq \|x\|_0^2,$$

so $\|T\| \leq 1$ and $\|T\|_A^2 \leq \alpha_{\mu, \lambda, \varepsilon}$ where $\alpha_{\mu, \lambda, \varepsilon} = \frac{(\mu + \varepsilon)(\lambda + \varepsilon)}{(\mu - \varepsilon)^2}$.

Since \mathcal{H}_* is of type **H**, we conclude that

$$\|T\|_B^2 \leq \mathbf{H}(\alpha_{\mu, \lambda, \varepsilon}),$$

whence

$$\begin{aligned} M_\varepsilon(\lambda) - \varepsilon' &\leq \|e_\lambda\|_*^2 = \|Te_\mu\|_*^2 \leq \mathbf{H}(\alpha_{\mu, \lambda, \varepsilon}) \|e_\mu\|_*^2 \\ &\leq \mathbf{H}(\alpha_{\mu, \lambda, \varepsilon}) (m_\varepsilon(\mu) + \varepsilon'). \end{aligned} \tag{3.4}$$

In particular, since ε' was arbitrary, and $m_\varepsilon(\lambda) \leq \|e_\lambda\|_*^2 \leq \|B\|$, we find that

$$M_\varepsilon(\lambda) - m_\varepsilon(\lambda) \leq [\mathbf{H}(\alpha_{\mu, \lambda, \varepsilon}) - 1] \|B\|.$$

By assumption, \mathbf{H} is continuous and $\mathbf{H}(1) = 1$. Hence, as $\varepsilon \downarrow 0$, the functions $M_\varepsilon(\lambda)$ diminish monotonically, converging uniformly to a function $h_*(\lambda)$ which is also the uniform limit of the family $m_\varepsilon(\lambda)$. It is clear that h_* is continuous, and since $m_\varepsilon \leq h_* \leq M_\varepsilon$, we have $h_* = h$ almost everywhere with respect to ν_A . The relation (3.3) now follows for $h = h_*$ by letting ε and ε' tend to zero in (3.4). \square

A partial converse to Theorem II is found below, see Theorem 6.3.

Remark 3.1 (The Non-Separable Case) Now consider the case when \mathcal{H}_0 is non-separable. (By regularity this means that also \mathcal{H}_1 and \mathcal{H}_* are non-separable.)

First assume that the operator A is bounded. Let \mathcal{H}'_0 be a separable reducing subspace for A such that the restriction A' of A to \mathcal{H}'_0 has the same spectrum as A . The space \mathcal{H}'_0 reduces B by Donoghue's lemma; by Theorem II the restriction B' of B to \mathcal{H}'_0 satisfies $B' = h'(A')$ for some continuous function h' satisfying (3.3) on $\sigma(A)$. Let \mathcal{H}''_0 be any other separable reducing subspace, where (as before) $B'' = h''(A'')$. Then $\mathcal{H}'_0 \oplus \mathcal{H}''_0$ is a separable reducing subspace on which $B = h(A)$ for some third continuous function h on $\sigma(A)$. Then $h(A') \oplus h(A'') = h'(A') \oplus h''(A'')$ and by continuity we must have $h = h' = h''$ on $\sigma(A)$. The function h thus satisfies $B = h(A)$ as well as the estimate (3.3).

If A is unbounded, we replace A by its compression to $P_n \mathcal{H}_0$ where P_n is the spectral projection of A corresponding to the spectral set $[0, n] \cap \sigma(A)$, $n = 1, 2, \dots$. The same reasoning as above shows that B appears as a continuous function of A on $\sigma(A) \cap [0, n]$. Since n is arbitrary, we find that $B = h(A)$ for a function h satisfying (3.3).

3.2 Geometric Interpolation

Now consider the particular case when \mathcal{H}_* is of exponent θ , viz. of type $\mathbf{H}(t) = t^\theta$ with respect to $\overline{\mathcal{H}}$. We write $B = h(A)$ where h is the continuous function provided by Theorem II (and Remark 3.1 in the non-separable case).

Fix a point $\lambda_0 \in \sigma(A)$ and let $C = h(\lambda_0)\lambda_0^{-\theta}$. The estimate (3.3) then implies that $h(\lambda) \leq C\lambda^\theta$ and $h(\mu) \geq C\mu^\theta$ for all $\lambda, \mu \in \sigma(A)$. We have proved the following theorem.

Theorem 3.2 ([27, 40]) *If \mathcal{H}_* is an exact interpolation Hilbert space of exponent θ relative to $\overline{\mathcal{H}}$, then $B = h(A)$ where $h(\lambda) = C\lambda^\theta$ for some positive constant C .*

Theorem 3.2 says that $\mathcal{H}_* = \mathcal{H}_\theta$ up to a constant multiple of the norm, where \mathcal{H}_θ is the space defined in (1.7). In the guise of operator inequalities: for any fixed positive operators A and B , the condition

$$T^*T \leq M_0 \quad , \quad T^*AT \leq M_1A \quad \Rightarrow \quad T^*BT \leq M_0^{1-\theta} M_1^\theta B$$

is equivalent to that $B = A^\theta$.

It was observed in [27] that \mathcal{H}_θ also equals to the complex interpolation space $C_\theta(\overline{\mathcal{H}})$. For the sake of completeness, we supply a short proof of this fact in the appendix.

Remark 3.3 An exact quadratic interpolation method, the *geometric mean* was introduced earlier by Pusz and Woronowicz [33] (it corresponds to the $C_{1/2}$ -method). In [40], Uhlmann generalized that method to a method (the *quadratic mean*) denoted QI_t where $0 < t < 1$; this method is quadratic and of exponent t .

In view of Theorem 3.2 and the preceding remarks we can conclude that $QI_\theta(\overline{\mathcal{H}}) = C_\theta(\overline{\mathcal{H}}) = \mathcal{H}_\theta$ for any regular Hilbert couple $\overline{\mathcal{H}}$. We refer to [40] for several physically relevant applications of this type of interpolation.

Finally, we want to mention that in [32] Peetre introduces the “Riesz method of interpolation”; in Section 5 he also defines a related method “QM” which comes close to the complex $C_{1/2}$ -method.

3.3 Donoghue’s Theorem

The exact quadratic interpolation spaces relative to a Hilbert couple were characterized by Donoghue in the paper [14]. We shall here prove the following equivalent version of Donoghue’s result (see [2, 3]).

Theorem III *An intermediate Hilbert space \mathcal{H}_* relative to $\overline{\mathcal{H}}$ is an exact interpolation space if and only if there is a positive radon measure ϱ on $[0, \infty]$ such that*

$$\|x\|_*^2 = \int_{[0, \infty]} (1 + t^{-1}) K(t, x; \overline{\mathcal{H}}) d\varrho(t).$$

Equivalently, \mathcal{H}_ is exact interpolation relative to $\overline{\mathcal{H}}$ if and only if the corresponding operator B can be represented as $B = h(A)$ for some function $h \in P'$.*

The statements that all norms of the given form are exact quadratic interpolation norms have already been shown (see Sect. 1.2). There remains to prove that there are no others.

Donoghue’s original formulation of the result, as well as other equivalent forms of the theorem, is found in Sect. 6 below. Our present approach follows [2] and is based on K -monotonicity.

Remark 3.4 The condition that \mathcal{H}_* be exact interpolation with respect to $\overline{\mathcal{H}}$ means that \mathcal{H}_* is of type \mathbf{H} where $\mathbf{H}(t) = \max\{1, t\}$. In view of Theorem II (and Remark 3.1), this means that we can represent $B = h(A)$ where h is *quasi-concave* on $\sigma(A) \setminus \{0\}$,

$$h(\lambda) \leq h(\mu) \max\{1, \lambda/\mu\}, \quad \lambda, \mu \in \sigma(A) \setminus \{0\}. \tag{3.5}$$

In particular, h is locally Lipschitzian on $\sigma(A) \cap \mathbf{R}_+$.

Remark 3.5 A related result concerning *non-exact* quadratic interpolation was proved by Ovchinnikov [30] using Donoghue’s theorem. Cf. also [4].

3.4 The Proof for Simple Finite-Dimensional Couples

Similar to our approach to Calderón’s problem, our strategy is to reduce Theorem III to a case of “simple couples.”

Theorem 3.6 *Assume that $\mathcal{H}_0 = \mathcal{H}_1 = \mathbf{C}^n$ as sets and that all eigenvalues $(\lambda_i)_1^n$ of the corresponding operator A are of unit multiplicity. Consider a third Hermitian norm $\|x\|_*^2 = \langle Bx, x \rangle_0$ on \mathbf{C}^n . Then \mathcal{H}_* is exact interpolation with respect to $\overline{\mathcal{H}}$ if and only if $B = h(A)$ where $h \in P'$.*

Remark 3.7 The lemma says that the class of functions h on $\sigma(A)$ satisfying

$$T^*T \leq 1 \quad , \quad T^*AT \leq A \quad \Rightarrow \quad T^*h(A)T \leq h(A), \quad (T \in M_n(\mathbf{C})) \quad (3.6)$$

is precisely the set $P'|\sigma(A)$ of restrictions of P' -functions to $\sigma(A)$. In this way, the condition (3.6) provides an operator-theoretic solution to the interpolation problem by positive Pick functions on a finite subset of \mathbf{R}_+ .

Proof of Theorem 3.6 We already know that the spaces \mathcal{H}_* of the asserted form are exact interpolation relative to $\overline{\mathcal{H}}$ (see Sects. 1.2 and 1.3).

Now let \mathcal{H}_* be any exact quadratic interpolation space. By Donoghue’s lemma and the argument in Sect. 2.3, we can for an appropriate positive sequence $\lambda = (\lambda_i)_1^n$ identify $\overline{\mathcal{H}} = \overline{\ell}_2^n(\lambda)$, $A = \text{diag}(\lambda_i)$, and $B = h(A)$ where h is some positive function defined on $\sigma(A) = \{\lambda_i\}_1^n$.

Our assumption is that $\ell_2^n(h(\lambda))$ is exact interpolation relative to $\overline{\ell}_2^n(\lambda)$. We must prove that $h \in P'|\sigma(A)$. To this end, write

$$k_{\lambda_i}(t) = \frac{(1+t)\lambda_i}{1+t\lambda_i},$$

and recall that (see Lemma 1.1)

$$K\left(t, x; \overline{\ell}_2^n(\lambda)\right) = \left(1+t^{-1}\right)^{-1} \sum_1^n |x_i|^2 k_{\lambda_i}(t).$$

Let us denote by C the algebra of continuous complex functions on $[0, \infty]$ with the supremum norm $\|u\|_\infty = \sup_{t>0} |u(t)|$. Let $V \subset C$ be the linear span of the k_{λ_i}

for $i = 1, \dots, n$. We define a positive functional ϕ on V by

$$\phi\left(\sum_1^n a_i k_{\lambda_i}\right) = \sum_1^n a_i h(\lambda_i).$$

We claim that ϕ is a *positive functional*, i.e., if $u \in V$ and $u(t) \geq 0$ for all $t > 0$, then $\phi(u) \geq 0$.

To prove this let $u = \sum_1^n a_i k_{\lambda_i}$ be non-negative on \mathbf{R}_+ and write $a_i = |x_i|^2 - |y_i|^2$ for some $x, y \in \mathbf{C}^n$. The condition that $u \geq 0$ means that

$$\begin{aligned} (1 + t^{-1}) K\left(t, x; \overline{\ell}_2^n(\lambda)\right) &= \sum_{i=1}^n |x_i|^2 k_{\lambda_i}(t) \\ &\geq \sum_{i=1}^n |y_i|^2 k_{\lambda_i}(t) \\ &= (1 + t^{-1}) K\left(t, y; \overline{\ell}_2^n(\lambda)\right), \quad t > 0. \end{aligned} \tag{3.7}$$

Since $\overline{\ell}_2^n(\lambda)$ is an exact Calderón couple (by Theorem 2.4), the space $\ell_2^n(h(\lambda))$ is exact K -monotonic. In other words, (3.7) implies that

$$\|x\|_{\ell_2^n(h(\lambda))} \geq \|y\|_{\ell_2^n(h(\lambda))},$$

i.e.,

$$\phi(u) = \sum_1^n \left(|x_i|^2 - |y_i|^2\right) h(\lambda_i) \geq 0.$$

The asserted positivity of ϕ is thereby proved.

Replacing λ_i by $c\lambda_i$ for a suitable positive constant c we can without losing generality assume that $1 \in \sigma(A)$, i.e., that the unit $\mathbf{1}(x) \equiv 1$ of the C^* -algebra C belongs to V . The positivity of ϕ then ensures that

$$\|\phi\| = \sup_{u \in V; \|u\|_\infty \leq 1} |\phi(u)| = \phi(\mathbf{1}).$$

Let Φ be a Hahn–Banach extension of ϕ to C and note that

$$\|\Phi\| = \|\phi\| = \phi(\mathbf{1}) = \Phi(\mathbf{1}).$$

This means that Φ is a positive functional on C (cf. [29], §3.3). By the Riesz representation theorem there is thus a positive Radon measure ϱ on $[0, \infty]$ such

that

$$\Phi(u) = \int_{[0, \infty]} u(t) d\varrho(t), \quad u \in C.$$

In particular

$$h(\lambda_i) = \phi(k_{\lambda_i}) = \Phi(k_{\lambda_i}) = \int_{[0, \infty]} \frac{(1+t)\lambda_i}{1+t\lambda_i} d\varrho(t), \quad i = 1, \dots, n.$$

We have shown that h is the restriction to $\sigma(A)$ of a function of class P' . □

3.5 The Proof of Donoghue's Theorem

We here prove Theorem III in full generality.

We remind the reader that if $S \subset \mathbf{R}_+$ is a subset, we write $P'|S$ for the convex cone of restrictions of P' -functions to S . We first collect some simple facts about this cone.

Lemma 3.8

- (i) The class $P'|S$ is closed under pointwise convergence.
- (ii) If S is finite and if $\lambda = (\lambda_i)_{i=1}^n$ is an enumeration of the points of S , then h belongs to $P'|S$ if and only if $\ell_2^n(h(\lambda))$ is exact interpolation with respect to the pair $\overline{\ell_2^n}(\lambda)$.
- (iii) If S is infinite, then a continuous function h on S belongs to $P'|S$ if and only if $h \in P'|\Lambda$ for every finite subset $\Lambda \subset S$.

Proof

- (i) Let h_n be a sequence in P' converging pointwise on S and fix $\lambda \in S$. It is clear that the boundedness of the numbers $h_n(\lambda)$ is equivalent to boundedness of the total masses of the corresponding measures ϱ_n on the compact set $[0, \infty]$. It now suffices to apply Helly's selection theorem.
- (ii) This is Theorem 3.6.
- (iii) Let Λ_n be an increasing sequence of finite subsets of S whose union is dense. Let $h_n = h|\Lambda_n$ where h is continuous on S . If $h_n \in P'|\Lambda_n$ for all n , then the sequence h_n converges pointwise on $\cup \Lambda_n$ to h . By part (i) we then have $h \in P'|\sigma(A)$. □

We can now finish the proof of Donoghue's theorem (Theorem III).

Let \mathcal{H}_* be exact interpolation with respect to $\overline{\mathcal{H}}$ and represent the corresponding operator as $B = h(A)$ where h satisfies (3.3). By the remarks after Theorem III, the function h is locally Lipschitzian.

In view of Lemma 3.8 we shall be done when we have proved that $\ell_2^n(h(\lambda))$ is exact interpolation with respect to $\overline{\ell_2^n}(\lambda)$ for all sequences $\lambda = (\lambda_i)_1^n \subset \sigma(A)$ of distinct points. Let us arrange the sequences in the increasing order: $0 < \lambda_1 < \dots < \lambda_n$.

Fix $\varepsilon > 0$, $\varepsilon < \min\{c, \lambda_1, 1/\lambda_n\}$ and let $E_i = [\lambda_i - \varepsilon, \lambda_i + \varepsilon] \cap \sigma(A)$; we assume that ε is sufficiently small that the E_i be disjoint. Let $M = \cup_1^n E_i$. We can assume that h has Lipschitz constant at most 1 on M .

Let \mathcal{M} be the reducing subspace of \mathcal{H}_0 corresponding to the spectral set M , and let \tilde{A} be the compression of A to \mathcal{M} . We define a function g on M by $g(\lambda) = \lambda_i$ on E_i . Then $|g(\lambda) - \lambda| < \varepsilon$ on $\sigma(\tilde{A})$, so

$$\|\tilde{A} - g(\tilde{A})\| \leq \varepsilon \quad , \quad \|h(\tilde{A}) - h(g(\tilde{A}))\| \leq \varepsilon. \tag{3.8}$$

Lemma 3.9 *Suppose that $A', A'' \in \mathcal{L}(\mathcal{M})$ satisfy $A', A'' \geq \delta > 0$ and $\|A' - A''\| \leq \varepsilon$. Then $\|T\|_{A''} \leq \sqrt{1 + 2\varepsilon/\delta} \max\{\|T\|, \|T\|_{A'}\}$ for all $T \in \mathcal{L}(\mathcal{M})$.*

Proof By definition, $\|T\|_{A'}$ is the smallest number $C \geq 0$ such that $T^*A'T \leq C^2A'$. Thus

$$\begin{aligned} T^*A''T &= T^*(A'' - A')T + T^*A'T \\ &\leq \|T\|^2 \varepsilon + \|T\|_{A'}^2 (A'' + (A' - A'')) \\ &\leq 2\varepsilon \max\{\|T\|^2, \|T\|_{A'}^2\} + \|T\|_{A'}^2 A'' \\ &\leq \max\{\|T\|^2, \|T\|_{A'}^2\} (1 + 2\varepsilon/\delta) A''. \end{aligned}$$

□

We can find $\delta > 0$ such that the operators \tilde{A} , $g(\tilde{A})$, $h(\tilde{A})$, and $h(g(\tilde{A}))$ are $\geq \delta$. Then by repeated use of Lemma 3.9,

$$\begin{aligned} \|T\|_{h(g(\tilde{A}))} &\leq \sqrt{1 + 2\varepsilon/\delta} \max\{\|T\|, \|T\|_{h(\tilde{A})}\} \\ &\leq \sqrt{1 + 2\varepsilon/\delta} \max\{\|T\|, \|T\|_{\tilde{A}}\} \\ &\leq (1 + 2\varepsilon/\delta) \max\{\|T\|, \|T\|_{g(\tilde{A})}\}, \quad T \in \mathcal{L}(\mathcal{M}). \end{aligned}$$

Let e_i be a unit vector supported by the spectral set E_i and define a space $\mathcal{V} \subset \mathcal{M}$ to be the n -dimensional space spanned by the e_i . Let A_0 be the compression of $g(\tilde{A})$ to \mathcal{V} , then

$$\|T\|_{h(A_0)} \leq (1 + 2\varepsilon/\delta) \max\{\|T\|, \|T\|_{A_0}\}, \quad T \in \mathcal{L}(\mathcal{V}). \tag{3.9}$$

Identifying \mathcal{V} with ℓ_2^n and A_0 with the matrix $\text{diag}(\lambda_i)$, we see that (3.9) is independent of ε . Letting ε diminish to 0 in (3.9) now gives that $\ell_2^n(h(\lambda))$ is exact interpolation with respect to $\overline{\ell_2^n}(\lambda)$. In view of Lemma 3.8, this finishes the proof of Theorem III. \square

4 Classes of Matrix Functions

In this section, we discuss the basic properties of interpolation functions: in particular, the relation to the well-known classes of monotone matrix functions. We refer to the books [12] and [34] for further reading on the latter classes.

4.1 Interpolation and Matrix Monotone Functions

Let A_1 and A_2 be positive operators in ℓ_2^n ($n = \infty$ is admitted). Suppose that $A_1 \leq A_2$ and form the following operators on $\ell_2^n \oplus \ell_2^n$:

$$T_0 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad A = \begin{pmatrix} A_2 & 0 \\ 0 & A_1 \end{pmatrix}.$$

It is then easy to see that $T_0^*T_0 \leq 1$ and that $T_0^*AT_0 = \begin{pmatrix} A_1 & 0 \\ 0 & 0 \end{pmatrix} \leq A$.

Now assume that a function h on $\sigma(A)$ belongs to the class C_A defined in Sect. 1.4, i.e., that h satisfies

$$T^*T \leq 1, \quad T^*AT \leq A \quad \Rightarrow \quad T^*h(A)T \leq h(A), \tag{4.1}$$

where T denotes an operator on ℓ_2^{2n} .

We then have $T_0^*h(A)T_0 \leq h(A)$, or

$$\begin{pmatrix} h(A_1) & 0 \\ 0 & 0 \end{pmatrix} \leq \begin{pmatrix} h(A_2) & 0 \\ 0 & h(A_1) \end{pmatrix}.$$

In particular, we find that $h(A_1) \leq h(A_2)$. We have shown that (under the assumptions above)

$$A_1 \leq A_2 \quad \Rightarrow \quad h(A_1) \leq h(A_2). \tag{4.2}$$

We now change our point of view slightly. Given a positive integer n , we let C_n denote the convex of positive functions h on \mathbf{R}_+ such that (4.1) holds for *all* positive operators A on ℓ_2^n and all $T \in \mathcal{L}(\ell_2^n)$.

Similarly, we let P'_n denote the class of all positive functions h on \mathbf{R}_+ such that $h(A_1) \leq h(A_2)$ whenever A_1, A_2 are positive operators on ℓ_2^n such that $A_1 \leq A_2$. We refer to P'_n as the cone of positive functions *monotone of order n* on \mathbf{R}_+ .

We have shown above that $C_{2n} \subset P'_n$.

In the other direction, assume that $h \in P'_{2n}$. Let A, T be bounded operators on ℓ_2^{2n} with $A > 0, T^*T \leq 1$ and $T^*AT \leq A$. Assume also that h be continuous. We will use the following lemma due to Hansen [19]. We recall the proof for completeness.

Lemma 4.1 ([19]) $T^*h(A)T \leq h(T^*AT)$.

Proof Put $S = (1 - TT^*)^{1/2}$ and $R = (1 - T^*T)^{1/2}$ and consider the $2n \times 2n$ matrix

$$U = \begin{pmatrix} T & S \\ R & -T^* \end{pmatrix}, \quad X = \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix}.$$

It is well known, and easy to check, that U is unitary and that

$$U^*XU = \begin{pmatrix} T^*AT & T^*AS \\ SAT & SAS \end{pmatrix}.$$

Next fix a number $\varepsilon > 0$, a constant $\lambda > 0$ (to be fixed), and form the matrix

$$Y = \begin{pmatrix} T^*AT + \varepsilon & 0 \\ 0 & 2\lambda \end{pmatrix}$$

which, provided that we choose $\lambda \geq \|SAS\|$, satisfies

$$Y - U^*XU = \begin{pmatrix} \varepsilon & -T^*AS \\ -SAT & 2\lambda - SAS \end{pmatrix} \geq \begin{pmatrix} \varepsilon & D \\ D^* & \lambda \end{pmatrix},$$

where we have written $D = -T^*AS$.

If we now also choose λ so that $\lambda \geq \|D\|^2/\varepsilon$, then we obtain for all $\xi, \eta \in \mathbf{C}^n$ that

$$\begin{aligned} \left\langle \begin{pmatrix} \varepsilon & D \\ D^* & \lambda \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \begin{pmatrix} \xi \\ \eta \end{pmatrix} \right\rangle &= \varepsilon\|\xi\|^2 + \langle D\eta, \xi \rangle + \langle D^*\xi, \eta \rangle + \lambda\|\eta\|^2 \\ &\geq \varepsilon\|\xi\|^2 - 2\|D\|\|\xi\|\|\eta\| + \lambda\|\eta\|^2 \geq 0. \end{aligned}$$

Hence $U^*XU \leq Y$ and as a consequence $U^*h(X)U = h(U^*XU) \leq h(Y)$, since h is matrix monotone of order $2n$. The last inequality means that

$$\begin{pmatrix} T^*h(A)T & T^*h(A)S \\ Sh(A)T & Sh(A)S \end{pmatrix} \leq \begin{pmatrix} h(T^*AT + \varepsilon) & 0 \\ 0 & h(2\lambda) \end{pmatrix},$$

so in particular $T^*h(A)T \leq h(T^*AT + \varepsilon)$. Since $\varepsilon > 0$ was arbitrary, and since h is assumed to be continuous, we conclude the lemma. \square

We now continue our discussion. Assuming that $T^*T \leq 1$ and $T^*AT \leq A$, and that $h \in P'_{2n}$ is continuous, we have $h(T^*AT) \leq h(A)$ [since $h \in P'_n$], so $T^*h(A)T \leq h(A)$ by Lemma 4.1. We conclude that $h \in C_n$.

To prove that $P'_{2n} \subset C_n$, we need to remove the continuity assumption on h made above. This is completely standard: let φ be a smooth positive function on \mathbf{R}_+ such that $\int_0^\infty \varphi(t) dt/t = 1$, and define a sequence h_k by $h_k(\lambda) = k^{-1} \int_0^\infty \varphi(\lambda^k/t^k) h(t) dt/t$. The class P'_{2n} is a convex cone, closed under pointwise convergence [12], so the functions h_1, h_2, \dots are of class P'_{2n} . They are furthermore continuous, so by the argument above, they are of class C_n . By Lemma 3.8, the cone C_n is also closed under pointwise convergence, so $h = \lim h_n \in C_n$.

To summarize, we have the inclusions $C_{2n} \subset P'_n, P'_{2n} \subset C_n$, and also $C_{n+1} \subset C_n, P'_{n+1} \subset P'_n$. In view of Theorem III, we have the identity $\cap_1^\infty C_n = P'$. The inclusions above now imply the following result, sometimes known as ‘‘L6wner’s theorem on matrix monotone functions.’’

Theorem 4.2 *We have $\cap_1^\infty P'_n = \cap_1^\infty C_n = P'$.*

The identity $\cap_1^\infty P'_n = P'$ says that a positive function h is monotone of all finite orders if and only if it is of class P' . The somewhat less precise fact that $P'_\infty = P'$ is interpreted as that the class of *operator monotone functions* coincides with P' .

The identity $C_\infty = P'$ is, except for notation, contained in the work of Foias and Lions, from [16]. See Sect. 6.4.

Note that the inclusion $P'_{2n} \subset C_n$ shows that a matrix monotone functions of order $2n$ can be interpolated by a positive Pick function at n points. Results of a similar nature, where it is shown, in addition, that an interpolating Pick function can be taken rational of a certain degree, are discussed, for example, in Donoghue’s book [14, Chapter XIII] or (more relevant in the present connection) in the paper [13].

It seems somewhat inaccurate to refer to the identity $\cap_1^\infty P'_n = P'$ as ‘‘L6wner’s theorem,’’ since L6wner discusses more subtle results concerning matrix monotone functions of a given finite order n . In spite of this, it is common nowadays to let ‘‘L6wner’s theorem’’ refer to this identity.

4.2 More on the Cone C_A

We can now give an short proof of the following result due to Donoghue [13].

Theorem 4.3 *For a positive function h on $\sigma(A)$ we define two positive functions \tilde{h} and h^* on $\sigma(A^{-1})$ by $\tilde{h}(\lambda) = \lambda h(1/\lambda)$ and $h^*(\lambda) = 1/h(1/\lambda)$. Then the following conditions are equivalent:*

- (i) $h \in C_A,$
- (ii) $\tilde{h} \in C_{A^{-1}},$
- (iii) $h^* \in C_{A^{-1}}.$

Proof Let \mathcal{H}_* be a quadratic intermediate space relative to a regular Hilbert couple $\overline{\mathcal{H}}$; let $B = h(A)$ be the corresponding operator. It is clear that \mathcal{H}_* is exact interpolation relative to $\overline{\mathcal{H}}$ if and only if \mathcal{H}_* is exact interpolation relative to the reverse couple $\overline{\mathcal{H}^{(r)}} = (\mathcal{H}_1, \mathcal{H}_0)$. The latter couple has corresponding operator A^{-1} and it is clear that the identity $\|x\|_*^2 = \langle h(A)x, x \rangle_0$ is equivalent to that $\|x\|_*^2 = \langle A^{-1}\tilde{h}(A^{-1})x, x \rangle_1$. We have shown the equivalence of (i) and (ii).

Next let $\overline{\mathcal{H}^*} = (\mathcal{H}_0^*, \mathcal{H}_1^*)$ be the dual couple, where we identify $\mathcal{H}_0^* = \mathcal{H}_0$. With this identification, \mathcal{H}_1^* becomes associated with the norm $\|x\|_{\mathcal{H}_1^*}^2 = \langle A^{-1}x, x \rangle_0$, and \mathcal{H}_*^* is associated with $\|x\|_{\mathcal{H}_*^*}^2 = \langle B^{-1}x, x \rangle_0$. It remains to note that \mathcal{H}_* is exact interpolation relative to $\overline{\mathcal{H}}$ if and only if \mathcal{H}_*^* is exact interpolation relative to $\overline{\mathcal{H}^*}$, proving the equivalence of (i) and (iii). \square

Combining with Theorem III, one obtains alternative proofs of the interpolation theorems for P' -functions discussed by Donoghue in the paper [13].

Remark 4.4 The exact quadratic interpolation spaces which are fixed by the duality, i.e., which satisfy $\mathcal{H}_*^* = \mathcal{H}_*$, correspond precisely to the class of P' -functions which are self-dual: $h^* = h$. This class was characterized by Hansen in the paper [20].

4.3 Matrix Concavity

A function h on \mathbf{R}_+ is called *matrix concave of order n* if we have Jensen's inequality

$$\lambda h(A_1) + (1 - \lambda)h(A_2) \leq h(\lambda A_1 + (1 - \lambda)A_2)$$

for all positive $n \times n$ matrices A_1, A_2 , and all numbers $\lambda \in [0, 1]$. Let us denote by Γ_n the convex cone of positive concave functions of order n on \mathbf{R}_+ . The fact that $\cap_n \Gamma_n = P'$ follows from the theorem of Kraus [23]. Following [2] we now give an alternative proof of this fact.

Proposition 4.5 *For all n we have the inclusion $C_{3n} \subset \Gamma_n \subset P'_n$. In particular $\cap_1^\infty \Gamma_n = P'$.*

Proof Assume first that $h \in C_{3n}$ and pick two positive matrices A_1 and A_2 . Define $A_3 = (1 - \lambda)A_1 + \lambda A_2$ where $\lambda \in [0, 1]$ is given, and define matrices A and T of order $3n$ by

$$A = \begin{pmatrix} A_3 & 0 & 0 \\ 0 & A_1 & 0 \\ 0 & 0 & A_2 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 0 & 0 \\ \sqrt{1 - \lambda} & 0 & 0 \\ \sqrt{\lambda} & 0 & 0 \end{pmatrix}.$$

It is clear that $T^*T \leq 1$ and

$$T^*AT = \begin{pmatrix} A_3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \leq A,$$

so, since $h \in C_{3n}$, we have $T^*h(A)T \leq h(A)$, or

$$\begin{pmatrix} (1-\lambda)h(A_1) + \lambda h(A_2) & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \leq \begin{pmatrix} h(A_3) & 0 & 0 \\ 0 & h(A_1) & 0 \\ 0 & 0 & h(A_2) \end{pmatrix}.$$

Comparing the matrices in the upper left corners, we find that $h \in \Gamma_n$.

Assume now that $h \in \Gamma_n$, and take positive definite matrices A_1, A_2 of order n with $A_1 \leq A_2$. Also pick $\lambda \in (0, 1)$. Then $\lambda A_2 = \lambda A_1 + (1 - \lambda)A_3$ where $A_3 = \lambda(1 - \lambda)^{-1}(A_2 - A_1)$. By matrix concavity, we then have

$$h(\lambda A_2) \geq \lambda h(A_1) + (1 - \lambda)h(A_3) \geq \lambda h(A_1),$$

where we used non-negativity to deduce the last inequality. Being concave, h is certainly continuous. Letting $\lambda \uparrow 1$ one thus finds that $h(A_1) \leq h(A_2)$. We have shown that $h \in P'_n$. □

For a further discussion of classes of convex matrix functions and their relations to monotonicity, we refer to the paper [21].

4.4 Interpolation Functions of Two Variables

In this section, we briefly discuss a class of interpolation functions of two matrix variables. We shall not completely characterize the class of such generalized interpolation functions here, but we hope that the following discussion will be of some use for a future investigation.

Let H_1 and H_2 be Hilbert spaces. We turn $H_1 \otimes H_2$ into a Hilbert space by defining the inner product on elementary tensors via $\langle x_1 \otimes x_2, x'_1 \otimes x'_2 \rangle := \langle x_1, x'_1 \rangle_1 \cdot \langle x_2, x'_2 \rangle_2$ (then extend via sesqui-linearity). Similarly, if T_i are operators on H_i , the tensor product $T_1 \otimes T_2$ is defined on elementary tensors via $(T_1 \otimes T_2)(x_1 \otimes x_2) = T_1x_1 \otimes T_2x_2$. It is then easy to see that if A_i are positive operators on H_i for $i = 1, 2$, then $A_1 \otimes A_2 \geq 0$ as an operator on the tensor product. Furthermore, we have $A_1 \otimes A_2 \leq A'_1 \otimes A'_2$ if $A_i \leq A'_i$ for $i = 1, 2$.

Given two positive definite matrices A_i of orders n_i and a function h on $\sigma(A_1) \times \sigma(A_2)$, we define a matrix $h(A_1, A_2)$ by

$$h(A_1, A_2) = \sum_{(\lambda_1, \lambda_2) \in \sigma(A_1) \times \sigma(A_2)} h(\lambda_1, \lambda_2) E_{\lambda_1}^1 \otimes E_{\lambda_2}^2$$

where E^j is the spectral resolution of the matrix A_j .

We shall say that h gives rise to exact interpolation relative to (A_1, A_2) , and write $h \in C_{A_1, A_2}$, if the condition

$$T_j^* T_j \leq 1 \quad , \quad T_j^* A_j T_j \leq A_j, \quad j = 1, 2 \tag{4.3}$$

implies

$$\begin{aligned} & h(A_1, A_2) + (T_1 \otimes T_2)^* h(A_1, A_2) (T_1 \otimes T_2) \\ & - (T_1 \otimes 1)^* h(A_1, A_2) (T_1 \otimes 1) - (1 \otimes T_2)^* h(A_1, A_2) (1 \otimes T_2) \geq 0. \end{aligned} \tag{4.4}$$

Taking $T_1 = T_2 = 0$ we see that $h \geq 0$ for all $h \in C_{A_1, A_2}$. It is also clear that C_{A_1, A_2} is a convex cone closed under pointwise convergence on the finite set $\sigma(A_1) \times \sigma(A_2)$.

If $h = h_1 \otimes h_2$ is an elementary tensor where $h_j \in C_{A_j}$ is a function of one variable, then (4.3) implies $T_j^* h_j(A_j) T_j \leq h_j(A_j)$, whence $(h_1(A_1) - T_1^* h_1(A_1) T_1) \otimes (h_2(A_2) - T_2^* h_2(A_2) T_2) \geq 0$, which implies (4.4). We have shown that $C_{A_1} \otimes C_{A_2} \subset C_{A_1, A_2}$.

Since for each $t \geq 0$ the P' -function $\lambda \mapsto \frac{(1+t)\lambda}{1+t\lambda}$ is of class C_{A_j} , we infer that every function representable in the form

$$h(\lambda_1, \lambda_2) = \iint_{[0, \infty]^2} \frac{(1+t_1)\lambda_1}{1+t_1\lambda_1} \frac{(1+t_2)\lambda_2}{1+t_2\lambda_2} d\rho(t_1, t_2) \tag{4.5}$$

with some positive Radon measure ρ on $[0, \infty]^2$ is in the class C_{A_1, A_2} .

We shall say that a function h on $\sigma(A_1) \times \sigma(A_2)$ has the *separate interpolation-property* if for each fixed $b \in \sigma(A_2)$ the function $\lambda_1 \mapsto h(\lambda_1, b)$ is of class C_{A_1} , and a similar statement holds for all functions $\lambda_2 \mapsto h(a, \lambda_2)$.

Lemma 4.6 *Each function of class C_{A_1, A_2} has the separate interpolation-property.*

Proof Let $T_2 = 0$ and take an arbitrary T_1 with $T_1^* T_1 \leq 1$ and $T_1^* A_1 T_1 \leq A_1$. By hypothesis,

$$(T_1 \otimes 1)^* h(A_1, A_2) (T_1 \otimes 1) \leq h(A_1, A_2).$$

Fix an eigenvalue b of A_2 and let y be a corresponding normalized eigenvector. Then for all $x \in H_1$ we have $\langle h(A_1, A_2)x \otimes y, x \otimes y \rangle = \langle h(A_1, b)x, x \rangle_{H_1}$ and $\langle (T_1 \otimes 1)^* h(A_1, A_2)(T_1 \otimes 1)x \otimes y, x \otimes y \rangle = \langle T_1^* h(A_1, b)T_1 x, x \rangle_{H_1}$ so

$$\langle T_1^* h(A_1, b)T_1 x, x \rangle_{H_1} \leq \langle h(A_1, b)x, x \rangle_{H_1}.$$

The functions $h(a, \lambda_2)$ can be treated similarly. \square

Example The function $h(\lambda_1, \lambda_2) = (\lambda_1 + \lambda_2)^{1/2}$ clearly has the separate interpolation-property for all A_1, A_2 . However, it is not representable in the form (4.5). Indeed, $\operatorname{Re}\{h(i, i) - h(-i, i)\} = 1$ while it is easy to check that $\operatorname{Re}\{h(\lambda_1, \lambda_2) - h(\bar{\lambda}_1, \lambda_2)\} \leq 0$ whenever $\operatorname{Im} \lambda_1, \operatorname{Im} \lambda_2 > 0$ and h is of the form (4.5).

Let us say that a function $h(\lambda_1, \lambda_2)$ defined on $\mathbf{R}_+ \times \mathbf{R}_+$ is an interpolation function (of two variables) if $h \in C_{A_1, A_2}$ for all A_1, A_2 . Lemma 4.6 implies that interpolation functions are separately real-analytic in $\mathbf{R}_+ \times \mathbf{R}_+$ and that the functions $h(a, \cdot)$ and $h(\cdot, b)$ are of class P' (cf. Theorem III).

The above notion of interpolation function is close to Korányi's definition of monotone matrix function of two variables: $f(\lambda_1, \lambda_2)$ is *matrix monotone* in a rectangle $I = I_1 \times I_2$ (I_1, I_2 intervals in \mathbf{R}) if $A_1 \leq A'_1$ (with spectra in I_1) and $A_2 \leq A'_2$ (with spectra in I_2) implies

$$f(A'_1, A'_2) - f(A'_1, A_2) - f(A_1, A'_2) + f(A_1, A_2) \geq 0.$$

Lemma 4.7 *Each interpolation function is matrix monotone in $\mathbf{R}_+ \times \mathbf{R}_+$.*

Proof Let $0 < A_i \leq A'_i$ and put $\tilde{A}_i = \begin{pmatrix} A'_i & 0 \\ 0 & A_i \end{pmatrix}$, $T_i = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. Since $T_i^* \tilde{A}_i T_i \leq \tilde{A}_i$, an interpolation function h will satisfy the interpolation inequality (4.4) with A_i replaced by \tilde{A}_i . Applying this inequality to vectors of the form $\begin{pmatrix} x_1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} x_2 \\ 0 \end{pmatrix}$ we readily obtain

$$\begin{aligned} & \langle h(A'_1, A'_2)x_1 \otimes x_2, x_1 \otimes x_2 \rangle - \langle h(A_1, A'_2)x_1 \otimes x_2, x_1 \otimes x_2 \rangle \\ & - \langle h(A'_1, A_2)x_1 \otimes x_2, x_1 \otimes x_2 \rangle + \langle h(A_1, A_2)x_1 \otimes x_2, x_1 \otimes x_2 \rangle \geq 0. \end{aligned}$$

The same result obtains with $x_1 \otimes x_2$ replaced by a sum $x_1 \otimes x_2 + x'_1 \otimes x'_2 + \dots$, i.e., h is matrix monotone. \square

Remark 4.8 Assume that f is of the form $f(\lambda_1, \lambda_2) = g_1(\lambda_1) + g_2(\lambda_2)$. Then f is matrix monotone for all g_1, g_2 and f is an interpolation function if and only if $g_1, g_2 \in P'$. In order to disregard "trivial" monotone functions of the above type, Korányi [22] imposed the normalizing assumption (a) $f(\lambda_1, 0) = f(0, \lambda_2) = 0$ for all λ_1, λ_2 .

It follows from Lemma 4.7 and the proof of [22, Theorem 4] that if h is a C^2 -smooth interpolation function, then the function

$$k(x_1, x_2; y_1, y_2) = \frac{h(x_1, x_2) - h(x_1, y_2) - h(y_1, x_2) + h(y_1, y_2)}{(x_1 - y_1)(x_2 - y_2)}$$

is *positive definite* in the sense that $\sum_m \sum_n k(x_m, y_m; x_n, y_n) \alpha_m \bar{\alpha}_n \geq 0$ for all finite sequences of positive numbers x_j, y_k and all complex numbers α_l . (The proof uses Löwner’s matrix.) Korányi uses essentially this positive definiteness condition (and condition (a) in the remark above) to deduce an integral representation formula for h as an integral of products of Pick functions. See Theorem 3 in [22]. However, in contrast to our situation, Korányi considers functions monotone on the rectangle $(-1, 1) \times (-1, 1)$, so this last result cannot be immediately applied. (It easily implies local representation formulas, valid in finite rectangles, but these representations do not appear to be very natural from our point of view.)

This is not the right place to attempt to extend Korányi’s methods to functions on $\mathbf{R}_+ \times \mathbf{R}_+$; it would seem more appropriate to give a more direct characterization of the classes C_{A_1, A_2} or of the class of interpolation functions. At present, we do not know if there is an interpolation function which is not representable in the form (4.5).

5 Proof of the K -Property

In this section we extend the result of Theorem 2.4 to obtain the full proof of Theorem I. The discussion is in principle not hard, but it does require some care to keep track of both norms when reducing to a finite-dimensional case.

Recall first that, by Lemma 2.3, it suffices to consider the diagonal case $\overline{\mathcal{H}} = \overline{\mathcal{K}}$.

To prove Theorem I we fix a regular Hilbert couple $\overline{\mathcal{H}}$; we must prove that it has the K -property (see Sect. 1.5). By Theorem 2.4, we know that this is true if $\overline{\mathcal{H}}$ is finite dimensional and the associated operator only has eigenvalues of unit multiplicity.

We shall use a weak* type compactness result ([2]). To formulate it, let $\mathcal{L}_1(\overline{\mathcal{H}})$ be the unit ball in the space $\mathcal{L}(\overline{\mathcal{H}})$. Moreover, let Σ_t be the sum $\mathcal{H}_0 + \mathcal{H}_1$ normed by $\|x\|_{\Sigma_t}^2 := K(t, x)$. Note that $\|\cdot\|_{\Sigma_t}$ is an equivalent norm on Σ and that $\Sigma_1 = \Sigma$ isometrically. We denote by $\mathcal{L}_1(\Sigma_t)$ the unit ball in the space $\mathcal{L}(\Sigma_t)$.

In view of Remark 1.4, one has the identity

$$\mathcal{L}_1(\overline{\mathcal{H}}) = \bigcap_{t \in \mathbf{R}_+} \mathcal{L}_1(\Sigma_t). \tag{5.1}$$

We shall use this to define a compact topology on $\mathcal{L}_1(\overline{\mathcal{H}})$.

Lemma 5.1 *The subset $\mathcal{L}_1(\overline{\mathcal{H}}) \subset \mathcal{L}_1(\Sigma)$ is compact relative to the weak operator topology inherited from $\mathcal{L}(\Sigma)$.*

Recall that the *weak operator topology* on $\mathcal{L}(H)$ is the weakest topology such that a net T_i converges to the limit T if the inner product $\langle T_i x, y \rangle_H$ converges to $\langle T x, y \rangle_H$ for all $x, y \in H$.

Proof of Lemma 5.1 The weak operator topology coincides on the unit ball $\mathcal{L}_1(\Sigma)$ with the weak*-topology, which is compact, due to Alaoglu’s theorem (see [29], Chap. 4 for details). It is clear that for a fixed $t > 0$, the subset $\mathcal{L}_1(\Sigma) \cap \mathcal{L}_1(\Sigma_t)$ is weak operator closed in $\mathcal{L}_1(\Sigma)$; hence it is also compact. In view of (5.1), the set $\mathcal{L}_1(\overline{\mathcal{H}})$ is an intersection of compact sets. Hence the set $\mathcal{L}_1(\overline{\mathcal{H}})$ is itself compact, provided that we endow it with the subspace topology inherited from $\mathcal{L}_1(\Sigma)$. \square

Denote by P_n the projections $P_n = E_{\sigma(A) \cap [n^{-1}, n]}$ on \mathcal{H}_0 where E is the spectral resolution of A and $n = 1, 2, 3, \dots$. Consider the couple

$$\overline{\mathcal{H}^{(n)}} = (P_n(\mathcal{H}_0), P_n(\mathcal{H}_1)),$$

the associated operator of which is the compression A_n of A to the subspace $P_n(\mathcal{H}_0)$. Note that the norms in the couple $\overline{\mathcal{H}^{(n)}}$ are equivalent, i.e., the associated operator A_n is bounded above and below.

We shall need two lemmas.

Lemma 5.2 *If $\overline{\mathcal{H}^{(n)}}$ has the K -property for all n , then so does $\overline{\mathcal{H}}$.*

Proof Note that $\|P_n\|_{\mathcal{L}(\overline{\mathcal{H}})} = 1$ for all n , and that $P_n \rightarrow 1$ as $n \rightarrow \infty$ relative to the strong operator topology on $\mathcal{L}(\Sigma)$. Suppose that $x^0, y^0 \in \Sigma$ are elements such that, for some $\rho > 1$,

$$K(t, y^0) < \frac{1}{\rho} K(t, x^0), \quad t > 0. \tag{5.2}$$

Then $K(t, P_n y^0) \leq K(t, y^0) < \rho^{-1} K(t, x^0)$. Moreover, the identity $K(t, P_n y^0) = \left\langle \frac{t A_n}{1+t A_n} P_n y^0, P_n y^0 \right\rangle_0$ shows that we have an estimate of the form $K(t, P_n y^0) \leq C_n \min\{1, t\}$ for $t > 0$ and large enough C_n (this follows since A_n is bounded above and below).

The functions $K(t, P_m x^0)$ increase monotonically, converging uniformly on compact subsets of \mathbf{R}_+ to $K(t, x^0)$ when $m \rightarrow \infty$. By concavity of the function $t \mapsto K(t, P_m x^0)$ we will then have

$$K(t, P_n y^0) < \frac{1}{\tilde{\rho}} K(t, P_m x^0), \quad t \in \mathbf{R}_+, \tag{5.3}$$

provided that m is sufficiently large, where $\tilde{\rho}$ is any number in the interval $1 < \tilde{\rho} < \rho$.

Indeed, let $A = \lim_{t \rightarrow \infty} K(t, P_n y^0)$ and $B = \lim_{t \rightarrow 0} K(t, P_n y^0)/t$. Take points $t_0 < t_1$ such that $K(t, P_n y^0) \geq A/\rho'$ when $t \geq t_1$ and $K(t, P_n y^0)/t \leq B\rho'$ when $t \leq t_0$. Here ρ' is some number in the interval $1 < \rho' < \rho$.

Next use (5.2) to choose m large enough that $K(t, P_m x^0) > \rho K(t, P_n y^0)$ for all $t \in [t_0, t_1]$. Then $K(t, P_m x^0) > (\rho/\rho')K(t, P_n y^0)$ for $t = t_1$, hence for all $t \geq t_1$, and $K(t, P_m x^0)/t > (\rho/\rho')K(t, P_n y^0)/t$ for $t = t_0$ and hence also when $t \leq t_0$. Choosing $\rho' = \rho/\tilde{\rho}$ now establishes (5.3).

Put $N = \max\{m, n\}$. If $\overline{\mathcal{H}}^{(N)}$ has the K -property, we can find a map $T_{nm} \in \mathcal{L}_1(\overline{\mathcal{H}})$ such that $T_{nm} P_m x^0 = P_n y^0$. (Define $T_{nm} = 0$ on the orthogonal complement of $P_N(\mathcal{H}_0)$ in Σ .) In view of Lemma 5.1, the maps T_{nm} must cluster at some point $T \in \mathcal{L}_1(\overline{\mathcal{H}})$. It is clear that $T x^0 = y^0$. Since $\rho > 1$ was arbitrary, we have shown that $\overline{\mathcal{H}}$ has the K -property. \square

Lemma 5.3 *Given $x^0, y^0 \in \mathcal{H}_0^{(n)}$ and a number $\epsilon > 0$ there exists a positive integer n and a finite-dimensional couple $\overline{\mathcal{V}} \subset \overline{\mathcal{H}}^{(n)}$ such that $x^0, y^0 \in \mathcal{V}_0 + \mathcal{V}_1$ and*

$$(1 - \epsilon)K(t, x; \overline{\mathcal{H}}) \leq K(t, x; \overline{\mathcal{V}}) \leq (1 + \epsilon)K(t, x; \overline{\mathcal{H}}), \quad t > 0, x \in \mathcal{V}_0 + \mathcal{V}_1. \tag{5.4}$$

Moreover, $\overline{\mathcal{V}}$ can be chosen so that all eigenvalues of the associated operator $A_{\overline{\mathcal{V}}}$ are of unit multiplicity.

Proof Let A_n be the operator associated with the couple $\overline{\mathcal{H}}^{(n)}$; thus $1/n \leq A_n \leq n$.

Take $\eta > 0$ and let $\{\lambda_i\}_1^N$ be a finite subset of $\sigma(A_n)$ such that $\sigma(A_n) \subset \cup_1^N E_i$ where $E_i = (\lambda_i - \eta/2, \lambda_i + \eta/2)$. We define a Borel function $w : \sigma(A_n) \rightarrow \sigma(A_n)$ by $w(\lambda) = \lambda_i$ on $E_i \cap \sigma(A_n)$, then $\|w(A_n) - A_n\|_{\mathcal{L}(\mathcal{H}_0)} \leq \eta$.

Let $k_t(\lambda) = \frac{t\lambda}{1+t\lambda}$. It is easy to check that the Lipschitz constant of the restriction $k_t \mid \sigma(A_n)$ is bounded above by $C_1 \min\{1, t\}$ where $C_1 = C_1(n)$ is independent of t . Hence

$$\|k_t(w(A_n)) - k_t(A_n)\|_{\mathcal{L}(\mathcal{H}_0)} \leq C_1 \eta \min\{1, t\}.$$

It follows readily that

$$|\langle (k_t(w(A_n)) - k_t(A_n))x, x \rangle_0| \leq C_1 \eta \min\{1, t\} \|x\|_0^2, \quad x \in P_n(\mathcal{H}_0).$$

Now let $c > 0$ be such that $A \geq c$. The elementary inequality $k_t(c) \geq (1/2) \min\{1, ct\}$ shows that

$$\langle k_t(A_n)x, x \rangle_0 \geq C_2 \min\{1, t\} \|x\|_0^2, \quad x \in P_n(\mathcal{H}_0),$$

where $C_2 = (1/2) \min\{1, c\}$. Combining these estimates, we deduce that

$$|\langle (k_t(w(A_n))x, x) - \langle k_t(A_n)x, x \rangle_0| \leq C_3 \eta \langle k_t(A_n)x, x \rangle_0, \quad x \in P_n(\mathcal{H}_0) \tag{5.5}$$

for some suitable constant $C_3 = C_3(n)$.

Now pick unit vectors e_i, f_i supported by the spectral sets $E_i \cap \sigma(A_n)$ such that x^0 and y^0 belong to the space \mathcal{W} spanned by $\{e_i, f_i\}_1^N$. Put $\mathcal{W}_0 = \mathcal{W}_1 = \mathcal{W}$ and define norms on those spaces by

$$\|x\|_{\mathcal{W}_0} = \|x\|_{\mathcal{H}_0} \quad , \quad \|x\|_{\mathcal{W}_1}^2 = \langle w(A)x, x \rangle_{\mathcal{H}_0} .$$

The operator associated with $\overline{\mathcal{W}}$ is then the compression of $w(A_n)$ to \mathcal{W}_0 , i.e.,

$$\|x\|_{\mathcal{W}_1}^2 = \langle A_{\overline{\mathcal{W}}}x, x \rangle_{\mathcal{W}_0} = \langle w(A_n)x, x \rangle_{\mathcal{H}_0} , \quad x \in \mathcal{W} .$$

Let $\epsilon = 2C_3\eta$ and observe that, by (5.5)

$$|K(t, x; \overline{\mathcal{W}}) - K(t, x; \overline{\mathcal{H}})| \leq (\epsilon/2)K(t, x; \overline{\mathcal{H}}), \quad f \in \mathcal{W}. \quad (5.6)$$

The eigenvalues of $A_{\overline{\mathcal{W}}}$ typically have multiplicity 2. To obtain unit multiplicity, we perturb $A_{\overline{\mathcal{W}}}$ slightly to a positive matrix $A_{\overline{\mathcal{V}}}$ such that $\|A_{\overline{\mathcal{W}}} - A_{\overline{\mathcal{V}}}\|_{\mathcal{L}(\mathcal{H}_0)} < \epsilon/2C_3$. Let $\overline{\mathcal{V}}$ be the couple associated with $A_{\overline{\mathcal{V}}}$, i.e., put $\mathcal{V}_i = \mathcal{W}$ for $i = 0, 1$ and

$$\|x\|_{\mathcal{V}_0} = \|x\|_{\mathcal{W}_0} \quad \text{and} \quad \|x\|_{\mathcal{V}_1}^2 = \langle A_{\overline{\mathcal{V}}}x, x \rangle_{\mathcal{V}_0} .$$

It is then straightforward to check that

$$|K(t, f; \overline{\mathcal{W}}) - K(t, f; \overline{\mathcal{V}})| \leq (\epsilon/2)K(t, f; \overline{\mathcal{H}}), \quad f \in \mathcal{W} .$$

Combining this with the estimate (5.6), one finishes the proof of the lemma. \square

Proof of Theorem I Given two elements $x^0, y^0 \in \Sigma$ as in (5.2) we write $x^n = P_n(x^0)$ and $y^n = P_n(y^0)$. By the proof of Lemma 5.2 we then have $K(t, y^n) \leq \tilde{\rho}^{-1}K(t, x^n)$ for large enough n , where $\tilde{\rho}$ is any given number in the interval $(1, \rho)$.

We then use Lemma 5.3 to choose a finite-dimensional sub-couple $\overline{\mathcal{V}} \subset \overline{\mathcal{H}}^{(n)}$ such that

$$\begin{aligned} K(t, y^n; \overline{\mathcal{V}}) &\leq (1 + \epsilon)K(t, y^n; \overline{\mathcal{H}}) \\ &< \tilde{\rho}^{-1}K(t, x^n; \overline{\mathcal{V}}) + \epsilon(K(t, x^n; \overline{\mathcal{H}}) + K(t, y^n; \overline{\mathcal{H}})) . \end{aligned}$$

Here $\epsilon > 0$ is at our disposal.

Choosing ϵ sufficiently small, we can arrange that

$$K(t, y^n; \overline{\mathcal{V}}) \leq K(t, x^n; \overline{\mathcal{V}}), \quad t > 0. \quad (5.7)$$

By Theorem 2.4, the condition (5.7) implies the existence of an operator $T' \in \mathcal{L}_1(\overline{\mathcal{V}})$ such that $T'x^n = y^n$. Considering the canonical inclusion and projection

$$I : \Sigma(\mathcal{V}) \rightarrow \Sigma(\mathcal{H}) \quad \text{and} \quad \Pi : \Sigma(\mathcal{H}) \rightarrow \Sigma(\mathcal{V}),$$

we have, by virtue of Lemma 5.3,

$$\| I \|_{\mathcal{L}(\overline{\mathcal{V}}; \overline{\mathcal{H}})}^2 \leq (1 - \epsilon)^{-1} \quad \text{and} \quad \| \Pi \|_{\mathcal{L}(\overline{\mathcal{H}}; \overline{\mathcal{V}})}^2 \leq 1 + \epsilon.$$

Now let $T = T_\epsilon := IT'\Pi \in \mathcal{L}(\overline{\mathcal{H}^{(n)}})$. Then $\| T \|^2 \leq \frac{1+\epsilon}{1-\epsilon}$ and $Tx^n = y^n$. As $\epsilon \downarrow 0$ the operators T_ϵ will cluster at some point $T \in \mathcal{L}_1(\overline{\mathcal{H}^{(n)}})$ such that $Tx^n = y^n$ (cf. Lemma 5.1).

We have shown that $\overline{\mathcal{H}^{(n)}}$ has the K -property. In view of Lemma 5.2, this implies that $\overline{\mathcal{H}}$ has the same property. The proof of Theorem I is therefore complete. \square

6 Representations of Interpolation Functions

6.1 Quadratic Interpolation Methods

Let us say that an interpolation method defined on regular Hilbert couples taking values in Hilbert spaces is a *quadratic interpolation method*. (Donoghue [14] used the same phrase in a somewhat wider sense, allowing the methods to be defined on non-regular Hilbert couples as well.)

If F is an exact quadratic interpolation method, and $\overline{\mathcal{H}}$ a Hilbert couple, then by Donoghue's theorem III there exists a positive Radon measure ϱ on $[0, \infty]$ such that $F(\overline{\mathcal{H}}) = \mathcal{H}_\varrho$, where the latter space is defined by the familiar norm $\|x\|_\varrho^2 = \int_{[0, \infty]} (1 + t^{-1}) K(t, x) d\varrho(t)$.

A priori, the measure ϱ could depend not only on F but also on the particular $\overline{\mathcal{H}}$. That ϱ is independent of $\overline{\mathcal{H}}$ can be realized in the following way. Let $\overline{\mathcal{H}'}$ be a regular Hilbert couple such that every positive rational number is an eigenvalue of the associated operator. Let B' be the operator associated with the exact quadratic interpolation space $F(\overline{\mathcal{H}'})$. There is then clearly a unique P' -function h on $\sigma(A')$ such that $B' = h(A')$, viz. there is a unique positive Radon measure ϱ on $[0, \infty]$ such that $F(\overline{\mathcal{H}'}) = \mathcal{H}'_\varrho$ (see Sect. 1.2 for the notation).

If $\overline{\mathcal{H}}$ is any regular Hilbert couple, we can form the direct sum $\overline{\mathcal{S}} = \overline{\mathcal{H}'} \oplus \overline{\mathcal{H}}$. Denote by \tilde{A} the corresponding operator and let $\tilde{B} = \tilde{h}(\tilde{A})$ be the operator corresponding to the exact quadratic interpolation space $F(\overline{\mathcal{S}})$. Then $\tilde{h}(\tilde{A}) = \tilde{h}(A') \oplus \tilde{h}(A) = h(A') \oplus \tilde{h}(A)$. This means that $\tilde{h}(A') = h(A')$, i.e., $\tilde{h} = h$. In particular, the operator B corresponding to the exact interpolation space $F(\overline{\mathcal{H}})$ is equal to $h(A)$. We have shown that $F(\overline{\mathcal{H}}) = \mathcal{H}_\varrho$. We emphasize our conclusion with the following theorem.

Theorem 6.1 *There is a one-to-one correspondence $\varrho \mapsto F$ between positive Radon measures and exact quadratic interpolation methods.*

We will shortly see that Theorem 6.1 is equivalent to the theorem of Foiaş and Lions [16]. As we remarked above, a more general version of the theorem, admitting for non-regular Hilbert couples, is found in Donoghue’s paper [14].

6.2 Interpolation Type and Reiteration

In this subsection, we prove some general facts concerning quadratic interpolation methods; we shall mostly follow Fan [15].

Fix a function $h \in P'$ of the form

$$h(\lambda) = \int_{[0, \infty[} \frac{(1+t)\lambda}{1+t\lambda} dQ(t).$$

It will be convenient to write $\overline{\mathcal{H}}_h$ for the corresponding exact interpolation space \mathcal{H}_Q . Thus, we shall denote

$$\|x\|_h^2 = \langle h(A)x, x \rangle_0 = \int_{[0, \infty[} (1+t^{-1}) K(t, x) dQ(t).$$

More generally, we shall use the same notation when h is any quasi-concave function on \mathbf{R}_+ , then $\overline{\mathcal{H}}_h$ is a quadratic interpolation space, but not necessarily exact.

Recall that, given a function \mathbf{H} of one variable, we say that \mathcal{H}_* is of type \mathbf{H} with respect to $\overline{\mathcal{H}}$ if $\|T\|_{\mathcal{L}(\mathcal{H}_i)}^2 \leq M_i$ implies $\|T\|_{\mathcal{L}(\mathcal{H}_*)}^2 \leq M_0 \mathbf{H}(M_1/M_0)$.

We shall say that a quasi-concave function h on \mathbf{R}_+ is of type \mathbf{H} if $\overline{\mathcal{H}}_h$ is of type \mathbf{H} relative to any regular Hilbert couple $\overline{\mathcal{H}}$. The following result somewhat generalizes Theorem 3.2. The class of functions of type \mathbf{H} clearly forms a convex cone.

Theorem 6.2 *Let h be of type \mathbf{H} , where (i) $\mathbf{H}(1) = 1$ and $\mathbf{H}(t) \leq \max\{1, t\}$, and (ii) \mathbf{H} has left and right derivatives $\theta_{\pm} = H'(1 \pm)$ at the point 1, where $\theta_- \leq \theta_+$. Then for any positive constant c ,*

$$\min \{ \lambda^{\theta_-}, \lambda^{\theta_+} \} \leq \frac{h(c\lambda)}{h(c)} \leq \max \{ \lambda^{\theta_-}, \lambda^{\theta_+} \}, \quad \lambda \in \mathbf{R}_+. \tag{6.1}$$

In particular, if $\mathbf{H}(t)$ is differentiable at $t = 1$ and $\mathbf{H}'(1) = \theta$, then $h(\lambda) = \lambda^\theta$, $\lambda \in \mathbf{R}_+$.

Proof Replacing A by cA , it is easy to see that if h is of type \mathbf{H} , then so is $h_c(t) = h(ct)/h(c)$. Fix $\mu > 0$ and consider the function $h_0(t) = h_c(\mu t)/h_c(\mu)$. By Theorem II, we have $h_0(t) \leq \mathbf{H}(t)$ for all t . Furthermore $h_0(1) = \mathbf{H}(1) = 1$ by (i). Since h_0 is differentiable, the assumption (ii) now gives $\theta_- \leq h'_0(1) \leq \theta_+$, or

$$\theta_- \leq \frac{\mu h'_c(\mu)}{h_c(\mu)} \leq \theta_+.$$

Dividing through by μ and integrating over the interval $[1, \lambda]$, one now verifies the inequalities in (6.1). \square

The following result provides a partial converse to Theorem II.

Theorem 6.3 ([15]) *Let $h \in P'$ and set $\mathbf{H}(t) = \sup_{s>0} h(st)/h(s)$. Then h is of type \mathbf{H} .*

Proof Let $T \in \mathcal{L}(\overline{\mathcal{H}})$ be a non-zero operator; put $M_j = \|T\|_{\mathcal{L}(\mathcal{H}_j)}^2$ and $M = M_1/M_0$. We then have (by Lemma 1.1)

$$\begin{aligned} \|Tx\|_h^2 &= \int_{[0,\infty]} (1+t^{-1}) K(t, Tx) d\varrho(t) \\ &\leq M_0 \int_{[0,\infty]} (1+t^{-1}) K(tM, x) d\varrho(t) \\ &= M_0 \int_{[0,\infty]} \left\langle \frac{(1+t)MA}{1+tMA} x, x \right\rangle_0 d\varrho(t) \\ &= M_0 \langle h(MA)x, x \rangle_0. \end{aligned}$$

Letting E be the spectral resolution of A , we have

$$\langle h(MA)x, x \rangle_0 = \int_0^\infty h(M\lambda) d \langle E_\lambda x, x \rangle_0.$$

Since $h(M\lambda)/h(\lambda) \leq \mathbf{H}(M)$, we conclude that

$$\|Tx\|_h^2 \leq M_0 \mathbf{H}(M) \int_0^\infty h(\lambda) d \langle E_\lambda x, x \rangle_0 = M_0 \mathbf{H}(M) \|x\|_h^2,$$

which finishes the proof. \square

Given a function h of a positive variable, we define a new function \tilde{h} by

$$\tilde{h}(s, t) = s h(t/s).$$

The following reiteration theorem is due to Fan.

Theorem 6.4 ([15]) *Let $h, h_0, h_1 \in P'$, and $\varphi(\lambda) = \tilde{h}(h_0(\lambda), h_1(\lambda))$. Then $\overline{\mathcal{H}}_\varphi = (\overline{\mathcal{H}}_{h_0}, \overline{\mathcal{H}}_{h_1})_h$ with equal norms. Moreover, $\overline{\mathcal{H}}_\varphi$ is an exact interpolation space relative to $\overline{\mathcal{H}}$.*

Proof Let $\overline{\mathcal{H}}'$ denote the couple $(\overline{\mathcal{H}}_{h_0}, \overline{\mathcal{H}}_{h_1})$. The corresponding operator A' then obeys

$$\|x\|_{\overline{\mathcal{H}}_{h_1}} = \|(A')^{1/2}x\|_{\mathcal{H}'_0} = \|\varphi_0(A)^{1/2}(A')^{1/2}x\|_0, \quad x \in \Delta(\overline{\mathcal{H}}').$$

On the other hand, $\|x\|_{\overline{\mathcal{H}}_{h_1}} = \|\varphi_1(A)^{1/2}x\|_0$, so

$$(A')^{1/2}x = \varphi_0(A)^{-1/2}\varphi_1(A)^{1/2}x, \quad x \in \Delta(\overline{\mathcal{H}}^t).$$

We have shown that $A' = \varphi_0(A)^{-1}\varphi_1(A)$, whence (by Lemma 1.1)

$$\begin{aligned} K(t, x; \overline{\mathcal{H}}^t) &= \left\langle \frac{t\varphi_0(A)^{-1}\varphi_1(A)}{1+t\varphi_0(A)^{-1}\varphi_1(A)}x, x \right\rangle_{\mathcal{H}'_0} \\ &= \left\langle \frac{t\varphi_1(A)}{1+t\varphi_0(A)^{-1}\varphi_1(A)}x, x \right\rangle_{\mathcal{H}'_0}. \end{aligned} \tag{6.2}$$

Now let the function $h \in P'$ be given by

$$h(\lambda) = \int_{[0, \infty]} \frac{(1+t)\lambda}{1+t\lambda} d\varrho(t),$$

and note that the function $\varphi = \tilde{h}(h_0, h_1)$ is given by

$$\varphi(\lambda) = \int_{[0, \infty]} \frac{(1+t)h_1(\lambda)}{1+th_1(\lambda)/h_0(\lambda)} d\varrho(t).$$

Combining with (6.2), we find that

$$\begin{aligned} \|x\|_{\overline{\mathcal{H}}_h}^2 &= \int_{[0, \infty]} (1+t^{-1}) K(t, x; \overline{\mathcal{H}}^t) d\varrho(t) \\ &= \int_0^\infty \left[\int_{[0, \infty]} \frac{(1+t)h_1(\lambda)}{1+th_1(\lambda)/h_0(\lambda)} d\varrho(t) \right] d\langle E_\lambda x, x \rangle_0 = \|x\|_{\overline{\mathcal{H}}_\varphi}^2. \end{aligned}$$

This finishes the proof of the theorem. □

Combining with Donoghue’s theorem III, one obtains the following, purely function-theoretic corollary. Curiously, we are not aware of a proof which does not use interpolation theory.

Corollary 6.5 ([15]) *Suppose that $h \in P'$ and that $h_0, h_1 \in P'|F$, where F is some closed subset of \mathbf{R}_+ . Then the function $\varphi = \tilde{h}(h_0, h_1)$ is also of class $P'|F$.*

6.3 Donoghue’s Representation

Let $\overline{\mathcal{H}}$ be a regular Hilbert couple. In Donoghue’s setting, the principal object is the space $\Delta = \mathcal{H}_0 \cap \mathcal{H}_1$ normed by $\|x\|_\Delta^2 = \|x\|_0^2 + \|x\|_1^2$. In the following, all involutions are understood to be taken with respect to the norm of Δ .

We express the norms in the spaces \mathcal{H}_i as

$$\|x\|_0^2 = \langle Hx, x \rangle_\Delta \quad \text{and} \quad \|x\|_1^2 = \langle (1 - H)x, x \rangle_\Delta,$$

where H is a bounded positive operator on Δ , $0 \leq H \leq 1$. The regularity of $\overline{\mathcal{H}}$ means that neither 0 nor 1 is an eigenvalue of H .

To an arbitrary quadratic intermediate space \mathcal{H}_* there corresponds a bounded positive injective operator K on Δ such that

$$\|x\|_*^2 = \langle Kx, x \rangle_\Delta.$$

It is then easy to see that \mathcal{H}_* is exact interpolation if and only if, for bounded operators T on Δ , the conditions $T^*HT \leq H$ and $T^*(1 - H)T \leq 1 - H$ imply $T^*KT \leq K$. It is straightforward to check that the relations between H, K and the operators A, B used in the previous sections are

$$H = \frac{1}{1 + A}, \quad A = \frac{1 - H}{H}, \quad K = \frac{B}{1 + A}, \quad B = \frac{K}{H}. \tag{6.3}$$

(It follows from the proof of Lemma 1.2 that H and K commute.)

By Theorem III we know that \mathcal{H}_* is an exact interpolation space if and only if $B = h(A)$ for some $h \in P'$. By (6.3), this is equivalent to that $K = k(H)$ where

$$k(H) = \frac{h(A)}{1 + A} = H h\left(\frac{1 - H}{H}\right).$$

In its turn, this means that

$$\begin{aligned} k(\lambda) &= \lambda \int_{[0, \infty]} \frac{(1 + t)(1 - \lambda)/\lambda}{1 + t(1 - \lambda)/\lambda} d\varrho(t) \\ &= \int_{[0, \infty]} \frac{(1 + t)\lambda(1 - \lambda)}{\lambda + t(1 - \lambda)} d\varrho(t), \quad \lambda \in \sigma(H), \end{aligned}$$

where ϱ is a suitable Radon measure. Applying the change of variables $s = 1/(1 + t)$ and defining a positive Radon measure μ on $[0, 1]$ by $d\mu(s) = d\varrho(t)$, we arrive at the expression

$$k(\lambda) = \int_0^1 \frac{\lambda(1 - \lambda)}{(1 - s)(1 - \lambda) + s\lambda} d\mu(s), \quad \lambda \in \sigma(H), \tag{6.4}$$

which gives the representation exact quadratic interpolation spaces originally used by Donoghue in [14].

6.4 *J-Methods and the Foias-Lions Theorem*

We define the (quadratic) J -functional relative to a regular Hilbert couple $\overline{\mathcal{H}}$ by

$$J(t, x) = J(t, x; \overline{\mathcal{H}}) = \|x\|_0^2 + t \|x\|_1^2, \quad t > 0, x \in \Delta(\overline{\mathcal{H}}).$$

Note that $J(t, x)^{1/2}$ is an equivalent norm on Δ and that $J(1, x) = \|x\|_\Delta^2$.

Given a positive Radon measure ν on $[0, \infty]$, we define a Hilbert space $J_\nu(\overline{\mathcal{H}})$ as the set of all elements $x \in \Sigma(\overline{\mathcal{H}})$ such that there exists a measurable function $u : [0, \infty] \rightarrow \Delta$ such that

$$x = \int_{[0, \infty]} u(t) d\nu(t) \quad (\text{convergence in } \Sigma) \quad (6.5)$$

and

$$\int_{[0, \infty]} \frac{J(t, u(t))}{1+t} d\nu(t) < \infty. \quad (6.6)$$

The norm in the space $J_\nu(\overline{\mathcal{H}})$ is defined by

$$\|x\|_{J_\nu}^2 = \inf_u \int_{[0, \infty]} \frac{J(t, u(t))}{1+t} d\nu(t) \quad (6.7)$$

over all u satisfying (6.5) and (6.6).

The space (6.7) was (with different notation) introduced by Foias and Lions in the paper [16], where it was shown that there is a unique minimizer $u(t)$ of the problem (6.7), namely

$$u(t) = \varphi_t(A)x \quad \text{where} \quad \varphi_t(\lambda) = \frac{1+t}{1+t\lambda} \left(\int_{[0, \infty]} \frac{1+s}{1+s\lambda} d\nu(s) \right)^{-1}. \quad (6.8)$$

Inserting this expression for u into (6.7), one finds that

$$\|x\|_{J_\nu}^2 = \langle h(A)x, x \rangle_0$$

where

$$h(\lambda)^{-1} = \int_{[0, \infty]} \frac{1+t}{1+t\lambda} d\nu(t). \quad (6.9)$$

It is easy to verify that the class of functions representable in the form (6.9) for some positive Radon measure ν coincides with the class P' . We have thus arrived at the following result.

Theorem 6.6 *Every exact quadratic interpolation space \mathcal{H}_* can be represented isometrically in the form $\mathcal{H}_* = J_\nu(\overline{\mathcal{H}})$ for some positive Radon measure ν on $[0, \infty]$. Conversely, any space of this form is an exact quadratic interpolation space.*

In the original paper [16], Foias and Lions proved the less precise statement that each exact quadratic interpolation method F can be represented as $F = J_\nu$ for some positive Radon measure ν .

6.5 The Relation Between the K - and J -Representations

The assignment $K_\varrho = J_\nu$ gives rise to a non-trivial bijection $\varrho \mapsto \nu$ of the set of positive Radon measures on $[0, \infty]$. In this bijection, ϱ and ν are in correspondence if and only if

$$\int_{[0, \infty]} \frac{(1+t)\lambda}{1+t\lambda} d\varrho(t) = \left(\int_{[0, \infty]} \frac{1+t}{1+t\lambda} d\nu(t) \right)^{-1}.$$

As an example, let us consider the geometric interpolation space (where $c_\theta = \pi / \sin(\pi\theta)$)

$$\|x\|_\theta^2 = \langle A^\theta x, x \rangle_0 = c_\theta \int_0^\infty t^{-\theta} K(t, x) \frac{dt}{t}.$$

The measure ϱ corresponding to this method is $d\varrho_\theta(t) = \frac{c_\theta t^{-\theta}}{1+t} dt$. On the other hand, it is easy to check that

$$\lambda^\theta = \left(\int_0^\infty \frac{1+t}{1+t\lambda} d\nu_\theta(t) \right)^{-1} \quad \text{where} \quad d\nu_\theta(t) = \frac{c_\theta t^\theta}{1+t} \frac{dt}{t}.$$

We leave it to the reader to check that the norm in \mathcal{H}_θ is the infimum of the expression

$$c_\theta \int_0^\infty t^\theta J(t, u(t)) \frac{dt}{t}$$

over all representations

$$x = \int_0^\infty u(t) \frac{dt}{t}.$$

We have arrived at the Hilbert space version of Peetre’s J -method of exponent θ . The identity $J_{\nu_\theta} = K_{\varrho_\theta}$ can now be recognized as a sharp (isometric) Hilbert space version of the equivalence theorem of Peetre, which says that the standard K_θ and J_θ -methods give rise to equivalent norms on the category of Banach couples (see [7]).

The problem of determining the pairs ϱ, ν having the property that the K_ϱ and J_ν methods give equivalent norms was studied by Fan in [15, Section 3].

6.6 Other Representations

As we have seen in the preceding subsections, using the space \mathcal{H}_0 to express all involutions and inner products leads to a description of the exact quadratic interpolation spaces in terms of the class P' . If we instead use the space Δ as the basic object, we get Donoghue's representation for interpolation functions. Similarly, one can proceed from any fixed interpolation space \mathcal{H}_* to obtain a different representation of interpolation functions.

6.7 On Interpolation Methods of Power p

Fix a number $p, 1 < p < \infty$. We shall write $L_p = L_p(X, \mathcal{A}, \mu)$ for the usual L_p -space associated with an arbitrary but fixed (σ -finite) measure μ on a measure space (X, \mathcal{A}) . Given a positive measurable weight function w , we write $L_p(w)$ for the space normed by

$$\|f\|_{L_p(w)}^p = \int_X |f(x)|^p w(x) d\mu(x).$$

We shall write $\bar{L}_p(w) = (L_p, L_p(w))$ for the corresponding weighted L_p couple. Note that the conditions imposed mean precisely that $\bar{L}_p(w)$ be separable and regular.

Let us say that an exact interpolation functor F defined on the totality of separable, regular weighted L_p -couples and taking values in the class of weighted L_p -spaces is of power p .

Define, for a positive Radon measure ϱ on $[0, \infty]$, an exact interpolation functor $F = K_\varrho(p)$ by the definition

$$\|f\|_{F(\bar{L}_p(w))}^p := \int_{[0, \infty]} (1 + t^{-\frac{1}{p-1}})^{p-1} K_p(t, f; \bar{L}_p(w)) d\varrho(t).$$

We contend that F is of power p .

Indeed, it is easy to verify that

$$K_p(t, f; \bar{L}_p(w)) = \int_X |f(x)|^p \frac{tw(x)}{(1 + (tw(x))^{-\frac{1}{p-1}})^{p-1}} d\mu(x),$$

so Fubini’s theorem gives that

$$\|f\|_{F(\overline{L}_p(w))}^p = \int_X |f(x)|^p h(w(x)) d\mu(x),$$

where

$$h(\lambda) = \int_{[0,\infty]} \frac{(1+t^{\frac{1}{p-1}})^{p-1} \lambda}{(1+(t\lambda)^{\frac{1}{p-1}})^{p-1}} d\varrho(t), \quad \lambda \in w(X). \tag{6.10}$$

We have shown that $F(\overline{L}_p(w)) = L_p(h(w))$, so F is indeed of power p .

Let us denote by $\mathcal{K}(p)$ the totality of positive functions h on \mathbf{R}_+ representable in the form (6.10) for some positive Radon measure ϱ on $[0, \infty]$.

Further, let $\mathcal{I}(p)$ denote the class of all (exact) *interpolation functions of power p* , i.e., those positive functions h on \mathbf{R}_+ having the property that for each weighted L_p couple $\overline{L}_p(w)$ and each bounded operator T on $\overline{L}_p(w)$, it holds that T is bounded on $L_p(h(w))$ and

$$\|T\|_{\mathcal{L}(L_p(h(w)))} \leq \|T\|_{\mathcal{L}(\overline{L}_p(w))}.$$

The class $\mathcal{I}(p)$ is in a sense the natural candidate for the class of “operator monotone functions on L_p -spaces.” The class $\mathcal{I}(p)$ clearly forms a convex cone; it was shown by Peetre [31] that this cone is contained in the class of concave positive functions on \mathbf{R}_+ (with equality if $p = 1$).

We have shown that $\mathcal{K}(p) \subset \mathcal{I}(p)$. By Theorem 6.1, we know that equality holds when $p = 2$. For other values of p it does not seem to be known whether the class $\mathcal{K}(p)$ exhausts the class $\mathcal{I}(p)$, but one can show that we would have $\mathcal{K}(p) = \mathcal{I}(p)$ provided that each finite-dimensional L_p -couple $\overline{\ell}_p^n(\lambda)$ has the K_p -property (or equivalently, the K -property, see (2.2)). Naturally, the latter problem (about the K_p -property) also seems to be open, but some comments on it are found in Remark 2.9.

Let ν be a positive Radon measure on $[0, \infty]$. In [16], Foias and Lions introduced a method, which we will denote by $F = J_\nu(p)$ in the following way. Define the J_p -functional by

$$J_p(t, f; \overline{L}_p(\lambda)) = \|f\|_0^p + t \|f\|_1^p, \quad f \in \Delta, t > 0.$$

We then define an intermediate norm by

$$\|f\|_{F(\overline{L}_p(\lambda))}^p := \inf \int_{[0,\infty]} (1+t)^{-\frac{1}{p-1}} J_p(t, u(t); \overline{L}_p(\lambda)) d\nu(t),$$

where the infimum is taken over all representations

$$f = \int_{[0, \infty]} u(t) \, d\nu(t)$$

with convergence in Σ . It is straightforward to see that the method F so defined is exact; in [16] it is moreover shown that it is of power p . More precisely, it is there proved that

$$\|f\|_{F(\overline{L}_p(\lambda))}^p = \int_X |f(x)|^p h(w(x)) \, d\mu(x),$$

where

$$h(\lambda)^{-\frac{1}{p-1}} = \int_{[0, \infty]} \frac{(1+t)^{\frac{1}{p-1}}}{(1+t\lambda)^{\frac{1}{p-1}}} \, d\nu(t), \quad \lambda \in w(X). \tag{6.11}$$

Let us denote by $\mathcal{J}(p)$ the totality of functions h representable in the form (6.11). We thus have that $\mathcal{J}(p) \subset \mathcal{I}(p)$. In view of our preceding remarks, we conclude that if all weighted L_p -couples have the K_p property, then necessarily $\mathcal{J}(p) \subset \mathcal{K}(p)$. Note that $\mathcal{J}(2) = \mathcal{K}(2)$ by Theorem 6.6.

Appendix: The Complex Method is Quadratic

Let $S = \{z \in \mathbf{C}; 0 \leq \operatorname{Re} z \leq 1\}$. Fix a Hilbert couple $\overline{\mathcal{H}}$ and let \mathcal{F} be the set of functions $S \rightarrow \Sigma$ which are bounded and continuous in S , analytic in the interior of S , and which maps the line $j + i\mathbf{R}$ into \mathcal{H}_j for $j = 0, 1$. Fix $0 < \theta < 1$. The norm in the complex interpolation space $C_\theta(\overline{\mathcal{H}})$ is defined by

$$\|x\|_{C_\theta(\overline{\mathcal{H}})} = \inf \{ \|f\|_{\mathcal{F}}; f(\theta) = x \}. \tag{*}$$

Let \mathcal{P} denote the set of polynomials $f = \sum_1^N a_i z^i$ where $a_i \in \Delta$. We endow \mathcal{P} with the inner product

$$\langle f, g \rangle_{M_\theta} = \sum_{j=0,1} \int_{\mathbf{R}} \langle f(j + it), g(j + it) \rangle_j P_j(\theta, t) \, dt,$$

where $\{P_0, P_1\}$ is the Poisson kernel for S ,

$$P_j(\theta, t) = \frac{e^{-\pi t} \sin \theta \pi}{\sin^2 \theta \pi + (\cos \theta \pi - (-1)^j e^{-\pi t})^2}.$$

Let M_θ be the completion of \mathcal{P} with this inner product. It is easy to see that the elements of M_θ are analytic in the interior of S , and that evaluation map $f \mapsto f(\theta)$ is continuous on M_θ . Let N_θ be the kernel of this functional and define a Hilbert space \mathcal{H}_θ by

$$\mathcal{H}_\theta = M_\theta / N_\theta.$$

We denote the norm in \mathcal{H}_θ by $\|\cdot\|_\theta$.

Proposition A.1 $C_\theta(\overline{\mathcal{H}}) = \mathcal{H}_\theta$ with equality of norms.

Proof Let $f \in \mathcal{F}$. By the Calderón lemma in [7, Lemma 4.3.2], we have the estimate

$$\log \|f(\theta)\|_{C_\theta(\overline{\mathcal{H}})} \leq \sum_{j=0,1} \int_{\mathbf{R}} \log \|f(j+it)\|_j P_j(\theta, t) dt.$$

Applying Jensen’s inequality, this gives that

$$\|f(\theta)\|_{C_\theta(\overline{\mathcal{H}})} \leq \left(\sum_{j=0,1} \int_{\mathbf{R}} \|f(j+it)\|_j^2 P_j(\theta, t) dt \right)^{1/2} = \|f\|_{M_\theta}.$$

Hence $\mathcal{H}_\theta \subset C_\theta(\overline{\mathcal{H}})$ and $\|\cdot\|_{C_\theta(\overline{\mathcal{H}})} \leq \|\cdot\|_\theta$. On the other hand, for $f \in \mathcal{P}$ one has the estimates

$$\|f(\theta)\|_\theta \leq \|f\|_{M_\theta} \leq \sup\{\|f(j+it)\|_j; t \in \mathbf{R}, j = 0, 1\} = \|f\|_{\mathcal{F}},$$

whence $C_\theta(\overline{\mathcal{H}}) \subset \mathcal{H}_\theta$ and $\|\cdot\|_{C_\theta(\overline{\mathcal{H}})} \geq \|\cdot\|_\theta$. □

It is well known that the method C_θ is of exponent θ (see, e.g., [7]). We have shown that C_θ is an exact quadratic interpolation method of exponent θ .

Complex Interpolation with Derivatives

In [15, pp. 421–422], Fan considers the more general complex interpolation method $C_{\theta(n)}$ for the n :th derivative. This means that in (*), one consider representations $x = \frac{1}{n!} f^{(n)}(\theta)$ where $f \in \mathcal{F}$; the complex method C_θ is thus the special case $C_{\theta(0)}$. It is shown in [15] that, for $n \geq 1$, the $C_{\theta(n)}$ -method is represented, up to equivalence of norms, by the quasi-power function $h(\lambda) = \lambda^\theta / (1 + \frac{\theta(1-\theta)}{n} |\log \lambda|)^n$. The complex method with derivatives was introduced by Schechter [37]; for more details on that method, we refer to the list of references in [15].

References

1. J. Agler, J.E. McCarthy, N. Young, Operator monotone functions and Löwner functions of several variables. *Ann. Math.* **176**, 1783–1826 (2012)
2. Y. Ameer, The Calderón problem for Hilbert couples. *Ark. Mat.* **41**, 203–231 (2003)
3. Y. Ameer, A new proof of Donoghue’s interpolation theorem. *J. Funct. Space Appl.* **3**, 253–265 (2004)
4. Y. Ameer, A note on a theorem of Sparr. *Math. Scand.* **94**, 155–160 (2004)
5. Y. Ameer, M. Cwikel, On the K -divisibility constant for some special, finite-dimensional Banach couples. *J. Math. Anal. Appl.* **360**, 130–155 (2009)
6. N. Aronszajn, W. Donoghue, On exponential representations of functions. *J. Anal. Math.* **5**, 321–388 (1956–57)
7. J. Bergh, J. Löfström, *Interpolation Spaces, an Introduction* (Springer, Berlin, 1976)
8. Y.A. Brudnyi, N.Y. Krugljak, *Interpolation Functors and Interpolation Spaces* (North Holland, Amsterdam, 1991)
9. A.P. Calderón, Intermediate spaces and interpolation, the complex method. *Studia Math.* **24**, 113–190 (1964)
10. A.P. Calderón, Spaces between L_1 and L_∞ and the theorem of Marcinkiewicz. *Studia Math.* **26**, 273–299 (1966)
11. M. Cwikel, Monotonicity properties of interpolation spaces. *Ark. Mat.* **14**, 213–236 (1976)
12. W. Donoghue, *Monotone Matrix Functions and Analytic Continuation* (Springer, Berlin, 1974)
13. W. Donoghue, The theorems of Loewner and Pick. *Israel J. Math.* **4**, 153–170 (1966)
14. W. Donoghue, The interpolation of quadratic norms. *Acta Math.* **118**, 251–270 (1967)
15. M. Fan, Quadratic interpolation and some operator inequalities. *J. Math. Inequal.* **5**, 413–427 (2011)
16. C. Foiaş, J.L. Lions, Sur certains théorèmes d’interpolation. *Acta Sci. Math.* **22**, 269–282 (1961)
17. C. Foiaş, S.C. Ong, P. Rosenthal, An interpolation theorem and operator ranges. *Integr. Equ. Oper. Theory* **10**, 802–811 (1987)
18. P.R. Halmos, Quadratic interpolation. *J. Oper. Theory* **7**, 303–305 (1982)
19. F. Hansen, An operator inequality. *Math. Ann.* **246**, 249–250 (1980)
20. F. Hansen, Selfadjoint means and operator monotone functions. *Math. Ann.* **256**, 29–35 (1981)
21. O. Heinävaara, Local characterizations for the matrix monotonicity and convexity of fixed order. *Proc. Am. math. Soc.* **146**, 3791–3799 (2018)
22. A. Korányi, On some classes of analytic functions of several variables. *Trans. Am. Math. Soc.* **101**, 520–554 (1961)
23. F. Kraus, Über konvexe Matrixfunktionen. *Math. Z.* **41**, 18–42 (1936)
24. J.L. Lions, Espaces intermédiaires entre espaces Hilbertiens et applications. *Bull. Math. de la Soc. Sci. Math. Phys. de la R. P. R.* **2**, 419–432 (1958)
25. J.L. Lions, E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications*, vol. 1 (Springer, New York, 1972)
26. K. Löwner, Über monotone Matrixfunktionen. *Math. Z.* **38**, 177–216 (1934)
27. J.E. McCarthy, Geometric interpolation of Hilbert spaces. *Ark. Mat.* **30**, 321–330 (1992)
28. B. Mityagin, An interpolation theorem for modular spaces. *Mat. Sbornik* **66**, 473–482 (1965)
29. G.J. Murphy, *C^* -Algebras and Operator Theory* (Academic, New York, 1991)
30. V.I. Ovchinnikov, The method of orbits in interpolation theory. *Math. Rep.* **1**, 349–515 (1984)
31. J. Peetre, On interpolation functions III. *Acta Sci. Math. (Szeged)* **30**, 235–239 (1969)
32. J. Peetre, Two new interpolation methods based on the duality map. *Acta Math.* **173**, 73–91 (1979)
33. W. Pusz, S.L. Woronowicz, Functional calculus for sesquilinear forms and the purification map. *Rep. Math. Phys.* **8**, 159–170 (1975)

34. M. Rosenblum, J. Rovnyak, *Hardy Classes and Operator Theory* (Dover, Mineola, 1997)
35. A. Sedaev, Description of interpolation spaces for the pair $(L_p(a_0), L_p(a_1))$. *Soviet Math. Dokl.* **14**, 538–541 (1973)
36. A. Sedaev, E.M. Semenov, On the possibility of describing interpolation spaces in terms of the K -functional of Peetre. *Optimizacija* **4**, 98–114 (1971)
37. M. Shechter, Complex interpolation. *Compos. Math.* **18**, 117–147 (1967)
38. G. Sparr, Interpolation of weighted L^p spaces. *Studia Math.* **62**, 229–271 (1978)
39. G. Sparr, A new proof of Löwner's theorem. *Math. Scand* **47**, 266–274 (1980)
40. A. Uhlmann, Relative entropy and the Wigner–Yanase–Dyson–Lieb concavity in an interpolation theory. *Commun. Math. Phys.* **54**, 21–32 (1977)
41. H. Vasudeva, On monotone matrix functions of two variables. *Trans. Am. Math. Soc.* **176**, 305–318 (1973)