



Riikka Takala

Recommendations

Level I

There are insufficient data to support a Level I recommendation for this topic.

Level II

Hypoxemia and hypotension increase morbidity and mortality after TBI.

Level III

Securing the airway, maintaining adequate blood pressure, oxygenation and ventilation are all key factors in the prevention of secondary brain injury. After stabilization, rapid transport of patients with severe TBI to a neurosurgical centre is crucial.

9.1 Overview

Traumatic brain injury (TBI) is a worldwide health problem with an incidence rate of 262 per 100,000/year. Most TBIs are mild, approximately 70–80%, with the rest being classified as

moderate or severe TBIs. Severe TBI is associated with a high mortality, 30–40% (Maas et al. 2017).

TBI is a dynamic process where the primary injury occurs at the time of insult. Secondary injuries start to develop immediately after the insult and may continue to progress even days or weeks after the insult. The aim of treatment and care is to prevent and decrease secondary injuries. Quick and correct emergency treatment is therefore a corner stone of TBI care. Severe TBI patients should be transferred to a hospital with neurosurgical facilities as quickly as possible.

Tips, Tricks and Pitfalls

- Secondary injuries start to develop within minutes of the insult.
- Hypotension and hypoxia increase morbidity and mortality.
- Hyper- and hypoventilation increase mortality.
- Repeat neurological assessment often: GCS and pupils.
- Continuous monitoring: ECG, SpO₂, blood pressure.
- GCS < 9 patients need to be intubated.

R. Takala (✉)
Perioperative Services, Intensive Care Medicine
and Pain Management, Turku University Hospital,
Turku, Finland

9.2 Summary of Prehospital Treatment

Hypoxemia ($\text{SpO}_2 < 90\%$) and hypotension ($\text{RR}_{\text{sys}} < 120 \text{ mmHg}$) increase morbidity and mortality. They must be avoided and promptly treated. The rule of ABC (ensure protected airway and provide adequate oxygenation and ventilation (breathing) and circulation) applies to the emergency care of TBI.

Patients who have impaired consciousness (Glasgow Coma Scale ≤ 8) are unable to maintain their airway, may have impaired respiratory drive and are at risk of aspiration. Endotracheal intubation and controlled ventilation are therefore needed. Intubation is also needed if the patient has better consciousness than GCS of 9, but has concomitant multitrauma and supplemental oxygen does not correct hypoxemia. One must always assume that head-injured patients also have an injured neck, until otherwise proven. Normoventilation is recommended (etCO_2 35–40 mmHg, 4.5–5.5 kPa) as both hyper- and hypoventilation increase mortality. However, if there are signs of cerebral herniation (i.e. pupil dilatation, Cushing's triad), short lasting hyperventilation may be considered. Hypertonic saline or mannitol may be used to decrease increased intracranial pressure (ICP) and in situations with cerebral herniation.

Hypotension is treated with isotonic or hypertonic crystalloids and vasopressors such as noradrenaline or phenylephrine.

9.3 Background

9.3.1 Prehospital Assessments

9.3.1.1 Oxygenation and Blood Pressure

Hypoxemia, defined as peripheral oxygen saturation (SpO_2) $< 90\%$, and hypotension (previously defined as systolic blood pressure (SBP) $< 90 \text{ mmHg}$) are both associated with high mortality and morbidity. A recent study including 13,151 patients observed that 20.7% and 28.1% patients died if they had only hypotension or hypoxemia, respectively, and mortality was

43.9% if both conditions existed (Spaite et al. 2017a).

Of those patients who did not have hypotension or hypoxemia, only 5.6% died. Updated Brain Trauma Foundation (BTF, www.brain-trauma.org) guidelines from 2017 recommends systolic blood pressure values higher than 90 mmHg. They recommend that SBP should be kept at $\geq 100 \text{ mmHg}$ for patients 50–69 years and at $\geq 110 \text{ mmHg}$ or higher for patients 15–49 or >70 years old. In a TBI patient material consisting of over 3800 patients, aged 10 years or older, there was a linear association between systolic blood pressure, ranging from 40 to 119 mmHg, and death. An increase of systolic blood pressure of 10 mmHg was associated with an 18.8% decreased adjusted odds of death (Spaite et al. 2017b). A bimodal distribution of death has been observed in TBI patients, optimal SBP being between 120 and 140 mmHg (Zafar et al. 2011).

Cerebral perfusion pressure (CPP) is calculated as: mean arterial pressure (MAP) – intracranial pressure (ICP). BTF recommends that CPP should be kept between 60 and 70 mmHg.

Blood pressure should be monitored frequently if it is measured noninvasively. If possible, an arterial cannulation and continuous blood pressure measurement would be preferable as it is less prone to disturbances during the transportation and allows simple sampling for blood gas analysis. Electrocardiography and SpO_2 must also be monitored continuously.

9.3.1.2 Glasgow Coma Scale

The most common tool for assessing level of consciousness is the Glasgow Coma Scale (GCS) (Table 9.1). Chapter 3 discusses this topic further, and www.glasgowcomascale.org demonstrates in detail how it should be performed. GCS is a clinically important tool, and it should be performed immediately after the ABC evaluation. When evaluating GCS, hypoxaemia, hypoventilation, hypotension and hypoglycaemia must be corrected, otherwise GCS is not reliable. There are also confounding factors that may decrease the summed GCS score, such as alcohol and narcotic drugs, spinal cord injury, orbital swelling and dysphasia. Repeated and documented GCS are valuable means for monitoring the trend of con-

Table 9.1 GCS

| GCS | |
|------------------------------------|---|
| <i>Eye opening</i> | |
| Spontaneous | 4 |
| Speech | 3 |
| Pain | 2 |
| None | 1 |
| <i>Verbal response</i> | |
| Oriented | 5 |
| Confused | 4 |
| Inappropriate | 3 |
| Incomprehensible | 2 |
| None | 1 |
| <i>Motor response</i> | |
| Obeys command | 6 |
| Localize pain | 5 |
| Flexor withdrawal (normal flexion) | 4 |
| Abnormal flexion | 3 |
| Extension | 2 |
| None | 1 |

sciousness and for identifying possible worsening in neurological status. GCS is known to correlate with the severity of TBI. However, GCS at the scene is not such a reliable outcome predictor as GCS obtained upon admission, and motor score has the best predictive power (Marmarou et al. 2007; Lesko et al. 2013). GCS is performed by evaluating the patient's eye opening (1–4 points), verbal response (1–5 points) and best motor response (1–6 points) to speech or painful stimulus. When testing eye opening to pain, a peripheral stimulus should be used as the grimace associated with central pain may cause eye closure. If patients do not obey commands and motor response to painful stimulus is used, then central stimulus should be used. This is done by pressing the supraorbital notch. However, in cases of facial fractures adjacent to supraorbital notch, a peripheral stimulus is safer and is performed by applying slowly increasing pressure to nail tip. The points of each area are recorded separately, and the summed GCS score must be documented.

9.3.1.3 Pupils and Their Reactivity

Chapter 3 elaborates on this topic in detail. Evaluating and documenting the size of the pupils (in mm) and their reactivity to light are also essential as any changes in them give valuable

information. Normal pupils may have a 1 mm diameter difference. A bright flashlight is used to assess the light reactivity. In a normal situation, the pupil constricts to a light and so does the contralateral pupil (direct and consensual reaction). Both the right and left pupil findings must be documented.

Before pupil evaluation, hypoxemia, hypoventilation and hypotension must be treated.

An important sign of uncal herniation is a unilaterally dilated and unreactive (fixed) pupil. In this situation, the ipsilateral anterior temporal lobe is pushed through the tentorium and the uncus of the temporal lobe compresses the oculomotor nerve (III cranial nerve). Damage to the medulla results in bilaterally dilated and fixed pupils. Acute pupil dilation is considered a neurological emergency.

Orbital trauma or a direct trauma to the eye may injure short ciliary nerve and pupillary sphincter muscle, respectively, also resulting in a dilated pupil. Documentation of traumas to the aforementioned areas is therefore important.

9.4 Prehospital Treatment

9.4.1 Airway, Ventilation and Oxygenation

A TBI patient with GCS ≤ 8 must be intubated as they are not able to maintain their airway and may have low respiratory drive, and they are at risk of aspiration. Short-acting anaesthetic agents should be used for intubation and maintaining sedation during the transport as the neurological status is often reassessed when the patient arrives to the emergency department. Short-acting opioids, propofol and short-acting benzodiazepines are suitable for intubation. Ketamine has become popular in the emergency medicine due to its stable haemodynamical properties, but there are currently not enough data regarding its safety in TBI patients, used as a sole anaesthetic agent. Some data shows that, when it is used in conjunction with other anaesthetics such as midazolam or propofol and patients are already mechanically ventilated, it does not increase ICP (Chang et al. 2013). In addition, hallucinations associated with

ketamine administration may impair reliable neurological assessment upon admission to the hospital. However, in haemodynamically unstable patients with multiple trauma, ketamine is probably a safe choice as propofol is known to decrease blood pressure.

Prehospital intubation of TBI patients has resulted in conflicting results regarding their survival and neurological outcome. These contradictory results seem to arise from the experience and skills of the personnel. Data supports that unexperienced and untrained personnel should not intubate TBI patients (Bossers et al. 2015). The correct endotracheal tube placement must be ensured by pulmonary auscultation and end-tidal CO₂ (etCO₂) measurement. Endotracheal tubes should be secured with drapes, and the head of the bed should be elevated 15–30° and neck and head positioned in neutral position to ensure cerebral venous return. Of note, semi-rigid collars may also obstruct cerebral venous return.

Normoventilation (etCO₂ 35–40 mmHg, 4.5–5.5 kPa) is recommended as both hypoventilation and hyperventilation result in increased mortality (Davis et al. 2010). Hypoxaemia is corrected by administering supplemental oxygen. SpO₂ is aimed to be maintained at >90%. Hyperoxia may be harmful in critically ill patients and normoxia is suggested in TBI patients, although high-quality studies on the topic are still lacking.

9.4.2 Fluid Resuscitation

Isolated head trauma does not lead to hypovolaemia but may cause hypotension (Mahoney et al. 2003). However, if the patient has concomitant high cervical injury, this may cause neurogenic shock with hypotension and bradycardia. In addition, TBI with concomitant multitrauma with haemorrhage may result in hypovolaemia and hypotension. Both result in reduced cerebral perfusion pressure and oxygen delivery, predisposing the already injured and vulnerable brain to secondary injuries. Fluid resuscitation aims to restore oxygen delivery and adequate cerebral perfusion pressure and cerebral blood flow. Isotonic or hypertonic crystalloids (Gantner et al.

2014) are suitable for fluid resuscitation in the TBI patients. Hypertonic crystalloids can also be used as they decrease the ICP, but they do not improve the outcome of the patients (Cooper et al. 2004). The use of colloids and albumin are not recommended (Myburgh et al. 2007; Reinhart et al. 2012; Cooper et al. 2013; Oddo et al. 2018) as they may increase the mortality in TBI patients. Glucose containing fluids should be avoided, unless the patient has hypoglycaemia (B-glucose <5 mmol/L). If fluid therapy does not increase the blood pressure, vasoactive agents should be used. Noradrenaline and phenylephrine are recommended as continuous infusion.

9.4.3 Cerebral Herniation

Frequent neurological assessment is important as the patient's clinical status can worsen rapidly due to ongoing processes such as mass lesions, cerebral oedema or hydrocephalus. Clinical signs of increased ICP and cerebral herniation are loss of consciousness or progressive neurologic deterioration (at least 2 points decrease in GCS), one or both pupils dilated and unreactive to light, Cushing's triad (hypertension and bradycardia) and extension or no reaction to painful stimulus.

Sedation and analgesia must be sufficiently provided, preferably with a continuous infusion.

Hyperosmolar therapy with either mannitol or hypertonic saline is recommended. They both increase osmotic gradient, which induces water to move from cerebral tissue to the extracellular space, thus reducing cerebral oedema. Mannitol may cause electrolyte disturbances and hypovolemia due to excessive diuresis (Torre-Healy et al. 2012). Hypertonic saline increases cardiac output, blood pressure and cerebral blood flow (Torre-Healy et al. 2012; Tyagi et al. 2007). Hypertonic saline may be more effective and have longer lasting property to reduce ICP than mannitol (Kamel et al. 2011; Mortazavi et al. 2012).

Hyperventilation constricts cerebral arterial vessels and reduces cerebral flow (CBF) and ICP. CBF is often reduced in the acute phase of TBI and further decrease in CBF may lead to

cerebral ischemia. In a small study of TBI patients, hyperventilation decreased ICP and improved CPP but increased hypoperfused brain volume (Coles et al. 2002). Hyperventilation worsens the neurological outcome in TBI patients (Stocchetti and Maas 2014). However, short-term hyperventilation can be useful in cases of imminent herniation and may outweigh its harmful effects. In such a case, hyperventilation should be titrated to etCO_2 of 30–35 mmHg, 3.9–4.6 kPa.

In patients with concomitant multitrauma, increased ICP can also result from increased venous pressure due to pneumothorax or abdominal injury. Pneumothorax must be treated promptly at the scene with chest drain or with thoracosentesis if drainage is not possible.

9.4.4 Seizure Treatment

TBI patients are at risk of epileptic seizures, which may cause and worsen secondary injuries. They must be promptly treated. Intravenous propofol or benzodiazepines such as lorazepam or midazolam are first-line choices. Of antiepileptic drugs, phosphenytoine and levetiracetam are currently the most widely used. They both seem to prevent early seizures in TBI, but in long-term use, phosphenytoine may worsen neurological outcome (Szaflarski et al. 2010; Bhullar et al. 2014).

9.4.5 Hypothermia

Hypothermia has been considered as neuroprotective, and several studies have assessed its role in the outcome of TBI patients. The recent POLAR study demonstrated that early prophylactic hypothermia in TBI patients does not improve neurological outcome or reduce mortality (Cooper et al. 2018), and its use is not recommended.

9.5 Transport

TBI patients should be transported to Level I or Level II trauma centres with 24 h coverage of neurosurgery, anaesthesiology, radiology (CT),

traumatology and intensive care. There are no prospective randomized studies comparing “scoop and run” with “stay and play” on the outcome of TBI patients, and both the approaches are used in Nordic countries (Cnossen et al. 2018).

Severe TBI patients seem to benefit from a rapid transfer directly to a neurotrauma centre, and during transport all efforts must be done to prevent secondary injuries.

9.6 Paediatric TBI

Paediatric patients are not small adults, and the aetiology and pathology of paediatric TBI differ from that of adults. However, like in adults, hypotension and hypoxia increase mortality in paediatric TBI patients. Accordingly, assessment, treatment approaches and principals, such as avoiding hypoxemia and hypotension, are applied in the same way as in adults.

Paediatric patients should be transferred to Level I or II trauma centres familiar with paediatric TBI patients or to trauma centres familiar with paediatric TBI patients.

9.6.1 Oxygenation and Blood Pressure

Paediatric SpO_2 target is also $>90\%$, whereas blood pressure targets are age dependent. In the paediatric population, hypotension is defined according to American College of Surgeons as follows:

Age 0–28 days, SBP < 60 mmHg, 1–12 months, SBP < 70 mmHg, 1–10 years, SBP $< 70 + 2 \times \text{age in years}$, >10 years, SBP < 90 mmHg. Current paediatric TBI guidelines from 2019 state that in children minimal CPP is 40 mmHg and threshold 40–50 mmHg should be considered, with infants at the lower end and adolescents at the upper end (Kochanek et al. 2019).

However, it may be that these hypotension and CPP thresholds in paediatric TBI patients are too low (Suttipongkaset et al. 2018), and they should

Table 9.2 Paediatric GCS

| PGCS | |
|-----------------------------|---|
| <i>Eye opening</i> | |
| Spontaneous | 4 |
| Speech | 3 |
| Pain | 2 |
| None | 1 |
| <i>Verbal response</i> | |
| Coos, babbles | 5 |
| Irritable cries | 4 |
| Cries to pain | 3 |
| Moans to pain | 2 |
| None | 1 |
| <i>Motor response</i> | |
| Normal spontaneous movement | 6 |
| Withdraws to touch | 5 |
| Withdraws to pain | 4 |
| Abnormal flexion | 3 |
| Extension | 2 |
| None | 1 |

be more age specific (Allen et al. 2014), but data is currently too limited.

9.6.2 Paediatric GCS

Just like in adults, in paediatric patients GCS is assessed after stabilization. In children over 2 years old, adult GCS can be employed, whereas the paediatric GCS is used in pre-verbal children (Table 9.2).

References

- Allen BB, Chiu Y-L, Gerber LM, Ghajar J, Greenfield JP. Age-specific cerebral perfusion pressure thresholds and survival in children and adolescents with severe traumatic brain injury. *Pediatr Crit Care Med.* 2014;15:62–70.
- Bhullar IS, Johnson D, Paul JP, Kerwin AJ, Tepas JJ III, Frykberg ER. More harm than good. Antiseizure prophylaxis after traumatic brain injury does not decrease seizure rates but may inhibit functional recovery. *J Trauma Acute Care Surg.* 2014;76:54–61.
- Bossers SM, Schwarte LA, Loer SA, Twisk JWR, Boer C, Schober P. Experience in prehospital endotracheal intubation significantly influences mortality of patients with severe traumatic brain injury: a systematic review and meta-analysis. *PLoS One.* 2015;10:e0141034–26.
- Chang LC, Raty SR, Ortiz J, Bailard NS, Mathew SJ. The emerging use of ketamine for anesthesia and sedation in traumatic brain injuries. *CNS Neurosci Ther.* 2013;19:390–5.
- Crossen MC, van der Brande R, Lingsma HF, Polinder S, Lecky F, Maas AIR. Prehospital trauma care among 68 European neurotrauma centers: results of the CENTER-TBI provider profiling questionnaires. *J Neurotrauma.* 2018. <https://doi.org/10.1089/neu.2018.5712>.
- Coles JP, Minhas PS, Fryer TD, Smielewski P, Aigbirihio F, Donovan T, et al. Effect of hyperventilation on cerebral blood flow in traumatic head injury: clinical relevance and monitoring correlates. *Crit Care Med.* 2002;30:1950–9.
- Cooper DJ, Myles PS, McDermott FT, Murray LJ, Laidlaw J, Cooper G, et al. Prehospital hypertonic saline resuscitation of patients with hypotension and severe traumatic brain injury—a randomized controlled trial. *JAMA.* 2004;291:1350–7.
- Cooper DJ, Myburgh J, Heritier S, Finfer S, Bellomo R, Billot L, et al. Albumin resuscitation for traumatic brain injury: is intracranial hypertension the cause of increased mortality? *J Neurotrauma.* 2013;30(7):512–8.
- Cooper DJ, Nichol AD, Bailey M, Bernard S, Cameron PA, Pili-Floury S, et al. Effect of early sustained prophylactic hypothermia on neurologic outcomes among patients with severe traumatic brain injury. *JAMA.* 2018;320:2211–20.
- Davis DP, Peay J, Sise MJ, Kennedy F, Simon F, Tominaga G, et al. Prehospital airway and ventilation management: a trauma score and injury severity score-based analysis. *J Trauma.* 2010;69:294–301.
- Gantner D, Moore EM, Cooper DJ. Intravenous fluids in traumatic brain injury. *Curr Opin Crit Care.* 2014;20:385–9.
- Kamel H, Navi BB, Nakagawa K, Hemphill JC III, Ko NU. Hypertonic saline versus mannitol for the treatment of elevated intracranial pressure: a meta-analysis of randomized clinical trials. *Crit Care Med.* 2011;39:554–9.
- Kochanek PM, Tasker RC, Carney N, Totten AM, Adelson PD, Selden NR, et al. Guidelines for the management of pediatric severe traumatic brain injury, third edition. *Pediatr Crit Care Med.* 2019;20:S1–S82.
- Lesko MM, Jenks T, O'Brien SJ, Childs C, Bouamra O, Woodford M, Lecky F. Comparing model performance for survival prediction using total Glasgow Coma Scale and its components in traumatic brain injury. *J Neurotrauma.* 2013;30:17–22.
- Maas AIR, Menon DK, Adelson PD, Andelic N, Bell MJ, Belli A, et al. The lancet neurology commission traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol.* 2017;16:1–62.
- Mahoney EJ, Biffi WL, Harrington DT, Cioffi WG. Isolated brain injury as a cause of hypotension in the blunt trauma patient. *J Trauma.* 2003;55(6):1065–9.
- Marmarou A, Lu J, Butcher I, McHugh GS, Murray GD, Steyerberg EW, et al. Prognostic value of the Glasgow Coma Scale and pupil reactivity in traumatic brain injury assessed pre-hospital and on

- enrollment: an IMPACT analysis. *J Neurotrauma*. 2007;24:270–80.
- Mortazavi MM, Romeo AK, Deep A, Griessenauer CJ, Shoja MM, Tubbs RS, Fisher W. Hypertonic saline for treating raised intracranial pressure: literature review with meta-analysis. *J Neurosurg*. 2012;116:210–21.
- Myburgh J, Cooper DJ, Finfer S, Bellomo R, Norton R, Bishop N, et al. Saline or albumin for fluid resuscitation in patients with traumatic brain injury. *N Engl J Med*. 2007;357:874–84.
- Oddo M, Poole D, Helbok R, Meyfroidt G, Stocchetti N, Bouzat P, et al. Fluid therapy in neurointensive care patients: ESICM consensus and clinical practice recommendations. *Intensive Care Med*. 2018;44:449–63.
- Reinhart K, Perner A, Sprung CL, Jaeschke R, Schortgen F, Johan Groeneveld AB, et al. Consensus statement of the ESICM task force on colloid volume therapy in critically ill patients. *Intensive Care Med*. 2012;38:368–83.
- Spaite DW, Hu C, Bobrow BJ, Chikani V, Barnhart B, Gaither JB, et al. The effect of combined out-of-hospital hypotension and hypoxia on mortality in major traumatic brain injury. *Ann Emerg Med*. 2017a;69:62–72.
- Spaite DW, Hu C, Bobrow BJ, Chikani V, Sherrill D, Barnhart B, et al. Mortality and prehospital blood pressure in patients with major traumatic brain injury. *JAMA Surg*. 2017b;152:360–9.
- Stocchetti N, Maas AIR. Traumatic intracranial hypertension. *N Engl J Med*. 2014;370:2121–30.
- Suttipongkaset P, Chaikittisilpa N, Vavilala MS, Lele AV, Watanitanon A, Chandee T, Krishnamoorthy V. Blood pressure thresholds and mortality in pediatric traumatic brain injury. *Pediatrics*. 2018;142:e20180594–9. <https://doi.org/10.1542/peds.2018-0594>.
- Szaflarski JP, Sangha KS, Lindsell CJ, Shutter LA. Prospective, randomized, single-blinded comparative trial of intravenous levetiracetam versus phenytoin for seizure prophylaxis. *Neurocrit Care*. 2010;12:165–72.
- Torre-Healy A, Marko NF, Weil RJ. Hyperosmolar therapy for intracranial hypertension. *Neurocrit Care*. 2012;17:117–30.
- Tyagi R, Donaldson K, Loftus CM, Jallo J. Hypertonic saline: a clinical review. *Neurosurg Rev*. 2007;30:277–90.
- Zafar SN, Millham FH, Chang Y, Fikry K, Alam HB, King DR, et al. Presenting blood pressure in traumatic brain injury: a bimodal distribution of death. *J Trauma*. 2011;71:1179–84.