

# *Drosophila* Eye as a Model to Study Regulation of Growth Control: The Discovery of Size Control Pathways



Shilpi Verghese, Indrayani Waghmare, Shree Ram Singh, and Madhuri Kango-Singh

## Introduction

In the biological sense, the term growth has intricate ramifications that we have only started to comprehend. Growth is the overall increase in cell mass or size of a tissue or organism (Conlon and Raff 1999; Cook and Tyers 2007; Edgar 1999; Raff 1996). Growth may be due to increase in cell number resulting from cell division (cell proliferation), increase in cellular mass without cell division (cell enlargement), or release of more extracellular matrix (cell accretion). These processes are intimately linked, and it is clear that if coordinated growth has to occur in an organism, it is necessary for various biological pathways to interact and relay appropriate signals to proper cell types. Growth regulation is precisely controlled and affected by several intrinsic and extrinsic factors (Cooper 2004; Crickmore and Mann 2008;

---

Shilpi Verghese and Indrayani Waghmare contributed equally to this work.

---

S. Verghese

Department of Cell Biology, Emory University School of Medicine, Atlanta, GA, USA

I. Waghmare

Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN, USA

S. R. Singh

Mouse Cancer Genetics Program, National Cancer Institute, Frederick, MD, USA

M. Kango-Singh (✉)

Department of Biology, University of Dayton, Dayton, OH, USA

Center for Tissue Regeneration and Engineering at Dayton (TREND), University of Dayton, Dayton, OH, USA

Premedical Program, University of Dayton, Dayton, OH, USA

Integrative Science and Engineering Center (ISE), University of Dayton, Dayton, OH, USA

e-mail: [mkangosingh1@udayton.edu](mailto:mkangosingh1@udayton.edu)

© Springer Nature Switzerland AG 2020

A. Singh, M. Kango-Singh (eds.), *Molecular Genetics of Axial Patterning, Growth and Disease in Drosophila Eye*,

[https://doi.org/10.1007/978-3-030-42246-2\\_8](https://doi.org/10.1007/978-3-030-42246-2_8)

Grebien et al. 2005; Johnston and Gallant 2002). The intrinsic factors mainly involve synthesis and secretion of signals or ligands, which bind to their cognate receptors to relay downstream signals. These signals consist of a variety of molecules such as hormones, mitogens, apoptosis-inducing signals, patterning and axis-determining signals, etc. which eventually determine organ size and tissue homeostasis (Johnston and Gallant 2002; Mitchison et al. 1997; Montagne 2000; Tumaneng et al. 2012a). Growth of a tissue or organ is impacted not only by cell division but also by regulated cell death (apoptosis or programmed cell death) (Bangs and White 2000; Jacobson et al. 1997; Martin et al. 2009; Oldham et al. 2000a; Richardson and Kumar 2002; Rusconi et al. 2000).

In this chapter, we will focus on growth regulation in imaginal discs (epithelial sacs that are precursors of adult appendages) in *Drosophila melanogaster* (Bergantinos et al. 2010; Bryant 1978, 1987, 2001; Bryant and Schmidt 1990). The obvious advantages that *Drosophila* has to offer as a model organism include short life cycle, high fecundity, low-cost maintenance, and lack of redundancy in genome (Bier 2005; Blair 2003; Boutros and Ahringer 2008; Pagliarini et al. 2003; St Johnston 2002; Vidal and Cagan 2006). Furthermore, the sophisticated fly genetics provides great deal of versatility in terms of designing experiments. The plethora of knowledge thus generated through exhausting efforts of scientists has not only revealed to us classic information about how growth occurs but has also led to better understanding of growth-related diseases such as cancer.

## ***Drosophila* Eye as a Model to Study Regulation of Growth**

The compound eyes of *Drosophila* arise from the eye-antennal imaginal discs, a monolayer epithelial sheet of cells that is responsible for the development of the eyes, the antennae, the ocelli, and a major part of the adult head cuticle. Each eye of the adult fruit fly on an average consists of about 800 ommatidia (Wolff and Ready 1993). Ommatidia arise from a set of 19 precursor cells that are generated by spatially and temporally coordinated cellular processes such as cell proliferation, cell differentiation, and cell death in the eye imaginal discs. Eighteen of these cells contribute to the eye per se, whereas the 19th cell gives rise to a sensory bristle (Cagan 1993). A key feature that distinguishes the eye from the rest of the organs is its ability to perceive light and relay the signal to distinct areas in the brain called the optic lobes. The eye imaginal discs arise from about 50 primordial cells that express the *Drosophila* PAX 6 gene *eyeless* (*ey*) during mid to late embryogenesis. Two such discs develop in each larva and differentiate into two compound eyes, antennae, ocelli, and the head cuticle in the adult.

Much is known about the regulation of growth and differentiation of the eye-antennal imaginal discs (Baker 2001; Cagan 1993; Dominguez and Casares 2005; Hafen 1991; Kramer and Cagan 1994; Kumar 2001). Until the second larval instar of development, the cells of the eye-antennal discs proliferate without differentiation (Baker 2001; Wolff and Ready 1993). During the second instar stage, a unique

process of cell differentiation begins in the eye-antennal disc that paves the way for formation of photoreceptor neurons in the posterior region of the eye-antennal imaginal disc (Wolff and Ready 1993). The differentiation occurs in the wake of a so-called morphogenetic furrow—a front marked by apical constriction of epithelial cells in response to complex developmental signaling from the Hedgehog, Dpp, Wg, and EGFR pathways (Acquisti et al. 2009; Chen and Chien 1999; Firth et al. 2010; Harvey et al. 2001; Kango-Singh et al. 2003; Penton et al. 1997). Posterior to the morphogenetic furrow, the cells begin to acquire particular photoreceptor cell fates and organize into ommatidial clusters.

Anterior to the furrow, cells divide asynchronously and do not differentiate; however, in the morphogenetic furrows, cells are arrested in the G1 phase of the cell cycle, synchronize, and either start to differentiate into photoreceptor cells as they leave the furrow or undergo one additional round of cell division, referred to as the second mitotic wave (SMW) before differentiating into the remaining photoreceptor, cone, pigment, and bristle cells (Baker 2001; Dickson and Hafen 1993; Wolff and Ready 1993). The cells posterior to the morphogenetic furrow enter G1 arrest caused by Dpp (*decapentaplegic*) signaling that is maintained by the *roughex* (*rx*) gene, which negatively regulates G1-S transition. The cells that are temporarily trapped in the G1 phase begin differentiation with specification of the R8 (photoreceptor) cell due to expression of the proneural protein Atonal (*Ato*) (Baker et al. 1996; Chen and Chien 1999; Daniel et al. 1999; Dominguez 1999; Greenwood and Struhl 1999; Jarman et al. 1994). R8 recruits other photoreceptor cells—R2, R3, R4, and R5—to form a cluster of five photoreceptor precursors. Once specified, these cells never enter cell cycle or cell division again. All other non-specified cells reenter cell cycle only once—a process referred to as the second mitotic wave (SMW) (Anon 2003; Baker 2001; de Nooij and Hariharan 1995). Cells in SMW undergo G2/M phase that is mediated through local signaling from Spitz (*Spi*). Binding of *Spi* to its cognate receptor EGFR in precursor cells causes activation of downstream *string* (*Bakal*) that completes the G2-M transition during mitosis. Local *Spi*-EGFR signaling also plays an important role limiting the progression of SMW. For instance, on an average, the *Spi* signal from one pre-cluster can span to a length of seven cells only causing these cells to divide, whereas the remaining cells remain arrested in G2 phase and fail to divide (Baker 2001; Brumby and Richardson 2003) (de Nooij and Hariharan 1995; Jarman et al. 1994; Price et al. 2002) (Wolff and Ready 1991). The progression of the morphogenetic furrow is complete by the mid-third instar of larval development, and the eye-antennal disc is fully grown to about 50,000 cells (Kumar 2009; Kumar and Moses 2000, 2001; Sun 2007).

Following development in larval stages, supernumerary cells are eliminated via apoptosis during pupal development. This event is mediated through Notch signaling (Bonini and Fortini 1999; Burke and Basler 1997; Sawamoto and Okano 1996; Treisman and Heberlein 1998; Zipursky 1989). By contrast, survival of pupal cells is brought about by EGFR expression that mediates its cell survival function through suppressing the transcriptional activity of the pro-apoptotic gene *head involution defective* (*hid*) (Bonini and Fortini 1999). In addition, survival signals emanating from cone or primary pigment cells in each ommatidium play a role in survival and


proliferation of secondary and tertiary pigment cells and secondary bristle organs (Cagan 1993, 2009; Rubin 1989; Singh et al. 2012; Tsachaki and Sprecher 2012; Yamamoto 1993). During metamorphosis, the two eye-antennal imaginal discs fuse at the dorsal midline to form the fly head with three ocelli, two antennae, and compound eyes. Thus, the eye-antennal disc is ideal for the study of organogenesis, morphogenesis, pattern formation, and several cell biological processes including the regulation of cell cycle, cell death, cell junctions and adhesion, transport of molecules, cell signaling, and metabolism. Recently, the eye discs have been used as an experimental system for genetic screens to discover postembryonic lethality and for screening small molecule inhibitors in chemical and drug screens.

## The Mosaic Analysis Systems and the *Drosophila* Eye

Mutagenesis screen is a very well-established tool for gene discovery in flies [for review, see (Bellen et al. 2011, 1989; Blair 2003; Pfeiffer et al. 2010; St Johnston 2002; Venken and Bellen 2012; Xu and Rubin 1993)]. Over the years, the **mosaic techniques** have evolved to include the FLP-FRT, eyFlp, EGUF, eyFlp cl w+, Flp-out clones, and MARCM [for review, see (Blair 2003; St Johnston 2002)]. One of the first tissue-specific mosaic systems was developed in the eye-antennal discs where the mosaic clones were restricted to the eye-antennal discs by virtue of expression of the Flippase gene under the control of the eyeless promoter (commonly referred to as the “*ey-FLP* system”) (Newsome et al. 2000). This tissue-specific system was further refined by the development of the “cell-lethal” system, where effects of loss of function of a gene could be surveyed more clearly because the wild-type twin clones are eliminated due to the presence of *cell-lethal* mutations (the *cell-lethal FLP-FRT* system) (Newsome et al. 2000). We focus on the genetic screens performed about 10–12 years ago (simultaneously in our labs) that lead to the identification of many new genes that were shown to belong to the two major growth regulatory networks: the Hippo pathway and the TSc-ToR pathway.

## Genetic Screens for Genes That Regulate Growth: The “Big-Head” and “Pin-Head” Mutations

Barry Dickson’s group (Newsome et al. 2000) improved the traditional FLP-FRT approach developed in the Rubin Lab (Xu and Rubin 1993), to allow generation of essentially mutant eye discs by eliminating the wild-type twin clone via a *cell-lethal* mutation (the *cell-lethal FLP-FRT* system) (Fig. 1). This so-called “cell-lethal” approach allows the mutant clones to grow to their highest potential due to elimination of competitive interactions between the mutant cells and their wild-type neigh-



**Fig. 1** Mutagenesis schemes for eye-specific mosaics lead to the identification of several Hippo and Tsc-TOR pathway mutants. (a) Modified mutagenesis scheme, (b) typical phenotypes of Hippo and Tsc-TOR pathway mutant from the mutagenesis screen


bors. Using this system, several groups carried out mutagenesis screens in flies (on the X, 2L, 2R, 3L, 3R chromosomes) and found mutations that affected patterning, growth, cell death, and differentiation [for review, see (St Johnston 2002)].

Of special interest were gene mutations which caused a remarkable effect on growth without disrupting the patterning process (Conlon and Raff 1999; Johnston and Gallant 2002; Mitchison et al. 1997; Oldham et al. 2000a; Raff 1996; Su and O’Farrell 1998; Tumaneng et al. 2012a). Characterization of these mutants revealed the mechanisms that regulate growth and tissue size by controlling cell number (Hippo pathway) (Zhao et al. 2011b) or cell size (InR/TSC-TOR pathway) (Kim and Guan 2011; Loewith 2011; Montagne 2000; Potter et al. 2003; Soulard et al. 2009) in a developing organ. Typically, loss-of-function mutations in positive regulators of these pathways caused development of enlarged heads that showed overgrowth—referred to as the “big-head” mutations (Hafen 2004; Oldham and Hafen 2003; Pan 2007, 2010). In contrast, loss-of-function mutations in negative regulators of these pathways caused reduction in head size and development of smaller organs, which may be due to cell death or reduction in cell size, and were referred to as the “pin-head” mutations.

## The Hippo Signaling Pathway

The Hippo signaling pathway was first discovered in flies following characterization of “big-head” mutants identified from genetic screens [for review, see (Edgar 2006; Pan 2007; Saucedo and Edgar 2007)]. Analysis of the loss-of-function phenotypes revealed that a fundamental function of the Hippo pathway was the regulation of organ size (Boggiano and Fehon 2012; Harvey and Hariharan 2012; Schroeder and Halder 2012; Staley and Irvine 2012). Interestingly, the pathway received its name just after some growth regulatory genes [*warts* (*wts*), *salvador* (*sav*, aka *shar-pie*, *shrp*)] were characterized. Warts (*wts*) was named based on the bumpy “warts-like” phenotype of the mutant cells in mitotic (mosaic) clones on the body of the adult flies that were reminiscent of the warts on toads (Justice et al. 1995). Another group led by Xu et al. (1995) also independently found *warts* in the initial FLP/FRT-based screen and named it *large tumor suppressor* (*lats*) (Xu et al. 1995). Two independent groups identified the gene encoding the adaptor protein Salvador (*Sav*) (aka *Shar-pie*, *Shrp* after the dog species of the same name as the mutant flies showed a characteristic phenotype of folded dark cuticle on the overgrown heads) from complementation groups isolated from the big-head genetic screens (Kango-Singh et al. 2002; Tapon et al. 2002). Interestingly, both *Wts* and *Sav* regulated growth by suppressing proliferation and promoting apoptosis. Hippo was the name given to another complementation group from the “big-head” screens that showed a phenotype that was very similar to *Wts* and *Sav* (Harvey et al. 2003; Jia et al. 2003; Pantalacci et al. 2003; Udan et al. 2003; Wu et al. 2003).

Molecular analysis of the three genes revealed that *Wts* and *Hpo* genes encode for serine-threonine (S-T) kinases, whereas *Sav* is a WW domain-containing adaptor protein (Kango-Singh et al. 2002; Tapon et al. 2002). By this time, it was clear that Warts, Salvador, and Hippo all show similar loss-of-function phenotypes and control organ size by a common signaling pathway that promotes apoptosis and restricts cell proliferation (Edgar 2006; O'Neill and Kolch 2005; Rothenberg and Jan 2002), and the pathway got its name from the last member of this trio of genes. A complete pathway that relays a growth regulatory signal from the plasma membrane to the nucleus has emerged over the last decade. Although genetic mutagenesis screens led to the initial discovery of this pathway, several components were identified by other genetic screening strategies and biochemical approaches (e.g., yeast two-hybrid screens, TAP-TAG-based protein interaction assays) [for review, see (Halder and Johnson 2011; Kango-Singh and Singh 2009; Staley and Irvine 2012; Tumaneng et al. 2012a; Varelas and Wrana 2012)]. Today the Hippo pathway has grown to a large network of tumor suppressor genes that function upstream and downstream of the three initial members of the Hippo pathway (aka the core kinase cascade) that control several aspects of tissue homeostasis. Overall, the Hippo signaling pathway is a key size regulatory pathway that controls organ size in flies and vertebrates, and misregulation of Hippo signaling is implicated in several diseases including cancer [for review, see (Harvey and Hariharan 2012; Schroeder and Halder 2012; Staley and Irvine 2012; Zhao et al. 2011b)] (Fig. 2).



**Fig. 2** Schematic representation of the Hippo pathway in *Drosophila melanogaster*. (a) Hippo pathway is downregulated in response to extracellular signals. Hippo (Hpo, #3206) fails to get phosphorylated and does not phosphorylate Warts (Wts). Inactive Wts cannot phosphorylate Yorkie (Yki) and allows Yki to enter the nucleus to bind cognate transcription factors and induce expression of target genes. (b) Hippo pathway is activated by stress, wherein Hippo (Hpo, #3206) is phosphorylated and in turn phosphorylates Warts (Wts) with the help of adaptor proteins Salvador (Sav) and Mats. Activated Wts phosphorylates Yorkie (Yki) and prevents it from entering the nucleus, thus preventing transcription of target genes. In addition, cell death is induced when the pathway is hyperactivated

### Regulation by Core Kinase Cascade of the Hippo Pathway

The molecular analysis of the three initial members of the Hippo pathway in *Drosophila* revealed that Hpo codes for a S-T kinase of the mammalian sterile-20 family of kinases (Harvey et al. 2003; Jia et al. 2003; Pantalacci et al. 2003; Udan et al. 2003; Wu et al. 2003) and can physically associate with the WW domain-containing adaptor protein Sav (Harvey et al. 2003; Jia et al. 2003; Pantalacci et al. 2003; Udan et al. 2003; Wu et al. 2003). Wts is a S-T kinase protein of the DMPK family that associates with another adaptor protein Mob as tumor suppressor (Mats) (Justice et al. 1995; Lai et al. 2005; Shimizu et al. 2008; Wei et al. 2007; Xu et al. 1995). Loss of function of these genes in genetic mosaics revealed strong over-growth phenotype caused by increased cell proliferation and diminished sensitivity to apoptosis. Hyperactivation of the pathway by overexpression of Hpo, Sav, Wts, or Mats leads to formation of smaller organs due to increased apoptosis (Harvey et al. 2003; Pantalacci et al. 2003; Udan et al. 2003; Wei et al. 2007; Wu et al. 2003). Biochemical analysis showed that the Hpo kinase phosphorylates and can physically associate with Sav, Wts, and Mats to form protein complexes in vitro (Wei et al. 2007) (Fig. 2). However, Hpo associates with its cognate adaptor protein Sav

to form the Hpo-Sav complex for efficient activation of the downstream kinase Wts (Huang et al. 2005; Wu et al. 2003). Wts itself associates with Mats to form the downstream Wts-Mats complex of the core kinase cascade of the Hippo pathway (Wei et al. 2007). Association of these adaptor proteins is known to stimulate the catalytic activity of the Hpo and Wts kinases (Dong et al. 2007; Pan 2007; Wei et al. 2007). Moreover, phosphorylation of Mats by the Hpo kinase increases its affinity for the Wts kinase (Dong et al. 2007; Pan 2007, 2010; Wei et al. 2007). Wts is activated by autophosphorylation and phosphorylation by Hpo kinase. Activated Wts associates with Mats (thus Mats cannot simultaneously associate with Hpo and Wts), which acts as a coactivator for the kinase activity of Wts (Dong et al. 2007; Huang et al. 2005; Oh and Irvine 2008, 2009). A major output of the core kinase cascade is to inhibit the growth-promoting activity of Yorkie (Yki), the *Drosophila* homolog of the mammalian YAP oncogene that acts as a transcriptional coactivator (Dong et al. 2007; Huang et al. 2005) (Fig. 2). Yorkie (Yki) was identified via a yeast two-hybrid screen as an interactor of Warts. Overexpression of Yki phenocopies the loss of function of *hpo*, *sav*, *wts*, and *mats* (all genes of the core kinase cascade) and causes overgrowth (Dong et al. 2007; Wei et al. 2007). Loss of function of *yki* results in formation of smaller organs due to induction of cell death (Huang et al. 2005).

Yki activity is regulated by controlling its subcellular localization via phosphorylation-dependent and phosphorylation-independent interactions with the core kinase cascade of the Hippo pathway (Oh and Irvine 2008, 2010; Ren et al. 2010b). Yki associates with Wts, and one mechanism by which the Wts kinase restricts Yki activity is via phosphorylation at Ser168 that creates a 14-3-3 protein-binding site (Goulev et al. 2008; Peng et al. 2009; Ren et al. 2010b; Wu et al. 2008; Zhang et al. 2008b, 2009a). Interestingly, only phosphorylated forms of Yki can associate with 14-3-3 proteins. Yki is phosphorylated at multiple sites (e.g., Ser 111 and S250), which increase Yki activity making it less sensitive to Hpo/Wts-mediated inhibition. These phosphorylation events act in parallel to phospho-Yki/14-3-3-mediated mechanisms and inhibit Yki nuclear localization and activity. It is suggested that nuclear export is required for shuttling Yki to the nucleus in response to Hpo signaling, and binding of 14-3-3 proteins is thought to impede nuclear import and/or promote nuclear export, thereby facilitating nucleocytoplasmic shuttling of target proteins (Brunet et al. 2002; Kumagai and Dunphy 1999). Nuclear transport of Yki depends on its binding with cognate transcription factors as Yki does not have an intrinsic nuclear localization signal (NLS) (Goulev et al. 2008; Zhang et al. 2008a, b) (Fig. 2). Currently, it is unclear if binding of 14-3-3 proteins to Yki prevents its binding with cognate transcription factors or masks the nuclear localization signals or promotes export from the nucleus. Nevertheless, coactivator Yki/YAP is the critical downstream regulatory target of the Hpo kinase cascade, and regulation of its subcellular localization is the primary mechanism by which the Hippo pathway influences target gene expression (Goulev et al. 2008; Huang et al. 2005; Oh and Irvine 2008, 2009, 2010; Oh et al. 2009; Peng et al. 2009; Ren et al. 2010b).

Yki (like Sav) is a WW domain-containing protein and interacts with the PPXY motifs in Wts (Huang et al. 2005; Kango-Singh et al. 2002; Tapon et al. 2002).



Besides Wts, the WW domains of Yki interact with the PPXY motifs present in other components of Hippo signaling pathway like Expanded (Ex), Hpo, WW domain-binding protein 2 (Wpb2), and Myopic (Gilbert et al. 2011) to regulate Hippo signaling via phosphorylation-independent mechanisms (Badouel et al. 2009; Gilbert et al. 2011; Oh et al. 2009; Zhang et al. 2011b). Another protein that acts via its WW domains is Kibra which associates with the PPXY motifs in Ex (and binds Mer in a WW domain-independent manner) (Baumgartner et al. 2010; Genevet et al. 2010). The identification of multiple proteins that act through the interaction between WW domains and PPXY motifs in the Hippo pathway suggests that these motif-specific interactions are important for regulation of Hippo signaling [reviewed in (Sudol 2010; Sudol and Harvey 2010)].

## Yki Activity and Regulation of Expression of Target Genes


Hyperactivation of the pathway, for example, by overexpression of Hpo, leads to phosphorylation and activation of Hpo and Wts with the help of adaptor proteins Sav and Mats. Wts, in turn, phosphorylates the transcriptional coactivator Yki, which associates with 14-3-3 proteins and remains sequestered in the cytoplasm (Dong et al. 2007; Huang et al. 2005; Oh and Irvine 2008; Oh et al. 2009; Ren et al. 2010b). Analysis of adult and imaginal disc phenotypes reveals that overexpression of Hpo results in induction of ectopic apoptosis early in development in imaginal disc cells due to induction of caspase-dependent cell death (Hamaratoglu et al. 2006; Harvey et al. 2003; Udan et al. 2003; Verghese et al. 2012a). In mammalian cells, activation of MST1/2 and hyper-phosphorylation of YAP2 by MST2 and LATS1 kinase lead to activation of cell death. MST1/2 are known targets of caspases. Furthermore, YAP1/2 are known to interact with p73 via a PDZ domain in YAP and induce apoptotic target genes (Bertini et al. 2009; Sudol 2010; Sudol and Harvey 2010). However, these mechanisms of regulating apoptosis may not be conserved in flies because the site for caspase cleavage is not conserved in *Drosophila* Hpo (Wu et al. 2003), and *Drosophila* Yki does not have the conserved PDZ domain (Sudol and Harvey 2010). Nevertheless, Hpo overexpression in flies induces apoptosis through an alternate mechanism that does not involve caspase cleavage or p73. Recently, it was shown that the effector caspase Dronc (*Drosophila* homolog of mammalian caspase-9) is induced in conditions when Hippo pathway is hyperactivated. Further, using reporter genes, it was shown that dronc transcription is induced during gain-of-function and downregulated during loss-of-function conditions of the Hippo pathway, suggesting that *dronc* is a transcriptional target of the Hippo pathway (Verghese et al. 2012a). However, the molecular mechanism by which Yki interacts with Dronc remains unclear. Both phosphorylation-dependent (e.g., with 14-3-3 by phosphorylation-dependent mechanisms) and phosphorylation-independent mechanisms (binding with Hpo, Wts, or Ex) result in cytoplasmic retention of Yki in multiple protein complexes. Thus, the possibility remains that hyperactivation of Hippo pathway releases Yki from one or more cytoplasmic com-

plexes to allow its binding to transcription factors and shuttle into the nucleus to induce *dronc* transcription. Alternatively, hyperactivation of the Hippo pathway involves a transcriptional repressor that acts together with or independent of Yki to control *dronc* expression. Thus, although it is clear that hyperactivation of the Hippo pathway leads to induction of apoptosis, the molecular mechanisms underlying this process are yet unidentified.

When the pathway is downregulated, the genes of the core kinase cascade act as tumor suppressors by suppressing the growth-promoting activity of Yki (Fig. 2). Under these conditions, Yorkie can partner with transcription factors like the TEAD family protein, Scalloped (Sd), and enter the nucleus and cause transcription of target genes which regulate cell proliferation and apoptosis. Sd was identified as the transcriptional factor of the pathway via yeast two-hybrid screen and in vitro Yki activity assays (luciferase assay) (Goulev et al. 2008; Wu et al. 2008; Zhang et al. 2008b). Sd is required for wing development (Campbell et al. 1992; Liu et al. 2000), whereas Yki is required for regulating growth of all imaginal disc cells. Other transcription factors that bind Yki to regulate growth via Hippo signaling have since been discovered. These include Mothers Against Dpp (Mad) (Alarcon et al. 2009; Oh and Irvine 2010; Peng et al. 2009), Homothorax (Hth), and Teashirt (Tsh) (Peng et al. 2009). Mad is a known transcription factor within the Dpp/TGF $\beta$  signaling pathway, and Mad and Hth were shown to control the activity of the *bantam* miRNA (Alarcon et al. 2009; Peng et al. 2009). Mad, Hth, and Tsh are known transcription factors that respond to other signals and are required for patterning of imaginal discs during development.

Yki activity is controlled by the upstream signals (Grusche et al. 2010; Oh and Irvine 2010) (Fig. 2). A large number of target genes have been identified over the past decade, which include the cell cycle regulators E2F1 and *cyclins E, A, B, and D*; the growth promoter *Myc* and cell survival-promoting miRNA *bantam*; genes regulating cell death like the *Drosophila inhibitor of apoptosis diap1, hid, and dronc*; and cytoskeletal proteins like F-actin, which drive cell proliferation and cell survival (Fig. 3) (Goulev et al. 2008; Harvey et al. 2003; Huang et al. 2005; Jia et al. 2003; Kango-Singh et al. 2002; Neto-Silva et al. 2010; Nolo et al. 2006; Pantalacci et al. 2003; Peng et al. 2009; Tapon et al. 2002; Thompson and Cohen 2006; Udan et al. 2003; Wu et al. 2003, 2008; Zhang et al. 2008a; Ziosi et al. 2010). Yki also controls the expression of several upstream components of the Hippo pathway like Ex, Mer, Kibra, Crumbs (Crb) and Four-jointed (Fjose et al. 1984) by a negative feedback loop (Cho et al. 2006; Fjose et al. 1984; Genevet et al. 2009, 2010; Hamaratoglu et al. 2006). Recently, Yki was shown to affect the expression of components of other signaling pathways, such as ligands for the Notch, Wnt, EGFR, and Jak-Stat pathways (Cho et al. 2006; Karpowicz et al. 2010; Ren et al. 2010a; Shaw et al. 2010; Staley and Irvine 2010, 2012; Zhang et al. 2009a). These interactions suggest that Hippo pathway interacts with the major signal transduction pathways, and these points of contact between different pathways may play an important role in controlling correct tissue sizes and maintaining homeostasis (Fig. 3).

Genetic and biochemical studies thus provide a basic premise for how Yki activity is modulated when Hippo signaling is down- or upregulated (Halder and Johnson



**Fig. 3** Hippo pathway target genes regulate cell proliferation and apoptosis: (a–d, #6887) *GMRGAL4 UASHPo* third instar eye-antennal imaginal disc showing effect on target proteins upon pathway hyperactivation in the GMR domain. (a) Cyc E is downregulated, (b) DIAP-1 levels remain unaffected, and (c) Drice is activated (Drice is the homolog of *Drosophila* Caspase3\* and is a readout of active Dronc). (d) Dronc is upregulated in the GMR domain upon Hpo overexpression. (e) Loss-of-function clones of *ft* (GFP negative) made with *yw* *hsFLP*; *UbiGFP* [*hsFLP*; *FRT40A ft<sup>td</sup>/FRT40A ubiGFP*] show upregulation of Cyc E in the mutant cells. This effect is very strong in the region of the second mitotic wave (SMW). (f–h) *GMRGAL4 UASYki* third instar eye-antennal imaginal discs. (f) DIAP-1 is upregulated, (g) Caspase3\* staining is not observed, and (h) Dronc is downregulated in the GMR domain consistent with overproliferation and no apoptosis

2011; Harvey and Hariharan 2012; Schroeder and Halder 2012; Staley and Irvine 2012). Studies in imaginal discs and other cell types like intestinal stem cells and fat cells revealed that Hippo signaling is needed in all cell types to regulate growth and that the activity of the pathway is modulated to achieve tissue homeostasis (Halder et al. 2012; Halder and Johnson 2011; Harvey and Hariharan 2012; Tumaneng et al. 2012a; Zhao et al. 2008a, 2010a). Whether Hippo signaling pathway is regulated by other global instructive signals (e.g., morphogen gradients) or if the pathway is constitutively active remains unknown. However, several inputs that communicate a growth regulatory signal to the core kinase cascade have been identified. We will discuss the key inputs and their connection to the core kinase cascade in the following sections.

## Upstream Regulators of the Hippo Pathway

Since the discovery of the core kinase cascade, several upstream regulators of the Hippo pathway were identified (Table 1). These discoveries highlighted two remarkable properties of the Hippo pathway—one, that the Hippo pathway is a signaling network with multiple points of signal integration rather than a linear system of

**Table 1** Hippo pathway components and their biological roles

	Gene name, <i>symbol</i> [Chr]	Nature of protein	Role	References
Upstream regulators	Crumbs <i>Crb</i> [3]	Protein kinase C binding	Organization of adherens junction, establishment of cell polarity, photoreceptor and rhabdome development	Fan et al. (2003), Pichaud and Desplan (2001), Tepass et al. (1990)
	Expanded <i>ex</i> [2]	Protein binding	Compound eye, photoreceptor cell differentiation, negative regulation of Hippo signaling cascade	Maitra et al. (2006), Pellock et al. (2007), Badouel et al. (2009), McCartney et al. (2000)
	Merlin <i>Mer</i> [1]	Protein binding	Regulation of programmed cell death, negative regulator of Hippo signaling	Pellock et al. (2007), Hamaratoglu et al. (2006)
	Kibra <i>Kibra</i> [3]	Protein binding	Compound eye morphogenesis, regulation of Hippo signaling cascade	Ling et al. (2010), Genevet et al. (2010), Yu et al. (2010), Baumgartner et al. (2010)
Fat branch	Fat <i>ft</i> [2]	Cell adhesion molecule binding	Establishment of planar polarity, negative regulation of growth, imaginal disc growth	Yang et al. (2002), Mao et al. (2006), Torok et al. (1993), Garoia et al. (2000), Matakatsu and Blair (2006)
	Low fat <i>lft</i> [2]	Protein binding	Wing morphogenesis	Mao et al. (2009)
	Dachs <i>D</i> [2]	ATPase activity (predicted nature)	Establishment of ommatidial planar polarity, positive regulation of growth	Mao et al. (2006)
	Dachsous <i>Ds</i> [2]	Cell adhesion molecule binding	Eye morphogenesis, establishment of cell polarity, cell proliferation	Baena-Lopez et al. (2005), Clark et al. (1995)
	Four-jointed <i>Fj</i> [2]	Wnt-protein binding: protein kinase activity	Imaginal disc growth, establishment of planar polarity	Villano and Katz (1995), Bosveld et al. (2012)
	Scribbled <i>Scrib</i> [2]	Protein binding	Establishment of ommatidial planar polarity, negative regulation of imaginal disc growth	Courbard et al. (2009), Zeitler et al. (2004), Verghese et al. (2012)
	Zyxin <i>Zyx</i> [4]	Protein binding	Positive regulation of imaginal disc growth	Rauskolb et al. (2011)
	Approximated <i>App</i> [3]	Protein-cysteine S-palmitoyltransferase activity (predicted nature)	Establishment of body hair or bristle planar orientation	Matakatsu and Blair (2008)
	Discs overgrown <i>Deco</i> [3]	Kinase activity	Establishment of ommatidial planar polarity, positive regulation of cell growth	Strutt et al. (2006), Klein et al. (2006), Guan et al. (2007)

Core kinase cascade	Warts Wts [3]	Protein binding, kinase activity	Negative regulation of cell proliferation, R8 cell fate specification	Justice et al. (1995), Mikeladze-Dvali et al. (2005)
	Mob as tumor suppressor Mats [3]	Protein binding	Cell proliferation	Lai et al. (2005)
	Hippo Hpo [2]	Protein binding; serine/threonine kinase activity	Negative regulation of cell proliferation, R8 cell fate specification	Udan et al. (2003), Mikeladze-Dvali et al. (2005)
	Salvador Sav [3]	Protein binding	Negative regulation of cell proliferation, R8 cell fate specification	Kango-Singh et al. (2002), Mikeladze-Dvali et al. (2005)
	Ajuba Jub [1]	Ligand-dependent nuclear receptor binding	Positive regulation of organ growth	Das Thakur et al. (2010)
	Tao Tao [1]	Serine/threonine kinase activity	Negative regulation of organ growth	Poon et al. (2011)
	Echinoid Ed [2]	Protein binding	Negative regulation of Hippo signaling cascade	Yue et al. (2012)
	Pez Pez [2]	Protein tyrosine phosphatase activity	Negative regulation of Hippo signaling cascade	Poembacher et al. (2012)
	d-STRIPAK PP2A Pp2A-29B [2]	Serine/threonine phosphatase activity	Centrosome organization	Dobbelaere et al. (2008)
	Ras association family member Rassf [3]	Protein binding	Negative regulation of signal transduction	Polesello et al. (2006)
Other regulators	Par-6 Par-6 [1]	Protein binding	Cell adhesion	Kiger et al. (2003)
	Atypical protein kinase C a-PKC [2]	Protein binding; serine/threonine kinase activity	Compound eye retinal cell programmed cell death, establishment of epithelial cell planar polarity	Ogawa et al. (2009), Kaplan et al. (2011)
	Stardust Sdt [1]	Protein binding	Zonula adherens assembly	Nam and Choi (2003), Bachmann et al. (2001)
				(continued)

**Table 1** (continued)

	Gene name, <i>symbol</i> [Chr]	Nature of protein	Role	References
	Lethal 2 giant larvae L2gl [2]	Myosin II binding; myosin binding	Cell competition in a multicellular organism, establishment of epithelial cell planar polarity	Tamori et al. (2010), Kaplan and Tolwinski (2010)
	Myopic Mop [3]	Protein tyrosine phosphatase activity	Regulation of growth	Gilbert et al. (2011)
Transcription factors/ coactivators	Patj dPatj [3]	Protein kinase C binding	Adherens junction organization	Nam and Choi (2006)
	Yorkie Yki [2]	Protein binding; transcription coactivator activity	Cell competition in a multicellular organism, cell proliferation	Ziosi et al. (2010), Huang et al. (2005), Thompson and Cohen (2006)
	Scalloped Sd [1]	Transcription factor binding	Compound eye morphogenesis	Garg et al. (2007)
	Homothorax Hth [3]	Protein binding; transcription factor	Compound eye photoreceptor fate determination	Wernet et al. (2003)
	Teashirt Tsh [2]	Transcription factor activity	Eye-antennal disc development	Singh et al. (2004)
	Wpb2 Wbp2 [3]	Transcription factor binding	Positive regulation of imaginal disc growth	Zhang et al. (2011a, b)
	Mothers against dpp Mad [2]	Transcription factor activity	Compound eye morphogenesis, negative regulation of gene expression	Cordero et al. (2007), Anderson et al. (2006)

epistatic genes (Fig. 2), and two, the interactions between various protein complexes (at the signal integration points) may play a decisive role in shaping the outcome, i.e., Yki activity levels. Although our understanding of the network is incomplete in both these areas, it is clear that signaling interactions within this pathway are shaped by several distinct inputs.

### ***Fat Signaling and the Hippo Pathway***

*fat* (*ft*) alleles were spontaneous mutations first described by Mohr (1923, 1929). Subsequent analysis of mutations in the *ft* locus revealed both viable and lethal alleles, of which the null alleles are larval lethal and show hyperplastic overgrowth of imaginal discs thereby acting as tumor suppressor genes (Bryant et al. 1988). Molecular cloning of *ft* revealed that it codes for a transmembrane protein, which is an atypical cadherin (Mahoney et al. 1991). Loss of *ft* affects two distinct aspects of imaginal disc growth and development, restriction of cell proliferation and generation of correctly oriented cells within the epithelial sheet, phenotypes that were mapped to two distinct signaling pathways—the Hippo and the planar cell polarity (PCP) pathway (see (Cho 2006 #659) (Brittle et al. 2010; Matakatsu and Blair 2006, 2008, 2012)). Ft is ubiquitously expressed; however, its functions are regulated by two genes, Dachous (Ds) and Fj, which are expressed in gradients in developing tissues (Matakatsu and Blair 2004; Reddy and Irvine 2008). Ds is another protocadherin in flies that acts as the ligand for Ft for both the Hippo and PCP pathways [reviewed in (Thomas and Strutt 2012)]. Fj is a Golgi-localized kinase that phosphorylates the extracellular cadherin domains of Ft and Ds to promote their binding (Ishikawa et al. 2008; Simon et al. 2010). Phosphorylation of Fat by Fj increases its affinity to Ds, while phosphorylation of Ds reduces its affinity to Ft. One way in which Fat regulates growth and PCP is based on the slope and vector of the Ds and Fj gradients (Halder and Johnson 2011; Willecke et al. 2008; Zecca and Struhl 2010) (Fig. 2).

Several years after Ft was discovered, it was realized that the growth regulatory functions of Fat were tied to the Hippo pathway (Bennett and Harvey 2006; Cho et al. 2006; Silva et al. 2006; Willecke et al. 2006). Loss of *ft* in mutant clones phenocopied the loss-of-function phenotypes of genes within the core kinase cascade of the Hippo pathway. Imaginal discs containing somatic clones of *ft* mutant cells continued to proliferate when normal cells had stopped, thereby forming large overgrown discs. Transcriptional targets of Hippo pathway are induced within the *ft* mutant cells, a phenotype similar to loss of function of positive regulators of Hippo pathway (e.g., *wts*, *Hpo*, *sav*, *mats*). Ft affects the levels and localization of Hippo pathway components, including Wts, Ex, and Yki (Bennett and Harvey 2006; Cho et al. 2006; Oh and Irvine 2008; Silva et al. 2006; Tyler and Baker 2007; Willecke et al. 2006). Ft influences Hippo signaling independent of other upstream regulators like *expanded*, *merlin* (*mer*), and *kibra* which form a heteromeric complex (Ex-Mer-Kibra) and other genes like the Tao-1 kinase (Boggiano et al. 2011; Poon et al.

2011) that act upstream of Hpo (Boggiano and Fehon 2012). However, several other genes were recently identified that specifically act downstream of Ft and integrate with the Hippo pathway by influencing the activity of the downstream kinase Wts. Thus, the Fat branch of the Hippo pathway has emerged that independently influences Wts activity and tissue growth (Halder and Johnson 2011; Kango-Singh and Singh 2009; Reddy and Irvine 2008; Staley and Irvine 2012) (Fig. 2).

Several components of the Ft branch influence the intracellular domain of Ft—the region critical for transducing the signal within cells (Fig. 2). These include the *Drosophila* Discs overgrown (Dco, #6929), a homolog of casein kinase I, which phosphorylates the Ft intracellular cytoplasmic domain in a Ds-dependent manner (Cho et al. 2006; Feng and Irvine 2009; Sopko et al. 2009), and the unconventional myosin Dachs (D) (Cho et al. 2006; Cho and Irvine 2004; Mao et al. 2006). Loss of function of *dco*<sup>3</sup>, a hypomorphic allele, in homozygous discs and in somatic clones results in tissue overgrowth and shows elevated levels of Fj and Diap-1 (Bryant and Schmidt 1990; Feng and Irvine 2009; Guan et al. 2007). Dco binds to the cytoplasmic domain of Fat, and in *dco* mutants, Fat intracellular domains fail to phosphorylate. Ds enriches availability of Fat at the point of cell contacts by forming *cis*-dimers with Fat. This promotes the transphosphorylation of Fat by Dco. Lowfat is a novel protein that interacts with the intracellular domains of Fat and Ds and stabilizes the Fat-Ds interaction (Mao et al.). Lowfat was identified in a genome-wide yeast two-hybrid screen as a Fat- and Ds-interacting protein (Mao et al. 2006, 2009). In addition, the palmitoyltransferase Approximated (App) acts downstream of Ft, and Ft regulates the localization of D to the membrane through APP (Matakatsu and Blair 2008). Recently, the apical-basal polarity gene *scribble* (*scrib*) (Verghese et al. 2012b) and the LIM-domain protein *zyxin 102* (*zyx*) (Rauskolb et al. 2011) were shown to act in the Fat branch of Hippo signaling pathway (Bennett and Harvey 2006; Cho et al. 2006; Meignin et al. 2007; Polesello and Tapon 2007; Reddy et al. 2010; Silva et al. 2006; Willecke et al. 2006).

The differences in Ds and Fj expression between neighboring cells stimulate Yki activity, whereas the vector property of the gradients affects PCP signaling. Localization of D to the membrane is regulated by Fj, Ds, and Ft (Cho et al. 2006; Mao et al. 2006; Rogulja et al. 2008; Willecke et al. 2008). D controls Yki activity by two alternative mechanisms: the first involves posttranslational effects of Ft on Wts, and the second involves the localization of Ex to the subapical membrane (Bennett and Harvey 2006). The apical-basal polarity gene *scrib* and the atypical myosin D are responsible for partitioning the growth regulatory signal from Ft to downstream genes. Genetic epistasis experiments placed Ft upstream of D and the apical regulator of the pathway—Expanded (Ex) (Cho et al. 2006; Mao et al. 2006; Verghese et al. 2012b). D can reverse the effects of loss of *ft* on growth and expression of Fat target genes like Wg, Serrate, and Fj (Mao et al. 2006). *Scrib* was also placed upstream of D and Ex and downstream of Ft based on genetic epistasis experiments (Verghese et al. 2012b) (Fig. 2). When Ft is inactive, D is regulated by Approximated (App) (Matakatsu and Blair 2008). App posttranscriptionally modifies D and affects its localization at the apical cell cortex. Hence, App functions in the Hippo pathway by affecting the availability of D at the apical cell cortex. When



Ft is activated, D is released from App and binds to Zyxin (Zyx), which in turn interacts with Wts and stabilizes Wts activity (Rauskolb et al. 2011). Zyx binds to D; genetic epistasis experiments placed Zyx downstream of Ft and Dco and upstream of Wts (Feng and Irvine 2007, 2009; Rauskolb et al. 2011). Thus, influencing Wts stability is a primary mechanism by which Ft controls growth via Hippo signaling (Fig. 2). However, the other input via Ex remains less clear although there is clearly an input from Ft to Ex that also contributes to the Fat-branch-related phenotypes and regulation of the Hippo signaling pathway. Whether Fat signaling simultaneously signals through Ex (and the core kinase cascade) and D or the signals downstream of Ft are partitioned to allow maximum and more efficient signal transduction to the core kinase cascade remains unknown. Currently, the possibility that certain extracellular signals preferentially transmit the signal to Ex or D downstream of Ft has not been addressed.

### ***Apical Membrane Proteins of the Hippo Pathway***

Over the last 5 years, it has become clear that membrane-localized proteins are an intrinsic part of the Hippo signaling pathway (Genevet and Tapon 2011; Grusche et al. 2011; Halder et al. 2012; Schroeder and Halder 2012) (Table 1). Among these are the cell polarity proteins and proteins required for maintaining the cytoskeleton (Fig. 2). The FERM domain-containing adaptor proteins Ex and Merlin (Mer) were among the earliest Hippo pathway components that were known to localize to the apical membrane (Hamaratoglu et al. 2006; McCartney et al. 2000). Ex and Mer act upstream of the Hpo kinase and regulate pathway activation (Hamaratoglu et al. 2006). Loss of *mer* and *ex* together in somatic clones caused dramatic overproliferation of cells leading to overgrowths. These effects were synergistic because loss of function of *ex* or *mer* alone does not cause similar defects. These genes function together to control proliferation by regulating expression of transcriptional targets of Hippo pathway (e.g., cyclin E and DIAP1). Expanded can also regulate the pathway by independently interacting with Yki and sequestering it in the cytoplasm (Badouel et al. 2009; Oh et al. 2009).

Another protein that binds Ex and Mer and acts upstream of Hpo is the WW and C2 domain-containing adaptor protein Kibra. Ex, Mer, and Kibra form a complex at the apical membrane in epithelial cells, which then activates the downstream core kinase cascade (Baumgartner et al. 2010; Cho et al. 2006; Genevet et al. 2010; Hamaratoglu et al. 2006; Pellock et al. 2007; Tyler and Baker 2007) (Fig. 2). Kibra was identified via a genome-wide screen in *Drosophila* and in S2 cells for candidates that modified Yki activity (Baumgartner et al. 2010; Genevet et al. 2010; Yu et al. 2010). Genetic epistasis experiments placed Kibra upstream of Hpo and Yorkie. Kibra affects the phosphorylation of Hpo and Yorkie. Kibra acts synergistically with Ex and Mer to regulate Wts phosphorylation, and Kibra binds to Sav and Hpo in a Sav-dependent manner (Baumgartner et al. 2010; Genevet et al. 2010).

Cell polarity genes have been well characterized in flies and mammalian model systems, and recent studies reveal a role for cell polarity genes in the regulation of Hippo signaling (Table 1, Fig. 2) (Genevet and Tapon 2011; Grusche et al. 2010; Grzeschik et al. 2007, 2010a; b; Schroeder and Halder 2012). Crumbs (Crb), a transmembrane protein, is the upstream regulator that regulates Ex activity (Chen et al. 2010; Ling et al. 2010; Robinson et al. 2010). Crb is required for proper localization of Ex. Crb regulates Yki activity by interacting with Expanded (Chen et al. 2010; Grzeschik et al. 2010a; Robinson et al. 2010). Crb was found through a genetic screen, and loss and gain of function of Crb cause overgrowth of tissues and upregulation of the Hippo pathway target genes. Echinoid (Ed) is another upstream regulator of the Hippo pathway that like kibra interacts with both Ex and Yki (Baumgartner et al. 2010; Genevet et al. 2010; Yu et al. 2010; Yue et al. 2012). Cells mutant for *ed* cause mislocalization of Sav from the subapical membrane without affecting Ex or Mer localization. Ed also interacts physically with Hpo, Ex, Mer, and Kibra (Yue et al. 2012).

F-actin acts as an upstream regulator of the Hippo pathway (Fig. 2). Increased levels of F-actin inhibit the pathway, and activation of Hippo pathway inhibits F-actin accumulation (Fernandez et al. 2011; Richardson 2011; Sansores-Garcia et al. 2011). Tao-1 phosphorylates Hpo and acts upstream of Hpo at T195 (Boggiano and Fehon 2012; Boggiano et al. 2011; Poon et al. 2011). RNAi knockdown of Kibra, Ex, and Mer (KEM) resulted in a significant decrease of endogenous Hpo protein in the membrane fraction (Boggiano and Fehon 2012; Boggiano et al. 2011; Poon et al. 2011). Thus, the apical proteins regulate Hpo at least in part by bringing the latter to the membrane, where Hpo may be activated via mechanisms yet to be determined.

## Negative Regulators of the Hippo Pathway

Several members of the Hippo pathway were identified based on their effects on tissue growth, and the loss-of-function phenotypes of these components showed dramatic outgrowths and benign lesions in fly epithelia (Table 1). It was clear that additional components that keep this pathway in check (e.g., phosphatases or kinase inhibitors) must exist, as Hippo activity would need to be modulated both positively and negatively for maintaining tissue homeostasis. Thus, the search for negative regulators began that yielded many important and critical regulators of the Hippo pathway. Among the first genes identified in this category was the Ras Association Family (RASSF) gene, *dRASSF1* (Polesello et al. 2006) (Fig. 2). The dRASSF protein negatively regulates the pathway by inhibiting the phosphorylation of Hpo, thus interrupting the Hpo kinase from signaling to the downstream kinase Wts (Polesello et al. 2006; Scheel and Hofmann 2003). Other inhibitors that act by dephosphorylating Hpo are the phosphatases—striatin-interacting phosphatase (STRIPAK) and protein phosphatase 2A (PP2A) (Ribeiro et al. 2010) (Fig. 2). A second mechanism of inhibition of Yki activity was identified by the *Drosophila* Ajuba family gene,

*djub* (Das Thakur et al. 2010) (Fig. 2). Loss of *djub* in mutant clones in imaginal discs caused reduced proliferation and increased apoptosis, akin to *yki* mutant clones. Genetic interaction studies showed that *djub* acts downstream of Hpo but upstream of Yki and Wts (Das Thakur et al. 2010). Furthermore, Djub can physically associate with Wts and Sav and influence the signaling activity of Yki. Thus, *djub* negatively regulates the Hippo signaling by interfering with Yki phosphorylation and its subcellular localization (Das Thakur et al. 2010). Recently, another negative regulator, *myopic* (Bonner and Boulianne 2011), was identified in a genetic screen for conditional growth suppressors (Gilbert et al. 2011) (Fig. 2). *mop* encodes the *Drosophila* homolog of human *His domain protein tyrosine phosphatase* gene (*HD-PTP* or *PTPN23*) (Toyooka et al. 2000). *mop* mutant cells show overgrowth phenotypes due to a block in cell death. This growth is accompanied by upregulation of a subset of Yki transcriptional targets but not the antiapoptotic gene *diap1*. *mop* interacts genetically with *yki* and acts downstream of *wts* but at the level of *ex* and *yki*. Myopic PPxY motifs bind conserved residues in the WW domains of the transcriptional coactivator Yorkie, and Myopic colocalizes with Yorkie at endosomes (Gilbert et al. 2011). Thus, several negative regulators of the Hippo pathway are now known; however, much remains unknown about their mechanism of action and their influence on growth regulation during development (Tables 1 and 2).

## Hippo Pathway Cross-Talks with Other Pathways

Hippo pathway is known to interact with other pathways to regulate growth (Table 2). In mice it has been shown that Mst2 interacts with Raf-1 of the ERK/MAPK pathway (Graves et al. 1998). Raf-1 inhibits dimerization of Mst2 and recruits a phosphatase to dephosphorylate Mst2, thereby inactivating it, a function independent of

**Table 2** Pathways known to interact with the Hippo network


Pathway interactions	Responses	References
JNK pathway	Cell competition, compensatory proliferation, regeneration, cytoskeletal integrity, tumorigenesis	Chen et al. (2012), Sun et al. (2011), Densham et al. (2009), Enomoto et al. (2012)
Wingless pathway	Growth control	Verelas et al. (2010)
EGFR pathway	Growth control	Herranz et al. (2012)
Decapentaplegic pathway	Growth control	Rogulja et al. (2008)
Hedgehog pathway	Growth control, neuronal differentiation	Kagey et al. (2012), Lin et al. (2012)
Notch pathway	Neural stem cell maintenance, polar cell fate during oogenesis, cell differentiation, proliferation	Li et al. (2012), Chen et al. (2011), Yu et al. (2008)
TSC-TOR pathway		Latest paper from Tapon

the MAPK pathway (O'Neill and Kolch 2005). More recently, many points of intersection between Hippo and other signaling pathways have come to light. For example, in the last 5 years, Hippo pathway was shown to interact with JNK pathway to regulate compensatory proliferation, regeneration, and tumor progression (Chen et al. 2012; Doggett et al. 2011; Grzeschik et al. 2010a; Staley and Irvine 2010; Sun and Irvine 2010, 2011; Tyler et al. 2007; Varelas et al. 2010a). Furthermore, Hippo pathway interacts with Wingless/Wnt pathways in flies and mammals (Varelas et al. 2010a, b). Hippo pathway restricts Wnt/beta-catenin signaling by promoting an interaction between TAZ and DVL in the cytoplasm. TAZ inhibits the CK1delta/epsilon-mediated phosphorylation of DVL, thereby inhibiting Wnt/beta-catenin signaling (Azzolin et al. 2012; Tsai et al. 2012; Varelas et al. 2010a). In *Drosophila*, Hippo signaling modulates Wg target gene expression (Varelas et al. 2010a, b). More connections of Hippo signaling with pathways that control morphogenetic patterning and growth have been uncovered which include the discovery of the regulation of TGF beta/SMAD complexes by YAP/TAZ in mammalian models and Yki in flies (Chan et al. 2011; Meignin et al. 2007; Polesello and Tapon 2007; Rogulja et al. 2008; Sudol and Harvey 2010; Varelas et al. 2010b). Dpp (Decapentaplegic) signaling interacts with D to maintain Fj and Ds gradient in order to regulate proliferation in the wing (Rogulja et al. 2008). Hippo pathway also intersects the PI3K/TOR pathway via multiple interactions (Bellosta and Gallant 2010; Collak et al. 2012; Karni et al. 2008; Kim et al. 2010; Mills et al. 2008; Sekido 2008; Strassburger et al. 2012; Tumaneng et al. 2012a, b; Wehr et al. 2013), with G-protein-coupled receptor (GPCR) signaling (Yu et al. 2012) and receptor tyrosine kinase signaling (Gadd et al. 2012; Garami et al. 2003). In fact, the web of interactions has grown exponentially over the last few years such that oftentimes the Hippo pathway is sometimes referred to as a network or superhighway (Barry and Camargo 2013) (Fig. 4).

## Mammalian Hippo Pathway

Hippo pathway is responsible for regulating organ size and is involved in regeneration (Bertini et al. 2009; Hiemer and Varelas 2013; Hong and Guan 2012; Liu et al. 2012a). The core kinase pathway is highly conserved in mammals (Hong and Guan 2012; Liu et al. 2012a; Zhao et al. 2008a). In vertebrate models, the core kinase cascade consists of Mst1/2 (Hpo homolog) and Lats1/2 (Wts homolog) along with their adaptor proteins WW45 (Sav) and MOB1 (Mats homolog), which control growth by regulating phosphorylation of YAP (Yki homolog) (Hong and Guan 2012; Liu et al. 2012a; Zhao et al. 2008a). Ft1-4 (Ft homolog), Dchs1-2 (Ds homolog), and Fjx1 (Fj homolog) are known to regulate planar cell polarity; however, their connection to other Hippo pathway components still needs to be explored (Brittle et al. 2010; Hiemer and Varelas 2013; Skouloudaki et al. 2009; Sopko et al. 2009; Zhao et al. 2007).

The other downstream components like Dco and Lowfat homolog have not been shown yet to function within the Hippo pathway (Sopko et al. 2009; Zhang et al.



**Fig. 4** Hippo pathway is linked to many biological and developmental processes. Hippo signaling has been shown to participate in generating myriad cellular responses that are aimed at attaining tissue homeostasis in addition to regulating organ size. Thus, the role of Hippo signaling is implicated not only during organ development but also in differentiated tissues. Further, tumorigenesis has also been attributed to dysregulation of Hippo signaling placing it in the global network of regulatory mechanisms required for proper growth

2008a, 2011a; Zhao et al. 2010a). However, Dco homolog CK1δ/ε has been shown to be involved in YAP/TAZ degradation (Zhao et al. 2010b). Neurofibromatosis type II (NF2), the Mer homolog, is the most extensively studied upstream regulator in mammals (Sekido 2011; Striedinger et al. 2008; Zhang et al. 2009b; Zhao et al. 2007). NF2 interacts with CD44 and adherens junction to relay the signal downstream to other Hippo pathway components during contact inhibition (Li et al. 2012; Morrison et al. 2001; Zhao et al. 2007). KIBRA is known to interact with Lats2 to promote its phosphorylation (Zhang et al. 2012). It also protects Lats2 from proteosomal degradation by preventing its ubiquitination. KIBRA is also the transcriptional target of Hippo pathway (Angus et al. 2012; Ishiuchi and Takeichi 2012; Visser-Grieve et al. 2012; Xiao et al. 2011). Angiomotin family (AMOT) interacts with its PPxY domain to YAP WW domain and TAZ PDZ domain independent of the upstream components. This interaction inhibits the activity of YAP/TAZ (Chan et al. 2011; Paramasivam et al. 2011; Skouloudaki and Walz 2012; Wang et al. 2009, 2012a; Zhao et al. 2011a). Ex1/FRMD6/Willin (Ex homolog) interacts with upstream Hippo pathway components like Mer (Angus et al. 2012; Ishiuchi and Takeichi 2012; Visser-Grieve et al. 2012). Crb interacts with YAP/TAZ and promotes its phosphorylation, which is

dependent on cell density and at the same time inhibits TGF- $\beta$  SMAD pathway (Varelas et al. 2010b). Unlike *Drosophila* RASSF1, mammalian RASSF homologs activate MST1/2 (Avruch et al. 2012; Guo et al. 2007; Hergovich 2012; Hwang et al. 2007; Kim et al. 2003; Polesello et al. 2006; Ribeiro et al. 2010; Schagdarsurengin et al. 2010; Seidel et al. 2007).

NPHP4, a known cilia-associated protein that is mutated in the severe degenerative renal disease nephronophthisis, acts as a potent negative regulator of mammalian Hippo signaling (Habbig et al. 2011, 2012). NPHP4 directly interacted with the kinase Lats1 and inhibited Lats1-mediated phosphorylation of the Yes-associated protein (YAP) and TAZ (transcriptional coactivator with PDZ-binding domain), leading to derepression of these protooncogenic transcriptional regulators. Moreover, NPHP4 induced release from 14-3-3 binding and nuclear translocation of YAP and TAZ, promoting TEA domain (TEAD)/TAZ/YAP-dependent transcriptional activity (Habbig et al. 2011). ITCH interacts with LATS to negatively regulate its stability (Ho et al. 2011; Salah et al. 2011; Wang et al. 2012a).  $\alpha$ -Catenin interacts with YAP and affects its stability by stabilizing the YAP/14-3-3 complex to restrict YAP activity and by preventing PP2A to interact with YAP (Azzolin et al. 2012; Schlegelmilch et al. 2011; Silvis et al. 2011; Tsai et al. 2012) (Varelas 2010 #1830; Konsavage 2013 #3450; Mauviel et al. 2012 #3755). Zona occludens-2 (ZO-2) promotes the pro-apoptotic function of YAP (Oka et al. 2010). The ASPP (apoptosis-stimulating protein of p53) family of proteins can function in the nucleus to modulate the transcriptional activity of p53, with ASPP1 and ASPP2 contributing to the expression of apoptotic target genes (Vigneron et al. 2010). ASPP increases YAP/TAZ nuclear availability by preventing LATS interaction with YAP/TAZ (Vigneron et al.). Similarly, PP1A interacts with ASPP1 to dephosphorylate TAZ leading to increased TAZ nuclear availability (Liu et al. 2010, 2011).

In mammalian cell lines, E-cadherin acts as an upstream regulator of the pathway, which activates the pathway in response to contact inhibition. YAP and TAZ interact with several transcriptional factors. YAP/TAZ interacts with TEAD1/4 and Runx2. TAZ interacts with thyroid transcription factor-1, peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ), Tbx5, Pax3, and Smad2/3/4. Yap interacts with p73 to mediate its pro-apoptotic functions. Various target genes are as follows: *CTGF*, *AREG*, *BIRC5-2*, and *GLI-2* (Liu et al. 2012b; Zhao et al. 2008a, 2010a). YAP1 interacts with sonic hedgehog pathway to promote the proliferation of cerebellar granule neuron precursors (CGNPs). TAZ inhibits Wnt signaling by inhibiting the phosphorylation of dishevelled (DVL) by CK1 $\delta$ e. YAP/TAZ has also been shown to interact with SMAD to regulate tumorigenesis (Zhang et al. 2011a; Zhao et al. 2011b).

## The Insulin Receptor Signaling Pathway: Regulation of Cell Size

The pin-head screens showed a large number of mutations that primarily caused decreased growth due to formation of smaller cells (Oldham et al. 2000a; Stocker and Hafen 2000). These mutants were subsequently categorized into two well-

studied signaling pathways: the insulin/phosphoinositide 3-kinase (PI3K) pathway and the TOR (target of rapamycin) pathway. Using genetic and biochemical strategies, the epistatic and molecular interactions were elucidated for genes that comprise these pathways.


## The Regulation of Cell Size and Not Cell Numbers

### *The PI3K Pathway*

*Drosophila* has one insulin/IGF (insulin-like growth factor) receptor homolog known as dInR (Chen et al. 1996; Fernandez et al. 1995) and several insulin-like peptides (dILPs) (Brogiolo et al. 2001). These together control the carbohydrate metabolism and growth in flies (Ikeya et al. 2002; Rulifson et al. 2002). Through a mechanism that involves phosphorylation of its carboxy-terminal end, the dInR recruits downstream signaling molecules without the need for adaptor proteins. The signaling also involves the insulin receptor substrate (IRS) protein Chico, which contains a phosphotyrosine-binding domain (PTB) that facilitates its binding with activated dInR (Bohni et al. 1999; Poltilove et al. 2000). Subsequently, the pathway functions by activating the PI3K pathway, via activation of the *Drosophila* PI3K-Dp110 and its adaptor subunit Dp60 (Leevers 2001; Leevers et al. 1996; Weinkove et al. 1999). Dp110/Dp60 heterodimers are recruited to the plasma membrane following the binding of p60 SH2 domain to phosphorylated dInr and Chico, which allows the PI3K access to the phosphoinositide substrates in the plasma membrane. This sets up a signaling cascade in which PIP3 transduces the signal to downstream effectors that contain the PIP3-binding PH domains and causes relocalization of these proteins to the plasma membrane (Fig. 5).

In flies, two such effectors exist—which are the *Drosophila* homolog of phosphoinositide-dependent kinase 1 (PDK1) and its substrate AKT aka protein kinase B (PKB). PDK1 localizes to the membrane during low levels of PI3K activity via its affinity to PIP3, whereas AKT requires high levels of PI3K activity to become membrane localized, through a process involving binding of PIP3 to its PH domain and phosphorylation by PDK1 (Vanhaesebroeck and Alessi 2000). In flies, the activity of dAkt is reduced in the absence of Dp110, and co-expression of dPDK1 and dAKT activates dAKT and induces growth (Cho et al. 2001; Radimerski et al. 2002b; Rintelen et al. 2001) (Fig. 5).

A negative regulator of the PI3K activity is the lipid phosphatase PTEN, which removes the 3' phosphate from three phosphoinositides generated by PI3K (Gao et al. 2000; Goberdhan et al. 1999; Huang et al. 1999) (Fig. 5). Genetic interaction studies support the model where PTEN directly antagonizes PI3K. Loss of PTEN leads to overgrowths due to increased levels of PIP3 (Oldham et al. 2002). Recently, the FOXO family of transcription factors was identified as the target that enabled AKT to regulate growth (Tran et al. 2003). AKT-mediated phosphorylation of FOXO antagonizes its transcriptional activity by creating a 14-3-3 binding site that leads to



**Fig. 5** Model depicting regulation of INR/TOR signaling pathway governed by nutritional status in *Drosophila*. Cellular growth in part is also dependent on the availability of nutrients. This aspect of growth regulation is mainly regulated by the insulin/TOR signaling pathway. Some of the well-studied players of the pathway include phosphatidylinositide 3-kinase and Akt that integrate upstream signaling from growth factor receptors and relay it to TSC1 and TSC2 to regulate ribosomal and protein biosynthesis in addition to actin organization. Other energy-sensing and amino acid-sensing mechanisms are also thought to interact with the core TSC/TOR pathway. However, the exact role or the mechanism by which this takes place remains largely unknown

cytoplasmic sequestration of FOXO (Brunet et al. 1999, 2002; Burgering and Kops 2002). *Drosophila* has one FOXO family transcription factor (dFOXO)—which functions downstream of AKT. Interestingly, loss of function of dFOXO has no apparent effect on cell size or growth as flies homozygous mutant for dFOXO are viable and normal in size (Junger et al. 2003).

The loss of function of Dp110, p60, chico, dINR, dPDK1, and dAKT shows similar effects on cell size and tissue growth (Fig. 5). For example, twin-spot analysis revealed that loss-of-function clones of mutations in these genes are smaller than the corresponding wild-type twin clones that lead to formation of smaller structures (Bohni et al. 1999; Brogiolo et al. 2001; Rintelen et al. 2001; Verdu et al. 1999; Weinkove et al. 1999). Overexpression of PI3K pathway components like Dp110 leads to increased insulin/PI3K signaling and a corresponding increase in cell size, cell number, and tissue growth (Goberdhan et al. 1999; Huang et al. 1999; Leever et al. 1996). Overall, changes in levels of insulin/PI3K signaling have profound effects on organ and organismal size due to effects on cell growth and cell division throughout development and affect the final body/organ size (Fig. 5).

### ***The TSC-TOR Pathway***

Two target of rapamycin (TOR) genes, *TOR1* and *TOR2*, were initially identified in yeast and were shown to be kinases that regulate growth in all organisms by acting as nutrient sensors that couple signaling to nutrient availability (for review, see



Neufeld 2003; Gingras et al. 2001). *Drosophila* TOR (dTOR) promotes growth by stimulating translation via promoting the activity of the *Drosophila* S6Kinase (Montagne et al. 1999) and inhibiting the *Drosophila* 4E-BP1 (a homolog of the eukaryotic translation initiator 4E)—the translational inhibitor of eIF4E, which is a part of the translation initiation complex (Gingras et al. 2001; Lasko 2000). Hyperphosphorylation of d4E-BP1, which is in part controlled by the TOR kinase, relieves its interaction with eIF4E leading to translation initiation.

TOR signaling is negatively regulated by a complex formed by the tuberous sclerosis complex tumor suppressors, TSC1 and TSC2 (Marygold and Leever 2002) (Fig. 5). Mutations in TSC1/2 cause formation of large cells and are implicated in the inherited benign hamartomas observed in the tuberous sclerosis patients (Kandt 2002; Montagne et al. 2001). The *Drosophila Tsc1/2* genes show similar effects on cell size and were identified by several groups in the *eyFLP cell lethal* screens as mutants with overgrown heads (Gao and Pan 2001; Potter et al. 2001; Tapon et al. 2001). Loss of *Tsc1/2* causes increased growth, whereas overexpression of TSC1/2 causes reduced growth due to slow cell cycle progression in the mutant cells. Growth regulation via TSC1/2 happens through preventing dS6K activation via dTOR (Gao et al. 2002; Radimerski et al. 2002a, b). Another important component of this pathway is the GTPase *Rheb*, which is a target of TSC (Saucedo et al. 2003; Stocker et al. 2003; Zhang et al. 2003). The Rheb-GTP levels play a central role in regulating the activity of TOR pathway and the TOR protein that exists in two large multimeric complexes in the cell, viz., the rapamycin-sensitive TORC1 complex and the rapamycin-resistant TORC2 complex (Hara et al. 2002; Kim et al. 2002, 2003; Loewith et al. 2002; Sarbassov et al. 2004).

The TORC1 complex consists of TOR, Raptor, and LST8; and responds to the presence of growth factors and nutrients to control protein synthesis (Fig. 5). The small GTPase protein Rheb (Ras homolog enriched in the brain) is a direct activator of TORC1 (Long et al. 2004; Saucedo et al. 2003; Stocker et al. 2003), and the tuberous sclerosis (TSC) complex (TSC1/TSC2) negatively regulates TORC1 by functioning as a GTPase-activating protein (GAP) for Rheb (Potter and Xu 2001; Zhang et al. 2003). Growth factors such as insulin or insulin-like growth factors (IGFs) activate TORC1 signaling upstream of the TSC1/TSC2 (TSC1/2) complex through the insulin receptor (InR)/phosphoinositide 3-kinase (PI3K)/AKT signaling pathway (Inoki et al. 2002; Potter et al. 2002). TORC1 also senses nutrient availability. Amino acids regulate TORC1 through mechanisms independent or downstream of TSC complex, and recently the Rag small GTPases have been shown to interact with TOR and promote TORC1 activity by controlling its subcellular localization (Nellist et al. 2008; Sancak et al. 2010).

TORC2 complex (Fig. 5) consists of TOR, Rictor, Sin1 (stress-activated map kinase-interacting protein 1), and LST8 and phosphorylates and activates several AGC family kinases, including AKT, serum and glucocorticoid-regulated kinase (SGK), and protein kinase C (PKC), and thereby regulates cell survival, cell cycle progression, and metabolism (Pearce et al. 2010) (Li 2010 #8573; Gao 2010 #8574). In contrast to TORC1, little is known about the upstream activators of mTORC2. Although the general mechanisms have not been accepted, PI3K, TSC, and Rheb

have been shown to regulate TORC2 activity, and Rictor has been identified as a substrate of S6 kinase (S6K), suggesting possible regulation of TORC2 through the TORC1 pathway (Dibble et al. 2009; Treins et al. 2010; Yang et al. 2006). Nevertheless, it is generally thought that growth factors may control TORC2, either directly or indirectly (Zinzalla et al. 2011). TORC2 has been proposed to function independent of amino acid availability (Jacinto et al. 2006); however, recent findings show that amino acids may also activate TORC2 (Tato et al. 2011).

The central role of TOR in cell growth has been largely attributed to TORC1, but mounting evidence points to a role for TORC2 as well in this basic cellular process. For instance, TORC2 localizes in polysomal fractions and associates with ribosomal proteins, indicating a potential role for TORC2 in protein synthesis and maturation (Cybulski and Hall 2009; Zinzalla et al. 2011). *lst8* knockout flies are viable but small, similar to *rictor* mutants but dissimilar to flies with *tor* or *rheb* mutations, which are lethal (Avruch et al. 2009; Liao et al. 2008; Wang et al. 2012b). Neither loss nor overexpression of LST8 affected the kinase activity of TORC1 toward S6K or autophagy, whereas the kinase activity of TORC2 toward AKT was completely lost in the *lst8* mutants (Avruch et al. 2009; Liao et al. 2008; Wang et al. 2012b).

In terms of effects of TOR signaling on growth phenotypes in *Drosophila*, loss of dTOR leads to a decrease in larvae size; however, the larvae fail to mature and die before reaching adulthood. In mosaic *Drosophila*, loss of dTOR leads to a decrease in cell size while maintaining the general organization of the tissue (Oldham et al. 2000b; Zhang et al. 2000). However, it is less clear how cell size is regulated downstream of mTOR. One of the most potent candidates in this regulation is S6K. In *Drosophila*, knockout of *S6K* results in high rates of embryonic lethality. In the surviving adults, however, there is a decrease in body size. Knockdown of either dPTEN or dTSC1 is sufficient to increase cell size; however, a double knockdown of dPTEN and dTSC1 has additive effects on cell size regulation. This suggests that in *Drosophila*, the pathways may have independent components in the regulation of cell size (Gao and Pan 2001). It may also highlight the differences in the regulation of TSC2 by AKT in *Drosophila* as seen by mutations of the AKT phosphorylation sites on TSC2 (Dong and Pan 2004; Pan et al. 2004). Loss of either dPTEN or dTSC1 can lead to increases in cell size; however, a report has suggested that only knockdown of dTSC1 leads to increases in dS6K (Radimerski et al. 2002a), whereas other reports have also seen increases in dS6K with the knockdown of dPTEN (Sarbasov et al. 2004; Yang et al. 2006). It is possible that dTSC1 regulates cell size in a dTOR-dependent manner, whereas dPTEN partially regulates cell size in a dTOR-independent manner (Radimerski et al. 2002b).

In conclusion, the TOR signaling pathway is a complex network of cell size regulators that is also implicated in tumorigenesis and cell survival (Fig. 5). Several pathways interact and intersect with the TOR pathway at multiple points upstream and downstream of TOR.

## Growth Regulation: A Network of Tumor Suppressors

Overall, growth control occurs through the Hippo and TSC-TOR pathways in conjunction with pathways regulating pattern formation during development. These pathways intersect in complicated signaling networks in all cell types and coordinately regulate overall growth of an organism. Our progress in understanding of these pathways has led the way to find molecules and interactions important for regenerative growth and wound healing—phenomena that have been well documented but not well understood at the molecular level for a long time. In addition, the establishment of these growth regulatory networks has led many insights in the fields of cancer (e.g., the underlying genetics and biology link between hamartomas and TSC genes; schwannomas and NF2; YAP and hepatocellular carcinoma, TAZ and breast cancer, etc.). In the future, it will be interesting to learn about the regulation of these pathways by extracellular and intracellular mechanisms, an area expected to expand rapidly with our increased understanding of the integration points in the circuitry of these networks.

## References

- Acquisti C, Kumar S, Elser JJ (2009) Signatures of nitrogen limitation in the elemental composition of the proteins involved in the metabolic apparatus. *Proc Biol Sci* 276:2605–2610
- Alarcon C, Zaromytidou AI, Xi Q, Gao S, Yu J, Fujisawa S, Barlas A, Miller AN, Manova-Todorova K, Macias MJ et al (2009) Nuclear CDKs drive Smad transcriptional activation and turnover in BMP and TGF-beta pathways. *Cell* 139:757–769
- Angus L, Moleirinho S, Herron L, Sinha A, Zhang X, Nestrata M, Dholakia K, Prystowsky MB, Harvey KF, Reynolds PA et al (2012) Willin/FRMD6 expression activates the Hippo signaling pathway kinases in mammals and antagonizes oncogenic YAP. *Oncogene* 31:238–250
- Anon (2003) INGN 201: Ad-p53, Ad5CMV-p53, Adenoviral p53, INGN 101, p53 gene therapy – Introgen, RPR/INGN 201. *BioDrugs* 17:216–222
- Avruch J, Long X, Ortiz-Vega S, Rapley J, Papageorgiou A, Dai N (2009) Amino acid regulation of TOR complex 1. *Am J Physiol Endocrinol Metab* 296:E592–E602
- Avruch J, Zhou D, Fitamant J, Bardeesy N, Mou F, Barrufet LR (2012) Protein kinases of the Hippo pathway: regulation and substrates. *Semin Cell Dev Biol* 23:770–784
- Azzolin L, Zanconato F, Bresolin S, Forcato M, Basso G, Bicciato S, Cordenonsi M, Piccolo S (2012) Role of TAZ as mediator of Wnt signaling. *Cell* 151:1443–1456
- Bachmann A, Schneider M, Theilenberg E, Grawe F, Knust E (2001) *Drosophila* Stardust is a partner of Crumbs in the control of epithelial cell polarity. *Nature* 414(6864):638–643
- Badouel C, Gardano L, Amin N, Garg A, Rosenfeld R, Le Bihan T, McNeill H (2009) The FERM-domain protein expanded regulates Hippo pathway activity via direct interactions with the transcriptional activator Yorkie. *Dev Cell* 16:411–420
- Baker NE (2001) Cell proliferation, survival, and death in the *Drosophila* eye. *Semin Cell Dev Biol* 12:499–507
- Baker NE, Yu S, Han D (1996) Evolution of proneural atonal expression during distinct regulatory phases in the developing *Drosophila* eye. *Curr Biol* 6:1290–1301
- Bangs P, White K (2000) Regulation and execution of apoptosis during *Drosophila* development. *Dev Dyn* 218:68–79

- Barry ER, Camargo FD (2013) The Hippo superhighway: signaling crossroads converging on the Hippo/Yap pathway in stem cells and development. *Curr Opin Cell Biol* 25(2):247–253
- Baumgartner R, Poernbacher I, Buser N, Hafen E, Stocker H (2010) The WW domain protein Kibra acts upstream of Hippo in *Drosophila*. *Dev Cell* 18:309–316
- Bellen HJ, O’Kane CJ, Wilson C, Grossniklaus U, Pearson RK, Gehring WJ (1989) P-element-mediated enhancer detection: a versatile method to study development in *Drosophila*. *Genes Dev* 3:1288–1300
- Bellen HJ, Levis RW, He Y, Carlson JW, Evans-Holm M, Bae E, Kim J, Metaxakis A, Savakis C, Schulze KL et al (2011) The *Drosophila* gene disruption project: progress using transposons with distinctive site specificities. *Genetics* 188:731–743
- Bellosta P, Gallant P (2010) Myc function in *Drosophila*. *Genes Cancer* 1:542–546
- Bennett FC, Harvey KF (2006) Fat cadherin modulates organ size in *Drosophila* via the Salvador/Warts/Hippo signaling pathway. *Curr Biol* 16:2101–2110
- Bergantinos C, Vilana X, Corominas M, Serras F (2010) Imaginal discs: Renaissance of a model for regenerative biology. *BioEssays* 32:207–217
- Bertini E, Oka T, Sudol M, Strano S, Blandino G (2009) YAP: at the crossroad between transformation and tumor suppression. *Cell Cycle* 8:49–57
- Bier E (2005) *Drosophila*, the golden bug, emerges as a tool for human genetics. *Nat Rev Genet* 6:9–23
- Blair SS (2003) Genetic mosaic techniques for studying *Drosophila* development. *Development* 130:5065–5072
- Boggiano JC, Fehon RG (2012) Growth control by committee: intercellular junctions, cell polarity, and the cytoskeleton regulate Hippo signaling. *Dev Cell* 22:695–702
- Boggiano JC, Vanderzalm PJ, Fehon RG (2011) Tao-1 phosphorylates Hippo/MST kinases to regulate the Hippo-Salvador-Warts tumor suppressor pathway. *Dev Cell* 21:888–895
- Bohni R, Riesgo-Escovar J, Oldham S, Brogiolo W, Stocker H, Andrus BF, Beckingham K, Hafen E (1999) Autonomous control of cell and organ size by CHICO, a *Drosophila* homolog of vertebrate IRS1-4. *Cell* 97:865–875
- Bonini NM, Fortini ME (1999) Surviving *Drosophila* eye development: integrating cell death with differentiation during formation of a neural structure. *BioEssays* 21:991–1003
- Bonner JM, Boulianne GL (2011) *Drosophila* as a model to study age-related neurodegenerative disorders: Alzheimer’s disease. *Exp Gerontol* 46:335–339
- Bosveld F, Bonnet I, Guirao B, Tlili S, Wang Z, Petitalot A, Marchand R, Bardet PL, Marcq P, Graner F, Bellaïche Y (2012) Mechanical control of morphogenesis by Fat/Dachsous/Four-jointed planar cell polarity pathway. *Science* 336(6082):724–727
- Boutros M, Ahringer J (2008) The art and design of genetic screens: RNA interference. *Nat Rev Genet* 9:554–566
- Brittle AL, Repiso A, Casal J, Lawrence PA, Strutt D (2010) Four-jointed modulates growth and planar polarity by reducing the affinity of dachsous for fat. *Curr Biol* 20:803–810
- Brogiolo W, Stocker H, Ikeya T, Rintelen F, Fernandez R, Hafen E (2001) An evolutionarily conserved function of the *Drosophila* insulin receptor and insulin-like peptides in growth control. *Curr Biol* 11:213–221
- Brumby AM, Richardson HE (2003) Scribble mutants cooperate with oncogenic Ras or Notch to cause neoplastic overgrowth in *Drosophila*. *EMBO J* 22:5769–5779
- Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, Anderson MJ, Arden KC, Blenis J, Greenberg ME (1999) Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 96:857–868
- Brunet A, Kanai F, Stehn J, Xu J, Sarbassova D, Frangioni JV, Dalal SN, DeCaprio JA, Greenberg ME, Yaffe MB (2002) 14-3-3 transits to the nucleus and participates in dynamic nucleocytoplasmic transport. *J Cell Biol* 156:817–828
- Bryant PJ (1978) Pattern formation in imaginal discs. *Ashburner, Wright, 1978–1980 c*, 230–335
- Bryant PJ (1987) Experimental and genetic analysis of growth and cell proliferation in *Drosophila* imaginal discs. In Loomis WF (ed), pp. 339–372

- Bryant PJ (2001) Growth factors controlling imaginal disc growth in *Drosophila*. Novartis Found Symp 237:182–194; discussion 194–202
- Bryant PJ, Schmidt O (1990) The genetic control of cell proliferation in *Drosophila* imaginal discs. J Cell Sci Suppl 13:169–189
- Bryant PJ, Huettner B, Held LI Jr, Ryerse J, Szidonya J (1988) Mutations at the fat locus interfere with cell proliferation control and epithelial morphogenesis in *Drosophila*. Dev Biol 129:541–554
- Burgering BM, Kops GJ (2002) Cell cycle and death control: long live Forkheads. Trends Biochem Sci 27:352–360
- Burke R, Basler K (1997) Hedgehog signaling in *Drosophila* eye and limb development – conserved machinery, divergent roles? Curr Opin Neurobiol 7:55–61
- Cagan R (1993) Cell fate specification in the developing *Drosophila* retina. Dev Suppl 119:19–28
- Cagan R (2009) Principles of *Drosophila* eye differentiation. Curr Top Dev Biol 89:115–135
- Campbell S, Inamdar M, Rodrigues V, Raghavan V, Palazzolo M, Chovnick A (1992) The scalloped gene encodes a novel, evolutionarily conserved transcription factor required for sensory organ differentiation in *Drosophila*. Genes Dev 6:367–379
- Chan SW, Lim CJ, Chen L, Chong YF, Huang C, Song H, Hong W (2011) The Hippo pathway in biological control and cancer development. J Cell Physiol 226:928–939
- Chen CK, Chien CT (1999) Negative regulation of atonal in proneural cluster formation of *Drosophila* R8 photoreceptors. Proc Natl Acad Sci USA 96:5055–5060
- Chen C, Jack J, Garofalo RS (1996) The *Drosophila* insulin receptor is required for normal growth. Endocrinology 137:846–856
- Chen CL, Gajewski KM, Hamaratoglu F, Bossuyt W, Sansores-Garcia L, Tao C, Halder G (2010) The apical-basal cell polarity determinant Crumbs regulates Hippo signaling in *Drosophila*. Proc Natl Acad Sci USA 107:15810–15815
- Chen CL, Schroeder MC, Kango-Singh M, Tao C, Halder G (2012) Tumor suppression by cell competition through regulation of the Hippo pathway. Proc Natl Acad Sci U S A 109(2):484–489
- Cho E, Irvine KD (2004) Action of fat, four-jointed, dachsous and dachs in distal-to-proximal wing signaling. Development 131:4489–4500
- Cho KS, Lee JH, Kim S, Kim D, Koh H, Lee J, Kim C, Kim J, Chung J (2001) *Drosophila* phosphoinositide-dependent kinase-1 regulates apoptosis and growth via the phosphoinositide 3-kinase-dependent signaling pathway. Proc Natl Acad Sci USA 98:6144–6149
- Cho E, Feng Y, Rauskolb C, Maitra S, Fehon R, Irvine KD (2006) Delineation of a Fat tumor suppressor pathway. Nat Genet 38:1142–1150
- Clark HF, Brentrup D, Schneitz K, Bieber A, Goodman C, Noll M (1995) Dachsous encodes a member of the cadherin superfamily that controls imaginal disc morphogenesis in *Drosophila*. Genes Dev 9(12):1530–1542
- Collak FK, Yagiz K, Luthringer DJ, Erkaya B, Cinar B (2012) Threonine-120 phosphorylation regulated by phosphoinositide-3-kinase/Akt and mammalian target of rapamycin pathway signaling limits the antitumor activity of mammalian sterile 20-like kinase 1. J Biol Chem 287:23698–23709
- Conlon I, Raff M (1999) Size control in animal development. Cell 96:235–244
- Cook M, Tyers M (2007) Size control goes global. Curr Opin Biotechnol 18:341–350
- Cooper S (2004) Control and maintenance of mammalian cell size. BMC Cell Biol 5:35
- Cordero JB, Larson DE, Craig CR, Hays R, Cagan R (2007) Dynamic decapentaplegic signaling regulates patterning and adhesion in the *Drosophila* pupal retina. Development 134(10):1861–1871
- Courbard JR, Djiane A, Wu J, Mlodzik M (2009) The apical/basal-polarity determinant Scribble cooperates with the PCP core factor Stbm/Vang and functions as one of its effectors. Dev Biol 333(1):67–77
- Crickmore MA, Mann RS (2008) The control of size in animals: insights from selector genes. BioEssays 30:843–853

- Cybulski N, Hall MN (2009) TOR complex 2: a signaling pathway of its own. *Trends Biochem Sci* 34:620–627
- Daniel A, Dumstrei K, Lengyel JA, Hartenstein V (1999) The control of cell fate in the embryonic visual system by atonal, tailless and EGFR signaling. *Development* 126:2945–2954
- Das Thakur M, Feng Y, Jagannathan R, Seppa MJ, Skeath JB, Longmore GD (2010) Ajuba LIM proteins are negative regulators of the Hippo signaling pathway. *Curr Biol* 20:657–662
- de Nooij JC, Hariharan IK (1995) Uncoupling cell fate determination from patterned cell division in the *Drosophila* eye. *Science* 270:983–985
- Dibble CC, Asara JM, Manning BD (2009) Characterization of Rictor phosphorylation sites reveals direct regulation of mTOR complex 2 by S6K1. *Mol Cell Biol* 29:5657–5670
- Dickson B, Hafen E (1993) Genetic dissection of eye development in *Drosophila*. In: Bate M, Martinez Arias A (eds) *The development of Drosophila melanogaster*, vol II. Cold Spring Harbor Laboratory Press, New York, pp 1327–1362
- Doggett K, Grusche FA, Richardson HE, Brumby AM (2011) Loss of the *Drosophila* cell polarity regulator scribbled promotes epithelial tissue overgrowth and cooperation with oncogenic Ras-Raf through impaired Hippo pathway signaling. *BMC Dev Biol* 11:57
- Dominguez M (1999) Dual role for Hedgehog in the regulation of the proneural gene atonal during ommatidia development. *Development* 126:2345–2353
- Dominguez M, Casares F (2005) Organ specification-growth control connection: new in-sights from the *Drosophila* eye-antennal disc. *Dev Dyn* 232:673–684
- Dong J, Pan D (2004) Tsc2 is not a critical target of Akt during normal *Drosophila* development. *Genes Dev* 18:2479–2484
- Dong J, Feldmann G, Huang J, Wu S, Zhang N, Comerford SA, Gayyed MF, Anders RA, Maitra A, Pan D (2007) Elucidation of a universal size-control mechanism in *Drosophila* and mammals. *Cell* 130:1120–1133
- Edgar BA (1999) From small flies come big discoveries about size control. *Nat Cell Biol* 1:E191–E193
- Edgar BA (2006) From cell structure to transcription: Hippo forges a new path. *Cell* 124:267–273
- Enomoto M, Igaki T (2011) Deciphering tumor-suppressor signaling in flies: Genetic link between Scribble/Dlg/Lgl and the Hippo pathways. *J Genet Genomics* 38(10):461–470
- Fan SS, Chen MS, Lin JF, Chao WT, Yang VC (2003) Use of gain-of-function study to delineate the roles of crumbs in *Drosophila* eye development. *J Biomed Sci* 10(6 Pt. 2):766–773
- Feng Y, Irvine KD (2007) Fat and expanded act in parallel to regulate growth through warts. *Proc Natl Acad Sci USA* 104:20362–20367
- Feng Y, Irvine KD (2009) Processing and phosphorylation of the Fat receptor. *Proc Natl Acad Sci USA* 106:11989–11994
- Fernandez R, Tabarini D, Azpiazu N, Frasch M, Schlessinger J (1995) The *Drosophila* insulin receptor homolog: a gene essential for embryonic development encodes two receptor isoforms with different signaling potential. *EMBO J* 14:3373–3384
- Fernandez BG, Gaspar P, Bras-Pereira C, Jezowska B, Rebelo SR, Janody F (2011) Actin-capping protein and the Hippo pathway regulate F-actin and tissue growth in *Drosophila*. *Development* 138:2337–2346
- Firth LC, Bhattacharya A, Baker NE (2010) Cell cycle arrest by a gradient of Dpp signaling during *Drosophila* eye development. *BMC Dev Biol* 10:28
- Fjose A, Polito LC, Weber U, Gehring WJ (1984) Developmental expression of the white locus of *Drosophila melanogaster*. *EMBO J* 3:2087–2094
- Gadd S, Beezhold P, Jennings L, George D, Leuer K, Huang CC, Huff V, Tognon C, Sorensen PH, Triche T et al (2012) Mediators of receptor tyrosine kinase activation in infantile fibrosarcoma: a Children's oncology group study. *J Pathol* 228:119–130
- Gao X, Pan D (2001) TSC1 and TSC2 tumor suppressors antagonize insulin signaling in cell growth. *Genes Dev* 15:1383–1392
- Gao X, Neufeld TP, Pan D (2000) *Drosophila* PTEN regulates cell growth and proliferation through PI3K-dependent and -independent pathways. *Dev Biol* 221:404–418

- Gao X, Zhang Y, Arrazola P, Hino O, Kobayashi T, Yeung RS, Ru B, Pan D (2002) Tsc tumour suppressor proteins antagonize amino-acid-TOR signalling. *Nat Cell Biol* 4:699–704
- Garami A, Zwartkruis FJ, Nobukuni T, Joaquin M, Rocco M, Stocker H, Kozma SC, Hafen E, Bos JL, Thomas G (2003) Insulin activation of Rheb, a mediator of mTOR/S6K/4E-BP signaling, is inhibited by TSC1 and 2. *Mol Cell* 11:1457–1466
- Garg A, Srivastava A, Davis MM, O’Keefe SL, Chow L, Bell JB (2007) Antagonizing scalloped with a novel vestigial construct reveals an important role for scalloped in *Drosophila melanogaster* leg, eye and optic lobe development. *Genetics* 175(2):659–669
- Garoia F, Guerra D, Pezzoli MC, Lopez-Varea A, Cavicchi S, Garcia-Bellido A (2000) Cell behaviour of *Drosophila* fat cadherin mutations in wing development. *Mech Dev* 94(1–2):95–109
- Genevet A, Tapon N (2011) The Hippo pathway and apico-basal cell polarity. *Biochem J* 436:213–224
- Genevet A, Polesello C, Blight K, Robertson F, Collinson LM, Pichaud F, Tapon N (2009) The Hippo pathway regulates apical-domain size independently of its growth-control function. *J Cell Sci* 122:2360–2370
- Genevet A, Wehr MC, Brain R, Thompson BJ, Tapon N (2010) Kibra is a regulator of the Salvador/Warts/Hippo signaling network. *Dev Cell* 18:300–308
- Gilbert MM, Tipping M, Veraksa A, Moberg KH (2011) A screen for conditional growth suppressor genes identifies the *Drosophila* homolog of HD-PTP as a regulator of the oncoprotein Yorkie. *Dev Cell* 20:700–712
- Gingras AC, Raught B, Sonenberg N (2001) Regulation of translation initiation by FRAP/mTOR. *Genes Dev* 15:807–826
- Goberdhan DC, Paricio N, Goodman EC, Mlodzik M, Wilson C (1999) *Drosophila* tumor suppressor PTEN controls cell size and number by antagonizing the Chico/PI3-kinase signaling pathway. *Genes Dev* 13:3244–3258
- Goulev Y, Fauny JD, Gonzalez-Marti B, Flagiello D, Silber J, Zider A (2008) SCALLOPED interacts with YORKIE, the nuclear effector of the hippo tumor-suppressor pathway in *Drosophila*. *Curr Biol* 18:435–441
- Graves JD, Gotoh Y, Draves KE, Ambrose D, Han DK, Wright M, Chernoff J, Clark EA, Krebs EG (1998) Caspase-mediated activation and induction of apoptosis by the mammalian Ste20-like kinase Mst1. *EMBO J* 17:2224–2234
- Grebien F, Dolznig H, Beug H, Mullner EW (2005) Cell size control: new evidence for a general mechanism. *Cell Cycle* 4:418–421
- Greenwood S, Struhl G (1999) Progression of the morphogenetic furrow in the *Drosophila* eye: the roles of Hedgehog, Decapentaplegic and the Raf pathway. *Development* 126:5795–5808
- Grusche FA, Richardson HE, Harvey KF (2010) Upstream regulation of the hippo size control pathway. *Curr Biol* 20:R574–R582
- Grusche FA, Degoutin JL, Richardson HE, Harvey KF (2011) The Salvador/Warts/Hippo pathway controls regenerative tissue growth in *Drosophila melanogaster*. *Dev Biol* 350:255–266
- Grzeschik NA, Amin N, Secombe J, Brumby AM, Richardson HE (2007) Abnormalities in cell proliferation and apico-basal cell polarity are separable in *Drosophila* lgl mutant clones in the developing eye. *Dev Biol* 311:106–123
- Grzeschik NA, Parsons LM, Allott ML, Harvey KF, Richardson HE (2010a) Lgl, aPKC, and Crumbs regulate the Salvador/Warts/Hippo pathway through two distinct mechanisms. *Curr Biol* 20:573–581
- Grzeschik NA, Parsons LM, Richardson HE (2010b) Lgl, the SWH pathway and tumorigenesis: it’s a matter of context & competition! *Cell Cycle* 9:3202–3212
- Guan J, Li H, Rogulja A, Axelrod JD, Cadigan KM (2007) The *Drosophila* casein kinase I epsilon/delta Discs overgrown promotes cell survival via activation of DIAP1 expression. *Dev Biol* 303:16–28
- Guo C, Tommasi S, Liu L, Yee JK, Dammann R, Pfeifer GP (2007) RASSF1A is part of a complex similar to the *Drosophila* Hippo/Salvador/Lats tumor-suppressor network. *Curr Biol* 17:700–705

- Habbig S, Bartram MP, Muller RU, Schwarz R, Andriopoulos N, Chen S, Sagmuller JG, Hoehne M, Burst V, Liebau MC et al (2011) NPHP4, a cilia-associated protein, negatively regulates the Hippo pathway. *J Cell Biol* 193:633–642
- Habbig S, Bartram MP, Sagmuller JG, Griessmann A, Franke M, Muller RU, Schwarz R, Hoehne M, Bergmann C, Tessmer C et al (2012) The ciliopathy disease protein NPHP9 promotes nuclear delivery and activation of the oncogenic transcriptional regulator TAZ. *Hum Mol Genet* 21:5528–5538
- Hafen E (1991) Patterning by cell recruitment in the *Drosophila* eye. *Curr Opin Genet Dev* 1:268–274
- Hafen E (2004) Interplay between growth factor and nutrient signaling: lessons from *Drosophila* TOR. *Curr Top Microbiol Immunol* 279:153–167
- Halder G, Johnson RL (2011) Hippo signaling: growth control and beyond. *Development* 138:9–22
- Halder G, Dupont S, Piccolo S (2012) Transduction of mechanical and cytoskeletal cues by YAP and TAZ. *Nat Rev Mol Cell Biol* 13(9):591–600
- Hamaratoglu F, Willecke M, Kango-Singh M, Nolo R, Hyun E, Tao C, Jafar-Nejad H, Halder G (2006) The tumour-suppressor genes NF2/Merlin and expanded act through Hippo signalling to regulate cell proliferation and apoptosis. *Nat Cell Biol* 8:27–36
- Hara K, Maruki Y, Long X, Yoshino K, Oshiro N, Hidayat S, Tokunaga C, Avruch J, Yonezawa K (2002) Raptor, a binding partner of target of rapamycin (TOR), mediates TOR action. *Cell* 110:177–189
- Harvey KF, Hariharan IK (2012) The hippo pathway. *Cold Spring Harb Perspect Biol* 4:a011288
- Harvey NL, Daish T, Mills K, Dorstyn L, Quinn LM, Read SH, Richardson H, Kumar S (2001) Characterization of the *Drosophila* caspase, DAMM. *J Biol Chem* 276:25342–25350
- Harvey KF, Pflieger CM, Hariharan IK (2003) The *Drosophila* Mst ortholog, Hippo, restricts growth and cell proliferation and promotes apoptosis. *Cell* 114:457–467
- Hergovich A (2012) Mammalian Hippo signalling: a kinase network regulated by protein-protein interactions. *Biochem Soc Trans* 40:124–128
- Herranz H, Hong X, Cohen SM (2012) Mutual repression by bantam miRNA and Capicua links the EGFR/MAPK and Hippo pathways in growth control. *Curr Biol* 22(8):651–657
- Hiemer SE, Varelas X (2013) Stem cell regulation by the Hippo pathway. *Biochim Biophys Acta* 1830:2323–2334
- Ho KC, Zhou Z, She YM, Chun A, Cyr TD, Yang X (2011) Itch E3 ubiquitin ligase regulates large tumor suppressor 1 stability [corrected]. *Proc Natl Acad Sci USA* 108:4870–4875
- Hong W, Guan KL (2012) The YAP and TAZ transcription co-activators: key downstream effectors of the mammalian Hippo pathway. *Semin Cell Dev Biol* 23:785–793
- Huang H, Potter CJ, Tao W, Li DM, Brogiolo W, Hafen E, Sun H, Xu T (1999) PTEN affects cell size, cell proliferation and apoptosis during *Drosophila* eye development. *Development* 126:5365–5372
- Huang J, Wu S, Barrera J, Matthews K, Pan D (2005) The Hippo signaling pathway coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the *Drosophila* Homolog of YAP. *Cell* 122:421–434
- Hwang E, Ryu KS, Paakkonen K, Guntert P, Cheong HK, Lim DS, Lee JO, Jeon YH, Cheong C (2007) Structural insight into dimeric interaction of the SARAH domains from Mst1 and RASSF family proteins in the apoptosis pathway. *Proc Natl Acad Sci USA* 104:9236–9241
- Ikeya T, Galic M, Belawat P, Nairz K, Hafen E (2002) Nutrient-dependent expression of insulin-like peptides from neuroendocrine cells in the CNS contributes to growth regulation in *Drosophila*. *Curr Biol* 12:1293–1300
- Inoki K, Li Y, Zhu T, Wu J, Guan KL (2002) TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling. *Nat Cell Biol* 4:648–657
- Ishikawa HO, Takeuchi H, Haltiwanger RS, Irvine KD (2008) Four-jointed is a Golgi kinase that phosphorylates a subset of cadherin domains. *Science* 321:401–404
- Ishiyuchi T, Takeichi M (2012) Nectins localize Willin to cell-cell junctions. *Genes Cells* 17:387–397



- Jacinto E, Facchinetti V, Liu D, Soto N, Wei S, Jung SY, Huang Q, Qin J, Su B (2006) SIN1/MIP1 maintains rictor-mTOR complex integrity and regulates Akt phosphorylation and substrate specificity. *Cell* 127:125–137
- Jacobson MD, Weil M, Raff MC (1997) Programmed cell death in animal development. *Cell* 88:347–354
- Jarman AP, Grell EH, Ackerman L, Jan LY, Jan YN (1994) Atonal is the proneural gene for *Drosophila* photoreceptors. *Nature* 369:398–400
- Jia J, Zhang W, Wang B, Trinko R, Jiang J (2003) The *Drosophila* Ste20 family kinase dMST functions as a tumor suppressor by restricting cell proliferation and promoting apoptosis. *Genes Dev* 17:2514–2519
- Johnston LA, Gallant P (2002) Control of growth and organ size in *Drosophila*. *BioEssays* 24:54–64
- Junger MA, Rintelen F, Stocker H, Wasserman JD, Vegh M, Radimerski T, Greenberg ME, Hafen E (2003) The *Drosophila* forkhead transcription factor FOXO mediates the reduction in cell number associated with reduced insulin signaling. *J Biol* 2:20
- Justice RW, Zilian O, Woods DF, Noll M, Bryant PJ (1995) The *Drosophila* tumor suppressor gene warts encodes a homolog of human myotonic dystrophy kinase and is required for the control of cell shape and proliferation. *Genes Dev* 9:534–546
- Kagey JD, Brown JA, Moberg KH (2012) Regulation of Yorkie activity in *Drosophila* imaginal discs by the Hedgehog receptor gene patched. *Mech Dev* 129(9–12):339–349
- Kandt RS (2002) Tuberous sclerosis complex and neurofibromatosis type 1: the two most common neurocutaneous diseases. *Neurol Clin* 20:941–964
- Kango-Singh M, Singh A (2009) Regulation of organ size: insights from the *Drosophila* Hippo signaling pathway. *Dev Dyn* 238:1627–1637
- Kango-Singh M, Nolo R, Tao C, Verstreken P, Hiesinger PR, Bellen HJ, Halder G (2002) Shar-pei mediates cell proliferation arrest during imaginal disc growth in *Drosophila*. *Development* 129:5719–5730
- Kango-Singh M, Singh A, Henry Sun Y (2003) Eyeless collaborates with Hedgehog and Decapentaplegic signaling in *Drosophila* eye induction. *Dev Biol* 256:49–60
- Kaplan NA, Colosimo PF, Liu X, Tolwinski NS (2011) Complex interactions between GSK3 and aPKC in *Drosophila* embryonic epithelial morphogenesis. *PLoS One* 6(4):e18616
- Kaplan NA, Tolwinski NS (2010) Spatially defined Dsh-Lgl interaction contributes to directional tissue morphogenesis. *J Cell Sci* 123(18):3157–3165
- Karni R, Hippo Y, Lowe SW, Krainer AR (2008) The splicing-factor oncoprotein SF2/ASF activates mTORC1. *Proc Natl Acad Sci USA* 105:15323–15327
- Karpowicz P, Perez J, Perrimon N (2010) The Hippo tumor suppressor pathway regulates intestinal stem cell regeneration. *Development* 137:4135–4145
- Kiger A, Baum B, Jones S, Jones M, Coulson A, Echeverri C, Perrimon N (2003) A functional genomic analysis of cell morphology using RNA interference. *J Biol* 2(4):27
- Klein TJ, Jenny A, Djiane A, Mlodzik M (2006) CKIepsilon/discs overgrown promotes both Wnt-Fz/beta-catenin and Fz/PCP signaling in *Drosophila*. *Curr Biol* 16(13):1337–1343
- Kim J, Guan KL (2011) Amino acid signaling in TOR activation. *Annu Rev Biochem* 80:1001–1032
- Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, Tempst P, Sabatini DM (2002) mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. *Cell* 110:163–175
- Kim DH, Sarbassov DD, Ali SM, Latek RR, Guntur KV, Erdjument-Bromage H, Tempst P, Sabatini DM (2003) GbetaL, a positive regulator of the rapamycin-sensitive pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol Cell* 11:895–904
- Kim D, Shu S, Coppola MD, Kaneko S, Yuan ZQ, Cheng JQ (2010) Regulation of proapoptotic mammalian ste20-like kinase MST2 by the IGF1-Akt pathway. *PLoS One* 5:e9616
- Konsavage WM, Yochum GS (2013) Intersection of Hippo/YAP and Wnt/beta-catenin signaling pathways. *ACTA BIOCH BIOPH SIN* 45(2):71–79

- Kramer H, Cagan RL (1994) Determination of photoreceptor cell fate in the *Drosophila* retina. *Curr Opin Neurobiol* 4:14–20
- Kumagai A, Dunphy WG (1999) Binding of 14-3-3 proteins and nuclear export control the intracellular localization of the mitotic inducer Cdc25. *Genes Dev* 13:1067–1072
- Kumar JP (2001) Signalling pathways in *Drosophila* and vertebrate retinal development. *Nat Rev Genet* 2:846–857
- Kumar JP (2009) The molecular circuitry governing retinal determination. *Biochim Biophys Acta* 1789:306–314
- Kumar JP, Moses K (2000) Cell fate specification in the *Drosophila* retina. *Results Probl Cell Differ* 31:93–114
- Kumar JP, Moses K (2001) Eye specification in *Drosophila*: perspectives and implications. *Semin Cell Dev Biol* 12:469–474
- Lai ZC, Wei X, Shimizu T, Ramos E, Rohrbaugh M, Nikolaidis N, Ho LL, Li Y (2005) Control of cell proliferation and apoptosis by mob as tumor suppressor, mats. *Cell* 120:675–685
- Lasko P (2000) The *Drosophila melanogaster* genome: translation factors and RNA binding proteins. *J Cell Biol* 150:F51–F56
- Leevers SJ (2001) Growth control: invertebrate insulin surprises! *Curr Biol* 11:R209–R212
- Leevers SJ, Weinkove D, MacDougall LK, Hafen E, Waterfield MD (1996) The *Drosophila* phosphoinositide 3-kinase Dp110 promotes cell growth. *EMBO J* 15:6584–6594
- Li W, Cooper J, Karajannis MA, Giancotti FG (2012) Merlin: a tumour suppressor with functions at the cell cortex and in the nucleus. *EMBO Rep* 13:204–215
- Li L, Edgar BA, Grewal SS (2010a) Nutritional control of gene expression in *Drosophila* larvae via TOR, Myc and a novel cis-regulatory element. *BMC Cell Biol* 11:7
- Li L, Kim E, Yuan H, Inoki K, Goraksha-Hicks P, Schiesher RL, Neufeld TP, Guan KL (2010b) Regulation of mTORC1 by the Rab and Arf GTPases. *J Biol Chem* 285(26):19705–19709
- Liao XH, Majithia A, Huang X, Kimmel AR (2008) Growth control via TOR kinase signaling, an intracellular sensor of amino acid and energy availability, with crosstalk potential to proline metabolism. *Amino Acids* 35:761–770
- Ling C, Zheng Y, Yin F, Yu J, Huang J, Hong Y, Wu S, Pan D (2010) The apical transmembrane protein Crumbs functions as a tumor suppressor that regulates Hippo signaling by binding to expanded. *Proc Natl Acad Sci USA* 107:10532–10537
- Liu X, Grammont M, Irvine KD (2000) Roles for scalloped and vestigial in regulating cell affinity and interactions between the wing blade and the wing hinge. *Dev Biol* 228:287–303
- Liu CY, Zha ZY, Zhou X, Zhang H, Huang W, Zhao D, Li T, Chan SW, Lim CJ, Hong W et al (2010) The hippo tumor pathway promotes TAZ degradation by phosphorylating a phosphodegron and recruiting the SCF{beta}-TrCP E3 ligase. *J Biol Chem* 285:37159–37169
- Liu C, Huang W, Lei Q (2011) Regulation and function of the TAZ transcription co-activator. *Int J Biochem Mol Biol* 2:247–256
- Liu AM, Wong KF, Jiang X, Qiao Y, Luk JM (2012a) Regulators of mammalian Hippo pathway in cancer. *Biochim Biophys Acta* 1826:357–364
- Liu H, Jiang D, Chi F, Zhao B (2012b) The Hippo pathway regulates stem cell proliferation, self-renewal, and differentiation. *Protein Cell* 3:291–304
- Loewith R (2011) A brief history of TOR. *Biochem Soc Trans* 39:437–442
- Loewith R, Jacinto E, Wullschleger S, Lorberg A, Crespo JL, Bonenfant D, Oppliger W, Jenoe P, Hall MN (2002) Two TOR complexes, only one of which is rapamycin sensitive, have distinct roles in cell growth control. *Mol Cell* 10:457–468
- Long X, Muller F, Avruch J (2004) TOR action in mammalian cells and in *Caenorhabditis elegans*. *Curr Top Microbiol Immunol* 279:115–138
- Mahoney PA, Weber U, Onofrechuk P, Biessmann H, Bryant PJ, Goodman CS (1991) The fat tumor suppressor gene in *Drosophila* encodes a novel member of the cadherin gene superfamily. *Cell* 67:853–868

- Maitra S, Kulikauskas RM, Gavilan H, Fehon RG (2006) The tumor suppressors Merlin and Expanded function cooperatively to modulate receptor endocytosis and signaling. *Curr Biol* 16(7):702–709
- Mao Y, Rauskolb C, Cho E, Hu WL, Hayter H, Minihan G, Katz FN, Irvine KD (2006) Dachs: an unconventional myosin that functions downstream of Fat to regulate growth, affinity and gene expression in *Drosophila*. *Development* 133:2539–2551
- Mao Y, Kucuk B, Irvine KD (2009) *Drosophila* lowfat, a novel modulator of Fat signaling. *Development* 136:3223–3233
- Martin FA, Perez-Garijo A, Morata G (2009) Apoptosis in *Drosophila*: compensatory proliferation and undead cells. *Int J Dev Biol* 53:1341–1347
- Marygold SJ, Leever SJ (2002) Growth signaling: TSC takes its place. *Curr Biol* 12:R785–R787
- Matakatsu H, Blair SS (2004) Interactions between Fat and Dachsous and the regulation of planar cell polarity in the *Drosophila* wing. *Development* 131:3785–3794
- Matakatsu H, Blair SS (2006) Separating the adhesive and signaling functions of the Fat and Dachsous protocadherins. *Development* 133:2315–2324
- Matakatsu H, Blair SS (2008) The DHHC palmitoyltransferase approximated regulates Fat signaling and Dachs localization and activity. *Curr Biol* 18:1390–1395
- Matakatsu H, Blair SS (2012) Separating planar cell polarity and Hippo pathway activities of the protocadherins Fat and Dachsous. *Development* 139:1498–1508
- Mauviel A, Nallet-Staub F, Varelas X (2012) Integrating developmental signals: a hippo in the (path)way. *Oncogene* 31(14):1743–1756
- McCartney BM, Kulikauskas RM, LaJeunesse DR, Fehon RG (2000) The Neurofibromatosis-2 homologue, Merlin, and the tumor suppressor expanded function together in *Drosophila* to regulate cell proliferation and differentiation. *Development* 127:1315–1324
- Meignin C, Alvarez-Garcia I, Davis I, Palacios IM (2007) The salvador-warts-hippo pathway is required for epithelial proliferation and axis specification in *Drosophila*. *Curr Biol* 17:1871–1878
- Mikeladze-Dvali T, Wernet M, Desplan C (2005) Warts and Melted regulate subset-specific opsin expression in R8 photoreceptors. *A Dros Res Conf* 46:396C
- Mills JR, Hippo Y, Robert F, Chen SM, Malina A, Lin CJ, Trojahn U, Wendel HG, Charest A, Bronson RT et al (2008) mTORC1 promotes survival through translational control of Mcl-1. *Proc Natl Acad Sci USA* 105:10853–10858
- Mitchison JM, Novak B, Sveiczler A (1997) Size control in the cell cycle. *Cell Biol Int* 21:461–463
- Mohr OL (1923) Modifications of the sex-ratio through a sex-linked semi-lethal in *Drosophila melanogaster*. *Studia Mendeliana*, Brunn 266–287
- Mohr OL (1929) Exaggeration and inhibition phenomena encountered in the analysis of an autosomal dominant. *Z. Indukt Abstamm. VererbLehre* 50:113–200
- Montagne J (2000) Genetic and molecular mechanisms of cell size control. *Mol Cell Biol Res Commun* 4:195–202
- Montagne J, Stewart MJ, Stocker H, Hafen E, Kozma SC, Thomas G (1999) *Drosophila* S6 kinase: a regulator of cell size. *Science* 285:2126–2129
- Montagne J, Radimerski T, Thomas G (2001) Insulin signaling: lessons from the *Drosophila* tuberous sclerosis complex, a tumor suppressor. *Sci STKE* 2001:pe36
- Morrison H, Sherman LS, Legg J, Banine F, Isacke C, Haipek CA, Gutmann DH, Ponta H, Herrlich P (2001) The NF2 tumor suppressor gene product, merlin, mediates contact inhibition of growth through interactions with CD44. *Genes Dev* 15:968–980
- Nam SC, Choi KW (2003) Interaction of Par-6 and Crumbs complexes is essential for photoreceptor morphogenesis in *Drosophila*. *Development* 130(18):4363–4372
- Nam SC, Choi KW (2006) Domain-specific early and late function of Dpatj in *Drosophila* photoreceptor cells. *Dev Dyn* 235(6):1501–1507
- Nellist M, Sancak O, Goedbloed M, Adriaans A, Wessels M, Maat-Kievit A, Baars M, Dommering C, van den Ouweland A, Halley D (2008) Functional characterisation of the TSC1-TSC2 com-

- plex to assess multiple TSC2 variants identified in single families affected by tuberous sclerosis complex. *BMC Med Genet* 9:10
- Neto-Silva RM, de Beco S, Johnston LA (2010) Evidence for a growth-stabilizing regulatory feedback mechanism between Myc and Yorkie, the *Drosophila* homolog of Yap. *Dev Cell* 19:507–520
- Neufeld TP (2003) Body building: regulation of shape and size by PI3K/TOR signaling during development. *Mech Dev* 120(11):1283–96. Review. PMID: 14623438
- Newsome TP, Asling B, Dickson BJ (2000) Analysis of *Drosophila* photoreceptor axon guidance in eye-specific mosaics. *Development* 127:851–860
- Nolo R, Morrison CM, Tao C, Zhang X, Halder G (2006) The bantam microRNA is a target of the hippo tumor-suppressor pathway. *Curr Biol* 16:1895–1904
- O'Neill E, Kolch W (2005) Taming the Hippo: Raf-1 controls apoptosis by suppressing MST2/Hippo. *Cell Cycle* 4:365–367
- Ogawa H, Ohta N, Moon W, Matsuzaki F (2009) Protein phosphatase 2A negatively regulates aPKC signaling by modulating phosphorylation of Par-6 in *Drosophila* neuroblast asymmetric divisions. *J Cell Sci* 122(18):3242–3249
- Oh H, Irvine KD (2008) In vivo regulation of Yorkie phosphorylation and localization. *Development* 135:1081–1088
- Oh H, Irvine KD (2009) In vivo analysis of Yorkie phosphorylation sites. *Oncogene* 28:1916–1927
- Oh H, Irvine KD (2010) Yorkie: the final destination of Hippo signaling. *Trends Cell Biol* 20:410–417
- Oh H, Reddy BV, Irvine KD (2009) Phosphorylation-independent repression of Yorkie in Fat-Hippo signaling. *Dev Biol* 335(1):188–197
- Oka T, Remue E, Meerschaert K, Vanloo B, Boucherie C, Gfeller D, Bader G, Sidhu S, Vandekerckhove J, Gettemans J et al (2010) Functional complex between YAP2 and ZO-2 is PDZ domain dependent, regulates YAP2 nuclear localization and signaling. *Biochem J* 432(3):461–472
- Oldham S, Hafen E (2003) Insulin/IGF and target of rapamycin signaling: a TOR de force in growth control. *Trends Cell Biol* 13:79–85
- Oldham S, Bohni R, Stocker H, Brogiolo W, Hafen E (2000a) Genetic control of size in *Drosophila*. *Philos Trans R Soc Lond Ser B Biol Sci* 355:945–952
- Oldham S, Montagne J, Radimerski T, Thomas G, Hafen E (2000b) Genetic and biochemical characterization of dTOR, the *Drosophila* homolog of the target of rapamycin. *Genes Dev* 14:2689–2694
- Oldham S, Stocker H, Laffargue M, Wittwer F, Wymann M, Hafen E (2002) The *Drosophila* insulin/IGF receptor controls growth and size by modulating PtdInsP(3) levels. *Development* 129:4103–4109
- Pagliarini RA, Quinones AT, Xu T (2003) Analyzing the function of tumor suppressor genes using a *Drosophila* model. *Methods Mol Biol* 223:349–382
- Pan D (2007) Hippo signaling in organ size control. *Genes Dev* 21:886–897
- Pan D (2010) The hippo signaling pathway in development and cancer. *Dev Cell* 19:491–505
- Pan D, Dong J, Zhang Y, Gao X (2004) Tuberous sclerosis complex: from *Drosophila* to human disease. *Trends Cell Biol* 14:78–85
- Pantalacci S, Tapon N, Leopold P (2003) The Salvador partner Hippo promotes apoptosis and cell-cycle exit in *Drosophila*. *Nat Cell Biol* 5:921–927
- Paramasivam M, Sarkeshik A, Yates JR 3rd, Fernandes MJ, McCollum D (2011) Angiotensin family proteins are novel activators of the LATS2 kinase tumor suppressor. *Mol Biol Cell* 22:3725–3733
- Pearce LR, Komander D, Alessi DR (2010) The nuts and bolts of AGC protein kinases. *Nat Rev Mol Cell Biol* 11:9–22
- Pellock BJ, Buff E, White K, Hariharan IK (2007) The *Drosophila* tumor suppressors expanded and Merlin differentially regulate cell cycle exit, apoptosis, and Wingless signaling. *Dev Biol* 304:102–115

- Peng HW, Slattery M, Mann RS (2009) Transcription factor choice in the Hippo signaling pathway: homothorax and yorkie regulation of the microRNA bantam in the progenitor domain of the *Drosophila* eye imaginal disc. *Genes Dev* 23:2307–2319
- Penton A, Selleck SB, Hoffmann FM (1997) Regulation of cell cycle synchronization by decapentaplegic during *Drosophila* eye development. *Science* 275:203–206
- Pfeiffer BD, Ngo TT, Hibbard KL, Murphy C, Jenett A, Truman JW, Rubin GM (2010) Refinement of tools for targeted gene expression in *Drosophila*. *Genetics* 186:735–755
- Pichaud F, Desplan C (2001) A new visualization approach for identifying mutations that affect differentiation and organization of the *Drosophila* ommatidia. *Development* 128(6):815–826
- Poernbacher I, Baumgartner R, Marada SK, Edwards K, Stocker H (2012) *Drosophila* Pez acts in hippo signaling to restrict intestinal stem cell proliferation. *Curr Biol* 22(5):389–396
- Polesello C, Tapon N (2007) Salvador-warts-hippo signaling promotes *Drosophila* posterior follicle cell maturation downstream of notch. *Curr Biol* 17:1864–1870
- Polesello C, Huelsmann S, Brown NH, Tapon N (2006) The *Drosophila* RASSF homolog antagonizes the hippo pathway. *Curr Biol* 16:2459–2465
- Poltilov RM, Jacobs AR, Haft CR, Xu P, Taylor SI (2000) Characterization of *Drosophila* insulin receptor substrate. *J Biol Chem* 275:23346–23354
- Poon CL, Lin JI, Zhang X, Harvey KF (2011) The sterile 20-like kinase Tao-1 controls tissue growth by regulating the Salvador-Warts-Hippo pathway. *Dev Cell* 21:896–906
- Potter CJ, Xu T (2001) Mechanisms of size control. *Curr Opin Genet Dev* 11:279–286
- Potter CJ, Huang H, Xu T (2001) *Drosophila* Tsc1 functions with Tsc2 to antagonize insulin signaling in regulating cell growth, cell proliferation, and organ size. *Cell* 105:357–368
- Potter CJ, Pedraza LG, Xu T (2002) Akt regulates growth by directly phosphorylating Tsc2. *Nat Cell Biol* 4:658–665
- Potter CJ, Pedraza LG, Huang H, Xu T (2003) The tuberous sclerosis complex (TSC) pathway and mechanism of size control. *Biochem Soc Trans* 31:584–586
- Price DM, Jin Z, Rabinovitch S, Campbell SD (2002) Ectopic expression of the *Drosophila* Cdk1 inhibitory kinases, Wee1 and Myt1, interferes with the second mitotic wave and disrupts pattern formation during eye development. *Genetics* 161:721–731
- Radimerski T, Montagne J, Hemmings-Mieszczak M, Thomas G (2002a) Lethality of *Drosophila* lacking TSC tumor suppressor function rescued by reducing dS6K signaling. *Genes Dev* 16:2627–2632
- Radimerski T, Montagne J, Rintelen F, Stocker H, van der Kaay J, Downes CP, Hafen E, Thomas G (2002b) dS6K-regulated cell growth is dPKB/dPI(3)K-independent, but requires dPDK1. *Nat Cell Biol* 4:251–255
- Raff MC (1996) Size control: the regulation of cell numbers in animal development. *Cell* 86:173–175
- Rauskolb C, Pan G, Reddy BV, Oh H, Irvine KD (2011) Zyxin links fat signaling to the hippo pathway. *PLoS Biol* 9:e1000624
- Reddy BV, Irvine KD (2008) The Fat and Warts signaling pathways: new insights into their regulation, mechanism and conservation. *Development* 135:2827–2838
- Reddy BV, Rauskolb C, Irvine KD (2010) Influence of fat-hippo and notch signaling on the proliferation and differentiation of *Drosophila* optic neuroepithelia. *Development* 137:2397–2408
- Ren F, Wang B, Yue T, Yun EY, Ip YT, Jiang J (2010a) Hippo signaling regulates *Drosophila* intestine stem cell proliferation through multiple pathways. *Proc Natl Acad Sci USA* 107:21064–21069
- Ren F, Zhang L, Jiang J (2010b) Hippo signaling regulates Yorkie nuclear localization and activity through 14-3-3 dependent and independent mechanisms. *Dev Biol* 337:303–312
- Ribeiro PS, Josue F, Wepf A, Wehr MC, Rinner O, Kelly G, Tapon N, Gstaiger M (2010) Combined functional genomic and proteomic approaches identify a PP2A complex as a negative regulator of Hippo signaling. *Mol Cell* 39:521–534
- Richardson HE (2011) Actin up for Hippo. *EMBO J* 30:2307–2309
- Richardson H, Kumar S (2002) Death to flies: *Drosophila* as a model system to study programmed cell death. *J Immunol Methods* 265:21–38

- Rintelen F, Stocker H, Thomas G, Hafen E (2001) PDK1 regulates growth through Akt and S6K in *Drosophila*. Proc Natl Acad Sci USA 98:15020–15025
- Robinson BS, Huang J, Hong Y, Moberg KH (2010) Crumbs regulates Salvador/Warts/Hippo signaling in *Drosophila* via the FERM-domain protein expanded. Curr Biol 20:582–590
- Rogulja D, Rauskolb C, Irvine KD (2008) Morphogen control of wing growth through the Fat signaling pathway. Dev Cell 15:309–321
- Rothenberg ME, Jan YN (2002) Salvador – the persistence of proliferation. Cancer Cell 2:171–173
- Rubin GM (1989) Development of the *Drosophila* retina: inductive events studied at single cell resolution. Cell 57:519–520
- Rulifson EJ, Kim SK, Nusse R (2002) Ablation of insulin-producing neurons in flies: growth and diabetic phenotypes. Science 296:1118–1120
- Rusconi JC, Hays R, Cagan RL (2000) Programmed cell death and patterning in *Drosophila*. Cell Death Differ 7:1063–1070
- Salah Z, Melino G, Aqeilan RI (2011) Negative regulation of the Hippo pathway by E3 ubiquitin ligase ITCH is sufficient to promote tumorigenicity. Cancer Res 71:2010–2020
- Sancak Y, Bar-Peled L, Zoncu R, Markhard AL, Nada S, Sabatini DM (2010) Ragulator-Rag complex targets mTORC1 to the lysosomal surface and is necessary for its activation by amino acids. Cell 141:290–303
- Sansores-Garcia L, Bossuyt W, Wada K, Yonemura S, Tao C, Sasaki H, Halder G (2011) Modulating F-actin organization induces organ growth by affecting the Hippo pathway. EMBO J 30:2325–2335
- Sarbassov DD, Ali SM, Kim DH, Guertin DA, Latek RR, Erdjument-Bromage H, Tempst P, Sabatini DM (2004) Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. Curr Biol 14:1296–1302
- Saucedo LJ, Edgar BA (2007) Filling out the Hippo pathway. Nat Rev Mol Cell Biol 8:613–621
- Saucedo LJ, Gao X, Chiarelli DA, Li L, Pan D, Edgar BA (2003) Rheb promotes cell growth as a component of the insulin/TOR signalling network. Nat Cell Biol 5:566–571
- Sawamoto K, Okano H (1996) Cell-cell interactions during neural development: multiple types of lateral inhibitions involved in *Drosophila* eye development. Neurosci Res 26:205–214
- Schagdarsurengin U, Richter AM, Hornung J, Lange C, Steinmann K, Dammann RH (2010) Frequent epigenetic inactivation of RASSF2 in thyroid cancer and functional consequences. Mol Cancer 9:264
- Scheel H, Hofmann K (2003) A novel interaction motif, SARAH, connects three classes of tumor suppressor. Curr Biol 13:R899–R900
- Schlegelmilch K, Mohseni M, Kirak O, Pruszk J, Rodriguez JR, Zhou D, Kreger BT, Vasioukhin V, Avruch J, Brummelkamp TR et al (2011) Yap1 acts downstream of alpha-catenin to control epidermal proliferation. Cell 144:782–795
- Schroeder MC, Halder G (2012) Regulation of the Hippo pathway by cell architecture and mechanical signals. Semin Cell Dev Biol 23(7):803–811
- Seidel C, Schagdarsurengin U, Blumke K, Wurl P, Pfeifer GP, Hauptmann S, Taubert H, Dammann R (2007) Frequent hypermethylation of MST1 and MST2 in soft tissue sarcoma. Mol Carcinog 46:865–871
- Sekido Y (2008) Molecular biology of malignant mesothelioma. Environ Health Prev Med 13:65–70
- Sekido Y (2011) Inactivation of Merlin in malignant mesothelioma cells and the Hippo signaling cascade dysregulation. Pathol Int 61:331–344
- Shaw RL, Kohlmaier A, Polesello C, Veelken C, Edgar BA, Tapon N (2010) The Hippo pathway regulates intestinal stem cell proliferation during *Drosophila* adult midgut regeneration. Development 137:4147–4158
- Shimizu T, Ho LL, Lai ZC (2008) The mob as tumor suppressor gene is essential for early development and regulates tissue growth in *Drosophila*. Genetics 178:957–965
- Silva E, Tsatskis Y, Gardano L, Tapon N, McNeill H (2006) The tumor-suppressor gene fat controls tissue growth upstream of expanded in the hippo signaling pathway. Curr Biol 16:2081–2089

- Silvis MR, Kreger BT, Lien WH, Klezovitch O, Rudakova GM, Camargo FD, Lantz DM, Seykora JT, Vasioukhin V (2011) Alpha-catenin is a tumor suppressor that controls cell accumulation by regulating the localization and activity of the transcriptional coactivator Yap1. *Sci Signal* 4:ra33
- Simon MA, Xu A, Ishikawa HO, Irvine KD (2010) Modulation of fat: dachsous binding by the cadherin domain kinase four-jointed. *Curr Biol* 20:811–817
- Singh A, Kango-Singh M, Choi KW, Sun YH (2004) Dorso-ventral asymmetric functions of teashirt in *Drosophila* eye development depend on spatial cues provided by early DV patterning genes. *Mech Dev* 121(4):365–370
- Singh A, Tare M, Puli OR, Kango-Singh M (2012) A glimpse into dorso-ventral patterning of the *Drosophila* eye. *Dev Dyn* 241:69–84
- Skouloudaki K, Walz G (2012) YAP1 recruits c-Abl to protect angiominin-like 1 from Nedd4-mediated degradation. *PLoS One* 7:e35735
- Skouloudaki K, Puetz M, Simons M, Courbard JR, Boehlke C, Hartleben B, Engel C, Moeller MJ, Englert C, Bollig F et al (2009) Scribble participates in Hippo signaling and is required for normal zebrafish pronephros development. *Proc Natl Acad Sci USA* 106:8579–8584
- Sopko R, Silva E, Clayton L, Gardano L, Barrios-Rodiles M, Wrana J, Varelas X, Arbouzova NI, Shaw S, Saburi S et al (2009) Phosphorylation of the tumor suppressor fat is regulated by its ligand Dachsous and the kinase discs overgrown. *Curr Biol* 19:1112–1117
- Soulard A, Cohen A, Hall MN (2009) TOR signaling in invertebrates. *Curr Opin Cell Biol* 21:825–836
- St Johnston D (2002) The art and design of genetic screens: *Drosophila melanogaster*. *Nat Rev Genet* 3:176–188
- Staley BK, Irvine KD (2010) Warts and Yorkie mediate intestinal regeneration by influencing stem cell proliferation. *Curr Biol* 20:1580–1587
- Staley BK, Irvine KD (2012) Hippo signaling in *Drosophila*: recent advances and insights. *Dev Dyn* 241:3–15
- Stocker H, Hafen E (2000) Genetic control of cell size. *Curr Opin Genet Dev* 10:529–535
- Stocker H, Radimerski T, Schindelhof B, Wittwer F, Belawat P, Daram P, Breuer S, Thomas G, Hafen E (2003) Rheb is an essential regulator of S6K in controlling cell growth in *Drosophila*. *Nat Cell Biol* 5:559–565
- Strassburger K, Tiebe M, Pinna F, Breuhahn K, Teleman AA (2012) Insulin/IGF signaling drives cell proliferation in part via Yorkie/YAP. *Dev Biol* 367:187–196
- Striedinger K, VandenBerg SR, Baia GS, McDermott MW, Gutmann DH, Lal A (2008) The neurofibromatosis 2 tumor suppressor gene product, merlin, regulates human meningioma cell growth by signaling through YAP. *Neoplasia* 10:1204–1212
- Strutt H, Price MA, Strutt D (2006) Planar polarity is positively regulated by casein kinase Iepsilon in *Drosophila*. *Curr Biol* 16(13):1329–1336
- Su TT, O'Farrell PH (1998) Size control: cell proliferation does not equal growth. *Curr Biol* 8:R687–R689
- Sudol M (2010) Newcomers to the WW domain-mediated network of the Hippo tumor suppressor pathway. *Genes Cancer* 1:1115–1118
- Sudol M, Harvey KF (2010) Modularity in the Hippo signaling pathway. *Trends Biochem Sci* 35(11):627–633
- Sun Q (2007) The mechanism of pattern formation in the developing *Drosophila* retina. *Sci China C Life Sci* 50:120–124
- Sun G, Irvine KD (2010) Regulation of Hippo signaling by Jun kinase signaling during compensatory cell proliferation and regeneration, and in neoplastic tumors. *Dev Biol* 350(1):139–151
- Sun G, Irvine KD (2011) Regulation of Hippo signaling by Jun kinase signaling during compensatory cell proliferation and regeneration, and in neoplastic tumors. *Dev Biol* 350(1):139–151
- Tamori Y, Bialucha CU, Tian AG, Kajita M, Huang YC, Norman M, Harrison N, Poulton J, Ivanovitch K, Disch L, Liu T, Deng WM, Fujita Y (2010) Involvement of Lgl and Mahjong/VprBP in cell competition. *PLoS Biol* 8(7)

- Tapon N, Ito N, Dickson BJ, Treisman JE, Hariharan IK (2001) The *Drosophila* tuberous sclerosis complex gene homologs restrict cell growth and cell proliferation. *Cell* 105:345–355
- Tapon N, Harvey K, Bell D, Wahrer D, Schiripo T, Haber D, Hariharan I (2002) Salvador promotes both cell cycle exit and apoptosis in *Drosophila* and is mutated in human cancer cell lines. *Cell* 110:467
- Tato I, Bartrons R, Ventura F, Rosa JL (2011) Amino acids activate mammalian target of rapamycin complex 2 (mTORC2) via PI3K/Akt signaling. *J Biol Chem* 286:6128–6142
- Tepass U, Theres C, Knust E (1990) Crumbs encodes an EGF-like protein expressed on apical membranes of *Drosophila* epithelial cells and required for organization of epithelia. *Cell* 61(5):787–799
- Thomas C, Strutt D (2012) The roles of the cadherins Fat and Dachshous in planar polarity specification in *Drosophila*. *Dev Dyn* 241:27–39
- Thompson BJ, Cohen SM (2006) The Hippo pathway regulates the bantam microRNA to control cell proliferation and apoptosis in *Drosophila*. *Cell* 126:767–774
- Torok T, Tick G, Alvarado M, Kiss I (1993) P-lacW insertional mutagenesis on the second chromosome of *Drosophila melanogaster*: isolation of lethals with different overgrowth phenotypes. *Genetics* 135(1):71–80
- Toyouka S, Ouchida M, Jitsumori Y, Tsukuda K, Sakai A, Nakamura A, Shimizu N, Shimizu K (2000) HD-PTP: a novel protein tyrosine phosphatase gene on human chromosome 3p21.3. *Biochem Biophys Res Commun* 278:671–678
- Tran H, Brunet A, Griffith EC, Greenberg ME (2003) The many forks in FOXO's road. *Sci STKE* 2003:RE5
- Treins C, Warne PH, Magnuson MA, Pende M, Downward J (2010) Rictor is a novel target of p70 S6 kinase-1. *Oncogene* 29:1003–1016
- Treisman JE, Heberlein U (1998) Eye development in *Drosophila*: formation of the eye field and control of differentiation. *Curr Top Dev Biol* 39:119–158
- Tsachaki M, Sprecher SG (2012) Genetic and developmental mechanisms underlying the formation of the *Drosophila* compound eye. *Dev Dyn* 241:40–56
- Tsai BP, Hoverter NP, Waterman ML (2012) Blending hippo and WNT: sharing messengers and regulation. *Cell* 151:1401–1403
- Tumaneng K, Russell RC, Guan KL (2012a) Organ size control by Hippo and TOR pathways. *Curr Biol* 22:R368–R379
- Tumaneng K, Schlegelmilch K, Russell RC, Yimlamai D, Basnet H, Mahadevan N, Fitamant J, Bardeesy N, Camargo FD, Guan KL (2012b) YAP mediates crosstalk between the Hippo and PI(3)K-TOR pathways by suppressing PTEN via miR-29. *Nat Cell Biol* 14:1322–1329
- Tyler DM, Baker NE (2007) Expanded and fat regulate growth and differentiation in the *Drosophila* eye through multiple signaling pathways. *Dev Biol* 305:187–201
- Tyler DM, Li W, Zhuo N, Pellock B, Baker NE (2007) Genes affecting cell competition in *Drosophila*. *Genetics* 175:643–657
- Udan RS, Kango-Singh M, Nolo R, Tao C, Halder G (2003) Hippo promotes proliferation arrest and apoptosis in the Salvador/Warts pathway. *Nat Cell Biol* 5(10):914–920
- Vanhaesebroeck B, Alessi DR (2000) The PI3K-PDK1 connection: more than just a road to PKB. *Biochem J* 346(Pt 3):561–576
- Varelas X, Wrana JL (2012) Coordinating developmental signaling: novel roles for the Hippo pathway. *Trends Cell Biol* 22:88–96
- Varelas X, Miller BW, Sopko R, Song S, Gregorieff A, Fellouse FA, Sakuma R, Pawson T, Hunziker W, McNeill H, Wrana JL, Attisano L (2010) The Hippo pathway regulates Wnt/beta-catenin signaling. *Dev Cell* 18(4):579–591
- Varelas X, Samavarchi-Tehrani P, Narimatsu M, Weiss A, Cockburn K, Larsen BG, Rossant J, Wrana JL (2010b) The crumbs complex couples cell density sensing to Hippo-dependent control of the TGF-beta-SMAD pathway. *Dev Cell* 19:831–844
- Venken KJ, Bellen HJ (2012) Genome-wide manipulations of *Drosophila melanogaster* with transposons, Flp recombinase, and PhiC31 integrase. *Methods Mol Biol* 859:203–228



- Verdu J, Buratovich MA, Wilder EL, Birnbaum MJ (1999) Cell-autonomous regulation of cell and organ growth in *Drosophila* by Akt/PKB. *Nat Cell Biol* 1:500–506
- Vergheze S, Bedi S, Kango-Singh M (2012a) Hippo signalling controls Dronc activity to regulate organ size in *Drosophila*. *Cell Death Differ* 19(10):1664–1676
- Vergheze S, Waghmare I, Kwon H, Hanes K, Kango-Singh M (2012b) Scribble acts in the *Drosophila* fat-hippo pathway to regulate warts activity. *PLoS One* 7:e47173
- Vidal M, Cagan RL (2006) *Drosophila* models for cancer research. *Curr Opin Genet Dev* 16:10–16
- Vigneron AM, Ludwig RL, Vousden KH (2010) Cytoplasmic ASPP1 inhibits apoptosis through the control of YAP. *Genes Dev* 24:2430–2439
- Villano JL, Katz FN (1995) Four-jointed is required for intermediate growth in the proximal-distal axis in *Drosophila*. *Development* 121(9):2767–2777
- Visser-Grievé S, Hao Y, Yang X (2012) Human homolog of *Drosophila* expanded, hEx, functions as a putative tumor suppressor in human cancer cell lines independently of the Hippo pathway. *Oncogene* 31:1189–1195
- Wang K, Degerny C, Xu M, Yang XJ (2009) YAP, TAZ, and Yorkie: a conserved family of signal-responsive transcriptional coregulators in animal development and human disease. *Biochem Cell Biol* 87:77–91
- Wang C, An J, Zhang P, Xu C, Gao K, Wu D, Wang D, Yu H, Liu JO, Yu L (2012a) The Nedd4-like ubiquitin E3 ligases target angiomin/p130 to ubiquitin-dependent degradation. *Biochem J* 444:279–289
- Wang T, Blumhagen R, Lao U, Kuo Y, Edgar BA (2012b) LST8 regulates cell growth via target-of-rapamycin complex 2 (TORC2). *Mol Cell Biol* 32:2203–2213
- Wehr MC, Holder MV, Gailite I, Saunders RE, Maile TM, Ciirdaeva E, Instrell R, Jiang M, Howell M, Rossner MJ et al (2013) Salt-inducible kinases regulate growth through the Hippo signaling pathway in *Drosophila*. *Nat Cell Biol* 15(1):61–71
- Wei X, Shimizu T, Lai ZC (2007) Mob as tumor suppressor is activated by Hippo kinase for growth inhibition in *Drosophila*. *EMBO J* 26:1772–1781
- Weinkove D, Neufeld TP, Twardzik T, Waterfield MD, Leever SJ (1999) Regulation of imaginal disc cell size, cell number and organ size by *Drosophila* class I(A) phosphoinositide 3-kinase and its adaptor. *Curr Biol* 9:1019–1029
- Wernet MF, Labhart T, Baumann F, Mazzoni EO, Pichaud F, Desplan C (2003) Homothorax switches function of *Drosophila* photoreceptors from color to polarized light sensors. *Cell* 115(3):267–279
- Willecke M, Hamaratoglu F, Kango-Singh M, Udan R, Chen CL, Tao C, Zhang X, Halder G (2006) The Fat Cadherin acts through the Hippo tumor-suppressor pathway to regulate tissue size. *Curr Biol* 16(21):2090–2100
- Willecke M, Hamaratoglu F, Sansores-Garcia L, Tao C, Halder G (2008) Boundaries of Dachous Cadherin activity modulate the Hippo signaling pathway to induce cell proliferation. *Proc Natl Acad Sci USA* 105:14897–14902
- Wolff T, Ready DF (1991) The beginning of pattern formation in the *Drosophila* compound eye: the morphogenetic furrow and the second mitotic wave. *Development* 113:841–850
- Wolff T, Ready DF (1993) Pattern formation in the *Drosophila* retina. In: Bate M, Martinez Arias A (eds) *The development of Drosophila melanogaster*. Cold Spring Harbor Laboratory Press, New York, pp 1277–1325
- Wu S, Huang J, Dong J, Pan D (2003) Hippo encodes a Ste-20 family protein kinase that restricts cell proliferation and promotes apoptosis in conjunction with salvador and warts. *Cell* 114:445–456
- Wu S, Liu Y, Zheng Y, Dong J, Pan D (2008) The TEAD/TEF family protein Scalloped mediates transcriptional output of the Hippo growth-regulatory pathway. *Dev Cell* 14:388–398
- Xiao L, Chen Y, Ji M, Dong J (2011) KIBRA regulates Hippo signaling activity via interactions with large tumor suppressor kinases. *J Biol Chem* 286:7788–7796
- Xu T, Rubin GM (1993) Analysis of genetic mosaics in developing and adult *Drosophila* tissues. *Development* 117:1223–1237

- Xu T, Wang W, Zhang S, Stewart RA, Yu W (1995) Identifying tumor suppressors in genetic mosaics: the *Drosophila* *lats* gene encodes a putative protein kinase. *Development* 121:1053–1063
- Yamamoto D (1993) Positive and negative signaling mechanisms in the regulation of photoreceptor induction in the developing *Drosophila* retina. *Review. Genetica* 88:153–164
- Yang CH, Axelrod JD, Simon MA (2002) Regulation of frizzled by fat-like cadherins during planar polarity signaling in the *Drosophila* compound eye. *Cell* 108(5):675–688
- Yang Q, Inoki K, Kim E, Guan KL (2006) TSC1/TSC2 and Rheb have different effects on TORC1 and TORC2 activity. *Proc Natl Acad Sci USA* 103:6811–6816
- Yu J, Zheng Y, Dong J, Klusza S, Deng WM, Pan D (2010) Kibra functions as a tumor suppressor protein that regulates Hippo signaling in conjunction with Merlin and expanded. *Dev Cell* 18:288–299
- Yu FX, Zhao B, Panupinthu N, Jewell JL, Lian I, Wang LH, Zhao J, Yuan H, Tumaneng K, Li H et al (2012) Regulation of the Hippo-YAP pathway by G-protein-coupled receptor signaling. *Cell* 150:780–791
- Yue T, Tian A, Jiang J (2012) The cell adhesion molecule echinoid functions as a tumor suppressor and upstream regulator of the Hippo signaling pathway. *Dev Cell* 22:255–267
- Zecca M, Struhl G (2010) A feed-forward circuit linking wingless, fat-dachsous signaling, and the warts-hippo pathway to *Drosophila* wing growth. *PLoS Biol* 8:e1000386
- Zeitler J, Hsu CP, Dionne H, Bilder D (2004) Domains controlling cell polarity and proliferation in the *Drosophila* tumor suppressor Scribble. *J Cell Biol* 167(6):1137–1146
- Zhang H, Stallock JP, Ng JC, Reinhard C, Neufeld TP (2000) Regulation of cellular growth by the *Drosophila* target of rapamycin dTOR. *Genes Dev* 14:2712–2724
- Zhang Y, Gao X, Saucedo LJ, Ru B, Edgar BA, Pan D (2003) Rheb is a direct target of the tuberous sclerosis tumour suppressor proteins. *Nat Cell Biol* 5:578–581
- Zhang J, Smolen GA, Haber DA (2008a) Negative regulation of YAP by LATS1 underscores evolutionary conservation of the *Drosophila* Hippo pathway. *Cancer Res* 68:2789–2794
- Zhang L, Ren F, Zhang Q, Chen Y, Wang B, Jiang J (2008b) The TEAD/TEF family of transcription factor Scalloped mediates Hippo signaling in organ size control. *Dev Cell* 14:377–387
- Zhang L, Yue T, Jiang J (2009a) Hippo signaling pathway and organ size control. *Fly (Austin)* 3:68–73
- Zhang X, Milton CC, Humbert PO, Harvey KF (2009b) Transcriptional output of the Salvador/warts/hippo pathway is controlled in distinct fashions in *Drosophila melanogaster* and mammalian cell lines. *Cancer Res* 69:6033–6041
- Zhang X, George J, Deb S, Degoutin JL, Takano EA, Fox SB, Bowtell DD, Harvey KF (2011a) The Hippo pathway transcriptional co-activator, YAP, is an ovarian cancer oncogene. *Oncogene* 30:2810–2822
- Zhang X, Milton CC, Poon CL, Hong W, Harvey KF (2011b) Wbp2 cooperates with Yorkie to drive tissue growth downstream of the Salvador-Warts-Hippo pathway. *Cell Death Differ* 18:1346–1355
- Zhang L, Iyer J, Chowdhury A, Ji M, Xiao L, Yang S, Chen Y, Tsai MY, Dong J (2012) KIBRA regulates aurora kinase activity and is required for precise chromosome alignment during mitosis. *J Biol Chem* 287:34069–34077
- Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, Xie J, Ikenoue T, Yu J, Li L et al (2007) Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control. *Genes Dev* 21:2747–2761
- Zhao B, Lei QY, Guan KL (2008a) The Hippo-YAP pathway: new connections between regulation of organ size and cancer. *Curr Opin Cell Biol* 20(6):638–646
- Zhao B, Li L, Lei Q, Guan KL (2010a) The Hippo-YAP pathway in organ size control and tumorigenesis: an updated version. *Genes Dev* 24:862–874
- Zhao B, Li L, Tumaneng K, Wang CY, Guan KL (2010b) A coordinated phosphorylation by Lats and CK1 regulates YAP stability through SCF(beta-TRCP). *Genes Dev* 24:72–85
- Zhao B, Li L, Lu Q, Wang LH, Liu CY, Lei Q, Guan KL (2011a) Angiomotin is a novel Hippo pathway component that inhibits YAP oncoprotein. *Genes Dev* 25:51–63

- Zhao B, Tumaneng K, Guan KL (2011b) The Hippo pathway in organ size control, tissue regeneration and stem cell self-renewal. *Nat Cell Biol* 13:877–883
- Zinzalla V, Stracka D, Oppliger W, Hall MN (2011) Activation of mTORC2 by association with the ribosome. *Cell* 144:757–768
- Ziosi M, Baena-Lopez LA, Grifoni D, Frolidi F, Pession A, Garoia F, Trotta V, Bellosta P, Cavicchi S (2010) dMyc functions downstream of Yorkie to promote the supercompetitive behavior of hippo pathway mutant cells. *PLoS Genet* 6:e1001140
- Zipursky SL (1989) Molecular and genetic analysis of *Drosophila* eye development: sevenless, bride of sevenless and rough. *Trends Neurosci* 12:183–189