



Coagulation

23

Ernest E. Moore

KPI title: Patients requiring an emergent operation on warfarin therapy should have an international normalized ratio (INR) measured at the time of initial assessment

Description	INR measurement
Rationale	An emergent operation can be complicated with bleeding which may be accentuated due to warfarin therapy because of inadequate concentrations of clotting factors II, VII, IX, and X.
Target	100% of patients undergoing an emergent operation on warfarin therapy have a preoperative INR measured.
KPI collection frequency	Annually
KPI reporting frequency	Annually
KPI calculation	Numerator: number of patients undergoing an emergent operation on warfarin therapy who have a preoperative INR measured Denominator: number of patients undergoing an emergent operation on warfarin therapy
Reporting aggregation	Hospital 1
Data source(s)	Emergency surgery database ICU database

The original version of the book was revised: Affiliation of one of the editors, Ernest E. Moore, has been updated. The correction to the book is available at https://doi.org/10.1007/978-3-030-49363-9_26

E. E. Moore (✉)
Ernest E Moore Shock Trauma Center, Denver Health, Denver, CO, USA
e-mail: Ernest.Moore@dhha.org

© World Society of Emergency Surgery and Donegal Clinical and Research Academy 2020 153
M. Sugrue et al. (eds.), *Resources for Optimal Care of Emergency Surgery*,
Hot Topics in Acute Care Surgery and Trauma,
https://doi.org/10.1007/978-3-030-49363-9_23