

Chapter 9

EDTA Chelation for Calcific Band Keratopathy



Alanna S. Nattis and Richard J. Nattis

Indications

Calcific band keratopathy causing significant visual disturbance, irritation, pain, and destruction of ocular surface

Essential Steps

1. Application of topical anesthetic to cornea
2. EDTA application to cornea
3. Removal of calcific keratopathy by Weck-Cel sponge, cotton-tipped applicator, or blade
4. Application of antibiotic and cycloplegic eye drops to eye
5. Placement of bandage contact lens

Complications

- Infection
- Inflammation
- Irregular corneal surface causing visual disturbances
- Persistence of corneal irritation
- Corneal abrasion

A. S. Nattis (✉)
SightMD, Babylon, NY, USA

R. J. Nattis
Good Samaritan Hospital, West Islip, NY, USA

- Corneal perforation
- Persistent corneal epithelial defect
- Recurrence of band keratopathy

Template Operative Dictation

Preoperative diagnosis: Calcific band keratopathy (*OD/OS*)

Procedure: EDTA chelation (*OD/OS*)

Postoperative diagnosis: *Same*

Indication: This ____-year-old (*male/female*) was found to have symptomatic calcific band keratopathy leading to symptoms of ocular discomfort and visual disturbance (*OD/OS*). After a detailed review of the risks, benefits, and alternatives, the patient elected to undergo the procedure.

Description of the procedure: The patient was identified in the holding area, and the (*right/left*) eye was marked with a marking pen. The patient was brought into the minor procedure room and placed in the supine position. Topical anesthetic was instilled $\times 2$ into the operative eye. The (*right/left*) eye was prepped and draped in the usual sterile fashion. The operating microscope was centered over the (*right/left*) eye, and an eyelid speculum was placed. A proper time-out was performed verifying correct patient, procedure, site, positioning, and special equipment prior to starting the case.

2% EDTA was drawn up into a tuberculin syringe and was used to saturate a Weck-Cel sponge. The saturated Weck-Cel sponge was applied to the area of band keratopathy for 1–2 min. The Weck-Cel sponge was then removed from the eye, and the band keratopathy was debrided using a Tooke spatula. Larger calcific plaques were removed using forceps. Application of EDTA was repeated in an area of persistent calcific keratopathy. This was then followed by repeated debridement until the calcium was sufficiently cleared from the ocular surface.

After all visible calcific band keratopathy was removed from the cornea, balanced saline solution was used to copiously irrigate the ocular surface. Following one drop each of cyclopentolate and an antibiotic, the eyelid speculum was removed. A bandage contact lens was placed on the eye. The patient tolerated the procedure well and was carefully monitored for any adverse reactions.

Additional Resource

<https://www.youtube.com/watch?v=BOeaT9YgcB0>.