

Opers for Higher States of the Quantum Boussinesq Model



Davide Masoero and Andrea Raimondo

Abstract We study the ODE/IM correspondence for all the states of the quantum Boussinesq model. We consider a particular class of third order linear ordinary differential operators and show that the generalised monodromy data of such operators provide solutions to the Bethe Ansatz equations of the Quantum Boussinesq model.

Keywords ODE/IM correspondence · Quantum Boussinesq · Opers · Bethe Ansatz

1 Introduction

The quantum Boussinesq model (Bazhanov et al. 2002) is a 2 dimensional conformal field theory with a \mathcal{W}_3 symmetry, and it can be exactly solved via the Bethe Ansatz equations. This model can be realised as the quantisation of a \mathfrak{sl}_3 Drinfeld-Sokolov hierarchy, or as the continuum limit of a \mathfrak{sl}_3 XXZ chain. It belongs to a large family of theories which are known as \mathfrak{g} -quantum KdV models; they exist for any Kac Moody algebra \mathfrak{g} (Feigin and Frenkel 1996) (in the present case $\mathfrak{g} = \widehat{\mathfrak{sl}}_3$), and in the simplest case, namely $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, the Hamiltonian structure of such a theory is the quantisation of the second Poisson structure of the classical KdV equation (Bazhanov et al. 1996).

According to the celebrated ODE/IM correspondence (Dorey and Tateo 1999, 2000; Bazhanov et al. 2001, 2004; Dorey et al. 2007; Feigin and Frenkel 2011;

D. Masoero (✉)

Grupo de Física Matemática da Universidade de Lisboa, Campo Grande, Edifício C6,
1749-016 Lisboa, Portugal
e-mail: dmasoero@gmail.com

Faculdade de Ciências da Universidade de Lisboa, Campo Grande,
1749-016 Lisbon, Portugal

A. Raimondo

Dipartimento di Matematica e Applicazioni, Università di Milano-Bicocca,
Via Roberto Cozzi 55, 20125 Milano, Italy
e-mail: andrea.raimondo@unimib.it

© Springer Nature Switzerland AG 2020

F. Nijhoff et al. (eds.), *Asymptotic, Algebraic and Geometric Aspects of Integrable Systems*, Springer Proceedings in Mathematics & Statistics 338, https://doi.org/10.1007/978-3-030-57000-2_5

Masoero et al. 2016, 2017; Masoero and Raimondo 2020; Kotousov and Lukyanov 2019) to every state of the \mathfrak{g} quantum KdV model there corresponds a unique \mathfrak{g}^L oper (here \mathfrak{g}^L is the Langlands dual of \mathfrak{g}) whose generalised monodromy data provide the solution of the Bethe Ansatz equations of that state.

In our previous paper (Masoero and Raimondo 2020) we constructed theopers corresponding to higher states of the \mathfrak{g} quantum KdV model, for any \mathfrak{g} untwisted affinization of a simply laced Lie algebra. This was done by following the definition given in Feigin and Frenkel (2011); solutions to the Bethe Ansatz were obtained based on our previous works (Masoero et al. 2016, 2017).

In this note we provide explicit and simpler formulas foropers corresponding to higher states of the quantum Boussinesq model, by specialising the results of Masoero and Raimondo (2020) to the case $\mathfrak{g} = \widehat{\mathfrak{sl}}_3$. This serves two purposes: we illustrate the general theory and its somehow heavy machinery in terms of familiar and simple objects, and we find formulas which are much closer to the original work on higher statesopers of the $\widehat{\mathfrak{sl}}_2$ -quantum KdV model (Bazhanov et al. 2004), where higher states are conjectured to correspond to Schrödinger operators with a *monster potential*.

As the result of the present paper, we conjecture that the level N states of the quantum Boussinesq model correspond to the following third order differential operators:

$$L = \partial_z^3 - \left(\sum_{j=1}^N \left(\frac{3}{(z-w_j)^2} + \frac{k}{z(z-w_j)} \right) + \frac{\bar{r}^1}{z^2} \right) \partial_z + \sum_{j=1}^N \left(\frac{3}{(z-w_j)^3} + \frac{a_j}{z(z-w_j)^2} + \frac{2(k+3)a_j - k^2}{3z^2(z-w_j)} \right) + \frac{\bar{r}^2}{z^3} + \frac{1}{z^2} + \lambda z^k, \quad (1)$$

where $-3 < k < -2$, and $\bar{r}^1, \bar{r}^2 \in \mathbb{C}$, and where the $2N$ complex variables $\{a_\ell, w_\ell\}_{\ell=1, \dots, N}$, satisfy the following system of $2N$ algebraic equations

$$a_\ell^2 - ka_\ell + k^2 + 3k - 3\bar{r}^1 = \sum_{\substack{j=1, \dots, N \\ j \neq \ell}} \left(\frac{9w_\ell^2}{(w_\ell - w_j)^2} + \frac{3kw_\ell}{w_\ell - w_j} \right), \quad (2a)$$

$$Aa_\ell + B - 9(k+2)w_\ell = \sum_{\substack{j=1 \\ j \neq \ell}}^N \left(\frac{18(k - a_\ell - a_j)w_\ell^3}{(w_\ell - w_j)^3} + \frac{(12k + 9k^2 - (63 + 6k)a_j - 9ka_\ell)w_\ell^2}{(w_\ell - w_j)^2} + \frac{(9k + 16k^2 + 6(k^2 + 10k + 6)a_j - 5ka_\ell)w_\ell}{w_\ell - w_j} \right). \quad (2b)$$

The parameters A, B are given by

$$A = 14k^2 + 50k - 8\bar{r}^1 + 45,$$

$$B = 27(\bar{r}^1 - \bar{r}^2) - k(7k^2 + 7k + 9\bar{r}^2 - 13\bar{r}^1 + 9),$$

and the additional singularities w_j , $j = 1 \dots, N$ are assumed to be pairwise distinct and nonzero. The system of algebraic equation (2) is equivalent to the requirement that the monodromy around the singular point $z = w_j$ is trivial for all $j = 1 \dots N$, independently on the parameter λ .

The correspondence among the free parameters $\lambda, \bar{r}^1, \bar{r}^2, k$ of the above equations and the free parameters $c, (\Delta_2, \Delta_3), \mu$ (respectively the central charge, the highest weight, the spectral parameter) of the Quantum Boussinesq model, as constructed in Bazhanov et al. (2002) (more about this below), goes as follows:

$$c = -\frac{3(4k+9)(3k+5)}{k+3}, \tag{3a}$$

$$\Delta_2 = \frac{(\bar{r}^1 - 8)k^2 + 6(\bar{r}^1 - 5)k + 9\bar{r}^1 - 27}{9(k+3)}, \tag{3b}$$

$$\Delta_3 = \frac{(k+3)^{3/2}}{27}(\bar{r}^1 - \bar{r}^2), \tag{3c}$$

$$\lambda = -i \Gamma(-k-2)^3 \mu^3, \tag{3d}$$

where $\Gamma(s)$ denotes the Γ function with argument s . Moreover the integer N , which is the number of additional regular singularities in (1), coincides with the level of the state. Hence, system (2) is expected to possess $p_2(N)$ solutions, where $p_2(N)$ is the number of bi-coloured partitions of N .

The paper is organised as follows. In Sect. 2 we introduce the quantum KdV opers, following (Masoero and Raimondo 2020) (which in turns builds on Feigin and Frenkel 2011), and derive from the general theory of the formulas (1) and (2). In Sect. 3 we review the construction of solutions of the Bethe Ansatz equations as generalised monodromy data, following Masoero et al. (2016), Masoero and Raimondo (2020). Finally, in Sect. 4 we briefly summarise the construction of the quantum Boussinesq model provided in Bazhanov et al. (2002).

This work deals with differential equations and representation theory. We omit many proofs of the analytic results, which can be found in greater generality in Masoero and Raimondo (2020). However, we do provide all details of the algebraic calculations.

2 Quantum KdV Opers

In this section we introduce the Quantum KdV opers, as defined in Feigin and Frenkel (2011), in the special case $\mathfrak{g} = \widehat{\mathfrak{sl}}_3$, and derive the third order scalar differential oper-

ator (1). The reader should refer to Masoero and Raimondo (2020), and references therein for more details.

We begin by introducing some theory of the algebra $\mathfrak{sl}_3(\mathbb{C})$,¹ which we realise as the Lie algebra of traceless 3 by 3 matrices (in such a way that it coincides with its first fundamental representation, also known as standard representation). The algebra has the decomposition $\mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$, where \mathfrak{n}_- are lower diagonal matrices, \mathfrak{h} is the Cartan subalgebra of traceless diagonal matrices, and \mathfrak{n}_+ are upper diagonal matrices. The subalgebra $\mathfrak{b}_+ := \mathfrak{h} \oplus \mathfrak{n}_+$ is called the Borel subalgebra. We provide an explicit basis of \mathfrak{b}_+ as follows

$$h_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad h_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad (4)$$

$$e_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad e_\theta = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (5)$$

We introduce three further elements, the sum of the negative Chevalley generators of the Lie algebra $f \in \mathfrak{n}_-$ (principal nilpotent element), the dual of the Weyl vector $\rho^\vee \in \mathfrak{h}$, and the dual of the highest root $\theta^\vee \in \mathfrak{h}$. We have:

$$f = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad \rho^\vee = \theta^\vee = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \quad (6)$$

The unipotent group $\mathcal{N} = \{\exp y, y \in \mathfrak{n}_+\}$ acts on \mathfrak{sl}_3 via the formula

$$\exp y . g = g + \sum_{k \geq 1} \frac{(\text{ad}_y)^k . g}{k!}, \quad \text{ad}_y . g := [y, g],$$

and the affine subspace $f + \mathfrak{b}_+$ is preserved by the action. Following Kostant (1978), and given a vector subspace $\mathfrak{s} \subset \mathfrak{n}_+$, we say that the affine subspace $f + \mathfrak{s}$ is a *transversal space* if

1. The orbit of $f + \mathfrak{s}$ under the action of \mathcal{N} coincides with $f + \mathfrak{b}_+$.
2. For each $s \in \mathfrak{s}$, then $\exp y . (f + s) \notin f + \mathfrak{s}$ unless $y = 0$.

The subspace $\mathfrak{s} = \mathbb{C}e_1 \oplus \mathbb{C}e_\theta$ satisfies the above hypotheses² and the transversal space $f + \mathfrak{s}$ is the space of companion matrices:

¹For sake of simplicity we prefer to work with \mathfrak{sl}_3 -opers, instead of $\widehat{\mathfrak{sl}}_3$ -opers. We do that by considering the loop algebra variable λ as a free complex parameter. More about this in Masoero and Raimondo (2020), Sect. 4.

²As an example, the Cartan subalgebra \mathfrak{h} satisfies the first but not the second hypothesis above.

$$f + \mathfrak{s} = \left\{ \begin{pmatrix} 0 & a & b \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \mid a, b \in \mathbb{C} \right\}.$$

We fix this choice for the rest of the paper.

2.1 Opers

We denote by K the field of rational functions in the variable z , and we define

1. $\mathfrak{g}(K)$, $\mathfrak{b}_+(K)$, $\mathfrak{n}_+(K)$ the Lie algebras of rational functions with values in \mathfrak{g} , \mathfrak{b}_+ , \mathfrak{n}_+ respectively.
2. The space of (global meromorphic) \mathfrak{g} -valued connections $\text{conn}(K) = \{\partial_z + g, g \in \mathfrak{g}(K)\}$.
3. The subset $\text{op}(K) = \{\mathcal{L} = \partial_z + f + b, b \in \mathfrak{b}_+(K)\} \subset \text{conn}(K)$.
4. The group of unipotent Gauge transformations $\mathcal{N}(K) = \{\exp y, y \in \mathfrak{n}_+(K)\}$, acting on $\text{conn}(K)$ via the formula

$$\exp y \cdot (\partial_z + g) = \partial_z - \sum_{k \geq 0} \frac{1}{(k+1)!} (\text{ad}_y)^k \frac{dy}{dz} + \exp y \cdot g. \quad (7)$$

Note that the above action preserves the subset $\text{op}(K)$.

5. The space of \mathfrak{sl}_3 opers as $\text{Op}(K) = \text{op}(K)/\mathcal{N}(K)$.

The space of opers $\text{Op}(K)$ admits a very explicit description once a transversal space $f + \mathfrak{s}$ is fixed: any element in $\text{op}(K)$ is Gauge equivalent to a unique connection of the form $\partial_z + f + s$, $s \in \mathfrak{s}(K)$. Hence we have a bijection

$$\text{Op}(K) \cong \{\partial_z + f + s, s \in \mathfrak{s}(K)\}.$$

We call $\partial_z + f + s$ the *canonical form* of any oper Gauge equivalent to it.

2.2 Opers and Scalar ODEs

It is a standard and elementary result that the space of \mathfrak{sl}_3 opers coincides with the space of third order linear scalar differential operators (with principal symbol equal to 1 and vanishing sub-principal symbol). Indeed, for what we have said so far, any oper has a unique representative of the form

$$\mathcal{L} = \partial_z + f + v_1(z)e_1 + v_2(z)e_\theta,$$

where v_1, v_2 are a pair of (arbitrary) rational functions. In the first fundamental representation, this oper takes the form

$$\mathcal{L} = \partial_z + \begin{pmatrix} 0 & v_1(z) & v_2(z) \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \quad (8)$$

If $\{\epsilon_1, \epsilon_2, \epsilon_3\}$ is the standard basis of \mathbb{C}^3 , and given $\psi = \mathbb{C} \rightarrow \mathbb{C}^3$, with $\psi(z) = \psi_1(z)\epsilon_1 + \psi_2(z)\epsilon_2 + \psi_3(z)\epsilon_3$, then the matrix first order equation

$$\mathcal{L}\psi(z) = 0,$$

is easily seen to be equivalent to the following scalar ODE for the third coefficient $\Psi := \psi_3$

$$(\partial_z^3 - v_1\partial_z + v_2)\Psi(z) = 0. \quad (9)$$

We will use this scalar representation in the rest of the paper.

2.3 (Ir)Regular Singularities

Let \mathcal{L} be an oper in the canonical form (8), and $w \in \mathbb{C}$ a pole of v_1 or v_2 , so that

$$\begin{aligned} v_1 &= \bar{s}_1(z-w)^{-\delta_1} + o((z-w)^{-\delta_1}), \\ v_2 &= \bar{s}_2(z-w)^{-\delta_2} + o((z-w)^{-\delta_2}) \end{aligned}$$

for some $\bar{s}^1, \bar{s}^2 \neq 0$ and some $\delta_1, \delta_2 \in \mathbb{Z}$. We define (Masoero and Raimondo 2020)

– The *slope* of the singular point $w \in \mathbb{C}$ as

$$\mu = \max \left\{ 1, \max \left\{ \frac{\delta_1}{2}, \frac{\delta_2}{3} \right\} \right\} \in \mathbb{Q}.$$

– The *principal coefficient* of the singular point w as

$$f - \rho^\vee + \bar{s}^1 e_1 + \bar{s}^2 e_2 = \begin{pmatrix} -1 & \bar{s}^1 & \bar{s}^2 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} \quad \text{if } \mu = 1,$$

and

$$f + \bar{s}^1 e_1 + \bar{s}^2 e_2 = \begin{pmatrix} 0 & \bar{s}^1 & \bar{s}^2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \text{if } \mu > 1.$$

As proved in Masoero and Raimondo (2020), the singularity is *regular* (in the sense of linear connections) if $\mu = 1$ and *irregular* if $\mu > 1$.

Remark 1 In the case when $w = \infty$, we write $v_1 = z^{\delta_1} + o(z^{\delta_1})$, and $v_2 = \bar{s}_2 z^{\delta_2} + o(z^{\delta_2})$ for some $\bar{s}^1, \bar{s}^2 \neq 0$, and $\delta_1, \delta_2 \in \mathbb{Z}$, and define the slope of $w = \infty$ as $\mu = \max\{1, \max\{\frac{\delta_1}{2}, \frac{\delta_2}{3}\} + 2\}$. The principal coefficient is defined as above.

2.4 \mathfrak{sl}_3 -Quantum KdV Opers

We define \mathfrak{sl}_3 -quantum KdV opers following Feigin and Frenkel (2011). To this aim we fix $-3 < k < -2$ and $\bar{r}^1, \bar{r}^2 \in \mathbb{C}$ and write

$$\mathcal{L}(z, \lambda) = \mathcal{L}_{G, \mathfrak{s}}(z, \lambda) + s(z), \quad s \in K(\mathfrak{s}). \tag{10}$$

Here $\mathcal{L}_{G, \mathfrak{s}}$ is the ground state oper

$$\mathcal{L}_{G, \mathfrak{s}}(z, \lambda) = \partial_z + \begin{pmatrix} 0 & \bar{r}^1/z^2 & \bar{r}^2/z^3 + z^{-2} + \lambda z^k \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \tag{11}$$

We notice that $\mathcal{L}_{G, \mathfrak{s}}(z, \lambda)$ has two singular points: $z = 0$ is a regular singularity with principal coefficient

$$\begin{pmatrix} -1 & \bar{r}^1 & \bar{r}^2 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix},$$

while $z = \infty$ is an irregular singularity, with slope $\mu = \frac{4}{3}$ and principal coefficient

$$\begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

As it will be reviewed in the next section, one can obtain solutions of the Bethe Ansatz equations by considering the differential equation $\mathcal{L}_{G, \mathfrak{s}}\psi = 0$: more precisely these are obtained as coefficients of the expansion of the subdominant solution at $+\infty$ in terms of a distinguished basis of solutions defined at $z = 0$.

In Bazhanov et al. (2004), Bazhanov, Lukyanov and Zamolodchikov proved that in the case $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, the ground state oper could be modified without altering the above global structure, so that the modified equations yield (different) solutions of the same Bethe Ansatz equations (as coefficients of the same expansion). Feigin and Frenkel (2011) extended these idea to the case of a general Kac-Moody algebra, and conjectured that the higher level opers could be uniquely specified by imposing on the \mathfrak{s} -valued function s the 4 conditions below. These conditions were shown

to sufficient (Masoero and Raimondo 2020), and are expected to be necessary for generic values of the parameters \hat{k} , \bar{r}^1 , \bar{r}^2 (Feigin and Frenkel 2011). We say that the oper $\mathcal{L}(z, \lambda)$ of the form (10) is a \mathfrak{sl}_3 -quantum KdV oper if it satisfies the following 4 assumptions:

Assumption 1 The slope and principal coefficient at 0 do not depend on s .

Assumption 2 The slope and principal coefficient at ∞ do not depend on s .

Assumption 3 All additional singular points are regular and the corresponding principal coefficients are conjugated to the element $f - \rho^\vee - \theta^\vee \in f + \mathfrak{h}$.

Assumption 4 All additional singular points have trivial monodromy for every $\lambda \in \mathbb{C}$.

The following proposition, which is Proposition 4.7 in Masoero and Raimondo (2020) specialised to the case of $\mathfrak{g} = \mathfrak{sl}_3$, is a first characterisation of the Quantum KdV oper; it shows that they have the form (1).

Proposition 1 *An operator $\mathcal{L}(z, \lambda)$ of the form (10) satisfies the first three Assumptions if and only if there exists a (possibly empty) arbitrary finite collection of non-zero mutually distinct complex numbers $\{w_j\}_{j \in J} \subset \mathbb{C}^\times$ and a collection of numbers $\{a_{11}^{(j)}, a_{21}^{(j)}, a_{22}^{(j)}\}_{j \in J} \subset \mathbb{C}$, such that $\mathcal{L}(z, \lambda)$ has the form*

$$\mathcal{L}(z, \lambda) = \partial_z + \begin{pmatrix} 0 & W_1 & W_2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (12)$$

where

$$W_1(z) = \frac{\bar{r}^1}{z^2} + \sum_{j \in J} \left(\frac{3}{(z - w_j)^2} + \frac{a_{11}^{(j)}}{z(z - w_j)} \right), \quad (13a)$$

$$W_2(z, \lambda) = \frac{\bar{r}^2}{z^3} + \frac{1}{z^2} + \lambda z^k + \sum_{j \in J} \left(\frac{3}{(z - w_j)^3} + \frac{a_{21}^{(j)}}{z(z - w_j)^2} + \frac{a_{22}^{(j)}}{z^2(z - w_j)} \right). \quad (13b)$$

Note that when J is empty then (12) reduces to the ground state oper (11). If J is not empty, then we set $J = \{1, \dots, N\}$, for some $N \in \mathbb{Z}_+$. In order to fully characterise the \mathfrak{sl}_3 -quantum KdV oper, we must impose the fourth and last Assumption on the oper of the form (12), namely the triviality of the monodromy about all the additional singularities w_j , $j = 1 \dots N$. We notice that the oper of the form (12) depend on the $4N$ complex parameters $\{a_{11}^{(j)}, a_{21}^{(j)}, a_{22}^{(j)}, w_j\}_{j \in 1 \dots N}$. We will show in the following subsection that the trivial monodromy conditions are equivalent to a complete system of $4N$ algebraic equations, which in turn are equivalent to (1) and (2).

2.5 Trivial Monodromy Conditions

We fix $\ell \in 1 \dots N$ and study under which conditions the oper $\mathcal{L}(z, \lambda)$ of the form (12) has trivial monodromy about w_ℓ . As we showed in Masoero and Raimondo (2020), Assumption 3 (more precisely, the fact that θ^\vee is a co-root) implies that the monodromy about w_ℓ is trivial if and only if it is trivial in at least one irreducible (nontrivial) representation. In other words, it is necessary and sufficient that the monodromy at $z = w_\ell$ is trivial for the solutions of the equation $\mathcal{L}(z, \lambda)\Psi = 0$ in the standard representation.

To this aim we write the above equation in the scalar form

$$(\partial_z^3 - W_1 \partial_z + W_2)\Phi(z) = 0, \quad (14)$$

and use the method of the Frobenius expansion, that is we look for solutions of the form

$$\Phi^{(\beta)}(z) = \sum_{m \geq 0} \Phi_m(z - w_\ell)^{\beta+m}. \quad (15)$$

Writing the Laurent expansion of (14) at w_ℓ as

$$W_1(z) = \sum_{m=0}^{+\infty} q_{1m}^{(\ell)}(z - w_\ell)^{m-2}, \quad q_{10}^{(\ell)} = 3, \quad (16a)$$

$$W_2(z) = \sum_{m=0}^{+\infty} q_{2m}^{(\ell)}(z - w_\ell)^{m-3}, \quad q_{20}^{(\ell)} = 3, \quad (16b)$$

expanding the Eq. (15) in powers of $z - w_\ell$, and equating to zero term-by-term we obtain

$$\begin{aligned} \Phi_0^{(\beta)} P(\beta) &= 0, \\ P(\beta + r) \Phi_r^{(\beta)} &= \sum_{m=1}^r \left((\beta + r - m) q_{1m}^{(\ell)} - q_{2m}^{(\ell)} \right) \Phi_{r-m}^{(\beta)}, \end{aligned} \quad (17)$$

where the indicial polynomial $P(\beta) = (\beta - 3)(\beta - 1)(\beta + 1)$. The roots of the indicial polynomial, $\beta = -1, 1, 3$, are known as *indices*. Since the indices are integers, the monodromy matrix has a unique eigenvalue, 1, with algebraic multiplicity 3, and the monodromy is trivial if and only if the recursion (17) has a solution for all the indices. Indeed, in such a case, $\Phi^\beta(e^{2\pi i} z) = \Phi^\beta(z)$ for $\beta = -1, 1, 3$; otherwise logarithmic terms must be added to the series (15) and the monodromy is not diagonalizable (Wasow 2018).

We analyse the recursion (17) separately for the three indices.

The recursion (17) for the index $\beta = 3$ admits always a unique solution, since $P(3 + r) \neq 0, \forall r \geq 1$.

In the case $\beta = 1$, we have that $P(\beta + r) = 0$, $r \geq 1$ if and only if $r = 2$. Hence the recursion is over-determined. Computing the first two terms we obtain

$$\begin{aligned} -3\Phi_1^{(1)} &= (q_{11}^{(\ell)} - q_{21}^{(\ell)}) \Phi_0^{(1)}, \\ 0 \times \Phi_2^{(1)} &= (2q_{11}^{(\ell)} - q_{21}^{(\ell)}) \Phi_1^{(1)} + (q_{12}^{(\ell)} - q_{22}^{(\ell)}) \Phi_0^{(1)}. \end{aligned}$$

It follows that the recursion for the index $\beta = 1$ has at least one solution if and only if

$$q_{12}^{(\ell)} - q_{22}^{(\ell)} = \frac{2}{3} (q_{11}^{(\ell)})^2 - q_{11}^{(\ell)} q_{21}^{(\ell)} + \frac{1}{3} (q_{21}^{(\ell)})^2. \quad (18)$$

Finally, the Frobenius method for the index $\beta_2 = -1$ gives

$$\begin{aligned} 3\Phi_1^{(-1)} &= -(q_{11}^{(\ell)} + q_{21}^{(\ell)}) \Phi_0^{(-1)}, \\ 0 \times \Phi_2^{(-1)} &= -q_{21}^{(\ell)} \Phi_1^{(-1)} - (q_{12}^{(\ell)} + q_{22}^{(\ell)}) \Phi_0^{(-1)}, \\ -3\Phi_3^{(-1)} &= (q_{11}^{(\ell)} - q_{21}^{(\ell)}) \Phi_2^{(-1)} - q_{22}^{(\ell)} \Phi_1^{(-1)} - (q_{13}^{(\ell)} + q_{23}^{(\ell)}) \Phi_0^{(-1)}, \\ 0 \times \Phi_4^{(-1)} &= (2q_{11}^{(\ell)} - q_{21}^{(\ell)}) \Phi_3^{(-1)} + (q_{12}^{(\ell)} - q_{22}^{(\ell)}) \Phi_2^{(-1)} - q_{23}^{(\ell)} \Phi_1^{(-1)} - (q_{14}^{(\ell)} + q_{24}^{(\ell)}) \Phi_0^{(-1)}, \end{aligned}$$

and we obtain the following constraints

$$\begin{aligned} q_{12}^{(\ell)} + q_{22}^{(\ell)} &= \frac{1}{3} q_{21}^{(\ell)} q_{11}^{(\ell)} + \frac{1}{3} (q_{21}^{(\ell)})^2, \\ q_{14}^{(\ell)} + q_{24}^{(\ell)} &= (2q_{11}^{(\ell)} - q_{21}^{(\ell)}) \left(-\frac{1}{9} q_{22}^{(\ell)} (q_{11}^{(\ell)} + q_{21}^{(\ell)}) + \frac{1}{3} (q_{13}^{(\ell)} + q_{23}^{(\ell)}) \right) \\ &\quad + \frac{1}{3} q_{23}^{(\ell)} (q_{11}^{(\ell)} + q_{21}^{(\ell)}). \end{aligned}$$

Combining these with (18) we obtain the following characterisation: the monodromy about w_ℓ is trivial if and only if the following system of 3 equations

$$q_{12}^{(\ell)} = \frac{1}{3} \left((q_{11}^{(\ell)})^2 - q_{11}^{(\ell)} q_{21}^{(\ell)} + (q_{21}^{(\ell)})^2 \right), \quad (19)$$

$$q_{22}^{(\ell)} = \frac{1}{3} q_{11}^{(\ell)} (2q_{21}^{(\ell)} - q_{11}^{(\ell)}), \quad (20)$$

$$\begin{aligned} q_{14}^{(\ell)} + q_{24}^{(\ell)} &= \frac{1}{3} q_{13}^{(\ell)} (2q_{11}^{(\ell)} - q_{21}^{(\ell)}) + q_{11}^{(\ell)} q_{23}^{(\ell)} \\ &\quad + \frac{1}{27} q_{11}^{(\ell)} (2q_{11}^{(\ell)} - q_{21}^{(\ell)}) (q_{11}^{(\ell)} - 2q_{21}^{(\ell)}) (q_{11}^{(\ell)} + q_{21}^{(\ell)}). \end{aligned} \quad (21)$$

In order to proceed further we write explicitly the coefficients q 's, which appear in the above equations, in terms of the parameters of the opers (12)

$$\begin{aligned}
 q_{10}^\ell &= 3, & q_{20}^\ell &= 3, & q_{11}^{(\ell)} &= \frac{a_{11}^{(\ell)}}{w_\ell}, & q_{21}^{(\ell)} &= \frac{a_{21}^{(\ell)}}{w_\ell}, \\
 q_{12}^{(\ell)} &= \frac{\bar{r}^1 - a_{11}^{(\ell)}}{w_\ell^2} + \sum_{\substack{j=1 \\ j \neq \ell}}^N \left(\frac{3}{(w_\ell - w_j)^2} + \frac{a_{11}^{(j)}}{w_\ell(w_\ell - w_j)} \right), & q_{22}^{(\ell)} &= \frac{a_{22}^{(\ell)} - a_{21}^{(\ell)}}{w_\ell^2}, \\
 q_{13}^{(\ell)} &= \frac{a_{11}^{(\ell)} - 2\bar{r}^1}{w_\ell^3} - \sum_{\substack{j=1 \\ j \neq \ell}}^N \left(\frac{6}{(w_\ell - w_j)^3} + \frac{a_{11}^{(j)}}{w_\ell(w_\ell - w_j)^2} + \frac{a_{11}^{(j)}}{w_\ell^2(w_\ell - w_j)} \right), \\
 q_{23}^{(\ell)} &= \frac{\bar{r}^2 + a_{21}^{(\ell)} - 2a_{22}^{(\ell)} + w_\ell}{w_\ell^3} + \lambda w_\ell^k + \sum_{\substack{j=1 \\ j \neq \ell}}^N \left(\frac{3}{(w_\ell - w_j)^3} + \frac{a_{21}^{(j)}}{w_\ell(w_\ell - w_j)^2} + \frac{a_{22}^{(j)}}{w_\ell^2(w_\ell - w_j)} \right), \\
 q_{14}^{(\ell)} &= \frac{3\bar{r}^1 - a_{11}^{(\ell)}}{w_\ell^4} + \sum_{\substack{j=1 \\ j \neq \ell}}^N \left(\frac{9}{(w_\ell - w_j)^4} + \frac{a_{11}^{(j)}}{w_\ell(w_\ell - w_j)^3} + \frac{a_{11}^{(j)}}{w_\ell^2(w_\ell - w_j)^2} + \frac{a_{11}^{(j)}}{w_\ell^3(w_\ell - w_j)} \right), \\
 q_{24}^{(\ell)} &= \frac{3a_{22}^{(\ell)} - a_{21}^{(\ell)} - 3\bar{r}^2 - 2w_\ell}{w_\ell^4} + \lambda k w_\ell^{k-1} \\
 &\quad - \sum_{\substack{j=1 \\ j \neq \ell}}^N \left(\frac{9}{(w_\ell - w_j)^4} + \frac{2a_{21}^{(j)}}{w_\ell(w_\ell - w_j)^3} + \frac{a_{21}^{(j)} + a_{22}^{(j)}}{w_\ell^2(w_\ell - w_j)^2} + \frac{2a_{22}^{(j)}}{w_\ell^3(w_\ell - w_j)} \right).
 \end{aligned}$$

We notice that while Eqs. (19) and (20) do not depend on λ , Eq. (21) is a first-order polynomial in λ . Since the trivial monodromy conditions must hold for any λ , Eq. (21) consists of a pair of independent constraints: both the constant part in λ and the linear part in λ are required to vanish independently. The vanishing of the part of (21) which is linear in λ reads:

$$q_{11}^{(\ell)} w_\ell^k - k w_\ell^{k-1} = 0, \quad \text{or} \quad q_{11}^{(\ell)} = \frac{k}{w_\ell},$$

from which we obtain

$$a_{11}^{(\ell)} = k, \quad \ell = 1, \dots, N. \quad (22)$$

Making use of the explicit expression of the q 's in terms of the a 's, as given above, and denoting

$$a_\ell = a_{21}^{(\ell)}, \quad \ell = 1, \dots, N, \quad (23)$$

from (20) we obtain

$$a_{22}^{(\ell)} = \frac{2}{3}(k+3)a_\ell - \frac{k^2}{3}, \quad \ell = 1, \dots, N. \quad (24)$$

Substituting (22) and (24) into the expression for the q 's found above, then from (19) we obtain (2a), while the vanishing of the constant (in λ) coefficient of (21) is equivalent to (2b).

We have thus arrived to the following result: an \mathfrak{sl}_3 Quantum KdV oper is equivalent to a scalar third order differential operator of the form (1) such that its coefficients satisfy the system of algebraic equations (2).

2.6 The Dual Representation. Formal Adjoint Operator

Before we proceed further with our analysis, and we construct solutions to the Bethe Ansatz equations, we introduce a second representation of the algebra \mathfrak{sl}_3 . This is called the second fundamental representation or dual representation, and we denote it by \mathbb{C}^{3*} . If $\{\epsilon_1, \epsilon_2, \epsilon_3\}$ is the standard basis of \mathbb{C}^3 as above, we denote by $\{\epsilon_1^*, \epsilon_2^*, \epsilon_3^*\}$, the corresponding dual basis in \mathbb{C}^{3*} such that $\langle \epsilon_i^*, \epsilon_j \rangle = \delta_{ij}$. In these basis, the matrices representing the elements $h_1, h_2, \theta^\vee, \rho^\vee, e_1, e_2, e_\theta, f$ read

$$h_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad h_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \rho^\vee = \theta^\vee = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

$$e_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad e_\theta = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

As we have discussed before, the general \mathfrak{sl}_3 oper can be written the canonical form as the connection $\mathcal{L} = \partial_z + f + v_1 e_1 + v_2 e_\theta$, for an arbitrary pair of rational functions $v_1, v_2 \in K$. In the dual representation, we thus have

$$\mathcal{L} = \partial_z + \begin{pmatrix} 0 & 0 & -v_2 \\ 1 & 0 & v_1 \\ 0 & 1 & 0 \end{pmatrix}.$$

We showed that in the standard representation \mathbb{C}^3 the connection \mathcal{L} is equivalent to the scalar third order operator (9). In the dual representation the same oper is equivalent to a different scalar operator, namely to its formal adjoint. Let $\psi^* =: \mathbb{C} \rightarrow \mathbb{C}^{3*}$, with $\psi^*(z) = \psi_1^*(z)\epsilon_1^* + \psi_2^*(z)\epsilon_2^* + \psi_3^*(z)\epsilon_3^*$, satisfy $\mathcal{L}\psi^* = 0$ in the dual representation. Then $\Psi^*(z) := \psi_3^*(z)$ satisfies the scalar ODE

$$(-\partial_z^3 + v_1 \partial_z + (v_2 + v_1'))\Psi^*(z) = 0, \quad (25)$$

which is the formal adjoint of the Eq. (9).

The following standard isomorphisms (of \mathfrak{sl}_3 -modules) will be needed later to derive the Bethe Ansatz equations: $\bigwedge^2 \mathbb{C}^3 \cong \mathbb{C}^{3*}$ and $\bigwedge^2 \mathbb{C}^{3*} \cong \mathbb{C}^3$. Explicitly,

$$\iota(\epsilon_1 \wedge \epsilon_2) = \epsilon_1^*, \quad \iota(\epsilon_1 \wedge \epsilon_3) = \epsilon_2^*, \quad \iota(\epsilon_2 \wedge \epsilon_3) = \epsilon_3^*, \quad (26)$$

$$\iota(\epsilon_1^* \wedge \epsilon_2^*) = \epsilon_1, \quad \iota(\epsilon_1^* \wedge \epsilon_3^*) = \epsilon_2, \quad \iota(\epsilon_2^* \wedge \epsilon_3^*) = \epsilon_3. \quad (27)$$

The above isomorphisms imply that if $\psi(z), \phi(z)$ are solutions of $\mathcal{L}\psi(z) = 0$, for $\psi : \mathbb{C} \rightarrow \mathbb{C}^3$ in the standard representation then $\iota(\psi(z) \wedge \phi(z))$ is a solution of the dual equation $\mathcal{L}\psi^*(z) = 0$, with $\psi^* : \mathbb{C} \rightarrow \mathbb{C}^{3*}$; and conversely.

In the present paper we prefer to work with solutions of the equations in the scalar form (9) and (25). Recall that the solution of the equations in the scalar form is just the third component of the solution of the vector equation. If $\psi(z) = \psi_1(z)\epsilon_1 + \psi_2(z)\epsilon_2 + \psi_3(z)\epsilon_3$ and $\phi(z) = \phi_1(z)\epsilon_1 + \phi_2(z)\epsilon_2 + \phi_3(z)\epsilon_3$, then a simple calculation shows that

$$\langle \iota(\psi \wedge \phi), \epsilon_3 \rangle = Wr[\psi_3, \phi_3]$$

where $Wr[\cdot, \cdot]$ denotes the usual Wronskian $Wr[f(z), g(z)] = f(z)g'(z) - f'(z)g(z)$. Similarly, for $\psi^*(z) = \psi_1^*(z)\epsilon_1^* + \psi_2^*(z)\epsilon_2^* + \psi_3^*(z)\epsilon_3^*$ and $\phi^*(z) = \phi_1^*(z)\epsilon_1^* + \phi_2^*(z)\epsilon_2^* + \phi_3^*(z)\epsilon_3^*$ we have

$$\langle \epsilon_3^*, \iota(\psi^* \wedge \phi^*) \rangle = Wr[\psi_3^*, \phi_3^*].$$

To prove the above relations, it is sufficient to note that from the matrix first order equations $\mathcal{L}\psi(z) = 0, \mathcal{L}\psi^*(z) = 0$ we obtain the identities $\psi_2(z) = -\psi_3'(z)$ and $\psi_2^* = -\psi_3^{*\prime}(z)$. We have thus shown that the Wronskian of two solutions of (9) satisfies (25), and conversely the Wronskian of two solutions of (25) satisfies (9).

2.7 Relation with Previous Works

The ground state \mathfrak{sl}_3 -quantum KdV oper, given by Eq. (11), was also considered – in the scalar form – by Dorey and Tateo (2000), and by Bazhanov et al. (2002), who wrote the following third order scalar operator

$$\tilde{L}(x, E) = \partial_x^3 + \frac{\tilde{w}_1}{x^2} \partial_x + \frac{\tilde{w}_2}{x^3} + x^{3M} - E, \quad (28)$$

with $\tilde{w}_1 = \tilde{\ell}_1 \tilde{\ell}_2 + \tilde{\ell}_1 \tilde{\ell}_3 + \tilde{\ell}_2 \tilde{\ell}_3 - 2, \tilde{w}_2 = -\tilde{\ell}_1 \tilde{\ell}_2 \tilde{\ell}_3$ and where the $\tilde{\ell}_i$'s are constrained by the equation $\tilde{\ell}_1 + \tilde{\ell}_2 + \tilde{\ell}_3 = 3$. In addition, in our previous paper (Masoero et al. 2016) we considered the ground state oper in the following form

$$\mathcal{L}(x, E) = \partial_x + \begin{pmatrix} \ell_1/x & 0 & x^{3M} - E \\ 1 & (\ell_2 - \ell_1)/x & 0 \\ 0 & 1 & -\ell_2/x \end{pmatrix} \quad (29)$$

for arbitrary $\ell_1, \ell_2 \in \mathbb{C}$ and $M > 0$. We now show that the differential operators (11), (28), and (29) are equivalent under appropriate change of coordinates and Gauge transformations, once the parameters are correctly identified. To show that the differential operators (28) and (29) are equivalent, we write the operator (28) in the oper form

$$\partial_x + \begin{pmatrix} 0 & -\tilde{w}_1/x^2 & \tilde{w}_2/x^3 + x^{3M} - E \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \quad (30)$$

It is then a simple computation to show that (30) and (29) are Gauge equivalent if we set $\tilde{\ell}_1 = -\ell_1 + 2$, $\tilde{\ell}_2 = \ell_1 - \ell_2 + 1$ and $\tilde{\ell}_3 = \ell_2$.

Next we show the equivalence between (29) and (11). As observed in Feigin and Frenkel (2011), after the change of variable

$$z = \varphi(x) = \left(\frac{k+3}{3} \right)^3 x^{\frac{3}{k+3}}, \quad k = -\frac{3M+2}{1+M}, \quad (31)$$

the operator (29) reads

$$\mathcal{L}_G(z, \lambda) = \partial_z + \begin{pmatrix} r_1/z & 0 & z^{-2} + \lambda z^k \\ 1 & (r_2 - r_1)/z & 0 \\ 0 & 1 & -r_2/z \end{pmatrix}, \quad (32)$$

where $\lambda \in \mathbb{C}$ and $r_1, r_2 \in \mathbb{C}$ are defined by the relations

$$E = -\left(\frac{k+3}{3} \right)^{3(k+2)} \lambda, \quad \ell_i = \frac{3}{k+3}(r_i - 1) + 1, \quad i = 1, 2. \quad (33)$$

It is again a simple computation to show that the opers (32) and (11) are Gauge equivalent provided the coefficients $r^1, r^2, \bar{r}^1, \bar{r}^2$ satisfy the following relations

$$\begin{cases} \bar{r}^1 = (r^1)^2 - r^1 r^2 + (r^2)^2 - r^1 - r^2, \\ \bar{r}^2 = r^1 r^2 (r^1 - r^2) + r^2 (2r^2 - r^1 - 2). \end{cases} \quad (34)$$

2.8 Weyl Group Symmetry

The parametrisation (34) of \bar{r}^1, \bar{r}^2 in terms of r^1 and r^2 will be very convenient when discussing the behaviour of solutions of $\mathcal{L}(z, \lambda)\psi = 0$ in a neighbourhood of $z = 0$. The Weyl group of \mathfrak{sl}_3 – which is isomorphic to the group of permutations of three elements, S_3 – is a symmetry of the map (34), once its action on the parameters r^1, r^2 , which is called the *dot action*, is properly defined:

$$\sigma \cdot \begin{pmatrix} r^1 \\ r^2 \end{pmatrix} = \begin{pmatrix} -r^2 + 2 \\ -r^1 + 2 \end{pmatrix}, \quad \tau \cdot \begin{pmatrix} r^1 \\ r^2 \end{pmatrix} = \begin{pmatrix} -r^2 + 2 \\ r^1 - r^2 + 1 \end{pmatrix}. \quad (35)$$

We let the reader verify that σ, τ generate the group S_3 (in particular $\sigma^2 = 1, \tau^3 = 1$) and that the above action is a symmetry of (34). This phenomenon is studied in great detail and generality in Masoero and Raimondo (2020), Sect. 5.

3 The Bethe Ansatz Equations

In this section we construct solutions of the Bethe Ansatz equations as generalised monodromy data of Quantum KdV opers, $\mathcal{L}(z, \lambda)$. As proved in Sect. 2, these are opers of the form

$$\mathcal{L}(z, \lambda) = \partial_z + \begin{pmatrix} 0 & W_1 & W_2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (36)$$

where

$$W_1(z) = \frac{\bar{r}^1}{z^2} + \sum_{j \in J} \left(\frac{3}{(z - w_j)^2} + \frac{k}{z(z - w_j)} \right), \quad (37a)$$

$$W_2(z, \lambda) = \frac{\bar{r}^2}{z^3} + \frac{1}{z^2} + \lambda z^k + \sum_{j \in J} \left(\frac{3}{(z - w_j)^3} + \frac{a_j}{z(z - w_j)^2} + \frac{2(k + 3)a_j - k^2}{3z^2(z - w_j)} \right), \quad (37b)$$

and where $\{a_j, w_j\}_{j=1, \dots, N}$ satisfy the system of equations (2). We follow Masoero et al. (2016), Masoero and Raimondo (2020) closely and the reader should refer to these papers for all missing proofs. Any finite dimensional representation V of \mathfrak{sl}_3 defines the ODE

$$\mathcal{L}(z, \lambda)\psi = 0, \quad \Psi : \mathbb{C} \rightarrow V.$$

Since the monodromy of $\mathcal{L}(z, \lambda)$ about w_j is trivial for any j , then the solutions of the above equation are, for fixed λ , analytic functions on the universal cover of \mathbb{C}^* , minus the lift of the points $w_j, j \in 1 \dots N$. We denote such a domain by $\widehat{\mathbb{C}}$. As it was originally observed by Dorey and Tateo, the appearance of the Bethe Ansatz equations is due to a discrete symmetry which acts on both the variable z and the parameter λ . It is therefore necessary to consider solutions $\psi(z, \lambda)$ as analytic functions of both variables z and λ . More precisely for our purpose $\psi(z, \cdot)$ is assumed to be an entire function of λ . We thus define a solution to be an analytic map $\psi : \widehat{\mathbb{C}} \times \mathbb{C} \rightarrow V$ which satisfies the equation $\mathcal{L}(z, \lambda)\psi(z, \lambda) = 0$ for every (z, λ) .

The space of solutions, which we denote by $V(\lambda)$, is an infinite dimensional vector space which, as we showed in Masoero and Raimondo (2020), is simply isomorphic

to $V \otimes O_\lambda$, where O_λ is the ring of entire functions of the variable λ . This means that an O_λ -basis of the space of solutions has cardinality $\dim V$.

3.1 Twisted Opers

Let $\hat{k} = -k - 2$, so that $0 < \hat{k} < 1$. For any $t \in \mathbb{R}$ we define the twisted operator and twisted solution:

$$\mathcal{L}^t(z, \lambda) := \mathcal{L}(e^{2i\pi t} z, e^{2i\pi t \hat{k}} \lambda), \quad (38)$$

$$\psi_t(z, \lambda) = e^{2i\pi t \rho^\vee} \psi(e^{2\pi i t} z, e^{2\pi i t \hat{k}} \lambda). \quad (39)$$

Taking into account the oper change of variables (Masoero and Raimondo 2020), then from (36) we explicitly have

$$\mathcal{L}^t(z, \lambda) = \partial_z + f + e^{4\pi i t} W_1(e^{2\pi i t} z) e_1 + e^{6\pi i t} W_2(e^{2\pi i t} z, e^{2\pi i t \hat{k}} \lambda) e_\theta,$$

and one easily see that the function $\psi_t(z, \lambda)$ satisfies $\mathcal{L}^t(z, \lambda)\psi_t(z, \lambda) = 0$. A crucial property of the oper (36) is the following Dorey-Tateo discrete symmetry:

$$\mathcal{L}^{t=1}(z, \lambda) = \mathcal{L}(z, \lambda), \quad (40)$$

which leads us to consider the following (O_λ -linear) monodromy operator

$$M : V(\lambda) \rightarrow V(\lambda), \quad M(\psi(z, \lambda)) = e^{2i\pi \rho^\vee} \psi(e^{2\pi i} z, e^{2\pi i \hat{k}} \lambda). \quad (41)$$

In the case \mathfrak{sl}_3 , we just need to consider the equations $\mathcal{L}^t(z, \lambda)\psi = 0$ for the standard representation and its dual. More precisely, the standard representation at 0 twist, and the dual representation at twist $t = \frac{1}{2}$

$$\mathcal{L}(z, \lambda)\psi(z, \lambda) = 0, \quad \psi : \widehat{\mathbb{C}} \times \mathbb{C} \rightarrow \mathbb{C}^3, \quad (42)$$

$$\mathcal{L}^{\frac{1}{2}}(z, \lambda)\psi^*(z, \lambda) = 0, \quad \psi^* : \widehat{\mathbb{C}} \times \mathbb{C} \rightarrow \mathbb{C}^{3*}. \quad (43)$$

By a slight abuse of notation we denote $\mathbb{C}^3(\lambda)$ the space of solutions of the first equation, and by $\mathbb{C}^{3*}(\lambda)$ the space of solutions of the latter equations, as well as the solutions of the same equations in the equivalent scalar form

$$(\partial_z^3 - W_1(z)\partial_z + W_2(z, \lambda))\Psi(z, \lambda) = 0, \quad (44)$$

$$(\partial_z^3 - W_1(-z)\partial_z + W_2(-z, e^{\pi i \hat{k}} \lambda) - W_1'(-z))\Psi^*(z, \lambda) = 0. \quad (45)$$

Since the solution of the equations in the scalar form is the third component of a solution of the equation in the matrix form, and since $\rho^\vee \epsilon_3 = -\epsilon_3$, $\rho^\vee \epsilon_3^* = -\epsilon_3^*$, the twist for solutions of the above scalar ODEs is defined as follows

$$\Psi_t(z, \lambda) = e^{-2i\pi t} \Psi(e^{2\pi i t} z, e^{2\pi i t \hat{k}} \lambda), \quad \Psi_t^*(z, \lambda) = e^{-2i\pi t} \Psi^*(e^{2\pi i t} z, e^{2\pi i t \hat{k}} \lambda).$$

Equation (45) is the adjoint equation to (44) twisted by $t = \frac{1}{2}$; and conversely, Eq. (44) is the adjoint equation to (45) twisted by $t = \frac{1}{2}$. As we recalled in Sect. 2.6, the Wronskian of two solutions of a scalar ODE solves the adjoint equation. It follows that

1. If $\Psi(z, \lambda), \Phi(z, \lambda) \in \mathbb{C}^3(\lambda)$ then

$$Wr[\Psi_{-\frac{1}{2}}(z, \lambda), \Phi_{\frac{1}{2}}(z, \lambda)] \in \mathbb{C}^{3*}(\lambda),$$

2. If $\Psi^*(z, \lambda), \Phi^*(z, \lambda) \in \mathbb{C}^{3*}(\lambda)$, then

$$Wr[\Psi_{-\frac{1}{2}}^*(z, \lambda), \Phi_{\frac{1}{2}}^*(z, \lambda)] \in \mathbb{C}^3(\lambda).$$

3.2 The Eigenbasis of the Monodromy Operator. Expansion at $z = 0$

The point $z = 0$ is a regular singularity for the Eqs. (44) and (45), but it is also a branch point of the potential W_2 , because of the term $\lambda z^{\hat{k}}$. It follows that the standard Frobenius series cannot provide solution of the above equations at $z = 0$. A generalised Frobenius series, introduced in Masoero and Raimondo (2020), does however the job. The latter is defined as

$$\Phi^{(\beta)}(z, \lambda) = z^\beta \sum_{m \geq n \geq 0} c_{m,n} z^m \zeta^n, \quad c_{0,0} = 1, \quad \zeta = \lambda z^{-\hat{k}}, \quad (46)$$

where the indices β are computed as in the standard Frobenius method: if the equation reads

$$\left(\partial_z^3 + \frac{a + o(1)}{z^2} \partial_z + \frac{b + o(1)}{z^3} \right) \Psi(z) = 0,$$

the indices are the roots of the indicial polynomial $P(\beta) = \beta^3 - 3\beta^2 + (2 + a)\beta + b$. The following facts are proved in Masoero and Raimondo (2020), Proposition 5.1. For every finite dimensional representation V of \mathfrak{sl}_3 , and under some genericity assumptions³ on the triple $(\hat{k}, \bar{r}^1, \bar{r}^2)$, we have:

³The genericity assumptions imply that the monodromy operator M is diagonal and no logarithmic terms appear in the generalised Frobenius series.

1. The series (46) converges to a solution $\Phi^{(\beta)}(z, \lambda) \in V(\lambda)$.
2. $M\Phi^{(\beta)}(z, \lambda) = e^{2\pi i\beta}\Phi^{(\beta)}(z, \lambda)$, where M is the monodromy operator defined in (41).
3. The collection of the solutions $\Phi^{(\beta)}(z, \lambda)$ for all indices β forms an O_λ -basis of $V(\lambda)$.

In the cases under our study, namely Eqs. (44) and (45), the indicial polynomials are, respectively, given by

$$\begin{aligned} P(\beta) &= \beta^3 - 3\beta^2 + (2 - \bar{r}^1)\beta + \bar{r}^2, \\ P^*(\beta) &= \beta^3 - 3\beta^2 + (2 - \bar{r}^1) + 2\bar{r}^1 - \bar{r}^2. \end{aligned}$$

Using (34), then we obtain the factorizations

$$\begin{aligned} P(\beta) &= (\beta - r^2)(\beta - 1 + r^2 - r^1)(\beta - 2 + r^1), \\ P^*(\beta) &= (\beta - r^1)(\beta - 1 + r^1 - r^2)(\beta - 2 + r^2), \end{aligned}$$

so that the indices are given by

$$\beta_1 = r^2, \quad \beta_2 = r^1 - r^2 + 1, \quad \beta_3 = -r^1 + 2, \quad (47a)$$

$$\beta_1^* = -r^2 + 2, \quad \beta_2^* = r^2 - r^1 + 1, \quad \beta_3^* = r^1. \quad (47b)$$

We denote by

$$\{\Phi^{(\beta_1)}(z, \lambda), \Phi^{(\beta_2)}(z, \lambda), \Phi^{(\beta_3)}(z, \lambda)\}, \quad (48a)$$

$$\{\Phi^{(\beta_1^*)}(z, \lambda), \Phi^{(\beta_2^*)}(z, \lambda), \Phi^{(\beta_3^*)}(z, \lambda)\}, \quad (48b)$$

the corresponding solutions of (44) and (45) respectively. Recall that the Weyl group acting by the dot action (35) on r^1, r^2 , provides a group of symmetries of \bar{r}^1, \bar{r}^2 , hence it leaves the indicial polynomial invariant, permuting its roots.⁴ The (induced) action of the generators σ, τ of the Weyl group, see (35), on the indices (47) is provided by the following permutations:

$$\sigma(\beta_i) = \beta_{\sigma(i)}, \quad \tau(\beta_i) = \beta_{\tau(i)}, \quad i = 1, 2, 3, \quad (49a)$$

$$\sigma(\beta_i^*) = \beta_{\sigma(i)}^*, \quad \tau(\beta_i^*) = \beta_{\tau(i)}^*, \quad i = 1, 2, 3, \quad (49b)$$

where

$$\sigma(1, 2, 3) := (3, 2, 1), \quad \tau(1, 2, 3) := (2, 3, 1). \quad (50)$$

Comparing the asymptotic behaviour at $z = 0$, we deduce the following 6 quadratic identities among the (properly normalised) $\Phi^{(\beta)}$'s and $\Phi^{(\beta^*)}$'s. Let $s \in S_3$, then (we

⁴Many authors fix r^1, r^2 by imposing the conditions $\Re\beta_1 > \Re\beta_2 > \Re\beta_3$, or equivalently $\Re\beta_3^* > \Re\beta_2^* > \Re\beta_1^*$ (Bazhanov et al. 2002; Dorey et al. 2007; Masoero et al. 2016).

can find a normalisation of the solutions $\Phi^{(\beta_i)}$, $\Phi^{(\beta_i^*)}$ such that):

$$Wr[\Phi_{\mp\frac{1}{2}}^{(\beta_{s(1)})}, \Phi_{\pm\frac{1}{2}}^{(\beta_{s(2)})}] = (-1)^{p(s)} e^{\pm i\pi s(\gamma)} \Phi^{(\beta_{s(3)})}, \quad (51a)$$

$$Wr[\Phi_{\mp\frac{1}{2}}^{(\beta_{s(3)})}, \Phi_{\pm\frac{1}{2}}^{(\beta_{s(2)})}] = (-1)^{p(s)} e^{\pm i\pi s(\gamma^*)} \Phi^{(\beta_{s(1)})}, \quad (51b)$$

where $p(s)$ is the parity of $s \in S_3$, and where

$$s(\gamma) = \beta_{s(2)} - \beta_{s(1)}, \quad s(\gamma^*) = \beta_{s(2)}^* - \beta_{s(3)}^*, \quad (52)$$

with the $\beta_{s(i)}$ and $\beta_{s(i)}^*$ defined by the relations (47) and (49).

3.3 Sibuya Solutions. Expansion at $z = \infty$

We let $q(z, \lambda)$ be the Puiseux series of $(z^{-2}(1 + \lambda z^{-\hat{k}}))^{\frac{1}{3}}$ truncated after terms of z^{-1} , and $S(z, \lambda)$ be its primitive

$$q(z, \lambda) = z^{-\frac{2}{3}} \left(1 + \sum_{l=0}^{\lfloor \frac{1}{3\hat{k}} \rfloor} c_l \lambda^l z^{-l\hat{k}} \right), \quad S(z, \lambda) = \int^z q(y, \lambda) dy, \quad (53)$$

where c_l are the coefficients of Taylor series expansion at $y = 0$ of $(1 - y)^{\frac{1}{3}}$, and $\int^z y^l dy = \frac{z^{l+1}}{l+1}$, $l \neq -1$, $\int^z \frac{1}{y} = \log z$.

The Sibuya, or subdominant, solution of the Eqs. (44) and (45) is uniquely defined by the following asymptotics

$$\Psi(z, \lambda) = z^{\frac{2}{3}} e^{-S(z, \lambda)} (1 + o(1)), \quad \text{as } z \rightarrow +\infty, \quad (54a)$$

$$\Psi^*(z, \lambda) = z^{\frac{2}{3}} e^{-S(z, \lambda)} (1 + o(1)), \quad \text{as } z \rightarrow +\infty. \quad (54b)$$

Moreover we have that

$$\Psi'(z, \lambda) = -e^{-S(z, \lambda)} (1 + o(1)), \quad \text{as } z \rightarrow +\infty, \quad (55a)$$

$$\Psi^{*'}(z, \lambda) = -e^{-S(z, \lambda)} (1 + o(1)), \quad \text{as } z \rightarrow +\infty. \quad (55b)$$

The Sibuya solutions Ψ, Ψ^* satisfy the following properties

- It is the solution (unique up to a multiplicative constant) with the fastest decrease as $z \rightarrow +\infty$.
- The asymptotic formulas (54) hold true on the sector $|\arg z| \leq \pi + \varepsilon$, for some $\varepsilon > 0$ (Masoero et al. 2016). In other words, if we continue analytically $\Psi(z, \lambda), \Psi^*$

(z, λ) as well as the functions $q(z, \lambda)$ and $S(z, \lambda)$ past the negative real semi-axis, the asymptotic formulas still hold.

- The solutions $\Psi(z, \lambda)$, $\Psi^*(z, \lambda)$ are entire functions of λ , i.e. $\Psi(z, \lambda) \in \mathbb{C}^3(\lambda)$ and $\Psi^*(z, \lambda) \in \mathbb{C}^{3*}(\lambda)$.
- Finally, and most importantly, the solutions $\Psi(z, \lambda)$, $\Psi^*(z, \lambda)$ satisfy the so-called Ψ -system

$$Wr[\Psi_{-\frac{1}{2}}(z, \lambda), \Psi_{\frac{1}{2}}(z, \lambda)] = \Psi^*(z, \lambda), \quad (56a)$$

$$Wr[\Psi_{-\frac{1}{2}}^*(z, \lambda), \Psi_{\frac{1}{2}}^*(z, \lambda)] = \Psi(z, \lambda). \quad (56b)$$

The latter identities can be checked by comparing the asymptotic expansion of the left and right hand side as $z \rightarrow +\infty$.

The Ψ -system is the last necessary ingredient to construct solutions of the Bethe Ansatz equations.

3.4 $\tilde{Q}\tilde{Q}$ System and the Bethe Ansatz

As we have shown, the solutions $\{\Phi^{(\beta_1)}(z, \lambda), \Phi^{(\beta_2)}(z, \lambda), \Phi^{(\beta_3)}(z, \lambda)\}$ obtained in (48a) provide an O_λ basis of $\mathbb{C}^3(\lambda)$, and the Sibuya solution $\Psi(z, \lambda)$ belongs to the same space. It follows that there exists a unique triplet of entire functions $Q_i(\lambda) \in O_\lambda$, for $i = 1, 2, 3$, such that

$$\Psi(z, \lambda) = \tilde{Q}_1(\lambda)\Phi^{(\beta_1)}(z, \lambda) + \tilde{Q}_2(\lambda)\Phi^{(\beta_2)}(z, \lambda) + \tilde{Q}_3(\lambda)\Phi^{(\beta_3)}(z, \lambda). \quad (57a)$$

Similarly, we have that

$$\Psi^*(z, \lambda) = Q_1^*(\lambda)\Phi^{(\beta_1^*)}(z, \lambda) + Q_2^*(\lambda)\Phi^{(\beta_2^*)}(z, \lambda) + Q_3^*(\lambda)\Phi^{(\beta_3^*)}(z, \lambda), \quad (57b)$$

for a unique triplet of entire functions $Q_i^*(\lambda) \in O_\lambda$, with $i = 1, 2, 3$. Substituting the expansions (57) in the Ψ -system (56) and making use of the relations (51) we obtain the following quadratic relations among the coefficients Q 's and Q^* 's, which is known as $\tilde{Q}\tilde{Q}$ -system. For each $s \in S_3$ we have

$$\begin{aligned} (-1)^{p(s)} Q_{s(3)}^*(\lambda) &= e^{i\pi s(\gamma)} Q_{s(1)}(e^{-i\pi\hat{k}}\lambda) Q_{s(2)}(e^{i\pi\hat{k}}\lambda) \\ &\quad - e^{-i\pi s(\gamma)} Q_{s(1)}(e^{i\pi\hat{k}}\lambda) Q_{s(2)}(e^{-i\pi\hat{k}}\lambda), \end{aligned} \quad (58a)$$

$$\begin{aligned} (-1)^{p(s)} Q_{s(3)}(\lambda) &= e^{i\pi s(\gamma^*)} Q_{s(1)}^*(e^{-i\pi\hat{k}}\lambda) Q_{s(2)}^*(e^{i\pi\hat{k}}\lambda) \\ &\quad - e^{-i\pi s(\gamma^*)} Q_{s(1)}^*(e^{i\pi\hat{k}}\lambda) Q_{s(2)}^*(e^{-i\pi\hat{k}}\lambda), \end{aligned} \quad (58b)$$

where $p(s)$ is the parity of s , and the phases $s(\gamma)$, $s(\gamma^*)$ are defined in (52).

Remark 2 System (58) was shown by Frenkel and Hernandez (2018) to be a universal system of relations in the commutative Grothendieck ring $K_0(\mathcal{O})$ of the category \mathcal{O} of representations of the Borel subalgebra of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_3)$.

Finally, the Bethe Ansatz equations is a pair of functional relations for each one of the six pairs of Q functions of the form $\{Q_{s(1)}Q_{s(3)}^*\}$, $s \in S_3$. Let λ_s denote an arbitrary zero of the function $Q_{s(1)}$, and λ_s^* an arbitrary zero of $Q_{s(3)}^*$. Evaluating the above relations at $e^{\pm i\pi\hat{k}}\lambda_s$ we obtain the Bethe Ansatz equations

$$\begin{aligned} -e^{2i\pi s(\gamma)} \frac{Q_{s(1)}(e^{2i\pi\hat{k}}\lambda_s)}{Q_{s(1)}(e^{-2i\pi\hat{k}}\lambda_s)} &= \frac{Q_{s(3)}^*(e^{i\pi\hat{k}}\lambda_s)}{Q_{s(3)}^*(e^{i\pi\hat{k}}\lambda_s)}, \\ -e^{2i\pi s(\gamma^*)} \frac{Q_{s(3)}^*(e^{2i\pi\hat{k}}\lambda_s^*)}{Q_{s(3)}^*(e^{-2i\pi\hat{k}}\lambda_s^*)} &= \frac{Q_{s(1)}(e^{i\pi\hat{k}}\lambda_s^*)}{Q_{s(1)}(e^{-i\pi\hat{k}}\lambda_s^*)}. \end{aligned}$$

It is believed that each one of the 6 Bethe Ansatz equations is strong enough to characterise all of the Q 's and Q^* 's, by means of the so-called Destri-De Vega equations.

4 Quantum Boussinesq Model

The quantum Boussinesq model has been described in great detail by Bazhanov et al. (2002), from which the notation of the present section is taken and to which we refer for further details. The model is defined by considering a highest weight representation $\mathcal{V}_{\Delta_2, \Delta_3}$ of the Zamolodchikov's \mathcal{W}_3 -algebra (Zamolodchikov 1985), and it is characterized by 4 parameters: the central charge $-\infty < c < 2$, the highest weight $(\Delta_2, \Delta_3) \in \mathbb{C}^2$, and the spectral parameter $\mu \in \mathbb{C}$. For generic values of c, Δ_2, Δ_3 , the representation $\mathcal{V}_{\Delta_2, \Delta_3}$ is irreducible, a condition we assume from now on. Let $\mathbf{L}_n, \mathbf{W}_n, n \in \mathbb{Z}$, denote the generators of the \mathcal{W}_3 algebra as in Bazhanov et al. (2002), Sect. 2. The highest weight fixes a ground state $|\Delta_2, \Delta_3\rangle \in \mathcal{V}_{\Delta_2, \Delta_3}$, satisfying $\mathbf{L}_n|\Delta_2, \Delta_3\rangle = \mathbf{W}_n|\Delta_2, \Delta_3\rangle = 0$ for $n > 0$, and

$$\mathbf{L}_0|\Delta_2, \Delta_3\rangle = \Delta_2|\Delta_2, \Delta_3\rangle \quad \mathbf{W}_0|\Delta_2, \Delta_3\rangle = \Delta_3|\Delta_2, \Delta_3\rangle.$$

The \mathcal{W}_3 -module $\mathcal{V}_{\Delta_2, \Delta_3}$ admits the level decomposition

$$\mathcal{V}_{\Delta_2, \Delta_3} = \bigoplus_{N=0}^{\infty} \mathcal{V}_{\Delta_2, \Delta_3}^{(N)}, \quad \mathbf{L}_0\mathcal{V}_{\Delta_2, \Delta_3}^{(N)} = (\Delta_2 + N)\mathcal{V}_{\Delta_2, \Delta_3}^{(N)}.$$

The ground state $|\Delta_2, \Delta_3\rangle$ has level zero, the higher states are obtained by the action of products of the lowering operators $\mathbf{L}_n, \mathbf{W}_n, n < 0$. More precisely, let $\{\nu_1 \dots, \nu_l, \bar{\nu}_1 \dots, \bar{\nu}_k\}$, with $\nu_j, \bar{\nu}_k \in \mathbb{N}$, be a bicoloured integer partition of the integer

N , namely $\nu_j \leq \nu_{j+1}$, $\bar{\nu}_j \leq \bar{\nu}_{j+1}$ and $\sum_j \nu_j + \sum_j \bar{\nu}_j = N$; to any such a partition one associates a state of level N by the formula $\prod_j \mathbf{L}_{-\nu_j} \prod_j \mathbf{W}_{-\bar{\nu}_j} |0\rangle$.

The integrable structure of the quantum Boussinesq model can be conveniently encoded in the so-called \mathbf{Q} -operators (Bazhanov et al. 2002, Sect. 2), from which the quantum integrals of motion of the model can be obtained. The \mathbf{Q} -operators are more precisely operator-valued functions $\mathbf{Q}_i(t)$, $\bar{\mathbf{Q}}_i(t)$, $i = 1, 2, 3$, depending on the parameter $t = \mu^3$, where μ is the spectral parameter of the quantum model.⁵ The level subspaces $\mathcal{V}_{\Delta_2, \Delta_3}^{(N)}$ are invariant with respect to the action of the \mathbf{Q} -operators,

$$\mathbf{Q}_i(t) : \mathcal{V}_{\Delta_2, \Delta_3}^{(N)} \rightarrow \mathcal{V}_{\Delta_2, \Delta_3}^{(N)}, \quad \bar{\mathbf{Q}}_i(t) : \mathcal{V}_{\Delta_2, \Delta_3}^{(N)} \rightarrow \mathcal{V}_{\Delta_2, \Delta_3}^{(N)},$$

and in particular (for $N = 0$), the ground state $|\Delta_2, \Delta_3\rangle$ is an eigenvector for the \mathbf{Q} -operators:

$$\begin{aligned} \mathbf{Q}_i(t)|\Delta_2, \Delta_3\rangle &= P_i^{(vac)}(t)|\Delta_2, \Delta_3\rangle, \\ \bar{\mathbf{Q}}_i(t)|\Delta_2, \Delta_3\rangle &= \bar{P}_i^{(vac)}(t)|\Delta_2, \Delta_3\rangle. \end{aligned}$$

As proved in Bazhanov et al. (2002), Sect. 5, the \mathbf{Q} -operators (and therefore their eigenvalues) satisfy the system of quadratic relations

$$c_1 \bar{\mathbf{Q}}_1(t) = \mathbf{Q}_2(qt) \mathbf{Q}_3(q^{-1}t) - \mathbf{Q}_3(qt) \mathbf{Q}_2(q^{-1}t), \quad (59a)$$

$$c_1 \mathbf{Q}_1(t) = \bar{\mathbf{Q}}_3(qt) \bar{\mathbf{Q}}_2(q^{-1}t) - \bar{\mathbf{Q}}_2(qt) \bar{\mathbf{Q}}_3(q^{-1}t), \quad (59b)$$

$$c_2 \bar{\mathbf{Q}}_2(t) = \mathbf{Q}_3(qt) \mathbf{Q}_1(q^{-1}t) - \mathbf{Q}_1(qt) \mathbf{Q}_3(q^{-1}t), \quad (59c)$$

$$c_2 \mathbf{Q}_2(t) = \bar{\mathbf{Q}}_1(qt) \bar{\mathbf{Q}}_3(q^{-1}t) - \bar{\mathbf{Q}}_3(qt) \bar{\mathbf{Q}}_1(q^{-1}t), \quad (59d)$$

$$c_3 \bar{\mathbf{Q}}_3(t) = \mathbf{Q}_1(qt) \mathbf{Q}_2(q^{-1}t) - \mathbf{Q}_2(qt) \mathbf{Q}_1(q^{-1}t), \quad (59e)$$

$$c_3 \mathbf{Q}_3(t) = \bar{\mathbf{Q}}_2(qt) \bar{\mathbf{Q}}_1(q^{-1}t) - \bar{\mathbf{Q}}_1(qt) \bar{\mathbf{Q}}_2(q^{-1}t), \quad (59f)$$

where

$$q = e^{i\pi g}, \quad (60a)$$

$$c_1 = e^{i\pi(p_1 - \sqrt{3}p_2)} - e^{-i\pi(p_1 - \sqrt{3}p_2)}, \quad (60b)$$

$$c_2 = e^{-2i\pi p_1} - e^{2i\pi p_1}, \quad (60c)$$

$$c_3 = e^{i\pi(p_1 + \sqrt{3}p_2)} - e^{-i\pi(p_1 + \sqrt{3}p_2)}, \quad (60d)$$

and the parameter g , p_1 , p_2 are related to c , Δ_2 , Δ_3 by the identities (Bazhanov et al. 2002, Sect. 3)

$$c = 50 - 24(g + g^{-1}), \quad \Delta_2 = \frac{p_1^2 + p_2^2}{g} + \frac{c - 2}{24}, \quad \Delta_3 = \frac{2p_2(p_2^2 - 3p_1^2)}{(3g)^{3/2}}. \quad (61)$$

⁵The spectral parameter μ is denoted λ in Bazhanov et al. (2002).

4.1 From (58) to (59)

We now prove that the $Q\tilde{Q}$ system (58) and the system (59) are equivalent. As a by-product we deduce the explicit relations (3) among the parameters of the opers, $\tilde{r}^1, \tilde{r}^2, k, \lambda$, and the parameters of the quantum theory, $c, \Delta_2, \Delta_3, \mu$. More precisely, we derive (3a), (3b) and (3c) while (3d) can be found in Bazhanov et al. (2002).

Let $Q_i, Q_i^*, i = 1, 2, 3$ be the functions defined by the expansions (57), satisfying the $Q\tilde{Q}$ -system (58). Assume that $Q_i(0) \neq 0$ and $Q_i^*(0) \neq 0, i = 1, 2, 3$. Recall the definition of the indices $\beta_i, \beta_i^*, i = 1, 2, 3$ as given in (47). Then, a direct calculation shows that the functions

$$P_i(t) = t^{\beta_i} \frac{Q_i(t)}{Q_i(0)}, \quad P_i^*(t) = t^{\beta_i^*} \frac{Q_i^*(t)}{Q_i^*(0)}, \quad i = 1, 2, 3$$

satisfy (59), with the parameters g, p_1, p_2 appearing in (60) related to the parameters \hat{k}, r^1, r^2 by the relations

$$g = 1 - \hat{k} = k + 3, \quad p_1 = \frac{r^1}{2} + \frac{r^2}{2} - 1, \quad p_2 = \frac{\sqrt{3}}{2}(r^1 - r^2). \quad (62)$$

Substituting the above equation into (61) and using (34) we obtain (3a)–(3c).

Acknowledgements The authors are partially supported by the FCT Project PTDC/MAT-PUR/30234/2017 ‘Irregular connections on algebraic curves and Quantum Field Theory’. D. M. is supported by the FCT Investigator grant IF/00069/2015 ‘A mathematical framework for the ODE/IM correspondence’. The authors thank Dipartimento di Matematica dell’Università degli studi di Genova for the kind hospitality.

References

- Bazhanov, V.V., Hibberd, A., Khoroshkin, S.: Integrable structure of W3 conformal field theory, quantum Boussinesq theory and boundary affine Toda theory. Nucl. Phys. B **622**(3), 475–547 (2002)
- Bazhanov, V.V., Lukyanov, S.L., Zamolodchikov, A.B.: Integrable structure of conformal field theory, quantum KdV theory and thermodynamic Bethe ansatz. Commun. Math. Phys. **177**(2) (1996)
- Bazhanov, V.V., Lukyanov, S.L., Zamolodchikov, A.B.: Spectral determinants for Schrodinger equation and Q operators of conformal field theory. J. Statist. Phys. **102**, 567–576 (2001)
- Bazhanov, V.V., Lukyanov, S.L., Zamolodchikov, A.B.: Higher-level eigenvalues of Q-operators and Schrodinger equation. Adv. Theor. Math. Phys. **7**, 711 (2004)
- Dorey, P., Dunning, C., Masoero, D., Suzuki, J., Tateo, R.: Pseudo-differential equations, and the Bethe ansatz for the classical Lie algebras. Nucl. Phys. B **772**(3), 249–289 (2007)
- Dorey, P., Dunning, C., Tateo, R.: The ODE/IM correspondence. J. Phys. A: Math. Theor. **40**(32) (2007)
- Dorey, P., Tateo, R.: Anharmonic oscillators, the thermodynamic Bethe ansatz, and nonlinear integral equations. J. Phys. A **32**, L419–L425 (1999)

- Dorey, Patrick, Tateo, Roberto: Differential equations and integrable models: the $SU(3)$ case. *Nucl. Phys. B* **571**(3), 583–606 (2000)
- Feigin, B., Frenkel, E.: Integrals of motion and quantum groups. In: *Integrable systems and quantum groups*, pp. 349–418. Springer (1996)
- Feigin, B., Frenkel, E.: Quantization of soliton systems and Langlands duality. In: *Exploring new structures and natural constructions in mathematical physics*. *Adv. Stud. Pure Math.*, vol. 61, pp. 185–274. Math. Soc. Japan, Tokyo (2011)
- Frenkel, E., Hernandez, D.: Spectra of quantum Kdv hamiltonians, Langlands duality, and affineopers. *Commun. Math. Phys.* **362**(2), 361–414 (2018)
- Kostant, Bertram: On whittaker vectors and representation theory. *Inventiones mathematicae* **48**, 101–184 (1978)
- Kotousov, G. A., Lukyanov, S. L.: Bethe state norms for the heisenberg spin chain in the scaling limit (2019). [arXiv:1906.07081](https://arxiv.org/abs/1906.07081)
- Masoero, D., Raimondo, A.: Opers for higher states of quantum KdV models. *Commun. Math. Phys.* **378**, 1–74 (2020)
- Masoero, D., Raimondo, A., Valeri, D.: Bethe ansatz and the spectral theory of affine lie algebra-valued connections I. The simply-laced case. *Comm. Math. Phys.* **344**(3), 719–750 (2016)
- Masoero, D., Raimondo, A., Valeri, D.: Bethe ansatz and the spectral theory of affine lie algebra-valued connections II: the non simply-laced case. *Comm. Math. Phys.* **349**(3), 1063–1105 (2017)
- Wasow, W.: *Asymptotic Expansions for Ordinary Differential Equations*. Courier Dover Publications (2018)
- Zamolodchikov, A. B.: Infinite extra symmetries in two-dimensional conformal quantum field theory. *Teoret. Mat. Fiz.* **65**(3), 347–359 (1985)