



Stefan Wirth

Abbreviations

ALT	Alanine aminotransferase
Anti-HBc IgG	Immune globulin G
Anti-HBc IgM	Immune globulin G
Anti-HBc	Antibody against HBcAg
Anti-HBc	Antibody against HBcAg
Anti-HBc	Antibody against HBcAg
Anti-HBe	Antibody against HBeAg
Anti-HBs	Antibody against HBsAg
Anti-HDV	Antibody against delta virus
cccDNA	Covalently closed circular DNA
DAA	Direct-acting antiviral
DNA	Deoxyribonucleic acid
EMA	European Medicines Agency
FDA	Food and Drug Administration
HBcAg	Hepatitis B core antigen
HBeAg	Hepatitis e antigen
HBsAg	Hepatitis B surface antigen
HBV	Hepatitis B virus
HCC	Hepatocellular carcinoma
HCV	Hepatitis C virus
HIV	Human immune deficiency virus
IgG	Immune globulin G
IL2	Interleukin 2
IL28B	Interferon lambda 3 gene
NCTP	Sodium/taurocholate cotransporting polypeptide
ORF	Open reading frame
RNA	Ribonucleic acid
SVR	Sustained viral response
TNF- α	Tumor necrosis factor alpha

Chronic Hepatitis B

Introduction

Hepatitis B virus (HBV) infection remains a global health burden with estimated 250 million people chronically infected worldwide. Nevertheless, since HBV vaccine has become available for more than 25 years and many countries introduced vaccination programs as a prevention strategy on a regular basis for young infants, significant reduction of the incidence of acute hepatitis B in children and adolescents has been observed. Unprotected, approximately 90% of HBV-infected infants and 20–25% of those infected in preschool age will develop chronic infection decreasing to a chronicity rate of around 5% for adolescents and adults [1–3]. Despite a rather benign spontaneous course of the disease during childhood and adolescence, there is a considerable lifetime risk of progressive liver disease, liver cirrhosis, and the development of a hepatocellular carcinoma (HCC), which may eventually reduce life expectancy. Thus, careful long-term monitoring has to be performed, and appropriate treatment options, which unfortunately are not entirely curative at present, have to be considered.

Pathogenesis of Chronic HBV Infection

HBV belongs to a DNA virus family called hepadnaviruses. It contains a partially double-stranded DNA genome with about 3200 nucleotides. The minus strand covers four overlapping open reading frames (ORFs): S, for the surface gene encoding three envelope proteins (hepatitis B surface antigen, HBsAg); C, for the core gene encoding the core protein (hepatitis B core antigen, HBcAg); X, for the regulatory X gene; and P, for the polymerase gene encoding the viral DNA polymerase. By using multiple start codons, HBV is able to encode more than one protein from an ORF. After hepatocyte entry by the NCTP-receptor, the viral envelope is

S. Wirth (✉)
HELIOS University Hospital Wuppertal, Department of Pediatrics,
Witten/Herdecke University, Wuppertal, Germany
e-mail: stefan.wirth@helios-gesundheit.de

removed, and the nucleocapsid reaches the nucleus, where the double strand will be completed and converted into a covalently closed circular DNA (cccDNA). This is an important step, because the majority of cccDNA is then organized into nucleosomes forming the viral minichromosome, which is serving as template for the synthesis of the viral mRNA. The transcripts are translated into the viral proteins, and simultaneously reverse transcription leads to the synthesis of a complete minus strand of HBV DNA. The plus strand can then be synthesized again, and the molecule circularizes. Thus, the replication of HBV is similar to that of a retrovirus. The proteins are synthesized and assembled at the endoplasmic reticulum and eventually discharged by vesicular transport as a Dane particle which contains the complete virus. The cccDNA plays a key role in viral persistence, viral reactivation after treatment withdrawal, and drug resistance. It accumulates in the nucleus of the hepatocyte as a stable minichromosome organized by histone and nonhistone viral and cellular proteins [4, 5]. Persistent HBV replication is associated with a high frequency of integration of HBV sequences into the human host liver cell genome. Enhanced DNA replication and DNA damage occurring during chronic inflammation with cycles of cell death and regeneration increase the availability of DNA ends in host genomic DNA and promote the process of viral integration [6]. Furthermore, it is presumed that certain altered cells are susceptible to the development of additional genetic and epigenetic changes that may lead to the development of malignant cell transformation and HCC.

For the understanding of the different phases during the course of the chronic disease, it is important to realize that the virus itself is not primarily pathogenic to the hepatocyte. The mechanism of cell death is generally accepted to be the result of a cytotoxic T-lymphocyte-mediated immune response of the host to the virus. Additionally, it has been shown that some HBV proteins may be able to induce apoptosis. During the transition from the immune-tolerant to the immune-active phase, a shift from the hepatitis e antigen (HBeAg)-specific Th2 cell tolerance to Th1 cell activation may recognize HBV-related epitopes on hepatocytes resulting in secretion of cytokines such as interleukin (IL)-2 and tumor necrosis factor alpha (TNF- α) and thus activating inflammation [7].

Epidemiology

Overall, the number of infected people is decreasing. Nevertheless, there are still high endemic countries in Asia, Africa, and some parts of South America with an HBsAg prevalence of more than 8%. The Arabian region, parts of the Eastern hemisphere, and Greenland show an HBV prevalence of 2–7%, and in the Western countries the rate is below

2% [8, 9]. Global immunization programs have been established in many countries, and the HBV infection rate has declined worldwide. Vertical transmission has become the main route of infection; nevertheless, in some areas, HBV may also be a predominant disease in adolescents and adults due to high-risk sexual behavior and drug abuse [10]. Unfortunately, up to 2–15% of perinatal HBV infection of antibody against HBeAg (anti-HBe)- and HBeAg-positive mothers cannot be prevented by active and passive immunization due to intrauterine infection, vaccine failure, or HBsAg escape mutants [11, 12]. Thus, passive and active immunization has to be started immediately after birth in all newborns from HBsAg-positive mothers. HBeAg-positive mothers can be considered to receive treatment with the nucleoside analogues telbivudine and tenofovir in late pregnancy to decrease viral load [13, 14]. After complete immunization, there are no objections against breastfeeding. In countries with blood donor screening and serum testing, parenteral transmission does no longer play a significant role. The HBsAg prevalence in children is estimated between 0.02% and 0.03% in Western countries and the USA, 0.14% in Brazil, and 0.5% in Taiwan after immunization [1, 15]. Given HBsAg prevalence in pregnant women of 0.4% in Western Europe and an HBV transmission rate of 5–10% despite complete vaccination, 20–40 newborns in 100,000 births may be infected and become a chronic carrier state.

Ten HBV genotypes (A–J) have been documented showing a distinct distribution. Genotypes A and D are predominant in North America, Europe, and India, and genotypes B and C are mostly found in Asian countries. To date, routine determination of genotype is not yet recommended because treatment options are not adjusted to genotypes. Nevertheless, since there is line of evidence that genotypes C and D may be associated with more aggressive liver disease, this might become significant during the long-term follow-up [16].

Diagnostics

Chronic hepatitis B infection is defined as a repeatedly positive HBsAg test result within 6 months. Apart from the aminotransferases, HBeAg, anti-HBe, anti-HBcIgG, and quantitative HBV DNA have to be determined to confirm chronic hepatitis B and to classify the present stage. Additionally, antibody against delta virus (anti-HDV) should be tested to exclude concomitant hepatitis D. It is recommended to perform an ultrasound examination including liver stiffness assessment for baseline findings. Since chronic hepatitis B usually is a mild disease in terms of inflammation in childhood, histological examination by liver biopsy is not mandatory. However, in subjects who are suspicious of progressive liver disease or cirrhosis or if an impact on therapeutic decisions is identifiable, liver biopsy may be a reasonable completion.

Natural History

There are four natural stages of chronic hepatitis B infection: immune tolerance stage, immune reactive or immune clearance stage, inactive HBsAg carrier stage, and reactivation stage. The immune-tolerant phase is also named chronic hepatitis B infection and the immune-active phase chronic hepatitis B. As fifth and last stage, viral elimination with antibody against HBsAg (anti-HBs) seroconversion, which is a rather rare event occurring not more than 0.5% annually in children, could be denominated [7, 17]. Figure 63.1 illustrates the different phases of chronic hepatitis B.

Children who have HBV infection acquired perinatally or in the first months have an initial tolerance stage which is characterized by the presence of HBsAg, HBeAg, and extremely high HBV DNA levels (10^{7-9} virions/ml) and normal aminotransferases. The duration of the immune-tolerant phase is not predictable and may last 1–4 decades. Asian patients use to have a longer immune-tolerant stage. T helper (Th) cell immune tolerance is generated by HBeAg functioning as an immunoregulatory protein mostly already transplacentally transmitted. This kind of induced immune tolerance may explain the high chronicity rate and the younger the individuals are when they get infected. Usually, only minimal inflammatory activity is detectable in the liver tissue in this phase. Even in adult patients, no severe progression is expected during the immune-tolerant stage [18]. Although antiviral therapy is still not recommended for immune-tolerant subjects, they should be carefully monitored to duly recognize progression to immune-active phase.


With time, a nonspecific increase of inflammatory activity or a decrease of HBeAg serum concentration, which may be due to emerging mutants in the core promotor or precore region resulting in a lower HBeAg production, may activate

HBeAg-specific T cell clones. In this immune reactive phase, HBeAg remains positive, and aminotransferases rise. HBV DNA remains high or stays at a little lower level. During this time, progression to liver fibrosis or cirrhosis may occur. However, liver cirrhosis rate is not expected to exceed 3–5% until reaching adulthood [19–21]. In the immune reactive phase treatment has to be considered.

A key event in the natural course of chronic hepatitis B is the HBeAg/anti-HBe seroconversion which occurs unpredictably for the single individual during the immune-active phase. Anti-HBe seroconversion is associated with a significant decrease of viral replication and normalization of aminotransferases reflecting the biochemical and histological remission of inflammatory activity. In some studies, the annual seroconversion rate depends on the route of infection and the ethnic origin. Whereas the mean seroconversion rate in non-Asian children ranges between 8% and 15%, seroconversion in Asian children was considerably lower with approximately 5% per year [7, 20]. Anti-HBe seroconversion is followed by the inactive HBsAg carrier state with persistently normal aminotransferases and a low viral load. Viral replication is considered low when HBV DNA serum concentrations remain below 2000 IU/ml. Some carriers may be lucky and develop anti-HBs antibodies indicating viral elimination and cure of the disease. The estimated incidence of this rare event in children is 0.05–0.8% per year in endemic areas with predominantly perinatal HBV transmission [7, 15].

Approximately 20–30% of inactive HBsAg carriers will experience spontaneous reactivation during long-term follow-up. Those episodes may cause progressive liver damage. The reactivation phase is characterized by the presence of anti-HBe and elevated aminotransferases. HBV DNA levels rise over 2000 IU/ml. This status is also named HBeAg-

Fig. 63.1 Illustration of the different phases of chronic hepatitis B. *HBV-DNA* hepatitis B virus deoxyribonucleic acid, *DNA-PCR* deoxyribonucleic acid polymerase chain reaction, *HBV* hepatitis B virus, *HBsAg* hepatitis B surface antigen, *Anti-HBc* antibody against HBcAg, *ALT* alanine aminotransferase, *HBeAg* hepatitis e antigen, *Anti-HBs* antibody against HBsAg



negative hepatitis B. However, reactivation rarely occurs during childhood and adolescents.

Another particular condition warrants mention: occult HBV infection. It is defined as the existence of HBV DNA in serum among HBsAg-negative patients and can be classified into seropositive and seronegative with respect to the presence of anti-HBs or antibody against HBcAg (anti-HBc) antibodies. Possible explanations are low levels of viral replication activity or the emergence of HBV variants in the a-determinant of the S-gene. Occult hepatitis B is most common in endemic regions and seems rare with 1.4% [22]. However, prevalence may rise considerably in immunized children from HBsAg-positive mothers. One study reported a prevalence of 28% in this special group [23].

Long-Term Prognosis

Individuals with chronic HBV infection are at risk to develop long-term sequelae such as end-stage liver disease including liver cirrhosis, hepatic failure, and HCC. Progression strongly correlates with the disease activity in terms of viral replication level, inflammatory activity, HBsAg levels, HBV genotypes, and HBeAg/anti-HBe status. Strong risk factors for developing liver cirrhosis and HCC are higher age, male, presence of HBeAg, HBV DNA levels $>10^4$ copies/ml, HBsAg serum concentrations $>10^3$ IU/ml, and alanine aminotransferase (ALT) >45 IU/l [24, 25]. Progression to liver cirrhosis in children is under 5% until adulthood, and data of the Asian region report 0.01–0.003% of individuals with chronic hepatitis B to be expected developing HCC in childhood [26, 27]. In general, anti-HBe seroconversion significantly reduces the risk of developing HCC. The time at which anti-HBe seroconversion occurs is important. A study in adults investigating the 15-year cumulative incidences of HBeAg-negative hepatitis demonstrated that cirrhosis and HCC increased with increasing age of HBeAg seroconversion [28]. The lowest risk was observed in patients with anti-HBe seroconversion under the age of 30 (cirrhosis 7%, HCC 2.1%) and highest in individuals older than 40 years (cirrhosis 42.9%, HCC 7.7%). The hazard ratio for HBeAg-negative hepatitis, cirrhosis, and HCC was 2.95, 17.6, and 5.22, respectively, in the older compared with the younger group. The authors concluded that patients with HBeAg seroconversion before age 30 have an excellent prognosis, whereas patients with delayed HBeAg seroconversion after age 40 have significantly higher incidences of HBeAg-negative hepatitis, cirrhosis, and HCC. An additional precondition is persistently normal ALT levels [29]. Since children have a high probability to experience anti-HBe seroconversion until adulthood, the overall risk of developing severe liver disease in later life seems limited. Nevertheless, there remain a considerable number of patients with immune tolerance or

inflammatory activity that needs careful and professional monitoring.

Relevance of Genotypes and Mutants

During the replication cycle, HBV polymerase is acting as a reverse transcriptase without proofreading function. Therefore, mutant viral genomes are regularly emerging in a considerable number particularly during the high replicative status. Peculiar requirements such as replication modalities, selection pressure, and changing immunological conditions may select variants and strongly influence the predominant HBV quasispecies in an infected individual. Generally, a change of the primarily determined genotype is possible during long-term course and ranges between 2.8% and 19% usually associated with anti-HBe seroconversion [26, 30]. It is not yet known if there is any clinical impact at all. In adults, genotype C infection rather than genotype B is associated with a delayed anti-HBe seroconversion and a higher risk of developing HCC. Genotype D tends to proceed more severely and shows delayed anti-HBe seroconversion compared with genotype A. Precore and basic core promotor (BCP) mutants are frequently associated with HBeAg-negative hepatitis, and HBsAg escape mutants are now increasingly observed in association with primarily vaccinated children. The typical precore point mutant is the G1896A stop codon preventing the production of HBeAg. It emerges typically around the time of anti-HBe seroconversion and may be associated with a decreased risk of developing HCC compared with the wild type. But it can also be found in patients with HBeAg-negative hepatitis. Depending on European or Asian regions, precore mutants have been detected between 8% and 50% in HBeAg-negative children. The BCP mutants A1762T/G1764A prevail to be associated with an increased risk for HCC. But, finally, the data remain controversial [7, 16, 31, 32].

Treatment

Since there is no definite curative medical treatment available to date, it has to be defined what the aim of antiviral treatment should be in dependence on age group and phase of chronic hepatitis B. There is no doubt that one major goal is to reduce the risk of progressive liver disease and long-term sequelae such as liver cirrhosis, hepatic decompensation, and HCC and eventually to achieve the same life expectancy compared with healthy individuals of the same age. Unfortunately, anti-HBs seroconversion can only be reached in 5–10% at the most under current medical treatment strategies. Thus, the most important task in the treatment of children and adolescents is to achieve anti-HBe

seroconversion at the earliest possible time associated with suppressed viral replication and decreased liver inflammation followed by persistent presence of anti-HBe, undetectable HBV DNA, and preferably aminotransferase values less than half of the upper limit of normal. Children with HBeAg-positive hepatitis should be monitored every 6 months with physical examination, measurement of laboratory parameters such as aminotransferases, hepatitis B serology, alpha-fetoprotein, and ultrasound of the liver. After anti-HBe seroconversion, follow-up visits can be performed for lifetime on an annual basis [15, 17, 33].

The decision to treat should be based on age, phase of HBV infection determined by ALT level, HBeAg/anti-HBe status, liver histology, coexisting diseases, and expectable compliance. The response rate in patients in the immune-tolerant phase is very low. Thus, treatment of children with normal aminotransferases is not recommended. Recently published clinical trials in immune-tolerant children and adults combining a nucleoside analogue and peg-alpha interferon did not achieve a higher anti-HBe seroconversion rate compared to the control group [34, 35]. However, treatment should be considered when aminotransferases rise and transition to the immune-active phase is recognized. Children and adolescents who have persistently elevated ALT levels for more than 6 months should be offered treatment. Currently, seven treatment options are approved for hepatitis B in adults, including two formulations of conventional and pegylated interferon as an immunomodulatory therapy and five nucleos(t)ide analogues (lamivudine, adefovir dipivoxil, entecavir, telbivudine, and tenofovir disoproxil) with strong reduction of viral replication. Approval for children and adolescents depends on the region. Large trials have been performed in children for lamivudine, adefovir, entecavir, and tenofovir. Entecavir has been authorized from 2 years onward, and adefovir and tenofovir disoproxil have been approved for subjects older than 12 years of age in the USA and Europe [36–38]. Predictors of response may be increased ALT levels, relatively low HBV DNA levels, and infection with genotype A or B. The main problem of all clinical trials with nucleos(t)ide analogues is the duration of medical treatment of not more than 96 weeks. Although a high proportion of treated patients will experience a significant decrease of viral load, the anti-HBe seroconversion rate cannot be expected to exceed 25%. After ceasing treatment, a reactivation of viral replication to baseline levels can be observed. A 24-week course of alpha interferon yields an approximately 10% higher anti-HBe seroconversion rate. Anti-HBs seroconversion rate is limited to single cases with nucleos(t)ide analogues and may range between 6% and 10% in patients with alpha interferon. There is no doubt that these results are dissatisfying with respect to our primary goal of anti-HBe seroconversion. Another interesting fact is that alpha interferon treatment only accelerates anti-HBe seroconversion in

successfully treated individuals but does not enhance the absolute number of responders [39]. Extending the treatment with nucleos(t)ide analogues for several years will result in an anti-HBe seroconversion rate of 40–50% [33]. However, there are no long-term data in children with regard to side effects. In the case of anti-HBe seroconversion, treatment should be maintained for 12 months, because the treatment-induced anti-HBe-positive status may be instable and reactivation may occur [40, 41].

In view of the present data and experience, there is a remarkable counseling conflict between the choice of drug and the duration of treatment, given that anti-HBe seroconversion remains the essential goal. Antiviral drug resistance is a major limitation to the long-term success of antiviral treatment. For this reason, lamivudine with a 5-year resistance rate of 70% has been considered obsolete just as adefovir which does often not sufficiently suppress viral replication. Nevertheless, at least for smaller children, lamivudine can be used as an approved drug for a limited time. Telbivudine has also a considerable resistance risk and is not approved. Entecavir and tenofovir do not show significant resistances after years of treatment. Tenofovir disoproxil may be associated with an increase in serum creatinine levels after 3–5 years of therapy. Decrease of bone mineral density has also been reported. These side effects might occur less frequent with the new tenofovir alafenamide. Oral treatment with nucleos(t)ide analogues is quite comfortable but needs a real true commitment to the treatment, and alpha interferon may have sometimes restrictive side effects but with the advantage of a defined duration.

Thus, the decision which treatment option to choose is not that easy and has to be achieved in agreement with the patient and the parents. Alpha interferon is particularly appropriate for those children and adolescents who are reluctant to commit to a long duration of treatment and are not in the pubertal growth spurt. Nowadays, peg-alpha interferon should be recommended for 48 weeks. Nucleos(t)ide analogues are most appropriate for patients with contraindications to interferon, after liver transplantation with an anti-HBc-positive donor or under immune suppressive treatment. It is most important that they are willing to commit to a treatment for several, probably 3–5, years, maybe longer. Entecavir and tenofovir have the best profile in terms of safety, efficacy, and drug resistance. For younger patients, entecavir seems actually the preferable option.

Children and adolescents with a HBeAg-negative hepatitis should be treated with a nucleos(t)ide analogue if ALT levels are elevated, and HBV DNA concentration is above 20,000 IU/ml to prevent progressive liver disease [42]. During long-term treatment with nucleos(t)ide analogues, HBV DNA, HBeAg/anti-HBe status, and aminotransferase levels should be monitored every 3 months. Very low or negative HBV DNA concentrations are important preconditions to avoid drug resistance.

Prevention

Vaccination is the most effective procedure in order to prevent infection with the HBV. Active and passive immunization is well established in newborns of HBsAg-positive mothers. The first injections have to be administered within 12–24 h after birth to achieve a seroprotective response in 90–95% when two monthly follow-up active vaccinations are completed. Very-low-birth-weight preterm infants should receive a total of four doses. HBeAg-positive mothers can be treated with a nucleoside analogue (telbivudine, tenofovir) during the last trimester of pregnancy to reduce the risk of vertical transmission [14].

In many countries, routine active HBV vaccination is implemented in the vaccination schedule of all infants. Postvaccination testing for a protective anti-HBs concentration (> 100 IU/l) is not routinely recommended. If indicated, the best time would be approximately 2–3 months after the last vaccination. Revaccination is indicated in subjects with an anti-HBs titer <10 IU/l. In the majority of nonresponders, three more vaccinations will induce protective response. According to present experiences, protective anti-HBs response will be maintained for more than 15 years [13, 41].

Chronic Hepatitis C

Introduction

Hepatitis C virus (HCV) infection is a frequent course of chronic liver disease, and approximately 71 million people are estimated to be chronically infected worldwide. The number of chronically infected children up to the age of 15 is estimated with approximately 13 million. Unfortunately, to date, no preventive vaccination could be developed. Despite a normally benign course of the disease during childhood and adolescence, there is a considerable lifetime risk of progressive liver disease, liver cirrhosis, and the development of a HCC, which may eventually reduce life expectancy. Remarkable advances have been made in therapeutic approaches so far, and considerable rates of cure have been yielded with the former interferon-based treatment standard of care. In the last years, interferon-free treatment regimen using direct-antiviral agents (DAA) has become the new standard of care, yielding excellent results for viral elimination. DAAs are now also available for children and adolescents.

Pathogenesis of Chronic Hepatitis C Infection

HCV is a positive-stranded RNA virus within the *Flaviviridae* family. It forms its own genus *Hepacivirus*,

and there are six main genotypes. The viral genome encodes nine proteins including its own RNA polymerase. Because of the high error rate of the virus-specific RNA polymerase, many variants may be produced. So-called quasispecies represent the high variability of the virus which allows a survival advantage to the virus. Replication of HCV starts with the binding to hepatocytes and entry which is a rather complex procedure. RNA is released into the cytoplasm and translated in the rough endoplasmic reticulum. A 3000-amino-acid-long polypeptide arises and is then cleaved into ten different products. Membranous replication vesicles are induced, and HCV assembly is accomplished and released with the help of very-low-density lipoprotein (VLDL) synthesis. Chronic hepatitis C in children is associated with a variety of histological patterns, mostly considered as mild and slow progressive. Nevertheless, significant fibrosis or cirrhosis may occur but is not expected to exceed 4% until reaching adulthood. Need for liver transplantation is very rare as is the development of HCC [43]. Little information is available about the host response to the virus. Cluster of differentiation 4 (CD4) + lymphocytes seem to be involved. Infants with the rs 12,979,860 CC genotype for the IL28B polymorphism tend to experience a higher spontaneous viral elimination [44–46].

Epidemiology

The prevalence of HCV infection in children in developed countries ranges between 0.1% and 0.4%. For adults, prevalence rates are 0.4–3% in North America and Western Europe and higher in Eastern Europe and Middle East. Egypt has the highest prevalence with 9%, almost exclusively genotype 4 [47, 48]. Central and East Asia and North Africa are estimated to have a prevalence between 3.6% and 3.8% [49]. During the last 15 years, the predominant route of viral hepatitis C transmission has become vertical infection. Contamination through blood products is exceedingly rare in developed countries but may remain an issue in developing countries. The rate of perinatal transmission from an HCV-RNA-positive mother ranges from 2% to 5%. Out of this group, a considerable number of infants received the infection probably already in utero [50]. Concomitant HIV infection may increase the risk of HCV transmission. Breastfeeding does not promote viral transmission and is allowed. The HCV prevalence in pregnant women from North America and Central Europe was reported between 0.16% and 0.53%. Assumed a perinatal transmission rate of 2–4%, 8–10 newborns in 100,000 births per year may be infected and become chronically infected during the first year of life. Viral clearance in vertically infected children seems to be dependent on the genotype and was reported to range from 2.4% to 25%. In contrast, children infected with genotype 3 had a higher

spontaneous clearance rate compared to individuals with genotype 1. Beyond the age of 5 years, spontaneous viral elimination becomes less likely [51, 52]. In view of the fact that more and more patients can be cured with DAAs, it is expected that the prevalence of chronic hepatitis C infection will decrease significantly within the next 5 years.

Diagnosis

Serologic testing for anti-HCV antibodies is the appropriate screening test for HCV. The next diagnostic step is the determination of quantitative HCV RNA and the genotype. The most prevalent genotype in pediatric trials performed in Western countries was genotype 1 (ca. 74%) followed by genotype 3 (ca. 14%) and 2 (ca. 9%). Genotype 4 had the lowest prevalence (ca. 3%) [53]. It is useful to perform an ultrasound examination including liver stiffness assessment for the baseline report. As chronic hepatitis C usually is a histologically mild disease with low inflammatory activity in childhood, liver biopsy is not mandatory. However, in subjects, who are suspicious of progressive liver disease or cirrhosis or if there is an impact on therapeutic decisions, liver biopsy may be a reasonable measure [53].

Several studies have demonstrated that certain host polymorphisms (e.g., CC) located upstream of the IL28B (interferon lambda 3) gene are associated with a higher sustained viral response rate to combination treatment with peg-alpha interferon and ribavirin. There is also an association with spontaneous clearance of HCV. So far, the determination of IL28B polymorphisms has not been used routinely and is becoming far less important in the treatment with DAAs.

Natural History

Normally, HCV infection is asymptomatic. Histological findings are usually mild, and the risk of severe complications until the infected individuals are reaching adulthood is low. Not more than 5% of children and adolescents will have evidence of advanced liver fibrosis or cirrhosis. Liver transplantation units from the USA have been reported on 133 transplanted children due to chronic hepatitis C during a time span of 13 years. In a lifetime, the risk of developing liver cirrhosis is about 20%, and the risk of HCC based on liver cirrhosis is estimated to be 2–5% [45]. These data are from adults, and there are no long-term follow-up studies in vertically infected patients. Natural history is also affected by other medical and social factors. Overweight children with liver steatosis are at greater risk for progressive liver disease. Risky behaviors and alcohol misuse worsen the long-term prognosis. Similar course and progression of hepatitis C were reported in former pediatric patients with successfully treated malignant disease

after three decades of observation with about 20% spontaneous clearance and up to 5% liver cirrhosis. During the chronic course, ALT levels may be normal or intermittently elevated. Only few patients show persistent markedly elevated aminotransferases. Also the HCV RNA serum concentration may considerably fluctuate but without immediate prognostic relevance. Spontaneous resolution beyond the preschool age is quite rare and may occur in up to 10% of adolescents [45, 48, 54]. Some extrahepatic manifestations may be associated with chronic hepatitis C such as glomerulonephritis and possibly cognitive deficits or developmental delay [55]. However, in adult patients, the clinical effects of reported symptoms such as fatigue, depression, or marginal poorer learning efficiency were rather limited [56]. In conclusion, early acquired chronic hepatitis C is a clinically and histologically silent and hidden condition. Nevertheless, it may become insidious. Although the rate of developing liver cirrhosis until adulthood is low, activation beyond the second decade of life is likely. A large Danish study in adults revealed that in patients with chronic HCV infection, the 8-year risk of liver-related death was 5.5% compared with 2.0% in individuals who cleared the infection [57]. However, patients who have eventually proceeded to compensated liver cirrhosis have a dubious prognosis. In a follow-up study of cirrhotic patients over more than 10 years, HCC developed in 32%, and the annual mortality rate was 4% [58]. Thus, the literal aim of therapeutic interventions in children and adolescents is not the treatment of an ongoing liver disease but the prevention of a future one by early eradication of the infection.

Treatment

The primary goal of HCV therapy is to cure the infection, which is reflected by persistently negative HCV RNA in serum and normalized aminotransferases. Sustained viral response (SVR) is defined as an undetectable HCV RNA level 24 weeks after cessation of treatment. The interferon-based treatment regimens included the combination of pegylated alpha interferon with ribavirin and a protease inhibitor. They were the approved and established standard of care in adults until the year 2014. With these regimens, considerable sustained viral response rates could be achieved ranging from more than 65% for genotype 1 to over 80% for genotype 2 and 3 patients [59]. Then, the interferon-free era started with the introduction of direct antiviral agents (DAA) focusing on genotypes 1 and 4 [60]. Genotypes 2 and 3 were also treated in different combinations over 12 to 24 weeks. The most important substance was sofosbuvir. However, more and more combinations were tested and approved, and numerous drugs were on the market only for a few years [61]. Table 63.1 summarizes the currently mostly used preparations with respect to the approval status in adults and children.

Table 63.1 Direct antiviral agents for the treatment of chronic hepatitis C

Substances	Registered trade name	Genotype	Trials in children	Approved >12 years of age ^a	Approved >3 years of age ^a
Sofosbuvir/Ledipasvir	Harvoni	1, 4	Yes	Yes	Yes
Ribavirin/Sofosbuvir ^a	Ribavirin, Sovaldi	2, 3	Yes	Yes	Yes
Ritonavir/Ombitasvir/Dasabuvir/Ribavirin ^a	Viekirax, Exviera	1, 4	Yes	Not intended	Not intended
Daclatasvir/Sofosbuvir ^b	Daklinza, Sovaldi		Yes	Not intended	Not intended
Grazoprevir/Elbasvir	Zepatier	1, 4	Yes	In question	In question
Sofosbuvir/Velpatasvir	Epclusa	1–6	Yes	Yes	Yes, > 6 years
Sofosbuvir/Velpatasvir/Voxilaprevir ^c	Vosevi	1–6	Yes	Not intended	Not intended
Glecaprevir/Pibrentasvir ^c	Maviret	1–6	Yes	Yes	Yes

^a Approval by FDA and EMA

^b No longer in use in adults

^c Duration of treatment 8 weeks in non-cirrhotic patients

Experiences with the treatment of children with chronic hepatitis C started in the early 1990s. Nineteen studies using recombinant alpha interferon were published between 1992 and 2003. A meta-analysis of trials with alpha interferon monotherapy showed a wide range of viral response (0–76%). Based on an increasing number of trials in adults, ribavirin was also added to alpha interferon treatment trials for children. Between the years 2000 and 2005, six studies were published showing a sustained viral response rate from 27% to 64% [52]. It became clear that genotype-2- and genotype-3-infected individuals responded much better. Alpha interferon-2b in combination with ribavirin was then approved by the FDA. Trials with peg-alpha interferon and ribavirin followed in the next years, and both peg-alpha interferon-2b and peg-alpha interferon-2a have been approved by FDA and EMA 2008/2009 and 2011/2012 in combination with ribavirin for children. Peg-alpha interferon and ribavirin therapies in treatment-naïve children and adolescents yield a sustained viral response rate in approximately 50% of adequately treated genotype-1-infected patients. Thus, this option could theoretically be offered to all interested individuals. In patients infected with genotype 2 or 3, treatment for 24 weeks has a response rate of more than 90% [62–64]. It could be used in children from 3 years and peg-alpha interferon-2a from 5 years onward, but it is also no longer recommended. Since spontaneous viral elimination in vertically infected subjects may occur within the first 3 to 4 years, the start of therapy was agreed after the age of three.

Treatment management of children with chronic hepatitis C infection is formed by the attitude of the medical attendant regarding the need of therapeutic intervention with respect to a generally slow progressive disease. In general, also for children and adolescents, interferon-based treatment options are considered obsolete, despite approval. Adverse events during treatment were frequent, and the duration was 24 to 48 weeks in dependence on the genotype. Under the aspect of health prevention for a long lifetime, all children with a measureable level of HCV RNA should be treated. The level

neither of aminotransferases nor of HCV RNA predicts the long-term outcome of the disease. Also, liver histology is not a helpful entry criterion for indicating treatment, because children generally do not have severe lesions. With the development of the new DAAs, there is no reason not to treat children and adolescents after diagnosis of chronic hepatitis C from 3 years of age onward [65, 66]. The drugs are well tolerated and have only an exceptionally low rate of side effects.

Three groups of DAAs were differentiated in three groups: NS3/4A protease inhibitors, NS5a inhibitors, and polymerase inhibitors. Protease inhibitors have the ending “previr,” NS5a inhibitors have the ending “asvir,” and polymerase inhibitors are ending with “buvir.” Sofosbuvir is one of the most potent polymerase inhibitors. It is important to combine at least two of different groups to avoid developing rapid drug resistance.

For the pediatric population, several clinical trials have been performed. Due to the approval regulations, the first groups included were between 13 and 18 years of age, followed by the younger ages. The combination of sofosbuvir and ledipasvir is approved from 3 years onward and suitable for the treatment of genotypes 1 and 4. It is administered for 12 weeks [67–69]. For genotypes 2 and 3, sofosbuvir and ribavirin are approved. Genotype 2 has to be treated for 12 weeks and genotype 3 for 24 weeks [70, 71], which can now be considered second line. The most recent approval is the combination of glecaprevir and pibrentasvir, which is pangenotypic (genotypes 1–6) and has the additional advantage of an 8-week treatment duration [72]. With Sofobuvir/Velpatasvir there are now two pangenotypic active drugs available. This panel seems sufficient to successfully addressing the pediatric chronic HCV infections. The overall sustained viral response rate was extremely high and achieved in most of the trials 100%. That was also true for individuals who were treatment experienced with a previous interferon-based therapy. Relapses were observed only in few isolated cases. It is recommended to determine HCV RNA 4 weeks after the start of treatment. Most patients are already negative at that time. The most important checkpoint is SVR

24, 24 weeks after cessation of treatment, as HCV RNA negativity is associated with long-term elimination. The safety and tolerability profile of the DAAs is also excellent. The vast majority of adverse events were mild and unrelated to the medication. Headache, fatigue, or signs of upper respiratory tract infections were registered up to about 15% [72]. In adult trials, there was no difference to the side effects of the placebo arm. Treatment does not appear to have an impact on growth and development.

Overall, early treatment of the pediatric population with chronic hepatitis C will decrease the pool of infected individuals and not only alleviate the development of a progressive liver disease but also prevent further transmission in the hope to eradicate the disease in this age group.

References

- Nel E, Sokol RJ, Comparcola D, et al. Viral hepatitis in children. *J Pediatr Gastroenterol Nutr.* 2012;55:500–5.
- McMahon BJ, Alward WL, Hall DB, et al. Acute hepatitis B virus infection: relation of age to the clinical expression of disease and subsequent development of the carrier state. *J Infect Dis.* 1985;151:599–603.
- Chatzidaki V, Kouroumalis E, Galanakis E. Hepatitis B virus acquisition and pathogenesis in childhood: host genetic determinants. *J Pediatr Gastroenterol Nutr.* 2011;52:3–8.
- Trautwein C. Mechanisms of hepatitis B virus graft reinfection and graft damage after liver transplantation. *J Hepatol.* 2004;41:362–9.
- Levrero M, Pollicino T, Petersen J, et al. Control of cccDNA function in hepatitis B virus infection. *J Hepatol.* 2009;51:581–92.
- Bonilla Guerrero R, Roberts LR. The role of hepatitis B virus integrations in the pathogenesis of human hepatocellular carcinoma. *J Hepatol.* 2005;42:760–77.
- Shi YH, Shi CH. Molecular characteristics and stages of chronic hepatitis B virus infection. *World J Gastroenterol.* 2009;15:3099–105.
- Hadziyannis SJ. Natural history of chronic hepatitis B in Euro-Mediterranean and African countries. *J Hepatol.* 2011;55:183–91.
- Jefferies M, Rauff B, Rashid H, Lam T, Rafiq S. Update on global epidemiology of viral hepatitis and preventive strategies. *World J Clin Cases.* 2018;6:589–99.
- Mahtab MA, Rahman S, Khan M, Karim F. Hepatitis B virus genotypes: an overview. *Hepatobiliary Pancreat Dis Int.* 2008;7:457–64.
- Lee C, Gong Y, Brok J, Boxall EH, Gluud C. Effect of hepatitis B immunisation in newborn infants of mothers positive for hepatitis B surface antigen: systematic review and meta-analysis. *BMJ.* 2006;332:328–36.
- Lee C, Gong Y, Brok J, Boxall EH, Gluud C. Hepatitis B immunisation for newborn infants of hepatitis B surface antigen-positive mothers. *Cochrane Database Syst Rev.* 2006;2:CD004790.
- Han GR, Cao MK, Zhao W, et al. A prospective and open-label study for the efficacy and safety of telbivudine in pregnancy for the prevention of perinatal transmission of hepatitis B virus infection. *J Hepatol.* 2011;55:1215–21.
- Aslam A, Campoverde Reyes KJ, Malladi VR, Ishtiaq R, Lau DTY. Management of chronic hepatitis B during pregnancy. *Gastroenterol Rep (Oxf).* 2018;6:257–62.
- Paganelli M, Stephenne X, Sokal EM. Chronic hepatitis B in children and adolescents. *J Hepatol.* 2012;57:885–96.
- Wai CT, Fontana RJ. Clinical significance of hepatitis B virus genotypes, variants, and mutants. *Clin Liver Dis.* 2004;8:321–52. vi
- Sokal EM, Paganelli M, Wirth S, et al. Management of chronic hepatitis B in childhood: ESPGHAN clinical practice guidelines: consensus of an expert panel on behalf of the European Society of Pediatric Gastroenterology, Hepatology and Nutrition. *J Hepatol.* 2013;59(4):814–29.
- Hui CK, Leung N, Yuen ST, et al. Natural history and disease progression in Chinese chronic hepatitis B patients in immune-tolerant phase. *Hepatology.* 2007;46:395–401.
- Bortolotti F. Treatment of chronic hepatitis B in children. *J Hepatol.* 2003;39(Suppl 1):S200–5.
- Popalis C, Yeung LT, Ling SC, Ng V, Roberts EA. Chronic hepatitis B virus (HBV) infection in children: 25 years' experience. *J Viral Hepat.* 2013;20:e20–6.
- Iorio R, Giannattasio A, Cirillo F, D'Alessandro L, Vegnente A. Long-term outcome in children with chronic hepatitis B: a 24-year observation period. *Clin Infect Dis.* 2007;45:943–9.
- Minuk GY, Sun DF, Uhanova J, et al. Occult hepatitis B virus infection in a North American community-based population. *J Hepatol.* 2005;42:480–5.
- Shahmoradi S, Yahyapour Y, Mahmoodi M, et al. High prevalence of occult hepatitis B virus infection in children born to HBsAg-positive mothers despite prophylaxis with hepatitis B vaccination and HBIG. *J Hepatol.* 2012;57:515–21.
- Lee MH, Yang HI, Liu J, et al. Prediction models of long-term Cirrhosis and hepatocellular carcinoma risk in chronic hepatitis B patients: risk scores integrating host and virus profiles. *Hepatology.* 2013;58:546–54.
- Tseng TC, Liu CJ, Yang HC, et al. Serum hepatitis B surface antigen levels help predict disease progression in patients with low hepatitis B virus loads. *Hepatology.* 2013;57:441–50.
- Ni YH, Chang MH, Wang KJ, et al. Clinical relevance of hepatitis B virus genotype in children with chronic infection and hepatocellular carcinoma. *Gastroenterology.* 2004;127:1733–8.
- Chang MH, You SL, Chen CJ, et al. Decreased incidence of hepatocellular carcinoma in hepatitis B vaccines: a 20-year follow-up study. *J Natl Cancer Inst.* 2009;101:1348–55.
- Chen JD, Yang HI, Iloeje UH, et al. Carriers of inactive hepatitis B virus are still at risk for hepatocellular carcinoma and liver-related death. *Gastroenterology.* 2010;138:1747–54.
- Tai DI, Lin SM, Sheen IS, et al. Long-term outcome of hepatitis B e antigen-negative hepatitis B surface antigen carriers in relation to changes of alanine aminotransferase levels over time. *Hepatology.* 2009;49:1859–67.
- Wirth S, Bortolotti F, Brunert C, et al. Hepatitis B virus genotype change in children is closely related to HBeAg/anti-HBe seroconversion. *J Pediatr Gastroenterol Nutr.* 2013;57:363–6.
- Kao JH. Hepatitis B viral genotypes: clinical relevance and molecular characteristics. *J Gastroenterol Hepatol.* 2002;17:643–50.
- Alavian SM, Carman WF, Jazayeri SM. HBsAg variants: diagnostic-escape and diagnostic dilemma. *J Clin Virol.* 2013;57:201–8.
- Kwon H, Lok AS. Hepatitis B therapy. *Nat Rev Gastroenterol Hepatol.* 2011;8:275–84.
- Feld JJ, Terrault NA, Lin HS, et al. Entecavir and Peginterferon alfa-2a in adults with hepatitis B e antigen-positive immune-tolerant chronic hepatitis B virus infection. *Hepatology.* 2019;69:2338–48.
- Rosenthal P, Ling SC, Belle SH, et al. Combination of Entecavir/Peginterferon alfa-2a in children with hepatitis B e antigen-positive immune tolerant chronic hepatitis B virus infection. *Hepatology.* 2019;69:2326–37.
- Murray KF, Szenborn L, Wysocki J, et al. Randomized, placebo-controlled trial of tenofovir disoproxil fumarate in adolescents with chronic hepatitis B. *Hepatology.* 2012;56:2018–26.
- Jonas MM, Little NR, Gardner SD. Long-term lamivudine treatment of children with chronic hepatitis B: durability of therapeutic responses and safety. *J Viral Hepat.* 2008;15:20–7.

38. Jonas MM, Mizerski J, Badia IB, et al. Clinical trial of lamivudine in children with chronic hepatitis B. *N Engl J Med.* 2002;346:1706–13.
39. Bortolotti F, Iorio R, Nebbia G, et al. Interferon treatment in children with chronic hepatitis C: long-lasting remission in responders, and risk for disease progression in non-responders. *Dig Liver Dis.* 2005;37:336–41.
40. Sokal EM, Kelly DA, Mizerski J, et al. Long-term lamivudine therapy for children with HBeAg-positive chronic hepatitis B. *Hepatology.* 2006;43:225–32.
41. EASL clinical practice guidelines: management of chronic hepatitis B virus infection. *J Hepatol.* 2012;57:167–85.
42. Papatheodoridis GV, Manesis EK, Manolakopoulos S, et al. Is there a meaningful serum hepatitis B virus DNA cutoff level for therapeutic decisions in hepatitis B e antigen-negative chronic hepatitis B virus infection? *Hepatology.* 2008;48:1451–9.
43. Robinson JL, Doucette K. The natural history of hepatitis C virus infection acquired during childhood. *Liver Int.* 2012;32:258–70.
44. Bartenschlager R, Cosset FL, Lohmann V. Hepatitis C virus replication cycle. *J Hepatol.* 2010;53:583–5.
45. Mack CL, Gonzalez-Peralta RP, Gupta N, et al. NASPGHAN practice guidelines: diagnosis and management of hepatitis C infection in infants, children, and adolescents. *J Pediatr Gastroenterol Nutr.* 2012;54:838–55.
46. Ruiz-Extremera A, Munoz-Gamez JA, Salmeron-Ruiz MA, et al. Genetic variation in interleukin 28B with respect to vertical transmission of hepatitis C virus and spontaneous clearance in HCV-infected children. *Hepatology.* 2011;53:1830–8.
47. Ghany MG, Strader DB, Thomas DL, Seeff LB. Diagnosis, management, and treatment of hepatitis C: an update. *Hepatology.* 2009;49:1335–74.
48. Wirth S, Kelly D, Sokal E, et al. Guidance for clinical trials for children and adolescents with chronic hepatitis C. *J Pediatr Gastroenterol Nutr.* 2011;52:233–7.
49. Mohd Hanafiah K, Groeger J, Flaxman AD, Wiersma ST. Global epidemiology of hepatitis C virus infection: new estimates of age-specific antibody to HCV seroprevalence. *Hepatology.* 2013;57:1333–42.
50. Mok J, Pembrey L, Tovo PA, Newell ML. When does mother to child transmission of hepatitis C virus occur? *Arch Dis Child Fetal Neonatal Ed.* 2005;90:F156–60.
51. Bortolotti F, Indolfi G, Zancan L, et al. Management of chronic hepatitis C in childhood: the impact of therapy in the clinical practice during the first 2 decades. *Dig Liver Dis.* 2011;43:325–9.
52. Bortolotti F, Verucchi G, Camma C, et al. Long-term course of chronic hepatitis C in children: from viral clearance to end-stage liver disease. *Gastroenterology.* 2008;134:1900–7.
53. Wirth S. Current treatment options and response rates in children with chronic hepatitis C. *World J Gastroenterol.* 2012;18:99–104.
54. Batsis ID, Wasuwanich P, Karnsakul WW. The management of hepatitis B and hepatitis C in children. *Minerva Pediatr.* 2019;71:59–75.
55. Rodrigue JR, Balistreri W, Haber B, et al. Impact of hepatitis C virus infection on children and their caregivers: quality of life, cognitive, and emotional outcomes. *J Pediatr Gastroenterol Nutr.* 2009;48:341–7.
56. McAndrews MP, Farcnik K, Carlen P, et al. Prevalence and significance of neurocognitive dysfunction in hepatitis C in the absence of correlated risk factors. *Hepatology.* 2005;41:801–8.
57. Omland LH, Krarup H, Jepsen P, et al. Mortality in patients with chronic and cleared hepatitis C viral infection: a nationwide cohort study. *J Hepatol.* 2010;53:36–42.
58. Sangiovanni A, Prati GM, Fasani P, et al. The natural history of compensated cirrhosis due to hepatitis C virus: a 17-year cohort study of 214 patients. *Hepatology.* 2006;43:1303–10.
59. Aghemo A, De Francesco R. New horizons in hepatitis C antiviral therapy with direct-acting antivirals. *Hepatology.* 2013;58:428–38.
60. Afdhal N, Zeuzem S, Kwo P, et al. Ledipasvir and sofosbuvir for untreated HCV genotype 1 infection. *N Engl J Med.* 2014;370:1889–98.
61. Cornberg M, Honer zu Siederdisen C, Maasoumy B, Manns MP. New direct-acting antiviral agents for the treatment of chronic hepatitis C in 2014. *Internist (Berl)* 2014;55:390–400.
62. Druyts E, Thorlund K, Wu P, et al. Efficacy and safety of pegylated interferon alpha-2a or alpha-2b plus ribavirin for the treatment of chronic hepatitis C in children and adolescents: a systematic review and meta-analysis. *Clin Infect Dis.* 2013;56:961–7.
63. Sokal EM, Bourgois A, Stephenne X, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection in children and adolescents. *J Hepatol.* 2010;52:827–31.
64. Wirth S, Ribes-Koninckx C, Calzado MA, et al. High sustained virologic response rates in children with chronic hepatitis C receiving peginterferon alfa-2b plus ribavirin. *J Hepatol.* 2010;52:501–7.
65. Clemente MG, Antonucci R, Sotgiu G, et al. Present and future management of viral hepatitis B and C in children. *Clin Res Hepatol Gastroenterol.* 2020;44:801–9.
66. Karnsakul W, Schwarz KB. Management of Hepatitis C Infection in children in the era of direct-acting antiviral agents. *J Viral Hepat.* 2019;26:1034–9.
67. Balistreri WF, Murray KF, Rosenthal P, et al. The safety and effectiveness of ledipasvir-sofosbuvir in adolescents 12–17 years old with hepatitis C virus genotype 1 infection. *Hepatology.* 2017;66:371–8.
68. Schwarz KB, Rosenthal P, Murray KF, et al. Ledipasvir-Sofosbuvir for 12 weeks in children 3 to <6 years old with chronic hepatitis C. *Hepatology.* 2020;71:422–30.
69. Murray KF, Balistreri WF, Bansal S, et al. Safety and efficacy of Ledipasvir-Sofosbuvir with or without ribavirin for chronic hepatitis C in children ages 6–11. *Hepatology.* 2018;68:2158–66.
70. Wirth S, Rosenthal P, Gonzalez-Peralta RP, et al. Sofosbuvir and ribavirin in adolescents 12–17 years old with hepatitis C virus genotype 2 or 3 infection. *Hepatology.* 2017;66:1102–10.
71. Rosenthal P, Schwarz KB, Gonzalez-Peralta RP, et al. Sofosbuvir and ribavirin therapy for children aged 3 to <12 years with hepatitis C virus genotype 2 or 3 infection. *Hepatology.* 2020;71:31–43.
72. Jonas MM, Squires RH, Rhee SM, et al. Pharmacokinetics, safety, and efficacy of Glecaprevir/Pibrentasvir in adolescents with chronic hepatitis C virus: part 1 of the DORA study. *Hepatology.* 2020;71:456–62.