

## Chapter 7

# Special Supplement – List of Reported Interspecific Hybrids Within the Genus *Nicotiana*



Several investigators compiled comprehensive lists of *Nicotiana* hybrids. The earliest modern inventory was that by East (1928) who listed 65 interspecific combinations. It was followed by the list published by Kostoff (1943) who reported on 181 hybrids known to him. The last complete lists of interspecific *Nicotiana* hybrids with references were published by Goodspeed (1945, 1954). In the latter publication, the references to particular hybrids were scattered among the entries of the book's subject index. Together with several new combinations added a few years later (Goodspeed and Thompson, 1959) the number of documented interspecific hybrids stood at 243. The numbers of *Nicotiana* hybrids that were communicated later (Smith, 1968, 300 hybrids) and Apparao and Ramavarma (1974, 340 hybrids) were not supported by references to particular hybrid combinations. For the lack of more accurate data those estimations, although long outdated, have been non-the-less frequently quoted up to the present time. As more than 70 years have elapsed since Goodspeed's last inventory, the author of this volume thought it advisable to prepare a long-overdue update. Although the list below obviously exceeds the topical delimitations of this volume, annexing this updated inventory to a regular monograph on a closely related subject seems to be a good practical solution since publishing self-contained lists of this kind is not an accepted custom. The list also serves as an extension to the last column of Table 7.1.

In all probability, the list is not complete. It drew mostly on reports that were accessible through the web. The author faced the same problem as he did while working on other parts of this review: difficulty in searching materials published in remote parts of the globe or in exotic languages or/and scripts. Conversely, many of the listed hybrids have been repeatedly reported in many publications. Regardless of reporting frequency, only the first reports, a maximum of seven, were cited for each hybrid combination.

In earlier reports, hybrids that failed to reach the blooming stage did not qualify as successful hybrids and usually were listed separately. The list in this volume was extended by adding two, possibly debatable, categories:

**Table 7.1** Interspecific hybrids within the genus *Nicotiana* reported by the year 2023

No.	Hybrid combination	Species 1	Species 2	Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
1.	<i>N. alata</i>	<i>N. bonariensis</i>		F, M	Avery (1938), Kostoff (1940), Goodspeed (1945)
2.	<i>N. alata</i>	<i>N. langsdorffii</i>		F, M	Focke (1881), East and Hayes (1912), East (1916)
3.	<i>N. alata</i>	<i>N. longiflora</i>		F, M	East and Hayes (1912), Avery (1938), Goodspeed (1945), Náf (1958)
4.	<i>N. alata</i>	<i>N. mutabilis</i>		F, M	Stehmann et al. (2002)
5.	<i>N. alata</i>	<i>N. plumbaginifolia</i>		F, M	Goodspeed (1945), Takenaka (1950), Engvild (1950)
6.	<i>N. alata</i>	<i>N. sanderae</i>		F, M	D. Kostoff after Bradley and Goodspeed (1943)
7.	<i>N. alata</i>	<i>N. forgetiana</i>		F, M	East and Hayes (1912), Sorrentino et al. (2006)
8.	<i>N. alata</i>	<i>N. rastroensis</i> <sup>1</sup>		M	Descorbeth (2004), Descorbeth and McClure (2005)
9.	<i>N. alata</i>	<i>N. tabacum</i>		M	East and Hayes (1912), Kostoff (1930), Kostoff (1943), Ternovsky (1962)
10.	<i>N. alata</i>	<i>N. glauca</i>		M	Kostoff (1943), Goodspeed (1945), Takenaka (1953a)
11.	<i>N. alata</i>	<i>N. noctiflora</i>		M	Kostoff (1940), Goodspeed (1945), Náf (1958)
12.	<i>N. alata</i>	<i>N. knightiana</i>		M	Takenaka (1965), Takanashi and Marubashi (2017)
13.	<i>N. alata</i>	<i>N. paniculata</i>		F, M	Kostoff (1943), Goodspeed (1945), Náf (1958), Takenaka (1961)
14.	<i>N. alata</i>	<i>N. quadrivalvis</i>		M	Kostoff (1943), Goodspeed (1945)
15.	<i>N. alata</i>	<i>N. rustica</i>		F, M	Whitaker (1934), Zhukov (1939), Goodspeed (1945), Kehr (1951), Kehr and Smith (1954), Wolf (1965)
16.	<i>N. alata</i>	<i>N. amplexicaulis</i>		M	Gopinath et al. (1965, 1970)
	<i>N. alata</i>	<i>N. debneyi</i>		F	Zenkeler et al. (1981), Bartkowiak (1987)
18.	<i>N. alata</i>	<i>N. gossei</i>		F, M	Kostoff (1940), Goodspeed (1945), Takenaka (1962b), Gopinath et al. (1965, 1970)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
19.	<i>N. alata</i>	<i>N. maritima</i>	M	Kostoff (1943), Goodspeed (1945)
20.	<i>N. alata</i>	<i>N. suaveolens</i>	M	Kostoff (1943), Goodspeed (1945)
21.	<i>N. alata</i>	<i>N. velutina</i>	M	Bradley and Goodspeed (1943)
22.	<i>N. alata</i>	<i>N. eastii</i>	M	Kostoff (1940), Goodspeed (1945)
23.	<i>N. alata</i>	<i>N. tomentosiformis</i>	M	Goodspeed, 1945
24.	<i>N. alata</i>	<i>N. sylvestris</i>	M, asexual	Berbeć et al. (1976), Fluhr, 1983
25.	<i>N. alata</i>	<i>N. undulata</i>	M	Takanashi and Marubashi (2017)
26.	<i>N. bonariensis</i>	<i>N. langsdorffii</i>	F, M	Muntzing, 1935, Avery (1938), Kostoff (1943), Goodspeed (1945), Näf (1958)
27.	<i>N. bonariensis</i>	<i>N. longiflora</i>	F, M	Avery (1938), Goodspeed (1945, 1954)
28.	<i>N. bonariensis</i>	<i>N. mutabilis</i>	M	Stehmann et al. 2002
29.	<i>N. bonariensis</i>	<i>N. plumbaginifolia</i>	F	Kostoff (1943), Goodspeed (1945), Näf (1958)
30.	<i>N. bonariensis</i>	<i>N. rastroensis</i> <sup>1</sup>	F, M	Lee et al. (2008)
31.	<i>N. bonariensis</i>	<i>N. forgetiana</i>	F	Butenko & Luneva, 1966
32.	<i>N. bonariensis</i>	<i>N. sanderae</i>	F, M	Kostoff (1943)
33.	<i>N. bonariensis</i>	<i>N. tabacum</i>	M?	Stavely 1979, Busconi et al. (2010)
34.	<i>N. bonariensis</i>	<i>N. glauca</i>	F	Williams and Pandey 1975 <sup>5</sup>
35.	<i>N. bonariensis</i>	<i>N. quadrivalvis</i>	M (l)	Näf (1958)
36.	<i>N. bonariensis</i>	<i>N. rustica</i>	M	Näf (1958)
37.	<i>N. bonariensis</i>	<i>N. suaveolens</i>	M	Näf (1958)
38.	<i>N. forgetiana</i>	<i>N. langsdorffii</i>	F	East and Hayes (1912), White (1914)
39.	<i>N. forgetiana</i>	<i>N. mutabilis</i>	M	Stehmann et al. (2002)
40.	<i>N. forgetiana</i>	<i>N. plumbaginifolia</i>	M	Näf (1958), Takenaka (1965)
41.	<i>N. forgetiana</i>	<i>N. sanderae</i>	F, M	Sorrentino et al. (2006)
42.	<i>N. forgetiana</i>	<i>N. rastroensis</i> <sup>1</sup>	F, M	Lee et al. (2008)
43.	<i>N. forgetiana</i>	<i>N. tabacum</i>	M	Näf (1958), Ahuja (1962)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
44.	<i>N. forgetiana</i>	<i>N. glauca</i>	F, M	Williams and Pandey (1975)
45.	<i>N. forgetiana</i>	<i>N. quadrivalvis</i>	M (l)	Näf (1958)
46.	<i>N. forgetiana</i>	<i>N. rustica</i>	M	Näf (1958)
47.	<i>N. langsdorffii</i>	<i>N. longiflora</i>	M	Goodspeed, 1945, Takenaka et al. (1956), Avery, 1962
48.	<i>N. langsdorffii</i>	<i>N. mutabilis</i>	M	Stehmann et al. (2002)
49.	<i>N. langsdorffii</i>	<i>N. plumbaginifolia</i>	F, M	Kostoff (1943), Goodspeed, 1945, 1954
50.	<i>N. langsdorffii</i>	<i>N. sanderae</i>	F, M	East, 1935 Smith (1937), Kostoff, 1943, Goodspeed (1945, 1954)
51.	<i>N. langsdorffii</i>	<i>N. rastroensis</i> <sup>1</sup>	F, M	Lee et al. (2008)
52.	<i>N. langsdorffii</i>	<i>N. tabacum</i>	M	East & Hayes, 1912, East (1935), Hu, 1956, Takenaka (1962b), Burk, 1972
53.	<i>N. langsdorffii</i>	<i>N. glauca</i>	F, M	premendelian, East (1928, 1935), Whitaker (1934), Goodspeed (1954)
54.	<i>N. langsdorffii</i>	<i>N. noctiflora</i>	M	Kostoff (1943), Goodspeed (1945, 1954)
55.	<i>N. langsdorffii</i>	<i>knightiana</i>	M	Takanashi and Marubashi (2017)
56.	<i>N. langsdorffii</i>	<i>N. paniculata</i>	M	premendelian, East and Hayes (1912), Whitaker (1934), Kostoff (1939b), Kehr and Smith, 1954, Takenaka (1962a), Takenaka and Yoneda (1964)
57.	<i>N. langsdorffii</i>	<i>N. miersii</i>	M	Kostoff (1940, 1943), Goodspeed (1945, 1954), Kehr (1951), Kehr and Smith (1954), Wolf (1965)
58.	<i>N. langsdorffii</i>	<i>N. quadrivalvis</i>	F	East and Hayes (1912), Kostoff, 1943, Näf (1958), Tezuka (2012)
59.	<i>N. langsdorffii</i>	<i>N. repanda</i>	M	Schweppenhauser et al. (1963)
60.	<i>N. langsdorffii</i>	<i>N. suaveolens</i>	M	premendelian, East (1928), Kostoff (1939b, 1943), Smith (1958), Smith and Abashian (1963), Wolf (1965)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
61.	<i>N. langsdorffii</i>	<i>N. rustica</i>	M	East and Hayes (1912), Kostoff (1930), Whitaker (1934), Kehr (1951), Kehr and Smith (1954), Wolf (1965)
62.	<i>N. langsdorffii</i>	<i>N. otophora</i>	?	Berbeć et al. (1976)
63.	<i>N. langsdorffii</i>	<i>N. glutinosa</i>	M	East (1935)
64.	<i>N. langsdorffii</i>	<i>N. undulata</i>	M	Goodspeed (1945), Takanashi and Marubashi (2017)
65.	<i>N. langsdorffii</i>	<i>N. obtusifolia</i>	F (l)	Christoff (1928)
66.	<i>N. longiflora</i>	<i>N. sanderae</i>	F, M	Malloch and Malloch (1924), Brieger (1929), Kostoff (1943), Takenaka (1955)
67.	<i>N. longiflora</i>	<i>N. plumbaginifolia</i>	F, M	Avery (1938), Kostoff (1943), Goodspeed (1945), Murthy et al. (1998)
68.	<i>N. longiflora</i>	<i>N. tabacum</i>	M, F	Malloch (1924), Kostoff (1943), Takenaka (1962d), Sievert (1972) <sup>3</sup> , Sarala et al. (2023)
69.	<i>N. longiflora</i>	<i>N. glauca</i>	F, M	East (1935), Kostoff (1943), Kehr (1951), Takenaka (1952a, 1953a, 1965), Murthy et al. (1998)
70.	<i>N. longiflora</i>	<i>N. paniculata</i>	F	Christoff (1928), Takenaka (1962a)
71.	<i>N. longiflora</i>	<i>N. quadrivalvis</i>	M (l)	Tezuka (2012)
72.	<i>N. longiflora</i>	<i>N. repanda</i>	M	Schweppenhauser et al. (1963), Davis et al. (1988)
73.	<i>N. longiflora</i>	<i>N. rustica</i>	M	McCray (1933)
74.	<i>N. longiflora</i>	<i>N. amplexicaulis</i>	not indicated	Gopinath et al. (1970)
75.	<i>N. longiflora</i>	<i>N. debneyi</i>	M	Naf (1958), Ahuja (1968)
76.	<i>N. longiflora</i>	<i>N. excelsior</i>	M	Takenaka (1965), Murthy et al. (1998)
77.	<i>N. longiflora</i>	<i>N. gossei</i>	M	Goodspeed (1945), Takenaka et al. (1956), Takenaka (1962b), Takenaka and Yoneda (1964)
78.	<i>N. longiflora</i>	<i>N. ingulba</i>	M	Kostoff (1943)
79.	<i>N. longiflora</i>	<i>N. megalosiphon</i>	M	Goodspeed (1945, 1954)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
80.	<i>N. longiflora</i>	<i>N. suaveolens</i>	M	East (1935), Kostoff (1943), Goodspeed (1945, 1954), Takenaka (1952b), Takenaka and Yoneda (1964)
81.	<i>N. longiflora</i>	<i>N. eastii</i>	M	Kostoff (1940), Goodspeed (1945)
82.	<i>N. longiflora</i>	<i>N. sylvestris</i>	F	Savelli (1926)
83.	<i>N. longiflora</i>	<i>N. glutinosa</i>	F	Murthy et al. (1998)
84.	<i>N. longiflora</i>	<i>N. undulata</i>	M	Goodspeed (1945)
85.	<i>N. mutabilis</i>	<i>N. rastroensis</i> <sup>1</sup>	M	Lee et al. (2008)
86.	<i>N. plumbaginifolia</i>	<i>N. sanderae</i>	F, M	Goodspeed (1945), Kehr and Smith (1952), Takenaka (1953c, 1955)
87.	<i>N. plumbaginifolia</i>	<i>N. rastroensis</i> <sup>1</sup>	F, M	Lee et al. (2008)
88.	<i>N. plumbaginifolia</i>	<i>N. tabacum</i>	F, M	Pal and Nath (1936), Kostoff (1943), Kincaid (1949), Moav and Cameron (1960), Naf (1958), Burk (1960)
89.	<i>N. plumbaginifolia</i>	<i>N. glauca</i>	F, M	McCray (1932), Kostoff (1943), Ramanujam and Joshi (1942), Kehr and Smith (1952), Takenaka (1951a, 1953a)
90.	<i>N. plumbaginifolia</i>	<i>N. paniculata</i>	M	Christoff (1928) (l), Lilienfeld (1953), Naf (1958), Takenaka (1962a),
91.	<i>N. plumbaginifolia</i>	<i>N. quadrivalvis</i>	F	Christoff (1928) (l), Tezuka (2012) (l)
92.	<i>N. plumbaginifolia</i>	<i>N. rustica</i>	M	Kostoff (1943), Takenaka and Yoneda (1964)
93.	<i>N. plumbaginifolia</i>	<i>N. debneyi</i>	M	Kehr (1951), Kehr and Smith (1952, 1954), Smith and Abashian (1963)
94.	<i>N. plumbaginifolia</i>	<i>N. excelsior</i>	M	Sarala et al. (2008), Murthy and Rao (2008)
95.	<i>N. plumbaginifolia</i>	<i>N. gossei</i>	M	Takenaka et al. (1956), Takenaka (1962b), Murthy and Rama Prasad (2000)
96.	<i>N. plumbaginifolia</i>	<i>N. maritima</i>	F, M	Goodspeed (1945)
97.	<i>N. plumbaginifolia</i>	<i>N. megalosiphon</i>	M	Goodspeed (1945)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
98.	<i>N. plumbaginifolia</i>	<i>N. suaveolens</i>	M	Goodspeed ( <a href="#">1945</a> ), Kehr and Smith ( <a href="#">1952</a> ), Takenaka ( <a href="#">1952b</a> , <a href="#">1962b</a> ), Näf ( <a href="#">1958</a> )
99.	<i>N. plumbaginifolia</i>	<i>N. umbratica</i>	F, M	Gangadevi et al. ( <a href="#">1987</a> )
100.	<i>N. plumbaginifolia</i>	<i>N. eastii</i>	M	Goodspeed ( <a href="#">1945</a> )
101.	<i>N. plumbaginifolia</i>	<i>N. sylvestris</i>	asexual	Lin and Chen ( <a href="#">1990</a> ), Cseplo et al. ( <a href="#">1983</a> )
102.	<i>N. plumbaginifolia</i>	<i>N. otophora</i>	asexual	Lin and Chen ( <a href="#">1990</a> )
103.	<i>N. plumbaginifolia</i>	<i>N. obtusifolia</i>	M	Murthy et al. ( <a href="#">1998</a> )
104.	<i>N. sanderae</i>	<i>N. tabacum</i>	F, M	Christoff ( <a href="#">1928</a> ), Kostoff ( <a href="#">1930</a> ), East ( <a href="#">1935</a> ), Kehr and Smith ( <a href="#">1954</a> ), Ternovsky ( <a href="#">1962</a> )
105.	<i>N. sanderae</i>	<i>N. glauca</i>	M	Kostoff ( <a href="#">1930</a> ), Brieger and Forster ( <a href="#">1942</a> ), Goodspeed ( <a href="#">1945</a> )
106.	<i>N. sanderae</i>	<i>N. noctiflora</i>	M	Kostoff ( <a href="#">1935</a> , <a href="#">1940</a> ), Goodspeed ( <a href="#">1945</a> )
107.	<i>N. sanderae</i>	<i>N. paniculata</i>	F, M	Brieger ( <a href="#">1929</a> ), East ( <a href="#">1935</a> ), Goodspeed ( <a href="#">1954</a> ), Kehr ( <a href="#">1951</a> ), Kehr and Smith ( <a href="#">1954</a> )
108.	<i>N. sanderae</i>	<i>N. repanda</i>	M	Takenaka ( <a href="#">1953c</a> , <a href="#">1955</a> )
109.	<i>N. sanderae</i>	<i>N. miersii</i>	M	Näf ( <a href="#">1958</a> )
110.	<i>N. sanderae</i>	<i>N. rustica</i>	M	Kostoff ( <a href="#">1930</a> ), Whitaker ( <a href="#">1934</a> ), Goodspeed ( <a href="#">1945</a> ), Kehr ( <a href="#">1951</a> ), Kehr and Smith ( <a href="#">1954</a> )
111.	<i>N. sanderae</i>	<i>N. amplexicaulis</i>	M	Gopinath et al. ( <a href="#">1965</a> )
112.	<i>N. sanderae</i>	<i>N. debneyi</i>	asexual	Patel et al. ( <a href="#">2011</a> )
113.	<i>N. sanderae</i>	<i>N. gossei</i>	M	Goodspeed ( <a href="#">1945</a> )
114.	<i>N. sanderae</i>	<i>N. maritima</i>	M	Goodspeed ( <a href="#">1945</a> , <a href="#">1954</a> )
115.	<i>N. sanderae</i>	<i>N. suaveolens</i>	M	Goodspeed ( <a href="#">1945</a> ), Takenaka ( <a href="#">1955</a> )
116.	<i>N. sanderae</i>	<i>N. eastii</i>	M	Kostoff ( <a href="#">1939b</a> , <a href="#">1940</a> )
117.	<i>N. sanderae</i>	<i>N. sylvestris</i>	M	Kostoff ( <a href="#">1943</a> )
118.	<i>N. sanderae</i>	<i>N. quadrivalvis</i>	M	Whitaker ( <a href="#">1934</a> ), Kostoff ( <a href="#">1943</a> ), Kehr ( <a href="#">1951</a> ), Kehr and Smith ( <a href="#">1954</a> ), Wolf ( <a href="#">1965</a> )

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
119.	<i>N. tabacum</i>	<i>N. glauca</i>	F, M	premendelian, East (1935)
120.	<i>N. tabacum</i>	<i>N. noctiflora</i>	F	Palakarcheva (1975), Dorossiev et al. (1978)
121.	<i>N. tabacum</i>	<i>N. petunioides</i>	F	Gisquet et al. (1940)
122.	<i>N. tabacum</i>	<i>N. benavidesii</i>	F	Goodspeed (1945)
123.	<i>N. tabacum</i>	<i>N. cordifolia</i>	M	Durbin and Uchytíl (1977), Burk and Durbin (1978)
124.	<i>N. tabacum</i>	<i>N. knightiana</i>	F, M	Goodspeed (1945), Tanaka (1961), Takenaka (1962d), Berbeć et al. (1982)
125.	<i>N. tabacum</i>	<i>N. paniculata</i>	F, M	premendelian (East (1928)), Kostoff (1943), Goodspeed (1954), Takenaka (1962d)
126.	<i>N. tabacum</i>	<i>N. raimondii</i>	F, M	Kostoff (1943), Goodspeed (1954), Burk et al. (1982), Berbeć (1988)
127.	<i>N. tabacum</i>	<i>N. solanifolia</i>	F	Goodspeed (1945), Takenaka (1956a), b
128.	<i>N. tabacum</i>	<i>N. acuminata</i>	F	Gentcheff (Kostoff (1943)), Palakarcheva (1981), Iwai et al. (1986)
129.	<i>N. tabacum</i>	<i>N. pauciflora</i>	F	Gentscheff (1931), Kostoff (1943), Goodspeed (1945)
130.	<i>N. tabacum</i>	<i>N. clevelandii</i>	M	Kehr and Smith (1952), Kaul (1988)
131.	<i>N. tabacum</i>	<i>N. quadrivalvis</i>	F, M	premendelian (East (1928)), East and Hayes (1912), Kostoff (1943)
132.	<i>N. tabacum</i>	<i>N. nesophila</i>	M	Reed and Collins (1978, 1980), Huesing et al. (1989)
133.	<i>N. tabacum</i>	<i>N. nudicaulis</i>	F, M	Gentscheff (1931), Kostoff (1943)
134.	<i>N. tabacum</i>	<i>N. repanda</i>	F, M	Foster (1943), Kincaid (1949), Pittarelli and Stavely (1975), Nagao (1982 <sup>6</sup> )
135.	<i>N. tabacum</i>	<i>N. stocktonii</i>	M,	Wong (1975), Reed and Collins (1978, 1980)
136.	<i>N. tabacum</i>	<i>N. rustica</i>	F, M	premendelian, East and Hayes (1912), Eghis (1927), Savelli (1927), East (1935)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
137.	<i>N. tabacum</i>	<i>N. africana</i>	F, M	Gerstel et al. (1979), Nikova and Zagorska (1990)
138.	<i>N. tabacum</i>	<i>N. amplexicaulis</i>	M, F	Wark (1970), Berbeć and Doroszewska (1981), Larkina (1980), Sarala et al. (2023)
139.	<i>N. tabacum</i>	<i>N. benthamiana</i>	F, M	Subhashini et al. (1986), DeVerna et al. (1987), Nikova et al. (1991), Zaitlin and Mundell ((2006)), Mihaylova-Kroumová et al. (2020), Sarala et al. (2023)
140.	<i>N. tabacum</i>	<i>N. cavicola</i>	M	Nikova et al. (2006)
141.	<i>N. tabacum</i>	<i>N. debneyi</i>	M	Kostoff (1943), Goodspeed (1954), Takenaka (1956a)
142.	<i>N. tabacum</i>	<i>N. excelsior</i>	M	Wark (1970), Nikova (1986), Murthy et al. (2014)
143.	<i>N. tabacum</i>	<i>N. exigua</i>	M	Wark (1970), Kobus (1971)
144.	<i>N. tabacum</i>	<i>N. fragrans</i>	M	Durbin and Uchytil (1977)
145.	<i>N. tabacum</i>	<i>N. goodspeedii</i>	M	Butenko et al. (1970), Wark (1970)
146.	<i>N. tabacum</i>	<i>N. gossei</i>	M, F	Valleau (1952), Takenaka (1962d), Sievert (1972) <sup>4</sup> , Adachi and Inoue (1995), Murthy et al. (2014), Sarala et al. (2023)
147.	<i>N. tabacum</i>	<i>N. hesperis</i>	M	Kubo (1985)
148.	<i>N. tabacum</i>	<i>N. ingulba</i>	M	Butenko et al. (1970), Nikova et al. (1998)
149.	<i>N. tabacum</i>	<i>N. maritima</i>	M	Wark (1970), Palakarcheva (1975)
150.	<i>N. tabacum</i>	<i>N. megalosiphon</i>	M	Clayton (1950), Takenaka (1962d), Hranov (1970), Sarala et al. (2023)
151.	<i>N. tabacum</i>	<i>N. occidentalis</i>	M	Butenko et al. (1970), Ternovsky et al. (1972), Wong (1975)
152.	<i>N. tabacum</i>	<i>N. rosulata</i>	F	Ternovsky et al. (1976), Ternovsky and Larkina (1978), Larkina (2015, 2017)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
153.	<i>N. tabacum</i>	<i>N. rotundifolia</i>	asexual	Ilcheva et al. (2001)
154.	<i>N. tabacum</i>	<i>N. simulans</i>	M	Kubo (1985), Tatemichi (1990), Ma et al. (2020)
155.	<i>N. tabacum</i>	<i>N. suaveolens</i>	F, M	pre mendelian (East (1928)), Kostoff (1943)
156.	<i>N. tabacum</i>	<i>N. umbratica</i>	M	Murthy et al. (2014), Sarala et al. (2023)
157.	<i>N. tabacum</i>	<i>N. velutina</i>	M	Wark (1970), Ma et al. (2020)
158.	<i>N. tabacum</i>	<i>N. wuttkei</i>	M	Laskowska and Berbeć (2012)
159.	<i>N. tabacum</i>	<i>N. eastii</i>	M	Chaplin & Mann, 1961
160.	<i>N. tabacum</i>	<i>N. sylvestris</i>	F, M	East and Hayes (1912), Bellair (1913), Malinowski (1916), Sachs-Skalińska (1917)
161.	<i>N. tabacum</i>	<i>N. kawakamii</i>		Ohashi (1985)
162.	<i>N. tabacum</i>	<i>N. otophora</i>	F	Goodspeed (1945), Ar-Rushdi (1955), Takenaka (1962)
163.	<i>N. tabacum</i>	<i>N. setchellii</i>	F, M	Greenleaf (1941), Goodspeed (1945, 1954)
164.	<i>N. tabacum</i>	<i>N. tomentosa</i>	F, M	Goodspeed and Clausen (1928), McCray (1932), Kostoff (1943)
165.	<i>N. tabacum</i>	<i>N. tomentosiformis</i>	F, M	Brieger (1928), Breisser (1934), Lehmann (1936), Kostoff (1943)
166.	<i>N. tabacum</i>	<i>N. obtusifolia</i>	F, M	Takenaka (1951b, 1956a), Chung et al. (1988), Choi and Lee (1991), Liu and Marubashi (2014)
167.	<i>N. tabacum</i>	<i>N. palmeri</i>	F	Goodspeed (1945), Berbeć et al. (1982)
168.	<i>N. tabacum</i>	<i>N. arentsii</i>	asexual	DeVerna (1984)
169.	<i>N. tabacum</i>	<i>N. glutinosa</i>	F, M	pre mendelian (East (1928)), first report by Koelreuter (Mayr (1986)), East (1935),
170.	<i>N. tabacum</i>	<i>N. undulata</i>	M	Kehr and Smith (1952), Takenaka (after Goodspeed and Thompson (1959))

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
171.	<i>N. glauca</i>	<i>N. noctiflora</i>	F, M	Goodspeed (1934), Kostoff (1943), Goodspeed (1945)
172.	<i>N. glauca</i>	<i>N. benavidesii</i>	F	Goodspeed (1945)
173.	<i>N. glauca</i>	<i>N. paniculata</i>	F, M	East (1935), Lehmann (1936), Kostoff (1943), Poddubnaya-Arnoldi and Ludkina (1945), Kehr and Smith (1954)
174.	<i>N. glauca</i>	<i>N. raimondii</i>	M	Goodspeed (1954)
175.	<i>N. glauca</i>	<i>N. solanifolia</i>	M	Kostoff (1940), Goodspeed (1945)
176.	<i>N. glauca</i>	<i>N. attenuata</i>	M	Gentscheff (1931)
177.	<i>N. glauca</i>	<i>N. miersii</i>	M	Kostoff (1943)
178.	<i>N. glauca</i>	<i>N. clevelandii</i>	F	Kehr and Smith (1952)
179.	<i>N. glauca</i>	<i>N. quadrivalvis</i>	M	McCray (1932), East (1935) Kostoff (1943), Goodspeed (1945), Kehr and Smith (1952), Meiselman et al. (1961)
180.	<i>N. glauca</i>	<i>N. nudicaulis</i>	M	Goodspeed (1934, 1945)
181.	<i>N. glauca</i>	<i>N. rustica</i>	M	East (1935), Bolsunov (1937), Modilevsky (1937), Kostoff (1939c), Goodspeed (1945), Ilin (1948)
182.	<i>N. glauca</i>	<i>N. debneyi</i>	F, M	Kehr (1951), Lautz (1957), Smith and Abashian (1963)
183.	<i>N. glauca</i>	<i>N. goodspeedii</i>	F	Horton (1981)
184.	<i>N. glauca</i>	<i>N. gossei</i>	M, F	Akada and Hirai (1986) (somatic), Murthy et al. (1998), Murthy and Subbarao (2004), Sarala et al. (2008)
185.	<i>N. glauca</i>	<i>N. megalosiphon</i>	F	Goodspeed (1945)
186.	<i>N. glauca</i>	<i>N. simulans</i>	F	Horton (1981)
187.	<i>N. glauca</i>	<i>N. suaveolens</i>	F, M	East (1935), Goodspeed (1945), Naf (1958), Smith & Abashian, 1963 (amphidiploid synthetic species <i>N. x flindersiensis</i> )

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
188.	<i>N. glauca</i>	<i>N. umbratica</i>	M	Gangadevi et al. ((1987))
189.	<i>N. glauca</i>	<i>N. velutina</i>	F	Jeanes (1999)
190.	<i>N. glauca</i>	<i>N. eastii</i>	M	Kostoff (1943)
191.	<i>N. glauca</i>	<i>N. sylvestris</i>	F	Ternovsky (1962), Takenaka (1965)
192.	<i>N. glauca</i>	<i>N. otophora</i>	F	Takenaka (1953b)
193.	<i>N. glauca</i>	<i>N. tomentosa</i>	F, M	Kostoff (1932), East (1935), Lehmann (1936), Smith (1942)
194.	<i>N. glauca</i>	<i>N. tomentosiformis</i>	F, M	Kostoff (1932), Lehmann (1936), Goodspeed (1945)
195.	<i>N. glauca</i>	<i>N. obtusifolia</i>	M	Murthy et al. (1998)
196.	<i>N. glauca</i>	<i>N. glutinosa</i>	F, M	Kostoff (1930), East (1935), Goodspeed (1945)
197.	<i>N. glauca</i>	<i>N. thyrsiflora</i>	M	Goodspeed (1945, 1954)
198.	<i>N. noctiflora</i>	<i>N. petunioides</i>	F	Goodspeed and Thompson (1959)
199.	<i>N. noctiflora</i>	<i>N. paniculata</i>	M	Goodspeed (1934), Kostoff (1943), Goodspeed (1945)
200.	<i>N. noctiflora</i>	<i>N. rustica</i>	M	Kostoff (1943), Náf (1958), Takenaka and Yoneda (1964), Kehr and Smith (1954), Wolf (1965), Busconi et al. (2010)
201.	<i>N. petunioides</i>	<i>N. corymbosa</i>	M	Goodspeed and Thompson (1959)
202.	<i>N. petunioides</i>	<i>N. miersii</i>	M	Goodspeed (1945)
203.	<i>N. benavidesii</i>	<i>N. paniculata</i>	M	Goodspeed (1945, 1954)
204.	<i>N. benavidesii</i>	<i>N. raimondii</i>	F	Goodspeed (1945)
205.	<i>N. benavidesii</i>	<i>N. solanifolia</i>	F	Goodspeed (1945)
206.	<i>N. benavidesii</i>	<i>N. rustica</i>	M	Goodspeed (1945)
207.	<i>N. benavidesii</i>	<i>N. glutinosa</i>	F	Goodspeed (1945)
208.	<i>N. benavidesii</i>	<i>N. undulata</i>	M	Goodspeed (1945)
209.	<i>N. cordifolia</i>	<i>N. knightiana</i>	M	Goodspeed (1945, 1954)
210.	<i>N. cordifolia</i>	<i>N. raimondii</i>	F, M	Goodspeed (1945, 1954)
211.	<i>N. cordifolia</i>	<i>N. solanifolia</i>	F, M	Goodspeed (1954)
212.	<i>N. cordifolia</i>	<i>N. undulata</i>	M	Goodspeed (1945)
213.	<i>N. knightiana</i>	<i>N. raimondii</i>	M	Durbin and Uchytil (1977)
214.	<i>N. knightiana</i>	<i>N. solanifolia</i>	M	Durbin and Uchytil (1977)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (I) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
215.	<i>N. knightiana</i>	<i>N. rustica</i>	F	Takenaka (1958, 1959, 1962c)
216.	<i>N. knightiana</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1985)
217.	<i>N. knightiana</i>	<i>N. sylvestris</i>	asexual	Menczel et al. (1978)
218.	<i>N. paniculata</i>	<i>N. raimondii</i>	F, M	Kostoff (1940), Goodspeed (1945)
219.	<i>N. paniculata</i>	<i>N. solanifolia</i>	F, M	Goodspeed (1945), Durbin and Uchytíl (1977)
220.	<i>N. paniculata</i>	<i>N. miersii</i>	F, M	Kostoff (1943), Goodspeed (1945)
221.	<i>N. paniculata</i>	<i>N. pauciflora</i>	F	Goodspeed (1934)
222.	<i>N. paniculata</i>	<i>N. quadrivalvis</i>	F	premendelian (East (1928)), East and Hayes (1912)
223.	<i>N. paniculata</i>	<i>N. nudicaulis</i>	F (I)	Yamada et al. (1999) (invি- able type III)
224.	<i>N. paniculata</i>	<i>N. rustica</i>	F, M	premendelian (East (1928)), East and Hayes (1912)
225.	<i>N. paniculata</i>	<i>N. gossei</i>	F (I)	Yamada et al. (1999)
226.	<i>N. paniculata</i>	<i>N. suaveolens</i>	F	premendelian (East (1928))
227.	<i>N. paniculata</i>	<i>N. umbratica</i>	F	Subhashini and Gopinath (1974)
228.	<i>N. paniculata</i>	<i>N. velutina</i>	M	Bradley and Goodspeed (1943)
229.	<i>N. paniculata</i>	<i>N. otophora</i>	F	Takenaka (1953b)
230.	<i>N. paniculata</i>	<i>N. tomentosa</i>	F	Lehmann (1936)
231.	<i>N. paniculata</i>	<i>N. tomentosiformis</i>	F, M (I)	Lehmann (1936)
232.	<i>N. paniculata</i>	<i>N. glutinosa</i>	F, M	premendelian (East (1928))
233.	<i>N. paniculata</i>	<i>N. undulata</i>	F, M	Fatalizade (1939), Kostoff (1943), Ternovsky (1962)
234.	<i>N. raimondii</i>	<i>N. solanifolia</i>	F, M	Kostoff (1940, 1943), Goodspeed (1945)
235.	<i>N. raimondii</i>	<i>N. undulata</i>	M	Goodspeed (1945)
236.	<i>N. solanifolia</i>	<i>N. quadrivalvis</i>	M	Goodspeed (1934), Kostoff (1943), Goodspeed (1945)
237.	<i>N. solanifolia</i>	<i>N. rustica</i>	M	Goodspeed (1934, 1945)
238.	<i>N. solanifolia</i>	<i>N. suaveolens</i>	M	East (1928), Kostoff (1943)
239.	<i>N. solanifolia</i>	<i>N. otophora</i>		Berbeć et al. (1976)
240.	<i>N. solanifolia</i>	<i>N. undulata</i>	M	Ternovsky (1962)
241.	<i>N. acuminata</i>	<i>N. corymbosa</i>	M	Goodspeed and Thompson (1959)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
242.	<i>N. acuminata</i>	<i>N. pauciflora</i>	F, M	Kostoff (1943), Goodspeed and Thompson (1959), Takenaka (1965)
243.	<i>N. acuminata</i>	<i>N. undulata</i>	M	Gray (1978)
244.	<i>N. attenuata</i>	<i>N. miersii</i>	F, M	Kostoff (1943), Goodspeed and Thompson (1959), Pearse et al. (2006) Krügel (2010)
245.	<i>N. attenuata</i>	<i>N. clevelandii</i>	M	Goodspeed (1945)
246.	<i>N. attenuata</i>	<i>N. quadrivalvis</i>	M	Goodspeed (1945)
247.	<i>N. attenuata</i>	<i>N. rustica</i>	M	pre mendelian (East (1928)), Patel (1960). Goodspeed (1945), Tezuka (2012) (lethal)
248.	<i>N. attenuata</i>	<i>N. maritima</i>	M	Goodspeed (1945)
249.	<i>N. attenuata</i>	<i>N. obtusifolia</i>	M	Anssour et al. (2009), Navarro-Quezada et al. (2013), Navarro-Quezada (2014)
250.	<i>N. attenuata</i>	<i>N. undulata</i>	M	Gentscheff (1931), Kostoff (1943), Goodspeed (1954)
251.	<i>N. corymbosa</i>	<i>N. linearis</i>	F	Goodspeed and Thompson (1959)
252.	<i>N. miersii</i>	<i>N. clevelandii</i>	F	Goodspeed (1945)
253.	<i>N. miersii</i>	<i>N. quadrivalvis</i>	F	Kostoff (1940, 1943), Goodspeed (1945)
254.	<i>N. miersii</i>	<i>N. rustica</i>	M	East (1928), Goodspeed (1945), Patel (1960)
255.	<i>N. miersii</i>	<i>N. megalosiphon</i>	F	Kostoff (1940), Goodspeed (1945)
256.	<i>N. miersii</i>	<i>N. suaveolens</i>	F	Goodspeed (1934), Kostoff (1940), Goodspeed (1945)
257.	<i>N. pauciflora</i>	<i>N. rustica</i>	M	Goodspeed (1945)
258.	<i>N. clevelandii</i>	<i>N. quadrivalvis</i>	F, M	Masuta et al. (1993), Kiraly and Schoeltz (1995)
259.	<i>N. clevelandii</i>	<i>N. nudicaulis</i>	F	Goodspeed (1945)
260.	<i>N. clevelandii</i>	<i>N. debneyi</i>	M	Kehr and Smith (1952), Lautz (1957), Smith and Abashian (1963), Smith et al. (1970)
261.	<i>N. clevelandii</i>	<i>N. exigua</i>	F	Kostoff (1943), Goodspeed (1945)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
262.	<i>N. clevelandii</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1987)
263.	<i>N. clevelandii</i>	<i>N. glutinosa</i>	F, M	Kehr and Smith (1952), Lautz (1957), Smith and Abashian (1963), Smith et al. (1970)
264.	<i>N. quadrivalvis</i>	<i>N. nudicaulis</i>	F, M	Goodspeed (1945)
265.	<i>N. quadrivalvis</i>	<i>N. rustica</i>	F, M	premendelian (East (1928)), Maryanovich (1939), Fardy and Hitier (1945)
266.	<i>N. quadrivalvis</i>	<i>N. amplexicaulis</i>	F	Gopinath et al. (1965, 1970)
267.	<i>N. quadrivalvis</i>	<i>N. benthamiana</i>	F	Kostoff (1943), Goodspeed (1945), Apparao and Ramavarma (1974)
268.	<i>N. quadrivalvis</i>	<i>N. debneyi</i>	F	Kehr and Smith (1952)
269.	<i>N. quadrivalvis</i>	<i>N. excelsior</i>	M	Zaitlin and Mundell (2006), Mihaylova-Kroumovova et al. (2020)
270.	<i>N. quadrivalvis</i>	<i>N. megalosiphon</i>	F	Kostoff (1943), Goodspeed (1945), Takenaka (1959)
271.	<i>N. quadrivalvis</i>	<i>N. suaveolens</i>	F, M	premendelian (East (1928)), Goodspeed and Clausen (1927), East (1935), Fardy and Hitier (1945), Näf (1958)
272.	<i>N. quadrivalvis</i>	<i>N. eastii</i>	M	Goodspeed (1945)
273.	<i>N. quadrivalvis</i>	<i>N. sylvestris</i>	F	Goodspeed (1934), Kostoff (1943), Goodspeed (1945)
274.	<i>N. quadrivalvis</i>	<i>N. tomentosa</i>	F	East (1935), Kostoff (1943), Goodspeed (1945)
275.	<i>N. quadrivalvis</i>	<i>N. tomentosiformis</i>	F	Kostoff (1943), Goodspeed (1945)
276.	<i>N. quadrivalvis</i>	<i>N. glutinosa</i>	F, M	premendelian (East (1928)), East (1935)
277.	<i>N. quadrivalvis</i>	<i>N. undulata</i>	asexual	Razdan (2003)
278.	<i>N. nesophila</i>	<i>N. repanda</i>	F	Goodspeed (1945)
279.	<i>N. nesophila</i>	<i>N. stocktonii</i>	F	Goodspeed (1945)
280.	<i>N. nesophila</i>	<i>N. umbratica</i>	M	Raju et al. (2008), Murthy et al. (2014)
281.	<i>N. nesophila</i>	<i>N. sylvestris</i>	F	Unnikrishnan et al. (1981)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
282.	<i>N. nudicaulis</i>	<i>N. repanda</i>	F	Kostoff (1943)
283.	<i>N. nudicaulis</i>	<i>N. rustica</i>	F, M	Kostoff (1943), Goodspeed (1954)
284.	<i>N. nudicaulis</i>	<i>N. debneyi</i>	F, M	Goodspeed (1934, 1945)
285.	<i>N. nudicaulis</i>	<i>N. exigua</i>	F, M	Kostoff (1943), Goodspeed (1945)
286.	<i>N. nudicaulis</i>	<i>N. suaveolens</i>	F, M	Kostoff (1943), Goodspeed (1945)
287.	<i>N. nudicaulis</i>	<i>N. sylvestris</i>	F	McCray (1932), Kostoff (1943)
288.	<i>N. nudicaulis</i>	<i>N. tomentosa</i>	F	Kostoff (1940, 1943), Goodspeed (1945, 1954)
289.	<i>N. nudicaulis</i>	<i>N. tomentosiformis</i>	F	Brieger (1929), Goodspeed (1945, 1954)
290.	<i>N. nudicaulis</i>	<i>N. obtusifolia</i>	F, M	Kostoff (1943)
291.	<i>N. nudicaulis</i>	<i>N. palmeri</i>	F	Kostoff (1943), Goodspeed (1945)
292.	<i>N. nudicaulis</i>	<i>N. glutinosa</i>	F	Christoff (1928), Kostoff (1943), Goodspeed (1945)
293.	<i>N. repanda</i>	<i>N. stocktonii</i>	F, M	Kostoff (1943)
294.	<i>N. repanda</i>	<i>N. africana</i>	F	Murthy and Rama Prasad (2000), Murthy and Subhashini (2000)
295.	<i>N. repanda</i>	<i>N. benthamiana</i>	M	Subhashini et al. (1998), Raju et al. (2008, 2009), Murthy et al. (2014)
296.	<i>N. repanda</i>	<i>N. debneyi</i>	M (l)	Yamada et al. (1999)
297.	<i>N. repanda</i>	<i>N. exigua</i>	M	Goodspeed (1945)
298.	<i>N. repanda</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1987), Murthy and Rama Prasad (2000)
299.	<i>N. repanda</i>	<i>N. sylvestris</i>	F	Kostoff (1940), Goodspeed (1945), Burk and Dropkin (1961)
300.	<i>N. repanda</i>	<i>N. tomentosiformis</i>	F (l)	Kobori and Marubashi (2004)
301.	<i>N. repanda</i>	<i>N. palmeri</i>	F	Goodspeed (1945), Takenaka (1958, 1959), Takenaka, 1962c, Schweppenhauser et al. (1963)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
302.	<i>N. repanda</i>	<i>N. glutinosa</i>	M	Kehr and Smith (1952)
303.	<i>N. stocktonii</i>	<i>N. sylvestris</i>	F (l)	Muraida and Marubashi (2015)
304.	<i>N. stocktonii</i>	<i>N. tomentosiformis</i>	F (l)	Muraida and Marubashi (2015)
305.	<i>N. rustica</i>	<i>N. benthamiana</i>	F	Sarala et al. (2023)
306.	<i>N. rustica</i>	<i>N. exigua</i>	F	Bolsunov (1970)
307.	<i>N. rustica</i>	<i>N. gossei</i>	F	Takenaka (1965)
308.	<i>N. rustica</i>	<i>N. megalosiphon</i>	F	Takenaka (1965)
309.	<i>N. rustica</i>	<i>N. suaveolens</i>	F	pre mendelian (East (1928)), Christoff (1938), Goodspeed (1945)
310.	<i>N. rustica</i>	<i>N. sylvestris</i>	asexual	Gleddie et al. (1983)
311.	<i>N. rustica</i>	<i>N. kawakamii</i>	F	Japan Tobacco Inc. (2022)
312.	<i>N. rustica</i>	<i>N. tomentosa</i>	F	Japan Tobacco Inc. (2022)
313.	<i>N. rustica</i>	<i>N. tomentosiformis</i>	F	Kostoff (1929)
314.	<i>N. rustica</i>	<i>N. palmeri</i>	F	Kostoff (1930), Goodspeed (1945)
315.	<i>N. rustica</i>	<i>N. glutinosa</i>	F	Lehmann (1936), Calitz and Milne (1962), Bannikova (1965)
316.	<i>N. rustica</i>	<i>N. undulata</i>	F	Kostoff (1940, 1943); Goodspeed (1945)
317.	<i>N. rustica</i>	<i>N. wigandoides</i>	F	Goodspeed (1934, 1945)
318.	<i>N. africana</i>	<i>N. benthamiana</i>	M	Iizuka et al. (2012)
319.	<i>N. africana</i>	<i>N. fragrans</i>	M	Gerstel et al. (1979)
320.	<i>N. africana</i>	<i>N. gossei</i>	M	Murthy and Subhashini (2000)
321.	<i>N. amplexicaulis</i>	<i>N. excelsior</i>	F	Gopinath et al. (1970), Gangadevi et al. (1987)
322.	<i>N. amplexicaulis</i>	<i>N. gossei</i>	not indicated	Gopinath et al. (1970)
323.	<i>N. amplexicaulis</i>	<i>N. megalosiphon</i>	F	Gopinath et al. (1965, 1970)
324.	<i>N. amplexicaulis</i>	<i>N. occidentalis</i>	M	Gopinath et al. (1965, 1970)
325.	<i>N. amplexicaulis</i>	<i>N. suaveolens</i>	F	Gopinath et al. (1965, 1970)
326.	<i>N. amplexicaulis</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1987)
327.	<i>N. amplexicaulis</i>	<i>N. velutina</i>	F	Gopinath et al. (1965, 1970)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
328.	<i>N. amplexicaulis</i>	<i>N. glutinosa</i>	F	Gopinath et al. (1965, 1970)
329.	<i>N. benthamiana</i>	<i>N. cavivola</i>	F	Williams (1975)
330.	<i>N. benthamiana</i>	<i>N. debneyi</i>	F	Bradley and Goodspeed (1943), Goodspeed (1945), Apparao and Ramavarma (1974)
331.	<i>N. benthamiana</i>	<i>N. excelsior</i>	F, M <sup>2</sup>	Fitzmaurice (2002), Tezuka et al. (2021b)
332.	<i>N. benthamiana</i>	<i>N. fragrans</i>	F	Williams (1975)
333.	<i>N. benthamiana</i>	<i>N. gossei</i>	M	Goodspeed (1945), Takenaka (1965), Durbin and Uchytíl (1977)
334.	<i>N. benthamiana</i>	<i>N. ingulba</i>	M	Apparao and Ramavarma (1974), Ramavarma and Apparao (1976)
335.	<i>N. benthamiana</i>	<i>N. megalosiphon</i>	F	Goodspeed (1945), Apparao and Ramavarma (1974)
336.	<i>N. benthamiana</i>	<i>N. suaveolens</i>	M	Wheeler (1945), Goodspeed (1945), Apparao and Ramavarma (1974)
337.	<i>N. benthamiana</i>	<i>N. umbratica</i>	M	Gangadevi et al. (1987)
338.	<i>N. benthamiana</i>	<i>N. velutina</i>	F	Horton (1981)
339.	<i>N. benthamiana</i>	<i>N. glutinosa</i>	F	Apparao and Ramavarma (1974)
340.	<i>N. burbridgeae</i>	<i>N. velutina</i>	F	Symon (1984)
341.	<i>N. cavicola</i>	<i>N. debneyi</i>	M	Williams (1975)
342.	<i>N. cavicola</i>	<i>N. umbratica</i>	F, M	Williams (1975), Gangadevi et al. (1987)
343.	<i>N. debneyi (forsteri)</i>	<i>N. exigua</i>	F, M	Kostoff (1943)
344.	<i>N. debneyi (forsteri)</i>	<i>N. fragrans</i>	F	Iizuka et al. (2012), Tezuka et al. (2021a)
345.	<i>N. debneyi (forsteri)</i>	<i>N. goodspeedii</i>	M	Goodspeed (1945)
346.	<i>N. debneyi (forsteri)</i>	<i>N. maritima</i>	F, M	Kostoff (1943), Goodspeed (1945)
347.	<i>N. debneyi (forsteri)</i>	<i>N. megalosiphon</i>	M	Goodspeed (1945, 1954)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
348.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. occidentalis</i>	F	Gangadevi et al. (1987)
349.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. simulans</i>	M	Gangadevi et al. (1987)
350.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. suaveolens</i>	F	Kostoff (1943), Williams (1975)
351.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. umbratica</i>	F	Williams (1975), Gangadevi and Satyanarayana (1982)
352.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. velutina</i>	F, M	Gillham et al. (1977)
353.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. sylvestris</i>	F, M (l)	Yamada et al. (1999), Tezuka et al. (2007)
354.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. otophora</i>	F	Takenaka (1959), Takenaka (1962c)
355.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. tomentosiformis</i>	F	Takenaka (1965), Tezuka et al. (2007)
356.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. obtusifolia</i>	F, M	Kostoff (1940, 1943), Goodspeed (1945)
357.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. palmeri</i>	F, M	Kostoff (1943), Goodspeed (1945)
358.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. thyrsiflora</i>		Ohashi et al. (1992)
359.	<i>N. debneyi</i> ( <i>forsteri</i> )	<i>N. glutinosa</i>	F, M	Dusseau et al. (1944), Fardy and Hitier (1945), Kehr and Smith (1952)
360.	<i>N. excelsior</i>	<i>N. gossei</i>	M, F	Goodspeed (1945), Murthy et al. (1998)
361.	<i>N. excelsior</i>	<i>N. suaveolens</i>	M	Näf (1958), Takenaka (1965)
362.	<i>N. excelsior</i>	<i>N. velutina</i>	F	Bradley and Goodspeed (1943)
363.	<i>N. excelsior</i>	<i>N. obtusifolia</i>	M	Murthy et al. (1998)
364.	<i>N. excelsior</i>	<i>N. glutinosa</i>	F	Murthy & Rao, 2008
365.	<i>N. exigua</i>	<i>N. goodspeedii</i>	F	Wheeler (1945), Goodspeed (1954)
366.	<i>N. exigua</i>	<i>N. gossei</i>	M	Goodspeed (1945)
367.	<i>N. exigua</i>	<i>N. maritima</i>	F, M	Kostoff (1940), Goodspeed (1945)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
368.	<i>N. exigua</i>	<i>N. megalosiphon</i>	F	Wheeler (1945), Goodspeed (1945)
369.	<i>N. exigua</i>	<i>N. suaveolens</i>	F	Kostoff (1940), Goodspeed (1945)
370.	<i>N. exigua</i>	<i>N. umbratrica</i>	F	Gangadevi et al. (1987)
371.	<i>N. exigua</i>	<i>N. velutina</i>	F, M	Kostoff (1943), Goodspeed (1954)
372.	<i>N. exigua</i>	<i>N. wuttkei</i>	M (l)	Laskowska and Berbeć (2003)
373.	<i>N. fragrans</i>	<i>N. umbratrica</i>	M	Williams (1975)
374.	<i>N. fragrans</i>	<i>N. glutinosa</i>	M	Williams (1975)
375.	<i>N. goodspeedii</i>	<i>N. megalosiphon</i>	F	Goodspeed (1945)
376.	<i>N. goodspeedii</i>	<i>N. rotundifolia</i>	F	Goodspeed (1945)
377.	<i>N. goodspeedii</i>	<i>N. suaveolens</i>	F	Wheeler (1945), Goodspeed (1945), Takenaka (1965)
378.	<i>N. goodspeedii</i>	<i>N. umbratrica</i>	F	Gangadevi et al. (1987)
379.	<i>N. gossei</i>	<i>N. maritima</i>	M	Wheeler (1945), Goodspeed (1954)
380.	<i>N. gossei</i>	<i>N. megalosiphon</i>	F	Kostoff (1943), Goodspeed (1945, 1954)
381.	<i>N. gossei</i>	<i>N. suaveolens</i>	M	Kostoff (1943), Goodspeed (1945), Takenaka (1952b)
382.	<i>N. gossei</i>	<i>N. occidentalis</i>	M	Wheeler (1945), Goodspeed (1945, 1954)
383.	<i>N. gossei</i>	<i>N. umbratrica</i>	F	Gangadevi et al. (1987)
384.	<i>N. gossei</i>	<i>N. velutina</i>	M	Wheeler (1945), Goodspeed (1945)
385.	<i>N. gossei</i>	<i>N. eastii</i>	M	Kostoff (1943), Goodspeed (1945, 1954)
386.	<i>N. gossei</i>	<i>N. glutinosa</i>	M	Murthy et al. (1998)
387.	<i>N. hesperis</i>	<i>N. umbratrica</i>	F	Gangadevi et al. (1987)
388.	<i>N. ingulba</i>	<i>N. megalosiphon</i>	F	Ramavarma and Apparao (1976)
389.	<i>N. ingulba</i>	<i>N. velutina</i>	F	Ramavarma and Apparao (1976)
390.	<i>N. ingulba</i>	<i>N. glutinosa</i>	F	Subhashini and Unnikrishnan (1978)
391.	<i>N. maritima</i>	<i>N. megalosiphon</i>	F	Goodspeed (1945, 1954)
392.	<i>N. maritima</i>	<i>N. suaveolens</i>	F, M	Kostoff (1939a), Goodspeed (1945)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
393.	<i>N. maritima</i>	<i>N. velutina</i>	F, M	Kostoff (1943), Goodspeed (1945)
394.	<i>N. maritima</i>	<i>N. wuttkei</i>	M	Laskowska and Berbeć (2003)
395.	<i>N. maritima</i>	<i>N. eastii</i>	M	Kostoff (1939a, 1940), Goodspeed (1945, 1954)
396.	<i>N. maritima</i>	<i>N. undulata</i>	F	Goodspeed (1945)
397.	<i>N. megalosiphon</i>	<i>N. suaveolens</i>	F, M	Kostoff (1940), Goodspeed (1945), Durbin and Uchytíl (1977)
398.	<i>N. megalosiphon</i>	<i>N. umbratica</i>	F	Subhashini (1975), Gangadevi et al. (1987)
399.	<i>N. megalosiphon</i>	<i>N. velutina</i>	M	Wheeler (1945), Goodspeed (1945)
400.	<i>N. megalosiphon</i>	<i>N. eastii</i>	M	Kostoff (1943), Goodspeed (1945)
401.	<i>N. megalosiphon</i>	<i>N. glutinosa</i>	F, M	Satyanarayana and Subhashini (1964), Subhashini et al. (1985)
402.	<i>N. occidentalis</i>	<i>N. simulans</i>	F, M	Gangadevi et al. (1987)
403.	<i>N. occidentalis</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1987)
404.	<i>N. occidentalis</i>	<i>N. sylvestris</i>	M(l), F	Tezuka and Marubashi (2012), Kawaguchi et al. (2021)
405.	<i>N. occidentalis</i>	<i>N. tomentosiformis</i>	F	Tezuka and Marubashi (2012), Kawaguchi et al. (2021)
406.	<i>N. occidentalis</i>	<i>N. undulata</i>	M	Takenaka (1965)
407.	<i>N. rosulata</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1987)
408.	<i>N. rotundifolia</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1987)
409.	<i>N. rotundifolia</i>	<i>N. velutina</i>	M	Bradley & Goodspeed, 1943, Goodspeed (1945)
410.	<i>N. simulans</i>	<i>N. umbratica</i>	F	Gangadevi et al. (1987)
411.	<i>N. suaveolens</i>	<i>N. velutina</i>	M	Goodspeed (1945, 1954), Chase et al. (2021)
412.	<i>N. suaveolens</i>	<i>N. wuttkei</i>	M (l)	Laskowska and Berbeć (2003)
413.	<i>N. suaveolens</i>	<i>N. eastii</i>	M	Kostoff (1939a), Goodspeed (1945, 1954)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
414.	<i>N. suaveolens</i>	<i>N. sylvestris</i>	asexual, F (l)	Fluhr (1983), Inoue et al. (1996)
415.	<i>N. suaveolens</i>	<i>N. tomentosiformis</i>	F	Inoue et al. (1996)
416.	<i>N. suaveolens</i>	<i>N. glutinosa</i>	F, M	pre mendelian (East (1928)), Dusseau et al. (1943), Goodspeed (1945), Gupta and Gupta (1973)
417.	<i>N. umbratica</i>	<i>N. velutina</i>	M	Gangadevi et al. (1987)
418.	<i>N. umbratica</i>	<i>N. kawakamii</i>	F	Japan Tobacco Inc. (2021)
419.	<i>N. umbratica</i>	<i>N. glutinosa</i>	F	Subhashini (1974)
420.	<i>N. velutina</i>	<i>N. wuttkei</i>	M	Laskowska and Berbeć (2003)
421.	<i>N. velutina</i>	<i>N. eastii</i>	F, M	Kostoff (1943)
422.	<i>N. sylvestris</i>	<i>N. kawakamii</i>	F	Ogura (1980)
423.	<i>N. sylvestris</i>	<i>N. otophora</i>	F, M	Goodspeed (1945, 1954), Takenaka (1954, 1956a)
424.	<i>N. sylvestris</i>	<i>N. setchellii</i>	F	Greenleaf (1942), Goodspeed (1945)
425.	<i>N. sylvestris</i>	<i>N. tomentosa</i>	F, M	Goodspeed (1934, 1945), East (1935), Takenaka (1956a)
426.	<i>N. sylvestris</i>	<i>N. tomentosiformis</i>	F, M	Kostoff (1930), Goodspeed (1945)
427.	<i>N. sylvestris</i>	<i>N. obtusifolia</i>	F	Liu and Marubashi (2014)
428.	<i>N. sylvestris</i>	<i>N. glutinosa</i>	F, M	Goodspeed (1934, 1945), East (1935), Fardy and Hitier (1945)
429.	<i>N. sylvestris</i>	<i>N. undulata</i>	cybrid	Aviv et al. (1984)
430.	<i>N. kawakamii</i>	<i>N. otophora</i>	F	Ogura (1980)
431.	<i>N. kawakamii</i>	<i>N. tomentosa</i>	F	Ogura (1980)
432.	<i>N. kawakamii</i>	<i>N. tomentosiformis</i>	F	Ogura (1980)
433.	<i>N. otophora</i>	<i>N. setchellii</i>	F, M	Goodspeed (1945), Bawolska et al. (1978)
434.	<i>N. otophora</i>	<i>N. tomentosa</i>	F	Goodspeed (1945)
435.	<i>N. otophora</i>	<i>N. tomentosiformis</i>	M	Gerstel and Burns (1974)
436.	<i>N. otophora</i>	<i>N. obtusifolia</i>	M (l)	Yamada et al. (1999)
437.	<i>N. otophora</i>	<i>N. glutinosa</i>	M	Goodspeed (1945)
438.	<i>N. setchellii</i>	<i>N. tomentosa</i>	M	Goodspeed (1945)
439.	<i>N. setchellii</i>	<i>N. tomentosiformis</i>	M	Clausen (1932), Goodspeed (1945)

(continued)

**Table 7.1** (continued)

No.	Hybrid combination		Species 1 used as: F (female parent); M (male parent) (l) denotes lethal hybrid	Reported by:
	Species 1	Species 2		
440.	<i>N. setchelli</i>	<i>N. obtusifolia</i>	M	Goodspeed (1945)
441.	<i>N. setchellii</i>	<i>N. palmeri</i>	M	Goodspeed (1945)
442.	<i>N. tomentosa</i>	<i>N. tomentosiformis</i>	F	Goodspeed (1945, 1954)
443.	<i>N. tomentosa</i>	<i>N. obtusifolia</i>	M	Goodspeed (1934, 1945), Kostoff (1943)
444.	<i>N. tomentosa</i>	<i>N. palmeri</i>	M	Kostoff (1943), Goodspeed (1954)
445.	<i>N. tomentosa</i>	<i>N. glutinosa</i>	M	Goodspeed (1934), Greenleaf (1941), Goodspeed (1945)
446.	<i>N. tomentosiformis</i>	<i>N. obtusifolia</i>	M	Goodspeed and Bradley (1942), Kostoff (1943), Goodspeed (1954), Liu and Marubashi (2014)
447.	<i>N. tomentosiformis</i>	<i>N. palmeri</i>	M	McCray (1933), Kostoff (1943)
448.	<i>N. tomentosiformis</i>	<i>N. glutinosa</i>	M	Kostoff (1932), Goodspeed (1945, 1954)
449.	<i>N. obtusifolia</i>	<i>N. palmeri</i>	F, M	Goodspeed (1945), Takenaka (1959, 1962c)
450.	<i>N. obtusifolia</i>	<i>N. glutinosa</i>	M, F	Krishnamoorthy and Bhat (1957), Appa Rao and Krishnamurthy (1963), Murthy et al. (1998)
451.	<i>N. arenstii</i>	<i>N. undulata</i>	M	Goodspeed (1945, 1954)
452.	<i>N. arenstii</i>	<i>N. wigandoides</i>	F	Goodspeed (1945)
453.	<i>N. glutinosa</i>	<i>N. wigandoides</i>	F	Elvers (1934), Goodspeed (1945)
454.	<i>N. glutinosa</i>	<i>N. undulata</i>	M	Goodspeed (1954)
455.	<i>N. undulata</i>	<i>N. wigandoides</i>	F	Goodspeed (1945, 1954)

<sup>1</sup>hybrid seeds were obtained but plants were not attempted to be grown therefrom.

<sup>2</sup>amphidiploid form of that hybrid is known as *Nicotiana excelsiana*.

<sup>3</sup>amphidiploid 4x (*N. longiflora* x *N. tabacum*) included in the study

<sup>4</sup>amphidiploid 4x *N. tabacum* x *N. gossei* included in the study

<sup>5</sup>regular sexual hybrid mentioned once but with no details

<sup>6</sup>somatic sterile aneuploids

1. five reports on the intrasectional hybrids with *N. rastroensis* – the hybrids did not develop beyond the seed stage and seeds were not sown. The author assumes that in this particular case successful fertilization and viable seeds may substitute for an actual living hybrid plant since there has been no report on post-zygotic crossability barriers between 9-chromosome species of the section Alatae;
2. reporting on 12 lethal hybrids that may have not actually been grown past the seedling stage. The argument is that those hybrids mostly belong to lethality type II, the crossability barrier that has been repeatedly demonstrated to be removable by tissue culture and, more recently, by chemical treatment and genetic engineering. Those two categories put aside, the inventory still includes 434 viable hybrid combinations including a few that may have not grown to full maturity.

## References

- Adachi, A., & Inoue, M. (1995). Temperature-response of interspecific hybrid plants between *Nicotiana gossei* Domin and *N. tabacum* L., obtained from ovule culture. *Japanese Journal of Breeding, suppl 2*, 287.
- Ahuja, M. R. (1962). A cytogenetic study of heritable tumors in *Nicotiana* species hybrids. *Genetics, 47*, 865–880.
- Ahuja, M. R. (1968). An hypothesis and evidence concerning the genetic components controlling tumour formation in *Nicotiana*. *Molecular & General Genetics, 103*, 176–184.
- Akada, S., & Hirai, A. (1986). Studies on the mode of separation of chloroplast genomes in parasexual hybrid calli. III. Random separation of two types of chloroplast genomes in a hybrid callus. *Japanese Journal of Genetics, 61*, 437–445.
- Anssour, S., Krugel, T., Sharbel, T. F., Saluz, H. P., Bonaventure, G., & Baldwin, I. T. (2009). Phenotypic, genetic and genomic consequences of natural and synthetic polyploidization of *Nicotiana attenuata* and *Nicotiana obtusifolia*. *Annals of Botany (Lond), 103*, 1207–1217. <https://doi.org/10.1093/aob/mcp058>
- Appa Rao, K., & Krishna Murthy, K. V. (1963). Studies on multiple polyploids in *Nicotiana*. *Genetica, 34*, 66–78.
- Apparao, K., & Ramavarma, K. T. (1974). Cytogenetics of the amphidiploid *Nicotiana benthamiana* × *N. Glutinosa*. *Cytologia, 39*, 379–382.
- Ar-Rushdi, A. H. (1955). Inheritance in *Nicotiana tabacum*. XXVI. Sterility genes from tomentosae species. *Journal of Genetics, 54*, 9–22.
- Avery, P. (1938). Cytogenetic evidences of *Nicotiana* phylesis in the *alata*-group. *University of California Publications in Botany, 18*, 153–194.
- Aviv, D., Arzee-Gonen, P., Bleichmann, S., & Galun, E. (1984). Novel alloplasmic *Nicotiana* plants by “donor-recipient” protoplast fusion: cybrids having *N. tabacum* or *N. sylvestris* nuclear genomes and either or both plastomes and chondriomes from alien species. *Molecular & General Genetics, 196*, 244–253.
- Bannikova, V. P. (1965). Cytoembryology of distant hybrids. II. Disturbances in the fertilization process on crossing *Nicotiana glutinosa* L. with *N. rustica*. *Ukrainian Botanical Journal, 22*, 40–46.
- Bartkowiak, E. (1987). Metody embriologii eksperimentalnej i hybrydyzacji somatycznej w otrzymywaniu mieszańców międzygatunkowych u roślin. *Postępy Biologii Komórki, 14*, 1–16.
- Bawolska, M., Kobus, I., & Jurzysta, A. (1978). A characteristic of spontaneous hybrid *Nicotiana setchellii* × *N. otophora*. *Genetica Polonica, 19*, 49–59.

- Bellair, G. (1913). Recroisées entre elles deux species qui sont dégagées d'un hybride n'obéissent plus à la loi Mendélienne de la dominance. *Comptes rendus et rapports: IVe Conférence internationale de génétique*, 201–203.
- Berbeć, A. (1988). Morphology, cytogenetics and resistance to potato virus Y of the amphidiploid *Nicotiana raimondii* Macbride × *N. tabacum* L. *Genetica Polonica*, 29, 41–52.
- Berbeć, A., & Doroszewska, T. (1981). Investigations of the interspecific hybrid *Nicotiana amplexicaulis* Burbidge × *Nicotiana tabacum*. *Genetica Polonica*, 22, 197–207.
- Berbeć, J., Berbeć, A., Glazewska, Z., Opoka, B., Kobus, I., & Czop, T. (1976). *Investigations on obtaining new varieties and breeding lines of Nicotiana tabacum resistant to virus leaf rib necrosis (potato virus Y) and virus degeneration of plants by Lycopersicum virus 3* (USDA Grant no. FG-Po-341 (JB-6) report no 2). Institute of Soils Science and Plant Cultivation, Pulawy, Poland (unpublished).
- Berbeć, J., Berbeć, A., Glazewska, Z., Opoka, B., Kobus, I., & Czop, T. (1982). *Investigations on obtaining new varieties and breeding lines of Nicotiana tabacum resistant to virus leaf rib necrosis (Potato Virus Y) and virus degeneration of plants by Lycopersicum virus 3* (USDA Grant no. FG-Po-341 (JB-6) Final Report). Institute of Soils Science and Plant Cultivation. (unpublished).
- Bolsunov, I. (1937). Sur les caractères des hybrides de première génération entre *N. rustica* et *N. glauca*. *Journal of Institution Botany Academic Science RSS Ukraine*, 11, 51–57.
- Bolsunov, I. (1970). New Nicotiana amphidiploid as a valuable starting material for the control of virulent blue mould lines. In *International Tobacco Scientific Congress, Hamburg 1970 Proceedings 1971* (p. 160)
- Bradley, M. V., & Goodspeed, T. H. (1943). Colchicine-induced Allo- and autopolyploidy in *Nicotiana*. *Proceedings of the National Academy of Sciences of the United States of America*, 29, 295–301.
- Breisser, E. (1934). Zytologisch-genetische Untersuchungen an der Bastarden *Nicotiana tabacum* × *N. rustica* und ihren Rueckkreuzungen zu *N. rustica* und *N. sylvestris* I. Die Nachkommen aus reduzierten Eizelle. *N. Zeitschrift fuer indukt. Abstamm. u. Vererbungsl*, 67, 115–151.
- Brieger, F. (1928). Über die Vermehrung der Chromosomenzahl bei dem Bastard *Nicotiana tabacum* L. × Rusbyi Britt. *Zeitschrift Inductive Abstam u Vererbungslehre*, 47, 1–53.
- Brieger, F. G. (1929). Vererbung bei Artbastarden unter besonderer Berücksichtigung der Gattung *Nicotiana*. *Der Zuechter*, 1, 140–152.
- Brieger, F. G., & Forster, R. (1942). Tumores em certos híbridos do gênero *Nicotiana*. *Bragantia*, 22, 2259–2274.
- Burk, L. G. (1960). Male-sterile flower anomalies in interspecific tobacco hybrids. *The Journal of Heredity*, 51, 27–31.
- Burk, L. G. (1972). Viable hybrids from monosomics *N. tabacum* by *N. langsdorffii*. *Tobacco Science*, 16, 43–45.
- Burk, L. G., & Dropkin, V. H. (1961). Response of *N. repanda*, *N. sylvestris* and their amphidiploid to root knot nematodes. *Plant Disease Report*, 45, 734–735.
- Burk, L. G., & Durbin, R. D. (1978). Reaction of *Nicotiana* species to tentoxin. *The Journal of Heredity*, 69, 117–120.
- Burk, L. G., Gooding, G. V., & Chaplin, J. F. (1982). Reaction of *Nicotiana* species and cultivars or breeding lines of *Nicotiana tabacum* to three strains of Potato Virus Y. *Tobacco Science*, 26, 85–88.
- Busconi, M., Reggi, S., Lorenzoni, C., & Fogher, C. (2010). *Interspecific crosses in the genus Nicotiana*. Proceedings of the 56th Italian Society of Agricultural Genetics Annual Congress, Post 7.02.
- Butenko, R. G., & Luneva, M. Z. (1966). Application of the sterile culture technique for the culturing of distant hybrids of *Nicotiana* (in Russian). *Fizjologija Rastenij*, 13, 733–736.
- Butenko, R. G., Ternovsky, M. F., Semenova, I. V., & Moiseeva, M. E. (1970). *5th International Tobacco Science Congress CORESTA Inf Bul 12*. Abstr A102.

- Calitz, P. C., & Milne, D. L. (1962). Reaction of *Nicotiana* species and species crosses to the root knot nematode, *Meloidogyne javanica*. *South African Journal of Agricultural Science*, 5, 123–126.
- Chaplin, J. F., & Mann, T. J. (1961). Interspecific hybridization, gene transfer and chromosomal substitution in *Nicotiana*. *North Carolina Agricultural Experiment Station Technical Bulletin*, 128.
- Chase, M. W., Fay, M. F., Nollet, F., & Christenhusz, M. (2021). *Nicotiana notha*. Solanaceae. *Curtis's Botanical Magazine*, 38. <https://doi.org/10.1111/curt.12401>
- Choi, S. J., & Lee, S. C. (1991). Interspecific hybrid of *Nicotiana trigonophylla* X *N. tabacum* through *in vitro* culture of fertilized ovules. *Journal of Korean Society of Tobacco Science*, 13, 52–58.
- Christoff, M. (1928). Cytological studies in the genus *Nicotiana*. *Genetics*, 13, 233–277.
- Christoff, M. (1938). *Wrkhu edin sluchai na mzhkka sterlnost pri Nicotiana rustica* (On a case of male sterility in *Nicotiana rustica*). Godisznik Sofijskiya University.
- Chung, C. S., Nakjima, T., & Takeda, G. (1988). Interspecific hybridization between *Nicotiana trigonophylla* Dun. and *N. tabacum* L. through ovule culture. *Japanese Journal of Breeding*, 38, 319–326.
- Clausen, R. E. (1932). Interspecific hybridization in *Nicotiana*. XIII. Further data as to the origin of *Nicotiana tabacum*. *Svensk Botanisk Tidskrift*, 26, 123–136.
- Clayton, E. E. (1950). Male sterile tobacco. *Journal of Heredity*, 4, 171–175.
- Clayton, E. E. (1954). Identifying disease resistance suited to interspecific transfer. *Journal of Heredity*, 45, 273–277.
- Cseplo, A., Nagy, F., & Maliga, P. (1983). *Rescue of a cytoplasmic lincomycin resistance factor from Nicotiana sylvestris into Nicotiana plumbaginifolia by protoplast fusion* (pp. 126–127). International Protoplast Symposium Poster Proceedings, Basel.
- Davis, E. L., Rich, J. R., & Gwynn, G. R. (1988). Reaction of selected *Nicotiana* spp. × *Nicotiana tabacum* crosses grown in microplots to 3 *Meloidogyne* spp. *Nematropica*, 18(2), 109–116.
- Descorbet, M. P. (2004). *Genotype and phenotype of Nicotiana alata × Nicotina rastroensis*. University of Missouri–Columbia. Office of Undergraduate Research.
- Descorbet, P., & McClure, B. (2005). Genotype and Phenotype of *Nicotiana alata* X *Nicotiana rastroensis*. In Vivo. *Cancer and the Environment 37th Annual MACUB Conference, Long Island University*, 26(2), 12.
- DeVerna, J. W. (1984). *In vitro-facilitated wide hybridization in Nicotiana*. PhD dissertation, University of Kentucky.
- DeVerna, J. W., Myers, J. R., & Collins, G. B. (1987). Bypassing prefertilization barriers to hybridization in *Nicotiana* using *in vitro* pollination and fertilization. *Theoretical and Applied Genetics*, 73, 665–671.
- Dorossiev, L., Palakarcheva, M., Stanoeva, M., & Petkova, M. (1978). Overcoming the sterility in  $F_1$  of interspecific hybrids of the genus *Nicotiana* using the methods of tissue culture. *Bul Spéc CORESTA Symposium Sofia*, 80–81. (P03).
- Durbin, R. D., & Uchytíl, T. F. (1977). Cytoplasmic inheritance of chloroplast coupling factor 1 subunits. *Biochemical Genetics*, 15, 1143–1146.
- Dusseau, A., Hitier, H., & Fardy, A. (1943). Transformation d'un hybride interspecifique sterile *Nicotiana suaveolens* Lehm (n=16) - a meiose asyndetique, en un hybride fertile rigoureusement autosyndetique par l'action de la colchicine. *Comptes Rendus de l'Académie des Sciences*, 217, 704–705.
- Dusseau, A., Fardy, A., & Hitier, H. (1944). *Nicotiana* polyploides: especies tetraploides et hybrides interspecificos amphiploides, obtenus par l'action de la colchicine. *Comptes Rendus de l'Académie des Sciences*, 218, 124–126.
- East, E. M. (1916). Inheritance in crosses between *Nicotiana langsdorffii* and *Nicotiana alata*. *Genetics*, 1, 311–333.
- East, E. M. (1928). The genetics of the genus *Nicotiana*. *Bibliography Genetics*, 4, 243–318.
- East, E. M. (1935). Genetic reactions in *Nicotiana*. I. Compatibility. *Genetics*, 20, 403–413.

- East, E. M., & Hayes, H. K. (1912). Heterozygosis in evolution and in plant breeding. *USDA Bulletin*, 243, 26–32.
- Eghis, S. A. (1927). Experiments on interspecific hybridation in the genus *Nicotiana* (in Russian with English summary). *Bulletin of Applied Botany, Leningrad*, 17, 151–190.
- Elvers, I. (1934). Interspecific hybridization in *Nicotiana*. XIV. The cytology of *F<sub>1</sub> glutinosa* × *tomentosa*. *University of California Publications in Botany*, 17, 341–354.
- Engvild, Y. (1950). Cytogenetic studies on the genus *Nicotiana*. *National Institute of Genetics Annual Report*, 1, 35.
- Fardy, A., & Hitier, H. (1945). Espèces tétraploïdes et hybrides interspécifiques amphidiploïdes et triples diploïdes ed *Nicotiana*, obtenus par l'action de la colchicine. *Mémorial du Serv d'Expl Industr des Tabacs et des Allumettes (SEITA) Ser B*, 1–117.
- Fatalizade, F. A. (1939). Acenaphthene-induced polyploidy in *Nicotiana*. *Doklady Akademii Nauk URSS*, 22, 180–183. (after Kostoff 1943).
- Fitzmaurice, W. P. (2002). *Interspecific Nicotiana hybrids and their progeny*. US Patent 6344597.
- Fluhr, R. (1983). The segregation of organelles and cytoplasmic traits in higher plant somatic fusion hybrids. *Protoplasts 1983, Experientia 46: Suppl*, Birkhauser Verlag, Basel:69–84.
- Focke, W. O. (1881). *Die Pflanzen-Mischlinge*. Borntraeger.
- Foster, H. H. (1943). Resistance in the genus *Nicotiana* to *Phytophthora parasitica* Dastur var. *nicotianae* Tucker. *Phytopathology*, 33, 403–404.
- Gangadevi, T., & Satyanarayana, K. V. (1982). Cytogenetic study of the interspecific cross of *Nicotiana debneyi* x *N. umbratica*. *Theoretical and Applied Genetics*, 63, 177–181.
- Gangadevi, T., Rao, P. N., Rao, B. H., et al. (1985). A study of morphology, cytology and sterility in interspecific hybrids and amphidiploids of *Nicotiana knightiana* x *N. umbratica*. *Theoretical and Applied Genetics*, 70, 330–332.
- Gangadevi, T., Panuganti, N. R., & Satyanarayana, K. V. (1987). Morphological and cytological studies of interspecific hybrids involving *N. umbratica* Burbidge. *Cytologia*, 52, 475–486.
- Gentscheff, G. (1931). Investigations on interspecific hybridisation in the genus *Nicotiana* (in Bulgarian). *Zemedlska Misl*, 2, 91–111.
- Gerstel, D. U., & Burns, J. A. (1974). Meiosis and staminal sterility in hybrids between *Nicotiana tomentosiformis* and *N. otophora*. *Tobacco Science*, 18, 157–159.
- Gerstel, D. U., Burns, J. A., & Burk, L. G. (1979). Interspecific hybridizations with an african tobacco, *Nicotiana africana* Merxm. *Journal of Heredity*, 70, 342–344.
- Gillham, F. E. M., Wark, D. C., & Harrigan, E. K. S. (1977). Disease resistant flue-cured tobacco breeding lines for north Queensland. I. Resistance to blue mould, *Peronospora tabacina*. *Australian Journal of Experimental Agriculture and Animal Husbandry*, 17, 652–558.
- Gisquet, P., Dufrenoy, J., & Dusseau, A. (1940). Hybrides interspécifiques de *Nicotiana tabacum* var. *Purpurea* × *N. petunoides* var. *Sylvestris*. *Science Genetics (Torino)*, 2, 67–78.
- Gleddie, S., Keller, W. A., Setterfield, G., & Wetter, L. R. (1983). Somatic hybridization between *Nicotiana rustica* and *N. sylvestris*. *Plant Cell, Tissue and Organ Culture*, 2, 269–283.
- Goodspeed, T. H. (1934). *Nicotiana* phylesis in the light of chromosome number, morphology and behaviour. *University of California Publications in Botany*, 17, 369–398.
- Goodspeed, T. H. (1945). Cytotaxonomy of *Nicotiana*. *Botanical Review*, 11, 533–592.
- Goodspeed, T. H. (1954). *The Genus Nicotiana*. Chronica Botanica Co.
- Goodspeed, T., & Bradley, M. (1942). Amphidiploidy. *Botanical Review*, 8, 272–316.
- Goodspeed TH, Clausen RE (1927) Interspecific hybridization in Nicotiana V. Cytological features of two *F<sub>1</sub>* hybrids made with *Nicotiana bigelovii* as a parent. University of California, Berkeley, University of California Press 11 (6)
- Goodspeed, T. H., & Clausen, R. E. (1928). Interspecific hybridization in *Nicotiana*. VIII. The *sylvestris-tomentosa-tabacum* triangle and its bearing on the origin of tobacco. *University of California Publications in Botany*, 11, 127–140.
- Goodspeed, T. H., & Thompson, M. C. (1959). *Cytotaxonomy of Nicotiana II*. University of California Botanical Garden.

- Gopinath, D. M., Krishnamurthy, K. V., & Krishnamurthy, A. S. (1965). Cytological studies in interspecific hybrids of *Nicotiana* involving a new Australian species, *Nicotiana amplexicaulis*. *Canadian Journal of Genetics and Cytology*, 7, 328–340.
- Gopinath, D. M., Krishnamurthy, K. V., & Krishnamurthy, A. S. (1970). A note on the cytogenetic status of *Nicotiana amplexicaulis* Burbidge. *Current Science*, 39, 410–411.
- Gray, J. C. (1978). Serological reaction of fraction I proteins from interspecific hybrids in the genus *Nicotiana*. *Plant Systematics and Evolution*, 129, 177–183.
- Greenleaf, W. H. (1941). Sterile amphidiploids: Their possible relation to the origin of *Nicotiana tabacum*. *The American Naturalist*, 75, 394–399.
- Greenleaf, W. H. (1942). Genic sterility in tabacum-like amphidiploids of *Nicotiana*. *Journal of Genetics*, 43, 69–96.
- Gupta, S. B., & Gupta, P. (1973). Selective somatic elimination of *Nicotiana glutinosa* chromosomes in the F<sub>1</sub> hybrids of *N. suaveolens* and *N. glutinosa*. *Genetics*, 73, 605–612.
- Horton, P. (1981). A taxonomic revision of *Nicotiana* (Solanaceae) in Australia. *Journal of the Adelaide Botanic Gardens*, 3, 1–56.
- Hranov, A. (1970). Some problems on crossability between *Nicotiana tabacum* L. and *N. megalosiphon* Heurk et Muell (in Bulgarian). *Bulgaria Tobacco*, 10, 45–48.
- Hu, C. (1956). Cytogenetic Studies in *Nicotiana*. XII. Reduction divisions in hybrids between *N. langsdorffii* and three other species. *Japanese Journal of Breeding*, 6, 117–121.
- Huesing, J., Jones, D., DeVerna, J., Myers, J., Collins, G., Severson, R., & Sisson, V. (1989). Biochemical investigations of antibiosis material in leaf exudate of wild *Nicotiana* species and interspecific hybrids. *Journal of Chemical Ecology*, 15, 1203–1217.
- Iizuka, T., Kuboyama, T., Marubashi, W., Oda, M., & Tezuka, T. (2012). *Nicotiana debneyi* has a single dominant gene causing hybrid lethality in crosses with *N. tabacum*. *Euphytica*, 186, 321–328.
- Ilcheva, V., San, L. H., Zagorska, N., & Dimitrov, B. (2001). Production of male sterile interspecific somatic hybrids between transgenic *N. tabacum* (bar) and *N. rotundifolia* (npt ii) and their identification by AFLP analysis. *In Vitro Cellular & Developmental Biology. Plant*, 37, 496–502.
- Ilin, G. (1948). Alkaloids of the amphidiploid *Nicotiana rustica* x *N. glauca*. *Doklady Akademii Nauk SSSR*, 59, 279–280.
- Inoue, E., Marubashi, W., & Niwa, M. (1996). Genomic factors controlling the lethality exhibited in the hybrid between *Nicotiana suaveolens* Lehm. and *N. tabacum* L. *Theoretical and Applied Genetics*, 93, 341–347.
- Iwai, S., Kishi, C., Nakata, K., & Kawashima, N. (1986). Production of *Nicotiana tabacum* x *N. acuminata* hybrid by ovule culture. *Plant Cell Reports*, 5, 403–404. [(Tob Abstr 31:639 (1987)].
- Japan Tobacco Inc. (2021). *Nicotiana* F<sub>1</sub> hybrid and its use. <https://patents.google.com/patent/JP6917478B2/en?inventor=%E4%BF%8A+%E5%B1%B1%E5%86%85>
- Japan Tobacco Inc. (2022). Genus *Nicotiana* F<sub>1</sub> hybrid and use thereof. <https://patents.google.com/patent/WO2022014699A1/en?inventor=%E4%BF%8A+%E5%B1%B1%E5%86%85>
- Jeanes, J. A. (1999). Solanaceae (in) Walsh NG. In T. J. Entwistle (Ed.), *Flora of Victoria vol 4, Cornaceae to Asteraceae*. Inkata Press.
- Kaul, M. H. L. (1988). *Male sterility in higher plants* (p. 10). Springer.
- Kawaguchi, K., Ohya, Y., Maekawa, M., Iizuka, T., Hasegawa, A., et al. (2021). Two *Nicotiana occidentalis* accessions enable gene identification for Type II hybrid lethality by the cross to *N. sylvestris*. *Science Reports*, 11, 17093. <https://doi.org/10.1038/s41598-021-96482-6>
- Kehr, A. E. (1951). Genetic tumors in *Nicotiana*. *The American Naturalist*, 99, 73–79.
- Kehr, A. E., & Smith, H. H. (1952). Multiple genome relationships in *Nicotiana*. *Cornell University Agricultural Experiment Station Memoir*, 311, 3–19.
- Kehr, A. E., & Smith, H. H. (1954). Genetic tumors in *Nicotiana* hybrids. *Brookhaven Symposia in Biology*, 6, 55–78.
- Kincaid, R. R. (1949). Three interspecific hybrids of tobacco. *Phytopathology*, 39, 284–287.

- Kiraly, L., & Schoeltz, J. E. (1995). The interaction of a single, recessive gene in *Nicotiana clevelandii* and gene VI of cauliflower mosaic virus results in systemic necrosis. *Phytopathology*, 85, 1152.
- Kobori, S., & Marubashi, W. (2004). Programmed death cell detected in interspecific hybrids of *Nicotiana repanda* x *N. tomentosiformis* expressing hybrid lethality. *Breeding Science*, 54, 347–350.
- Kobus, I. (1971). Investigations in the polyploids of *Nicotiana tabacum* L. and their hybrids with wild tobacco species. *Genetica Polonica*, 12, 323–328.
- Kostoff, D. (1929). Acquired immunity in plants. *Genetics*, 14, 37–77.
- Kostoff, D. (1930). Ontogeny, genetics, and cytology of *Nicotiana* hybrids. *Genetica*, 12, 33–139.
- Kostoff, D. (1932). Pollen abortion in species hybrids. *Cytologia*, 5, 337–339.
- Kostoff, D. (1935). On the Increase of Mutation Frequency following Inter Specific Hybridization in *Nicotiana*. *Current Science*, 3, 302–304.
- Kostoff, D. (1939a). The origin of the tetraploid *Nicotiana* from Bathurst. *Current Science*, 8, 110–112.
- Kostoff, D. (1939b). Abnormal mitosis in tobacco plants forming hereditary tumours. *Nature*, 144, 599.
- Kostoff, D. (1939c). Nicotine and citric acid content in the progeny of the allopolyploid hybrid *Nicotiana rustica* L. x *N. glauca* Grah. *Doklady Akademii Nauk SSSR*, 23, 121–124.
- Kostoff, D. (1940). Relation degrees and phylesis of certain *Nicotiana* species determined by cytogenetic analysis. *Genetica*, 22, 215–230.
- Kostoff, D. (1943). *Cytogenetics of the genus Nicotiana*. State Printing House.
- Krishnamoorthy, K. V., & Bhat, N. R. (1957). An amphidiploid *Nicotiana glutinosa* L. *Nicotiana trigonophylla* Dun. Hybrid. *Current Science*, 26, 120.
- Krügel, T. (2010). *Nicotiana* × *mierata* und *N. obtusata*: Zwei neue synthetische Hybriden. *Haussknechtia*, 12, 23–25.
- Kubo, T. (1985). Studies on hybrid breeding by means of cytoplasmic male sterility in tobacco. *Bulletin Iwata Tobacco Experiment Station*, 17, 69–138.
- Larkina, N. I. (1980). Overcoming incompatibility between *Nicotiana* species by means of *in vitro* pollination. *Plant Breed Abstract*, 50(8067), 690.
- Larkina, N. I. (2015). *Scientific principles of interspecific hybridization using maternal Nicotiana tabacum*. Monograph. All-Russian Institute of Tobacco and Makhorka. Krasnodar ISBN 978-5-93491-681-8.
- Larkina, N. I. (2017). Creation of amphidiploids – fertile interspecific *Nicotiana* hybrids. *Sciences of Europe. Biological Sciences*, 2, 95–105.
- Laskowska, D., & Berbeć, A. (2003). Preliminary study of the newly discovered tobacco species *Nicotiana wuttkei* Clarkson and Symon. *Genetic Resources and Crop Evolution*, 50, 835–839.
- Laskowska, D., & Berbeć, A. (2012). Production and characterization of amphihaploid hybrids between *Nicotiana wuttkei* Clarkson et Symon and *N. tabacum*. *Euphytica*, 183, 75–82.
- Lautz, W. (1957). Resistance to black shank of 51 species of *Nicotiana* and of 13 interspecific hybrids. *Plant Disease Report*, 41, 95–98.
- Lee, C. B., Page, L. E., McClure, B. A., & Holtsford, T. P. (2008). Post-pollination hybridization barriers in *Nicotiana* section *Alatae*. *Sexual Plant Reproduction*, 21, 183–195.
- Lehmann, H. (1936). On interspecific cross sterility in the genus *Nicotiana*. *Plant Breed Abstract*, 7(1936), abst 1023 (in German)
- Lilienfeld, F. A. (1953). Cytogenetical problems in *Nicotiana*. a. A new amphidiploid. *National Institute of Genetics Japanese Annual Report*, 4, 32–33.
- Lin, R. F., & Chen, C. C. (1990). Cytological studies of interspecific somatic hybrids in *Nicotiana*. *Botanical Bulletin of Academia Sinica*, 31, 179–187.
- Liu, H., & Marubashi, W. (2014). Species origin of genomic factors in *Nicotiana nudicaulis* Watson controlling hybrid lethality in interspecific hybrids between *N. nudicaulis* Watson and *N. tabacum* L. *PLoS One*, 9, e97004. <https://doi.org/10.1371/journal.pone.0097004>

- Ma, J., Hancock, W. G., Nifong, J. M., Kernodle, S. P., & Lewis, R. S. (2020). Identification and editing of a hybrid lethality gene expands the range of interspecific hybridization potential in *Nicotiana*. *Theoretical and Applied Genetics*, 133, 2915–2925. <https://doi.org/10.1007/s00122-020-03641-w>
- Malinowski, E. (1916). On the occurrence of new forms in the progeny of the hybrids *Nicotiana atropurpurea* × *Nicotiana sylvestris* (in Polish). *Comptes Rendus des Séances de la Société des Sciences et des Lettres de Varsovie*, 9, 827–864.
- Malloch, W. S., & Malloch, F. W. (1924). Species crosses in *Nicotiana* with special reference to *N. longiflora* × *N. tabacum*, *N. longiflora* × *N. sanderae*, *N. tabacum* × *N. glauca*. *Genetics*, 9, 261–291.
- Maryanovich, O. (1939). On the cytology and embryology of the hybrid F<sub>1</sub> *Nicotiana rustica* (variety Unterwalden) × *Nicotiana quadrivalvis* (in Russian). *Zhurnal Instytutu Botaniky Ukrayins'koyi Akademiy Nauk. USSR* (21–22 (29–30)):187–195.
- Masuta, C., Suzuki, M., Takanami, Y., & Koiwai, A. (1993). Yellow mosaic symptoms induced by Y satellite RNA of Cucumber Mosaic Virus is regulated by a single incompletely dominant gene in wild *Nicotiana* species. *Phytopathology*, 83, 411–413.
- Mayr, E. (1986). Joseph Gottlieb Koelreuter's contribution to biology. *Osiris*, 2, 135–176. University of Chicago Press.
- McCray, F. A. (1932). Compatibility of certain *Nicotiana species*. *Genetics*, 17, 621–636.
- McCray, F. A. (1933). Embryo development in *Nicotiana* species hybrids. *Genetics*, 18, 95–110.
- Meiselman, N., Gunckel, J. E., & Sparrow, A. H. (1961). The general morphology and growth responses of two species of *Nicotiana* and their interspecific hybrid after chronic gamma irradiation. *Radiation Botany*, 1, 69–79.
- Menczel, L., Lazar, G., & Maliga, P. (1978). Isolation of somatic hybrids by cloning *Nicotiana* heterokaryons in nurse cultures. *Planta*, 143, 29–32.
- Mihaylova-Kroumova, A. B., Atrouchine, I., Korenkov, V. D., & Wagner, J. G. (2020). *Patterns of inheritance of acylsugar acyl groups in selected Interspecific hybrids of genus Nicotiana* (p. 23). Kentucky Tobacco Research and Development Center Faculty Publications.
- Moav, R., & Cameron, D. R. (1960). Genetic instability in *Nicotiana* hybrids. I. The expression of instability in *N. tabacum* × *N. plumbaginifolia*. *American Journal of Botany*, 47, 87–93.
- Modilevsky, J. (1937). Cytogenetic investigation of the genus *Nicotiana*. III. Cytology and embryology of the hybrid F1 *N. rustica* L. × *N. glauca* Grah (in Russian). *Zhurnal Instytutu Botaniky Ukrayins'koyi Akademiy Nauk USSR*, 11, 59–71.
- Muntingz, A. (1935). Chromosome behaviour in some *Nicotiana* hybrids. *Hereditas*, 20, 251–271.
- Muraida, N., & Marubashi, W. (2015). Characterization of hybrid seedlings from crosses of *Nicotiana stocktonii* Brandegee × *N. tabacum* L. and *N. stocktonii* × progenitors of *N. tabacum*. *Plant Biotechnology*, 32, 139–147.
- Murthy, T. G. K., & Rama Prasad, G. (2000). Aphid resistance in some *Nicotiana* species and interspecific hybrids. *Tobacco Research*, 26, 52–56.
- Murthy, T. G. K., & Rao, R. V. S. (2008). *Maintenance of the genus Nicotiana* (pp. 18–20). Central Tobacco Research Institute Annual Report 2007–2008, Rajamundry, India.
- Murthy, T. G. K., & Subbarao, I. V. (2004). Some new interspecific hybrids in the genus *Nicotiana* – Characterization and utilization. *Tobacco Research*, 30, 33–41.
- Murthy, T. G. K., & Subhashini, U. (2000). Studies on some interspecific hybrids of *Nicotiana africana* Merxm. *Tobacco Research*, 26, 12–15.
- Murthy, T. G. K., Subhashini, U., & Subba Rao, I. V. (1998). New interspecific hybrids in the genus *Nicotiana* – Characterization and utilization. *Tobacco Symposium*, 8, abstr B10.
- Murthy, T. G. K., Sredhaar, U., & Siva, R. K. (2014). *Incorporation of aphid resistance from N. gossei. N. repanda, N x. umbratica-nesophila and N × benthamiana-repanda* (pp. 17–20). Central Tobacco Research Institute Annual Report 2013–2014, Rajamundry, India.
- Näf, U. (1958). Studies on tumour formation in *Nicotiana* hybrids I. The classification of parents into two etiologically significant groups. *Growth*, 22, 167–180.

- Navarro-Quezada, A. (2014). *Quantification of homeolog expression bias upon herbivory of synthetic and natural allopolyploids from N. attenuata x N. obtusifolia*. ICE Symposium MPI for Chemical Ecology, Jena, DE.
- Navarro-Quezada, A., Alejandro Perez, N., Gaquerel, E., & Baldwin, I. T. (2013). *Herbivore resistance in natural- and artificial Nicotiana attenuata x N. obtusifolia allopolyploids: A transcriptome and metabolome analysis*. Conference on Plant Genome Evolution, Elsevier.
- Nikova, V. M. (1986). *Nicotiana excelsior* Black as a source of cytoplasmic male sterility in *Nicotiana tabacum* L. *Comptes Rendus de L'Academie Bulgare des Sciencesopen*, 39, 105–107.
- Nikova, V. M., & Zagorska, N. A. (1990). Overcoming hybrid incompatibility between *Nicotiana africana* Merxm and *Nicotiana tabacum* and development of cytoplasmically male sterile tobacco forms. *Plant Cell, Tissue and Organ Culture*, 32, 71–75.
- Nikova, V. M., Zagorska, N. A., & Pundeva, R. S. (1991). Development of four tobacco male sterile sources using *in vitro* techniques. *Plant Cell, Tissue and Organ Culture*, 27, 289–295.
- Nikova, V., Vladova, R., Pundeva, R., Petkova, A., & Vladovska, A. (1998). *Study of Nicotiana ingulba* Black. and its crossability with *N. tabacum* L (p. 150). CORESTA Congr Bull Spec, Brighton. PPOST 12 (b).
- Nikova, V., Iancheva, A. M., Vladova, R., Pandeva, R., & Petkova, A. (2006). *Genetic improvement of Nicotiana tabacum by applying remote hybridization and biotechnological methods* (p. 07). CORESTA Congr Agro-Phyto Groups, Paris, APOST.
- Ogura, H. (1980). Cytogenetic studies on a recently identified *Nicotiana* species, *N. kawakamii* Ohashi. *Cytologia*, 45, 33–42.
- Ohashi, Y. (1985). Breeding studies of wild species related to tobacco, with special reference to disease resistance. *Bulletin Iwata Tobacco Experiment Station*, 17, 1–62.
- Ohashi, Y., Ohshima, M., & Itoh, H. (1992). Constitutive expression of stress-inducible genes, including pathogenesis-related 1 protein gene in a transgenic interspecific hybrid of *Nicotiana glutinosa* x *Nicotiana debneyi*. *Plant & Cell Physiology*, 33-2, 177–187.
- Pal, B. P., & Nath, P. (1936). A note on the sterile hybrid between *N. tabacum* and *N. plumbaginifolia*. *Indian Journal of Agricultural Sciences*, 6, 828–832.
- Palakarcheva, M. (1975). Interspecific hybridization in tobacco (in Bulgarian). *Bulgarian Tiutun*, 11, 30–34.
- Palakarcheva, M. (1981). Genetic peculiarities of tobacco breeding for disease resistance (in Bulgarian). *Genetics Selection*, 14, 277–284.
- Patel, M. S. (1960). *Indian tobacco. A monograph*. Indian Central Tobacco Committee.
- Patel, D., Power, J. B., Anthony, P., Badakshi, F., Hesslop-Harison, J. S., & Davey, M. R. (2011). Somatic hybrid plants of *Nicotiana × sanderae* (+) *N. debneyi* with fungal resistance to *Peronospora tabacina*. *Annals of Botany*, 108, 809–819.
- Pearse, I. S., Krugel, T., & Baldwin, I. T. (2006). Innovation in anti-herbivore defense systems during neopolyploidy – The functional consequences of instantaneous speciation. *The Plant Journal*, 47, 196–210.
- Pittarelli, G. W., & Stavely, J. R. (1975). Direct hybridization of *Nicotiana repanda* x *N. tabacum*. *The Journal of Heredity*, 66, 281–284.
- Poddubnaya-Arnoldi, V. A., & Ludkina, M. M. (1945). Embryogenesis in hybridization of distant species and polyploidy in the genus *Nicotiana* (in Russian). *Zhurnal Instytutu Botaniki Ukrains'koyi Akademiyi Nauk USSR*, 30, 195–216.
- Raju, K. S., Sheshumadav, M., & Murthy, T. G. K. (2008). Molecular diversity in the genus *Nicotiana* as revealed by randomly amplified polymorphic DNA. *Physiology and Molecular Biology of Plants*, 14, 377–382.
- Raju, K. S., Sheshumadav, M., Chandrasekhararao, C., & Murthy, T. G. K. (2009). Molecular diversity in genus *Nicotiana* as revealed by randomly amplified polymorphic DNA. *Indian Journal of Biotechnology*, 8, 61–66.
- Ramanujam, S., & Joshi, A. B. (1942). Interspecific hybridization in *Nicotiana*. A cytological study of the hybrid *N. glauca* Grah. X *N. plumbaginifolia*. *Indian Journal Of Genetics And Plant Breeding*, 2, 80–97.

- Ramavarma, K. T., & Apparao, K. (1976). Studies on interspecific hybrids in genus *Nicotiana* involving the species *Nicotiana ingulba*. *Tobacco Research*, 2, 137–140.
- Razdan, M. K. (2003). Chapter XII. Somatic hybridisation and cybridisation (in) Introduction to plant tissue Culture. *Science Publishers, Appendix*, 12(2), 174.
- Reed, S., & Collins, G. B. (1978). Interspecific hybrids in *Nicotiana* through in vitro culture of fertilized ovules. *The Journal of Heredity*, 69, 311–315.
- Reed, S., & Collins, G. B. (1980). Chromosome pairing relationship and black shank resistance in three *Nicotiana* interspecific hybrids. *The Journal of Heredity*, 71, 423–426.
- Sachs-Skalińska, M. (1917). Beitrag zur Cytologie des sterilen Bastardes *Nicotiana atropurpurea* × *Nicotiana silvestris*. *Comptes Rendus des Séances de la Société des Sciences et des Lettres de Varsovie*, 10(6).
- Sarala, K., Rao, R. V. S., Murthy, T. D. K., & Satyavani, J. V. R. (2008). Micropropagation of sterile and non-flowering *Nicotiana* lines. *Cytologia*, 73, 9–13.
- Sarala, K., Prabhakara Rao, K., & Sheshu Madhav, M. (2023). *Interspecific hybridization. Division of crop improvement* (pp. 11–14). ICAR -Central Tobacco Research Institute Technical Bulletin.
- Satyanarayana, K. V., & Subhashini, L. (1964). Interspecific hybridization and aneuploidy in the genus *Nicotiana*. *Indian Journal of Genetics and Plant Breeding*, 24, 264–271.
- Savelli, R. (1926). Un ibrido di tipo eccezionale nella F1 di *Nicotiana longiflora* Cav. X *N. sylvestris* Sp. Et Comes. *Arch Bot Sist Modena*, 2, 228–240.
- Savelli, R. (1927). Method of measuring the flowers of *Nicotiana* and biometric data on the hybrid *Nicotiana rustica* x. *N. tabacum*. *Arch Bot Sist Fitogeogr Genet*, 3, 95–106.
- Schweppenhauser, M. A., Raeber, J. G., & Daulton, R. A. C. (1963). *Resistance to the root knot nematode Meloidogyne javanica in the genus Nicotiana* (pp. 222–229). Proceedings of III World Tobacco Science Congress, Salisbury, S Rhodesia.
- Sievert, R. C. (1972). Resistance to anthracnose in the genus *Nicotiana*. *Tobacco Science*, 16, 32–34.
- Smith, H. H. (1937). Quantitative genetic interactions. I. Genome combinations of *Nicotiana longiflora* and *N. sanderae*. *Genetics*, 22, 361–375.
- Smith, H. H. (1942). Alkaloids in certain species and interspecific hybrids of *Nicotiana*. *Journal of Agricultural Research*, 347–359.
- Smith, H. H. (1958). Genetic plant tumors in *Nicotiana*. *Annals of the New York Academy of Sciences*, 71, 1163–1178.
- Smith, H. H. (1968). Recent cytogenetic studies in the genus *Nicotiana*. *Advances in Genetics*, 14, 1–43.
- Smith, H. H., & Abashian, D. V. (1963). Chromatographic investigations on the alkaloid content of *Nicotiana* species and interspecific combinations. *American Journal of Botany*, 50, 435–447.
- Smith, H. H., Hamill, D. E., Weaver, E. A., & Thompson, K. H. (1970). Multiple molecular forms of peroxidase and esterase among *Nicotiana* species and amphidiploids. *The Journal of Heredity*, 61, 203–212.
- Sorrentino, C., Del Piano, L., Crimaldi, M., Abet, M., & Cuciniello, A. (2006). *Constitutional of ornamental hybrids of Nicotiana* ssp. Proceedings of the 50th Italian Society of Agricultural Genetics Annual Congress, Ischia, Italy.
- Stavely, J. R. (1979). Disease resistance. In R. D. Durbin (Ed.), *Nicotiana. Procedures for Experimental Use* (U.S. Department of Agriculture Technical Bulletin 1586) (pp. 111–120).
- Stehmann, J. R., Semir, J., & Ippolito, A. (2002). *Nicotiana mutabilis* (Solanaceae), a new species from southern Brazil. *Kew Bulletin*, 57, 639–646.
- Subhashini, U. (1974). Study of the cytoplasmic effects in the interspecific cross *Nicotiana umbratica* Burbidge & *N. glutinosa* Linn. *Euphytica*, 23, 289–293.
- Subhashini, U. (1975). Interspecific Hybridization in the Genus *Nicotiana* Involving the Species *N. umbratica* B and *N. megalosiphon* H and M Arg. *Cytologia*, 40, 409–413.
- Subhashini, U., & Gopinath, D. M. (1974). Interspecific Hybridization in the Genus *Nicotiana* Involving a New Australian Species *N. umbratica*. *Cytologia*, 39, 403–409.

- Subhashini, U., & Unnikrishnan, M. (1978). Cytogenetic relationship of *Nicotiana ingulba* B. with *N. glutinosa* L. *Tobacco Research*, 2, 72–74.
- Subhashini, U., Venkateswarlu, T., Anjani, K., & Prasad, G. S. R. (1985). In vitro hybridization in an incompatible cross *Nicotiana glutinosa* times *Nicotiana megalosiphon*. *Theoretical and Applied Genetics*, 71, 545–549.
- Subhashini, U., Venkateswarlu, T., & Anjani, K. (1986). Embryo rescue in *Nicotiana* hybrids by in vitro culture. *Plant Science*, 43, 219–222.
- Subhashini, U., Murthy, T. G. K., & Siva Raju, K. (1998). Genetic transformation in *N. tabacum* using direct gene transfer from an amphidiploid (*N. benthamiana* × *N. repanda*). *Tobacco Symposium*, 7, abstr. B9.
- Symon, D. E. (1984). A new species of *Nicotiana* (Solanaceae) from Dalhousie Springs. *South Australian Journal of the Adelaide Botanic Gardens*, 7, 117–121.
- Takanashi, H., & Marubashi, W. (2017). Tumorigenesis inheritance from the putative progenitor species of *Nicotiana rustica*. *Plant Biotechnology*, 34, 79–87.
- Takenaka, Y. (1950). Cytogenetic studies on the Genus *Nicotiana*. *National Institute of Genetics Annual Report*, 1, 35.
- Takenaka, Y. (1951a). Cytogenetic studies on the genus *Nicotiana* II (c). Hybrid between *N. glauca* and *N. plumbaginifolia*. *National Institute of Genetics Annual Report*, 2, 28.
- Takenaka, Y. (1951b). Cytogenetic studies in *Nicotiana*. On three hybrids between *N. tabacum* and other species. *La Kromosoma*, 9–10, 384–392.
- Takenaka, Y. (1952a). Cytogenetic studies in the genus *Nicotiana*. IV. (a) Reduction divisions in hybrids between *N. glauca* and the *N. alata* group. *National Institute of Genetics Annual Report*, 3, 40–41.
- Takenaka, Y. (1952b). Cytogenetic studies in the genus *Nicotiana*. IV. (c) Reduction divisions in hybrids between *N. suaveolens* and three other species. *National Institute of Genetics Annual Report*, 3, 42.
- Takenaka, Y. (1953a). Cytogenetic studies of *Nicotiana*. III. Reduction divisions in three hybrids between *N. glauca* and the *alata* group. *Japanese Journal of Genetics*, 28, 155–162.
- Takenaka, Y. (1953b). Cytogenetic studies in the genus *Nicotiana*. V. (b) Reduction divisions in the hybrids between the *tomentosa* group and three other species. *National Institute of Genetics Annual Report*, 4, 34–35.
- Takenaka, Y. (1953c). Cytogenetic Studies on the genus *Nicotiana*. V. (c) Reduction divisions in hybrids between *N. sanderae* and three other species. *National Institute of Genetics Annual Report*, 4, 35–36.
- Takenaka, Y. (1954). Cytogenetic studies in the genus *Nicotiana*. VI. b. Reduction divisions in several hybrids between *Nicotiana* species. *National Institute of Genetics Annual Report*, 5, 64–65.
- Takenaka. (1955). Cytogenetic studies of Nicotiana. VIII. Reduction divisions in hybrids between *N. sanderae* and three other species. *Japanese Journal of Genetics*, 30, 236–241.
- Takenaka, Y. (1956a). Cytogenetic studies in *Nicotiana*. XIV. Reduction divisions in five interspecific hybrids. *Japanese Journal of Genetics*, 31, 155–161.
- Takenaka, Y. (1956b). Cytogenetic studies in Nicotiana. VI. On three hybrids between *N. tabacum* and other species. *Journal of Japanese Botany*, 68, 358–362.
- Takenaka, Y. (1959). Cytogenetic studies of the genus *Nicotiana* XII. *National Institute of Genetics Annual Report*, 10, 77–79.
- Takenaka, Y. (1961). Cytogenetic studies in the genus Nicotiana XIV. *National Institute of Genetics Annual Report*, 12, 68–69.
- Takenaka, Y. (1962a). Cytogenetic studies in Nicotiana, XIX. The reduction division in four hybrids between *N. paniculata* and other species of the section Alatae. *Japanese Journal of Breeding*, 12, 278–280.
- Takenaka, Y. (1962b). Cytogenetic Studies in *Nicotiana*, XV. Reduction divisions in five Interspecific hybrids between sections Alatae and Suaveolentes. *Japanese Journal of Genetics*, 37, 80–85.

- Takenaka, Y. (1962c). Cytogenetic Studies in *Nicotiana*. XVII. Reduction Divisions in Five Interspecific Hybrids. *Japanese Journal of Genetics*, 37, 343–347.
- Takenaka, Y. (1962d). Cytogenetic studies in *Nicotiana* XVI. Reduction divisions in six interspecific hybrids between *N. tabacum* and other six species. *Shokubutsugaku Zasshi*, 75(888), 237–241.
- Takenaka, Y. (1965). Cytogenetic studies in the genus *Nicotiana*. *National Institute of Genetics Japanese Annual Report*, 16, 75–76.
- Takenaka, Y., & Yoneda, Y. (1964). Tumor formation in interspecific hybrids of *Nicotiana*, especially in relation to injury. *National Institute of Genetics Annual Report*, 15, 87.
- Takenaka, Y., Amano, Y., & Tateoka, T. (1956). Cytogenetic studies in the genus *Nicotiana*. IX. *Japanese National Institute of Genetics Annual Report*, 7, 42–44.
- Tanaka, M. (1961). The effect of irradiated pollen grains on species crosses of *Nicotiana*. *Bulletin of the Hatano Tobacco Experiment Station*, 51, 1–38.
- Tatemichi, Y. (Ed.). (1990). *Illustrated book of the genus Nicotiana*. Plant Breeding and Genetics Research Laboratory, Japan Tobacco Inc.
- Ternovsky, M. F. (1962). *Summary and prospects of interspecific hybridization in the genus Nicotiana (in) Wide hybridization of plants* (pp. 331–345). Proceedings of the Conference on Wide Hybridization of Plants and Animals; collection of reports.
- Ternovsky, M. F., & Larkina, N. I. (1978). Hybridization of native tobacco cultivars with the wild species *Nicotiana rosulata*. *Soviet Genetics*, 14, 1038–1045.
- Ternovsky, M. F., Butenko, R. G., & Moiseeva, M. E. (1972). The use of tissue culture to overcome the barrier of incompatibility between species and sterility of interspecific hybrids. *Soviet Genetics*, 8, 27–33.
- Ternovsky, M. F., Shinkareva, I. K., & Larkina, N. I. (1976). Production of interspecific tobacco hybrids by the pollination of ovules *in vitro*. *Soviet Genetics*, 12, 1209–1213.
- Tezuka, T. (2012). Hybrid lethality in the Genus *Nicotiana*. In J. K. Mworia (Ed.), *Botany* (pp. 191–210). Rijeka InTech.
- Tezuka, T., & Marubashi, W. (2012). Genes in S and T subgenomes are responsible for hybrid lethality in interspecific hybrids between *Nicotiana tabacum* and *Nicotiana occidentalis*. *Online Plos One*. <https://doi.org/10.1371/journal.pone.0036204>
- Tezuka, T., Kuboyama, T., Matsuda, T., & Marubashi, W. (2007). Possible involvement of genes on the Q chromosome of *Nicotiana tabacum* in expression of hybrid lethality and programmed cell death during interspecific hybridization to *Nicotiana debneyi*. *Planta*, 226, 753–764.
- Tezuka, T., Kitamura, N., Imagawa, S., Hasegawa, A., et al. (2021a). Genetic Mapping of the HLA1 locus causing hybrid lethality in *Nicotiana* interspecific hybrids. *Plants*, 10, 2062. <https://doi.org/10.3390/plants10102062>
- Tezuka, T., Kitamura, N., Yanase, M., & Morikawa, T. (2021b). Evaluation of crossability between *Nicotiana benthamiana* and *Nicotiana excelsior*. *Agronomy*, 11(12), 2583. <https://doi.org/10.3390/agronomy11122583>
- Unnikrishnan, M., Subhashini, U., & Satyanarayana, K. V. (1981). Tissue culture in interspecific hybridization in *Nicotiana*. *Tobacco Research*, 7, 62–67.
- Valleau, W. D. (1952). Breeding tobacco for disease resistance. *Economic Botany*, 6, 69–102.
- Wark, D. C. (1970). Development of flue cured tobacco cultivars resistant to common strain of blue mold. *Tobacco Science*, 14, 147–150.
- Wheeler, H. M. (1945). A Contribution to the Cytology of the Australian-South Pacific Species of *Nicotiana*. *Proceedings of the National Academy of Sciences of the United States of America*, 31, 177–185.
- Whitaker, T. W. (1934). The occurrence of tumors on certain *Nicotiana* hybrids. *Journal of the Arnold Arboretum*, 15, 144–153.
- White, O. E. (1914). Studies of teratological phenomena in their relation to evolution and the problems of heredity. I. A study of certain floral abnormalities in *nicotiana* and their bearing on theories of dominance. *American Journal of Botany*, 1, 23–36.

- Williams, E. (1975). A new chromosome number in the Australian species *Nicotiana cavigola* L. (Burbidge). *New Zealand Journal of Botany*, 13, 811–812.
- Williams, E., & Pandey, K. K. (1975). Meiotic chromosome pairing in interspecific hybrids of *Nicotiana*. II. South American species hybrids: The influence of genotype on pairing. *New Zealand Journal of Botany*, 13, 611.
- Wolf, P. A. (1965). Hereditary abnormalities of tobacco. *Journal of the Elisha Mitchell Scientific Society*, 81(85–92), 144–172.
- Wong, Y. N. (1975). Genome analysis of tobacco interspecific hybrids. *Taiwan Tob Wine Monop Bur Tob Res Inst Res Rep*, 2, 22–31.
- Yamada, T., Marubashi, W., & Niwa, M. (1999). Detection of four lethality types in interspecific crosses among *Nicotiana* species through the use of three rescue methods for lethality. *Breeding Science*, 49, 203–210.
- Zaitlin, D., & Mundell, R. (2006). *Nicotiana hybrids and plant varieties for use in production of pharmaceuticals*. US Patent 20060236433 A1.
- Zenktelej, M., Slusarkiewicz Jarzina, A., & Wozna, J. (1981). Cytological investigations of hybrid plants of *Nicotiana alata* and *Nicotiana debneyi* obtained by *in vitro* pollination of ovules. *Bulletin de la Société des Amis des Sciences et des Lettres de Poznan, Serie D*, 21, 79–84.
- Zhukov, N. I. (1939). Hybrids of *Nicotiana rustica* L. x *N. alata* Link et Otto (in Russian). *Sb Nauch Issled Rabot WITIM*, 139, 236–244.