

Jos Houbraken and Paul S. Dyer

## 2.1 Introduction

The vast majority of fungal species are able to undergo sexual reproduction involving the formation of sexual spores via meiosis. This form of reproduction is thought to have many evolutionary advantages, and where present offers a valuable laboratory tool for experimental genetic analysis (Dyer and Paoletti 2005; Aanen and Hoekstra 2007; Lee et al. 2010). However, a surprisingly high minority of fungal species (approximately 20 %) are only known to reproduce by asexual (mitotic) means. This includes many species of industrial importance, notably several *Aspergillus* and *Penicillium* species (Dyer and Paoletti 2005; Dyer and O’Gorman 2011). The aim of this chapter is twofold. Firstly, to describe how sexual cycles may be induced in filamentous fungi including ‘fastidious’ species (Kwon-Chung and Sugui 2009), which require very specific conditions, and sexually ‘recalcitrant’ species, where a sexual cycle might not yet have been reported. Secondly,

to describe methods by which sexual progeny can be isolated, with further possibilities suggested for progeny analysis. The chapter will focus on filamentous ascomycete species (*Pezizomycotina*), which represent one of the largest groups in the fungal kingdom, and in particular on members of industrial importance. A final section is included to briefly describe ways in which the sexual cycle can be exploited for purposes including gene identification and localization, strain improvement, and gene complementation.

## 2.2 Sexual Reproduction and Breeding Systems in Filamentous Ascomycete Fungi

In 1820, microscopic sexual structures in fungi were reported for the first time in a culture of *Syzygites megalocarpus* (*Zygomycota*, *Mucorales*) (Ehrenberg 1820; Idnurm 2011). In 1904, Blakeslee showed that *S. megalocarpus* is a self-fertile species and also observed the existence of different “sexes” (‘mating types’) in *Rhizopus stolonifer* (reported as *Rhizopus nigricans*). Based on his findings he introduced the terms ‘homothallism’ for self-fertile (or self-compatible) and ‘heterothallism’ for self-incompatible (or obligate outcrossing) individuals (Blakeslee 1904). Thus, by definition, individuals of homothallic fungal species can complete the sexual cycle without the

---

J. Houbraken, Ph.D. (✉)  
Department of Applied and Industrial Mycology,  
CBS-KNAW Fungal Biodiversity Centre,  
Uppsalalaan 8, 3584 CT Utrecht, The Netherlands  
e-mail: [j.houbraken@cbs.knaw.nl](mailto:j.houbraken@cbs.knaw.nl)

P.S. Dyer, B.A., M.A., Ph.D.  
School of Life Sciences, University of Nottingham,  
Nottingham, NG7 2RD, UK  
e-mail: [Paul.Dyer@nottingham.ac.uk](mailto:Paul.Dyer@nottingham.ac.uk)

need for a mating partner, whereas individuals of heterothallic species require a mating partner of compatible mating type for sexual reproduction to occur. However, it is important to note that homothallic species are not restricted to self-fertility, as individuals normally retain the ability to out-cross under suitable conditions (Dyer et al. 1992; Burnett 2003; Cavindera and Trail 2012). In the case of heterothallic pezizomycete species there are normally only two mating types present. By convention these are now termed *MATI-1* and *MATI-2*, although for some species alternative established terminology such as *matA* and *matB* (e.g. in *Neurospora*) or plus '+' and minus '-' (e.g. in *Podospora*) are used (Dyer et al. 1992; Turgeon and Yoder 2000). Also some heterothallic species have an additional layer of sexual compatibility superimposed on the mating type. Individuals can either be male (M), female (F), or hermaphrodites (MF) with respect to their ability to form sexual mating structures such as ascogonia, protoperithecia, microconidia, and spermatia (Debuchy et al. 2010). For a cross to be successful not only must isolates of opposite mating type be present, but also one mating partner must be able to act as a male and the other as a female. However, this system appears to be restricted to certain taxonomic groupings such as *Fusarium* and *Magnaporthe* species (Gordon 1961; Takan et al. 2012). A third reproductive strategy occurring in filamentous fungi was later described, called 'pseudohomothallism' (or secondary homothallism) (Dodge 1957). Ascomycetous pseudohomothallic species (e.g. *Neurospora tetrasperma*, *Podospora anserina*) develop four spored asci in which most ascospores contain two nuclei, one of each mating type (Raju and Perkins 1994). A typical binucleate ascospore germinates to form a self-fertile mycelium due to the fact that the arising heterokaryic hyphae contains both *matA* and *matB* nuclei (Pöggeler 2001).

In heterothallic pezizomycete species sexually compatible haploid strains are normally morphologically indistinguishable and not differentiated into male and female sexes, the mating partners are instead distinguished only by their mating type (Bistis 1998). Extensive studies over the past 25 years have revealed that both sexual iden-

tity and later stages of sexual development are controlled in fungi by so called 'mating-type' (*MAT*) genes (Debuchy et al. 2010). In heterothallic pezizomycete fungi there is usually only one *MAT* locus at which between one and three *MAT* genes may be present. The DNA sequence of the *MAT* genes of isolates of opposite mating type, together with other non-coding sequence present at the *MAT* locus, is highly dissimilar although the regions flanking the *MAT* locus are highly conserved (Debuchy and Turgeon 2006). Because of the dissimilarity in sequence between the opposite *MAT* loci, they are referred to as 'idiomorphs' instead of alleles to emphasize that the sequences at the same locus are highly dissimilar (Metzenberg and Glass 1990). By convention, *MATI-1* mating-type isolates contain a *MAT* idiomorph which includes a *MATI-1* gene encoding a protein with a motif called the alpha box, whereas *MATI-2* mating-type isolates contain a *MAT* idiomorph which includes a *MATI-2* gene encoding a regulatory protein with a DNA-binding domain of the high mobility group (HMG) family. These two idiomorphs are designated *MATI-1* and *MATI-2*, respectively. Where more than one *MAT* gene is present in an idiomorph, each gene within an idiomorph is indicated by the idiomorph symbol followed by a dash and a number, e.g., *MATI-2-1* (Turgeon and Yoder 2000). The organization of *MAT* genes differs in homothallic species, where both alpha box *MATI-1* and HMG domain *MATI-2* genes are normally present in the same individual; this co-occurrence appears to confer the ability to self-fertilize. The alpha and HMG genes can be tightly linked at a single *MAT* locus or be present at two distinct *MAT* loci within the genome (Paoletti et al. 2007; Debuchy et al. 2010).

### 2.2.1 Use of *MAT* Genes as Diagnostic Tools for Induction of a Sexual Cycle

The discovery and characterization of mating-type genes from a diverse range of fungi has provided a major advance in the ability to induce sexual cycles of chosen fungal species in vitro.

Fungal *MAT* genes were first identified using molecular methods from the yeast *Saccharomyces cerevisiae* (Astell et al. 1981) and since then have been characterized from numerous filamentous ascomycete species (Debuchy and Turgeon 2006; Debuchy et al. 2010).

Although the *MAT* genes and idiomorphs show considerable sequence divergence overall, it has nevertheless been possible to identify partly conserved alpha box and HMG domain regions of the *MAT1-1* and *MAT1-2* genes, respectively, encoding homologous 65–80 amino acid regions of the MAT proteins. This has allowed the design of degenerate PCR primers [which can include the alternative base inosine (I) to avoid too high a rate of degeneracy] that can be used to amplify these regions of the *MAT* genes from species where genome sequence data is lacking. Due to sequence divergence within the Pezizomycotina, degenerate primers often need to be designed for groups of related fungal species because it has proved difficult to design all encompassing pezizomycete *MAT* degenerate primer sets (Dyer et al. 1995; Arie et al. 1997; Singh et al. 1999; Table 2.1). But once *MAT* amplicons have been obtained they can then be sequenced to confirm homology to known *MAT* genes, and if desired the arising sequence can be used to design specific primers (which are likely to be less prone to PCR artefacts than degenerate primers) for use as a *MAT* diagnostic tool to determine the mating type of isolates. It is noted that the *MAT1-2* gene sequence tends to be conserved to a higher extent than *MAT1-1* gene sequence, so it might be necessary to obtain *MAT1-1* idiomorph sequence via a chromosome walking approach based on inwards sequencing from the *SLA* and *APC* genes found in the conserved region bordering the *MAT* loci (Eagle 2009, C Eagle and PS Dyer unpublished results) or use of TAIL-PCR to amplify whole *MAT* idiomorph regions (Arie et al. 1997). More recently it has also been possible to use whole genome sequence to identify *MAT* genes by BLAST analysis and thereby design *MAT* diagnostic primers set directly. Such *MAT* diagnostic tests may use different primer pairs for the *MAT1-1* or *MAT1-2* genes, necessitating two rounds of PCR (see examples in Table 2.1).

Alternatively, multiplex PCR-based tests have been designed to allow mating type to be determined using a single PCR (Table 2.1). The latter diagnostic relies on the use of one primer binding to conserved sequence in the flanking regions of isolates of both mating type, together with two further primers that bind to sequence either present in the *MAT1-1* or *MAT1-2* idiomorph. The location of the latter two primers is designed such that differential size PCR products are generated according to whether isolates are of *MAT1-1* or *MAT1-2* genotype, thereby allowing the rapid and efficient determination of mating type [for further explanation of rationale see Dyer et al. (2001b) and Paoletti et al. (2005)].

The availability of a PCR-based mating-type detection method is a major aid when dealing with induction of sexual reproduction in heterothallic species. Before the introduction of these *MAT* diagnostic techniques, isolates had to be crossed in all combinations with each other and when successful mating occurred, mating-type tester strains were selected. With these tester strains, larger sets of isolates were then examined in order to determine their fertility and mating type. Whereas now with the use of *MAT*-specific primers, the mating type of isolates can be determined prior to crossing on agar media. It is therefore possible to set up, in a more efficient manner, directed crossings between isolates known to be of the opposite mating type. This significantly reduces the amount of effort needed to study for example the presence of a sexual cycle in presumed asexual species or the sexual fertility within a population. Indeed, this approach has led to major breakthroughs over recent years with the induction of a sexual cycle in a series of previously considered ‘asexual’ species when crosses were set up between known *MAT1-1* and *MAT1-2* isolates. A model example concerns the opportunistic human pathogen *Aspergillus fumigatus*. This species was described by Fresenius in 1863 and for more than 140 years was only known as an asexual organism (Samson et al. 2009). However, there was accumulating evidence for the presence of a cryptic (i.e., hidden, so far undescribed) sexual cycle. Firstly, studies of the population structure provided evidence for sexual

**Table 2.1** Degenerate and specific PCR primers used in mating-type diagnostic tests for a selection of heterothallic ascomycete species, especially in the *Eurotiales* and *Hypocreales*

Taxonomic grouping/ species	Primers (MAT1-1; 5'-3')	Primers (MAT1-2; 5'-3')	Remarks	Reference
Leotiomycete and eurotiomycete fungi	N/A	MAT 5-3 (F) AARIHCCIMGICCIIMYTAAT MAT 3-1 (R) CKIGGHIIRTAIYKRTAIIHGG	Degenerate primer pair used successfully in conjunction with hot start PCR to amplify <i>MAT1-2</i> sequence from leotiomycete and eurotiomycetes species.	Dyer et al. (1995), Singh et al. (1999), Paoletti et al. (2005)
Eurotiomycete and lecanoromycete fungi	N/A	MAT5-4 (F) AARRTICCIMGICCIICIAAYGC MAT3-2 (R) TTNCKIGGIGTRTAITGRTARTCNGG	Degenerate primer pair used successfully with hot start PCR to amplify <i>MAT1-2</i> sequence from eurotiomycete and lecanoromycete species.	Seymour et al. (2005)
Eurotiomycete fungi	MAT5-6 (F) GIMGICCIYTAAAYWSITTYATHGC MAT3-4 (R) ARRAAICKIARIATICCSWYTT	MAT5-7 (F) THSCIMGICCIICIAAYKSITTYAT MAT3-5 (R) TTICKIGGIIKRWAYIKRTARTYNGG	Degenerate PCR primer pairs used successfully with hot-start PCR to amplify <i>MAT</i> sequence from a range of <i>Aspergillus</i> and <i>Penicillium</i> species.	Eagle (2009), C Eagle and PS Dyer (unpublished results)
Loculoascomycete fungi	N/A	ChHMGI (F) AAGGCNCCNGYCCNATGAAC ChHMG2 (R) CTNNGGNGTGATYTTGTAATTNGG	Degenerate PCR primer pairs used successfully to amplify <i>MAT1-2</i> sequence from a range of loculoascomycete species.	Arite et al. (1997)

Pyrenomycete fungi	N/A	NcHMG1 (F) CCYCGYCCYCCYAAAYGCNTAYAT NcHMG2 (R) CGNGGRTTRTARCCGRRTNRGG	Degenerate PCR primer pairs used successfully to amplify <i>MAT1-2</i> sequence from a range of pyrenomycete species.	Arie et al. (1997)
<i>Aspergillus felis</i>	AFM1_F65655 (F) CCTYGACGMGATGGGTTGG	MAT2_F6086 (F) TCGACAAGATCAAAWICYCGTC	Annealing temperature 48 °C.	Barrs et al. (2013)
	MAT1_R6215 (R) TGTCAAAGARTCCAAAAGGAGG	MAT2_R6580 (R) CTTYTTGARCTCTTCYCGTACG		
	M1F (F) ATTGCCAATTTGGCCTTGAA M1R (R) TTGATGACCAATGCCACCAGA	M2F (F) GCATTCATCCTTATCGTCAGC M2R (R) GCTTCTTTTCGGATGGCTTGCGG	Multiplex.	Ramirez-Prado et al. (2008), Horn et al. (2011)
<i>Aspergillus fumigatus</i> , <i>A. lentulus</i>	AFM1 (F) CCTTGACGGATGGGGTGG AFM3 (R) CGGAAATCTGATGTCGCCACG	AFM2 (F) CGCTCCTCATCAGAAACAACCTCG AFM3 (R) CGGAAATCTGATGTCGCCACG	Multiplex; AFM3 is “common” primer and binds in the flanking region bordering both idiomorphs.	Paoletti et al. (2005), Swilaiman et al. (2013)
	AteM1F (F) GCGAGGCAGACATTCAGGAT AteM1R (R) CGAGGATGCCAATAAAACCAGC	AteM2F (F) TCTATCGGCAGCACCATCATCC AteM2R (R) CTTGTTGTGGTGGTTCGTTCT	Annealing temperature MAT1-1: 53.5 °C; MAT1-2: 55 °C.	Eagle (2009); C Eagle and P.S. Dyer (unpublished results)
<i>Aspergillus tubingensis</i>	M1F_Anig (F) GGTATCGCGAATGATGGAG M1R_Anig (R) CAGCGTCTTCAACGCATTC	MAT5-4 (F) AARRTICMIGICICICIAAYGC MAT3-2 (R) TTNCKIGIGITRITAITGRFARTCNGG	Annealing temperature MAT1-1: 62 °C; MAT1-2: 55 °C.	Horn et al. (2013), Rydholm et al. (2007)
	alpha1 (F) CTGGAGGAGCTTCTGCAGTAC alpha2 (R) GGAGTACGCCCTTCGCCGAG	HMG1 (F) CTCTTGTGGCAGGATGCTCT HMG2 (R) TTGCTGGTAGAGGGCAGTCT		Sugui et al. (2010)

(continued)

Table 2.1 (continued)

Taxonomic grouping/ species	Primers (MAT1-1; 5'-3')	Primers (MAT1-2; 5'-3')	Remarks	Reference
<i>Paecilomyces variotii</i>	MAT1-F1-VarSp (F) TATGCCCTCCTGGTGAGCTGG MAT1-R2-VarMar (R) GATCCCRGAYTTSGYCTTCTG	MAT2-F1Paec (F) AYCAYCAYCCCKATYGTCAAAGC MAT2-R1Paec (R) GYTTGCGYTTTATCTCTCTCYGC	Multiplex; annealing 58 °C.	Houbraken et al. (2008)
<i>Penicillium rubens</i> (reported as <i>P. chrysogenum</i> )	MAT-1-f (F) CTTCGTCCAITGAACTCTTTTATG MAT-1-r (R) ATCCCAAC CAGCCATCCTGAGAT	MAT-2-f (F) CCAAGT CTATCCACGAGGCTG MAT-2-r (R) GCAGGCAGTTGGCACGGGAAC		Hoff et al. (2008)
<i>Talaromyces derxii</i>	MAT 1-1b (F) CCACGTATAACGGGGCATC (R) CGGCTTGCCAMAGGTCTT	MAT 1-2b (F) GTGATAATGCTTSCGATAGAGAATG (R) GTTGGAGAGGAGGCGTTGAC	Touchdown 60–50 °C.	López-Villavicencio et al. (2010), Y. Yilmaz (personal communication)
<i>Trichoderma reesei</i>	MatA2-fw (F) CTCGAGAGGGATATACACCAG MatA2-rv (R) CTTCTACACGGATGCCAGA	Mat2fw (F) CAACACGTATGAAAGAGAGATG Mat2rv (R) ATTGGAACGGATCACCTTCTTG		Seidl et al. (2009)
<i>Fusarium keratoplasticum</i>	FS3MAT1-1 (F) ATGGCTTTCCGCAGTAAAGGA FS3MAT1-1 (R) CATGATAGGGCAGCAAAGAG	FS3MAT1-2 (F) GGGAATCTGAGAAAAGATACGTAC FS3MAT1-2 (R) CGGTACTGGTAGTCGGGGAT		Short et al. (2013)
<i>Fusarium culmorum</i>	FusALPHAfor (F) CGCCCTCKAAYGSCTTCATG FusALPHArev (R) GGARTARACYTTTAGCAATYAGGGC	FusHMGfor (F) CGACCTCCCAAAGCYTACAT FusHMGrev (R) TGGGCGGTACTGGTARTCRGG	Multiplex; annealing 55 °C.	Kerényi et al. (2004)
<i>Fusarium fujikuroi</i>	GFmat1a (F) GTTCAATCAAAGGGCAAGCG GFmat1b (R) TAAGCGCCCTTAAACGCCCTTC	GFmat2d (F) CTACGTTGAGAGCTGTACAG GFmat2c (R) AGCGTCAATTATTCGATCAAG		Steenkamp et al. (2000)

recombination based on the analysis of the association of alleles of five loci (Pringle et al. 2005) and sequence present at three intragenic regions (Paoletti et al. 2005). Secondly, it was possible to identify *MATI-1* and *MATI-2* isolates of *A. fumigatus* using genomic BLAST searching and degenerate PCR approaches, together with the presence of a series of genes related to sex within the genome (Galagan et al. 2005; Paoletti et al. 2005). A multiplex PCR-based *MAT* diagnostic test was then developed, and analysis of 290 worldwide clinical and environmental isolates revealed the presence of *MATI-1* and *MATI-2* genotypes in similar proportions (43 % and 57 %, respectively) (Paoletti et al. 2005). The presence of the two mating types in equal frequencies within a population is an indication of sexual reproduction (Milgroom 1996). In a subsequent study, analysis of a population of *A. fumigatus* strains from five locations in Dublin, Ireland, revealed an almost exact 1:1 ratio of *MATI-1*:*MATI-2* isolates (O’Gorman et al. 2009). Furthermore, 88 out of 91 isolates were genetically unique according to a RAPD-PCR DNA fingerprinting study, and a phylogenetic analysis demonstrated that the *MAT-1* and *MATI-2* isolates were interleaved when represented on a phylogenetic tree. This provided strong evidence for recent or extant sexual reproduction and led O’Gorman et al. (2009) to set up directed crosses between known *MATI-1* and *MATI-2* isolates on a range of media under a variety of different growth conditions. An exciting result was then obtained when it was found that a sexual cycle, leading to production of cleistothecia containing recombinant ascospores, could be induced when cultures were crossed on oatmeal agar in Parafilm-sealed Petri-dishes which were incubated at 30 °C in darkness for 6–12 months (O’Gorman et al. 2009). More recently, a “super-mater” pair of *A. fumigatus* isolates have been identified which produce abundant cleistothecia containing viable ascospores after only 4 weeks incubation under the same conditions (Sugui et al. 2011). There have subsequently been further examples of the discovery of sexual states in other supposedly ‘asexual’ fungal species using similar directed crosses between known *MATI-1*

and *MATI-2* isolates [reviewed by Dyer and O’Gorman (2012)]. These have included notably the description of a sexual state for the industrial workhorse *Trichoderma reesei* (Seidl et al. 2009), the penicillin producer *Penicillium chrysogenum* (Böhm et al. 2013), the opportunistic pathogen *Aspergillus lentulus* (Swilaiman et al. 2013), the aflatoxin producers *Aspergillus flavus* and *Aspergillus parasiticus* (Moore 2014), and the starter culture of blue veined cheese *Penicillium roqueforti* (Ropars et al. 2014; S Swilaiman, J Houbraeken, J Frisvad, R Samson and PS Dyer, unpublished results).

It should be cautioned that the presence of the *MAT* genes alone is insufficient to prove that a sexual stage exists. Given that several hundred other genes are also likely to be required for a functional sexual cycle to occur, it is possible that loss of function of any of these genes could result in reduced fertility or asexuality. For example, a series of over 75 genes have been identified which are required for sexual reproduction in the aspergilli, encompassing processes such as environmental sensing, mating, fruit body formation, and ascospore production. Such ‘sex-related’ genes might be predicted to accumulate deleterious mutations or even be lost in purely asexual species, in which there was no functional constraint on their conservation (Dyer 2007; Dyer and O’Gorman 2012).

---

### 2.3 Methods to Induce Sexual Reproduction in Filamentous Fungi

In this next section methods will be described to induce sexual reproduction in filamentous ascomycete species based on procedures that have been used to obtain successful mating in a variety of fungal species. The first step involves the selection of suitable strains; this is followed by *MAT* analysis; then selection of suitable agar media, inoculation procedures, and incubation conditions to induce a sexual cycle. Finally after formation of a sexual state, single ascospore isolates should be obtained and these should be examined for evidence of recombination. It is



noted that sexual reproduction for many homothallic species can be achieved fairly readily as there is no need for any mating step, so most of the following discussion will apply to heterothallic species which can be more demanding in their sexual requirements.

## 2.4 Materials

### 2.4.1 Solutions

Sterile distilled water or tap water.

Tween 80: 0.5 g/L, ddH<sub>2</sub>O to 1 L.

### 2.4.2 Agar Media Inducing Sexual Reproduction

Carrot agar (CA): Fresh washed, peeled, diced carrots (400 g) in 400 mL ddH<sub>2</sub>O. Autoclave at 121 °C for 15 min. After autoclaving, blend the carrots and add additional 500 mL ddH<sub>2</sub>O. Add ZnSO<sub>4</sub>·7H<sub>2</sub>O (0.01 g/L), CuSO<sub>4</sub>·5H<sub>2</sub>O (0.005 g/L), agar (20 g/L). Mix well and autoclave at 121 °C for 15 min.

Mixed cereal agar (MCA): Gerber mixed grain cereal (Gerber Products Co., Fremont, Michigan) (50 g/L), agar (20 g/L), in 1 L ddH<sub>2</sub>O. Mix well and autoclave at 121 °C for 15 min (McAlpin and Wicklow 2005).

Oatmeal agar (OA)<sup>1</sup>:

Version (1): Blend 30 g of oats and add 1 L ddH<sub>2</sub>O. Boil and let it stand for 1 h. Add ZnSO<sub>4</sub>·7H<sub>2</sub>O (0.01 g/L), CuSO<sub>4</sub>·5H<sub>2</sub>O (0.005 g/L), agar (20 g/L). Autoclave for 15 min, at 121 °C (Samson et al. 2010).

Version (2): Add 40 g of oats to 1 L tap water. Bring to the boil then lower the heat to just below

boiling point (i.e., bubbling gently) for a further 45 min. Then filter through two layers of cheese cloth and restore the volume of the solution to 1 L with tap water and mix thoroughly. Add agar (20 g/L). Autoclave for 15 min at 121 °C with slow cool down and slow release of pressure to prevent media loss (O’Gorman et al. 2009).

Potato dextrose agar (PDA): 200 g sliced potatoes are boiled in 1 L of ddH<sub>2</sub>O and sieved, add glucose (20 g/L), agar (20 g/L), ZnSO<sub>4</sub>·7H<sub>2</sub>O (0.01 g/L), CuSO<sub>4</sub>·5H<sub>2</sub>O (0.005 g/L). pH is approximately 5.6. Autoclave for 15 min, at 121 °C (Samson et al. 2010).

Tap water agar (TWA)<sup>2</sup>: Bacto™ agar (15 g/L) in 1 L tap water. Autoclave for 15 min, at 121 °C. Supplement with appropriate natural growth substrate (Dyer et al. 1993).

V8 agar (V8): V8® vegetable juice (Campbell) (175 mL/L), CaCO<sub>3</sub> (3 g/L), ZnSO<sub>4</sub>·7H<sub>2</sub>O (0.01 g/L), CuSO<sub>4</sub>·5H<sub>2</sub>O (0.005 g/L), agar (20 g/L) in 1 L ddH<sub>2</sub>O. Mix well and autoclave at 121 °C for 15 min (Samson et al. 2010).

## 2.5 Methods

### 2.5.1 Strain Selection

#### 2.5.1.1 Identification

The first step in inducing a sexual cycle is the requirement for the correct identification of isolates of the same biological species. In the past, fungal taxonomy was primary based on phenotypic and physiological characters. Nowadays molecular techniques like DNA sequencing are commonly applied for identification purposes. Such data has shown that many well-known ‘species’ are actually species complexes composed of closely related species that might be sexually fertile when crossing within a species, but sexually sterile when attempts are made to cross different species. For example, *P. chrysogenum* is a complex of five species, namely *P. chrysogenum sensu stricto*, *Penicillium rubens*, *Penicillium*

<sup>1</sup>Commercially made oatmeal agar (OA) is available from certain manufacturers, but in our experience this is not able to induce sex in demanding species. Instead, it is best to prepare OA in house. Different brands of oats can be used and these have can have an effect on the mating. Commonly used brands are Pinhead oatmeal (Odlums, Ireland) and Quaker Oats. For *Sordaria* and *Chaetomium* species and *Penicillium rubens* (*P. chrysogenum*) this medium needs to be supplemented with biotin (6.4 µg/L) to induce sex (Böhm et al. 2013).

<sup>2</sup>In our experience Bacto™ agar is less prone to condensation problems than some cheaper, less pure, commercial agars.



*allii-sativi*, *Penicillium tardochrysogenum*, and *Penicillium vanluykii* (Henk et al. 2011; Houbraken et al. 2012). Using these current taxonomic insights, the main penicillin producer is named *P. rubens* and in the study of Böhm et al. (2013) all *P. chrysogenum* strains that were able to reproduce sexually are actually *P. rubens*. The induction of the sexual cycle would have probably been much more difficult or even impossible if attempts had been made to cross different members of the *P. chrysogenum sensu lato* complex and might explain why the sexual stage had remained undiscovered up to that point. Similarly, it was discovered that the ‘single’ anamorphic species *Pseudocercospora herpotrichoides* is composed of two closely related, but intersterile, species *Tapesia (Oculimacula) yallundae* and *Tapesia acufiformis* (Dyer et al. 1996). Again, attempts to induce a sexual cycle by erroneously crossing isolates of the different species would have failed in this case.

Therefore, it is essential to verify that isolates to be used in crossing experiments are of the same biological species. The internal transcribed spacer regions (ITS) of the ribosomal gene have been used in many taxonomic studies for species identification and have been selected as fungal barcodes (Schoch et al. 2012). The main advantages of using the ITS locus for identification are the ease of amplification by PCR and the presence of a high number of ITS sequences in the public databases. However, the resolution of this locus is insufficient for species identification of all fungi. For example, closely related species belonging to the industrially important genera *Aspergillus*, *Penicillium*, *Fusarium*, and *Trichoderma* can share the same ITS barcode. In those cases, it is recommended to sequence other (protein coding) genes. There is no consensus which region to sequence and the choice largely depends on the genus/species being dealt with. Details on molecular and phylogenetic identification methods can be found in Crous et al. (2009).

### 2.5.1.2 Origin of Strains

Freshly isolated strains are generally more fertile than strains maintained for longer periods in culture collections, which can be prone to a ‘slow

decline’ in fertility following prolonged subculture (Dyer and Paoletti 2005). For example, *Paecilomyces variotii* strains isolated from heat-treated products proved to be fertile, while older isolates from a culture collection were unable to mate (Houbraken et al. 2008). Similar observations were found in the heterothallic *Histoplasma capsulatum*. This species lost fertility during laboratory passage and it was suggested that selective pressures may serve to maintain fertility in the environment (Kwong-Chung et al. 1974; Fraser et al. 2007). Furthermore, even when obtaining isolates from the field it is important to be aware that such isolates can exhibit a range of fertility due to various physiological and genetic factors (Dyer et al. 1992). Indeed the same ‘slow decline’ in sexual fertility observed during in vitro culture may be occurring in vivo in natural populations subject to strong selection pressure favoring asexual propagation (Dyer and Paoletti 2005). For example, Sugui et al. (2010) found that most attempted crosses involving the emerging agent of aspergillosis *Aspergillus (Neosartorya) udagawae* either failed to produce cleistothecia or produced ascospores which did not germinate. Similarly, Swilaiman et al. (2013) found that many clinical isolates of the opportunistic pathogen *A. lentulus* exhibited low fertility or were sterile in crosses. However, in both of these cases it was possible to detect isolates that successfully crossed to form cleistothecia with ascospores. This illustrates the fact that it is very important to select a number of representative field isolates for crossing studies to ensure that at least some representatives will exhibit sexual fertility if it is present.

### 2.5.1.3 Strain Typing

Before the start of the mating experiments, the isolates should ideally be typed using methods such as AFLP (Amplified Fragment Length Polymorphism), SSR (microsatellites or simple sequence repeats), RFLP (Restriction Fragment Length Polymorphism), RAPD (Random Amplified Polymorphic DNA), or MLST (MultiLocus Sequence Typing) DNA fingerprinting. The use of such typing methods allows the selection of independent (non-

clonal) strains for crossing purposes, avoiding the error of setting up repeated crosses with different isolates of the same field strain. Typing can also generate insights as to whether there is evidence of recombination among the strains.

### 2.5.2 Detection of MAT Genes

In order to increase the likelihood of success and reduce the number of crosses that need to be set up, it is recommended that the mating type of test strains be determined prior to crossing efforts. This enables directed crosses to be set up on agar media (see below) between *MATI-1* and *MATI-2* mating partners that are known to be potentially sexually compatible. For certain species or species groups, mating type-specific primers have been developed that amplify part of either the *MATI-1* or *MATI-2* gene. A selection of primers pairs already published for some important pezizomycete species is given in Table 2.1 together with details of some degenerate primers pairs that should be more broadly applicable to wide groups of species (Dyer et al. 1995; Arie et al. 1997; Singh et al. 1999; Paoletti et al. 2005; Seymour et al. 2005). Especially note that the use of hot-start PCR can greatly increase the chances of success; for example, Singh et al. (1999) were unable to amplify a *MATI-2* region from *T. yallundae* using standard PCR, but found that the use of hot-start PCR gave very strong amplification of the required product. In cases where no *MAT* amplicons are obtained, it is recommended that new degenerate primers are designed based on known *MAT* gene sequence of species that are phylogenetically closely related to the test species.

### 2.5.3 Agar Media

A large variety of agar media have been used for the induction of fungal sexual cycles. Generally, media based on natural substrates are more effective than synthetic media and which agar to use largely depends on the species or genus. Some species are fastidious and need specific nutrients, which often mimic their natural growth substrate.

For example, the sexual cycle of the cereal pathogen *T. yallundae* occurs in nature on straw stubble left after harvest (Dyer et al. 2001a) and attempts to induce the sexual cycle in vitro on a range of synthetic media failed. However, it was possible to induce the sexual cycle when *MATI-1* and *MATI-2* isolates were inoculated onto straw segments (especially those with nodes), which were kept moist by being placed on TWA (Dyer et al. 1993). Similarly, it was only possible to induce sex in *Thermomyces dupontii* (= *Talaromyces thermophilus*) on natural oat grains rather than synthetic agar media (Pitt 1979; Houbraken et al. 2014), and three dermatophytic *Trichophyton* species required growth on sterilized baby or rabbit hair (placed on agar) to induce sexual reproduction (Kawasaki et al. 2010). By contrast, for species such as *Neurospora crassa* it is possible to induce sex on fully synthetic media, one reason why this is used as a model organism (Perkins 1986). The diversity in nutrient requirements for sexual reproduction is illustrated well by members of the genus *Aspergillus*. Species such as the homothallic *Aspergillus nidulans* and *Aspergillus fischeri* reproduce sexually on a fairly wide range of media, including fully synthetic complete media and oatmeal agar (OA) (Paoletti et al. 2007). This contrasts with some heterothallic species which have more exacting demands. Members of the section *Flavi* (e.g. *A. flavus*, *A. parasiticus*) have only been successfully crossed on mixed cereal agar (MCA), while crosses of members of the section *Fumigati* (*A. fumigatus*, *A. lentulus*) have only proved fruitful on OA. In contrast to these two high water activity media, species belonging to section *Aspergillus* (Eurotium-type ascomata) require a low water activity medium (e.g., malt extract agar with 40% sucrose) for fruiting body formation (Dyer and O’Gorman 2012). In *Fusarium*, the standard medium to induce fruiting bodies, called perithecia, is carrot agar. In contrast, attempts with *Fusarium keratoplasticum* to use this standard medium were unsuccessful while crosses on V8 agar induced the sexual cycle in this species (Covert et al. 2007; Short et al. 2013). In the case of homothallic *Giberella zeae* sex can be induced by the gentle removal of surface mycelia, fol-

lowed by treatment with detergent solution (Cavindera and Trail 2012). One intriguing example is that of different *Cryptococcus* species. Nielsen et al. (2007) found that it was possible to induce the sexual cycle of *Cryptococcus neoformans* on pigeon guano media, but that this was not possible for the related species *Cryptococcus gattii*. It was suggested that the ability to undergo sexual reproduction on pigeon guano represented an evolutionary adaptation that allowed ancestral strains of *C. neoformans* to sweep the globe (Nielsen et al. 2007; Heitman et al. 2014). It is important to note that when making agar media there can be difference among ingredients of different suppliers. For example, different brands of yeast extracts are available and these can have a strong influence on the phenotype of the culture. Furthermore, agar media based on natural ingredients can vary between manufactures and labs, and even within one lab batch to batch differences can occur.

Some ascomycetes may require exogenous vitamins, minerals, or other natural materials for ascomata (ascocarp) production, and these are often not present in synthetic media. This might be one of the explanations why sexual reproduction is more often found on media made from natural substrates such as oatmeal, (mixed) cereals, and cornmeal agar. A fractionation study of V8 juice agar revealed that no single factor was responsible for its utility in inducing sex in *C. neoformans*, but rather the unique composition of V8 juice provided sustenance for sex, especially the copper content (Kent et al. 2008). Other studies have found it necessary to add compounds to a standard medium to induce sex; for example biotin was added to OA to stimulate mating in *P. rubens* (Böhm et al. 2013). Meanwhile, certain nutritional auxotrophs of *A. nidulans* can require supplementation of media to ensure sexual development, e.g., tryptophan, arginine, and *riboB2* mutants are self-sterile (Dyer and O’Gorman 2012) and heterologous expression of *pyrG* can result in reduced fertility (C Scazzocchio pers. comm.; Robellet et al. 2010). An overview of agar media used in mating experiments of selected heterothallic species

belonging to either the *Eurotiales* or *Hypocreales* as representative groupings is given in Table 2.2.

Thus, the best strategy when attempting to induce sexual reproduction in vitro is to trial a range of agar media, which should include some basal media supplemented with the natural growth substrate of the species in question.

#### 2.5.4 Incubation Conditions

Besides the nutrient availability, various other environmental factors such as light, temperature, and oxygen determine the success of mating experiments. These factors are genus and in some cases also species-specific. There are numerous reports in the literature of how different environmental conditions influence fungal sexual reproduction, and in the present chapter only some representative examples can be given. For example, *Fusarium perithecia* are formed in abundance under alternating 12 h/dark and 12 h/light cycles (with both fluorescent and near ultra violet light), whereas perithecia are absent when incubated in darkness (Table 2.2). Similarly the sexual cycle of *T. acutiformis* can be induced under near UV or white light, but not in darkness (Dyer et al. 1996). By contrast, it is necessary to incubate *Aspergillus* species in darkness to trigger sexual reproduction because light preferentially induces asexual sporulation, reflecting the natural ecology of many *Aspergillus* species (Mooney and Yager 1990; Han et al. 2003; Dyer and O’Gorman 2012). Incubation temperature also has a strong influence on sexual fertility. For example, Choi et al. (2009) showed that their *Fusarium fujikuroi* strains only produced perithecia at 23 °C and none were formed at 18, 26, and 28 °C. However, in *Fusarium graminearum*, the optimal temperature was 28.5 °C and *Fusarium circinatum* perithecia were more abundantly produced at 20 °C than at 25 °C (Table 2.2) (Tschanz et al. 1976; Covert et al. 1999). Meanwhile, in *Aspergillus* and *Penicillium*, oxygen limitation can induce sexual reproduction (Dyer and O’Gorman 2012), which can be achieved by sealing Petri dishes with Parafilm (Table 2.2).

**Table 2.2** Details of media and incubation conditions required to induce sexual reproduction in a selection of heterothallic fungal species

Species	Structure ascomata	Agar medium	Incubation		Inoculation method	Additional conditions	Reference
			temperature and time	zone			
<i>Aspergillus felis</i>	Neosartorya	Oatmeal agar	30 °C, 2–4 weeks	Barrage zone	Darkness, Petri dishes	Barrs et al. (2013)	
<i>Aspergillus flavus</i>	Petromyces	Mixed cereal agar	30 °C, 6–11 months	Mixed culture	Slants, in sealed plastic bags	Horn et al. (2009a)	
<i>Aspergillus fumigatus</i>	Neosartorya	Oatmeal agar	30 °C, 6 months	Barrage zone	Darkness, Parafilm-sealed Petri dishes	O’Gorman et al. (2009)	
<i>Aspergillus heterothallicus</i>	Emericella	Oatmeal agar	Room temperature, 2–3 weeks	Barrage zone		Raper and Fennell (1965)	
<i>Aspergillus lentulus</i>	Neosartorya	Oatmeal agar	28–30 °C, 3–7 weeks	Barrage zone	Darkness, Parafilm-sealed Petri dishes	Switainman et al. (2013)	
<i>Aspergillus nomius</i>	Petromyces	Mixed cereal agar	30 °C, 5–11 months	Mixed culture	Slants, in sealed plastic bags	Horn et al. (2011)	
<i>Aspergillus parasiticus</i>	Petromyces	Mixed cereal agar	30 °C, 6–9 months	Mixed culture	Slants, in sealed plastic bags	Horn et al. (2009b, c)	
<i>Aspergillus sclerotiticarbonarius</i>	Petromyces	Oatmeal agar	25 °C, 6–11 months	Barrage zone	Darkness, Parafilm-sealed Petri dishes	Darbyshir et al. (2013)	
<i>Aspergillus terreus</i>	Fennellia	Mixed cereal agar	37 °C, up to 6 months	Barrage zone	Darkness, 56 mm Petri dishes	Arabatzis and Velegraki (2013)	
<i>Aspergillus tubingensis</i>	Petromyces	Mixed cereal agar, Czapek agar	30 °C, 5–6 months	Mixed culture	Slants, in sealed plastic bags	Horn et al. (2013)	
<i>Aspergillus wyomingensis</i> , <i>A. udagawae</i>	Neosartorya	Oatmeal agar	25–30 °C, 4–5 weeks	Barrage zone		Nováková et al. (2014)	
<i>Paecilomyces variotii</i>	Byssochlamys	Potato dextrose agar	30 °C, 6–9 weeks	Barrage zone	Darkness, Petri dishes	Houbraken et al. (2008)	
<i>Penicillium rubens</i> (reported as <i>P. chrysogenum</i> )	Eupenicillium	Oatmeal agar supplemented with biotin (6.4 µg/L)	20 °C; 5 weeks	Barrage zone	Darkness, Parafilm-sealed Petri dishes	Böhm et al. (2013)	
<i>Talaromyces derxii</i>	Talaromyces	Oatmeal agar, malt extract agar	37 °C, 7–10 days	Barrage zone	Darkness	Takada and Udagawa (1988)	
<i>Trichoderma reesei</i>	Hypocrea	Optimal: Malt extract agar (3 % wt/vol; Merck); also potato dextrose agar	Optimal 20–22 °C; 7–10 days	Barrage zone	12-h light–dark cycle or daylight	Seidl et al. (2009)	

<i>Fusarium keratoplasticum</i>	Nectria	V-8 agar	Room temperature; 8 weeks	Barrage zone	Parafilm sealed; under 12 h cycles of direct fluorescent and UV light from 120 V bulbs	Short et al. (2013)
<i>Fusarium tucumaniae</i>	Nectria	Carrot agar	18 °C, 3–4 weeks	Barrage zone	Cool white, fluorescent bulbs	Covert et al. (2007)
<i>Fusarium fujikuroi</i>	Gibberella	V-8 agar, Carrot agar	23 °C	Fertilization	Alternating 12 h/light and 12 h/dark cycles with both FL/NUV and NUV light <sup>a</sup>	Choi et al. (2009)

A diverse range of conditions are evident even though species belonging to the genera *Aspergillus*, *Penicillium*, *Paecilomyces* and *Talaromyces* (members of *Eurotiales*), and *Fusarium* and *Trichoderma* species (both order *Hypocreales*), are phylogenetically related within their respective groupings

<sup>a</sup>Fluorescent (FL) and near ultra violet (NUV) light

Thus, the best strategy when attempting to induce sexual reproduction in vitro is to trial a range of growth conditions, which ideally might mimic those encountered in the wild when sexual reproduction occurs. This approach was used to induce sex in plant pathogenic *Tapesia* species which were known to sexually reproduce in the field in early spring in the UK on exposed straw stubble. It was found that incubation in vitro at low temperatures between 7 and 10 °C under white light could induce sex, but the sexual cycle was inhibited above these temperatures (Dyer et al. 1996).

## 2.5.5 Inoculations

For homothallic species, self fertilization can be induced using either point inoculation or spore spread methods (e.g. Paoletti et al. 2007; Todd et al. 2007; Cavindera and Trail 2012). In the case of heterothallic species, several methods have been described in the literature for crossing on agar media. We have summarized three different methods below. In the first, the “barrage zone” method, strains of opposite mating type are inoculated in close proximity on an agar medium. During incubation, the strains grow towards each other. Fruiting body formation then mainly occurs in the barrage zone (Fig. 2.1a, b), but sometimes also towards the centre or on the opposite periphery of the colony (Fig. 2.1c). Interestingly, this method can also be used to promote outcrossing in homothallic species, especially when using complementary auxotrophic strains [see Todd et al. (2007) and Cavindera and Trail (2012) for further details]. In the second, the “mixed culture” method, spore suspensions of the same concentration are prepared for isolates of opposite mating type. These suspensions are mixed together and then used to inoculate the agar medium. This results in intermingled growth of both partners from an early stage allowing close sexual interaction. Finally, in *Fusarium*, *Neurospora*, and other pyrenomycete fungi, a third “fertilization” crossing technique is also

sometimes applied. In this method, a strain of one mating type is cultured on an agar medium to allow development of protoperithecia. After growth, the culture is then fertilized (so-called ‘spermatization’) with a spore-suspension of the opposite mating partner. This fertilization method can also be used in heterothallic *Botrytis* species that produce sclerotia, which can be spermatized by being soaked with conidia of the opposite mating type (Faretra et al. 1988). Crosses with strains of the same mating type should be used as controls.

### 2.5.5.1 Barrage Zone Method

Cultures of opposite mating types are crossed in Petri dishes containing an agar medium inducing recombination. The following protocol is based on O’Gorman et al. (2009).


1. Prepare single spore isolates of each isolate and incubate under conditions inducing sporulation.
2. Harvest the spores of each isolate in sterile water containing 0.05 % Tween 80.
3. Inoculate 1.0–2.5 µL of each spore suspension (e.g., containing 500 spores) onto the agar surface about 4 cm apart and perpendicular to aliquots of spore-suspensions of the opposite mating type. This configuration created four interaction/barrage zones as colonies grew.
4. Seal, if required, with Petri dishes with Parafilm and incubate at conditions inducing recombination.
5. Regularly check the cultures on the production of fruiting bodies.

### 2.5.5.2 Mixed Culture Method

Agar slants are used in the protocol mentioned below and this method is useful when the agar medium needs be incubated for a long time. However, this method can be adopted for agar plates as well. In that case the agar media can be sealed with Parafilm in order to induce fruiting body formation and prevent drying out of the plates. The following protocol is based on Horn et al. (2009c).

1. Grow the fungal strains under conditions inducing sporulation.





**Fig. 2.1** Mating experiments between *P. variotii* strains. (a–c) Detail image of a potato dextrose agar plate where two *P. variotii* strains of opposite mating types were inoculated on each side of the Petri-dish. In a and b, ascomata are formed in the *middle*, in c, ascomata are also present on the opposite side of the colony. (d, e) Higher-

magnification view of the ascomata. (f) Micrograph of the ascomata showing asci with ascospores. (g) Similar as f, but also showing the presence of heat-sensitive conidia (arrows); isolation of the ascospores, which are heat-resistant, can be achieved by applying the “heat treatment method”. Scale bars = 10  $\mu$ m

- Harvest the spores in sterile water containing 0.05 % Tween 80.
- Dilute or concentrate (e.g., by centrifugation) the spore suspensions until a concentration of  $5 \times 10^5$  spores per mL is obtained.
- Mix the spore suspensions of strains of opposite mating types. Spore suspensions of single isolates can be used as negative controls.
- Inoculate the slants with 10  $\mu$ L spore suspensions on a medium inducing a sexual cycle (production of sclerotia).
- Incubate the slants with loose caps until sclerotia are produced.
- Enclose the caps of the slants and enclose the slants in sealed plastic bags to prevent desiccation.



8. Allow prolonged incubation and regularly check the cultures for the production of fruiting bodies.

### 2.5.5.3 Fertilization Method

The fertilization method below is derived from standard protocols used for *Fusarium* species (Klittich and Leslie 1988).

1. Grow the fungal strains of one mating type under conditions inducing sporulation.
2. Harvest the spores in sterile water containing 0.05 % Tween 80.
3. Spread the cultures of one mating type on a suitable agar media.
4. Incubate the agar plates until the agar is covered with fungal growth. Plates can be checked for evidence of ascogonia and protoperithecial formation.
5. Fertilize the cultures by dispensing 1 mL of spore-suspension in 0.05 % Tween 80 carrying at least  $5 \times 10^5$  conidia from the opposite strains.
6. Work the spore suspension into the mycelia with a glass rod until the suspension is absorbed. Self-fertilizations can be made by substituting sterilized 0.05 % Tween 80 solution for the spore suspension. Any excess conidial suspension can be removed using a sterile Pasteur pipette.
7. Seal, if required, with Petri dishes with Parafilm and incubate at conditions inducing recombination.
8. Regularly check the cultures for the production of fruiting bodies.

## 2.5.6 Single Ascospore Cultures

In heterothallic ascomycete species, a successful mating experiment will lead to production of ascomata (ascocarps) (Fig. 2.1d, e) containing asci and ascospores (Fig. 2.1f, g). Single ascospore cultures then need to be obtained for further analysis. The method of isolation of the ascospores depends on the way the ascospores are produced (e.g., in a closed ascoma or “open” perithecium) and other features of the species in question (e.g., degree of heat resistance). Four commonly used methods are described below.

### 2.5.6.1 Direct Isolation

The easiest method to obtain single ascospores cultures is by transferring ascospores directly from the fruiting body. This method can be applied in, e.g. *Fusarium*, where ascospores are produced in a perithecium and ooze out to form a prominent spore mass. This method can also be adapted for closed fruiting bodies as described by Todd et al. (2007) for *A. nidulans*.

1. Examine the agar plates using a stereomicroscope and isolate ascospores direct from a spore mass if present with a sharp needle. If necessary suspend the ascospores in 0.01 % Tween 20 or 0.05 % Tween 80 solution. In the case of closed fruiting bodies, these should be individually picked up and rolled across the surface of 4 % TWA plates to remove adhering conidia and hyphae [see Todd et al. (2007) for a detailed description]. The latter ascomata can then be transferred to a microfuge tube and crushed to release the ascospores. These should then be suspended in 0.01 % Tween 20 or 0.05 % Tween 80 solution.
2. Transfer the ascospores onto a clear agar medium (e.g. water agar), spreading or using a decimal dilution series if a spore suspension is used.
3. Incubate overnight (or longer as need be) at a temperature allowing germination of the ascospores.
4. Locate single ascospores using a dissecting microscope and transfer these or hyphal tips from germinating ascospores to separate agar plates. Transfers can be efficiently made using a fine, flattened platinum wire mounted on the end of glass tubing. Such wires can be flame-sterilized and cool down rapidly.

### 2.5.6.2 Heat Treatment

The second method is based on a heat treatment of the ascospores. In some genera, ascospores are markedly more heat-resistant than the (heat-sensitive) asexual spores. This feature can be used to generate single ascospore isolates free of contaminating conidia and hyphae. For some species this heat treatment step is also required to trigger ascospore germination; without this ascospores do not germinate at standard growth

temperatures (Perkins 1986; O’Gorman et al. 2009). This method has been applied for example in *Neurospora*, *Paecilomyces*, and *Aspergillus* sect. *Fumigati* species.

1. Transfer the fruiting bodies from the agar plate into microfuge tubes containing 0.05 % Tween 80 or 0.01 % Tween 20 (ideally buffered at pH 6 with, e.g., 0.01 M sodium phosphate).
2. Agitate the suspension with glass beads in order to break open the ascomata and asci, or simply crush the ascomata on the side of the tube before thoroughly mixing in the buffered Tween.
3. Check by microscopy whether the majority of ascospores are released into the suspension. If not, repeat step 2.
4. Heat-treat the suspension in a water bath or on a thermal cycler (time and temperature is species-dependent). Note that most conidia are eliminated by a treatment of 10 min at 60 °C.
5. Make a decimal dilution in 0.05 % Tween 80 or 0.01 % Tween 20.
6. Spread plate 0.1 mL of each dilution on agar plates (duplicate).
7. Incubate at a suitable temperature allowing ascospore germination.
8. After incubation, pick individual young germinating colonies from the highest dilutions (preferable) or locate single ascospores using a dissecting microscope as described above.

### 2.5.6.3 Ejected Ascospores

Many species forcibly eject ascospores from the ascomata. The fact that ascospores are shot out of the ascomata provides a convenient way to collect pure ascospores, free of contaminating hyphae and conidia (Dyer et al. 1993). However, some caution must be exercised as some species (notably lichen-forming fungi) eject ascospores in packets, so it can prove difficult to isolate individual ascospores. There are two related methods available according to how the ejected ascospores are trapped, as follows.

#### Method 1

1. Pour a thin layer of clear agar (e.g. water agar) into the top (i.e. uppermost) lid of a Petri dish.

2. Place the lid over a Petri dish containing either individual ascoma or sets of ascomata (transferred on agar to keep the cultures moist).
3. Incubate the cultures for a suitable period (e.g., overnight or up to 48 h) during which time ascospores will be ejected onto the overlying agar.
4. Inspect the lids containing the ascospores using a dissecting microscope.
5. Subculture either ascospores or hyphal tips from germinating ascospores as described above.

#### Method 2

1. Aliquot 3 mL of 0.05 % Tween 80 or 0.01 % Tween 20 into the bottom of a sterile 5 mL Petri dish.
2. Attach either an individual ascoma or sets of ascomata to the lid of the Petri dish using Vaseline (petroleum jelly), ensuring that some of the underlying growth media is included to prevent desiccation of the ascoma.
3. Incubate the cultures for a suitable period (e.g., overnight or up to 48 h) during which time ascospores will be ejected into the underlying liquid.
4. Spread plate 0.5 mL aliquots of the Tween solution onto 9 cm Petri dishes containing a suitable clear media, and leave the dishes left to dry.
5. Incubate further and inspect the plates using a dissecting microscope.
6. Subculture either ascospores or hyphal tips from germinating ascospores as described above.

### 2.5.6.4 Isolation from Sclerotial Fruiting Bodies

If the fruiting bodies are firm and sclerotial, the following method can be used.

1. Harvest the fruiting bodies by adding 10 mL 0.05 % Tween 80 to the culture (slant or plate).
2. Scrape of the agar surface with a transfer loop.
3. Filter the suspension through a 100 µm filter.
4. Transfer the retained sclerotia/stromata to a vial.
5. Vortex the suspension followed by decanting to remove residual conidia. Repeat this procedure at least five times.

6. Filter the suspension onto Whatman #4 filter paper.
7. Clean the stromata used for obtaining single-ascospore cultures further by vortexing 1–2 min in 10 mL sterile water containing 1 g glass beads (200–350  $\mu\text{m}$  diameter).
8. Remove from the beads.
9. Vortex and decant twice with water.
10. Dissect sclerotia/stromata with a microscalpel under the stereomicroscope.
11. Transfer the ascospores onto an agar medium (e.g. water agar) as described above.
12. Incubate overnight at a temperature allowing germination.
13. Locate single ascospores using a (dissecting) microscope and transfer to separate agar plates.

### 2.5.7 Analysis of Progeny

A selection of the collected single-ascospore isolates should be assessed, to confirm that recombination has occurred in the case of putative outcrossing. Ascospore analysis can also be used to confirm the breeding system of the species in question. Various DNA fingerprinting techniques, such as RAPD and AFLP analysis, together with use of MLST markers and segregation of the *MAT* locus itself have been used to evaluate variation in the parents and progeny of a cross (e.g. Murtagh et al. 2000; Seymour et al. 2005; O’Gorman et al. 2009; Horn et al. 2009c; Swilaiman et al. 2013). It would be expected that self-fertilization via a homothallic breeding system would lead to uniformity in the ascospore progeny, whereas heterothallism and outcrossing would lead to genetic variation among the offspring (Murtagh et al. 2000). Indeed, a 1:1 segregation of mating types among the ascospore progeny is a clear indication of the presence of a heterothallic breeding system. However, it is cautioned that some ascospore progeny will be identical due to the fact that each set of eight ascospores in an ascus is composed of four sets of sister ascospores, derived from a mitotic division post-meiosis and tetrad formation.

## 2.6 Utilization of the Sexual Cycle as a Tool for Gene Identification and Manipulation

Once a reliable method has been developed to induce the sexual cycle in a given fungal species, it can then be used as a valuable laboratory tool for a range of applications such as classical genetic analysis, strain improvement, and as a complement to modern genetic manipulation experiments. A comprehensive description of the applications and uses of the sexual cycle is beyond the scope of the present chapter. Instead brief mention will now be made of ways in which the sexual cycle can be used in the context of fungal gene identification and transformation relevant to current accompanying chapters.

### 2.6.1 Genetics of Traits of Interest and Gene Identification

For many studies of gene transformation, candidate genes will have been identified at the onset of studies. However, in some cases the gene(s) responsible for a particular trait (and associated phenotype) of interest might be unknown. When studying such a trait it is very useful to know at the onset of gene manipulation studies whether that trait has a monogenic (i.e., determined by a single gene) or polygenic (i.e., determined by several genes) basis. This can influence the design of subsequent experimental and gene transformation studies. The sexual cycle provides an ideal tool to determine the genetic basis of a trait of interest because crosses can be set up between parents which differ in that trait. The ascospore progeny can then be collected and assessed for the trait of interest, with different patterns predicted in the frequency of offspring according to the genetic basis of the trait. In the case of a monogenic trait determined by a single dominant gene, it would be expected that haploid ascospore progeny will show a 1:1 segregation pattern for that trait. This can ideally be confirmed by backcrossing to the relevant parent. Conversely, if a trait has a polygenic basis with

several genes segregating simultaneously then a more complex pattern of inheritance will be evident, with the progeny often failing to show distinct classes but rather a continuous distribution of phenotypes between the two parents (Caten 1979; Dyer et al. 2000).

Once the genetic basis of a trait is known, various techniques exploiting the sexual cycle are then available to locate and try and identify the specific gene or genes of interest, or other genetic causal factor(s). Examples include firstly the use of classical genetic mapping techniques. Genetic maps now available for almost 30 fungal species, although these vary in their coverage (Foulongne-Oriol 2012). By using two- and three-point crossing data it should be possible to locate the position of the gene between two known markers on a genetic map (Perkins 1986) and then chromosome walking and bioinformatic approaches can be used to try and identify specific genes. This topic has recently been reviewed by Foulongne-Oriol (2012) and examples of the use of mapping of Mendelian traits are provided in the review of Hall (2013). Secondly, a method termed ‘bulk segregant analysis’ (BSA) can be used to identify DNA marker(s) linked to a region of the genome responsible for a particular phenotype based on analysis of sexual progeny (Michelmore et al. 1991). This involves making pooled bulks of DNA from the progeny based on the presence or absence of the phenotype of interest. These pooled samples can either then be screened for the presence of DNA markers (e.g., PCR fingerprints) or subjected to next generation sequencing (NGS). In theory the only differences between the pools should arise from the genetic marker of interest together with regions of the genome linked to that marker. Examples of the application of BSA to filamentous fungi using molecular markers include the work of Chun and Lee (1999), Jurgenson et al. (2002), Jin et al. (2007), Lewis et al. (2007), and Dettman et al. (2010). Meanwhile, BSA has been applied in conjunction with NGS by Lambreghts et al. (2009), Pomraning et al. (2011), and Nowrousian et al. (2012). A third method for gene localization based on the analysis of sexual progeny is the technique of quantitative trait loci (QTL) analysis. The QTL

method is especially suitable for providing insights into the genetic basis of more complex polygenic traits and can provide an estimation of the number of genes contributing to a particular trait and the identification of regions of the genomes linked to a particular trait (Miles and Wayne 2008). Hall (2013) has described how QTL mapping can be applied to genetic analysis of *Neurospora* species, including a review of the various mapping methods, and readers are referred there for further details. Recent examples of QTL analysis in filamentous fungi include those of Christians et al. (2011) in *A. nidulans* and Turner et al. (2011) in *Neurospora*. This method has the pre-requirement of parents which differ genetically with respect to the trait of interest, together with the presence of a dense genetic map. Thus, QTL mapping is only applicable to certain studies. It is also cautioned that although both BSA and QTL approaches can involve considerable work, they most often end with the identification of a genome region of interest with various candidate genes, rather than the actual identification of specific genes. Thus, further work is normally required after such studies.

## 2.6.2 Gene Manipulation by Sexual Reproduction: Strain Improvement and Gene Complementation

Other chapters in this book describe how the genetic composition of fungi can be manipulated by various methods of gene transformation. Although often overlooked, the sexual cycle can be also used as an efficient method for gene transformation.

The sexual cycle can be used to combine together genes of interest by crossing parents with the individual gene(s) and then selecting for ascospore offspring showing the desirable mixture of genes. For example, it might be desirable in certain gene function studies to produce mutant strains with multiple gene deletions. This can be achieved by lengthy rounds of transformation and marker recycling (Yoon et al. 2011). However, the sexual cycle

can provide an efficient alternative method because strains with complementary gene deletions can simply be crossed together. The sexual progeny are then screened for the presence or absence of the genes (e.g. by PCR assay) and isolates containing the desired combinations of multiple gene deletion selected for further study. In a parallel fashion, Böhm et al. (2013) illustrated how such an approach could be used for industrial strain improvement, with sexual progeny of *P. chrysogenum* screened for isolates that exhibited high penicillin titre but lack of a contaminating secondary metabolite chrysogenin. More broadly, sexual crosses can be set up to generate novel genetic diversity, allowing the offspring to be screened for isolates exhibiting, for example, either enhanced or decreased production of a particular metabolite, a phenomenon known as ‘transgressive segregation’ (Rieseberg et al. 1999).

Finally, the sexual cycle can be used for gene complementation purposes. When gene deletion has been shown to lead to a particular loss (or gain) of function, it is often required to then return the original gene back to the mutant strain to show that the original phenotype can be restored. This can be problematic if a limited number of selectable markers are available. However, by crossing the gene deletion strain with a strain containing the wild-type allele, it is possible to reintroduce the wild-type gene (Paoletti et al. 2007). If a consistent correlation between the presence of the wild-type gene (this can be screened for by PCR) and the presence of the restored phenotype can be demonstrated, or progeny which recapitulate the original genotype can be shown to exhibit the wild-type phenotype, then the role of the gene has been proven, i.e., proof of gene function through gene restoration by sexual crossing.

---

## 2.7 Conclusions and Outlook

Many industrially and clinically important fungal species were once thought to reproduce only asexually. Using a selection of the methods listed above, it has been shown that certain of

these species may also reproduce sexually. For some other species a sexual state has still never been observed, although analyses of molecular markers indicate recombination. The term heterothallic is used for outcrossing species where a sexual state has been observed, and we suggest the term “proto-heterothallic” for such asexual species where genetic evidence, such as the presence of complementary *MAT* loci, indicates the presence of a sexual cycle. Similarly, for species with a homothallic mating type organization lacking a sexual state we propose the term “proto-homothallic”.

In this manuscript, we describe tools for the induction of a sexual state in heterothallic fungi. Based on the review, we show that no single protocol can secure induction of a sexual state in heterothallic fungi, not even in species belonging to the same genus, e.g., *Aspergillus*. There are likely to be more surprises waiting as we observe a ‘sexual revolution’ in fungi (Dyer and O’Gorman 2011). Recently, sclerotia production was induced in the proto-heterothallic species *Aspergillus niger*. The formation of sclerotia, thought to be sterile fruiting bodies in certain *Aspergillus* species, can be induced by inoculating *A. niger* onto fruits such as raisins, blueberries, cranberries, mulberries, apricot, prune, and mango on a CYA agar (Frisvad et al. 2014). By following (one of the) methods mentioned above and applying these specific growth conditions, a sexual state might be discovered in the near future for this biotechnologically important species. Finally, experimental crossings in vitro do not strictly reflect what happens in natura. For example, heat-resistant ascospores of the heterothallic *P. variotii* frequently spoil pasteurised fruit drinks and other food products, indicating a common occurrence of these ascospores and therefore the sexual state in nature. However, incubation times up to 6 weeks and specific agar media (PDA) are needed to induce sexual recombination in the lab (Houbraken et al. 2008).

**Acknowledgments** J.H. thanks Tineke van Doorn and Richard Summerbell for the discussions on the term proto-heterothallic.

## References

- Aanen DK, Hoekstra RF (2007) Why sex is good: on fungi and beyond. In: Heitman J, Kronstad JW, Taylor JW, Casselton LA (eds) Sex in fungi: molecular determination and evolutionary principles. ASM, Washington, pp 527–534
- Arabatzis M, Velegraki A (2013) Sexual reproduction in the opportunistic human pathogen *Aspergillus terreus*. *Mycologia* 105:71–79
- Arie T, Christiansen SK, Yoder OC, Turgeon BG (1997) Efficient cloning of ascomycete mating type genes by PCR amplification of the conserved MAT HMG box. *Fungal Genet Biol* 32:118–130
- Astell CR, Ahlstrom-Jonasson L, Smith M, Tatchell K, Nasmith KA, Hall BD (1981) The sequence of the DNAs coding for the mating-type loci of *Saccharomyces cerevisiae*. *Cell* 27:15–23
- Barrs VR, van Doorn TM, Houbraken J, Kidd SE, Martin P, Pinheiro DM, Richardson M, Varga J, Samson RA (2013) *Aspergillus felis* sp. nov., an emerging agent of invasive aspergillosis in humans, cats and dogs. *PLoS One* 8:e64871
- Bistis GN (1998) Physiological heterothallism and sexuality in euascomycetes: a partial history. *Fungal Genet Biol* 23:213–222
- Blakeslee AF (1904) Sexual reproduction in the Mucorineae. *Proc Natl Acad Sci U S A* 40:205–319
- Böhm J, Hoff B, O’Gorman CM, Wolfers S, Klux V, Binger D, Zadra I, Kürnsteiner H, Pöggeler S, Dyer PS, Kück U (2013) Sexual reproduction and mating-type-mediated strain development in the penicillin-producing fungus *Penicillium chrysogenum*. *Proc Natl Acad Sci U S A* 110:1476–1481
- Burnett J (2003) Fungal populations and species. Oxford University Press, New York
- Caten CE (1979) Quantitative genetic variation in fungi. In: Thompson JN, Thoday JM (eds) Quantitative genetic variation. Academic, New York, pp 35–59
- Cavindera B, Trail F (2012) Role of *Fig1*, a component of the low-affinity calcium uptake system, in growth and sexual development of filamentous fungi. *Eukaryot Cell* 11:978–988
- Choi H-W, Kim J-M, Hong S-K, Kim WG, Chun S-C, S-H Y (2009) Mating types and optimum culture conditions for sexual state formation of *Fusarium fujikuroi* isolates. *Mycobiology* 37:247–250
- Christians JK, Cheema MS, Vergara IA, Watt CA, Pinto LJ, Chen N, Moore MM (2011) Quantitative trait locus (QTL) mapping reveals a role for unstudied genes in *Aspergillus* virulence. *PLoS One* 6:e19325
- Chun SJ, Lee Y-H (1999) Genetic analysis of a mutation on appressorium formation in *Magnaporthe grisea*. *FEMS Microbiol Lett* 173:133–137
- Covert SF, Aoki T, O’Donnell K, Starkey D, Holliday A, Geiser DM, Cheung F, Town C, Strom A, Juba J, Scandiani M, Yang XB (2007) Sexual reproduction in the soybean sudden death syndrome pathogen *Fusarium tucumaniae*. *Fungal Genet Biol* 44:799–807
- Covert SF, Briley A, Wallace MM, McKinney VT (1999) Partial MAT-2 gene structure and the influence of temperature on mating success in *Gibberella circinata*. *Fungal Genet Biol* 28:43–54
- Crous PW, Verkley GJM, Groenewald JZ, Samson RA (eds) (2009) Fungal biodiversity. CBS laboratory manual series 1. Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands
- Darbyshir HL, van de Vondervoort PJI, Dyer PS (2013) Discovery of sexual reproduction in the black aspergill. *Fungal Genet Rep* 60(Suppl):687
- Debuchy R, Berteaux-Lecellier V, Silar P (2010) Mating systems and sexual morphogenesis in ascomycetes. In: Borkovich KA, Ebbole DJ (eds) Cellular and molecular biology of filamentous fungi. ASM, Washington, pp 501–535
- Debuchy R, Turgeon BG (2006) Mating-type structure, evolution, and function in euascomycetes. In: Kües U, Fischer R (eds) The mycota I: growth, differentiation and sexuality. Springer, Berlin, Germany, pp 293–323
- Dettman JR, Anderson JB, Kohn LM (2010) Genome-wide investigation of reproductive isolation in experimental lineages and natural species of *Neurospora*: identifying candidate regions by microarray-based genotyping and mapping. *Evolution* 64:694–709
- Dodge BO (1957) Rib formation in ascospores of *Neurospora* and questions of terminology. *Bull Torrey Bot Club* 84:182–188
- Dyer PS (2007) Sexual reproduction and significance of MAT in the aspergilli. In: Heitman J, Kronstad JW, Taylor JW, Casselton LA (eds) Sex in fungi: molecular determination and evolutionary principles. ASM, Washington, pp 123–142
- Dyer PS, Bateman GL, Wood HW (2001a) Development of apothecia of the eyespot pathogen *Tapesia* on cereal crop stubble residue in England. *Plant Pathol* 50:356–362
- Dyer PS, Furneaux PA, Douhan G, Murray TD (2001b) A multiplex PCR test for determination of mating type applied to the plant pathogens *Tapesia yallundae* and *Tapesia acuformis*. *Fungal Genet Biol* 33:173–180
- Dyer PS, Hansen J, Delaney A, Lucas JA (2000) Genetic control of resistance to the DMI fungicide prochloraz in the cereal eyespot pathogen *Tapesia yallundae*. *Appl Environ Microbiol* 66:4599–4604
- Dyer PS, Ingram DS, Johnstone K (1992) The control of sexual morphogenesis in the Ascomycotina. *Biol Rev* 67:421–458
- Dyer PS, Nicholson P, Lucas JA, Peberdy JF (1995) Genetic control of sexual compatibility in *Tapesia yallundae*, 79. Abstracts, eighteenth fungal genetics conference, Asilomar. University of California, USA
- Dyer PS, Nicholson P, Lucas JA, Peberdy JF (1996) *Tapesia acuformis* as a causal agent of eyespot disease of cereals and evidence for a heterothallic mating system using molecular markers. *Mycol Res* 100:1219–1226

- Dyer PS, Nicholson P, Rezanoor HN, Lucas JA, Peberdy JF (1993) Two-allele heterothallism in *Tapesia yallundae*, the teleomorph of the cereal eyespot pathogen *Pseudocercospora herpotrichoides*. *Physiol Mol Plant Pathol* 43:403–414
- Dyer PS, O’Gorman CM (2011) A fungal sexual revolution: *Aspergillus* and *Penicillium* show the way. *Curr Opin Microbiol* 14:649–654
- Dyer PS, O’Gorman CM (2012) Sexual development and cryptic sexuality in fungi: insights from *Aspergillus* species. *FEMS Microbiol Rev* 36:165–192
- Dyer PS, Paoletti M (2005) Reproduction in *Aspergillus fumigatus*: sexuality in a supposedly asexual species? *Med Mycol* 43(Suppl 1):S7–S14
- Eagle CE (2009) Mating-type genes and sexual potential in the Ascomycete genera *Aspergillus* and *Penicillium*. PhD thesis, University of Nottingham, UK
- Ehrenberg CG (1820) *Syzygites*, eine neue Schimmelgattung, nebst Beobachtungen über sichtbare Bewegung in Schimmeln. *Verhandl. Gesamte Naturf., Freunde, Berlin*, pp 98–109
- Faretra F, Antonacci E, Pollastro S (1988) Sexual behaviour and mating system of *Botryotinia fuckeliana*, teleomorph of *Botrytis cinerea*. *J Gen Microbiol* 134:2543–2550
- Foulongne-Oriol M (2012) Genetic linkage mapping in fungi: current state, applications, and future trends. *Appl Microbiol Biotechnol* 95:891–904
- Fraser JA, Stajich JE, Tarcha EJ, Cole GT, Inglis DO, Sil A, Heitman J (2007) Evolution of the mating type locus: insights gained from the dimorphic primary fungal pathogens *Histoplasma capsulatum*, *Coccidioides immitis*, and *Coccidioides posadasii*. *Eukaryot Cell* 6:622–629
- Fresenius G (1863) Beiträge zur Mykologie. H.L. Brönnner, Frankfurt, Germany
- Frisvad JC, Petersen LM, Lyhne K, Larsen TO (2014) Formation of sclerotia and production of indoloterpenes by *Aspergillus niger* and other species in section *Nigri*. *PLoS One* 9:e94857
- Galagan JE, Calvo SE, Cuomo C, Ma LJ, Wortman J, Batzoglou S, Lee SL, Batürkmen M, Spevak CC, Clutterbuck J, Kapitonov V, Jurka J, Scacciocchio C, Farman M, Butler J, Purcell S, Harris S, Braus GH, Draht O, Busch S, D’Enfert C, Bouchier C, Goldman GH, Bell-Pedersen D, Griffiths-Jones S, Doonan JH, Yu J, Vienken K, Pain A, Freitag M, Selker EU, Archer DB, Peñalva MA, Oakley BR, Momany M, Tanaka T, Kumagai T, Asai K, Machida M, Nierman WC, Denning DW, Caddick M, Hynes M, Paoletti M, Fischer R, Miller B, Dyer P, Sachs MS, Osmani SA, Birren B (2005) Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* 438:1105–1115
- Gordon WL (1961) Sex and mating type in relation to the production of perithecia by certain species of *Fusarium*. *Proc Can Phytopathol Soc* 28:11
- Hall C (2013) Quantitative genetics in *Neurospora*. In: Kasbekar DP, McCluskey K (eds) *Neurospora* genomics and molecular biology. Caister Academic, Norfolk, UK, pp 65–84
- Han KH, Lee DB, Kim JH, Kim MS, Han KY, Kim WS, Park YS, Kim HB, Han DM (2003) Environmental factors affecting development of *Aspergillus nidulans*. *J Microbiol* 41:34–40
- Heitman J, Carter DA, Dyer PS, Soll DR (2014) Sexual reproduction of human fungal pathogens. In: Casadevall A, Mitchell AP, Berman J, Kwon-Chung KJ, Perfect JR, Heitman J (eds) *Fungal pathogens*. Cold Spring Harbour Laboratory Press, New York
- Henk DA, Eagle CE, Brown K, van den Berg MA, Dyer PS, Peterson SW, Fisher MC (2011) Speciation despite globally overlapping distributions in *Penicillium chrysogenum*: the population genetics of Alexander Fleming’s lucky fungus. *Mol Ecol* 20:4288–4301
- Hoff B, Pöggeler S, Kück U (2008) Eighty years after its discovery, Fleming’s *Penicillium* strain discloses the secret of its sex. *Eukaryot Cell* 7:465–470
- Horn BW, Moore GG, Carbone I (2009a) Sexual reproduction in *Aspergillus flavus*. *Mycologia* 101:423–429
- Horn BW, Moore GG, Carbone I (2011) Sexual reproduction in aflatoxin-producing *Aspergillus nomius*. *Mycologia* 103:174–183
- Horn BW, Olarte RA, Peterson SW, Carbone I (2013) Sexual reproduction in *Aspergillus tubingensis* from section *Nigri*. *Mycologia* 105:1153–1163
- Horn BW, Ramirez-Prado J, Carbone I (2009b) The sexual state of *Aspergillus parasiticus*. *Mycologia* 101:275–280
- Horn BW, Ramirez-Prado JH, Carbone I (2009c) Sexual reproduction and recombination in the aflatoxin-producing fungus *Aspergillus parasiticus*. *Fungal Genet Biol* 46:169–175
- Houbraken J, Frisvad JC, Seifert KA, Overy DP, Tuthill DM, Valdez JG, Samson RA (2012) New penicillin-producing *Penicillium* species and an overview of section *Chrysogena*. *Persoonia* 29:78–100
- Houbraken J, Varga J, Rico-Munoz E, Johnson S, Samson RA (2008) Sexual reproduction as the cause of heat resistance in the food spoilage fungus *Byssosclamyces spectabilis* (anamorph *Paecilomyces variotii*). *Appl Environ Microbiol* 74:1613–1619
- Houbraken J, de Vries RP, Samson RA (2014) Modern taxonomy of biotechnologically important *Aspergillus* and *Penicillium* species. *Adv Appl Microbiol* 86:199–249
- Idnurm A (2011) Sex determination in the first-described sexual fungus. *Eukaryot Cell* 10:1485–1491
- Jin Y, Allen S, Baber L, Bhattarai EK, Lamb TM, Versaw WK (2007) Rapid genetic mapping in *Neurospora crassa*. *Fungal Genet Biol* 44:455–465
- Jurgenson JE, Zeller KA, Leslie JF (2002) Expanded genetic map of *Gibberella moniliformis* (*Fusarium verticillioides*). *Appl Environ Microbiol* 68:1972–1979
- Kawasaki M, Anzawa K, Wakasa A, Takeda K, Mochizuki T, Ishizaki H, Hemashettar B (2010) Matings among



- three teleomorphs of *Trichophyton mentagrophytes*. *Jpn J Med Mycol* 51:143–452
- Kent CR, Ortiz-Bermúdez P, Gilies SS, Hull CM (2008) Formulation of a defined V8 medium for induction of sexual development of *Cryptococcus neoformans*. *Appl Environ Biol* 74:6248–6253
- Kerényi Z, Moretti A, Waalwijk C, Olah B, Hornok L (2004) Mating type sequences in asexually reproducing *Fusarium* species. *Appl Environ Microbiol* 70:4419–4423
- Klittich CJR, Leslie JF (1988) Nitrate reduction mutants of *Fusarium moniliforme* (*Gibberella fujikuroi*). *Genetics* 118:417–423
- Kwon-Chung KJ, Sugui JA (2009) Sexual reproduction in *Aspergillus* species of medical or economic importance: why so fastidious? *Trends Microbiol* 17:481–487
- Kwon-Chung KJ, Weeks RJ, Larsh HW (1974) Studies on *Emmonsia capsulata* (*Histoplasma capsulatum*). II. Distribution of the two mating types in 13 endemic states of the United States. *Am J Epidemiol* 99:44–49
- Lambreghts R, Shi M, Belden WJ, deCaprio D, Park D, Henn MR, Galagan JE, Baştürkmen M, Birren BW, Sachs MS, Dunlap JC, Loros JL (2009) A high-density single nucleotide polymorphism map for *Neurospora crassa*. *Genetics* 181:467–781
- Lee SC, Ni M, Li W, Shertz C, Heitman J (2010) The evolution of sex: a perspective from the fungal kingdom. *Microbiol Mol Biol Rev* 74:298–340
- Lewis ZA, Shiver AL, Stiffler N, Miller MR, Johnson EA, Selker EU (2007) High-density detection of restriction-site-associated DNA markers for rapid mapping of mutated loci in *Neurospora*. *Genetics* 177:1163–1171
- López-Villavicencio M, Aguilera G, Giraud T, de Vienne DM, Lacoste S, Couloux A, Dupont J (2010) Sex in *Penicillium*: combined phylogenetic and experimental approaches. *Fungal Genet Biol* 47:693–706
- McAlpin CE, Wicklow DT (2005) Culture media and sources of nitrogen promoting the formation of stromata and ascocarps in *Petromyces alliaceus* (*Aspergillus* section *Flavi*). *Can J Microbiol* 51:765–771
- Metzenberg RL, Glass NL (1990) Mating type and mating strategies in *Neurospora*. *Bioessays* 12:53–59
- Michelmore RW, Paran I, Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proc Natl Acad Sci U S A* 88:9828–9832
- Miles CM, Wayne M (2008) Quantitative trait locus (QTL) analysis. *Nat Educ* 1:208
- Milgroom MG (1996) Recombination and the multilocus structure of fungal population. *Annu Rev Phytopathol* 34:457–477
- Mooney JL, Yager LN (1990) Light is required for conidiation in *Aspergillus nidulans*. *Genes Dev* 4:1473–1482
- Moore GG (2014) Sex and recombination in aflatoxigenic *Aspergilli*: global implications. *Front Microbiol* 5:32
- Murtagh GJ, Dyer PS, Crittenden PD (2000) Sex and the single lichen. *Nature* 404:564
- Nielsen KA, De Obaldia AL, Heitman J (2007) *Cryptococcus neoformans* mates on pigeon guano: implications for the realized ecological niche and globalization. *Eukaryot Cell* 6:949959
- Nowrousian M, Teichert I, Masloff S, Kück U (2012) Whole-genome sequencing of *Sordaria macrospora* mutants identifies developmental genes. *G3. Genes Genomes Genetics* 2:261–270
- Nováková A, Hubka V, Dudová Z, Matsuzawa T, Kubátová A, Yaguchi T, Kolařík M (2014) New species in *Aspergillus* section *Fumigati* from reclamation sites in Wyoming (U.S.A.) and revision of *A. viridinutans* complex. *Fung Div* 64(1):253–274. doi:10.1007/s13225-013-0262-5
- O’Gorman CM, Fuller HT, Dyer PS (2009) Discovery of a sexual cycle in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Nature* 457:471–474
- Paoletti M, Rydholm C, Schwier EU, Anderson MJ, Szakacs G, Lutzoni F, Debeauvais JP, Latgé JP, Denning DW, Dyer PS (2005) Evidence for sexuality in the opportunistic human pathogen *Aspergillus fumigatus*. *Curr Biol* 15:1242–1248
- Paoletti M, Seymour FA, Alcocer MJC, Kaur N, Calvo AM, Archer DB, Dyer PS (2007) Mating type and the genetic basis of self-fertility in the model fungus *Aspergillus nidulans*. *Curr Biol* 17:1384–1389
- Perkins DD (1986) Hints and precautions for the care, feeding and breeding of *Neurospora*. *Fungal Genet Newsl* 33:36–41
- Pitt JI (1979) The genus *Penicillium* and its teleomorph states *Eupenicillium* and *Talaromyces*. Academic, London, UK
- Pöggeler S (2001) Mating-type genes for classical strain improvements of ascomycetes. *Appl Microbiol Biotechnol* 56:589–601
- Pomraning KR, Smith KM, Freitag M (2011) Bulk segregant analysis followed by high-throughput sequencing reveals the *Neurospora* cell cycle gene, *ndc-1*, to be allelic with the gene for ornithine decarboxylase, *spe-1*. *Eukaryot Cell* 10:724–733
- Pringle A, Baker DM, Platt JL, Wares JP, Latgé JP, Taylor JW (2005) Cryptic speciation in the cosmopolitan and clonal human pathogenic fungus *Aspergillus fumigatus*. *Evolution* 59:1886–1899
- Raju NB, Perkins DD (1994) Diverse programs of ascus development in pseudohomothallic species of *Neurospora*, *Gelasinospora* and *Podospora*. *Dev Genet* 15:104–118
- Ramirez-Prado JH, Moore GG, Horn BW, Carbone I (2008) Characterization and population analysis of the mating-type genes in *Aspergillus flavus* and *Aspergillus parasiticus*. *Fungal Genet Biol* 45:1292–1299
- Raper KB, Fennell DI (1965) The genus *Aspergillus*. Williams & Wilkins Co, Baltimore
- Rieseberg LH, Archer MA, Wayne RK (1999) Transgressive segregation, adaptation and speciation. *Heredity* 83:363–372

- Robellet X, Oestreicher N, Guitton A, Vélot C (2010) Gene silencing of transgenes inserted in the *Aspergillus nidulans* *alcM* and/or *alcS* loci. *Curr Genet* 56:341–348
- Ropars J, López-Villavicencio M, Dupont J, Snirc A, Gillot G, Coton M, Jany J-L, Coton E, Giraud T (2014) Induction of sexual reproduction and genetic diversity in the cheese fungus *Penicillium roqueforti*. *Evol Appl* 7:433–441
- Rydholm C, Dyer PS, Lutzoni F (2007) DNA sequence characterization and molecular evolution of MAT1 and MAT2 mating-type loci of the self-compatible ascomycete mold *Neosartorya fischeri*. *Eukaryot Cell* 6:868–874
- Samson RA, Houbraken J, Thrane U, Frisvad JC, Andersen B (2010) Food and indoor fungi. CBS laboratory manual Series 2. CBS KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands
- Samson RA, Varga J, Dyer PS (2009) Morphology and reproductive mode of *Aspergillus fumigatus*. In: Latgé JP, Steinbach WJ (eds) *Aspergillus fumigatus* and Aspergillosis. ASM, Washington, pp 7–13
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Fungal Barcoding Consortium (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proc Natl Acad Sci U S A* 109:6241–6246
- Seidl V, Seibel C, Kubicek CP, Schmoll M (2009) Sexual development in the industrial workhorse *Trichoderma reesei*. *Proc Natl Acad Sci U S A* 106:13909–13914
- Seymour FA, Crittenden PD, Dickinson MJ, Paoletti M, Montiel D, Cho L, Dyer PS (2005) Breeding systems in the lichen-forming fungal genus *Cladonia*. *Fungal Genet Biol* 42:554–563
- Short DP, O'Donnell K, Thrane U, Nielsen KF, Zhang N, Juba JH, Geiser DM (2013) Phylogenetic relationships among members of the *Fusarium solani* species complex in human infections and the descriptions of *F. keratoplasticum* sp. nov. and *F. petrophilum* stat. nov. *Fungal Genet. Biol* 53:59–70
- Singh G, Dyer PS, Ashby AM (1999) Intra-specific and inter-specific conservation of mating-type genes from the discomycete plant-pathogenic fungi *Pyrenopeziza brassicae* and *Tapesia yallundae*. *Curr Genet* 36:290–300
- Steenkamp ET, Wingfield BD, Coutinho TA, Zeller KA, Wingfield MJ, Marasas WF, Leslie JF (2000) PCR-based identification of MAT-1 and MAT-2 in the *Gibberella fujikuroi* species complex. *Appl Environ Microbiol* 66:4378–4382
- Sugui JA, Losada L, Wang W, Varga J, Ngamskulrungrong P, Abu-Asab M, Chang YC, O'Gorman CM, Wickes BL, Nierman WC, Dyer PS, Kwon-Chung KJ (2011) Identification and characterization of an *Aspergillus fumigatus* “supermater” pair. *MBio* 2:e00234–11
- Sugui JA, Vinh DC, Nardone G, Shea YR, Chang YC, Zelazny AM, Marr KA, Holland SM, Kwon-Chung KJ (2010) *Neosartorya udagawae* (*Aspergillus udagawae*), an emerging agent of aspergillosis: how different is it from *Aspergillus fumigatus*? *J Clin Microbiol* 48:220–228
- Swilaiman SS, O'Gorman CM, Balajee SA, Dyer PS (2013) Discovery of a sexual cycle in *Aspergillus lentulus*, a close relative of *A. fumigatus*. *Eukaryot Cell* 12:962–969
- Takada M, Udagawa S (1988) A new species of heterothallic *Talaromyces*. *Mycotaxon* 31:417–425
- Takan JP, Chipili J, Muthumeenakshi S, Talbot NJ, Manyasa EO, Bandyopadhyay R, Sere Y, Nutsugah SK, Talhinhos P, Hossain M, Brown AE, Sreenivasaprasad S (2012) *Magnaporthe oryzae* populations adapted to finger millet and rice exhibit distinctive patterns of genetic diversity, sexuality and host interaction. *Mol Biotechnol* 50:145–158
- Todd RB, Davis MA, Hynes MJ (2007) Genetic manipulation of *Aspergillus nidulans*: meiotic progeny for genetic analysis and strain construction. *Nat Protoc* 2:811–821
- Turner E, Jacobson DJ, Taylor JW (2011) Genetic architecture of a reinforced, postmating reproductive isolation barrier between *Neurospora* species indicates evolution via natural selection. *PLoS Genet* 7:e1002204
- Tschanz AT, Horst RK, Nelson PE (1976) The effect of environment on sexual reproduction of *Gibberella zeae*. *Mycologia* 68:327–340
- Turgeon BG, Yoder OC (2000) Proposed nomenclature for mating type genes of filamentous ascomycetes. *Fungal Genet Biol* 31:1–5
- Yoon J, Maruyama J, Kitamoto K (2011) Disruption of ten protease genes in the filamentous fungus *Aspergillus oryzae* highly improves production of heterologous proteins. *Appl Microbiol Biotechnol* 89:747–759