

# ERRATUM

**Lawrence M. Friedman, Curt D. Furberg, David L. DeMets, David M. Reboussin and Christopher B. Granger**

© Springer International Publishing Switzerland 2015  
L.M. Friedman et al., *Fundamentals of Clinical Trials*,  
DOI 10.1007/978-3-319-18539-2\_23

---

DOI 10.1007/978-3-319-18539-2\_23

In the original publication on page vi, there is a typographical error in the print and online versions of this book. “Principal” was incorrectly spelled as “Principle.”

Corrections to chapter 17, page 381, follow, and these changes have been updated in the book.

---

The updated original online version for this chapter can be found at  
[10.1007/978-3-319-18539-2\\_17](https://doi.org/10.1007/978-3-319-18539-2_17)

© Springer International Publishing Switzerland 2015  
L.M. Friedman et al., *Fundamentals of Clinical Trials*,  
DOI 10.1007/978-3-319-18539-2\_23

## Chapter 17

# Statistical Methods Used in Interim Monitoring

p 381

Many different spending functions can be specified. The O'Brien–Fleming  $\alpha_1(t^*)$  and Pocock  $\alpha_2(t^*)$  type spending functions are specified as follows:

$$\begin{aligned}\alpha_1(t^*) &= 2 - 2\Phi\left(Z_{\alpha/2}/\sqrt{t^*}\right) && \sim \text{O'Brien-Fleming} \\ \alpha_2(t^*) &= \alpha \ln(1 + (e - 1)t^*) && \sim \text{Pocock} \\ \alpha_3(t^*) &= \alpha t^{*\theta} && \text{for } \theta > 0\end{aligned}$$

The spending function  $\alpha_3(t^*)$  spends alpha uniformly during the trial for  $\theta = 1$ , at a rate somewhat between  $\alpha_1(t^*)$  and  $\alpha_2(t^*)$ . Other spending functions have also been defined [75, 76].