

Adnan Aydiner  
Abdullah İğci  
Atilla Soran  
*Editors*

# Breast Disease

Diagnosis and Pathology

Volume 1

 Springer

---

# Breast Disease



---

Adnan Aydiner • Abdullah İğci  
Atilla Soran  
Editors

# Breast Disease

Diagnosis and Pathology

Volume 1

 Springer



*Editors*

Adnan Aydiner  
Medical Faculty  
Istanbul University  
Istanbul  
Turkey

Atilla Soran  
Magee-Womens Hospital of UPMC  
Pittsburg, CA  
USA

Abdullah İğci  
Institute of Oncology  
Istanbul University  
Istanbul  
Turkey

ISBN 978-3-319-22842-6      ISBN 978-3-319-22843-3 (eBook)  
DOI 10.1007/978-3-319-22843-3

Library of Congress Control Number: 2015955271

Springer Cham Heidelberg New York Dordrecht London  
© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer International Publishing AG Switzerland is part of Springer Science+Business Media  
([www.springer.com](http://www.springer.com))

---

## Preface

The goal of *Breast Disease Diagnosis and Pathology* is to provide a comprehensive, scholarly appraisal of contemporary basic science and diagnosis. Because of advances in molecular medicine and therapeutics, this appraisal requires a more extensive understanding of the basic science of oncology than was required in the past. This book is organized into 16 chapters, and a brief summary of their content is provided below. In addition, we highlight some of the various important points in this book.

The topic of *Benign Breast Diseases* covers a wide variety of conditions, both common and unusual, that may require additional work-up, excision, or surveillance. In addition, some benign conditions may confer an increased risk for future disease, and this risk should be explained to the patient during treatment for these entities. This section attempts to provide a basic understanding of some of the most frequently encountered benign breast conditions and various rare types, including current recommendations for work-up, management, differential diagnoses, and future surveillance. The specific conditions that are explored in this chapter include fibroadenomas, intraductal papillomas, lipomas, hamartomas, radial scars, and gynecomastia in males.

Age, family history, and both endogenous and exogenous ovarian hormone exposure have important effects on risk and have been incorporated into models that predict the individual *risk of breast cancer*. Diet, alcohol use, and other factors play smaller roles. BRCA mutation-associated breast cancer differs from sporadic breast cancer in that BRCA mutation carriers exhibit an increased risk of breast and ovarian cancer and differential sensitivity to chemotherapeutic agents. Because BRCA genetic testing is readily available, BRCA mutation status should be evaluated in high-risk women, including women who were diagnosed with breast cancer at an early age and women with a strong family history or triple-negative tumors. Given the high rate of contralateral breast cancer and ovarian cancer, mutation carriers with newly diagnosed breast cancer may choose to undergo contralateral prophylactic mastectomy or bilateral salpingo-oophorectomy. In addition, two selective estrogen-receptor modulators, tamoxifen and raloxifene, and aromatase inhibitors can be used to decrease the incidence of invasive breast cancer in women who are at high risk of this condition.

*Breast imaging* is an essential component of breast cancer diagnosis and guides surgery and treatment options. Imaging techniques, such as mammography, ultrasound (US), and magnetic resonance imaging (MRI), enable the

detection of breast cancer at earlier stages. Mammography remains the standard screening examination; however, additional imaging studies are useful in evaluating the breast. US is utilized primarily in the diagnostic setting to characterize mammographic or palpable findings and assess axillary lymph nodes. Supplemental US screening may also be useful in patients with an intermediate risk for developing breast cancer and dense breasts to increase cancer detection. In addition to mammography, high-risk patients may also have annual MRI or US screening if they are unable to undergo MRI. MRI is also performed to evaluate the extent of disease, the response to neoadjuvant chemotherapy, and silicone implant integrity. In addition, these imaging modalities are also used to guide percutaneous biopsy, enabling minimally invasive tissue diagnosis.

*Lobular carcinoma in situ* (LCIS) is a high-risk indicator lesion for and a non-obligate precursor of the development of invasive breast carcinoma. Loss of E-cadherin is the hallmark pathological feature of lobular entities. Effective clinical management of LCIS requires good communication among the radiologist, surgeon, pathologist, and medical oncologist and entails surgical excision, subsequent surveillance, and systemic and surgical strategies to reduce the risk of future invasive cancer.

*Ductal carcinoma in situ* (DCIS) is defined as abnormally proliferating malignant cells confined to the breast milk ducts by the basement membrane. DCIS is diagnosed most commonly as a mammographic abnormality but can occasionally present as a palpable breast mass. Overall survival after breast-conserving therapy is equivalent to that observed for mastectomy. Patients undergoing mastectomy for DCIS should have sentinel lymph node biopsy (SLNB). Immediate breast reconstruction should be considered for patients undergoing mastectomy. Endocrine therapy, such as tamoxifen, is offered for 5 years to women with estrogen receptor-positive DCIS.

The topic of *Biology and Genetics of Breast Cancer* covers a wide variety of molecular studies. Understanding the mechanisms of DNA alterations leading to carcinogenesis can provide crucial insights for resolving the development of malignant processes, such as growth, invasion, and metastasis. This chapter reviews hereditary and somatic genetic alterations, epigenetic misregulations, and miRNA signatures associated with breast cancer. This chapter also emphasizes the molecular profiles of breast cancer and critical signaling pathway alterations.

Human breast cancers depend on estrogen and/or progesterone for growth, and these effects are mediated through *estrogen receptors* (ERs) and *progesterone receptors* (PRs), respectively. The *human epidermal growth factor receptor 2* (HER2) gene encodes a member of the epidermal growth factor receptor family of receptor tyrosine kinases, and its amplification with resultant overexpression plays a major role in sustaining multiple pathways in cancer growth. ERs, PRs, and HER2 status are the most important molecular markers in the standard care of all primary and recurrent/metastatic breast cancer patients and play both predictive and prognostic roles. The responsiveness of a tumor to hormone therapy is an important parameter in breast cancer management in both the adjuvant and metastatic settings. Only breast cancers with HER2 amplification or overexpression respond to HER2-directed therapies. Tumor hormonal status

is prognostic for patient outcome and potential sites of metastasis. Hormonal receptor-positive disease represents an indolent and slowly growing tumor with longer time to disease recurrence. HER2 is a poor prognostic factor in the absence of HER2-directed therapies. Assessment of the ER/PR/HER2 status is an essential factor in the evaluation of every newly diagnosed breast cancer, and the standardization of assay methods is crucial.

Invasive breast carcinomas comprise a heterogeneous group of lesions that differ in their molecular and pathologic features and clinical behavior. Some patients experience long periods of disease-free survival, whereas others experience the rapid development of recurrence and metastases that are fatal within a few years of the initial diagnosis. Numerous factors in individual tumors can be evaluated to stratify patients into subsets with varying risks of recurrence and response to different therapy modalities. The *Prognostic and Predictive Factors of Invasive Breast Cancer* chapter describes the current standard prognostic and predictive factors of invasive breast carcinoma and discusses emerging data on molecular markers that can be considered in clinical practice.

Adjuvant chemotherapy and endocrine treatment decrease the mortality of early breast cancer. However, not all early breast cancer patients benefit equally from adjuvant endocrine treatment and/or chemotherapy. High-risk patients are classically identified based on clinicopathological factors, such as age, tumor size, histopathological grade, nodal status, hormone and HER2 receptor positivity, and menopausal status. However, for patients with early breast cancer, the use of these standard clinicopathological factors might not thoroughly reveal the individual risk of disease recurrence and the benefits from adjuvant systemic chemotherapy. Many patients with early breast cancer do not derive benefit from adjuvant systemic chemotherapy. Quantitative approaches for defining prognoses and individualizing treatments are required. In recent years, *molecular signatures of gene expression* have been correlated with breast cancer recurrence risk. Several tests for genomic expression have been developed and validated on specimens from previous phase III studies to improve the prognostication of early breast cancer patients and/or the prediction of adjuvant systemic treatment.

In clinical practice, although local recurrence or distant metastasis develops in some individuals who have been assessed as low risk despite treatment, some individuals with high-risk disease do not relapse despite systemic and local therapy. Therefore, oncologists must determine objective prognostic factors to identify early recurrence and metastasis in patients with breast cancer. Based on the presumption of residual disease, clinicians have recently attempted to identify micrometastases using *disseminated tumor cells* (DTCs) in the *bone marrow and circulating tumor cells* (CTCs) from the peripheral blood. DTCs are known as epithelial cells in the bone marrow, and they are also considered to be micrometastases in the bone marrow. DTCs are observed in approximately 30 % of early-stage breast cancer patients. Tumor cells that circulate in the peripheral blood of patients with cancer are referred to as CTCs. CTCs are cells that have entered the peripheral blood circulation after having detached from an existing primary tumor or its metastases. DTCs and CTCs can be used to predict progression-free and overall survival as well as response to treatment.

In the *Pathology of Breast Cancer* chapter, classification is based on the recent WHO classification of breast carcinoma, and specific gross and microscopic features of in situ and invasive breast carcinomas are explained. Morphological groups, grading of DCIS, and the necessary information that should be included in a surgical pathology report are discussed. Recent information regarding columnar cell lesions and flat epithelial atypia of the breast are discussed along with their clinical importance. Common forms of invasive carcinomas, such as invasive ductal carcinoma and invasive lobular carcinoma; special types; and rarer forms are also discussed along with their clinical consequences.

*Intraoperative pathological examination* may be performed for the rapid diagnosis of breast malignancy, the assessment of the surgical margins of breast-conserving excision specimens, and the pathological analysis of sentinel lymph nodes. The most commonly used methods for intraoperative pathological examination of breast lesions are cytological and frozen section examinations in addition to gross analysis. The pathological examinations of sentinel lymph nodes necessitate careful gross examination and serial and/or step sectioning. Immunostaining using antibodies against pancytokeratin can also be performed. Sentinel lymph node metastases should be clearly defined as macro- or micrometastases or isolated tumor cells. The differential diagnosis of subtypes of metastasis and mimickers is detailed.

*Fibroepithelial tumors* of the breast represent a heterogeneous group of biphasic tumors composed of a proliferation of epithelial and stromal components. Fibroadenomas and phyllodes tumors constitute the major entities. These tumors are among the most challenging diagnostic lesions for pathologists. It can be difficult to make a clear microscopic distinction between fibroadenomas and benign phyllodes tumors. No reliable morphological features or immunohistochemical markers that predict phyllodes tumors are available.

A variety of reactive and neoplastic lesions of the breast are characterized by spindle cell proliferation. The pathologist must be aware of the clinical, radiological, and morphological overlap between reactive and *neoplastic spindle cell lesions* of the breast. In addition, metaplastic (spindle cell) carcinoma is far more common than spindle cell sarcoma in the breast. Among the vascular lesions of the breast, angiosarcoma is more common and may appear very bland, simulating a hemangioma. Core biopsy samples must be evaluated very carefully to interpret spindle and vascular lesions. In general, excision is recommended due to morphological overlap, and clinicopathological correlation is necessary for a correct diagnosis.

We would like to dedicate this book to postgraduate physicians in training to become breast cancer specialists. Some of the recommendations are controversial and the subject of ongoing trials. We hope this book stimulates today's young doctors to contribute to the research on which future books will be based.

Istanbul, Turkey  
Istanbul, Turkey  
Pittsburgh, PA, USA

Adnan Aydiner, MD  
Abdullah İğci, MD  
Atilla Soran, MD

---

# Contents

<b>1 Breast Anatomy and Physiology</b> .....	1
Kandace P. McGuire	
<b>2 Benign Diseases of the Breast</b> .....	15
Edward R. Sauter	
<b>3 Benign Breast Tumors</b> .....	23
Emilia Josefa Borromeo Diego	
<b>4 Epidemiology, Risk Factors, and Prevention</b> .....	57
Soley Bayraktar and Banu K. Arun	
<b>5 Breast Imaging and Image-Guided Biopsy Techniques</b> .....	89
Marie Ganott, Brandy Griffith, and Scott M. Rudzinski	
<b>6 Lobular Carcinoma In Situ</b> .....	123
Priscilla McAuliffe	
<b>7 Ductal Carcinoma In Situ</b> .....	131
Priscilla McAuliffe	
<b>8 Biology and Genetics of Breast Cancer</b> .....	145
A. Lale Dogan	
<b>9 Clinical Aspects of Estrogen and Progesterone Receptors and ERBB2 Testing</b> .....	161
Ebru Sari and Suayib Yalcin	
<b>10 Prognostic and Predictive Factors of Invasive Breast Cancer</b> .....	187
Yun Wu and Aysegul A. Sahin	
<b>11 Gene Arrays, Prognosis, and Therapeutic Interventions</b> .....	207
Cagatay Arslan, M. Kadri Altundag, and Y. Yavuz Ozisik	
<b>12 Bone Marrow Micrometastases and Circulating Tumor Cells</b> .....	229
Saadettin Kilickap, M. Kadri Altundag, and Y. Yavuz Ozisik	
<b>13 Pathology of Breast Cancer</b> .....	241
Sitki Tuzlali	

---

<b>14 Intraoperative Pathological Examination of Breast Lesions</b> .....	267
Ekrem Yavuz	
<b>15 Fibroepithelial Tumors of the Breast</b> .....	283
Sennur Ilvan	
<b>16 Mesenchymal Tumors of the Breast</b> .....	289
Zerrin Calay	
<b>Index</b> .....	295

---

## Contributors

**M. Kadri Altundag, MD** Department of Medical Oncology,  
Hacettepe University Cancer Institute, Ankara, Turkey

**Cagatay Arslan, MD** Department of Medical Oncology,  
Izmir University, Faculty of Medicine, Medical Park Hospital,  
Karsiyaka, Izmir, Turkey

**Banu K. Arun, MD** Department of Breast Medical Oncology,  
The University of Texas MD Anderson Cancer Center,  
Houston, TX, USA

**Soley Bayraktar, MD, MBA** Department of Breast Medical Oncology,  
The University of Texas MD Anderson Cancer Center,  
Houston, TX, USA

Department of Medical Oncology, Mercy Cancer Center,  
Ardmore, OK, USA

**Zerrin Calay, MD** Department of Pathology, Cerrahpasa  
School of Medicine, Istanbul University, Istanbul, Turkey

**Emilia Josefa Borrromeo Diego, MD** Department of Surgery, Division of  
Surgical Oncology, Magee-Womens Hospital, University of Pittsburgh  
Medical Center, Pittsburgh, PA, USA

**A. Lale Dogan, MD** Department of Basic Oncology, Hacettepe  
Cancer Institute, Hacettepe University, Ankara, Turkey

**Marie Ganott, MD** Department of Radiology, Magee-Womens Hospital,  
University of Pittsburgh Medical Center, Pittsburgh, PA, USA

**Brandy Griffith, DO** Department of Radiology, University  
of Pittsburgh Medical Center Hamot, Erie, PA, USA

Breast Imaging Department, Erie, PA, USA

**Sennur Ilvan, MD** Department of Pathology, Cerrahpasa School  
of Medicine, Istanbul University, Fatih, Istanbul, Turkey

**Saadettin Kilickap, MD, MSc** Department of Medical Oncology,  
Hacettepe University Cancer Institute, Ankara, Turkey



**Priscilla McAuliffe, MD, PhD, FACS** Department of Surgery, Division of Surgical Oncology, Magee-Womens Hospital, University of Pittsburgh Medical Center, Pittsburgh, PA, USA

Magee-Women's Surgical Associates, Bethel Park, PA, USA

**Kandace P. McGuire, MD, FACS** Department of Surgery, Division of Surgical Oncology, Magee-Womens Hospital, University of Pittsburgh Medical Center, Pittsburgh, PA, USA

**Y. Yavuz Ozisik, MD** Department of Medical Oncology, Hacettepe University Cancer Institute, Ankara, Turkey

**Scott M. Rudzinski, MD** Windsong Radiology Group, Williamsville, NY, USA

**Aysegul A. Sahin, MD** Department of Pathology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

**Ebru Sari, MD** Department of Medical Oncology, Diskapi Yildirim Beyazit Treatment and Research Hospital, Ankara, Turkey

**Edward R. Sauter, MD, PhD** Oncology and Surgery, Cancer Treatment and Prevention Center, Health Science Center, University of Texas, Tyler, TX, USA

**Sitki Tuzlali, MD** Department of Pathology, Istanbul Medical Faculty, Istanbul University, Istanbul, Turkey

**Yun Wu, MD, PhD** Department of Pathology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

**Suayib Yalcin, MD** Department of Medical Oncology, Hacettepe University Cancer Institute, Ankara, Turkey

**Ekrem Yavuz, MD** Department of Pathology, Istanbul Medical Faculty, Istanbul University, Istanbul, Turkey

Kandace P. McGuire

---

## Abstract

Breast development begins in the 5th and 6th weeks of fetal development and continues through puberty. Errors during development can lead to abnormal development or complete failure of breast development. The breast comprises several structures that are both functional and supportive. Some of these structures do not fully develop until pregnancy and lactation and regress or involute after lactation and at menopause. The anatomy of the breast and axilla are important in oncological surgery and must be considered during surgical planning to ensure the proper treatment of breast cancer.

---

## Keywords

Anatomy • Physiology • Embryology • Development • Lactation • Involution • Poland's syndrome • Axilla • Mastectomy • Sentinel node biopsy

---

## Embryology/Development

During the 5th and/or 6th week of fetal development, the two bands of thickened ectoderm referred to as the ectodermal primitive milk streak develop between the groin and the axilla [1, 2]. This remains in the thorax to become the

mammary ridge, whereas the remainder regresses in the human development [2].

The breast develops from an ingrowth of the ectoderm into the mesoderm to form a breast bud [1]. The glandular portion of the breast develops from the ectoderm. During the 12th week of development, 16–24 secondary buds will form off the primary bud [3].

---

K.P. McGuire, MD, FACS  
Department of Surgery, Division of Surgical  
Oncology, Magee-Womens Hospital,  
University of Pittsburgh Medical Center,  
300 Halket St., Pittsburgh, PA 15213, USA  
e-mail: [mcguirek2@upmc.edu](mailto:mcguirek2@upmc.edu)

## Sequence of Development

The development of the breast, summarized in Table 1.1, follows a stepwise progression

**Table 1.1** Embryonic breast development by gestational week

Gestational week	Breast development
5–6	Primitive milk streak develops from the ectoderm
7–8	Thickening of the mammary anlage
	Invagination into the mesoderm
	Growth of breast buds
12–16	Mesenchymal cells differentiate into the smooth muscle of the nipple-areola
	Secondary breast buds develop and branch
16–20	Tips of breast buds become the secretory alveoli
	Secondary mammary anlage differentiates into hair follicles and sebaceous and sweat gland elements
	Apocrine glands develop into Montgomery glands
20–32	Breast buds canalize and become lactiferous/mammary ducts
32–40	Parenchymal differentiation; lobules/alveoli develop
	Proliferation of mesenchyme forms the nipple-areola complex
	Pigmentation of the nipple-areola complex

beginning at the 5th week post-conception and continuing until birth. In weeks five and six, the primitive milk streak develops from a thickened band of ectoderm. Following development of the primitive milk streak in weeks seven and eight, the mammary anlage will thicken, and the mesoderm will invaginate. Simultaneously, the breast buds begin to grow. This process continues until weeks 12 through 16, when mesenchymal cells begin to differentiate into the smooth muscle of the nipple and areola. Secondary breast buds will further develop and branch but remain solid structures during this time period.

At week 16, the tips of the buds become the secretory alveoli. The secondary mammary anlage differentiates into hair follicles and the sebaceous and sweat gland elements. Apocrine glands develop to form the Montgomery glands. Beginning at week 20 of development and continuing until week 32, the breast buds will canalize to form lactiferous/mammary ducts. These

ducts open into a shallow mammary pit, which will become the nipple-areola complex. In the final weeks before birth, weeks 32 through 40, parenchymal differentiation occurs. The lobules and alveoli complete development. Finally, the nipple-areola complex develops via proliferation of the mesenchyme and becomes pigmented (Fig. 1.1) [2, 3].

## Developmental Anomalies

The development of a normal breast requires perfect adherence to the sequence of development described above. Should development stray from this pattern, anomalies may occur. The three most common developmental anomalies of the female breast are: (1) supernumerary breasts or nipples (polymastia/polythelia), (2) underdevelopment or lack of development of the breast, and (3) nipple inversion.


Supernumerary breasts/nipples (polymastia/polythelia) can occur in both genders and are referred to as accessory if they occur along the milk line (former primitive milk streak). They are referred to as ectopic if they occur elsewhere [3].

Accessory nipples (polythelia) occur in 2.5 % of the population and are much more common than accessory breasts (polymastia). Polythelia most commonly occurs in the thorax, while polymastia most commonly occurs in the axilla [2, 3].

Underdevelopment or hypoplasia of the breast can occur unilaterally or bilaterally and is usually clinically insignificant. However, severe unilateral hypoplasia of the breast can occur and is usually associated with hypoplasia of the pectoral muscle (lacking the lower third of the muscle) and deformity of the rib cage. This defect is termed *Poland's syndrome* because it was first recognized by Dr. Alfred Poland in 1841. Associated abnormalities of the hand (syndactyly and/or hypoplasia of the phalanges) may be present [2].

Amastia or lack of breast development is exceedingly rare. Athelia, a lack of development of the nipple-areola complex, can also occur, as can amazia, a lack of breast development in the presence of a nipple-areola complex [2, 3].

**Fig. 1.1** Fully developed breast lobular unit (From Townsend et al. [5]. Reproduced with permission from Elsevier)



Failure of the mesenchyme of the nipple-areola to proliferate and elevate the mammary pit above the skin results in an inverted nipple. This failure can occur unilaterally or bilaterally and occurs in 4 % of infants, both male and female [3].

## Anatomy

### Breast

The adult female breast lies between the second and sixth/seventh ribs. The base of the breast spans from the sternal border medially to the midaxillary line laterally and is encompassed by the superficial and deep fascia of the chest wall. Two-thirds of the breast lies anterior to the pectoralis major; the remainder lies anterior to the serratus anterior. A prolongation of the upper outer quadrant of the breast, referred to as the tail of Spence, extends into the axilla [3, 4].

### Components of the Breast

**Skin** – The skin is the most superficial layer of the breast. The dermis merges with the superficial fascia [3].

**Superficial fascia** – This layer lies just beneath the skin. It is continuous with the superficial abdominal and cervical fascia. Along with the deep fascia, it envelops the breast parenchyma [3].


**Breast parenchyma** – The parenchyma is composed of three principal tissue types: glandular epithelium, fibrous stroma, and supporting structures and fat.

Glandular epithelium comprises approximately 10–15 % of the adult female breast. It is composed of 15–20 lobes, which are subsequently composed of several lobules. These lobules are referred to as terminal ductules or acini, the milk-producing glands. The major milk ducts are lined with two layers of cuboidal epithelium, while the minor ducts have a single layer. The ductal epithelium is entirely surrounded by myoepithelial cells that serve to propel milk forward through the ducts. These cells are surrounded by a continuous basement membrane. Invasion through this membrane distinguishes invasive cancer from in situ carcinoma. The ducts widen under the nipple-areola complex to form the lactiferous sinuses and then exit through 10–15 orifices in the nipple.

The fibrous stroma and supporting structures are most commonly referred to as the suspensory ligaments of Cooper. These ligaments are fibrous bands of connective tissue that travel through the breast and insert into the dermis. Tumor involvement and contraction of these bands are responsible for the puckering noted at the site of a palpable breast lump.

The remainder of the breast is composed of adipose tissue (fat). The proportion of fat to

**Fig. 1.2** Components of the breast



glandular tissue increases with age and is maximal in the postmenopausal breast (Fig. 1.2) [1, 4, 5].

*Nipple-areola complex* – As described above, each lobe of the breast leads to a ductal structure that then widens to form a large lactiferous duct (2–4 mm) that continues to form a sinus. The sinus is lined with stratified squamous epithelium. This sinus then narrows as it passes into the ampulla of the nipple (0.4–0.7 mm).

The areola comprises a combination of sebaceous, sweat, and accessory glands that form the Montgomery tubercles. Smooth muscle fibers are contained in the areola and extend into the nipple, and these fibers are responsible for nipple erection. Erection is stimulated by the sensory nerve endings and Meissner's corpuscles, which are located within the dermis of the nipple [1, 3].


*Deep fascia* – This layer is deep to the breast parenchyma and envelops the pectoralis major. It is continuous with the deep abdominal fascia caudally and spans from the sternum to the axilla laterally and to the clavicle cranially [3].

## Neurovascular Structures

### Arterial

The arterial blood supply to the breast comes primarily from three sources: (1) anterior perforators of the internal mammary artery (responsible for approximately 60 % of the breast, mostly medial and central); (2) branches from the axillary artery, such as the highest and lateral thoracic, and the thoracoacromial artery (responsible for approximately 30 % of the breast, mostly the upper outer quadrant); and (3) lateral branches of the intercostal arteries (Fig. 1.3) [1–3].

**Fig. 1.3** Arterial supply and venous drainage of the breast



### Venous

Venous drainage typically mimics the arterial supply. Thus, the primary venous drainage consists of: (1) internal mammary perforating branches, (2) tributaries of the axillary vein, and (3) branches of the intercostal veins (Fig. 1.3) [1].

### Nervous

The sensory nerve supply to the breast is principally derived from the lateral cutaneous branches of the third through sixth intercostal nerves. Cranially, some sensory innervation is supplied by cutaneous branches of the cervical plexus. The nipple-areola complex is innervated by the fourth intercostal nerve [1, 3].

### Lymphatic Structures

The superficial lymphatic plexus that drains the skin of the breast and the nipple-areola complex is often referred to as Sappey's plexus. Lymph flows from the skin to the subareolar plexus and then into the interconnected deep lymphatic plexus that drains the breast parenchyma via the lymphatic vessels associated with the lactiferous

ducts. Approximately 97 % of the lymphatics from the breast drain to the axilla; the remaining 3 % drains to the internal mammary lymph nodes.

The internal mammary chain is located between the first and sixth intercostal spaces along the border of the sternum. The nodes are medial to the internal mammary vessels in the first two intercostal spaces and then become lateral to the vessels in spaces 3–6 (Fig. 1.4) [2, 4, 5]. The anatomy of the axilla and axillary lymph nodes will be discussed in the following section.


### Axilla

The axilla is an important component of breast anatomy. Directly contiguous with the breast, the lymph nodes within the axilla provide a rich drainage basin for the breast. The borders of the axilla, which define the extent of axillary dissection, are as follows.

*Lateral* – The axillary fat pad and the bicipital groove of the humerus



**Fig. 1.4** Lymphatic drainage of the breast



*Medial* – The serratus anterior and the second to sixth ribs

*Superior* – The apex of the axilla bordered by the clavicle, the scapula, and the first rib

The apex of the axilla can also be defined by the costoclavicular ligament, which is also called Halsted's ligament. The axillary vein is the superior extent of the modified radical axillary dissection. Medial to Halsted's ligament, the axillary vein becomes the subclavian vein.

*Anterior* – The pectoralis (major and minor) and subclavius muscles and the clavipectoral fascia. The clavipectoral fascia envelops the subclavius and pectoralis minor and is often referred to as the costocoracoid membrane. The lateral band of the clavipectoral fascia between the first rib and the coracoid process is called the costocoracoid ligament.

*Posterior* – The scapula, the subscapularis, the latissimus dorsi, and the teres major. The axillary fascia lying across the base of the axillary pyramid will envelop the pectoralis major and then the latissimus dorsi. It forms the dome of the axilla. Occasionally, there


can be a muscular connection between this fascia and the clavipectoral fascia, which is referred to as the suspensory ligament of the axilla [3, 5, 6].

### Axillary Lymph Nodes

There are several groups of lymph nodes within the axilla. These nodes can be grouped as the apical or subclavicular nodes, which are located medial to the pectoralis minor muscle, or the axillary vein lymph nodes, which run along the axillary vein between the pectoralis minor and the humerus. The interpectoral or Rotter's nodes lie between the pectoralis major and minor muscles. The central axillary nodes are found beneath the border of the pectoralis major muscle and below the pectoralis minor. The external mammary nodes lie over the axillary tail of Spence. Intramammary lymph nodes and paramammary lymph nodes can also be found in the fat layer over the upper, outer quadrant of the breast (Fig. 1.5).

For surgical dissection purposes, there are three lymph node levels of the axilla, which are

**Fig. 1.5** Axillary lymph node groups (From Townsend et al. [5]. Reproduced with permission from Elsevier)



all defined by their relationship to the pectoralis minor muscle. Level I nodes are found lateral to the edge of the pectoralis minor. This level includes external mammary, subscapular, and lateral axillary lymph nodes. Level II nodes are located posterior to the pectoralis minor. This level includes the central axillary lymph nodes. Level III nodes are medial and superior to the pectoralis minor. This level includes the subclavicular or apical lymph nodes [2, 6].

### Structures Within the Axilla

The *axillary lymph nodes* are divided into several different groups and levels as described above and are variable in number. The maximum number identified and removed during a radical mastectomy is approximately 50, including the level I, II, and III axillary lymph nodes.

The *axillary vein* defines the superior border of the axilla during axillary dissection. It lies posterior and caudal to the brachial plexus. The axillary vein is often paired or branches during its course through the axilla.

The *thoracodorsal nerve/neurovascular bundle* innervates the latissimus dorsi and should be preserved during axillary dissection. It runs posterior to the axillary vein and medial to the subscapular vein.

The *long thoracic nerve/neurovascular bundle* innervates the serratus anterior and should be preserved during axillary dissection. If sacrificed, it will lead to “winging” of the scapula. It runs longitudinally over the serratus anterior and can be found during dissection in the axillary fat pad approximately 7 or 8 cm deep to the lateral edge of the pectoralis minor. As the long thoracic nerve/neurovascular bundle continues caudally, it will become more anterior.

The *intercostobrachial nerves* provide sensory innervation to the medial portion of the upper arm. These nerves run parallel to the axillary vein between the chest wall and the arm. One or more of these nerves run through the axillary fat pad and may be difficult to dissect away from lymph nodes. If sacrificed, either hypo- or hyperesthesia of the posterior axillary web and the medial/upper arm can result [4, 6].

## Physiology

### Physiological Breast Development

Breast development is stimulated by a variety of hormones that are upregulated during the beginning stages of puberty. Estrogen and



progesterone are the main hormones responsible for breast growth and development during this time. Estrogen stimulates ductal development; progesterone stimulates lobular development and epithelial differentiation.

At the onset of puberty, the hypothalamic-pituitary axis becomes less sensitive to the negative feedback of estrogen. This desensitization leads to an increase in gonadotropin-releasing hormone (GnRH) from the hypothalamus. This increase in GnRH stimulates the release of luteinizing hormone (LH) and follicle-stimulating hormone (FSH) from the anterior pituitary, which in turn leads to an increase in estrogen and progesterone release, thus stimulating breast development, among other developmental changes.

During pregnancy and lactation, prolactin is primarily responsible for upregulating hormone receptors and stimulating epithelial development and lactogenesis in the breast [1].

### **Abnormal Breast Development/ Gynecomastia**

Gynecomastia refers to male breast hypertrophy and can be caused by numerous factors.

1. **Physiological gynecomastia:** This can occur in the neonatal, pubertal, and senescent periods. Neonatal hypertrophy occurs in response to maternal estrogen. Pubertal hypertrophy occurs due to a relative excess of estradiol to testosterone. Senescent hypertrophy occurs in response to falling testosterone levels associated with aging. The enlargement is usually unilateral in puberty but bilateral in senescence. This usually does not require surgery unless the enlargement is associated with a mass by physical exam or mammogram, fails to regress, or is cosmetically unacceptable.
2. **Pathologic gynecomastia:** There are a number of pathological causes of gynecomastia, including true hermaphroditism, testicular tumors, adrenal cortical neoplasms, lung or hepatocellular carcinoma, endocrine disorders,

cirrhosis, and nutritional deficiencies (estrogen excess states). Hypogonadism, as observed in congenital syndromes such as Klinefelter (XXY) syndrome or ACTH deficiency, can also cause gynecomastia. Secondary testicular failure from trauma, radiation, or untreated cryptorchidism can also cause hypertrophy (androgen deficiency states). Renal failure and other systemic diseases can lead to gynecomastia, as can drugs that provide exogenous estrogen or stimulate estrogen synthesis (e.g., digoxin, estrogens, anabolic steroids, marijuana, and HCG) or that inhibit the activity or production of testosterone (e.g., cimetidine, ketoconazole, phenytoin, spironolactone, antineoplastic drugs, and diazepam). Some drugs, such as reserpine, theophylline, verapamil, tricyclic antidepressants, and furosemide, lead to gynecomastia through idiopathic mechanisms [3, 5].

### **Physiology of Puberty**

As described above, pubertal development of the breast (thelarche) begins with the stimulation of estrogen and progesterone production via the hypothalamic-pituitary axis. At this time, the breast is composed primarily of dense fibrous stroma and scattered ducts lined with epithelium. Estrogen stimulates growth of the ductal epithelium. Buds form off the terminal ductules and will eventually become breast lobules. Periductal connective tissue grows and becomes more elastic. Some studies suggest that while estrogen promotes growth of ducts, estrogen and progesterone synergistically promote full ductular-lobular-alveolar development in the breast.

There are three distinct types of breast lobules in the human breast, and the proportion of each is related to a woman's parity and hormonal status. During puberty, the breast develops mostly type I (virginal) lobules, which consist of a cluster of 11 alveolar buds around a terminal duct. These lobules have a much higher rate of proliferation than type 2 or 3 lobules.

## Physiology of the Menstrual Cycle

The postpubertal breast contains fat, stroma, lactiferous ducts, and lobular units. The menstrual cycle affects not only the uterus and uterine lining but also the breast. During the follicular phase (days 4–14), levels of estrogen increase, stimulating epithelial proliferation/sprouting and an increased mitotic rate. During the luteal phase (days 15–28), progesterone increases, while estrogen abates. At this time, mammary ducts dilate, and alveolar epithelial cells differentiate into secretory cells. Often, lipid droplets accumulate, and some intralumenal secretion occurs. During this time, estrogen also exerts a histamine-like effect on the breast parenchyma, resulting in increased blood flow and breast edema just prior to the onset of menses.

During this time, type I lobules continue to predominate. The increased rate of cellular proliferation in these lobules may partly explain the differences in breast cancer rates based on parity and age at first live birth.

## Physiology of Pregnancy

During pregnancy, there is a decrease in fibrous stroma along with an increase in new acini/lobules. In the first trimester, ducts sprout and branch, and lobules develop as estrogen increases. Breast enlargement is significant, with dilatation of superficial veins and breast edema. The nipple-areola complex darkens and begins to enlarge. Type 3 lobules (with an average of 80 acini) begin to develop during this time and are referred to as alveoli.

In the second trimester, levels of progesterone increase, as does lobular formation. The alveoli begin to form colostrum, which is composed of desquamated eosinophilic cells, plasma cells, leukocytes, and epithelial cells.

In the third trimester, the alveoli continue to produce colostrum. At this time, epithelial differentiation is completed, resulting in the

development of secretory cells that produce and secrete milk proteins. Oxytocin increases over the last trimester, resulting in the proliferation of myoepithelial cells surrounding the ductal structures, which propel the milk forward toward the nipple-areola complex.

## Physiology of Lactation

After birth, there is a sudden decrease in the levels of estrogen, progesterone, and placental lactogen, coupled with an increase in prolactin, which induces the production and secretion of milk. Hormonal levels reach their lowest levels at about the fifth postpartum day, with a concomitant decrease in prolactin-inhibiting factor (PIF). This decrease results in the secretion of prolactin. Along with additional growth factors, prolactin secretion results in the accumulation of colostrum and, subsequently, milk in the alveoli and ducts. Stimulation of the nipple-areola complex stimulates the release of oxytocin and the contraction of the myoepithelial cells surrounding the ductal system. Upon cessation of breast-feeding (weaning), levels of prolactin and oxytocin fall. Retained secretions are removed via phagocytosis. Atrophy of the glandular, ductal, and stromal elements is observed. The secretory cells responsible for milk production undergo apoptosis. However, the type 3 lobules persist.

## Physiology of Menopause

After menopause, the breast parenchyma regresses and is replaced by adipose tissue. This replacement occurs by involution of the ductal, glandular, and stromal elements/connective tissue of the breast. The ductal system remains but undergoes atresia, with collapse of the lobular units. Type 1 lobules again predominate, as in the nulliparous breast. The number of lymphatic channels through the breast parenchyma also decreases [2, 4, 5].

## Surgical/Oncological Considerations

### Tumor Location Within the Breast

The adult breast develops in a conical form, with epithelial/ductal tissue in each quadrant of the breast. The axillary tail of Spence, as discussed previously, is an extension of the upper outer quadrant of the breast over the axilla. Because of this extension, the upper outer quadrant contains significantly more epithelial tissue than the other quadrants. Thus, this quadrant is the most frequent site of breast neoplasms and harbors more than half of both benign and malignant tumors [3, 4, 7].


The location of the tumor within the breast can also affect the ability to perform breast conservation (segmental mastectomy). In general, segmental mastectomy can be performed with good cosmetic outcome when the tumor volume is less than 20 % of the volume of the breast [8–10]. However, this percentage can vary with tumor location. Tumors in the upper outer quadrant are much easier to resect with good cosmetic outcome because there is a great amount of surrounding tissue in the region. Tumors that lie in the lower quadrants, particularly the lower inner quadrant, have little surrounding parenchyma, and excisions in these regions can lead to significant retraction and poor cosmetic outcome after surgery and radiation are performed. Partial breast reconstruction techniques, such as small latissimus dorsi flaps and local advancement flaps, can replace volume, particularly in the outer quadrants. However, these techniques require more extensive surgery, and the patient may be better served by mastectomy and whole breast reconstruction in this situation [11–13]. Depending on tumor location, volume loss can also be addressed by oncoplastic surgical techniques ranging from simple local advancement flaps to concurrent reduction mammoplasty [14–16].

## Borders of Mastectomy

There are three different types of mastectomy, all with different extents of dissection:

1. Simple or total mastectomy – Several skin incisions can be made, including peri-areolar (skin-sparing) or elliptical. Dissection is performed along the superficial fascial plane superiorly to the clavicle; medially to the sternal edge; inferiorly to the inframammary fold, just cranial to the insertion of the rectus sheath; and laterally to the edge of pectoralis major muscle. The deep border of dissection is the deep fascial plane, just superficial to the pectoralis major muscle. This method of mastectomy removes nearly 100 % of the breast epithelial/stromal tissue while preserving the axillary fat pad and axillary lymph nodes.

This approach is used most often in modern practice and is often combined with immediate reconstruction. When plastic surgery is involved, it is imperative that a multidisciplinary approach to surgery be used and that both oncological and plastic surgeons are involved in incision planning. This is particularly true in the case of nipple-sparing mastectomy, in which several incisions can be used [17–23]. An important anatomic consideration is the blood supply to the nipple and areola, which can vary greatly from patient to patient. Nipples that derive most of their blood supply from the underlying parenchyma are likely to suffer partial or complete necrosis after nipple-sparing mastectomy, whereas the viability of those that derive blood supply from the surrounding skin will be largely unaffected. Blood flow can be assessed either preoperatively or intraoperatively with imaging systems that detect fluorescent dye (usually indocyanine green) injected intravenously. The resulting perfusion patterns can help guide incision planning and also identify candidates for nipple-sparing surgery [24] (Fig. 1.6).



**Fig. 1.6** Nipple perfusion patterns

2. Modified radical mastectomy – This operation can be performed through the same incisions as a total or simple mastectomy. The elliptical incision can be extended superolaterally toward the axilla to facilitate axillary dissection. The superior, medial, inferior, and deep borders of the dissection are the same as in a total mastectomy. However, the modified radical mastectomy involves the removal of levels I and II axillary lymph nodes; thus, the lateral border of dissection is the latissimus dorsi extending superiorly to the axillary vein.
3. Halsted radical mastectomy – This operation is rarely performed and very rarely described in current surgical texts and atlases. The

superior, medial, inferior, and lateral borders are the same as in a modified radical mastectomy. However, the deep dissection includes the pectoralis major and minor muscles. The axillary dissection includes levels I, II, and III of the axillary lymph nodes and is thus extended superior and medial to the axillary vein. This operation is performed only in the presence of locally advanced cancers that involve one or both pectoralis muscles [6].

### Sentinel Node Biopsy

Sentinel lymph node biopsy was originally described as a method for detecting the lymphatic

drainage of melanoma. It has been modified for use in breast cancer using the following method.

1. Isosulfan or methylene blue dye and/or technetium-99 are injected preoperatively into the superficial lymphatic plexus, either into the subareolar plexus or around the tumor.
2. This injection *can* be followed by lymphoscintigraphy to identify the area into which the radioactive dye has drained. Lymphoscintigraphy requires allowing the technetium 1 h or more to travel through the breast lymphatics into the axillary and/or internal mammary lymph nodes. As noted above, 97 % of the breast drains to the axillary region; thus, this step is not necessary. It can be helpful in inner quadrant tumors, which more commonly drain to the internal mammary chain, and in patients with previous breast surgery, which might interfere with the normal lymphatic drainage of the breast.
3. Once in the operating room, the radioactive-sensitive probe can be used to localize the area in the axilla with the highest concentration of technetium colloid.
4. An incision is made in this area through the skin, the subcutaneous tissue, and the clavicular fascia. Once the axillary fat pad is identified, the probe can be used to localize the lymph node(s) with the highest concentration of radioactive dye. Those lymph node(s) that are both “hot” (radioactive) and “blue” (have taken up the blue dye) should be removed and sent for pathologic analysis (frozen, touch prep, or permanent). If these “sentinel” lymph nodes show evidence of malignancy, then a full axillary dissection as described for modified radical mastectomy is performed at that time or during a separate operation.

This method is based upon the anatomy of the breast lymphatic system. As described previously in this chapter, the lymph flows from the skin to the subareolar plexus and then into the interconnected deep lymphatic plexus that drains the breast parenchyma via the lymphatic vessels associated with the lactiferous ducts. Therefore,

any lymphatic drainage from the breast must travel through both the superficial and deep lymphatic plexuses before leaving the breast, and an injection into the superficial lymphatic plexus will identify the main route of drainage for the breast. This drainage is standard and reproducible. Once the channels reach the axilla, they drain first to the “sentinel” lymph node(s) in either levels I or II of the axilla before draining to the remainder of the axilla. If no cancer is found in the sentinel node, there is a >95 % likelihood that no other cancer exists in the axilla [2, 4, 5, 25].

Sentinel lymph node biopsy has several advantages over axillary dissection in appropriately selected patients, and in fact, axillary dissection has become increasingly rare. Several landmark trials have established the efficacy of sentinel lymph node biopsy in the setting of breast cancer. The great advantage of sentinel lymph node biopsy is the reduction in risk of postoperative arm (and to some extent breast) lymphedema. The rich lymphatic network experiences less disruption. In early studies, most notably NSABP B-32 and ACOSOG Z0010, the lymphedema rates after sentinel lymph node biopsy varied from 8 to 12 %, whereas axillary dissection resulted in lymphedema rates of 14–42 % [26, 27]. However, for patients who require axillary dissection, techniques can identify the lymphatics that primarily drain the arm (outside of the level III lymph nodes, which are excluded in modern axillary dissection if they are not clinically involved due to the low incidence of involved nodes in this region and the high incidence of arm lymphedema after radical mastectomy) [28, 29]. The best-described technique is axillary reverse lymphatic mapping (ARM). This technique involves injecting a small amount of blue dye in the subcutaneous tissue of the volar surface of the upper arm prior to lymphatic surgery. The axillary lymph nodes that drain the arm can be identified and frequently preserved. This can result in a lymphedema rate of 2.4 % after ALND in which the identified arm lymphatics are preserved, much lower than previous reports [30–33].

## References

- Brunicaardi FC, Andersen DK, Billiar TR, Dunn DL, Hunter JG, Matthews JB, et al., editors. *Schwartz's principles of surgery*. 8th ed. New York: McGraw-Hill, Health Pub. Division; 2005.
- Harris JR, Lippman ME, Morrow M, Osborne KC, editors. *Diseases of the breast*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2009.
- Skandalakis JE, Colborn GL, Weidman TA, Foster J, Roger S, Kingsnorth AN, Skandalakis LJ, et al., editors. *Skandalakis surgical anatomy: the embryologic and anatomic basis of modern surgery*. Athens: McGraw-Hill Professional Publishing; 2004.
- Mulholland MW, Lillemoe KD, Doherty GM, Maier RV, Upchurch GR, editors. *Greenfield's surgery: scientific principles and practice*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2006.
- Townsend CM, Beauchamp RD, Evers BM, Mattox KL, editors. *Sabiston textbook of surgery*. Philadelphia: Saunders/Elsevier; 2008.
- Zollinger J, Robert M, Zollinger S, Robert M. *Zollinger's atlas of surgical operations*. 8th ed. New York McGraw-Hill: Medical Publishing Division; 2003.
- Sohn VY, Arthurs ZM, Sebesta JA, Brown TA. Primary tumor location impacts breast cancer survival. *Am J Surg*. 2008;195(5):641–4.
- Ozmen T, Polat AV, Polat AK, Bonaventura M, Johnson R, Soran A. Factors affecting cosmesis after breast conserving surgery without oncoplastic techniques in an experienced comprehensive breast center. *Surgeon J Royal Coll Surg Edinb Ireland*. 2014.
- Polat AV, Soran A, Andacoglu O, Kamali Polat A, McGuire K, Diego E, et al. The importance of pre-operative needle core breast biopsy results on resected tissue volume, margin status, and cosmesis. *J BUON Off J Balkan Union Oncol*. 2013;18(3):601–7.
- Taylor ME, Perez CA, Halverson KJ, Kuske RR, Philpott GW, Garcia DM, et al. Factors influencing cosmetic results after conservation therapy for breast cancer. *Int J Radiat Oncol Biol Phys*. 1995;31(4):753–64.
- Losken A, Hamdi M. Partial breast reconstruction: current perspectives. *Plast Reconstr Surg*. 2009;124(3):722–36.
- Munhoz AM, Montag E, Filassi JR, Gemperli R. Current approaches to managing partial breast defects: the role of conservative breast surgery reconstruction. *Anticancer Res*. 2014;34(3):1099–114.
- Song HM, Styblo TM, Carlson GW, Losken A. The use of oncoplastic reduction techniques to reconstruct partial mastectomy defects in women with ductal carcinoma in situ. *Breast J*. 2010;16(2):141–6.
- Munhoz AM, Montag E, Gemperli R. Oncoplastic breast surgery: indications, techniques and perspectives. *Gland Surg*. 2013;2(3):143–57.
- Silverstein MJ. How I, do it: oncoplastic breast-conservation surgery. *Ann Surg Oncol*. 2010;17 Suppl 3:242–4.
- Silverstein MJ, Mai T, Savalia N, Vaince F, Guerra L. Oncoplastic breast conservation surgery: the new paradigm. *J Surg Oncol*. 2014;110(1):82–9.
- Blechman KM, Karp NS, Levovitz C, Guth AA, Axelrod DM, Shapiro RL, et al. The lateral inframammary fold incision for nipple-sparing mastectomy: outcomes from over 50 immediate implant-based breast reconstructions. *Breast J*. 2013;19(1):31–40.
- Dent BL, Small K, Swistel A, Talmor M. Nipple-areolar complex ischemia after nipple-sparing mastectomy with immediate implant-based reconstruction: risk factors and the success of conservative treatment. *Aesthet Surg J Am Soc Aesthet Plast Surg*. 2014;34(4):560–70.
- Dutton W, Ghareeb PA, McClellan WT. The lazy lateral incision: an innovative approach to the skin-sparing mastectomy. *W V Med J*. 2013;109(6):30–3.
- Garwood ER, Moore D, Ewing C, Hwang ES, Alvarado M, Foster RD, et al. Total skin-sparing mastectomy: complications and local recurrence rates in 2 cohorts of patients. *Ann Surg*. 2009;249(1):26–32.
- Petit JY, Veronesi U, Rey P, Rotmensz N, Botteri E, Rietjens M, et al. Nipple-sparing mastectomy: risk of nipple-areolar recurrences in a series of 579 cases. *Breast Cancer Res Treat*. 2009;114(1):97–101.
- Rusby JE, Smith BL, Gui GP. Nipple-sparing mastectomy. *Br J Surg*. 2010;97(3):305–16.
- Proano E, Perbeck LG. Influence of the site of skin incision on the circulation in the nipple-areola complex after subcutaneous mastectomy in breast cancer. *Scand J Plast Reconstr Surg Hand Surg Nordisk plastikkirurgisk forening [and] Nordisk klubb for handkirurgi*. 1996;30(3):195–200.
- Wapnir I, Dua M, Kierny A, Paro J, Morrison D, Kahn D, et al. Intraoperative imaging of nipple perfusion patterns and ischemic complications in nipple-sparing mastectomies. *Ann Surg Oncol*. 2014;21(1):100–6.
- Cox CE. Lymphatic mapping in breast cancer: combination technique. *Ann Surg Oncol*. 2001;8(9 Suppl):67S–70.
- Ashikaga T, Krag DN, Land SR, Julian TB, Anderson SJ, Brown AM, et al. Morbidity results from the NSABP B-32 trial comparing sentinel lymph node dissection versus axillary dissection. *J Surg Oncol*. 2010;102(2):111–8.
- Wilke LG, McCall LM, Posther KE, Whitworth PW, Reintgen DS, Leitch AM, et al. Surgical complications associated with sentinel lymph node biopsy: results from a prospective international cooperative group trial. *Ann Surg Oncol*. 2006;13(4):491–500.
- DiSipio T, Rye S, Newman B, Hayes S. Incidence of unilateral arm lymphoedema after breast cancer: a systematic review and meta-analysis. *Lancet Oncol*. 2013;14(6):500–15.
- Barros AC, Andrade FE, Bevilacqua JL, Barros MA, Piatto JR, Santos DR, et al. Radicality effect of adding an interpectoral to a subpectoral approach for dissection of level III axillary lymph nodes in breast cancer. *Tumori*. 2013;99(4):500–4.

30. Boneti C, Badgwell B, Robertson Y, Korourian S, Adkins L, Klimberg V. Axillary reverse mapping (ARM): initial results of phase II trial in preventing lymphedema after lymphadenectomy. *Minerva Ginecol.* 2012;64(5):421–30.
31. Gennaro M, Maccauro M, Sigari C, Casalini P, Bedodi L, Conti AR, et al. Selective axillary dissection after axillary reverse mapping to prevent breast-cancer-related lymphoedema. *Eur J Surg Oncol.* 2013; 39(12):1341–5.
32. Kuusk U, Seyednejad N, McKeivitt EC, Dingee CK, Wiseman SM. Axillary reverse mapping in breast cancer: a Canadian experience. *J Surg Oncol.* 2014; 110(7):791–5.
33. Ochoa D, Korourian S, Boneti C, Adkins L, Badgwell B, Klimberg VS. Axillary reverse mapping: five-year experience. *Surgery.* 2014;156(5):1261–8.

Edward R. Sauter

---

### Abstract

Historically, physical examinations of the breast, conducted both by the individual (breast self-examination, BSE) and the healthcare provider (clinical breast examination, CBE), have been the primary tools for the diagnosis of breast cancer. With the advent of breast imaging (mammography, ultrasound, breast MRI), which is more sensitive than BSE and CBE in breast cancer detection, many of the observations historically described on physical examination are not common in the smaller cancers seen in current practice. Physical examination has become an adjunct to imaging for the early detection of breast cancer. Indeed, BSE does not increase the detection of cancer, and CBE is no longer used to diagnose breast cancer. Rather, it is used as a screening test that can identify areas with breast cancer. While CBE is less sensitive than mammography, it is nonetheless the primary mode of detecting the 15 % of breast cancers that are missed by mammography.

Signs and symptoms of benign breast conditions include nipple discharge, inflammatory conditions, and breast masses. The key for each of these symptoms is appropriate diagnosis and treatment. The most important consideration is to exclude the presence of cancer. Changes during pregnancy and lactation make evaluation of the breast more difficult, both because of enlargement and the increased ratio of glands to total breast volume. These changes make the breast more nodular and dense. Mammography sensitivity decreases, and for pregnant women, the radiation of mammography should be avoided whenever possible. Pregnancy-associated breast cancer is a unique problem because the cancer diagnosis

---

E.R. Sauter, MD, PhD  
Oncology and Surgery,  
Cancer Treatment and Prevention Center,  
Health Science Center, University of Texas,  
11937 US Hwy 271, Tyler, TX 75708, USA  
e-mail: [Edward.sauter@uthct.edu](mailto:Edward.sauter@uthct.edu)



may be delayed due to changes in the breast and risk of radiation to the fetus and also because these tumors appear more biologically aggressive than most breast cancers.

### Keywords

Benign breast disease • Breast imaging • Nipple discharge • Breast inflammation • Fat necrosis • Diabetic mastopathy

## Physical Examination of the Breast

Physical examination of the breast can be performed within the context of a clinical examination or by the individual. The former setting is often termed a clinical breast examination (CBE) and the latter a breast self-examination (BSE). Before the era of routine standardized breast imaging, physical examination was generally the primary method of diagnosing breast cancers. Breast imaging lacked standardization until the creation of the BI-RADS system in 1993 [1].

CBE is currently used as a screening test that can identify areas with breast cancer. While CBE is less sensitive than mammography, it is nonetheless the primary mode of detection for the 15 % of breast cancers that are missed by mammography [2].

BSE was promoted to allow women to identify their own cancers at an early stage. Although BSE was anticipated to work for many reasons, randomized trials of BSE with increasingly sophisticated procedures for retraining and sustaining BSE practice have demonstrated that although there is increased identification of benign breast abnormalities, there is no increased identification of cancer and no improvement in breast cancer-specific survival [2].

## Examination of the Breast

The purpose of a CBE is to detect changes in the consistency of the breast tissue. Other tests are needed if an area of asymmetry is found [2]. CBE includes visual inspection of the breast first with the patient sitting and then supine. With the patient sitting, the position and contour of the

breast are observed with changes in posture and arm position. Unexpected changes in breast contour should be further evaluated by breast palpation. Supine is the optimal patient position for breast palpation. Alterations in breast consistency should be further evaluated with imaging studies and/or biopsy. For women with pathological nipple discharge (PND) (discussed in more detail below), CBE can be used to localize the source of the discharge. In the absence of PND, CBE seeks changes in the visual appearance of the breasts and/or signs of local metastases, e.g., to regional lymph nodes. Signs of advanced cancer are rarely encountered during CBE of a truly asymptomatic patient.

Although a patient's personal and family history influences the probability that cancer will be found, history is not relevant to the CBE or when interpreting the CBE results because the majority of breast cancer occurs in women without known risk factors. Models composed of silicone or other materials are widely used to teach both CBE and BSE skills, although their benefits in early cancer detection are largely unproven.

## Signs and Symptoms of Breast Disease

Many of the features that students are taught to identify on CBE, such as skin dimpling, peau d'orange, hardness, and fixed mass, were first described before mammography and apply to advanced breast cancers, but these features are generally not applicable to the earlier-stage cancers that are typically encountered in current practice [2]. Other findings can occur with early-stage disease, such as PND, breast asymmetry, and

masses. Breast inflammation is most often benign but can be an indicator of cancer. The most important consideration in any breast abnormality is to exclude carcinoma. Inflammatory carcinoma can present as erythema with or without pain in the breast. Biopsy of the skin and any associated underlying mass should be performed. The breast is also often inflamed in response to radiation therapy to treat breast cancer. In this chapter, we will focus on breast symptoms that are often, although not always, related to benign disease, including mastalgia, nipple discharge, breast inflammation, and breast masses.

## PND

CBE does not seek to determine if nipple discharge can be elicited. Nipple discharge is of interest only if it is spontaneous. Fluid can be elicited by many women from their breast with massage. This fluid is not spontaneous nipple discharge (SND), which is of concern because of its potential implications related to disease, but rather intraductal fluid that is present in the breasts of all women and is often termed nipple aspirate fluid (NAF) because a modified breast pump is sometimes used to collect it [3]. However, if a woman reports noticing fluid on her bra that she did not elicit, particularly from one but not the other breast, this symptom should be further investigated, and pressure applied sequentially in a circular pattern can help localize the source of the spontaneous discharge.

SND is common, accounting for nearly 7 % of all breast symptoms. SND is most often physiological, particularly if bilateral and from multiple breast ducts. Carcinoma prevalence in women with SND varies primarily according to the criteria used to assess whether the discharge is physiological or pathological. The criteria usually include whether the discharge is from one or both breasts and from one or multiple nipple ducts and the characteristics of the discharge (bloody vs. nonbloody; clear or serous vs. white, yellow, or green). Additional criteria include the presence of an associated mass or an imaging abnormality. Because the criteria distin-

guishing pathological from physiological discharge vary from publication to publication, the published incidence of carcinoma among women with SND also varies [4].

PND is more likely when the discharge is unilateral and from one milk duct. The most common diagnosis in women with PND is papilloma. Among breast lesions, papilloma is unique in its frequent presentation as PND. NAF originates in the breast ducts similarly to SND but is less voluminous and can be obtained from essentially all nonlactating women after a learning period [4]. In contrast to NAF cytology, in which false positives are rare, PND cytology in patients with papillomas is occasionally falsely interpreted as containing malignant cells [5] because exfoliated cells from papillomas can appear quite abnormal when not viewed in the context of histological architecture. Although older studies reported that bloody nipple discharge was more commonly associated with cancer than nonbloody discharge, more recent studies challenge this belief [4].

## Breast Inflammation

The breast is most fundamentally an appendage of the skin. Many systemic inflammatory conditions present on the skin of the breast, including sarcoidosis, vasculitides, diabetes, and infections [6]. Sarcoidosis of the breast occurs in less than 1 % of individuals with sarcoid and usually presents as a breast mass, less often with skin dimpling and peau d'orange changes [7]. Primary sarcoidosis of the breast without systemic manifestations can occur but is uncommon. Giant cell (temporal) arteritis can manifest in the breast, typically presenting as painful breast masses. Systemic symptoms related to giant cell arteritis are usually present. Other vasculitides such as polyarteritis nodosa and Wegener's granulomatosis involving the breast have also been reported [7]. Diabetic mastopathy is most often observed in premenopausal women with type I diabetes and classically presents as a hard painless mass in one or both breasts [8]. Diagnosis requires biopsy, and treatment is mass excision. IgG4-related autoimmune syndrome can present in the breast, commonly as a tender breast mass [6].

Mastalgia (breast pain) accounts for two-thirds of all physician visits for breast symptoms but is not a risk factor for breast cancer [9]. The pain may be cyclical. Cyclic pain is most often related to the menstrual cycle, is bilateral and diffuse, and occurs during the luteal phase as rising progesterone levels increase the water content in the breast [9]. Noncyclic pain may be in one or both breasts and has been associated with a variety of medications, including oral contraceptives, female hormones, psychotropics, and cardiovascular medications. Larger-breasted woman may develop ligamentous pain if an adequate support bra is not worn. If the clinical workup, including history of diffuse pain, CBE, and mammography (in women over 40), suggests a benign etiology, treatment is generally supportive, most often starting with confirming that the woman is wearing a supportive, well-fitted brassiere. Acetaminophen or a nonsteroidal anti-inflammatory medication is often effective [9].

Fat necrosis generally presents as a painful mass in the breast. There may be multiple masses, with or without skin retraction. Fibrosis and calcification are common. Women with large breasts are at highest risk [7], and necrosis usually occurs following trauma. The trauma can include cyst aspiration, breast massage, mammography, radiation therapy, biopsy, implant removal, and reduction mammoplasty. A biopsy is often needed to exclude malignancy.

Mastitis occurs most commonly in women of childbearing age. It can occur during pregnancy and lactation and in women who are not pregnant or lactating [6], although in the latter group, most women report giving birth within 5 years of mastitis onset. The normal bacterial flora of breast tissue resembles that of the skin, and coagulase-negative staph and *Propionibacterium* species are the predominant organisms [6]. Mastitis is associated with pain, breast swelling, mass(es), and inflammation that resembles an abscess. Ipsilateral axillary lymph nodes are enlarged in approximately one-sixth of patients [7]. Mastitis frequently leads to surgical interventions to biopsy the lesion(s) to exclude malignancy and drainage procedures to treat the

inflammatory process. The interventions can lead to breast scarring and/or shrinkage. Resolution occurs in only approximately half of benign conditions. The various forms of benign inflammatory processes (periductal mastitis, Zuska's disease, comedomastitis, duct ectasia, mastitis obliterans, lactiferous fistula, and idiopathic granulomatous lobular mastitis) may be part of a common disease process termed mammary duct-associated inflammatory disease sequence (MDAIDS) [7]. Smoking is linked to MDAIDS, and severe disease occurs almost exclusively in heavy smokers; thus, the cessation of smoking is a very important part of the treatment process [7]. These conditions may result from lactiferous duct obstruction, resulting in duct distention, inflammation, and ultimately rupture. SND may be observed. Treatment requires excision of the affected ductal system. Simple incision and drainage are associated with a high rate of mastitis recurrence and breast scarring.

## Breast Masses and Breast Imaging

Approximately 70–80 % of the breast lesions detected by physical examination or imaging and biopsied [10] are benign. Most solid or complex cystic breast lesions should undergo biopsy. Possible exceptions are lesions in young women that are highly consistent with a fibroadenoma, as long as the lesion does not enlarge over time, and lesions that have been present for years and remain unchanged. Microcalcifications are biopsied based on whether they are considered suspicious by imaging criteria.

False-positive imaging can occur not only with breast cancer screening but also in the workup of other malignancies. Patients who undergo fluorodeoxyglucose (FDG) PET or PET CT for staging of cancers other than breast are occasionally found to have (18)F-FDG-avid breast lesions. When biopsied, these lesions are most often benign. Among the reasons for this increased uptake include acute and chronic inflammation, physiological lactation, and benign breast masses, including silicone granuloma, fat

necrosis, fibroadenoma, and postsurgical changes [11]. To decrease the number of false-positive biopsies, Adejolu et al. recommend correlative imaging, including mammography, sonography, or MRI.

---

## Changes During Pregnancy or Lactation

Normal pregnancy-related breast changes include growth and enlargement, tenderness and hyper-sensitivity, darkening of the skin of the nipple and areola, and enlargement of superficial veins near the skin surface. The nipple and areola enlarge. The breasts are more prone to leaking during pregnancy.

Most pregnancy-induced conditions of the breast that are not considered normal are nonetheless benign. These benign conditions include lactating adenoma, galactocele, gigantomastia, and benign nipple discharge [12]. Cancer must be excluded by a thorough workup, including breast biopsy if indicated. During lactation, the most common problems are inflammation and infection. Organisms from the infant are the usual source of breast infections during lactation. Continuing breastfeeding with an infection in the breast is recommended because it is not known to harm the infant, and keeping the breast empty of milk promotes infection resolution by draining the material that is facilitating bacterial growth.

Pregnancy-associated masses are usually discovered during patient self-examination. Ultrasound is the imaging modality of choice to further delineate the lesion and is often useful if a biopsy is indicated. Lactation should be suppressed prior to biopsy in nursing women to reduce the risk of abscess and milk fistula formation [13]. Fine-needle aspiration is less reliable during pregnancy and lactation due to the hyper-proliferative features in the tissue of the pregnant, lactating, or involuting breast [14].

During pregnancy and lactation, the breast can be affected by a variety of benign disorders, including inflammatory and infectious diseases, juvenile papillomatosis, and benign tumors. Fibroadenomas may manifest with growth or

infarction. Galactocele is the most commonly observed breast lesion during lactation [15]. It manifests as either a cystic mass with a fat-fluid level or as a pseudohamartoma. The tumors and diseases that affect the breasts during pregnancy and lactation are also observed in nonpregnant women but may have a different appearance. The sensitivity of mammography in pregnant and lactating women is decreased due to increased parenchymal density. Instead, ultrasonography is the most appropriate radiological method for evaluating breast masses in this setting and is particularly useful in the diagnosis and treatment of abscesses.

Three percent of breast carcinomas occur in women aged 35 or younger. Breast cancer is the leading cause of cancer-related death for women 15–29 years of age [16]. Studies of how reproductive factors influence the development of breast cancer are increasing our understanding of why carcinoma presents as an advanced stage among women who are pregnant or lactating. Specifically, concurrent or recent pregnancy is associated with increased aggressiveness of the tumor and poorer survival. More than 15 % of women younger than age 40 who develop breast cancer do so during pregnancy or lactation [17]. Pregnancy-associated breast cancer (PABC) is classically defined as breast cancer diagnosed during pregnancy or within the first 12 months postpartum. The average age of women with PABC is 32–38 years of age [18]. The incidence of PABC may increase as more women chose to postpone childbearing until their mid-to-late 30s. Pregnancy-related Burkitt's lymphoma characteristically manifests with bilateral and diffuse involvement of the breasts [15].

## Pathology of Benign Breast Disease

Fibroadenomas and disorders related to breast growth are the most common breast diseases in adolescent women. The assessment of breast disorders in adolescents generally involves CBE and, when needed, ultrasonography. Fibroadenomas can be treated conservatively unless they continue to grow. When the diagnosis

is secure and surgical removal is selected, enucleation is the procedure of choice. Breast abscess is mainly due to duct ectasia [19]. Phyllodes tumors (PTs) of the breast are biphasic neoplasms in which interactions between the epithelium and stroma are critical for tumor development and progression. Intratumoral genetic heterogeneity is common in PTs and may account for the reported lack of correlation between histological grading and clinical behavior [20].

Desmoids are benign, slow-growing fibroblastic neoplasms that are characterized by an infiltrative and locally aggressive growth pattern and frequent recurrence but no metastatic potential. Breast desmoids are rare and often misdiagnosed because they can mimic other breast lesions, including carcinoma. Desmoid tumors should be considered in the differential diagnosis of patients presenting with hard breast lumps [21].

Diabetic mastopathy is a proliferation of fibrous tissue in the breast that mimics a tumor. No imaging modality is entirely reliable in differentiating diabetic mastopathy from malignancy, and core biopsy is essential for accurate diagnosis when mammography and/or ultrasonography are indicative of potential malignancy [22]. Breast calcium deposits in the media of arterioles are more frequently detected in the mammograms of diabetic subjects and must be differentiated from suspicious breast microcalcifications.

Pseudoangiomatous stromal hyperplasia (PASH) is a benign, proliferative mesenchymal lesion of the breast that typically affects women of reproductive age [23]. PASH is frequently an incidental histological finding in breast biopsies. Rarely, it can present as a firm, painless breast mass. When presenting as a mass, it is well circumscribed, firm, and rubbery. Histologically, it demonstrates dense collagenous stroma. The most important differential diagnosis is angiosarcoma. When incidentally found, no treatment is required. When PASH forms a tumor mass, it is treated by local surgical excision with clear margins.

Papillary lesions of the breast are common and morphologically varied, ranging from benign to atypical to malignant. Cytological assessment is very challenging and often inconsistent with the histological assessment of the same lesion [24]. Completely excised papillary lesions have an excellent prognosis, whereas incompletely excised lesions may recur or persist as carcinoma. Complete excision is therefore recommended for all papillary lesions [24].

Ectopic breast tissue in axillary lymph nodes (ALNs) is a benign condition that must be differentiated from primary or metastatic carcinoma. Rarely, proliferative conditions such as an intraductal papilloma can occur in ectopic breast tissue [25].

The growing use of breast image-detected biopsies has led to increased diagnosis of benign breast disease (BBD). As a group, BBD is a known risk factor for breast cancer among both Caucasian and African American women [26]. When separated into individual pathological entities, BBD ranges from diagnoses with no increased cancer risk to those with a consistently documented increased cancer risk. The lesions of highest risk, which are sometimes referred to as “borderline” lesions, contain atypical changes and include atypical hyperplasia and lobular carcinoma in situ. Some also classify ductal carcinoma in situ as a borderline lesion, although others do not. These “borderline” lesions can be difficult to diagnose, particularly because biopsy sample size is often limited [27]. Borderline lesions are associated with an increased risk of neighboring malignancy, particularly when the biopsy sample is small. Some of the most challenging scenarios include the differentiation between atypical ductal hyperplasia and low-grade ductal carcinoma in situ, lobular neoplasia versus solid low-grade ductal carcinoma in situ, correctly classifying papillary lesions with atypia, and classifying the spectrum of columnar cell changes [27]. Consensus criteria and uniform terminology for the diagnosis of these lesions do not exist.

## Management of Palpable Breast Masses

In general, palpable breast masses are evaluated by CBE and imaging. Although imaging is not initially required, it does provide a more accurate assessment of mass size and shape and the involvement of the mass with surrounding structures. Most palpable breast masses should undergo biopsy, regardless of whether the lesion appears suspicious on breast imaging, because some cancers appear benign based on imaging criteria. Exceptions include simple cysts and tumors that are consistent with fibroadenoma in a young woman and do not continue to enlarge. Intramammary lymph nodes are generally benign but rarely contain tumor spread from a nearby primary breast cancer. Treatment of simple cysts is generally therapeutic. Bloody and/or recurrent cysts should generally be excised [28]. The increasing use of screening mammography, liability risks, and volume control legislation by the federal government poses a major challenge to clinicians to safely select patients for breast biopsy. Despite a normal mammogram, a palpable breast mass often requires aspiration or excisional biopsy. Careful clinical judgment must prevail if observation is elected. A biopsy should be performed on a clinically suspicious mass regardless of whether the mammogram is suspicious. Management of the patient with a nonpalpable mammographic abnormality requires a close working relationship among the surgeon, pathologist, and radiologist. Thoughtful clinical judgment and interdisciplinary cooperation promote an acceptable benign-to-malignant ratio for breast biopsies.

## References

1. Burnside ES, Sickles EA, Bassett LW, Rubin DL, Lee CH, Ikeda DM, et al. The ACR BI-RADS experience: learning from history. *J Am Coll Radiol JACR*. 2009;6(12):851–60.
2. Goodson WH. Clinical breast examination and breast self examination. In: Sauter ER, Daly MB, editors. *Breast Cancer: Risk Reduction and Early Detection*. New York: Springer; 2010.
3. Sauter ER, Ross E, Daly M, Klein-Szanto A, Engstrom PF, Sorling A, et al. Nipple aspirate fluid: a promising non-invasive method to identify cellular markers of breast cancer risk. *Br J Cancer*. 1997;76(4):494–501.
4. Sauter ER, Schlatter L, Lininger J, Hewett JE. The association of bloody nipple discharge with breast pathology. *Surgery*. 2004;136(4):780–5.
5. Dinkel HP, Gassel AM, Muller T, Lourens S, Rominger M, Tschammler A. Galactography and exfoliative cytology in women with abnormal nipple discharge. *Obstet Gynecol*. 2001;97(4):625–9.
6. Pereira FA, Mudgil AV, Macias ES, Karsif K. Idiopathic granulomatous lobular mastitis. *Int J Dermatol*. 2012;51(2):142–51.
7. Akcan A, Akyildiz H, Deneme MA, Akgun H, Aritas Y. Granulomatous lobular mastitis: a complex diagnostic and therapeutic problem. *World J Surg*. 2006;30(8):1403–9.
8. Boullu S, Andrac L, Piana L, Darmon P, Dutour A, Oliver C. Diabetic mastopathy, complication of type 1 diabetes mellitus: report of two cases and a review of the literature. *Diabetes Metab*. 1998;24(5):448–54.
9. Salzman B, Fleegle S, Tully AS. Common breast problems. *Am Fam Physician*. 2012;86(4):343–9.
10. Vandromme MJ, Umphrey H, Krontiras H. Image-guided methods for biopsy of suspicious breast lesions. *J Surg Oncol*. 2011;103(4):299–305.
11. Adejolu M, Huo L, Rohren E, Santiago L, Yang WT. False-positive lesions mimicking breast cancer on FDG PET and PET/CT. *AJR Am J Roentgenol*. 2012;198(3):W304–14.
12. Scott-Conner CEH. Diagnosing and managing breast disease during pregnancy and lactation. *Medscape Womens Health*. 1997;2(5):1.
13. Woo JC, Yu T, Hurd TC. Breast cancer in pregnancy: a literature review. *Arch Surg*. 2003;138(1):91–8; discussion 9.
14. Novotny DB, Maygarden SJ, Shermer RW, Frable WJ. Fine needle aspiration of benign and malignant breast masses associated with pregnancy. *Acta Cytol*. 1991;35(6):676–86.
15. Sabate JM, Clotet M, Torrubia S, Gomez A, Guerrero R, de las Heras P, et al. Radiologic evaluation of breast disorders related to pregnancy and lactation. *Radiogr Rev Publ Radiol Soc North Am Inc*. 2007;27 Suppl 1:S101–24.
16. Di Nubila B, Cassano E, Urban LA, Fedele P, Abbate F, Maisonneuve P, et al. Radiological features and pathological-biological correlations in 348 women with breast cancer under 35 years old. *Breast*. 2006;15(6):744–53.
17. White TT. Carcinoma of the breast and pregnancy; analysis of 920 cases collected from the literature and 22 new cases. *Ann Surg*. 1954;139(1):9–18.
18. Keyser EA, Staat BC, Fausett MB, Shields AD. Pregnancy-associated breast cancer. *Rev Obstet Gynecol*. 2012;5:94–9.



19. Duflos C, Plu-Bureau G, Thibaud E, Kuttenn F. Breast diseases in adolescents. *Endocr Dev.* 2012;22: 208–21.
20. Karim RZ, O'Toole SA, Scolyer RA, Cooper CL, Chan B, Selinger C, et al. Recent insights into the molecular pathogenesis of mammary phyllodes tumours. *J Clin Pathol.* 2013;66(6):496–505.
21. Taylor TV, Sosa J. Bilateral breast fibromatosis: case report and review of the literature. *J Surg Educ.* 2011;68(4):320–5.
22. Gouveri E, Papanas N, Maltezos E. The female breast and diabetes. *Breast.* 2011;20(3):205–11.
23. Virk RK, Khan A. Pseudoangiomatous stromal hyperplasia: an overview. *Arch Pathol Lab Med.* 2010; 134(7):1070–4.
24. Ueng SH, Mezzetti T, Tavassoli FA. Papillary neoplasms of the breast: a review. *Arch Pathol Lab Med.* 2009;133(6):893–907.
25. Dzodic R, Stanojevic B, Saenko V, Nakashima M, Markovic I, Pupic G, et al. Intraductal papilloma of ectopic breast tissue in axillary lymph node of a patient with a previous intraductal papilloma of ipsilateral breast: a case report and review of the literature. *Diagn Pathol.* 2010;5:17.
26. Cote ML, Ruterbusch JJ, Alesh B, Bandyopadhyay S, Kim E, Albashiti B, et al. Benign breast disease and the risk of subsequent breast cancer in African American women. *Cancer Prev Res.* 2012;5(12): 1375–80.
27. Masood S, Rosa M. Borderline breast lesions: diagnostic challenges and clinical implications. *Adv Anat Pathol.* 2011;18(3):190–8.
28. Winchester DP. Evaluation and management of breast abnormalities. *Cancer.* 1990;66(6 Suppl):1345–7.

Emilia Josefa Borrromeo Diego

---

## Abstract

The topic of benign breast diseases covers a wide variety of conditions, both common and unusual, that may require additional workup, excision, or surveillance. In addition, some benign conditions may confer an increased risk for future disease, and this risk should be explained to the patient during treatment for these entities. This section attempts to provide a basic understanding of some of the most frequently encountered benign breast conditions as well as some rare types, including current recommendations for workup, management, differential diagnoses, and future surveillance. The specific conditions that are explored in this chapter include fibroadenomas, intraductal papillomas, lipomas, hamartomas, radial scars, and gynecomastia in males.

---

## Keywords

Fibroadenoma • Phyllode tumor • Intraductal papilloma • Lipoma • Hamartoma • Radial scar • Granular cell tumor • Gynecomastia

---

## Fibroadenoma

### Definition

Fibroadenomas are benign breast tumors that are composed of epithelial and stromal elements arising from the terminal duct lobular unit. They are frequently diagnosed in young women, predominantly in the second or third decade of life, but can occur at any age [1].

---

E.J.B. Diego, MD  
Department of Surgery, Division of Surgical  
Oncology, Magee-Womens Hospital,  
University of Pittsburgh Medical Center,  
300 Halket Street, Suite 2601,  
Pittsburgh, PA 15213, USA  
e-mail: [diegoe@upmc.edu](mailto:diegoe@upmc.edu)



**Fig. 3.1** Ultrasound appearance of a fibroadenoma



## Clinical Presentation

Most fibroadenomas are likely asymptomatic; they were present in up to 8–10 % of women based upon autopsy findings of “normal breasts” in 1955 [2]. More recent reports indicate that this incidence may be as high as 25 % in asymptomatic women, with 13–20 % having multiple fibroadenomas [3].

If the fibroadenoma causes symptoms, the most common presentation is a firm, movable mass that does not adhere to the chest wall or the skin of the breast. These masses are frequently painless but can occasionally cause discomfort, particularly when they are larger or located in areas that are pressed upon, such as the underwire of a female brassiere.

Medical attention is frequently sought in scenarios of pain, rapid growth, cosmetic deformity, and fear of malignancy.

## Radiological Findings and Workup

The imaging workup for a fibroadenoma usually begins with an ultrasound because these masses are frequently detected in younger women for whom a suspicion of cancer is relatively low. For women who have a personal history or a family history of breast cancer, are over the age of 35, or


have symptoms that are not clinically congruent with a fibroadenoma, a bilateral baseline mammogram should be considered as an adjunctive test.

The characteristic ultrasonographic finding for a fibroadenoma is a round, oval, or lobular well-circumscribed hypoechoic mass [4]. On mammogram, these masses are distinct and occasionally have calcifications. The ultrasound is a more specific test than the mammogram for diagnosing fibroadenomas and should be considered the first imaging modality in young women presenting with a breast mass [5]. See Figs. 3.1 and 3.2.

Core needle biopsy should be obtained when the diagnosis is uncertain due to suspicious features on imaging or there is change in clinical findings (e.g., rapid growth) that may affect surgical planning.

## Pathology

The gross examination of a fibroadenoma reveals a firm, smooth, tan-whitish, lobulated mass that is well-margined and distinct from the surrounding breast tissue. Sectioning of the specimen demonstrates a homogeneous mass that can have a “bulging” appearance. These masses range in size from sub-centimeter to >4 cm. Fibroadenomas that are larger than 5 cm are



**Fig. 3.2** Mammogram of a fibroadenoma

termed giant fibroadenomas and juvenile giant fibroadenomas, specifically, when found in younger women [6]. Historically, the term Brodie's disease of the breast has been used to indicate large, excessively cellular, and long-neglected fibroadenomas [7]. See Fig. 3.3.

Microscopically, fibroadenomas have epithelial and stromal elements with smooth, well-circumscribed borders that can exhibit one of the two growth patterns, pericanalicular or intracanalicular, which pertain to the architecture of the ductal elements. When there is evidence of sclerosing adenosis, metaplasia, or hyperplasia, they are termed complex fibroadenomas [8]. See Figs. 3.4, 3.5, and 3.6.

Special types of rare fibroadenomas are classified based upon histological features [1]. Tubular adenomas or pure adenomas have prominent adenosis and very little stromal elements [2]. See Figs. 3.7 and 3.8. Lactational adenomas exhibit lactational changes in secretory glands in fibroadenomas of pregnant or breastfeeding women [9]. See Figs. 3.9 and 3.10.



**Fig. 3.3** Gross appearance of a fibroadenoma

## Management


Simple fibroadenomas that are asymptomatic and <3 cm, without evidence of growth, in a patient who has no personal or family risk factors for breast cancer can be observed safely. The incidence of malignant transformation is extremely low, and there is no increased lifetime risk of breast cancer [10].

Excision should be considered in patients who have evidence of growth, indeterminate histopathological findings (frequently reported as fibroepithelial lesions), complex fibroadenomas, symptoms such as pain or issues with cosmesis, or patient desire.


Surgical excision remains the most common method for removing fibroadenomas. These palpable lesions can be “shelled out” without excision of the surrounding normal breast tissue, and the incisions are planned such that they do not compromise the appearance of the breast postoperatively. The most frequent approaches to these lesions are through a periareolar or inframammary incision. Should the fibroadenoma be located a distance away from these sites, exposure and delivery of the fibroadenoma can be achieved using an anchoring stitch or Allis clamp to grasp the mass into view. When the fibroadenoma is non-palpable, intraoperative ultrasound guidance or wire localization aids excision.

Ultrasound-guided vacuum-assisted biopsy devices are emerging as a technique to excise

**Fig. 3.4** Low-power magnification view of an intracanalicular fibroadenoma



**Fig. 3.5** Low-power magnification view of a pericanalicular fibroadenoma




smaller fibroadenomas. Advantages to this approach include its ability to be performed as an office procedure with image evidence of complete removal of the mass, a smaller required incision, preservation of breast cosmesis, and limited complications if the targeted lesion is <3 cm [11]. However, long-term data regarding outcomes and recurrence are not available.

### Differential Diagnosis


It is important to distinguish between fibroadenomas and other mass lesions of the breast, particularly a phyllode tumor, due to differences in surgical approach, amount of breast tissue excised, and patient counseling regarding risks of recurrence and need for further treatment. Although the



**Fig. 3.6** Low-power magnification view of a mixed intracanalicular/pericanalicular fibroadenoma



**Fig. 3.7** Low-power magnification view of a tubular adenoma



phyllode tumor is beyond the scope of this section, it is discussed elsewhere in this book.

## Intraductal Papilloma


### Definition and Incidence

Intraductal papillomas are benign breast neoplasms that develop within a mammary duct and are composed of breast epithelium supported by underlying


stroma and a branching fibrovascular core. They are so named due to their microscopic appearance, and they exhibit papillary architecture.

Benign intraductal papillomas are a rare entity with an incidence of 2–3 % in the general population and are frequently diagnosed in women aged 30–55 [12]. By contrast, papillary lesions are observed in up to 5 % of breast core needle biopsies [13]. The presence of atypical features may differentiate benign intraductal papillomas from lesions that confer a higher risk of malignancy.

**Fig. 3.8** High-power magnification view of a tubular adenoma with prominent adenosis and few stromal elements



**Fig. 3.9** Low-power magnification view of a lactational adenoma



## Clinical Presentation

Intraductal papillomas are located in either the central or peripheral portion of the breast. They may be solitary or numerous; in the latter case, the term “papillomatosis” may be employed.


The most frequent symptom associated with a central papilloma is single-duct, spontaneous nipple discharge, which may be serous, greenish, or bloody. Among central or subareolar

papillomas, 50 % will be solitary, and up to 30 % will be associated with bloody nipple discharge [14]. Less frequently, they may present as a palpable mass.

By contrast, peripheral papillomas are very frequently asymptomatic and detected incidentally on breast imaging. Peripheral papillomas, particularly when they are multiple or associated with atypia, are also more likely to be associated with malignancy [15].



**Fig. 3.10** High-power magnification view of a lactational adenoma with lactational changes within the secretory glands



**Fig. 3.11** Ultrasound picture of a centrally located intraductal papilloma. There is an intraductal hypoechoic mass measuring 7 × 2 mm



## Imaging

Central/subareolar papillomas are frequently mammographically occult [16]. However, in the presence of symptoms of a subareolar mass or nipple discharge, an ultrasound and ductogram will commonly reveal the cause.

The ultrasound appearance of an intraductal papilloma is an intraductal mass or complex

cystic lesion and is often associated with a dilated duct. When a ductogram is performed, cannulation of the duct producing the pathological discharge will reveal an abnormality in 91 % of patients with a papilloma. The findings are either a completely obstructed duct, duct expansion and distortion, intraductal filling defects, duct ectasia, or wall irregularity [17]. See Figs. 3.11 and 3.12.

Mammography detects peripheral papillomas more frequently than central papillomas due to the asymptomatic nature of the former. They appear as architectural distortions, nodular densities, breast

masses with or without calcifications, or calcifications alone [18]. See Figs. 3.13 and 3.14.


## Pathology

Intraductal papillomas are typically small lesions, measured in millimeters, that are not visible on gross examination unless associated with an enlarged duct. However, they can grow to several centimeters in size and will appear as a mass growing into the duct.

Microscopically, peripheral and central papillomas are similarly composed of a stalk with a fibrovascular core and overlying myoepithelial and ductal epithelial cells. The ductal epithelium can also exhibit the same proliferative changes observed elsewhere in the breast. Hence, they may also be associated with ductal epithelial hyperplasia, atypical ductal hyperplasia, or ductal carcinoma in situ. See Figs. 3.15, 3.16, 3.17, 3.18, and 3.19.

## Management

Surgical excision is still recommended for all intraductal papillomas found on image-guided biopsy due to their variably reported upstage



**Fig. 3.12** Ductogram demonstrating partial filling of an approximately 1.5-cm span of a duct with numerous intraductal masses




**Fig. 3.13** Ultrasound picture of a peripherally located intraductal papilloma. There is an intraductal hypoechoic mass

rate of 0–29 % [19]. Several series have attempted to define the subset of patients with intraductal papillomas that can safely be observed. Characteristics such as the absence of associated calcifications [19], the microscopic size (complete papilloma excision on core nee-

dle biopsy) [20], the amount of tissue obtained at core biopsy [21], or the absence of atypia [22] have been investigated as predictive factors for a very low risk of upstaging. However, the small sample size in these studies precludes recommendations on a large scale. Table 3.1 categorizes contemporary studies with their reported upstage rates.

### Differential Diagnosis

Benign intraductal papillomas must be distinguished from intraductal papillomas with atypia, papillary ductal carcinomas, and encapsulated (intracystic) papillary carcinomas due to their differences in clinical behavior and management.




**Fig. 3.14** Peripheral papilloma on mediolateral oblique mammogram, seen as an area of asymmetry in the left breast upper quadrant

### Lipoma

#### Definition and Incidence


A lipoma is a benign mass consisting of bland-appearing adipose tissue and is the most common soft tissue tumor in the body, with a prevalence of 2.1 per 1000 people [26]. Lipomas have been




**Fig. 3.15** Low-power magnification view of an intraductal papilloma with ductal epithelial hyperplasia



**Fig. 3.16** High-power magnification view of an intraductal papilloma with ductal epithelial hyperplasia



**Fig. 3.17** Low-power magnification view of an intraductal papilloma with atypia




reported in virtually every area of the body, including the breast. Due to their relatively dull clinical course, much of the literature on breast lipomas are case reports on giant lipomas that pose an interesting scenario for workup and management. Therefore, the true incidence of breast lipomas in general may not be known, although the series of Lanng in 2004 reported an incidence of 4.6 % [27].


### Clinical Presentation

Breast lipomas are typically small, soft, and doughy or semi-firm, painless, mobile masses that are frequently well circumscribed. The term “giant” lipoma was coined by Sanchez et al. [28] and is used to describe a mass that has grown to at least 10 cm in size. Breast lipomas have been observed in both males and females.

**Fig. 3.18** High-power magnification view of an intraductal papilloma with atypia



**Fig. 3.19** A 400× view of an intraductal papilloma with atypia



Consultation is frequently sought due to symptoms such as pain or growth, concern for malignancy, or issues of cosmesis. See Figs. 3.20, 3.21, and 3.22.

As with any breast mass, the diagnostic workup includes breast imaging. Age, clinical

features, and personal and family history should dictate the level of suspicion and attendant workup, which will frequently include a mammogram and breast ultrasound. Further imaging should be performed based upon the results of these tests.



**Table 3.1** Intraductal papilloma studies and upstage rates

	<i>N</i>	Papilloma with atypia	Upstage to atypia, in situ or invasive carcinoma (%)
Jaffer et al. (2009) [23]	104	0	16.4
Brennan et al. (2012) [24]	75	25 (33 %)	6
Holley et al. (2012) [21]	128	0	18
Rizzo et al. (2012) [18]	276	49 (15 %)	28.5
Kibil et al. (2012) [12]	62	12 (19.4 %)	3
Fu et al. (2012) [25]	268	65 (24 %)	23

## Radiological Findings

A lipoma on mammography may have the appearance of a mass with a density similar to that of the surrounding breast fat, possibly with a very thin surrounding capsule [29]. Occasionally, benign-appearing calcifications may also be observed within the mass and may represent fat necrosis [30]. See Figs. 3.22 and 3.23.


Ultrasound findings of a lipoma reveal an isoechoic or slightly hyperechoic mass with a thin surrounding echogenic capsule. There is also typically no posterior acoustic enhancement or shadowing [31].

## Pathology

On gross examination of a breast lipoma, an encapsulated, smooth, fatty mass that may have lobulations is usually observed. Histologically,

**Fig. 3.20** Clinical picture of a man with gynecomastia and a right breast lipoma**Fig. 3.21** On viewing from the side, there is also evidence of a lipoma in this patient's back**Fig. 3.22** Mediolateral oblique mammogram views of a large lipoma demonstrating density very similar to the surrounding breast fat, a very thin surrounding capsule, and benign-appearing calcifications within the mass that represent fat necrosis

the specimen is composed of bland-appearing mature adipocytes. See Figs. 3.24 and 3.25.



**Fig. 3.23** Craniocaudal mammogram views of a large lipoma demonstrating density very similar to the surrounding breast fat, a very thin surrounding capsule, and benign-appearing calcifications within the mass representing fat necrosis

## Management


Surgical excision for lipomas is curative although rarely necessary except for symptom control due to large size, diagnostic uncertainty, radiological-pathological discordance, or patient desire.

The use of liposuction for smaller lipomas has been described and can achieve removal with reduced risk of complications due to smaller incisions [32]. However, the practice has not been extended to the treatment of larger lipomas due to the higher likelihood of incomplete removal, subsequent recurrence, and increased incidence of complications such as hematomas [26]. This technique should also not be performed in situations of diagnostic uncertainty due to the difficulty that will arise in the pathological evaluation of the specimen.

## Prognosis and Differential Diagnosis


As with lipomas in other locations, breast lipomas are benign and do not confer an increased risk of breast cancer when found in the breast. There is no current medical evidence to support the malignant degeneration of lipomas if left in situ.

Other considerations for soft tissue masses in the breast that may present similarly to lipomas



**Fig. 3.24** Low-power magnification view of a lipoma composed primarily of adipocytes with a capsule

**Fig. 3.25** High-power magnification view of a lipoma composed primarily of adipocytes with a capsule



include hibernomas, hematomas, hamartomas (discussed shortly), fat necrosis, and malignant processes such as sarcomas or other tumors of mesenchymal origin.

---

## Hamartoma

### Definition/Incidence

The earliest documented report on hamartomas in the breast was in the German literature by Prym in 1928, in which he described a benign mass called a “mastoma” [33]. Arrigoni et al. first used the term mammary hamartoma to refer to a well-circumscribed mass of benign breast tissue admixed with stromal tissue and fat that is without structural organization [34]. Hamartomas are considered rare breast tumors that encompass fibroadenolipomas, adenolipomas, chondrolipomas, and myoid hamartomas [35]. Literature referring to breast hamartomas is sparse, with 25 patients in the largest series reported to date [34].

As with lipomas, the true incidence of hamartomas is not known, but its diagnosis is increasing due to improvements in breast imaging. Breast hamartomas have also been reported in males [36, 37].

### Clinical Presentation


A breast hamartoma typically presents as a well-circumscribed, mobile, soft, and non-tender breast mass. The palpable nature of the mass is what prompts consultation with a physician. However, they are also occasionally asymptomatic and detected on routine imaging.

Hamartomas have been diagnosed in women of varied ages, without any predilection for pre- or postmenopausal groups. The age range of hamartomas among various series is as young as 11 and as old as 76 [34, 38].


### Imaging

On imaging, a hamartoma may be difficult to differentiate from other benign, solid tumors of the breast such as fibroadenomas.

Mammographic findings may include architectural distortions, asymmetric masses with mixed densities and pseudocapsules, or well-circumscribed nodules. Because hamartomas are masses of disorganized breast tissue with various stromal elements, they will typically have a density similar to that of the surrounding tissue and sometimes have been referred to as “breast-in-breast” lesions [39].



**Fig. 3.26** Mammogram of a palpable 3.6-cm macrolobulated mass composed primarily of dense fat but with scattered areas of soft tissue density within it. This mass was identified as a hamartoma in a female, with the triangle marker overlying the palpable mass



**Fig. 3.27** Exaggerated and magnified craniocaudal mammographic view of a palpable hamartoma in a female, with a triangle marker overlying the palpable mass

However, ultrasonographic findings may more specifically demonstrate a hypoechoic, homogenous mass with distinct borders and no posterior acoustic shadowing [40]. Similarly, they may also demonstrate heterogeneous internal echo patterns due to the differential amounts of breast and stromal tissue present in the tumor. Therefore, it is difficult to make a specific diagnosis of a hamartoma solely based on mammographic or ultrasound imaging. See Figs. 3.26, 3.27, and 3.28.

With the increasing use of MRI in breast imaging, hamartomas are more frequently being detected and characterized. Hamartomas exhibit a gradual enhancement pattern on time-signal intensity curves that differentiates them from malignant processes, which have more rapid enhancement patterns, particularly on dynamic contrast-enhanced MRIs [39].

## Pathology


The gross appearance of a hamartoma is a smooth, lobulated mass with variable amounts of fat and fibrous tissue on sectioning. See Fig. 3.29.

Hamartomas can be further classified by their cellular composition. Adenolipomas will have disorganized benign glandular, adipose, and stromal elements that form a mass with a pseudocapsule or compressed tissue at the borders [41]. Chondrolipomas will contain benign hyaline cartilage admixed with breast lobules and adipose tissue. Myoid hamartomas have an additional smooth muscle component [42, 43]. See Figs. 3.30, 3.31, and 3.32.

On occasion, other pathological changes are observed within hamartomas, such as apocrine



**Fig. 3.28** Ultrasound of a hamartoma demonstrating a 3-cm circumscribed lesion that was isoechoic to normal fatty and fibroglandular tissue



**Fig. 3.29** Gross appearance of a hamartoma

metaplasia, usual or papillary hyperplasia of ductal epithelium, pseudoangiomatous stromal hyperplasia, cysts, and adenosis [34, 44]. Rare cases of associated ductal carcinoma have also been reported [45, 46].

## Management

Fine needle aspiration has little role in the precise diagnosis of hamartomas due to the architectural features that are necessary to differentiate these lesions. However, a bland-appearing aspirate can suggest the benign nature of the hamartoma [47, 48].

Complete surgical excision is curative. There is no evidence for the need to obtain margins beyond the hamartoma, although rare cases of recurrence have been reported, presumably due to incomplete resection [34]. Pure breast hamartomas do not increase a patient's lifetime risk of breast cancer.


## Granular Cell Tumor

### Definition/Incidence


Granular cell tumors are rare, benign neoplasms of neural origin. Specifically, they are thought to be derived from Schwann cells. The lesion is still occasionally called an Abrikossoff tumor after the first description of a tongue mass in 1926 by Russian pathologist Aleksei Ivanovich Abrikosov [49]. They can arise in any organ of the body but most commonly occur in the skin, oral cavity, or digestive tract [50]. Approximately 5–8 % of all granular cell tumors are found in the breast [51]. Granular cell tumors represent an estimated 1 of every 1000 breast neoplasms [52].

Granular cell tumors have been observed in both male and female breasts, and reports indicate that African Americans may be more prone to develop granular cell tumors [51].

**Fig. 3.30** Low-power magnification view of an adenolipoma with disorganized benign glandular, adipose, and stromal elements that form a mass with a pseudocapsule or compressed tissue at the borders



**Fig. 3.31** High-power magnification view of an adenolipoma with disorganized benign glandular, adipose, and stromal elements that form a mass with a pseudocapsule or compressed tissue at the borders



### Clinical Presentation


The clinical presentation of granular cell tumors can mimic a malignancy. Reports of granular cell tumors indicate that they can occur in any quadrant of the breast as a hard, non-tender, mobile mass; however, they may have a specific predilection for the medial breast quadrants due to their perineural origin along the path of the supraclavicular nerve [53, 54]. They

infrequently cause overlying skin changes, fixation to the skin or chest wall, nipple retraction, or breast edema [55].

The most frequent age at presentation in women is the 40s or 50s, but the reported age range in Papalas' series is 19–70 years [56]. The youngest reported case in the literature is a 14-year-old girl [54]. In men, these tumors also tend to occur within the 40–50-year age group [51].



**Fig. 3.32** High-power magnification view of an adenolipoma with disorganized benign glandular, adipose, and stromal elements that form a mass with a pseudocapsule or compressed tissue at the borders




## Imaging

The imaging appearance of a granular cell tumor may not always be helpful in indicating its benign nature or positively identifying it. The mammographic appearance may include smooth, rounded, or lobulated opacities suggestive of a benign process or an indistinct spiculated mass more suspicious of malignancy [53, 55]. Associated microcalcifications or lymphadenopathy has not been noted with pure granular cell tumors. See Figs. 3.33 and 3.34.

Ultrasonographically, granular cell tumors can appear as homogeneous or heterogeneously hypoechoic masses with indistinct borders and posterior acoustic shadowing or as anechoic lesions [50–57]. See Fig. 3.35.

Despite the upsurge in MRI use for breast imaging in recent years, there is very little in the literature regarding the appearance of a granular cell tumor with this modality. In addition, the reported MRI findings of granular cell tumors are also variable. Irregular masses with low-to-high signal intensity on T1-weighted imaging, absent-to-high signal intensity on T2-weighted sequencing, and homogeneous or heterogeneous enhancement on



**Fig. 3.33** Craniocaudal mammographic view of a palpable granular cell tumor in a female, with a triangle marker overlying the palpable mass, which is at the inframammary crease



**Fig. 3.34** Mediolateral oblique mammographic view of a palpable granular cell tumor in a female, with a triangle marker overlying the palpable mass, which is at the inframammary crease

contrast-enhanced T1-weighted imaging have been observed, rendering the MRI a nonspecific tool in identifying granular cell tumors [51, 52, 57].

## Pathology

Gross examination of granular cell tumors reveals smooth-surfaced occasionally lobulated firm masses that are gray-white or tan in color. Lesions are generally less than 3 cm but grow up to 6 cm in size [58]. See Fig. 3.36.


Regardless of the site of excision, granular cell tumors appear pathologically similar on gross or histological examination. They are non-encapsulated tumors composed of polygonal cells that may be arranged in groups, sheets, or nests with a granular, eosinophilic cytoplasm and bland nuclei [59, 60]. See Figs. 3.37 and 3.38.

The histological characteristics of granular cell tumors explain the theories behind the perineural origin of the tumor due to its microscopic similarities to Schwann cells, specifically its positive cytoplasmic and nuclear staining for S-100 protein [58–60].

See Fig. 3.39.




**Fig. 3.35** Ultrasound picture of a granular cell tumor with a 2-cm hypoechoic, irregular mass involving the dermis



**Fig. 3.36** Gross appearance of a granular cell tumor, sectioned (Photo courtesy of Dr. Kandace McGuire)

**Fig. 3.37** Low-power magnification view of a granular cell tumor composed of polygonal cells that have eosinophilic cytoplasm and bland nuclei




Malignant granular cell tumors are rare, occurring in 1–2 % of all cases [49, 50, 60]. The histological criteria for malignant granular cell tumors include tumors >5 cm, areas of necrosis within the tumor, high mitotic activity, and nuclear pleomorphism [50, 60].

### Management


A needle biopsy is helpful for establishing the diagnosis of a granular cell tumor; however, wide local excision is still the recommended treatment to exclude coexisting malignant pathology. The



**Fig. 3.38** High-power magnification view of a granular cell tumor composed of polygonal cells that have eosinophilic cytoplasm and bland nuclei



**Fig. 3.39** High-power magnification view of a granular cell tumor composed of polygonal cells with positive cytoplasmic and nuclear staining for S-100 protein



lifetime recurrence of granular cell tumors after excision, even in the setting of positive and close margins, is extremely low [56].

There is no role for radiation or chemotherapy in the treatment of granular cell tumors. There is also no evidence of an increased lifetime risk of breast cancer in patients who have been diagnosed with a granular cell tumor.

---

## Radial Scar

### Definition/Incidence

A radial scar is a benign breast lesion of unknown origin [61, 62]. The term was first proposed by Dr. H. Hamperl, a pathologist at the University of

Bonn, in 1975 [63]. He described the lesion as consisting of a “hyalinized sclerotic center containing abundant elastic and elastoid masses. These radiate into the periphery and enclose lobuli which reveal epithelial proliferation varying from simple hyperplasia with epithelial villi to the rather rare true papillomas [63].”

The term radial scar conventionally denotes pathological size and is used for lesions measuring up to 9 mm, whereas larger lesions are called complex sclerosing lesions [64]. However, other terms have been used to identify this pathological finding, including rosette-like lesions, proliferative centers, borderline breast tumor, sclerosing papillary proliferations, complex compound heteromorphic lesions, sclerosing lesions, benign sclerosing ductal proliferations, non-encapsulated sclerosing lesions, infiltrating epitheliosis, indurative mastopathy, and proliferating center of Aschoff [65]. The current literature most commonly utilizes radial scar and complex sclerosing lesions.

The true prevalence of radial scars is largely unknown due to its asymptomatic nature. Postmortem studies of women of various ages (17–93 years old) estimate the incidence of radial scars as 7–28 % [66, 67]. In these reports, multiple radial scars were noted in over half of the women examined at autopsy, with bilateral radial scars in some cases. The estimated detection rate of radial scars as a result of mammogram screening is 0.03–0.09 % [68, 69].

## Clinical Presentation

Radial scars are now frequently detected in asymptomatic patients as a result of screening mammography but can also present as a painless, firm breast mass once it reaches a considerable size. The concern of carcinoma is always foremost when this clinical presentation is encountered, prompting a comprehensive breast workup and ultimately leading to surgical excision.

Radial scars can occur anywhere in the breast and have no particular predilection for any quadrant. In addition, radial scars or their larger

counterpart, complex sclerosing lesions, has not been reported to fixate to the chest wall or involve the overlying skin.


## Imaging

The mammographic appearance of radial scars has been well described in the literature according to criteria set forth by Tabar and Dean [70]. The observation of an architectural distortion with a central lucency, radiating, long thin spicules that vary in appearance on different projections in the absence of a palpable clinical finding suggests a radial scar. The absence of microcalcifications has also been noted as a feature of radial scars.

The similar appearance of a radial scar and carcinoma on mammography has led investigators to seek features that may aid the differentiation of these two entities on imaging. In addition, not all radial scars fit the classic criteria, with some lacking the features of a central lucency, others presenting solely with calcifications, and even others displaying similar features of conspicuity variation on different projections, precluding the ability to distinguish carcinoma from a radial scar [68, 71, 72]. These characteristics often lead to a mammographic classification of a suspicious lesion, which prompts further imaging and eventual biopsy. See Fig. 3.40.

Ultrasound provides little additional information for distinguishing radial scars from malignant lesions but can be utilized to more easily localize the lesion for surgical excision or facilitate percutaneous biopsy [61, 65, 68, 73]. The most frequent findings on ultrasound are hypoechoic masses or parenchymal distortion, but up to 1/3 of radial scars may not be visible by this modality [61, 73]. See Fig. 3.41.

MRI evaluation holds promise as an adjunctive test for distinguishing radial scars from malignancy, with the understanding that not all mammographically detected radial scars are visualized on MRI [61, 74]. However, pure radial scars that are appreciated on MRI may exhibit characteristics suggesting their benign nature. Several studies have been published to establish



**Fig. 3.40** Mediolateral oblique mammogram of a palpable radial scar. There is a spiculated mass with associated suspicious calcifications and a triangle marker denoting the area of the palpable mass


the high negative predictive value of the MRI for high-risk lesions, including radial scars specifically [75–77].

## Pathology

The gross pathology of a radial scar reveals a firm lesion with a pale core, irregular edges, and yellowish radiating streaks that appear to be infiltrating the surrounding normal breast tissue [65, 68]. These findings are also consistent with carcinomas; hence, gross examination alone is often not helpful in arriving at the diagnosis.

One clue that may point to a radial scar on gross examination is the finding of surrounding microcysts, which may be present in radial scars but are not typically seen in invasive disease [78].


Histologically, radial scars exhibit a fibroelastotic core with entrapped ducts and radiating ducts and lobules at varying levels of proliferation [65]. These will resemble the spokes in a wheel and are best appreciated on low-power magnification. Calcifications can also be appreciated within radial scars. In addition, radial scars can be associated with atypical lesions, lobular neoplasia, and in situ or invasive carcinomas. See Figs. 3.42 and 3.43.




**Fig. 3.41** Ultrasound findings of a radial scar denoting an irregularly shaped, irregularly marginated hypoechoic lesion, mimicking the appearance of a malignancy



**Fig. 3.42** Low-power magnification view of a radial scar with a fibroelastotic core and entrapped radiating ducts and lobules with variable levels of proliferation



**Fig. 3.43** High-power magnification view of a radial scar with a fibroelastotic core and entrapped radiating ducts and lobules with variable levels of proliferation




It is often challenging to differentiate radial scars from invasive carcinomas, particularly tubular types, due to their similar appearance. The addition of immunohistochemical staining for myoepithelial markers such as smooth muscle actin, calponin, smooth myosin heavy chain, or p63 can help distinguish these two entities. See Figs. 3.44 and 3.45.


### Management

Surgical excision is still justified because of the variable upstage rate of radial scars diagnosed on percutaneous biopsy and because no imaging modality has been proven to reliably guarantee the benign nature of such lesions. A current review of the literature indicates an upstage rate of 0–32 % (Table 3.2).

**Fig. 3.44** High-power magnification view of a radial scar with positive smooth muscle myosin heavy chain staining



**Fig. 3.45** High-power magnification view of a radial scar with positive p63 staining



There may be a trend toward observation and follow-up, particularly in completely excised microscopic radial scars, radial scars without atypia, or radial scars that on MRI are non-enhancing based upon more recent evidence regarding the low likelihood of malignancy in these settings [75, 77, 82, 84]. Our current institutional practice, however, is to surgically excise all radial scars found at percutaneous biopsy.

### Differential Diagnosis and Breast Cancer Risk

The differential diagnosis for radial scars diagnosed on imaging varies widely from carcinoma to fat necrosis and even postoperative changes. A thorough history and physical examination should help eliminate or rule out these other causes.



**Table 3.2** Studies of radial scars and rates of upstage to in situ or invasive carcinoma

	N=radial scar w/ and w/o atypia	Upstage to in situ or invasive carcinoma (%)
Brenner et al. (2002) [79]	157	8
Cawson et al. (2003) [80]	75	7
Lopez-Medina et al. (2006) [81]	43	19
Manfrin et al. (2008) [82]	117	32
Linda et al. (2009) [72]	62	8
Resetkova et al. (2011) [83]	19	0
Rahka et al. (2011) [84]	329	13
Lee et al. (2011) [77] (microscopic radial scars)	18	0
Bianchi et al. (2012) [85]	49	8
Toth et al. (2012) [69]	45	25
Linda et al. (2012) [75]	35	6
Morgan et al. (2012) [86]	67	9
Andacoglu et al. (2013) [87]	67	6

With regard to its classification as a high-risk lesion, the evidence in the literature regarding future breast cancer risk in patients diagnosed with radial scars remains contradictory. Patterson in 2004, Berg in 2008, and Bunting in 2011 all report no increased incidence in breast cancer in women diagnosed with radial scars over follow-up periods of 5–17 years compared with the normal risk population [62, 88, 89]. However, in 2008, Manfrin reported an increased risk of breast cancer in women diagnosed with radial scars as a function of age [82]. However, this study did not follow these patients on a longitudinal basis but instead assessed risk at the time of radial scar diagnosis and concluded that older age conferred a higher risk of breast cancer. More recently, an update from the Nurses' Health Study also concluded that radial scars appear to be an independent risk factor for breast cancer. They identified 460 cases of radial scar with a mean follow-up period of 9 years. The risks for breast cancer in women >50 years and the development of hormone-receptor-negative breast cancers among all women were higher [90].

## Gynecomastia

### Definition/Prevalence

Gynecomastia is defined as benign male breast enlargement that can be unilateral or bilateral, painless, or tender and is a result of glandular proliferation and local fat deposition. These characteristics differ from those of pseudogynecomastia, which is commonly observed in obese males and is characterized by excess fat deposition without glandular proliferation [91]. In addition to physical symptoms, gynecomastia can have a negative psychological impact on affected individuals, prompting patients to seek treatment [92, 93].

Gynecomastia is a common occurrence with three distinct peaks of incidence: the neonatal period, adolescence, and old age [91, 94, 95]. It is observed in approximately 75 % of neonates [96]. Braunstein's 1993 review of gynecomastia in the adolescent age group revealed an incidence of 4–69 % [94]. In older age, gynecomastia has been reported in up to 55 % of autopsies, 57 % of healthy older men, and 70 % of hospitalized elder men [94–96].

Considering the high prevalence and varied causes of gynecomastia, understanding the etiology of the patient's gynecomastia is the clinician's best approach for successful management.

### Etiology/Pathophysiology

The root cause of gynecomastia is an imbalance of estrogen and androgen levels in male breast tissue [91–97]. The male breast has both estrogen and androgen receptors. Estrogen promotes glandular proliferation, whereas androgen inhibits it. Disproportionate activity of estrogen relative to androgens in breast tissue leads to gynecomastia. This imbalance may have several origins, including increased circulating levels of estrogen produced by the adrenal glands or testes, increased peripheral aromatization of estrogen precursors, exposure to estrogen-like substances, medications that

cause the release of more estrogen than androgen from sex hormone-binding globulin (SHBG), decreased androgen production by the testes or altered metabolism, a medication-induced shift of androgens from their receptors, or androgen receptor defects [91].

In the neonatal period, gynecomastia is a result of transplacental transfer of maternal estrogens, as evidenced by a small breast bud that is transient and spontaneously resolves over time [96]. No further treatment is required beyond reassurance.

Adolescent gynecomastia is typically noted at puberty, with a peak age of onset at 13 or 14, and results from peripheral aromatization of circulating androgens [98]. There is a 30-fold increase in testosterone concentration versus a threefold increase in estrogen levels during the shift from prepuberty to puberty in boys, and a disproportionate increase in hormone levels during this phase can cause transient pubertal gynecomastia [99]. As with neonatal gynecomastia, this condition frequently resolves in 1–3 years, requiring only reassurance and surveillance as treatment. However, approximately 8 % of adolescent gynecomastia will continue into adulthood [92]. Many of these patients will seek treatment due to psychological distress, including anxiety, embarrassment, and concern for malignancy, that can be associated with persistent gynecomastia [100].

Adult gynecomastia can be due to a variety of causes, and understanding the specific etiology facilitates the identification of the appropriate treatment. Gynecomastia in older age can be idiopathic or physiological and due to medications, chronic diseases such as cirrhosis or renal disease, endocrine tumors, or endocrine dysfunction [91]. The decrease in plasma testosterone in older men in association with increased peripheral conversion of androgens to estrogens may cause physiological or so-called senile gynecomastia [101]. Tables 3.3 and 3.4 illustrate the other causes of gynecomastia, including medications that cause gynecomastia.

**Table 3.3** Causes of gynecomastia [91, 94, 101]

Idiopathic	–
Physiological	Neonatal period, adolescence, and older men
Medication induced	Antiandrogens Antibiotics Antihypertensives Chemotherapeutic agents Psychoactive agents Diuretics Cardiovascular drugs Gastrointestinal drugs Drugs of abuse
Endocrine dysfunction	Hypogonadism Hyperthyroidism Obesity
Endocrine tumors	Testicular tumors Adrenocortical tumors Tumors secreting ectopic $\beta$ -hCG Pituitary tumors
Chronic diseases	Renal disease Cirrhosis HIV

### Clinical Evaluation and Workup

Most cases of gynecomastia for which consultation is sought are either idiopathic (25 %), acute, or persistent gynecomastia in puberty (25 %) or due to medication (10–20 %) [94, 96–98]. Other causes include cirrhosis or malnutrition (8 %), hypogonadism (8 %), and renal disease (1 %) [94]. In approximately 3 % of cases, gynecomastia is attributable to a testicular tumor [94, 96].

As with other conditions, a thorough history and physical examination are imperative in the evaluation of gynecomastia. Specific attention to underlying disease or medications known to cause gynecomastia may help identify the problem, although half of cases are idiopathic or physiological.

A physical exam with the patient disrobed, in the sitting and supine position, and with hands raised and at the patient's side should be performed. In addition, the supraclavicular, infraclavicular, and axillary lymph node basins must be evaluated.

**Table 3.4** Commonly used medications that cause gynecomastia [91, 94, 97, 101]


Antiandrogens	Bicalutamide, flutamide, nilutamide, finasteride, dutasteride
Antibiotics	Ketoconazole, isoniazid (rare) [97]
Gastrointestinal drugs	Histamine <sub>2</sub> blockers, omeprazole
Diuretics	Spirolactone
Cardiac drugs	Amiodarone, amlodipine, captopril, enalapril, nifedipine, digoxin, reserpine, verapamil, diltiazem
Chemotherapeutic agents	Methotrexate, cyclophosphamide, carmustine, etoposide, melphalan, cisplatin, vincristine, actinomycin D, procarbazine
Antipsychotics	Diazepam, tricyclic antidepressants, haloperidol, phenothiazine, olanzapine, ziprasidone
Recreational drugs	Alcohol, amphetamines, heroin, cocaine, marijuana, anabolic androgens (abuse in athletes)
Antiretrovirals	Efavirenz, saquinavir, indinavir, nelfinavir, ritonavir, lopinavir, stavudine
Others	Phenytoin, penicillamine, statins, theophylline

A distinguishing feature of gynecomastia is concentric enlargement of the breast as opposed to other types of breast masses seen in males, which will more often have an eccentric growth pattern. This growth pattern can be determined during palpation of the nipple areolar complex, where a disk of breast tissue can be palpated in the subareolar region [91, 95, 101, 102]. Gynecomastia is frequently bilateral, although up to 50 % of cases are unilateral [95, 103]. This distribution is in contrast to pseudogynecomastia, which is observed in obese men and is characterized by excess fat deposition without concomitant ductal proliferation. In this situation, soft,


bilaterally enlarged breasts are palpated without any distinct breast tissue in the subareolar region [95, 102]. In addition, breast masses that are not centrally located or have associated lymphadenopathy, skin changes, or nipple retraction must raise the suspicion for diseases other than gynecomastia. Nipple discharge is also uncommon in true gynecomastia [91].

Gynecomastia is frequently painless, asymptomatic, and detected incidentally on routine physical exams. However, it can be associated with pain, particularly in the setting of recent-onset gynecomastia [96].

It is imperative to rule out the possibility of carcinoma, particularly in unilateral cases and in patients who have a family history of breast cancer. The mammographic and sonographic characteristics of gynecomastia are dependent on the acuity or chronicity of the condition and described as having a nodular, dendritic, or diffuse glandular pattern [29, 102, 104]. In recent-onset gynecomastia (<1 year), the florid phase of proliferation of ducts and stromal tissue mammographically reveals a “fan-shaped” subareolar density that merges into the surrounding tissue, and sonography reveals a hypoechoic subareolar mass surrounded by fatty tissue [102, 104]. This stage is reversible because fibrosis has not yet occurred. The dendritic pattern is observed in the more chronic phase of gynecomastia in which irreversible fibrosis has occurred. The flame or cone-shaped subareolar density can be seen infiltrating the deeper, surrounding fat and can even permeate the upper outer quadrants of the breast on mammogram [29, 104]. The diffuse glandular pattern is observed in patients who are treated with high-dose estrogen therapy and have a mammographic and sonographic appearance similar to the female dense breast [29]. Any imaging characteristics that are suspicious warrant percutaneous biopsy and radiologic-pathologic correlation, as in women. The sensitivity and specificity for a mammogram in detecting breast malignancies in males are 92 and 90 %, respectively [98, 102]. See Figs. 3.46, 3.47, and 3.48.



**Fig. 3.46** Magnified tangential mammogram of gynecomastia. There is a large asymmetric density without a discrete mass, suspicious calcifications, or architectural distortions



**Fig. 3.47** Magnified mediolateral mammogram of gynecomastia. There is a large asymmetric density without a discrete mass, suspicious calcifications, or architectural distortions

**Fig. 3.48** Ultrasound findings of gynecomastia. There is absence of a focal mass but findings of heterogeneous tissue with ill-defined hypoechoic areas



Adjunctive laboratory testing may be justified when imaging is benign, and there is a high suspicion that the gynecomastia is representative of an underlying pathology. Some of these tests include determining testosterone,  $\beta$ -hCG, luteinizing hormone, thyroid-stimulating hormone, and prolactin levels [91, 94, 95, 98]. Abnormal results should prompt consultation with an endocrinologist. In addition, liver and renal function testing may reveal chronic hepatic or kidney disease as the cause of the gynecomastia.

## Pathology

Gross examination of a male breast with gynecomastia reveals a gray-white, rubbery mass in the subareolar region. As previously mentioned, there are two phases of gynecomastia: the florid (reversible) phase and fibrotic (irreversible) phase [105].

The florid phase is typically observed in the first year of onset. Histologically, there is benign proliferation of ductal epithelium and stromal elements, with periductal inflammation and surrounding edema without evidence of fibrosis [102, 105]. In the absence of fibrosis, nonsurgical treatment may be successful, and gynecomastia may be reversible [95]. See Fig. 3.49.


In the fibrotic phase, which can be observed beyond 6 months, there is minimal ductal proliferation and hyalinized periductal tissue [106]. If treatment is desired in this phase, only surgical options are feasible.

## Management

Most patients with gynecomastia may experience a resolution of symptoms upon removal of the precipitating cause or treatment of the underlying condition.

When medication-induced gynecomastia precludes withdrawal of the drug due to medical necessity, tamoxifen therapy may reverse the condition. Small studies have indicated gynecomastia in up to 80 % of patients and even prevention if used in a prophylactic manner, particularly in the setting of treatment for prostate cancer [107–109]. The US Food and Drug Administration, however, has not approved tamoxifen for this indication upon writing of this text [97]. Therefore, before initiating therapy, the patient must be counseled regarding current evidence, and the risk/benefit ratio must be considered on an individual basis.

Another strategy that deserves comment in the treatment and prophylaxis of gynecomastia



**Fig. 3.49** High-power magnification view of gynecomastia with benign proliferation of ductal epithelium and stromal elements



induced by prostate cancer therapy is low-dose radiation. A few European studies have established the efficacy of radiotherapy in both treatment and prophylaxis, with minimal and transient side effects, making it an acceptable alternative for patients [96, 110]. However, large-scale studies and standardized regimens are lacking.

Surgery is still the mainstay of treatment for long-standing gynecomastia. It is the tenth most common procedure performed by plastic surgeons worldwide [106]. However, the types of procedures performed for gynecomastia are evolving and trending toward less invasive techniques with smaller incisions. The subcutaneous mastectomy with a periareolar or inframammary incision is still standard (with or without resection of excess skin), but the addition of liposuction may be beneficial in patients with a greater degree of ptosis [100, 103, 106].

Special thanks to Dr. Faye Gao of the Department of Pathology at Magee-Womens Hospital for preparation of the pathology pictures.

## References

- Kuijper A, Mommers E, van der Wall E, van Diest P. Histopathology of fibroadenoma of the breast. *Am J Clin Pathol.* 2001;115:736–42.
- Frantz VK, Pickren JW, Melcher GW, Auchincloss Jr H. Incidence of chronic cystic disease in so-called “normal breasts”: a study based on 225 postmortem examinations. *Cancer.* 1951;4(4):762–83.
- El-Wakeel H, Umpleby HC. Systematic review of fibroadenoma as a risk factor for breast cancer. *Breast.* 2003;12(5):302–7.
- Harvey JA, Nicholson BT, LoRusso AP, Cohen MA, Bovbjerg VE. Short-term follow-up of palpable breast lesions with benign imaging features: evaluation of 375 lesions in 320 women. *Am J Roentgenol.* 2009;193(6):1723–30.
- Zimmermann N, Ohlinger R. Diagnostic value of palpation, mammography and ultrasonography in the diagnosis of fibroadenoma: impact of breast density, patient age, ultrasonographic size, and palpability. *Ultraschall Med.* 2011;33(7):E151–7.
- Matz D, Kerivan L, Reintgen M, Akman K, Lozicki A, Causey T, et al. Breast preservation in women with giant juvenile fibroadenoma. *Clin Breast Cancer.* 2013;13(3):219–22.
- Aiken D. Brodie’s disease of the breast. *Br J Surg.* 1946;34:87–90.
- Sklair-Levy M, Sella T, Alweiss T, Cracium L, Libson E, Mally B. Incidence and management of complex fibroadenomas. *Am J Roentgenol.* 2008;190(1):214–8.
- Esposito N. Fibroepithelial lesions. In: Dabbs DJ, editor. *Breast pathology.* Philadelphia: Saunders; 2012. p. 233–51.
- Valdes EK, Boolbol SK, Cohen JM, Feldman S. Malignant transformation of a breast fibroadenoma to cystosarcoma phyllodes: case report and review of the literature. *Am Surg.* 2005;71(4):348–53.
- Luo HJ, Chen X, Tu G, Wang J, Wu CY, Yang GL. Therapeutic application of ultrasound-guided 8-gauge mammotome system in presumed benign breast lesions. *Breast J.* 2011;17(5):490–7.
- Kibil W, Hodorowicz-Zaniewska D, Popiela T, Kulig J. Vacuum-assisted core biopsy in diagnosis and treatment of intraductal papillomas. *Clin Breast Cancer.* 2012. doi:10.1016/j.clbc.2012.09.018.
- Jakate K, De Brot M, Goldberg F, Muradali D, O’Malley FP, Mulligan AM. Papillary lesions of the breast: impact of breast pathology subspecialization on core biopsy and excision diagnosis. *Am J Surg Pathol.* 2012;36(4):544–51.
- Rizzo M, Lund MJ, Oprea G, Schniederjan M, Wood WC, Mosunjac M. Surgical follow-up and clinical presentation of 142 breast papillary lesions diagnosed by ultrasound-guided core-needle biopsy. *Ann Surg Oncol.* 2008;15(4):1040–7.
- Lewis JT, Hartmann LC, Vierkant RA, Maloney SD, Pankratz S, Allers TM, et al. An analysis of breast cancer risk in women with single, multiple and atypical papilloma. *Am J Surg Pathol.* 2006;30(6):665–72.
- Esposito N. Papilloma and papillary lesions. In: Dabbs DJ, editor. *Breast pathology.* Philadelphia: Saunders; 2012. p. 252–62.
- Cardenosa G, Eklund GW. Benign papillary neoplasms of the breast: mammographic findings. *Radiology.* 1991;181:751–5.
- Rizzo M, Linebarger J, Lowe M, Pan L, Gabram S, Vasquez L, et al. Management of papillary breast lesions diagnosed on core-needle biopsy: clinical pathologic and radiologic analysis of 276 cases with surgical follow-up. *J Am Coll Surg.* 2012;214(3):280–7.
- Li X, Weaver O, Desouki MM, Dabbs D, Shyum S, Carter G, et al. Microcalcification is an important factor in the management of breast intraductal papillomas diagnosed on core biopsy. *Am J Clin Pathol.* 2012;138:789–95.
- Lee KA, Zuley ML, Chivukula M, Choski ND, Ganott MA, Sumkin JH. Risk of malignancy when microscopic radial scars and microscopic papillomas are found at percutaneous biopsy. *Am J Roentgenol.* 2012;198:W141–5.
- Holley SO, Appleton CM, Farria DM, Reichert VC, Warrick J, Allread DC, et al. Pathologic outcomes of nonmalignant papillary breast lesions diagnosed at imaging-guided core needle biopsy. *Radiology.* 2012;265(2):379–84.

22. Swapp RE, Glazebrook KN, Jones KN, Brandts HM, Reynolds C, Visscher DW, et al. Management of benign intraductal solitary papilloma diagnosed on core needle biopsy. *Ann Surg Oncol*. 2013; 20(6):1900–5.
23. Jaffer S, Bleiweiss NC. Excision is indicated for intraductal papilloma of the breast diagnosed on core needle biopsy. *Cancer*. 2009;115(13):2837–43.
24. Brennan SB, Corben A, Liberman L, Dershaw DD, Brogi E, Van Zee KJ, et al. Papilloma diagnosed at MRI-guided vacuum-assisted breast biopsy: is surgical excision still warranted? *AJR Am J Roengenol*. 2012;199:W512–9.
25. Fu CY, Chen TW, Hong ZJ, Chan DC, Young CY, Chen CJ, et al. Papillary breast lesions diagnosed by core biopsy require complete excision. *Eur J Surg Oncol*. 2012;38(11):1029–35.
26. Groh O, Klaas in't Hof K. Giant lipoma of the male breast: case report and review of the literature. *Eur J Plast Surg*. 2012;35(5):407–9.
27. Lannig C, Eriksen BO, Hoffmann J. Lipoma of the breast: a diagnostic dilemma. *Breast*. 2004;13(5):408–11.
28. Sanchez MR, Golomb FM, Moy JA, Potozkin JR. Giant lipoma: case report and review of the literature. *J Am Acad Dermatol*. 1993;28:266–8.
29. Lattin GE, Jessinger RA, Mattu R, Glassman LM. From the radiologic pathology archives: diseases of the male breast: radiologic-pathologic correlation. *Radiographics*. 2013;33(2):461–89.
30. Muttarak M, Chaiwaun B. Imaging of giant breast masses with pathological correlation. *Singapore Med J*. 2004;45(3):132–9.
31. Yitta S, Singer CI, Toth HB, Mercado CL. Sonographic appearance of benign and malignant male breast disease with mammographic and pathologic correlation. *J Ultrasound Med*. 2010;29(6):931–47.
32. Grossman J, Menes T, Lahat G, Gur E, Weiss J, Barnea Y. Use of the oncoplastic reduction pattern technique following removal of a giant breast lipoma. *Ann Plast Surg*. 2011;67(2):106–8.
33. Uchoa DGM, Cruz DB, Schaefer PG, Pegas KL, Cambuzzi E. Myofibroblastoma arising in mammary hamartoma: a case report. *Pathol Res Int*. 2010;2010:726829.
34. Tse GM, Law BK, Ma TK, Chan AB, Pang L-M, Chu WC, Cheung SH. Hamartoma of the breast: a clinicopathological review. *J Clin Pathol*. 2002;55(12):951–4.
35. Shin SJ, Rabban JT. Mesenchymal neoplasms of the breast. In: Dabbs DJ, editor. *Breast pathology*. Philadelphia: Saunders; 2012. p. 596–641.
36. Desphande A, Munshi M. Mammary hamartoma: report of two cases including one in a male breast, and review of the literature. *Indian J Pathol Microbiol*. 2004;47(4):511–5.
37. Gupta SS, Singh O, Hastir A, Arora G, Sabharwal G, Mishra H. Breast hamartoma with intrathoracic extension in a 13-year old boy. *J Cancer Res Ther*. 2010;6(1):86–8.
38. Chang HL, Lerwill MF, Goldstein AM. Breast hamartomas in adolescent females. *Breast J*. 2009;15(5):515–20.
39. Erdem G, Karakas HM, Isik B, Firat AK. Advanced MRI findings in patients with breast hamartomas. *Diagn Interv Radiol*. 2011;17(1):33–7.
40. Am D, Ramesar K, Allan S, Marr A, Howlett D. Breast hamartoma arising in axillary ectopic breast tissue. *Breast J*. 2010;16(4):433–4.
41. Birrell AL, Warren LR, Birrell SN. Misdiagnosis of massive breast asymmetry: giant hamartomas. *ANZ J Surg*. 2012;82(12):941–2.
42. Mizuta N, Sakaguchi K, Mizuta M, Imai A, Nakatsukasa K, Morita M, et al. Myoid hamartoma of the breast that proved difficult to diagnose: a case report. *World J Surg Oncol*. 2012;10:12.
43. Kajo K, Zubor P, Danko J. Myoid (muscular) hamartoma of the breast: case report and review of the literature. *Breast Care*. 2010;5(5):331–4.
44. Yu L, Yang W, Xu X. Myoid hamartoma of the breast: clinicopathologic analysis of a rare tumor indicating occasional recurrence potential. *Breast J*. 2011;17(3):322–4.
45. Ruiz-Tovar J, Reguero-Callejas ME, Alaez AB, Ramiro C, Rojo R, Collado MV, et al. Infiltrating ductal carcinoma and ductal carcinoma in situ associated with mammary hamartoma. *Breast J*. 2006;12(4):368–70.
46. Choi N, Ko ES. Invasive ductal carcinoma in a mammary hamartoma: case report and review of the literature. *Korean J Radiol*. 2010;11(6):687–91.
47. Herbert M, Schvimer M, Zehavi S, Mendlovic S, Karni T, Pappo I, et al. Breast hamartoma: fine-needle aspiration cytologic finding. *Cancer*. 2003;99(4):255–8.
48. Herbert M, Mendlovic S, Liokumovich P, Segal M, Zahavi S, Rath-Wolfson L, et al. Can hamartoma of the breast be distinguished from fibroadenoma using fine-needle aspiration cytology? *Diagn Cytopathol*. 2004;34(5):326–9.
49. Haikal F, Maceira JP, Dias EP, Ramos-e-Silva M. Histogenesis of Abrikossoff tumour of the oral cavity. *Int J Dent Hyg*. 2004;8(1):53–62.
50. Aoyama K, Kamio T, Hirano A, Seshimo A, Kameoka S. Granular cell tumors: a report of six cases. *World J Surg Oncol*. 2012. doi:10.1186/1477-7819-10-204.
51. Kim EY, Kang DK, Kim TH, Jung YS, Kim KS, Yim H. Granular cell tumor of the male breast: two case descriptions and brief review of the literature. *J Ultrasound Med*. 2011;30(9):1295–301.
52. Maki DD, Horne D, Damore II LJ, Jones C. Magnetic resonance appearance of granular cell tumors of the breast. *Clin Imaging*. 2009;33(5):395–7.
53. Pergel A, Yucel AF, Karaca AS, Aydin I, Sahin DA, Demirbag N. A therapeutic and diagnostic dilemma: granular cell tumor of the breast. *Case Rep Med*. 2011. doi:10.1155/2011/972168.
54. De Simone N, Aggon A, Christy C. Granular cell tumor of the breast: clinical and pathologic characteristics of a rare case in a 14-year-old girl. *J Clin Oncol*. 2011;29(22):e656–7.

55. Lee SJ, Mahoney MC. The protean manifestations of granular cell tumor of the breast. *Breast J*. 2011;17(2):201–3.
56. Papalas JA, Wylie JD, Dash RC. Recurrence risk and margin status in granular cell tumors of the breast. *Arch Pathol Lab Med*. 2011;135(7):890–5.
57. Scaranelo AM, Bukhanov K, Crystal P, Mulligan AM, O'Malley FP. Granular cell tumour of the breast: MRI findings and review of the literature. *Br J Radiol*. 2007;80(960):970–4.
58. Filipovski V, Banev S, Janevska V, Dukova B. Granular cell tumor of the breast: a case report and review of literature. *Cases J*. 2009. doi:10.4076/1757-1626-2-8551.
59. Rekhi B, Jambhekar NA. Morphologic spectrum, immunohistochemical analysis, and clinical features of a series of granular cell tumors of soft tissues: a study from a tertiary referral cancer center. *Ann Diagn Pathol*. 2010;14(3):162–7.
60. Aneiros-Fernandez J, Arias-Santiago S, Husein-El Ahmed H, O'Valle F, Siendones MIA, Aneiros-Cachaza J. Cutaneous granular cell tumor of the breast: a clinical diagnostic pitfall. *J Clin Med Res*. 2010;2(4):185–8.
61. Linda A, Zuiani C, Londero V, Cedolini C, Girometti R, Bazzocchi M. Magnetic resonance imaging of radial sclerosing lesions (radial scars) of the breast. *Eur J Radiol*. 2012;81(11):3201–7.
62. Bunting DM, Steel JR, Holgate CS, Watkins RM. Long term follow-up and risk of breast cancer after a radial scar or complex sclerosing lesion has been identified in a benign open breast biopsy. *Eur J Surg Oncol*. 2011;37(8):709–13.
63. Hamperl H. Radial scars (scarring) and obliterating mastopathy (author's transl). *Virchows Arch A Pathol Anat Histol*. 1975;369(1):55–68.
64. Page D, Anderson T. Radial scars and complex sclerosing lesions. In: Page D, Anderson T, editors. *Diagnostic histopathology of the breast*. Edinburgh: Churchill Livingstone; 1987. p. 89–103.
65. Babu ED, Power DM, Powis SJA. Radial scar. *Breast*. 2000;9(2):66–71.
66. Bhathal PS, Browm RW, Lesueur GC, Russell IS. Frequency of benign and malignant breast lesions in 207 consecutive autopsies in Australian women. *Br J Cancer*. 1985;51(2):271–8.
67. Nielsen M, Jensen J, Andersen JA. An autopsy study of radial scar in the female breast. *Histopathology*. 1985;9(3):287–95.
68. Kennedy M, Masterson AV, Kerin M, Flanagan F. Pathology and clinical relevance of radial scars: a review. *J Clin Pathol*. 2003;56(10):721–4.
69. Toth D, Sebo E, Sarkadi L, Kovacs I, Kiss C, Damajanovich L. Role of core needle biopsy in the treatment of radial scar. *Breast*. 2012;21(6):761–3.
70. Tabar L, Dean PB. *Teaching atlas of mammography*. 3rd ed. New York: Thieme Verlag Inc; 2001. p. 93–148.
71. Cawson JN, Nickson C, Evans J, Kavanagh AM. Variation in mammographic appearance between projections of small breast cancers compared with radial scars. *J Med Imaging Radiat Oncol*. 2010;54(5):415–20.
72. Linda A, Zuiani C, Furlan A, Londero V, Girometti R, Machin P, et al. Radial scars without atypia diagnosed at imaging-guided needle biopsy: how often is associated malignancy found at subsequent surgical excision, and do mammography and sonography predict which lesions are malignant? *AJR Am J Roentgenol*. 2010;194(4):1146–51.
73. Lee E, Wylie E, Metcalf C. Ultrasound imaging features of radial scars of the breast. *Australas Radiol*. 2007;51(3):240–5.
74. Strigel RM, Eby PR, DeMartini WB, Gutierrez RL, Allison KH, Peacock S, et al. Frequency, upgrade rates, and characteristics of high-risk lesions initially identified with breast MRI. *AJR Am J Roentgenol*. 2010;195(3):792–8.
75. Linda A, Zuiani C, Furlan A, Lorenzon M, Londero V, Girometti R, et al. Nonsurgical management of high-risk lesions diagnosed at core needle biopsy: can malignancy be ruled out safely with breast MRI? *AJR Am J Roentgenol*. 2012;198(2):272–80.
76. Perfetto F, Fiorentino F, Urbano F, Silecchia R. Adjunctive diagnostic value of MRI in the breast radial scar. *Radiol Med*. 2009;114(5):757–70.
77. Lee KA, Zuley ML, Chivukula M, Choksi ND, Ganott MA, Sumkin JH. Risk of malignancy when microscopic radial scars and microscopic papillomas are found at percutaneous biopsy. *AJR Am J Roentgenol*. 2011;198(2):w141–5.
78. Mohsin SK. Radial scar. In: Dabbs DJ, editor. *Breast pathology*. Philadelphia: Saunders; 2012. p. 293–306.
79. Brenner RJ, Jackman RJ, Parker SH. Percutaneous core needle biopsy of radial scars of the breast: when is excision necessary? *AJR Am J Roentgenol*. 2002;179(5):1179–84.
80. Cawson JN, Malara F, Kavanagh A, Hill P, Balasubramaniam G, Henderson M. Fourteen-gauge needle core biopsy of mammographically evident radial scars: is excision necessary? *Cancer*. 2003;97(2):345–51.
81. Lopez-Medina A, Cintora E, Mugica B, Operé E, Vela AC, Ibañez T. Radial scar diagnosed at stereotactic core needle biopsy: surgical biopsy findings. *Eur Radiol*. 2006;16(8):1803–10.
82. Manfrin E, Remo F, Reghellin D, Bonetti F. Risk of neoplastic transformation in asymptomatic radial scar. Analysis of 117 cases. *Breast Cancer Res Treat*. 2008;107(3):371–7.
83. Resetkova E, Edelweiss M, Albarracin CT, Yang WT. Management of radial sclerosing lesions of the breast using percutaneous vacuum-assisted core needle biopsy: recommendations for excision based on seven years of experience at a single institution. *Breast Cancer Res Treat*. 2011;127(2):335–43.
84. Rahka EA, Lee AHS, Jenkins JA, Murphy AE, Hamilton LJ, Ellis IO. Characterisation and outcome of breast needle core biopsy diagnosis of lesions of uncertain malignant potential (B3) in abnormalities detected by mammographic screening. *Int J Cancer*. 2011;129(6):1417–24.



85. Bianchi S, Giannotti E, Vanzi E, Marziali M, Abdulcadir D, Boeri C, et al. Radial scar without associated atypical epithelial proliferation on image guided 14-gauge needle core biopsy: analysis of 49 cases from a single-centre and review of the literature. *Breast*. 2012;21(2):159–64.
86. Morgan C, Shah ZA, Hamilton R, Wang J, Spigel J, Deleon W, et al. The radial scar of the breast diagnosed at core needle biopsy. *Proc (Bayl Univ Med Cent)*. 2012;25(1):3–5.
87. Andacoglu O, Kanbour-Shakir A, Teh YC, Bonaventura M, Ozbek U, Anello M, et al. Rationale of excisional biopsy after the diagnosis of benign radial scar on core biopsy. *Am J Clin Oncol*. 2013;36(1):7–11.
88. Patterson JA, Scott M, Anderson N, Kirk SJ. Radial scar, complex sclerosing lesion and risk of breast cancer. Analysis of 175 cases in northern Ireland. *Eur J Surg Oncol*. 2004;30(10):1065–8.
89. Berg JC, Visscher DW, Vierkant RA, Pankratz VS, Maloney SD, Lewis JT, et al. Breast cancer risk in women with radial scars in benign breast biopsies. *Breast Cancer Res Treat*. 2008;108(2):167–74.
90. Aroner SA, Collins LC, Connolly JL, Colditz GA, Schnitt SJ, Rosner BA, et al. Radial scars and subsequent breast cancer risk: results from the nurses' health studies. *Breast Cancer Res Treat*. 2013;139(1):277–85.
91. Barros AC, Sampaio Mde C. Gynecomastia: pathophysiology, evaluation and treatment. *Sao Paulo Med J*. 2012;130(3):187–97.
92. Nuzzi LA, Cerrato FE, Erickson CR, Webb ML, Rosen H, Walsh EM, et al. Psychosocial impact of adolescent gynecomastia: a prospective case-control study. *Plast Reconstr Surg*. 2013;131(4):890–6.
93. Colombo-Benkmann M, Buse B, Stern J, Herfarth C. Indications for and results of surgical therapy for male gynecomastia. *Am J Surg*. 1999;178(1):60–3.
94. Braunstein GD. Gynecomastia. *N Engl J Med*. 1993;328(1):490–5.
95. Braunstein GD. Gynecomastia. *N Eng J Med*. 2007;357(12):1229–37.
96. Gikas P, Mokbel K. Management of gynaecomastia: an update. *Int J Clin Pract*. 2007;61(7):1209–15.
97. Bowman JD, Kim H, Bustamante JJ. Drug-induced gynecomastia. *Pharmacotherapy*. 2012;32(12):1123–40.
98. Morcos RN, Kizy T. Gynecomastia: when is treatment indicated? *J Fam Pract*. 2012;61(12):719–25.
99. Bidlingmaier F, Knorr D. Plasma testosterone and estrogens in pubertal gynecomastia. *Z Kinderheilkd*. 1973;115(1):89–94.
100. Song YN, Wang YB, Huang R, He XG, Zhang JF, Zhang GQ, et al. Surgical treatment of gynecomastia: mastectomy compared to liposuction technique. *Ann Plast Surg*. 2014;73(3):275–8.
101. Beenken SW, Page DL, Bland KI. Gynecomastia. In: Bland KI, Copeland EM, editors. *The breast: comprehensive management of benign and malignant diseases*. 4th ed. Philadelphia: Saunders; 2009. p. 151–68.
102. Nguyen C, Kettler MD, Swirsky ME, Miller VI, Scott C, Krause R, et al. Male breast disease: pictorial review with radiologic-pathologic correlation. *Radiographics*. 2013;33(3):763–79.
103. Kasielska A, Antoszewski B. Surgical management of gynecomastia: an outcome analysis. *Ann Plast Surg*. 2013;71(5):471–5.
104. Charlot M, Beatrix O, Chateau F, Dubuisson J, Golfier F, Valette PJ, et al. Pathologies of the male breast. *Diagn Interv Imaging*. 2013;94(1):26–37.
105. Chivukula M, Dabbs DJ. Neoplasia of the male breast. In: Dabbs DJ, editor. *Breast pathology*. Philadelphia: Saunders; 2012. p. 642–64.
106. Lee JH, Kim IK, Kim TG, Kim YH. Surgical correction of gynecomastia with minimal scarring. *Aesthetic Plast Surg*. 2012;36(6):1302–6.
107. Serretta V, Altieri V, Morgia G, Serretta V, Morgia G, et al. A randomized trial comparing tamoxifen therapy vs. tamoxifen prophylaxis in bicalutamide-induced gynecomastia. *Clin Genitourin Cancer*. 2012;10(3):174–9.
108. James R, Ahmed F, Cunnick G. The efficacy of tamoxifen in the treatment of primary gynecomastia: an observational study of tamoxifen versus observation alone. *Breast J*. 2012;18(6):620–1.
109. Kunath F, Keck B, Antes G, Wullich B, Meerpohl JJ. Tamoxifen for the management of breast events induced by non-steroidal antiandrogens in patients with prostate cancer: a systematic review. *BMC Med*. 2012;10:96.
110. Neu B, Sautter V, Momm F, Melcher U, Seegenschmied H, Micke O, et al. Radiotherapy for prevention and therapy of gynecomastia due to antiandrogen treatment in prostate cancer patients: a patterns-of-care study. *Strahlenther Onkol*. 2011;187(12):771–7.

Soley Bayraktar and Banu K. Arun

---

## Abstract

Breast cancer is the most frequently diagnosed cancer in women in the United States, with an estimated 296,000 new cases and more than 40,000 deaths in 2013. Age, family history, and both endogenous and exogenous ovarian hormone exposure have important effects on risk and have been incorporated into models that predict individual risk of breast cancer; diet, alcohol use, and other factors play smaller roles. Inherited mutations in BRCA1, BRCA2, and CHEK2 play a role in the development of hereditary breast cancer, and high-risk individuals can be directly tested. BRCA mutation-associated breast cancer differs from sporadic breast cancer in that BRCA mutation carriers have a higher risk of breast and ovarian cancer and differential sensitivity to chemotherapeutic agents. As BRCA genetic testing is readily available, BRCA mutation status should be evaluated in high-risk women, including women who were diagnosed with breast cancer at an early age, who have a strong family history, or whose tumors were triple negative (TN). Given the high rate of contralateral breast cancer and ovarian cancer, mutation carriers with newly diagnosed breast cancer may choose to undergo contralateral prophylactic mastectomy or bilateral salpingo-oophorectomy. In addition, two selective estrogen receptor modulators, tamoxifen and raloxifene, and aromatase inhibitors can be used to decrease the incidence of invasive breast cancer in women who are at high risk of this condition.

---

S. Bayraktar, MD, MBA  
Department of Breast Medical Oncology, The  
University of Texas MD Anderson Cancer Center,  
1515 Holcombe Boulevard, Unit 1354, Houston, TX,  
77030-4009, USA

Department of Medical Oncology, Mercy Cancer  
Center, Ardmore, OK, USA  
e-mail: [soley.bayraktar@gmail.com](mailto:soley.bayraktar@gmail.com)

B.K. Arun, MD (✉)  
Department of Breast Medical Oncology, The  
University of Texas MD Anderson Cancer Center,  
1515 Holcombe Boulevard, Unit 1354, Houston, TX,  
77030-4009, USA  
e-mail: [barun@mdanderson.org](mailto:barun@mdanderson.org)

**Keywords**

Breast cancer • Prevention • Risk factors • BRCA • Hereditary breast cancer  
• Risk models • Prophylactic mastectomy • Prophylactic oophorectomy

**Epidemiology of Breast Cancer**

In the United States (USA), breast cancer is the most frequently diagnosed cancer in women. In 2014, an estimated 232,670 new cases of invasive breast cancer were expected to be diagnosed in women in the United States, along with 62,570 new cases of noninvasive (in situ) disease. Approximately 40,000 women in the United States were expected to die in 2014 from breast cancer, although death rates have been decreasing since 1989, with larger decreases in women under 50 (American Cancer Society, [www.cancer.org](http://www.cancer.org)). Breast cancer accounts for 29 % of all cancers diagnosed and 16 % of all cancer deaths in US women [1].

Globally but particularly in developed countries, breast cancer is a major public health problem, with one million new cases diagnosed annually [2]. The lowest incidence rates of breast cancer are in Asian countries (10–15 cases/100,000 women) [2, 3]. However, incidence rates have increased rapidly in countries such as Japan, where major lifestyle changes have occurred in the last 50 years. When women from Asia or other low-risk areas migrate to an area of high risk, they gradually assume the risk of the high-risk population [4]. Interestingly, the difference in breast cancer incidence between Asian and Western populations is primarily the result of much lower incidence rates of postmenopausal breast cancer in Asian countries because premenopausal breast cancer rates are similar in Asian and Western countries [5]. These data suggest that the underlying genetic factors that primarily contribute to breast cancer in young women are similar in both populations, but hormone exposure and lifestyle factors that vary widely between continents may play an important role in defining postmenopausal breast cancer risk.

In the United States, approximately 1 in 8 US women (approximately 12 %) will develop inva-



sive breast cancer over the course of her lifetime [6]. However, more than half of this risk is incurred after 60 years of age, and the risk of 1 in 8 is not reached until 110 years of age [7] (Fig. 4.1). In addition, risk is very heterogeneous across the population. Therefore, individual risk assessment is considerably more useful than population risk in the development of clinical management strategies.

Breast cancer incidence rates in the United States began decreasing in 2000 after increasing for the previous two decades. They decreased by 7 % from 2002 to 2003 alone. This decrease may be due in part to the reduced use of hormone replacement therapy (HRT) by women after the results of a large study called the Women's Health Initiative were published in 2002. These results suggested a connection between HRT and increased breast cancer risk. White women are slightly more likely to develop breast cancer than African-American women (Fig. 4.2). However, in women under 45, breast cancer is more common in African-American women than white women. Overall, African-American women are more likely to die of breast cancer. Asian, Hispanic, and Native-American women have a lower risk of developing and dying from breast cancer [6] (Fig. 4.3).

**Breast Cancer Risk Factors: Non-genetic and Inherited Genetic Factors**

The risk factors that are associated with the development of breast cancer are summarized in Table 4.1 and are discussed briefly in the next section. Age, reproductive factors, personal or family history of breast disease, genetic predisposition, and environmental factors have all been associated with an increased risk of developing female breast cancer.

**Fig. 4.1** Percent of new breast cancer cases by age group (Surveillance, Epidemiology, and End Results [SEER], 2011, <http://seer.cancer.gov/>) [8]




**Fig. 4.2** Incidence rates\* of female breast cancer by race and ethnicity, USA, 1999–2011. Combined data from the National Program of Cancer Registries as submitted to the Centers for Disease Control and Prevention (CDC) and from the Surveillance, Epidemiology, and End Results (SEER) program as submitted to the National Cancer Institute (NCI) in November 2013 [9].

\*Rates are per 100,000 and are age adjusted to the 2000 US standard population (19 age groups Census

P25-1130). Incidence rates are for state registries that meet USCS publication criteria for all years, 1999–2011. Incidence rates cover approximately 99 % of the US population.

\*Hispanic origin is not mutually exclusive with other race categories (white, black, Asian/Pacific Islander, American Indian/Alaska Native)



**Fig. 4.3** Death rates\* by race and ethnicity, USA, 1999–2011.

US mortality files, National Center for Health Statistics and Centers for Disease Control and Prevention (CDC) [10].

\*Rates are per 100,000 and are age adjusted to the 2000 US standard population (19 age groups Census

P25-1130). Death rates cover 100 % of the US population.

†Hispanic origin is not mutually exclusive with other race categories (white, black, Asian/Pacific Islander, American Indian/Alaska Native)

## Age

The risk of developing breast cancer increases with age. According to the Surveillance, Epidemiology, and End Results (SEER) database, the probability of a woman in the United States developing breast cancer is 1 in 8 over a lifetime: 1 in 202 from birth to age 39 years of age, 1 in 26 from 40 to 59 years, and 1 in 28 from 60 to 69 years [1]. Young women who develop breast cancer appear to have worse disease-free survival (DFS) and overall survival (OS) and present with more aggressive-appearing biological characteristics than older women [28].

## Personal History

A personal history of breast cancer is also a significant risk factor for the development of a

second ipsilateral or contralateral breast cancer. In fact, the most common cancer among breast cancer survivors is metachronous contralateral breast cancer [29]. Factors associated with an increased risk of a second breast cancer include an initial diagnosis of ductal carcinoma in situ (DCIS), stage IIB, hormone receptor-negative cancers, and young age [30].

## Breast Pathology

Proliferative breast disease is associated with an increased risk of breast cancer. Proliferative breast lesions without atypia, including usual ductal hyperplasia, intraductal papillomas, sclerosing adenosis, and fibroadenomas, confer only a small increased risk of breast cancer development, approximately 1.5–2 times that of



**Table 4.1** Risk factors for breast cancer

Risk factor	Category at risk	Comparison category	Relative risk
Alcohol intake [11]	2 drinks per day	Nondrinker	1.2
Body mass index [12]	80th percentile, age 55 or greater	20th percentile	1.2
Hormone replacement therapy with estrogen and progesterone [13]	Current user for at least 5 years	Never used	1.3
Radiation exposure [14, 15]	Repeated fluoroscopy Radiation therapy for Hodgkin's disease	No exposure	1.6 5.2
Early menarche [16]	Younger than 12 years	Older than 15 years	1.3
Late menopause [17]	Older than 55 years	Younger than 45	1.2–1.5
Age at first childbirth [18, 19]	Nulliparous or 1st child after 30	1st child before 20	1.7–1.9
Current age [20]	65 or older	Less than 65	5.8
Past history of breast cancer [21]	Invasive breast carcinoma	No history of invasive breast carcinoma	6.8
Other histologic findings [22]	Lobular carcinoma in situ Ductal carcinoma in situ	No abnormality detected	16.4 17.3
Breast biopsy [23]	Hyperplasia without atypia Hyperplasia with atypia Hyperplasia with atypia and positive family history	No hyperplasia	1.9 5.3 11
Cytology (fine needle aspiration) [24]	Proliferation without atypia Proliferation with atypia Proliferation with atypia and positive family history	No abnormality detected	2.5 4.9–5 18.1
Family history [25]	1st-degree relative 50 years or older with postmenopausal breast cancer 1st-degree relative with premenopausal breast cancer 2nd-degree relative with breast cancer Two 1st-degree relatives with breast cancer	No 1st- or 2nd-degree relative with breast cancer	1.8 3.3 1.5 3.6
Germline mutation [26]	Heterozygous for BRCA1, age <40 Heterozygous for BRCA1, age 60–69	Not heterozygous for BRCA1	200 15

Adapted from Singletary [27]

the general population [31, 32]. Atypical hyperplasia, including both ductal and lobular, which are usually incidentally found during screening mammography, confers a substantial increased risk of breast cancer. Women with atypia have an approximately 4.3 times greater risk of developing cancer compared with the general population [23, 33].

### Family History

A woman's risk of breast cancer is increased if she has a family history of the disease. In the

Nurses' Health Study follow-up, women with a mother diagnosed before age 50 had an adjusted relative risk (RR) of 1.69, and women with a mother diagnosed at 50 or older had an RR of 1.37 compared with women without a family history of breast cancer. A history of a sister with breast cancer was associated with an increased RR of 1.66 if the diagnosis was made prior to age 50 and a RR of 1.52 if diagnosed after age 50 compared with patients without a family history [33]. The highest risk is associated with an increasing number of first-degree

relatives diagnosed with breast cancer at a young age (younger than 50). Compared with women who had no affected relative, women who had one, two, or three or more affected first-degree relatives had risk ratios of 1.80, 2.93, and 3.90, respectively [34].

## **Endogenous Hormone Exposure and Reproductive Factors**

### **Early Menarche**

Early age at menarche is a risk factor among both pre- and postmenopausal women for developing breast cancer. Delay in menarche by 2 years is associated with a corresponding risk reduction of 10 % [35]. Within the European Prospective Investigation into Cancer and Nutrition cohort, women who had early menarche ( $\leq 13$  years) exhibited a nearly twofold increase in the risk of hormone receptor-positive tumors [36]. Among women with BRCA1 mutation-associated breast cancers, later age at menarche was associated with a later age at breast cancer diagnosis [37].

### **Parity and Age at First Full-Term Pregnancy**

Nulliparous women are at an increased risk for the development of breast cancer compared with parous women. Young age at first birth has an overall protective effect, whereas relatively advanced age at first birth confers a relative risk of breast cancer greater than that of a nulliparous woman. Compared with nulliparous women, the cumulative incidence of breast cancer in women experiencing their first birth at age 20, 25, and 35 years was 20 % lower, 10 % lower, and 5 % higher, respectively [38].

### **Breast-Feeding**

Evidence suggests that breast-feeding has a protective effect against the development of breast cancer. Breast-feeding may delay the return of regular ovulatory cycles and decrease endogenous sex hormone levels. There is an estimated 4.3 % reduction for every 1 year of breast-feeding [39].

### **Testosterone**

High endogenous sex hormone levels increase the risk of breast cancer in both premenopausal

and postmenopausal women. High levels of circulating testosterone in postmenopausal women have been linked to an increased risk of developing breast cancer (RR, 2.86–3.28) [40].

### **Age at Menopause**

Later onset of menopause has also been associated with increased breast cancer risk. Every year of delay in the onset of menopause confers a 3 % increase in risk, and every 5-year delay in the onset of menopause confers a 17 % increase in the risk of breast cancer [35, 41].

### **Exogenous Hormone Exposure**

Evidence suggests a relationship between the use of hormone replacement therapy (HRT) and breast cancer risk. Breast cancers related to HRT use are usually hormone receptor positive. When compared with patients who did not use HRT, breast cancer risk is higher in HRT users [42]. An international meta-analysis examining the risk of breast cancer with HRT found that in women who did not use HRT, the RR increased by a factor of 1.28 for each year older at menopause, comparable to the RR of 1.023 per year in women who used HRT or for those who ceased to use HRT up to 4 years previously [43].

In the Woman's Health Initiative randomized control trial, combined estrogen plus progestin in postmenopausal women with an intact uterus significantly increased the risk of breast cancer, delayed breast cancer detection and diagnosis, and significantly increased breast cancer mortality. The study was terminated early because of increased mortality in the combined estrogen plus progestin group. By contrast, the use of estrogen alone by postmenopausal women without a uterus did not interfere with breast cancer detection and significantly decreased the risk of breast cancer [44]. Data from the Nurses' Health Study, however, suggest that women who use unopposed postmenopausal estrogen increase their risk of breast cancer by 23 % at age 70 [45].

Timing and duration of HRT seem to be important factors associated with breast cancer risk as well. Breast cancer risk from exogenous hormone exposure is inversely associated with time from menopause. Women who initiate hor-

mone therapy closer to menopause have a higher breast cancer risk [46]. Long-term (>5 years) combined HRT use has been associated with the highest risk, whereas short-term use of combined estrogen–progestin therapy does not appear to confer a significantly increased risk (RR=1.023 per year) [43].

### **Lifestyle Factors**

Modifiable risk factors, including the excessive use of alcohol, obesity, and physical inactivity, account for 21 % of all breast cancer deaths worldwide [47].

### **Alcohol Consumption**

Alcohol consumption has been significantly associated with increased breast cancer risk at consumption levels as low as 5.0–9.9 g per day, which is equivalent to three to six drinks per week (RR=1.15; 95 % CI, 1.06–1.24; 333 cases/100,000 person-years). Binge drinking, but not frequency of drinking, is associated with breast cancer risk after controlling for cumulative alcohol intake. Alcohol intake both earlier and later in adult life is independently associated with risk [48].

### **Physical Activity**

Consistent physical activity reduces the risk of breast cancer in a dose-dependent manner, with modest activity conferring a 2 % decrease in risk and vigorous activity conferring a 5 % decrease in risk [49].

### **Obesity**

Obesity, specifically in postmenopausal women, also increases a woman's risk of breast cancer. In the EPIC multicenter prospective cohort study, postmenopausal women who did not use HRT had an elevated breast cancer risk with increasing weight, body mass index (BMI), and hip circumference [42]. In this cohort, the multivariate RR was 1.28 for overweight women (BMI 25.0–29.9) and obese women (BMI > 30.0) compared with women in the normal weight range. Lean women on HRT are incongruously at an increased risk of breast cancer (RR=2.04) compared with their overweight (1.93) and obese (1.39) counterparts [42].

Insulin resistance and hyperinsulinemia have been studied as risk factors for the comorbidities associated with obesity, including cardiovascular disease and diabetes. Insulin has anabolic effects on cellular metabolism, and human cancer cells overexpress the insulin receptor [50]. Hyperinsulinemia is an independent risk factor for breast cancer in non-diabetic postmenopausal women and may help explain the relationship between obesity and breast cancer [51].

### **Radiation**

Radiation exposure from various sources, including medical treatment and nuclear explosion, increases the risk of breast cancer. Radiation to the chest wall for treatment of childhood cancer increases the risk of breast cancer linearly with chest radiation dose [52]. Survivors of childhood cancers who received therapeutic radiation are at a dose-dependent risk for the development of breast cancer, and those treated for Hodgkin's disease are at highest risk (RR=7) [53]. Radiation effects on the development of female breast cancer were also demonstrated in Japan after the nuclear attacks on Hiroshima and Nagasaki [54] and positively correlated with age younger than 35 years at time of exposure. The incidence of breast cancer has also increased in areas of Belarus and Ukraine. A significant twofold increase was observed in the most contaminated areas around Chernobyl following the nuclear accident and manifested in women who were younger at the time of the exposure [55].

### **Mammographic Breast Density**

Mammographic breast density (MBD), alone or in combination with other risk factors, is associated with an increased risk of breast cancer [56–58]. Percentage dense area (PDA) is the most common measurement of mammographic density. A four- to sixfold greater risk of breast cancer has been reported in women for whom more than 75 % of the total area on mammogram is occupied by dense area [59]. In addition to PDA, the absolute dense area of the breast obtained during an assessment of PDA is an independent risk factor for breast cancer, and its inclusion in risk assessment tools has been proposed [60].

## Genetic Predisposition

Approximately 20–25 % of breast cancer patients have a positive family history, but only 5–10 % of breast cancer cases demonstrate autosomal dominant inheritance [61, 62]. Genetic predisposition alleles have been described in terms of clinical significance [63]. High-risk predisposition alleles conferring a 40–85 % lifetime risk of developing breast cancer include BRCA1 and BRCA2 mutations, mutations in TP53 resulting in Li–Fraumeni syndrome, phosphatase and tensin homolog (PTEN) resulting in Cowden syndrome, STK11 causing Peutz–Jeghers syndrome, neurofibromatosis (NF1), and (CDH-1) E-cadherin [64]. Half of the breast cancer predisposition syndromes are associated with mutations in BRCA1 and BRCA2. Women with BRCA1 or BRCA2 deleterious mutations have a significantly higher risk of developing breast cancer than the general population. Lifetime breast cancer risk ranges from 65 to 81 % for BRCA1 mutation carriers and 45 to 85 % for BRCA2 carriers [65–67]. Moderate-risk genes, including homozygous ataxia-telangiectasia (ATM) mutations [68], somatic mutations in tumor suppressor gene CHEK2, and the BRCA1 and BRCA2 modifier genes BRIP1 [69] and PALB2 [70], confer a 20–40 % lifetime risk of breast cancer. A study suggested an association between germline TP53 mutations and early-onset HER2-positive breast cancer [71]. Numerous low-risk common alleles have been identified, largely through genome-wide association studies [63], and the clinical implications of these mutations have not been determined.

## Genetic Testing and Management of Patients with Hereditary Breast Cancer

### Role of BRCA Genes

BRCA1 and BRCA2 function as tumor suppressor genes and are important in the maintenance of genomic stability through their role in DNA damage signaling and DNA repair. Both BRCA1 and BRCA2 have been implicated in mediating the repair of double-strand breaks by homologous recombination (HR) by interactions with RAD51.

Upon DNA damage, BRCA1 associates with RAD51 and localizes to the damaged region; BRCA1 is then phosphorylated. BRCA2 functions downstream of BRCA1 by forming a complex with RAD51. The primary function of BRCA2 is to facilitate HR [72]. Cells deficient for BRCA1 or BRCA2 are unable to repair double-strand breaks via error-free HR, resulting in repair via the error-prone non-homologous end-joining (NHEJ) pathway, which introduces chromosomal instability [73, 74]. During S-phase, the expression levels of BRCA1 and BRCA2 increase, indicating a function in maintaining genomic stability during the DNA replication process [75]. In addition to its role in HR, BRCA1 appears to have functions in DNA repair. BRCA1 is also part of the BRCA1-associated genome surveillance complex (BASC), which includes ATM, RAD50, MRE11, NBS1, and the mismatch repair proteins MLH1, PMS2, MSH2, and MSH6 [76]. BRCA1 is also involved in transcription-coupled excision repair, chromatin remodeling, and, together with BARD1, the ubiquitination process, through which proteins are tagged for degradation by the proteasome [72, 77].

A germline mutation in *BRCA1* or *BRCA2* only represents the first hit in the classical Knudson's two-hit hypothesis, whereas the second inactivating somatic mutation often involves deletion of the wild-type allele, termed loss of heterozygosity (LOH). LOH is present in the majority (80 %) of tumors arising from mutation carriers [78, 79]. By contrast, small somatic mutations involving a single or few bases are very rare [80]. Another somatic inactivation mechanism, epigenetic silencing by promoter methylation, has been reported for *BRCA1* in 9–13 % of sporadic breast tumors and up to 42 % of non-*BRCA1/BRCA2* hereditary breast tumors, leading to reduced *BRCA1* expression [81, 82]. By contrast, *BRCA1* promoter methylation is rare in tumors from *BRCA1* and *BRCA2* mutation carriers [83], and *BRCA2* promoter methylation in general is seldom observed in both sporadic and hereditary breast cancers [84]. Genetic testing for BRCA mutations is now widely available, and multiple professional societies have published

guidelines for testing and management. Genetic testing trends include utilization of multi-gene panels that take advantage of next-generation sequencing and testing for low- and moderate-penetrance susceptibility genes [85].

### **BRCA1- and BRCA2-Associated Breast Cancers**

Up to 10 % of breast cancers result from specific genetic mutations in BRCA1 and BRCA2, which are associated with hereditary breast/ovarian cancer syndrome; CHEK2 and p53, which are associated with Li–Fraumeni syndrome; and PTEN, which is associated with Cowden syndrome [86]. Families carrying genetic mutations in the above-mentioned genes exhibit an apparently dominant inheritance pattern and are often characterized by early age of onset and overrepresentation of ovarian, bilateral breast, and male breast cancers [87].

Early reports suggested that germline mutations in the genes BRCA1 and BRCA2 were responsible for the majority of hereditary breast cancers, although more recent studies have demonstrated that mutations in the two genes only account for 25–28 % of the family risk [88, 89]. However, additional BRCA1/BRCA2 mutations likely remain undetected by the screening methods used today. Women carrying a BRCA1 or BRCA2 germline mutation also have increased risk of developing ovarian cancer and fallopian tube cancer. In addition, BRCA1/BRCA2 mutation carriers also have increased risk of other cancer types, such as male breast cancer, prostate cancer, pancreatic cancer, gastrointestinal cancers (e.g., gall bladder, bile duct, and stomach), and melanoma [90, 91]. In a large study by the Consortium of Investigators of Modifiers of BRCA1/BRCA2 (CIMBA), the median age of breast cancer diagnosis was 40 years among BRCA1 and 43 years among BRCA2 mutation carriers [92].

Although germline mutations in BRCA1 and BRCA2 confer high risk of breast and ovarian cancers, the penetrance of these genes is incomplete. The risk of developing breast cancer by the age of 70 is 45–87 % in BRCA1 and BRCA2 mutation carriers. For ovarian cancer, the risk is 45–60 % among BRCA1 mutation carriers and

11–35 % among BRCA2 mutation carriers [66, 67, 93]. However, the penetrance depends on several different factors, including the type of mutation and exogenous factors.

The majority of invasive breast cancers that arise in BRCA1 and BRCA2 carriers are invasive ductal carcinomas (IDCs) (80 %) [94]. A higher frequency of BRCA1 tumors is classified as medullary carcinomas compared with sporadic tumors (9 % versus 2 %, respectively) [92, 95]. Notably, 11 % of medullary carcinomas carry BRCA1 germline mutations [96]. By contrast, an excess of invasive lobular and tubular carcinomas has been reported for BRCA2 tumors relative to BRCA1 tumors [95]. BRCA1 tumors are more frequently high grade compared with sporadic tumors [97]. Most BRCA2 tumors are grade 2/3 with high mitotic rates.

A recent study examining pathology data from 4325 BRCA1 and 2568 BRCA2 mutation carriers reported that 78 % of tumors arising in BRCA1 carriers were estrogen receptor (ER) negative, while only 23 % of tumors arising in BRCA2 mutation carriers were ER negative. Furthermore, HER2 overexpression was only observed in approximately 10 % of tumors from mutation carriers. Consequently, 69 % of the BRCA1 tumors were triple negative (TN), which was true for only 16 % of the BRCA2 tumors [92]. In contrast to BRCA1 tumors, BRCA2 tumors seem to be more similar to sporadic tumors with respect to the expression of IHC markers. Most BRCA2 breast tumors exhibit a luminal phenotype featuring overexpression of ER, progesterone receptor (PR), and cytokeratins CK8 and CK18 [98].

Recent studies have observed preinvasive lesions both in prophylactic mastectomy specimens from mutation carriers and in normal breast tissue adjacent to breast cancers [99]. Among BRCA1-/BRCA2-associated breast cancers, 59 % had at least one associated preinvasive lesion compared with 75 % of controls. Preinvasive lesions were more prevalent in BRCA2 mutation carriers than in BRCA1 mutation carriers (70 % versus 52 %, respectively). The most common preinvasive lesion in both groups was DCIS; 56 % of BRCA1-/BRCA2-associated breast cancers and 71 % of



the sporadic breast cancers had adjacent intra-ductal disease, respectively [99]. These findings suggest that BRCA1-/BRCA2-associated breast cancers progress through the same intermediate steps as sporadic breast cancers and that DCIS should be considered part of the BRCA1/BRCA2 tumor spectrum.

While most studies indicate a similar prognosis for women with hereditary breast cancers compared with age-matched women with sporadic breast cancers [100–106], other studies have reported worse survival outcomes [107–111]. Lee et al. [112] reported similar survival rates in BRCA1 mutation carriers with TN disease compared with non-carriers. Confirming those findings, Bayraktar et al. [113] observed a 50 % prevalence of deleterious BRCA1/BRCA2 mutations in high-risk women diagnosed with TN breast cancer. Overall prognosis of TN breast cancer in BRCA carriers and non-carriers was not significantly different within the first 5 years following initial diagnosis.

### Genetic Counseling

A cancer genetic counseling risk assessment typically involves collecting a three- to four-generation family medical history (pedigree), which should include information such as current age/age at death, personal cancer history for each individual (cancer type, age at diagnosis, and pathology and treatment, if known), environmental exposures/lifestyle factors, and ethnicity [114]. The information obtained about the individual's personal and family medical history should then be used to estimate her cancer risk and the likelihood of a hereditary cancer syndrome. The National Comprehensive Cancer Network (NCCN) has guidelines regarding who should be offered genetic testing on the basis of personal and family history. Women who meet one or more of the following familial/hereditary breast cancer risk criteria should be referred to a cancer genetic counselor for further evaluation: individuals from a family with known mutations that increase their risk of breast cancer (*BRCA1*, *BRCA2*, *CDH1*, *STK11*, and *TP53*) or genes associated with breast cancer, a family history of two or more primary breast cancers in a single

individual, two or more members with breast primaries on the same side of the family, first- or second-degree relative  $\leq 45$  years of age with breast cancer, one or more primary ovarian cancers on the same side of the family, family history of male breast cancer, or one or more family members on the same side of the family with an aggressive early-onset cancer in addition to breast cancer [115]. Recently, BRCA mutation testing for women with TN breast cancers who were younger than 50 years at diagnosis was found to be a cost-effective strategy and was adopted into current guidelines for genetic testing [116]. Moreover, women who have DCIS and a family history of ovarian cancer or who have BRCAPRO scores  $\geq 10$  % have a high rate of BRCA positivity regardless of age at diagnosis [117]. The prophylactic contralateral mastectomy rate among patients with DCIS who undergo BRCA genetic testing is high [118]. Factors associated with increased likelihood of prophylactic contralateral mastectomy among this group are age, BRCA positivity, and a family history of ovarian cancer. Therefore, high-risk patients with DCIS may be appropriate candidates for genetic testing for BRCA mutations in the presence of predictive factors even if they do not have invasive breast cancer.

### Risk Assessment Models

Several models have been developed to predict an individual's lifetime risk for a specific cancer and/or their risk of having a genetic mutation. Empiric models of breast cancer risk assessment include the Claus, Gail, Tyrer–Cuzick, and BRCAPRO models. Each model incorporates different risk factors and thus may be utilized to provide a range of risk estimates.

The Claus model estimates a woman's lifetime risk for breast cancer based on her family history of breast cancer in first- and/or second-degree relatives [119]; however, the model does not include any other risk factors. The American Cancer Society guidelines for recommending MRI include women with a lifetime risk for breast cancer of 20–25 % or greater, as defined by models that largely depend on family history, such as the Claus model [120]. These

guidelines, however, are based on trials that did not consider the Gail model or “tissue risks” (ADH, ALH/LCIS). Therefore, Hollingsworth and Stough suggest that the current guidelines do not account for risk factors that increase the likelihood of mammography failure; thus, a different approach may be needed to determine MRI candidates [121].

The Gail model estimates a woman’s risk for breast cancer based on her age and personal risk factors, such as age at menarche, age at first live birth, and biopsy history; however, this model inadequately utilizes family history because only first-degree relatives are considered [122]. This model has been extensively validated in US populations; however, limited data are available related to the use of this model internationally. Pastor-Barriuso et al. recently evaluated the predictive accuracy of the Gail model in a Spanish cohort and found that the original Gail model cannot be applied to populations with varying rates of invasive breast cancer; however, a recalibrated version of the model provides more unbiased risk estimates [123].

The Tyrer–Cuzick model (also known as IBIS) incorporates personal and family history risk factors and can be utilized to calculate a woman’s risk of breast cancer and the likelihood of a BRCA mutation [124]. The BOADICEA model was designed to predict the probability of BRCA1/BRCA2 mutation and to provide breast and ovarian cancer risk estimates, and it uniquely accounts for the possibility of genetic modifiers and the polygenic component of breast cancer risk [125–127]. Several studies have assessed the validation and accuracy of these risk prediction tools in specific, high-risk cohorts. The IBIS model was well calibrated for women of Marin County, California, a population with high rates of breast cancer [128]. In addition, the BOADICEA and IBIS risk models were validated in a high-risk population of Israeli women, which revealed that the BOADICEA model has better predictive value and accuracy for 10-year breast cancer risk than the IBIS model [129].

BRCAPRO predicts the probability of a BRCA mutation based on an individual’s personal and family history of cancer and can be utilized

for an affected or unaffected patient [130, 131]. Several validation studies in various populations have indicated that BRCAPRO may accurately measure the probability of identifying a BRCA mutation; thus, the tool is widely utilized in clinical practice [132–134]. One recent study, however, examined the accuracy of BRCAPRO specifically in patients with a personal history of ovarian cancer and reported that the model significantly underestimated BRCA1/BRCA2 mutations in individuals with high-grade serous ovarian cancer at BRCAPRO scores less than 40 %, suggesting a benefit of universal testing in this patient population [135]. Another recent study evaluated the performance of the BRCAPRO and BOADICEA models in Italian cancer genetics clinics and reported that using these models at the commonly used 10 % threshold for testing missed 25 % of carriers; these data question the strict utilization of these models for risk assessment [136]. A study of a German patient cohort supported the use of BRCAPRO and BOADICEA for decision making related to BRCA1/BRCA2 genetic testing; however, the model calibration may need to be improved for this specific population [137]. Ready et al. [138] demonstrated that the BRCAPRO model overestimated the relative contribution of bilateral breast cancer to the likelihood of detecting a BRCA1 or BRCA2 mutation. In that study, bilateral breast cancer did not appear to be a good indicator of mutation status, particularly for women whose age at first diagnosis was >40 years. Clinical providers should proceed with caution when strictly using risk prediction tools, such as BRCAPRO, to determine if an individual should undergo genetic testing. Considerations of demographics (e.g., ethnicity), cancer histology and tumor markers, relatives with cancer beyond 1st and 2nd degrees of relatedness, limited family structure, and the possibility of misreported or unknown family history should all be considered during risk assessment.

Currently, there is a lack of validated models to predict an individual’s lifetime risk for gynecological malignancies such as endometrial and ovarian cancer. Pfeiffer et al. recently derived and validated a model to predict a woman’s absolute

risk for developing breast, endometrial, and ovarian cancers [139]. The model incorporates several risk factors, such as parity, menopausal status, age at menopause, BMI, smoking history, and family history. The validation study demonstrated expected to observed cancer ratios of 1.00 (95 % CI, 0.96–1.04) for breast cancer and 1.08 (95 % CI 0.97–1.19) for ovarian cancer; however, the number of endometrial cancers was significantly overestimated. There is currently a significant lack of literature related to risk prediction models for endometrial and ovarian cancer. Further research is needed to develop tools that can be utilized for individualized risk assessment to allow management recommendations to be made in the context of a woman's personal and family history risk factors.

### **Risk-Reducing Surgical Interventions in Women with BRCA Mutation-Associated Breast Cancer**

#### **Prophylactic Contralateral Mastectomy**

In addition to the risk of ipsilateral recurrence (IPR), breast cancer patients with a deleterious BRCA1 mutation have up to a 43.4 % 10-year risk of contralateral breast cancer (CBC), while BRCA2 mutation carriers have up to a 34.6 % 10-year risk [140]. Importantly, studies have shown that prophylactic bilateral mastectomy results in up to a 97 % risk reduction of CBC [141–143]. The decision for prophylactic bilateral mastectomy vs. ipsilateral mastectomy should be based on the type of surgery the patient is undergoing for the treatment of the primary breast cancer diagnosis and, of course, the patient's choice. If the patient is undergoing lumpectomy for ipsilateral breast cancer, then prophylactic contralateral mastectomy may not be the best option for that patient considering cosmesis. For patients who do not want to have contralateral prophylactic mastectomy, breast cancer screening with MRI every 6 months should be offered, and tamoxifen can be used as a chemoprevention in ER-positive breast cancer types.

Similar to the risk of IPR, the risk of CBC in BRCA mutation carriers is increased in patients

who are diagnosed at a younger age [144–146]. However, the impact of other cancer therapies on the risk of CBC in BRCA mutation carriers is controversial. While some studies have reported a 50–60 % reduction in the risk of CBC with adjuvant chemotherapy [147, 148], other studies have shown no impact of adjuvant chemotherapy [141, 146, 149]. Likewise, some studies have reported that tamoxifen reduces the risk of CBC by 50–70 % in BRCA mutation carriers [148, 150, 151], whereas other studies have not reported a significant reduction [141, 146, 147, 149, 152]. However, most studies have reported that oophorectomy reduces the risk of CBC in BRCA mutation carriers by 50–70 %, with the greatest benefit observed if the surgery is performed before the age of 50 [141, 144, 148, 150].

The above findings justify the practice of offering the option for risk-reducing surgery to the intact breast (before any breast cancer diagnosis) in women with BRCA mutations. Despite the significant reduction in the risk of CBC associated with prophylactic contralateral mastectomy in BRCA mutation carriers, the procedure has not currently been found to improve survival, although studies have been limited by short follow-up [142, 153].

#### **Risk-Reducing Salpingo-Oophorectomy**

BRCA mutation carriers have a well-established enhanced risk of ovarian cancer [66, 154, 155]. Risk-reducing salpingo-oophorectomy (RRSO) clearly reduces the risk of tubal-ovarian cancer [156, 157] and also reduces breast cancer [158]. A meta-analysis of ten studies reported a significant reduction in the risk of CBC and ovarian cancer in BRCA mutation carriers who had undergone RRSO [159] (Table 4.2). Overall, RRSO was associated with a 51 % reduction in breast cancer risk (HR, 0.49; 95 % confidence interval (CI) 0.37–0.65). Similar risk reductions were observed in BRCA1 (HR 0.47; 95 % CI 0.35–0.64) and BRCA2 (HR 0.47; 95 % CI 0.26–0.84) mutation carriers. RRSO was also associated with a significant risk reduction of ovarian–fallopian tube cancer (HR 0.21; 95 % CI 0.12–0.39).

Because BRCA1 carriers more typically have hormone receptor-negative breast cancer, the

**Table 4.2** Use of pharmacological interventions for breast cancer risk reduction: ASCO Clinical Practice Guidelines 2014

<i>Intervention</i>
Pharmacological interventions for breast cancer risk reduction, including selective estrogen receptor (ER) modulators and aromatase inhibitors
<i>Key recommendations</i>
Tamoxifen (20 mg per day orally for 5 years) should be discussed as an option to reduce the risk of invasive breast cancer, specifically ER-positive breast cancer, in premenopausal or postmenopausal women age 35 years at increased risk of breast cancer or with lobular carcinoma in situ (LCIS). Tamoxifen is not recommended for use in women with a history of deep vein thrombosis, pulmonary embolus, stroke, or transient ischemic attack; during prolonged immobilization; or in women who are pregnant, may become pregnant, or are nursing mothers. Tamoxifen is not recommended in combination with hormone therapy
Raloxifene (60 mg per day orally for 5 years) should be discussed as an option to reduce the risk of invasive breast cancer, specifically ER-positive breast cancer, in postmenopausal women age $\geq 35$ years at increased risk of breast cancer or with LCIS. It should not be used for breast cancer risk reduction in premenopausal women. Raloxifene is not recommended for use in women with a history of deep vein thrombosis, pulmonary embolus, stroke, or transient ischemic attack or during prolonged immobilization
Exemestane (25 mg per day orally for 5 years) should be discussed as an alternative to tamoxifen or raloxifene to reduce the risk of invasive breast cancer, specifically ER-positive breast cancer, in postmenopausal women age $\geq 35$ years at increased risk of breast cancer or with LCIS or atypical hyperplasia. Exemestane should not be used for breast cancer risk reduction in premenopausal women
For tamoxifen and raloxifene, the most favorable risk–benefit profile is seen in women with the greatest risk of developing breast cancer
Discussions with patients and healthcare providers should include both the risks and benefits of each agent under consideration

Reprinted with permission. © (2013) “American Society of Clinical Oncology. All rights reserved” [160]

efficacy of RRSO in reducing the risk of ovarian and breast cancer has been questioned in the past. In a multicenter, prospective cohort study by Domchek et al. [153], salpingo-oophorectomy was associated with a reduction of ovarian cancer and ovarian cancer mortality, a reduction of subsequent breast cancer and breast cancer mortality,

and a reduction of overall mortality. In this study of 2482 women with BRCA mutations, RRSO was associated with a significant (85 %) reduction (HR=0.15; 95 % CI, 0.04–0.63) in the risk of ovarian cancer among BRCA1 mutation carriers with a history of breast cancer. Among BRCA2 mutation carriers with and without a history of breast cancer, no cases of ovarian cancer were observed after RRSO. Most importantly, RRSO was associated with a significant mortality benefit in BRCA mutation carriers, both in patients with and without history of breast cancer. In the first group, RRSO reduced all-cause mortality by 70 % (HR=0.3; 95 % CI, 0.17–0.52), and interestingly, breast cancer-specific mortality was reduced by 65 % (HR=0.35; 95 % CI, 0.19–0.67), despite a lack of benefit in breast cancer incidence.

The optimal timing of prophylactic oophorectomy is still controversial. However, the risk of second primary breast cancer development or ipsilateral breast cancer recurrence cumulatively increases with younger age at initial diagnosis (<40 vs. >50 years of age) [145]. A few studies have shown that the survival outcomes are better if prophylactic oophorectomy is performed before 40 years of age [161, 162]. However, surgical menopause in premenopausal women is associated with long-term side effects, including increased risks of osteoporosis, heart disease, high cholesterol, and hot flashes [163]. Therefore, the associated side effects and the resultant decreased quality of life should be weighed against the benefits of this procedure. In summary, both prophylactic bilateral mastectomy and RRSO result in a reduction of breast cancer in mutation carriers, and the risk of CBC in women with BRCA mutations who undergo both procedures is less than 2 % [142, 143, 164]. A detailed discussion with the patient regarding the surgical risk-reducing intervention and its long-term side effects is central to the management of mutation carriers.

### Cancer Risk Management Decisions of Women with BRCA Variants of Uncertain Significance

One result of BRCA genetic testing is a variant of uncertain significance (VUS). VUSs are changes

in the BRCA genes that may or may not be associated with an increased risk of cancer. Women with a VUS are a growing population, making up to 21 % of patients who undergo genetic testing. Because the cancer risks associated with VUSs are unknown, cancer risk management recommendations are difficult, and patients must make decisions regarding risk management for cancers that they may or may not be at risk for. A recent study described the risk management decisions (RMDs) undertaken by women who have a BRCA VUS [165]. Women who had a BRCA VUS and a personal history of breast cancer appeared to be more aggressive in their RMD than women who had a VUS but no personal history of breast or ovarian cancer. Given the prognostic uncertainty and high rate of reclassification for women with a VUS, individualizing counseling and directing efforts toward surveillance, chemoprevention, or salpingectomy are currently recommended.

### **Systemic Therapy Options for Women with BRCA Mutation-Associated Breast Cancer**

#### **Cytotoxic Chemotherapy**

Traditionally, for those who develop breast or ovarian cancer, systemic therapy has been selected similarly to those with sporadic cancers, and the choice of chemotherapy (adjuvant or neoadjuvant as appropriate), endocrine therapy, and radiation has been based on ER/PR/HER2 status, lymph node involvement, and the size of the tumor. Studies have reported survival outcomes of BRCA carriers that are equivalent to those of patients with sporadic breast cancer after appropriate treatment [102, 106]. However, the approach to treatment is changing based on the recent data suggesting unique patterns of sensitivity and resistance to systemic therapies in BRCA mutation-associated breast cancers [166–171].

Due to the involvement of the BRCA1/BRCA2 protein products in DNA repair mechanisms, BRCA mutational status may impact sensitivity to different chemotherapeutic agents [172–174]. In vitro studies have demonstrated that BRCA1-defective cell lines are sensitive to DNA-damaging

agents, such as platinum, and are relatively resistant to taxanes compared with BRCA-competent cell lines [175, 176]. Several subsequent clinical studies have supported these preclinical findings [166, 167, 169]. Byrski et al. reported a remarkable pathological complete response (pCR) rate of 80 % in a small prospective trial evaluating neoadjuvant cisplatin in BRCA1 mutation-associated breast cancer [166]. The promising neoadjuvant data with cisplatin initiated a randomized phase III trial comparing carboplatin to docetaxel in metastatic BRCA mutation-associated breast cancer (NCT00321633) and a smaller phase II trial evaluating cisplatin for metastatic BRCA1 mutation-associated breast cancer. Early results from the phase II trial have been encouraging, with 46 % of women achieving a complete response and 26 % of women achieving a partial response. However, a recent study from MD Anderson demonstrated that BRCA1 carriers had a high pCR to neoadjuvant anthracycline–taxane-based chemotherapy (pCR in 46 % of BRCA1 carriers vs. 22 % of non-carriers) [168]. Of note, the sensitivity to single-agent taxanes was low in this study. Interestingly, BRCA mutation status and ER negativity were independently associated with higher pCR rates. Similarly, in another study among women with metastatic breast cancer, hormone receptor-negative BRCA1 mutation-associated breast cancer patients had lower response rates and shorter time to progression with a taxane-containing regimen compared with hormone receptor-negative sporadic breast cancer controls [177]. Regarding the current efficacy of other cytotoxic agents and targeted therapy, most studies have been directed against TN breast cancer and have assessed the roles of combinations of ixabepilone and cetuximab, gemcitabine and erlotinib, and paclitaxel and cetuximab [178, 179]. The specific responses to these combinations in BRCA mutation-associated breast cancers have not been analyzed.

#### **PARP Inhibitors**

With advances in molecularly targeted therapy in solid tumors, an appealing targeted therapy for BRCA1/BRCA2 carriers, poly(ADP-ribose) polymerase (PARP) inhibitors, has also been



developed. PARP proteins play a role in single-strand DNA repair; when PARP is inhibited, single-strand breaks cannot be repaired, leading to double-strand breaks at the replication fork [180, 181]. Because BRCA1 and BRCA2 proteins are critical in double-strand DNA repair, combining PARP inhibition with tumors that have defective BRCA1 or BRCA2 proteins exerts a synergistic lethal effect [182, 183]. This hypothesis has been supported by *in vitro* studies showing enhanced cytotoxicity in BRCA1- and BRCA2-deficient cells compared with cells with wild-type BRCA proteins [184, 185].

In a phase I study of 60 patients, of whom 22 were BRCA carriers, patients were treated with two different dose levels of the PARP inhibitor olaparib, and the maximum tolerated dose (MTD) was determined to be 400 mg orally twice daily. The most common side effects were grade 1 or 2 nausea, vomiting, fatigue, dysgeusia, and anorexia. Myelosuppression (anemia or thrombocytopenia) was also observed in a few patients [171]. All patients had a partial response, according to the Response Evaluation Criteria in Solid Tumors (RECIST), with responses lasting 20–80 weeks in the 19 BRCA mutation carriers with ovarian, breast, or prostate cancer who could be evaluated for tumor response.

Phase II multicenter, multinational studies that examined breast and ovarian cancers independently were then conducted in mutation carriers. Both used the phase I MTD, 400 mg twice daily and 100 mg twice daily, because this dose was the lowest dose at which an antitumor effect was seen in the phase I trial. The primary end point for both studies was the objective response rate (ORR). Among breast cancer patients, the ORR for those in the 400-mg arm was 41 % (11 of 27), with an additional 44 % (12/27) of women achieving stable disease. For those with metastatic breast cancer on 100-mg olaparib, 22 % (6/27) had partial responses, and an additional 44 % (12/27) of the patients achieved stable disease. These results were particularly impressive because the patients had undergone a median of three prior chemotherapy regimens. Similar results were observed in the ovarian cancer study, with an ORR of 33 % (11/33) in the 400-mg arm.

The most common side effects were nausea and fatigue [186, 187]. Both studies demonstrated the efficacy and tolerability of olaparib in BRCA1 or BRCA2 mutation carriers with breast or ovarian cancer. These data resulted in a paradigm shift assuming that BRCA1 or BRCA2 carriers have differential susceptibility to systemic therapy compared with non-carriers. Multiple studies are ongoing and examining different PARP inhibitors with the hope that BRCA1/BRCA2 mutation status will provide a targeted group for these agents.

In a randomized phase II trial, the intravenous PARP inhibitor iniparib was evaluated in combination with gemcitabine and carboplatin for the treatment of women with TN metastatic breast cancer [188]. The arm that was treated with the combination regimen had greater clinical benefit and improved progression-free survival and overall survival when compared with the chemotherapy alone arm [189]. Unfortunately, the results of a randomized phase III trial comparing iniparib plus chemotherapy to chemotherapy alone in TN metastatic breast cancer were disappointing, with no demonstration of survival benefit when used as a first-line treatment in the metastatic setting. Of note, iniparib was not specifically evaluated in BRCA mutation-associated breast cancer in this study.

### Endocrine Therapy

Two studies have demonstrated that adjuvant use of tamoxifen reduces the risk of IPR or CBC in BRCA1/BRCA2 carriers, regardless of ER status [151, 190]. A small retrospective study comparing outcomes in early-stage BRCA mutation-associated and sporadic breast cancer treated with endocrine therapy observed a lower OS in BRCA carriers, suggesting relative resistance to adjuvant endocrine therapy with tamoxifen [191]. These results, however, require confirmation, and the use of adjuvant endocrine therapy is recommended in patients with BRCA mutation-associated ER-positive breast cancer. Currently, there are no data regarding outcomes with aromatase inhibitors as adjuvant endocrine therapy in BRCA mutation-associated breast cancer.

## Increased Surveillance

Current screening recommendations for the asymptomatic *BRCA* mutation carrier encompass examination, imaging, and laboratory evaluation. Surveillance for female carriers emphasizes screening techniques for breast and ovarian cancers.

There is general agreement that women with a higher lifetime risk of breast cancer, such as that conferred by a *BRCA* mutation, should undergo earlier and more frequent screening, with additional imaging modalities considered. A consolidated summary of current screening recommendations published by the National Comprehensive Cancer Network (NCCN), American Cancer Society (ACS), American College of Radiology (ACR), and other national organizations for the asymptomatic, female, *BRCA* mutation carrier includes the following [192, 193]:

- Monthly breast self-exam (BSE) beginning at the age of 18 years
- Semiannual clinical breast exam (CBE) beginning at the age of 25 years
- Alternating annual mammograms with annual breast magnetic resonance imaging (MRI) beginning at the age of 25–30 years or individualized based on the earliest age of cancer onset in the family [194]

While risk-reducing prophylactic oophorectomy is more effective in preventing ovarian cancer in these women compared to general population, some may not opt to pursue this intervention until after their childbearing years. In the absence of more effective screening methods, transvaginal ultrasound (TVU) and CA-125 levels continue to be recommended and endorsed by national organizations for women who are at high risk for hereditary breast and ovarian syndromes [195]. Current NCCN screening guidelines for *BRCA* mutation carriers who are not undergoing prophylactic oophorectomy include the following:

- Semiannual concurrent pelvic exam, TVU, and CA-125 antigen determination beginning at the age of 35 years or 5–10 years earlier

than the youngest age at which any family member was diagnosed with ovarian cancer

Mutations in the tumor suppressor genes *BRCA1* and *BRCA2* place male and female carriers at increased risk for a number of other cancers, notably pancreatic, melanoma, colorectal, and other gastrointestinal tumors. Further research is needed to define the association of specific *BRCA* mutations with these cancers. No expert consensus or evidence-based guidelines exist regarding screening for these cancers. Some literature and investigational studies support considering the following additional surveillance modalities [196–198]:

- Pancreatic: annual endoscopic ultrasound, beginning at the age of 50 years or 10 years prior to the earliest pancreatic cancer diagnosis in the family
- Melanoma: annual full body skin and ocular exam
- Colorectal: population screening guidelines, beginning at the age of 50 years and continuing until 75 years old
- Annual fecal occult blood testing
- Sigmoidoscopy every 5 years or colonoscopy every 10 years

## Breast Cancer Prevention

Due to the rising incidence of breast cancer and because several of the risk factors are non-modifiable, strategies for the primary prevention of breast cancer represent an important area of interest. In this section, we will review the different approaches directed at reducing the incidence of breast cancer.

### Pharmacotherapy (Chemoprevention)

The effects of various pharmacologic agents on the incidence of invasive breast cancer (IBC) and noninvasive breast cancer have been investigated in several prospective randomized clinical trials [199]. In this review, we will discuss the role of selective estrogen receptor modulators (SERMs) such as tamoxifen, raloxifene, arzoxifene, and

lasofoxifene and aromatase inhibitors (AIs) such as exemestane. Table 4.2 provides a summary of the prior (2009) ASCO guidelines and updated recommendations.

### Tamoxifen

#### National Surgical Adjuvant Breast and Bowel Project (NSABP) Breast Cancer Prevention Trial (P1)

The NSABP-P1 trial, which was initiated in 1992, randomized 13,388 women to receive either 20 mg of tamoxifen or a placebo daily for 5 years [200]. After a median follow-up of 54.6 months, a 49 % reduction in the risk of IBC was observed in the patients who were treated with tamoxifen (RR=0.51; 95 % confidence interval [CI], 0.39–0.66). The cumulative incidence of IBC through 69 months was 43.4 versus 22.2 per 1000 women in the placebo and tamoxifen groups, respectively. Tamoxifen was effective across all age groups, in patients with a history of LCIS or atypical ductal hyperplasia, and in those with any category of predicted 5-year risk. Tamoxifen reduced the occurrence of IBC in ER-positive tumors by 69 % (RR=0.31; 95 % CI, 0.22–0.45), but no significant difference in the occurrence of ER-negative tumors was observed.

The increased incidence of endometrial cancer, stroke, deep venous thrombosis, and cataracts was noted in patients treated with tamoxifen, with most cases occurring in women who were >50 years of age. All endometrial cancers in the tamoxifen group were International Federation of Gynecology and Obstetrics stage I. In 2005, the NSABP provided the 7-year follow-up results of the above study, which continued to demonstrate a reduced incidence of both IBC (RR=0.57; 95 % CI, 0.46–0.70) and noninvasive breast cancer (RR =0.63; 95 % CI, 0.45–0.89) [201]. A 32 % reduction in osteoporotic fractures was also noted with tamoxifen.

#### Italian Tamoxifen Prevention Study

The Italian Tamoxifen Prevention Study randomized 5408 women who had previously undergone a hysterectomy to receive tamoxifen or placebo [202]. The initial results of the trial failed to dem-

onstrate an overall benefit of tamoxifen after a median follow-up of 46 months; however, after 11 years of follow-up, the investigators observe a significant reduction in the incidence of ER-positive breast cancer among women at high risk (defined as women taller than 160 cm, with at least one intact ovary, with no full-term pregnancy before the age of 24 years, and younger than age 14 years at menarche) treated with tamoxifen compared to women not treated with tamoxifen (6.26 versus 1.50 per 1000 woman-years; RR=0.24; 95 % CI, 0.10–0.59) [203].

#### The Royal Marsden Hospital Tamoxifen Chemoprevention Trial

The Royal Marsden Hospital Tamoxifen Chemoprevention Trial, which randomized 2494 women aged 30–70 years who also had a family history of breast cancer to tamoxifen or placebo, failed to demonstrate a decreased incidence of ER-positive breast cancer (30 cases in the tamoxifen arm versus 39 in the placebo arm; hazard ratio [HR]=0.77; 95 % CI, 0.48–1.23) [204]. In 2007, the investigators provided an update to this trial with an extended follow-up of 20 years, which revealed a significant decrease in the risk of ER-positive breast cancer in the tamoxifen arm (23 cases in the tamoxifen arm and 47 in the placebo arm; HR=0.48; 95 % CI, 0.29–0.79) [205]. The adverse events observed with tamoxifen in the European trials were similar to the NSABP-P1 trial.

#### International Breast Cancer Intervention Study (IBIS-I)

Another trial testing the efficacy of tamoxifen among women at increased risk of breast cancer in the United Kingdom, Australia, and New Zealand was initiated in 1992 [206]. With a median follow-up of 49.6 months, the investigators determined that tamoxifen decreased the incidence of breast cancer by 32 % (RR=0.68; 95 % CI, 0.50–0.92). With further follow-up (up to 96 months), the incidence continued to be lower in the tamoxifen group (27 % reduction in IBC; RR=0.73; 95 % CI, 0.58–0.91) [207]. Similar to the NSABP-P1 experience, the benefit of tamoxifen was only observed in ER-positive tumors, and an increased risk of thromboembolic

events with tamoxifen was reported. However, in contrast to the NSABP-P1 results, the use of HRT for postmenopausal symptoms at the lowest possible dose was permitted in the trial, and the increased risk of endometrial cancer with tamoxifen was not significant.

In 2003, an overview of the abovementioned tamoxifen prevention trials was published, and there was no reduction in ER-negative IBC; however, there was a significant 48 % decrease in the incidence of ER-positive IBC [208]. The consensus of endometrial cancer and venous thromboembolic events had a RR of 2.4 and 1.9, respectively; women aged 50 years or older had an increased risk. Overall, there was no effect on all-cause mortality, but there was a high degree of heterogeneity across the various trials.

Several studies have demonstrated that tamoxifen decreases MBD [209–211]. A case-control study nested within the IBIS-I showed a 10 % or greater reduction in breast density at the 12- to 18-month mammogram in 46 % of women in the tamoxifen group [212]. These women had a 63 % reduction in breast cancer risk (odds ratio [OR]=0.37; 95 % CI, 0.20–0.69;  $P=0.002$ ). The women who experienced less than a 10 % reduction in breast density with tamoxifen had no risk reduction (OR=1.13; 95 % CI, 0.72–1.77;  $P=0.60$ ). Similar reductions in MBD in the placebo group were not associated with decreased risk of breast cancer; hence, the authors concluded that a 12- to 18-month change in MBD was a good predictor of the response to tamoxifen for the prevention of breast cancer.

### Raloxifene

Raloxifene is an oral, second-generation SERM that has estrogenic effects on the bone, lipid metabolism, and blood clotting and antiestrogenic effects on the breast and uterus. The US Food and Drug Administration (FDA) initially approved raloxifene for the prevention and treatment of osteoporosis in postmenopausal women [213].

#### The Multiple Outcomes of Raloxifene Evaluation (MORE) Trial

In this trial, 7705 postmenopausal women with osteoporosis were randomly assigned to receive

raloxifene (60 mg or 120 mg per day) or placebo [214]. The initial results of this trial reported a 30 % reduction in the risk of vertebral fractures, which was associated with an increase in bone mineral density in the spine and femoral neck, but the incidence of non-vertebral fractures was not significantly different. The incidence of IBC, which was a secondary end point of the study, was decreased by 76 % during the 3 years of treatment and by 72 % after 4 years of treatment with raloxifene. The number needed to treat (NNT) to prevent one case of breast cancer was 126 [215, 216]. Similar to the tamoxifen trials, the benefit of raloxifene was limited to ER-positive breast cancer, and an increased risk of venous thromboembolism was observed (RR=3.1; 95 % CI, 1.5–6.2). In contrast to tamoxifen, raloxifene did not increase the risk of endometrial cancer (RR=0.8; 95 % CI, 0.2–2.7).

#### The Continuing Outcomes Relevant to Evista (CORE) Trial

This double-blind, placebo-controlled study investigated the efficacy of an additional 4 years of raloxifene compared with placebo in decreasing the incidence of IBC in women who had participated in the MORE trial [217]. The primary breast cancer analysis included 5213 patients (3996 who had completed MORE when CORE began and 1217 who were still participating in MORE when CORE began). The 4-year incidences in the raloxifene group of IBC and ER-positive IBC were reduced by 59 % and 66 %, respectively. Over the 8 years of both trials, the incidences of IBC and ER-positive IBC were reduced by 66 % (HR=0.34; 95 % CI, 0.22–0.50) and 76 % (HR=0.24; 95 % CI, 0.15–0.40), respectively, in patients who received raloxifene.

#### The Study of Tamoxifen and Raloxifene (STAR) Trial (NSABP-P2)

This double-blind, randomized controlled trial included 19,747 postmenopausal women aged 35 years and older with an increased risk of breast cancer [218], which was defined as a personal history of LCIS or a 5-year predicted risk for IBC of at least 1.66 % as determined by the Gail model. Women with a history of cerebral

vascular accidents, transient ischemic attack, pulmonary embolism, deep venous thrombosis, uncontrolled diabetes, uncontrolled hypertension, or atrial fibrillation were excluded from the study. Women were randomly assigned to receive 20 mg of tamoxifen per day plus a placebo or 60 mg of raloxifene per day plus a placebo for a 5-year period. The primary end point was the development of biopsy-proven IBC. The secondary end points of the trial included the incidence of noninvasive breast cancer, uterine cancer, cardiovascular events, stroke, transient ischemic attack, pulmonary embolism, deep venous thrombosis, osteoporotic fractures, cataracts, life, and death from any cause. There was no difference between the effects of tamoxifen and raloxifene on the incidence of breast cancer. There were 163 cases of IBC in the women assigned to the tamoxifen group, compared with 168 cases in the raloxifene group. The rate per 1000 woman-years was 4.3 in the tamoxifen group and 4.4 in the raloxifene group (RR=1.02; 95 % CI, 0.82–1.28). The pathological characteristics of the tumors did not differ between the treatment groups with respect to the distribution by tumor size, nodal status, or ER level. The incidence of noninvasive breast cancer was lower in the tamoxifen group (1.51 per 1000 women) compared with the raloxifene group (2.11 per 1000 women); however, this difference did not reach significance. There were fewer cases of uterine malignancies in the raloxifene group (23 cases) compared with the tamoxifen group (36 cases), although this difference was also not significant. Similarly, no significant differences between the two groups were observed regarding the incidence of stroke, transient ischemic attack, and osteoporotic fractures at the hip, spine, and radius. However, a 30 % decrease in the incidence of pulmonary embolism and deep venous thrombosis was noted in the raloxifene arm (100 versus 141 events in the raloxifene versus tamoxifen groups, respectively; RR=0.70; 95 % CI, 0.54–0.91). Fewer women who received raloxifene developed cataracts (RR=0.79; 95 % CI, 0.68–0.92). Mortality was similar in the two groups (101 deaths in the tamoxifen group versus 96 in the raloxifene group; RR=0.94; 95 %

CI, 0.71–1.26). Based on the data from STAR and other raloxifene trials, the FDA approved raloxifene for the prevention of IBC in postmenopausal women who are at increased risk of breast cancer or in postmenopausal women with osteoporosis.

An updated analysis of the STAR trial was performed in 2010 with a median follow-up time of 81 months [219]. There continued to be no significant difference in the incidence of IBC between tamoxifen and raloxifene (RR=1.24; 95 % CI, 1.05–1.47). There were 137 cases of noninvasive breast cancer in the raloxifene group and 111 cases in the tamoxifen group (RR=1.22; 95 % CI, 0.95–1.59); as such, the difference between the two groups was smaller than that in the original report. In contrast to the initial study, there was a significant decrease in the risk of endometrial cancer with raloxifene (RR=0.55; 95 % CI, 0.36–0.83). In addition, significant reductions in the incidence of thromboembolic events (RR=0.75; 95 % CI, 0.60–0.93) and uterine hyperplasia (RR=0.19; 95 % CI, 0.12–10.29) were reported. No significant mortality differences between raloxifene and tamoxifen were noted.

#### **Additional SERMS**

The Postmenopausal Evaluation and Risk Reduction with Lasofoxifene (PEARL) study randomly assigned 8556 postmenopausal women with osteoporosis to receive a placebo or either 0.25 mg or 0.5 mg of lasofoxifene per day [220]. A significant reduction in the incidence of ER-positive breast cancer (HR=0.19; 95 % CI, 0.07–0.56) was reported in women assigned to 0.5 mg of lasofoxifene per day. In addition, the incidences of vertebral and non-vertebral fractures, coronary heart disease events, and stroke were also reduced in this group. A smaller effect on the incidence of ER-positive IBC was noted with 0.25 mg of lasofoxifene per day.

The investigational SERM arzoxifene has also been evaluated in postmenopausal women with breast cancer. The GENERATIONS trial was a large, multicenter, double-blind, placebo-controlled study that compared daily dosing of 20 mg of arzoxifene to placebo in



9354 postmenopausal women with osteoporosis or low bone mass [221, 222]. The median follow-up was 48 months. The incidence of IBC was decreased in women assigned to the arzoxifene group (22 cases versus 53 in the placebo group; HR=0.41; 95 % CI, 0.25–0.68). This reduction was primarily observed in women with ER-positive breast cancer, similar to the results for other SERMs.

### Role of Aromatase Inhibitors

High aromatase levels in breast tissues and high circulatory estrogen levels are known risk factors for IBC [223]. Anastrozole, letrozole, and exemestane decrease the circulating estrogen levels in postmenopausal women by inhibiting the enzyme aromatase, which catalyzes the conversion of androgens to estrogens. The role of AIs in the adjuvant treatment of postmenopausal women with receptor-positive IBC is well established [224]. A 37–55 % reduction in the incidence of CBC has been reported with the use of AIs in clinical trials [225–227]. The main side effects of AIs include arthralgia and accelerated bone resorption, and its overall safety profile is relatively more favorable compared with tamoxifen.

#### The NCIC CTG MAP.3 Trial

The NCIC CTG MAP.3 trial was a prospective trial that investigated the role of exemestane in reducing the incidence of IBC in postmenopausal women who were at increased risk [228]. This double-blind trial randomized 4560 postmenopausal women who had at least one risk factor (age >60 years, Gail 5-year risk score >1.66 %, prior atypical ductal or lobular hyperplasia or LCIS, or DCIS with mastectomy) to receive either 25 mg of exemestane per day or placebo. The median age of women who participated in the trial was 62.5 years, and the median Gail risk score was 2.3 %. The investigators reported a reduction in the incidence of IBC in women assigned to the exemestane group (11 cases) compared with those in the placebo group (32 cases) at a median follow-up of 35 months. A 65 % relative reduction in the annual incidence of IBC (0.19 % versus 0.55 %; HR=0.35; 95 % CI, 0.18–0.70;  $P=0.002$ ) with exemestane was

reported. The NNT to prevent one case of IBC with exemestane therapy was 94 in 3 years. The annual incidence of IBC plus DCIS (20 in the exemestane group and 44 in the placebo group) was 0.35 % and 0.77 % in the exemestane and placebo groups, respectively (HR=0.47; 95 % CI, 0.27–0.79). Adverse events were experienced by 88 % of women in the exemestane group and 85 % in the placebo group; hot flashes and arthritis were the most common adverse events in both groups. There were no significant differences between the two groups regarding secondary end points, such as new osteoporosis, skeletal fractures, cardiovascular events, and cancers other than IBC. No treatment-related deaths were reported. Women taking exemestane reported slightly worse menopause-related quality-of-life events compared with placebo (7 % more overall).

#### IBIS-II Trial

IBIS-II is a multicenter, randomized, double-blind, placebo-controlled phase III trial that evaluated the AI anastrozole in postmenopausal women at high risk for breast cancer (family history, atypical hyperplasia or LCIS, nulliparity or age  $\geq 30$  at first birth, mammographic opacity covering at least 50 % of the breast) [229]. Anastrozole (1 mg/day) was associated with a 53 % reduction in the incidence of IBC and DCIS (primary end point) compared with placebo after a median follow-up of 5 years (HR=0.47; 95 % CI, 0.32–68.0;  $P<0.0001$ ). Similar to most chemoprevention trials, the protective effect of anastrozole was observed in ER-positive IBC, with no significant effect in the ER-negative subgroup. The total mortality was 0.9 % for both arms. Interestingly, a reduction in the incidence of skin, gastrointestinal, and gynecological cancers as well as other cancers was noted in the anastrozole group (2 % versus 4 % in the placebo group; RR=0.58; 95 % CI, 0.39–0.85). A significant increase in the incidence of musculoskeletal events, such as aches and pain, vasomotor symptoms, dryness of the eyes, and hypertension, was observed in the anastrozole arm. Bone fractures occurred in 7.7 % of those on placebo compared with 8.5 % of women receiving anastrozole.

Based on the results of this trial, anastrozole may be an effective chemopreventive option for postmenopausal women.

Recently, a meta-analysis based on individual participant data from nine randomized prevention trials using tamoxifen, raloxifene, arzoxifene, and lasofoxifene was reported [230]. These trials included the Royal Marsden Hospital Tamoxifen Trial, IBIS-I, NSABP-P1, Italian Tamoxifen Prevention Study, MORE/CORE, RUTH, STAR, PEARL, and GENERATIONS. The median follow-up time was 65 months. Overall, a 38 % reduction in the incidence of breast cancer (including DCIS) was noted (HR=0.62; 95 % CI, 0.56–0.69), with the largest reduction in the first 5 years of follow-up compared with years 5–10. The estimated 10-year cumulative incidence was 6.3 % in the control group and 4.2 % in the SERM group. A total of 42 women would need to be treated to prevent one breast cancer event in the first 10 years of follow-up. A significant overall reduction of 31 % in the incidence of DCIS was reported, with a 38 % reduction in the tamoxifen trials but no effect for raloxifene.

The investigators noted a significant reduction in all breast cancers and ER-positive breast cancers with 0.5 mg of lasofoxifene per day compared with placebo; however, there was a nonsignificant increase in the incidence of ER-negative IBC (HR=1.43; 95 % CI, 0.43–1.66) and a nonsignificant decrease for DCIS (HR=0.76; 95 % CI, 0.26–2.21) with lasofoxifene (both 0.5 mg and 0.25 mg per day). Similarly, arzoxifene decreased overall IBC and ER-positive breast cancer incidences by 58 % and 70 %, respectively. No effect was noted on ER-negative breast cancers, while there was a small reduction in DCIS (HR=0.30; 95 % CI, 0.08–1.09). Overall, a higher rate of endometrial cancer was noted in women receiving a SERM compared with placebo (HR=1.56; 95 % CI, 1.13–2.14;  $P=0.007$ ). This increase was limited to the first 5 years of follow-up and primarily to the tamoxifen trials. No increase in the incidence of endometrial cancer was observed in the raloxifene trials. An increased risk was also observed with arzoxifene (HR=2.26; 95 % CI, 0.70–7.32;  $P=0.2$ ).

An overall increase in the incidence of venous thromboembolic events was noted, with both

tamoxifen and raloxifene demonstrating a similar risk (OR=1.60; 1.21–2.12;  $P=0.001$  versus OR=1.45; 1.18–1.76;  $P<0.0001$ ). The rate was higher for arzoxifene and lasofoxifene. Overall, no side effects of myocardial infarction, stroke, or transient ischemic attack were noted for SERMs. The authors reported a 34 % reduction in vertebral fractures and a smaller reduction of non-vertebral fractures.

### **American Society of Clinical Oncology (ASCO) Clinical Practice Guidelines**

In July 2013, ASCO updated its clinical practice guidelines for the use of pharmacological agents to reduce the incidence of breast cancer [160]. The recommendations included a discussion of the use of tamoxifen (20 mg per day) in women (35 years or older) who are at increased risk of breast cancer. In postmenopausal women, raloxifene (60 mg per day for 5 years) and exemestane (25 mg per day for 5 years) may be an alternative to tamoxifen. An increased risk of breast cancer was defined as a 5-year projected absolute risk of breast cancer of 1.66 % (using the National Institute of Cancer Breast Cancer Risk Assessment Tool [122] or an equivalent measure) or women with LCIS. The use of tamoxifen or raloxifene was not recommended for women with a history of deep venous thrombosis, pulmonary embolism, stroke, or transient ischemic attack; during prolonged immobilizations; in women who are pregnant or may become pregnant; or in nursing mothers. Discussions with patients and healthcare providers should include the risks and benefits of the agents under consideration.

Currently, there are no data from phase III randomized trials on the protective effect of raloxifene and AIs in BRCA1/BRCA2 mutation carriers. Anastrozole is currently being studied as a prevention agent in a large phase III trial, and IGFBP-1 is being evaluated as a surrogate end point biomarker in prospective breast chemoprevention studies [231]. There are limited data on the effectiveness of tamoxifen for the reduction of breast cancer risk in BRCA1/BRCA2 mutation carriers. In the NSABP-P1, 19 of the 288 women who developed breast cancer had

BRCA1/BRCA2 mutations. A significant effect on breast cancer risk was not observed with tamoxifen in women with BRCA1 (RR=1.67; 95 % CI, 0.32–10.70) or BRCA2 (RR=0.38; 95 % CI, 0.06–1.56) mutations [232].

Interest is now focused on developing agents with a broader spectrum of preventive activity, particularly with regard to ER-negative breast cancer subtypes. A number of phase I and II trials using tissue-derived surrogate end point biomarkers (SEBs) as outcomes have been implemented. These smaller trials address prevention not only of ER-negative but also ER-positive breast cancers because approximately 50 % of the latter are resistant to the estrogen-targeting drugs used in the large trials [233].

### The Role of Diet and Nutrition

The association between various dietary factors and the risk of breast cancer has been controversial due to the lack of randomized prospective studies. An international panel of the World Cancer Research Fund and the American Institute for Cancer Research concluded that alcohol intake increases the risk of breast cancer for all age groups [234]. Some of the mechanisms postulated include carcinogenic metabolites of alcohol, such as acetaldehyde or oxygen radicals, interference with folate or estrogen metabolism, and several nutrient deficiencies associated with alcohol intake [235]. Some studies have demonstrated a 10 % increase in the risk of breast cancer for every 10 g of alcohol consumed per day [236, 237]. Interestingly, the excess risk due to alcohol consumption may be reduced or mitigated by adequate folate consumption [11, 238, 239]. In addition, the role of dietary fat as a possible risk factor for IBC has been considerably investigated, and a nonsignificant increase in the rate of breast cancer (6–11 %) was reported in women who consume excess dietary fat [234]. In the Women's Health Initiative Randomized Controlled Dietary Modification trial, a nonsignificant decrease in breast cancer risk was noted (RR=0.91; 95 % CI, 0.83–1.02) in women with a reduced intake of animal fat [240]. Similarly, a large prospective study demonstrated a small increase in the risk of IBC with increased intake

of dietary fat [234]. Red meat intake has also been linked to breast cancer risk. A modest association between the two was reported in a meta-analysis of case-control and cohort studies; however, this association was not observed in a pooled analysis of prospective studies [241, 242]. An increased breast cancer risk was observed among women with high red meat intake in the UK Women's Cohort Study (12 % increase risk per 50-g increment of meat per day) [243]. The influence of BMI on the risk of breast cancer has also been well characterized. Women with a higher BMI are at a lower risk of breast cancer before menopause but have an increased risk in the postmenopausal stage [234]. The prospective Nurses' Health Study II, in which 116,000 women have been followed since 1989, has prespecified objectives to assess the role of risk factors such as dietary fiber, saturated and unsaturated fat, plasma levels of insulin-like growth factor, low-dose oral contraceptive pills, breast-feeding, and physical activity among younger nurses [244]. In summary, there is currently no conclusive evidence based on randomized controlled trials that a specific dietary intervention or weight loss will decrease the risk of developing IBC.

### References

1. Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin.* 2014;64:9–29.
2. Key TJ, Verkasalo PK, Banks E. Epidemiology of breast cancer. *Lancet Oncol.* 2001;2:133–40.
3. Global Cancer Facts and Figures. [www.cancer.org](http://www.cancer.org).
4. Ziegler RG, Hoover RN, Pike MC, Hildesheim A, Nomura AM, West DW, et al. Migration patterns and breast cancer risk in Asian-American women. *J Natl Cancer Inst.* 1993;85:1819–27.
5. King SE, Schottenfeld D. The “epidemic” of breast cancer in the U.S. – determining the factors. *Oncology.* 1996;10:453–62; discussion 62, 64, 70–2.
6. Cancer Facts and Figures ACS.
7. Breast Cancer Statistics. <http://seer.cancer.gov/statfacts/html/breast.html>.
8. Percent of New Breast Cancer Cases by Age Group (Surveillance, Epidemiology, and End Results [SEER], 2011). <http://seer.cancer.gov/statfacts/html/breast.html>. Accessed 26 June 2015.
9. Incidence Rates of Female Breast Cancer by Race and Ethnicity, U.S., 1999–2011. <http://www.cdc.gov/cancer/dcp/c/data/race.html>. Accessed 26 June 2015.

10. Death Rates by Race and Ethnicity, U.S., 1999–2011. <http://www.cdc.gov/cancer/dcp/data/race.html>. Accessed 26 June 2015.
11. Zhang SM, Lee IM, Manson JE, Cook NR, Willett WC, Buring JE. Alcohol consumption and breast cancer risk in the Women's Health Study. *Am J Epidemiol*. 2007;165:667–76.
12. Tretli S. Height and weight in relation to breast cancer morbidity and mortality. A prospective study of 570,000 women in Norway. *Int J Cancer*. 1989;44:23–30.
13. Nelson HD, Humphrey LL, Nygren P, Teutsch SM, Allan JD. Postmenopausal hormone replacement therapy: scientific review. *JAMA*. 2002;288:872–81.
14. Boice Jr JD, Preston D, Davis FG, Monson RR. Frequent chest X-ray fluoroscopy and breast cancer incidence among tuberculosis patients in Massachusetts. *Radiat Res*. 1991;125:214–22.
15. Clemons M, Loijens L, Goss P. Breast cancer risk following irradiation for Hodgkin's disease. *Cancer Treat Rev*. 2000;26:291–302.
16. Brinton LA, Schairer C, Hoover RN, Fraumeni Jr JF. Menstrual factors and risk of breast cancer. *Cancer Invest*. 1988;6:245–54.
17. Trichopoulos D, MacMahon B, Cole P. Menopause and breast cancer risk. *J Natl Cancer Inst*. 1972;48:605–13.
18. Brinton LA, Hoover R, Fraumeni Jr JF. Reproductive factors in the aetiology of breast cancer. *Br J Cancer*. 1983;47:757–62.
19. Brinton LA, Hoover R, Fraumeni Jr JF. Epidemiology of minimal breast cancer. *JAMA*. 1983;249:483–7.
20. Ries LAG EM, Kosary CL, et al., editors. SEER cancer statistics, review. Bethesda: National Cancer Institute. NIH Pub. No. 00–, 2789. 2000.
21. Hankey BF, Curtis RE, Naughton MD, Boice Jr JD, Flannery JT. A retrospective cohort analysis of second breast cancer risk for primary breast cancer patients with an assessment of the effect of radiation therapy. *J Natl Cancer Inst*. 1983;70:797–804.
22. Page DL, Dupont WD, Rogers LW, Landenberger M. Intraductal carcinoma of the breast: follow-up after biopsy only. *Cancer*. 1982;49:751–8.
23. Dupont WD, Parl FF, Hartmann WH, Brinton LA, Winfield AC, Worrell JA, et al. Breast cancer risk associated with proliferative breast disease and atypical hyperplasia. *Cancer*. 1993;71:1258–65.
24. Fabian CJ, Kimler BF, Zalles CM, et al. Short-term breast cancer prediction by random periareolar fine-needle aspiration cytology and the Gail risk model. *J Natl Cancer Inst*. 2000;92:1217–27.
25. Pharoah PD, Day NE, Duffy S, Easton DF, Ponder BA. Family history and the risk of breast cancer: a systematic review and meta-analysis. *Int J Cancer*. 1997;71:800–9.
26. Easton DF, Ford D, Bishop DT. Breast and ovarian cancer incidence in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. *Am J Hum Genet*. 1995;56:265–71.
27. Singletary SE. Rating the risk factors for breast cancer. *Ann Surg*. 2003;237(4):474–82.
28. Litton JK, Eralp Y, Gonzalez-Angulo AM, Broglio K, Uyei A, Hortobagyi GN, et al. Multifocal breast cancer in women <or =35 years old. *Cancer*. 2007;110:1445–50.
29. Inskip PD, Curtis RE. New malignancies following childhood cancer in the United States, 1973–2002. *Int J Cancer*. 2007;121:2233–40.
30. Buist DS, Abraham LA, Barlow WE, Krishnaraj A, Holdridge RC, Sickles EA, et al. Diagnosis of second breast cancer events after initial diagnosis of early stage breast cancer. *Breast Cancer Res Treat*. 2010;124:863–73.
31. Cote ML, Ruterbusch JJ, Alosch B, Bandyopadhyay S, Kim E, Albashiti B, et al. Benign breast disease and the risk of subsequent breast cancer in African American women. *Cancer Prev Res*. 2012;5:1375–80.
32. Hartmann LC, Sellers TA, Frost MH. Benign breast disease and the risk of breast cancer. *N Engl J Med*. 2005;353:229–37.
33. Colditz GA, Kaphingst KA, Hankinson SE, Rosner B. Family history and risk of breast cancer: nurses' health study. *Breast Cancer Res Treat*. 2012;133:1097–104.
34. Collaborative Group on Hormonal Factors in Breast Cancer. Familial breast cancer: collaborative reanalysis of individual data from 52 epidemiological studies including 58 wwbca, 986 women without the disease. *Lancet*. 2001;358:1389–99.
35. Hsieh CC, Trichopoulos D, Katsouyanni K, Yuasa S. Age at menarche, age at menopause, height and obesity as risk factors for breast cancer: associations and interactions in an international case-control study. *Int J Cancer*. 1990;46:796–800.
36. Ritte R, Lukanova A, Berrino F. Adiposity, hormone replacement therapy use and breast cancer risk by age and hormone receptor status: a large prospective cohort study. *Breast Cancer Res*. 2012;14:R76.
37. Bayraktar S, Amendola L, Gutierrez-Barrera AM, Hashmi SS, Amos C, Gambello M, et al. Clinicopathologic characteristics of breast cancer in BRCA-carriers and non-carriers in women 35 years of age or less. *Breast*. 2014;23:770–4.
38. Rosner B, Colditz GA, Willett WC. Reproductive risk factors in a prospective study of breast cancer: the Nurses' Health Study. *Am J Epidemiol*. 1994;139:819–35.
39. Collaborative Group on Hormonal Factors in Breast Cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet*. 2002;360:187–95.
40. Sieri S, Krogh V, Bolelli G. Sex hormone levels, breast cancer risk, and cancer receptor status in postmenopausal women: the ORDET cohort. *Cancer Epidemiol Biomarkers Prev*. 2009;18:169–76.

41. Kelsey JL, Gammon MD, John EM. Reproductive factors and breast cancer. *Epidemiol Rev.* 1993;15:36–47.
42. Lahmann PH, Hoffmann K, Allen N. Body size and breast cancer risk: findings from the European Prospective Investigation into Cancer And Nutrition (EPIC). *Int J Cancer.* 2004;111:762–71.
43. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52 wwbca, 411 women without breast cancer. Collaborative Group on Hormonal Factors in Breast Cancer. *Lancet.* 1997;350:1047–59.
44. Anderson GL, Manson J, Wallace R, Lund B, Hall D, Davis S, et al. Implementation of the Women's Health Initiative study design. *Ann Epidemiol.* 2003;13:S5–17.
45. Colditz GA, Rosner B. Cumulative risk of breast cancer to age 70 years according to risk factor status: data from the Nurses' Health Study. *Am J Epidemiol.* 2000;152:950–64.
46. Chlebowski RT, Manson JE, Anderson GL, Cauley JA, Aragaki AK, Stefanick ML, et al. Estrogen plus progestin and breast cancer incidence and mortality in the Women's Health Initiative Observational Study. *J Natl Cancer Inst.* 2013;105:526–35.
47. Danaei G, Vander Hoorn S, Lopez AD, Murray CJ, Ezzati M, Comparative Risk Assessment Collaborating Group. Causes of cancer in the world: comparative risk assessment of nine behavioural and environmental risk factors. *Lancet.* 2005;366:1784–93.
48. Chen WY, Rosner B, Hankinson SE, Colditz GA, Willett WC. Moderate alcohol consumption during adult life, drinking patterns, and breast cancer risk. *JAMA.* 2011;306:1884–90.
49. Wu Y, Zhang D, Kang S. Physical activity and risk of breast cancer: a meta-analysis of prospective studies. *Breast Cancer Res Treat.* 2013;137:869–82.
50. Milazzo G, Giorgino F, Damante G, Sung C, Stampfer MR, Vigneri R, et al. Insulin receptor expression and function in human breast cancer cell lines. *Cancer Res.* 1992;52:3924–30.
51. Gunter MJ, Hoover DR, Yu H. Insulin, insulin-like growth factor-I, and risk of breast cancer in postmenopausal women. *J Natl Cancer Inst.* 2009;101:48–60.
52. Henderson TO, Amsterdam A, Bhatia S. Systematic review: surveillance for breast cancer in women treated with chest radiation for childhood, adolescent, or young adult cancer. *Ann Intern Med.* 2010;152:444–55; W144–54.
53. Guibout C, Adjadj E, Rubino C, Shamsaldin A, Grimaud E, Hawkins M, et al. Malignant breast tumors after radiotherapy for a first cancer during childhood. *J Clin Oncol.* 2005;23:197–204.
54. Preston DL, Cullings H, Suyama A. Solid cancer incidence in atomic bomb survivors exposed in utero or as young children. *J Natl Cancer Inst.* 2008;100:428–36.
55. Pukkala E, Kesminiene A, Poliakov S. Breast cancer in Belarus and Ukraine after the Chernobyl accident. *Int J Cancer.* 2006;119:651–8.
56. Barlow WE, White E, Ballard-Barbash R. Prospective breast cancer risk prediction model for women undergoing screening mammography. *J Natl Cancer Inst.* 2006;98:1204–14.
57. Boyd NF, Guo H, Martin LJ, Sun L, Stone J, Fishell E, et al. Mammographic density and the risk and detection of breast cancer. *N Engl J Med.* 2007;356:227–36.
58. McCormack VA, dos Santos Silva I. Breast density and parenchymal patterns as markers of breast cancer risk: a meta-analysis. *Cancer Epidemiol Biomarkers Prev.* 2006;15:1159–69.
59. Pettersson A, Hankinson SE, Willett WC, Lagiou P, Trichopoulos D, Tamimi RM. Nondense mammographic area and risk of breast cancer. *Breast Cancer Res.* 2011;13:R100.
60. Rauh C, Hack CC, Häberle L, Hein A, Engel A, Schrauder MG, et al. Percent mammographic density and dense area as risk factors for breast cancer. *Geburtshilfe Frauenheilkd.* 2012;72:727–33.
61. Lynch HT, Lynch JF. Breast cancer genetics in an oncology clinic: 328 consecutive patients. *Cancer Genet Cytogenet.* 1986;22:369–71.
62. Margolin S, Johansson H, Rutqvist LE, Lindblom A, Fornander T. Family history, and impact on clinical presentation and prognosis, in a population-based breast cancer cohort from the Stockholm County. *Fam Cancer.* 2006;5:309–21.
63. Lalloo F, Evans DG. Familial breast cancer. *Clin Genet.* 2012;82:105–14.
64. Sharif S, Moran A, Huson SM, Iddenden R, Shenton A, Howard E, et al. Women with neurofibromatosis 1 are at a moderately increased risk of developing breast cancer and should be considered for early screening. *J Med Genet.* 2007;44:481–4.
65. Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, et al. Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. The Breast Cancer Linkage Consortium. *Am J Hum Genet.* 1998;62:676–89.
66. Antoniou A, Pharoah PD, Narod S. Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case Series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet.* 2003;72:1117–30.
67. King MC, Marks JH, Mandell JB, New York Breast Cancer Study Group. Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science.* 2003;302:643–6.
68. Thompson D, Duedal S, Kirner J, McGuffog L, Last J, Reiman A, et al. Cancer risks and mortality in heterozygous ATM mutation carriers. *J Natl Cancer Inst.* 2005;97:813–22.
69. Seal S, Thompson D, Renwick A, Elliott A, Kelly P, Barfoot R, et al. Truncating mutations in the Fanconi



- anemia J gene BRIP1 are low-penetrance breast cancer susceptibility alleles. *Nat Genet.* 2006;38:1239–41.
70. Wong MW, Nordfors C, Mossman D. BRIP1, PALB2, and RAD51C mutation analysis reveals their relative importance as genetic susceptibility factors for breast cancer. *Breast Cancer Res Treat.* 2011;127:853–9.
71. Melhem-Bertrandt A, Bojadzieva J, Ready KJ, Obeid E, Liu DD, Gutierrez-Barrera AM, et al. Early onset HER2-positive breast cancer is associated with germline TP53 mutations. *Cancer.* 2012;118:908–13.
72. Roy R, Chun J, Powell SN. BRCA1 and BRCA2: different roles in a common pathway of genome protection. *Nat Rev Cancer.* 2012;12:68–78.
73. Yuan SS, Lee SY, Chen G, Song M, Tomlinson GE, Lee EY. BRCA2 is required for ionizing radiation-induced assembly of Rad51 complex in vivo. *Cancer Res.* 1999;59:3547–51.
74. Zhong Q, Chen CF, Li S. Association of BRCA1 with the hRad50-hMre11-p95 complex and the DNA damage response. *Science.* 1999;285:747–50.
75. Rajan JV, Wang M, Marquis ST, Chodosh LA. Brca2 is coordinately regulated with Brca1 during proliferation and differentiation in mammary epithelial cells. *Proc Natl Acad Sci U S A.* 1996;93:13078–83.
76. Wang Y, Cortez D, Yazdi P, Neff N, Elledge SJ, Qin J. BASC, a super complex of BRCA1-associated proteins involved in the recognition and repair of aberrant DNA structures. *Genes Dev.* 2000;14:927–39.
77. Narod SA, Foulkes WD. BRCA1 and BRCA2: 1994 and beyond. *Nat Rev Cancer.* 2004;4:665–76.
78. Collins N, McManus R, Wooster R. Consistent loss of the wild type allele in breast cancers from a family linked to the BRCA2 gene on chromosome 13q12-13. *Oncogene.* 1995;10:1673–5.
79. Cornelis RS, Neuhausen SL, Johansson O. High allele loss rates at 17q12-q21 in breast and ovarian tumors from BRCA1-linked families. The Breast Cancer Linkage Consortium. *Genes Chromosomes Cancer.* 1995;13:203–10.
80. Khoo US, Ozelik H, Cheung AN. Somatic mutations in the BRCA1 gene in Chinese sporadic breast and ovarian cancer. *Oncogene.* 1999;18:4643–6.
81. Esteller M, Silva JM, Dominguez G, Bonilla F, Matias-Guiu X, Lerma E, et al. Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors. *J Natl Cancer Inst.* 2000;92:564–9.
82. Tapia T, Smalley SV, Kohen P, Muñoz A, Solis LM, Corvalan A, et al. Promoter hypermethylation of BRCA1 correlates with absence of expression in hereditary breast cancer tumors. *Epigenetics.* 2008;3:157–63.
83. Dworkin AM, Spearman AD, Tseng SY, Sweet K, Toland AE. Methylation not a frequent “second hit” in tumors with germline BRCA mutations. *Fam Cancer.* 2009;8:339–46.
84. Collins N, Wooster R, Stratton MR. Absence of methylation of CpG dinucleotides within the promoter of the breast cancer susceptibility gene BRCA2 in normal tissues and in breast and ovarian cancers. *Br J Cancer.* 1997;76:1150–6.
85. Rich TA, Woodson AH, Litton J, Arun B. Hereditary breast cancer syndromes and genetic testing. *J Surg Oncol.* 2014.
86. Foulkes WD. Inherited susceptibility to common cancers. *N Engl J Med.* 2008;359:2143–53.
87. Honrado E, Benitez J, Palacios J. The molecular pathology of hereditary breast cancer: genetic testing and therapeutic implications. *Mod Pathol.* 2005;18:1305–20.
88. Gerdes AM, Cruger DG, Thomassen M, Kruse TA. Evaluation of two different models to predict BRCA1 and BRCA2 mutations in a cohort of Danish hereditary breast and/or ovarian cancer families. *Clin Genet.* 2006;69:171–8.
89. Melchor L, Benitez J. The complex genetic landscape of familial breast cancer. *Hum Genet.* 2013;132:845–63.
90. Thompson D, Easton D. The genetic epidemiology of breast cancer genes. *J Mammary Gland Biol Neoplasia.* 2004;9:221–36.
91. Diez O, Osorio A, Duran M. Analysis of BRCA1 and BRCA2 genes in Spanish breast/ovarian cancer patients: a high proportion of mutations unique to Spain and evidence of founder effects. *Hum Mutat.* 2003;22:301–12.
92. Mavaddat N, Barrowdale D, Andrulis IL. Pathology of breast and ovarian cancers among BRCA1 and BRCA2 mutation carriers: results from the Consortium of Investigators of Modifiers of BRCA1/2 (CIMBA). *Cancer Epidemiol Biomarkers Prev.* 2012;21:134–47.
93. Chen S, Parmigiani G. Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol.* 2007;25:1329–33.
94. Atchley DP, Albarracin CT, Lopez A, Valero V, Amos CI, Gonzalez-Angulo AM, et al. Clinical and pathologic characteristics of patients with BRCA-positive and BRCA-negative breast cancer. *J Clin Oncol.* 2008;26:4282–8.
95. Marcus JN, Watson P, Page DL, Narod SA, Lenoir GM, Tonin P, et al. Hereditary breast cancer: pathobiology, prognosis, and BRCA1 and BRCA2 gene linkage. *Cancer.* 1996;77:697–709.
96. Eisinger F, Jacquemier J, Charpin C, Stoppa-Lyonnet D, Bressac-de Paillerets B, Peyrat JP, et al. Mutations at BRCA1: the medullary breast carcinoma revisited. *Cancer Res.* 1998;58:1588–92.
97. Lakhani SR, Gusterson BA, Jacquemier J, Sloane JP, Anderson TJ, van de Vijver MJ, et al. The pathology of familial breast cancer: histological features of cancers in families not attributable to mutations in BRCA1 or BRCA2. *Clin Cancer Res.* 2000;6:782–9.
98. Lakhani SR, Gusterson BA, Jacquemier J, Sloane JP, Anderson TJ, van de Vijver MJ, et al. The pathology

- of familial breast cancer: predictive value of immunohistochemical markers estrogen receptor, progesterone receptor, HER-2, and p53 in patients with mutations in BRCA1 and BRCA2. *J Clin Oncol.* 2002;20:2310–8.
99. Arun B, Vogel KJ, Lopez A, Hernandez M, Atchley D, Broglio KR, et al. High prevalence of preinvasive lesions adjacent to BRCA1/2-associated breast cancers. *Cancer Prev Res.* 2009;2:122–7.
  100. Kennedy RD, Quinn JE, Johnston PH, Harkin DP. BRCA1: mechanisms of inactivation and implications for management of patients. *Lancet.* 2002;360:1007–14.
  101. Verhoog LC, Brekelmans CT, Seynaeve C, Dahmen G, van Geel AN, Bartels CC. Survival in hereditary breast cancer associated with germline mutations of BRCA2. *J Clin Oncol.* 1999;17:3396–402.
  102. Rennert G, Bisland-Naggan S, Barnett-Griness O, Bar-Joseph N, Zhang S, Rennert HS, et al. Clinical outcomes of breast cancer in carriers of BRCA1 and BRCA2 mutations. *N Engl J Med.* 2007;357:115–23.
  103. Brekelmans CT, Tilanus-Linthorst MM, Seynaeve C, vd OA, Menke-Pluymers MB, Bartels CC. Tumour characteristics, survival and prognostic factors of hereditary breast cancer from BRCA2-, B. *Eur J Cancer.* 2007;43:867–76.
  104. Kriege M, Seynaeve C, Meijers-Heijboer H, Collee JM, Menke-Pluymers MB, Bartels CC. Distant disease-free interval, site of first relapse and post-relapse survival in B. *Breast Cancer Res Treat.* 2008;111:303–11.
  105. Bonadona V, Dussart-Moser S, Voirin N, Sinilnikova OM, Mignotte H, Mathevet P, Bremond A, Treilleux I, Martin A, Rostaing P. Prognosis of early-onset breast cancer based on BRCA1/2 mutation status in a French population-based cohort and review. *Breast Cancer Res Treat.* 2007;101:233–45.
  106. Bordeleau L, Panchal S, Goodwin P. Prognosis of BRCA-associated breast cancer: a summary of evidence. *Breast Cancer Res Treat.* 2010;119:13–24.
  107. Robson M, Levin D, Federici M, Satagopan J, Bogolminy F, Heerdt A, Borgen P, McCormick B, Hudis C, Norton L, Boyd J, Offit K. Breast conservation therapy for invasive breast cancer in Ashkenazi women with BRCA gene founder mutations. *J Natl Cancer Inst.* 1999;91:2112–7.
  108. Chappuis PO, Kapusta L, Bégin LR, Wong N, Brunet JS, Narod SA, Slingerland J, Foulkes WD. Germline BRCA1/2 mutations and p27(Kip1) protein levels independently predict outcome after breast cancer. *J Clin Oncol.* 2000;18:4045–52.
  109. Stoppa-Lyonnet D, Ansquer Y, Dreyfus H, Gautier C, Gauthier-Villars M, Bourstyn E, Clough KB, Magdelenat H, et al. The Institute Curie Breast Cancer Group, Familial invasive breast cancer: worse outcome related to BRCA1 mutations. *J Clin Oncol.* 2000;18:4053–9.
  110. Moller P, Evans DG, Reis MM, Gregory H, Anderson E, Maehle L, et al. Surveillance for familial breast cancer: differences in outcome according to BRCA mutation status. *Int J Cancer.* 2007;121:1017–20.
  111. Lee EH, Park SK, Park B. Effect of BRCA1/2 mutation on short-term and long-term breast cancer survival: a systematic review and meta-analysis. *Breast Cancer Res Treat.* 2010;122:11–25.
  112. Lee LJ, Alexander B, Schnitt SJ, Comander A, Gallagher B, Garber JE, et al. Clinical outcome of triple negative breast cancer in BRCA1 mutation carriers and noncarriers. *Cancer.* 2011;117:3093–100.
  113. Bayraktar S, Gutierrez-Barrera AM, Liu D, Tasbas T, Akar U, Litton JK, et al. Outcome of triple-negative breast cancer in patients with or without deleterious BRCA mutations. *Breast Cancer Res Treat.* 2011;130:145–53.
  114. Riley BD, Culver JO, Skrzynia C. Essential elements of genetic cancer risk assessment, counseling, and testing: updated recommendations of the National Society of Genetic Counselors. *J Genet Couns.* 2012;21:151–61.
  115. Genetic/Familial High-Risk Assessment: Breast and Ovarian. NCCN Guidelines. 2015. [www.nccn.org](http://www.nccn.org).
  116. Kwon JS, Gutierrez-Barrera AM, Young D, Sun CC, Daniels MS, Lu KH, et al. Expanding the criteria for BRCA mutation testing in breast cancer survivors. *J Clin Oncol.* 2010;28:4214–20.
  117. Bayraktar S, Elsayegh N, Gutierrez Barrera AM, Lin H, Kuerer H, Tasbas T, et al. Predictive factors for BRCA1/BRCA2 mutations in women with ductal carcinoma in situ. *Cancer.* 2012;118:1515–22.
  118. Elsayegh N, Kuerer HM, Lin H, Gutierrez Barrera AM, Jackson M, Muse KI, et al. Predictors that influence contralateral prophylactic mastectomy election among women with ductal carcinoma in situ who were evaluated for BRCA genetic testing. *Ann Surg Oncol.* 2014;21:3466–72.
  119. Claus EB, Risch N, Thompson WD. Autosomal dominant inheritance of early-onset breast cancer. Implications for risk prediction. *Cancer.* 1994;73:643–51.
  120. Saslow D, Boetes C, Burke W. American Cancer Society guidelines for breast screening with MRI as an adjunct to mammography. *CA Cancer J Clin.* 2007;57:75–89.
  121. Hollingsworth AB, Stough RG. An alternative approach to selecting patients for high-risk screening with breast MRI. *Breast J.* 2014;20:192–7.
  122. Gail MH, Brinton LA, Byar DP. Projecting individualized probabilities of developing breast cancer for white females who are being examined annually. *J Natl Cancer Inst.* 1989;81:1879–86.
  123. Pastor-Barriuso R, Ascunce N, Ederra M. Recalibration of the Gail model for predicting invasive breast cancer risk in Spanish women: a population-based cohort study. *Breast Cancer Res Treat.* 2013;138:249–59.
  124. Tyrer J, Duffy SW, Cuzick J. A breast cancer prediction model incorporating familial and personal risk factors. *Stat Med.* 2004;23:1111–30.

125. Antoniou AC, Pharoah PP, Smith P, Easton DF. The BOADICEA model of genetic susceptibility to breast and ovarian cancer. *Br J Cancer*. 2004;91:1580–90.
126. Antoniou AC, Durocher F, Smith P, Simard J, Easton DF. BRCA1 and BRCA2 mutation predictions using the BOADICEA and BRCAPRO models and penetrance estimation in high-risk French-Canadian families. *Breast Cancer Res*. 2006;8:R3.
127. Antoniou AC, Cunningham AP, Peto J, Evans DG, Lalloo F, Narod SA, et al. The BOADICEA model of genetic susceptibility to breast and ovarian cancers: updates and extensions. *Br J Cancer*. 2008;98:1457–66.
128. Powell M, Jamshidian F, Cheyne K, Nititham J, Prebil LA, Eremam R. Assessing breast cancer risk models in Marin County, a population with high rates of delayed childbirth. *Clin Breast Cancer*. 2014;14:212–20 e1.
129. Laitman Y, Simeonov M, Keinan-Boker L, Liphshitz I, Friedman E. Breast cancer risk prediction accuracy in Jewish Israeli high-risk women using the BOADICEA and IBIS risk models. *Genet Res*. 2013;95:174–7.
130. Berry DA, Parmigiani G, Sanchez J, Schildkraut J, Winer E. Probability of carrying a mutation of breast-ovarian cancer gene BRCA1 based on family history. *J Natl Cancer Inst*. 1997;89:227–38.
131. Parmigiani G, Berry D, Aguilar O. Determining carrier probabilities for breast cancer-susceptibility genes BRCA1 and BRCA2. *Am J Hum Genet*. 1998;62:145–58.
132. Antoniou AC, Cunningham AP, Peto J, Evans DG, Lalloo F, Narod SA, et al. Predicting the likelihood of carrying a BRCA1 or BRCA2 mutation: validation of BOADICEA, BRCAPRO, IBIS, Myriad and the Manchester scoring system using data from UK genetics clinics. *J Med Genet*. 2008;45:425–31.
133. Berry DA, Iversen Jr ES, Gudbjartsson DF. BRCAPRO validation, sensitivity of genetic testing of BRCA1/BRCA2, and prevalence of other breast cancer susceptibility genes. *J Clin Oncol*. 2002;20:2701–12.
134. Vogel KJ, Atchley DP, Erlichman J, Broglio KR, Ready KJ, Valero V, et al. BRCA1 and BRCA2 genetic testing in Hispanic patients: mutation prevalence and evaluation of the BRCAPRO risk assessment model. *J Clin Oncol*. 2007;25:4635–41.
135. Daniels MS, Babb SA, King RH, Urbauer DL, Batte BA, Brandt AC, et al. Underestimation of risk of a BRCA1 or BRCA2 mutation in women with high-grade serous ovarian cancer by BRCAPRO: a multi-institution study. *J Clin Oncol*. 2014;32:1249–55.
136. Varesco L, Viassolo V, Viel A, Gismondi V, Radice P, Montagna M, et al. Performance of BOADICEA and BRCAPRO genetic models and of empirical criteria based on cancer family history for predicting BRCA mutation carrier probabilities: a retrospective study in a sample of Italian cancer genetics clinics. *Breast*. 2013;22:1130–5.
137. Fischer C, Kuchenbacker K, Engel C. Evaluating the performance of the breast cancer genetic risk models BOADICEA, IBIS, BRCAPRO and Claus for predicting BRCA1/2 mutation carrier probabilities: a study based on 7352 families from the German Hereditary Breast and Ovarian Cancer Consortium. *J Med Genet*. 2013;50:360–7.
138. Ready KJ, Vogel KJ, Atchley DP, Broglio KR, Solomon KK, Amos C, et al. Accuracy of the BRCAPRO model among women with bilateral breast cancer. *Cancer*. 2009;115:725–30.
139. Pfeiffer RM, Park Y, Kreimer AR. Risk prediction for breast, endometrial, and ovarian cancer in white women aged 50 y or older: derivation and validation from population-based cohort studies. *PLoS Med*. 2013;10:e1001492.
140. Warner E, Plewes DB, Hill KA, Causer PA, Zubovits JT, Jong RA, et al. Surveillance of BRCA1 and BRCA2 mutation carriers with magnetic resonance imaging, ultrasound, mammography, and clinical breast examination. *JAMA*. 2004;292:1317–25.
141. Metcalfe K, Lynch HT, Ghadirian P, Tung N, Olivetto I, Warner E, et al. Contralateral breast cancer in BRCA1 and BRCA2 mutation carriers. *J Clin Oncol*. 2004;22:2328–35.
142. van Sprundel TC, Schmidt MK, Rookus MA. Risk reduction of contralateral breast cancer and survival after contralateral prophylactic mastectomy in BRCA1 or BRCA2 mutation carriers. *Br J Cancer*. 2005;93:287–92.
143. Metcalfe KA, Lubinski J, Ghadirian P. Predictors of contralateral prophylactic mastectomy in women with a BRCA1 or BRCA2 mutation: the Hereditary Breast Cancer Clinical Study Group. *J Clin Oncol*. 2008;26:1093–7.
144. Metcalfe K, Lynch HT, Ghadirian P, Tung N, Kim-Sing C, Olopade OI, et al. Risk of ipsilateral breast cancer in BRCA1 and BRCA2 mutation carriers. *Breast Cancer Res Treat*. 2011;127:287–96.
145. Graeser MK, Engel C, Rhiem K, Gadzicki D, Bick U, Kast K, et al. Contralateral breast cancer risk in BRCA1 and BRCA2 mutation carriers. *J Clin Oncol*. 2009;27:5887–92.
146. Malone KE, Begg CB, Haile RW, Borg A, Concannon P, Telhede L, et al. Population-based study of the risk of second primary contralateral breast cancer associated with carrying a mutation in BRCA1 or BRCA2. *J Clin Oncol*. 2010;28:2404–10.
147. Reding KW, Bernstein JL, Langholz BM, Bernstein L, Haile RW, Begg CB, et al. Adjuvant systemic therapy for breast cancer in BRCA1/BRCA2 mutation carriers in a population-based study of risk of contralateral breast cancer. *Breast Cancer Res Treat*. 2010;123:491–8.
148. Narod SA, Brunet JS, Ghadirian P. Tamoxifen and risk of contralateral breast cancer in BRCA1 and BRCA2 mutation carriers: a case-control study. *Hereditary Breast Cancer Clinical Study Group. Lancet*. 2000;356:1876–81.

149. Robson M, Svahn T, McCormick B. Appropriateness of breast-conserving treatment of breast carcinoma in women with germline mutations in BRCA1 or BRCA2: a clinic-based series. *Cancer*. 2005;103:44–51.
150. Pierce LJ, Levin AM, Rebbeck TR. Ten-year multi-institutional results of breast-conserving surgery and radiotherapy in BRCA1/2-associated stage I/II breast cancer. *J Clin Oncol*. 2006;24:2437–43.
151. Gronwald J, Tung N, Foulkes WD, Offit K, Gershoni R, Daly M, et al. Tamoxifen and contralateral breast cancer in BRCA1 and BRCA2 carriers: an update. *Int J Cancer*. 2006;118:2281–4.
152. Robson ME, Chappuis PO, Satagopan J, Wong N, Boyd J, Goffin JR, et al. A combined analysis of outcome following breast cancer: differences in survival based on BRCA1/BRCA2 mutation status and administration of adjuvant treatment. *Breast Cancer Res*. 2004;6:R8–17.
153. Domchek SM, Friebel TM, Singer CF, Evans DG, Lynch HT, Isaacs C, et al. Association of risk-reducing surgery in BRCA1 or BRCA2 mutation carriers with cancer risk and mortality. *JAMA*. 2010;304:967–75.
154. Struewing JP, Hartge P, Wacholder S. The risk of cancer associated with specific mutations of BRCA1 and BRCA2 among Ashkenazi Jews. *N Engl J Med*. 1997;336:1401–8.
155. Satagopan JM, Boyd J, Kauff ND, Robson M, Scheuer L, Narod S, et al. Ovarian cancer risk in Ashkenazi Jewish carriers of BRCA1 and BRCA2 mutations. *Clin Cancer Res*. 2002;8:3776–81.
156. Rebbeck TR, Lynch HT, Neuhausen SL, Narod SA, Van't Veer L, Garber JE, et al. Prophylactic oophorectomy in carriers of BRCA1 or BRCA2 mutations. *N Engl J Med*. 2002;346:1616–22.
157. Finch A, Beiner M, Lubinski J, Lynch HT, Moller P, Rosen B, et al. Salpingo-oophorectomy and the risk of ovarian, fallopian tube, and peritoneal cancers in women with a BRCA1 or BRCA2 Mutation. *JAMA*. 2006;296:185–92.
158. Kramer JVI, Chen BS. Prophylactic oophorectomy reduces breast cancer penetrance during prospective, long-term follow-up of BRCA1 mutation carriers. *J Clin Oncol*. 2005;23:8629–35.
159. Rebbeck TR, Kauff ND, Domchek SM. Meta-analysis of risk reduction estimates associated with risk-reducing salpingo-oophorectomy in BRCA1 or BRCA2 mutation carriers. *J Natl Cancer Inst*. 2009;101:80–7.
160. Visvanathan K, Hurley P, Bantug E. Use of pharmacologic interventions for breast cancer risk reduction: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol*. 2013;31(23):2942–62.
161. Eisen A, Lubinski J, Klijn J. Breast cancer risk following bilateral oophorectomy in BRCA1 and BRCA2 mutation carriers: an international case-control study. *J Clin Oncol*. 2005;23:7491–6.
162. Kurian AW, Sigal BM, Plevritis SK. Survival analysis of cancer risk reduction strategies for BRCA1/2 mutation carriers. *J Clin Oncol*. 2010;28:222–31.
163. Finch A, Narod SA. Quality of life and health status after prophylactic salpingo-oophorectomy in women who carry a BRCA mutation: a review. *Maturitas*. 2011;70:261–5.
164. Heemskerk-Gerritsen BA, Rookus MA, Aalfs CM, Ausems MG, Collée JM, Jansen L, et al. Improved overall survival after contralateral risk-reducing mastectomy in BRCA1/2 mutation carriers with a history of unilateral breast cancer: a prospective analysis. *Int J Cancer*. 2015;136(3):668–77.
165. Ready K, Gutierrez-Barrera AM, Amos C, et al. Cancer risk management decisions of women with BRCA1 or BRCA2 variants of uncertain significance. *Breast J*. 2011;17:210–2.
166. Byrski T, Gronwald J, Huzarski T, Grzybowska E, Budryk M, Stawicka M, et al. Response to neoadjuvant therapy with cisplatin in BRCA1-positive breast cancer patients. *Breast Cancer Res Treat*. 2009;115:359–63.
167. Byrski T, Gronwald J, Huzarski T, Grzybowska E, Budryk M, Stawicka M, et al. Pathologic complete response rates in young women with BRCA1-positive breast cancers after neoadjuvant chemotherapy. *J Clin Oncol*. 2010;28:375–9.
168. Arun B, Bayraktar S, Liu DD, Gutierrez Barrera AM, Atchley D, Puszta L, et al. Response to neoadjuvant systemic therapy for breast cancer in BRCA mutation carriers and noncarriers: a single-institution experience. *J Clin Oncol*. 2011;29:3739–46.
169. Silver DP, Richardson AL, Eklund AC, Wang ZC, Szallasi Z, Li Q, Juul N, et al. Efficacy of neoadjuvant Cisplatin in triple-negative breast cancer. *J Clin Oncol*. 2010;28:1145–53.
170. Wen J, Li R, Lu Y, Shupnik MA. Decreased BRCA1 confers tamoxifen resistance in breast cancer cells by altering estrogen receptor-coregulator interactions. *Oncogene*. 2009;28:575–86.
171. Fong PC, Boss DS, Yap TA. Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med*. 2009;361:123–34.
172. Iglehart JD, Silver DP. Synthetic lethality – a new direction in cancer-drug development. *N Engl J Med*. 2009;361:189–91.
173. Carey LA, Sharpless NE. PARP and cancer – if it's broke, don't fix it. *N Engl J Med*. 2011;364:277–9.
174. Kennedy RD, Quinn JE, Mullan PB, Johnston PG, Harkin DP. The role of BRCA1 in the cellular response to chemotherapy. *J Natl Cancer Inst*. 2004;96:1659–68.
175. Tassone P, Tagliaferri P, Perricelli A, Blotta S, Quaresima B, Martelli ML, et al. BRCA1 expression modulates chemosensitivity of BRCA1-defective HCC1937 human breast cancer cells. *Br J Cancer*. 2003;88:1285–91.
176. Chabaliat C, Lamare C, Racca C, Privat M, Valette A, Larminat F. BRCA1 downregulation leads to

- premature inactivation of spindle checkpoint and confers paclitaxel resistance. *Cell Cycle*. 2006;5:1001–7.
177. Kriege M, Jager A, Hoening MJ, Huijskens E, Blom J, van Deurzen CH, et al. The efficacy of taxane chemotherapy for metastatic breast cancer in BRCA1 and BRCA2 mutation carriers. *Cancer*. 2012;118:899–907.
178. Gholam D, Chebib A, Hauteville D, Bralet MP, Jasmin C. Combined paclitaxel and cetuximab achieved a major response on the skin metastases of a patient with epidermal growth factor receptor-positive, estrogen receptor-negative, progesterone receptor-negative and human epidermal growth factor receptor-2-negative (triple-negative) breast cancer. *Anticancer Drugs*. 2007;18:835–7.
179. Petrelli F, Cabiddu M, Ghilardi M, Barni S. Current data of targeted therapies for the treatment of triple-negative advanced breast cancer: empiricism or evidence-based? *Expert Opin Investig Drugs*. 2009;18:1467–77.
180. Ame JC, Spenlehauer C, de Murcia G. The PARP superfamily. *Bioessays*. 2004;26:882–93.
181. Dantzer F, de La Rubia G, Menissier-De Murcia J, Hostomsky Z, de Murcia G, Schreiber V. Base excision repair is impaired in mammalian cells lacking Poly(ADP-ribose) polymerase-1. *Biochemistry*. 2000;39:7559–69.
182. McCabe N, Turner NC, Lord CJ, et al. Deficiency in the repair of DNA damage by homologous recombination and sensitivity to poly(ADP-ribose) polymerase inhibition. *Cancer Res*. 2006;66:8109–15.
183. Ashworth A. A synthetic lethal therapeutic approach: poly(ADP) ribose polymerase inhibitors for the treatment of cancers deficient in DNA double-strand break repair. *J Clin Oncol*. 2008;26:3785–90.
184. Farmer H, McCabe N, Lord CJ. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature*. 2005;434:917–21.
185. Bryant HE, Schultz N, Thomas HD. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature*. 2005;434:913–7.
186. Tutt A, Robson M, Garber JE, Domchek SM, Audeh MW, Weitzel JN, et al. Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and advanced breast cancer: a proof-of-concept trial. *Lancet*. 2010;376:235–44.
187. Audeh MW, Carmichael J, Penson RT, Friedlander M, Powell B, Bell-McGuinn KM, et al. Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and recurrent ovarian cancer: a proof-of-concept trial. *Lancet*. 2010;376:245–51.
188. O’Shaughnessy J, Danso MA. A randomized phase III study of iniparib (BSI-201) in combination with gemcitabine/carboplatin (G/C) in metastatic triple-negative breast cancer (TNBC). ASCO 2011. Chicago. 2011.
189. O’Shaughnessy J, Osborne C, Pippen JE, Yoffe M, Patt D, Rocha C, et al. Iniparib plus chemotherapy in metastatic triple-negative breast cancer. *N Engl J Med*. 2011;364:205–14.
190. Metcalfe K, Lynch H, Gadirian P. Tamoxifen may reduce contralateral cancers in BRCA mutation carriers. ASCO Chicago: *J Clin Oncol*. 2011.
191. Wesolowski R SA, Tao J, Moore HC. Differential outcomes in patients treated with endocrine therapy for early or locally advanced breast cancer based on BRCA mutation status. In: *Oncol JC*, editor. ASCO 2009. Chicago. 2009.
192. ACSG Cancer Screening Guidelines. <http://www.cancer.org/healthy/findcancerearly/cancerscreening-guidelines/CAf,american-cancer-society-guidelines-for-the-early-,detection-of-cancer>. Accessed 6 Dec.
193. Lee CH, Dershaw DD, Kopans D. Breast cancer screening with imaging: recommendations from the Society of Breast Imaging and the ACR on the use of mammography, breast MRI, breast ultrasound, and other technologies for the detection of clinically occult breast cancer. *J Am Coll Radiol*. 2010;7:18–27.
194. Le-Petross HT, Whitman GJ, Atchley DP. Effectiveness of alternating mammography and magnetic resonance imaging for screening women with deleterious BRCA mutations at high risk of breast cancer. *Cancer*. 2011;117:3900–7.
195. NCCN GFH-RABaO, Available. [http://www.nccn.org/professionals/physician\\_gls/pdf/f\\_genetics\\_screening.pdf](http://www.nccn.org/professionals/physician_gls/pdf/f_genetics_screening.pdf). Accessed 9 Dec.
196. Brentnall TA. Cancer surveillance of patients from familial pancreatic cancer kindreds. *Med Clin North Am*. 2000;84:707–18.
197. Canto MI, Goggins M, Yeo CJ. Screening for pancreatic neoplasia in high-risk individuals: an EUS-based approach. *Clin Gastroenterol Hepatol*. 2004;2:606–21.
198. Cancer: [http://www.ahrq.gov/P-E-Af\\_clinic/uspstf08/colocancer/colors.htm](http://www.ahrq.gov/P-E-Af_clinic/uspstf08/colocancer/colors.htm). Accessed 2 Dec.
199. Gabriel EM, Jatoi I. Breast cancer chemoprevention. *Expert Rev Anticancer Ther*. 2012;12:223–8.
200. Fisher B, Costantino JP, Wickerham DL. Tamoxifen for prevention of breast cancer: report of the National Surgical Adjuvant Breast and Bowel Project P-1 Study. *J Natl Cancer Inst*. 1998;90:1371–88.
201. Fisher B, Costantino JP, Wickerham DL. Tamoxifen for the prevention of breast cancer: current status of the National Surgical Adjuvant Breast and Bowel Project P-1 study. *J Natl Cancer Inst*. 2005;97:1652–62.
202. Veronesi U, Maisonneuve P, Costa A. Prevention of breast cancer with tamoxifen: preliminary findings from the Italian randomised trial among hysterectomised women. Italian Tamoxifen Prevention Study. *Lancet*. 1998;352:93–7.
203. Decensi A, Robertson C, Rotmensz N. Effect of tamoxifen and transdermal hormone replacement therapy on cardiovascular risk factors in a prevention



- trial. Italian Chemoprevention Group. *Br J Cancer*. 1998;78:572–8.
204. Powles T, Eeles R, Ashley S. Interim analysis of the incidence of breast cancer in the Royal Marsden Hospital tamoxifen randomised chemoprevention trial. *Lancet*. 1998;352:98–101.
  205. Powles TJ, Ashley S, Tidy A, Smith IE, Dowsett M. Twenty-year follow-up of the Royal Marsden randomized, double-blinded tamoxifen breast cancer prevention trial. *J Natl Cancer Inst*. 2007;99:283–90.
  206. Cuzick J, Forbes J, Edwards R. First results from the International Breast Cancer Intervention Study (IBIS-I): a randomised prevention trial. *Lancet*. 2002;360:817–24.
  207. Cuzick J, Forbes JF, Sestak I, International Breast Cancer Intervention Study I Investigators. Long-term results of tamoxifen prophylaxis for breast cancer – 96-month follow-up of the randomized IBIS-I trial. *J Natl Cancer Inst*. 2007;99(4):272–82.
  208. Cuzick J, Powles T, Veronesi U, et al. Overview of the main outcomes in breast-cancer prevention trials. *Lancet*. 2003;361:296–300.
  209. Atkinson C, Warren R, Bingham SA, Day NE. Mammographic patterns as a predictive biomarker of breast cancer risk: effect of tamoxifen. *Cancer Epidemiol Biomarkers Prev*. 1999;8:863–6.
  210. Brisson J, Brisson B, Cote G, Maunsell E, Berube S, Robert J. Tamoxifen and mammographic breast densities. *Cancer Epidemiol Biomarkers Prev*. 2000;9:911–5.
  211. Cuzick J, Warwick J, Pinney E, Warren RM, Duffy SW. Tamoxifen and breast density in women at increased risk of breast cancer. *J Natl Cancer Inst*. 2004;96:621–8.
  212. Cuzick J, Warwick J, Pinney E. Tamoxifen-induced reduction in mammographic density and breast cancer risk reduction: a nested case–control study. *J Natl Cancer Inst*. 2011;103:744–52.
  213. Raloxifene hydrochloride. 2007. <http://www.fda.gov>.
  214. Ettinger B, Black DM, Mitlak BH. Reduction of vertebral fracture risk in postmenopausal women with osteoporosis treated with raloxifene: results from a 3-year randomized clinical trial. Multiple Outcomes of Raloxifene Evaluation (MORE) Investigators. *JAMA*. 1999;282:637–45.
  215. Cauley JA, Norton L, Lippman ME. Continued breast cancer risk reduction in postmenopausal women treated with raloxifene: 4-year results from the MORE trial. Multiple outcomes of raloxifene evaluation. *Breast Cancer Res Treat*. 2001;65:125–34.
  216. Cummings SR, Eckert S, Krueger KA. The effect of raloxifene on risk of breast cancer in postmenopausal women: results from the MORE randomized trial. Multiple Outcomes of Raloxifene Evaluation. *JAMA*. 1999;281:2189–97.
  217. Martino S, Cauley JA, Barrett-Connor E. Continuing outcomes relevant to Evista: breast cancer incidence in postmenopausal osteoporotic women in a randomized trial of raloxifene. *J Natl Cancer Inst*. 2004;96:1751–61.
  218. Vogel VG, Costantino JP, Wickerham DL. Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: the NSABP Study of Tamoxifen and Raloxifene (STAR) P-2 trial. *JAMA*. 2006;295:2727–41.
  219. Vogel VG, Costantino JP, Wickerham DL. Update of the National Surgical Adjuvant Breast and Bowel Project Study of Tamoxifen and Raloxifene (STAR) P-2 Trial: Preventing breast cancer. *Cancer Prev Res*. 2010;3:696–706.
  220. Goldstein SR, Neven P, Cummings S. Postmenopausal Evaluation and Risk Reduction with Lasofoxifene (PEARL) trial: 5-year gynecological outcomes. *Menopause*. 2011;18:17–22.
  221. Cummings SR, McClung M, Reginster JY. Arzoxifene for prevention of fractures and invasive breast cancer in postmenopausal women. *J Bone Miner Res*. 2011;26:397–404.
  222. Powles TJ, Diem SJ, Fabian CJ. Breast cancer incidence in postmenopausal women with osteoporosis or low bone mass using arzoxifene. *Breast Cancer Res Treat*. 2012;134:299–306.
  223. Goss PE, Strasser K. Aromatase inhibitors in the treatment and prevention of breast cancer. *J Clin Oncol*. 2001;19:881–94.
  224. Litton JK, Bevers TB, Arun BK. Exemestane in the prevention setting. *Ther Adv Med Oncol*. 2012;4:107–12.
  225. Baum M, Buzdar A, Cuzick J. Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early-stage breast cancer: results of the ATAC (Arimidex, Tamoxifen Alone or in Combination) trial efficacy and safety update analyses. *Cancer*. 2003;98:1802–10.
  226. Coombes RC, Hall E, Gibson LJ. A randomized trial of exemestane after two to three years of tamoxifen therapy in postmenopausal women with primary breast cancer. *N Engl J Med*. 2004;350:1081–92.
  227. Goss PE, Ingle JN, Martino S. A randomized trial of letrozole in postmenopausal women after five years of tamoxifen therapy for early-stage breast cancer. *N Engl J Med*. 2003;349:1793–802.
  228. Goss PE, Ingle JN, Ales-Martinez JE. Exemestane for breast-cancer prevention in postmenopausal women. *N Engl J Med*. 2011;364:2381–91.
  229. Cuzick J, Sestak I, Forbes JF. Anastrozole for prevention of breast cancer in high-risk postmenopausal women (IBIS-II): an international, double-blind, randomised placebo-controlled trial. *Lancet*. 2014;383:1041–8.
  230. Cuzick J, Sestak I, Bonanni B. Selective oestrogen receptor modulators in prevention of breast cancer: an updated meta-analysis of individual participant data. *Lancet*. 2013;381:1827–34.
  231. Arun B, Valero V, Liu D. Short-term biomarker modulation prevention study of anastrozole in women at increased risk for second primary breast cancer. *Cancer Prev Res*. 2012;5:276–82.
  232. King MC, Wieand S, Hale K. Tamoxifen and breast cancer incidence among women with inherited mutations

- in BRCA1 and BRCA2: National Surgical Adjuvant Breast and Bowel Project (NSABP-P1) Breast Cancer Prevention Trial. *JAMA*. 2001;286:2251–6.
233. Arun B, Dunn BK, Ford LG, Ryan A. Breast cancer prevention trials: large and small trials. *Semin Oncol*. 2010;37:367–83.
234. World Cancer Research Fund/American Institute for Cancer Research. Food N, physical activity, and the prevention of cancer: a global perspective. Washington DC: AICR; 2007.
235. Mahoney MC, Bevers T, Linos E, Willett WC. Opportunities and strategies for breast cancer prevention through risk reduction. *CA Cancer J Clin*. 2008;58:347–71.
236. Hamajima N, Hirose K, Tajima K. Alcohol, tobacco and breast cancer – collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with breast cancer and 95,067 women without the disease. *Br J Cancer*. 2002;87:1234–45.
237. Smith-Warner SA, Spiegelman D, Yaun SS. Alcohol and breast cancer in women: a pooled analysis of cohort studies. *JAMA*. 1998;279:535–40.
238. Larsson SC, Bergkvist L, Wolk A. Folate intake and risk of breast cancer by estrogen and progesterone receptor status in a Swedish cohort. *Cancer Epidemiol Biomarkers Prev*. 2008;17:3444–9.
239. Zhang SM, Willett WC, Selhub J. Plasma folate, vitamin B6, vitamin B12, homocysteine, and risk of breast cancer. *J Natl Cancer Inst*. 2003;95:373–80.
240. Prentice RL, Caan B, Chlebowski RT. Low-fat dietary pattern and risk of invasive breast cancer: the Women’s Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA*. 2006;295:629–42.
241. Boyd NF, Stone J, Vogt KN, Connelly BS, Martin LJ, Minkin S. Dietary fat and breast cancer risk revisited: a meta-analysis of the published literature. *Br J Cancer*. 2003;89:1672–85.
242. Missmer SA, Smith-Warner SA, Spiegelman D. Meat and dairy food consumption and breast cancer: a pooled analysis of cohort studies. *Int J Epidemiol*. 2002;31:78–85.
243. Taylor EF, Burley VJ, Greenwood DC, Cade JE. Meat consumption and risk of breast cancer in the UK Women’s Cohort Study. *Br J Cancer*. 2007;96:1139–46.
244. Willett WC. Nurses’ Health, Study II (NHS II): risk factors for breast cancer among younger nurses. Bethesda, MD: National Cancer Institute; [Accessed December 2013], <http://epi.grants.cancer.gov/Consortia/members/nhs2.html>.

---

# Breast Imaging and Image-Guided Biopsy Techniques

# 5

Marie Ganott, Brandy Griffith,  
and Scott M. Rudzinski

---

## Abstract

Breast imaging is an essential component of breast cancer diagnosis and guides surgery and treatment options. Imaging techniques such as mammography, ultrasound (US), and magnetic resonance imaging (MRI) enable the detection of breast cancer at earlier stages. Mammography remains the standard screening examination; however, additional imaging studies are useful in evaluating the breast. US is utilized primarily in the diagnostic setting to characterize mammographic or palpable findings and assess axillary lymph nodes. Supplemental screening US may also be useful in patients with intermediate risk for developing breast cancer and dense breasts to increase cancer detection. In addition to mammography, high-risk patients may also have annual MRI or screening US if they are unable to undergo MRI. MRI is also performed to evaluate the extent of disease, the response to neoadjuvant chemotherapy, and silicone implant integrity. In addition, these imaging modalities are also used to guide percutaneous biopsy, enabling minimally invasive tissue diagnosis. There is insufficient evidence to support the screening use of investigational imaging modalities, such as nuclear medicine breast imaging, and positron emission mammography. However, these topics are briefly discussed for completeness.

---

M. Ganott, MD (✉)  
Department of Radiology, Magee-Womens Hospital,  
University of Pittsburgh Medical Center,  
300 Halket Street, Pittsburgh, PA 15213, USA  
e-mail: [mganott@mail.magee.edu](mailto:mganott@mail.magee.edu)

B. Griffith, DO  
Department of Radiology, University of Pittsburgh  
Medical Center Hamot, 104 East 2nd Street,  
Erie, PA 16507, USA

Breast Imaging Department, Erie, PA, USA  
e-mail: [bmariegriffith@gmail.com](mailto:bmariegriffith@gmail.com)

S.M. Rudzinski, MD  
Windsong Radiology Group, 55 Sprindrift Drive,  
Williamsville, NY 14221, USA  
e-mail: [scottrudzinski@gmail.com](mailto:scottrudzinski@gmail.com)

### Keywords

3D mammogram • Aspiration • BI-RADS • Bracketing • Breast-specific gamma imaging (BSGI) • Computed tomography (CT) • Diagnostic • Elastography • High-risk screening • Magnetic resonance imaging (MRI) • Mammogram • Molecular breast imaging (MBI) • Nuclear medicine breast imaging • Positron emission mammography (PEM) • Radioactive seed • Stereotactic • Tomosynthesis • Ultrasound (US) • Wire localization

## Mammography

### Screening Mammography

Observational studies of screening mammography show that deaths due to breast cancer are reduced by 30–40 % among women who undergo screening mammography compared to those who do not [1, 2]. The American College of Radiology and American Cancer Society recommend beginning annual screening mammography at age 40 for women with an average risk of developing breast cancer [3, 4]. The probability of a mammogram detecting breast cancer in a woman age 40 years and older is much higher than the risk of mammography causing breast cancer. The cancer detection rate of screening mammography is approximately 2–7 per 1000 screened women, depending on the patient population [5]. Mammography provides the lowest dose of radiation among imaging modalities that utilize radiation to image the breasts.

Mammography can be performed using a film-screen or digital technique. The diagnostic accuracies of film-screen and digital mammography are similar in the general screening population. However, digital mammography is superior to film-screen mammography in women younger than 50 years, premenopausal or perimenopausal women, and women with dense breasts (heterogeneously or extremely dense) on mammography [6]. Digital mammography allows easier access to and storage of images and uses a lower dose of radiation [7]. Therefore, many breast imaging centers have converted to digital mammography despite the increased expense of digital mammographic units and the high-resolution monitors required for viewing images. The DMIST (Digital Mammographic Imaging Screening Trial) demon-


strated that breast doses in digital mammography were 22 % lower per view than those in film-screen mammography [7]. Two-view digital and film-screen mammograms have a mean average glandular dose of 3.7 and 4.7 mGy, respectively [8].

A screening mammographic examination is performed on an asymptomatic individual to detect clinically unsuspected breast cancer. The standard mammographic views are craniocaudal (CC) and mediolateral oblique (MLO) (Fig. 5.1). A specially trained technologist skilled in mammographic positioning obtains the X-ray images, while the patient's breast is tightly compressed between a compression plate and the image receptor. The images are then typically reviewed on special workstations designed for mammography if acquired digitally or on a light-box designed for viewing film if recorded on special mammography film. Computer-aided detection (CAD) is often utilized, in which application of a computer algorithm to the images marks potentially suspicious lesions and calcifications that may have been missed on the initial mammogram review.

### Diagnostic Mammography


A diagnostic mammogram is performed in several circumstances. It is utilized on a patient with clinical signs or symptoms of breast disease, on an individual for whom further evaluation has been requested due to an abnormal screening mammogram, or for follow-up of prior imaging findings [3]. In addition, women with a personal history of breast cancer who have been treated with breast conservation may choose to have diagnostic mammographic imaging. The patient waits in the department while the radiologist reviews the images; additional mammographic

**Fig. 5.1** Normal screening mammogram in a 45-year-old asymptomatic female. (a–d) Bilateral digital craniocaudal and mediolateral oblique views demonstrate heterogeneously dense breast tissue






**Fig. 5.2** A 60-year-old female with a palpable lump in the left breast. (a) CC, (b) MLO, and (c) magnified mediolateral mammograms demonstrate a spiculated mass in the upper inner breast corresponding to the palpable lump (*triangle marker*). (d) Ultrasound demonstrates an irregular hypoechoic mass (*arrows*). Percutaneous ultrasound-guided core biopsy revealed high-grade infiltrating duct carcinoma



views and/or ultrasound (US) may be obtained at that time to evaluate findings or symptoms. Because the appearance of normal breast tissue varies on mammography, comparison to prior mammograms is extremely valuable and can allow the radiologist to detect a subtle developing malignancy.


Additional mammographic views enable further characterization of a lesion. Mammographic views, including true lateral, rolled CC, exaggerated CC, tangential, and cleavage views, are used to help image more of the breast tissue and determine the location of a lesion. Spot compression views and tomosyn-

thesis evaluate the margins of a mass and help distinguish summation of normal tissue from a true mass. Spot compression uses a smaller paddle to focally compress the breast, spread out the overlapping tissue, and improve lesion conspicuity. Masses with microlobulated or indistinct margins are worrisome for malignancy, and masses with spiculated margins are highly suspicious for malignancy (Fig. 5.2). However, while circumscribed oval or round masses are usually benign, malignant tumors can have this appearance (Figs. 5.3 and 5.4). A mass is described as obscured when its margins are



**Fig. 5.3** A 51-year-old female with a palpable lump in the left upper outer quadrant. (a) Craniocaudal and (b) spot compression mediolateral mammograms of the left breast demonstrate a circumscribed oval mass corresponding to the

area of palpable concern (*triangle marker*). (c) Ultrasound demonstrates a circumscribed slightly hypoechoic mass (calipers) oriented parallel to the chest wall. Percutaneous ultrasound-guided core biopsy revealed a fibroadenoma



**Fig. 5.4** A 73-year-old asymptomatic female recalled from screening mammography. (a) Spot compression mediolateral left breast mammogram confirms a circumscribed dense mass. (b) Mediolateral tomosynthesis demonstrates this mass and a second circumscribed mass

(circles). (c) Ultrasound of the larger mass (calipers) demonstrates heterogeneous echotexture. Ultrasound-guided core biopsy of both masses demonstrated high-grade invasive duct carcinoma. Diffuse metastatic disease was found on subsequent PET/CT



hidden by glandular tissue and can only be adequately visualized with additional views or US.

Architectural distortion refers to radiating spicules from a central point without an obvious mass or may manifest as focal retraction of the breast tissue. Most commonly, architectural distortion is secondary to a surgical scar but can be a sign of a subtle invasive carcinoma, radial scar, ductal carcinoma in situ (DCIS), focal fibrosis, and other less common lesions. Distortion is best observed on magnified views or tomosynthesis images and is often subtle on US. A biopsy is performed, unless it is due to a surgical scar (Fig. 5.5).

Asymmetry is reported when an area of tissue density is seen on only one view and is usually an area of glandular tissue or a summation artifact from overlapping breast tissue. A focal asymmetry observed on both views but without the convex borders and conspicuity of a mass is also usually an area of glandular tissue. These areas are evaluated with additional views or tomosynthesis and compared with prior mammograms. If new when compared with prior studies, these areas are concerning for malignancy and require additional investigation, including US and potential biopsy (Fig. 5.6).




Magnification views are typically used to evaluate microcalcifications; however, these mammographic views may also be useful in characterizing the margins of a mass and architectural distortion. Certain calcification morphologies are typically benign, including skin, vascular, rim, dystrophic, round, large rodlike, and milk of calcium (Figs. 5.7 and 5.8). Circumscribed masses containing coarse popcorn-like calcifications are characteristic of calcifying fibroadenomas. Calcifications that are layered in a meniscal fashion on true lateral views represent milk of calcium in microcysts. Calcifications that are categorized as not classically benign include punctate, coarse heterogeneous, and amorphous, while fine pleomorphic and fine linear calcifications are often observed with malignancy. The distribution of calcifications is also an important factor in characterizing calcifications as suspicious or

benign. The distribution may be described as diffuse, regional, grouped or clustered, linear, and segmental. Pleomorphic and linear calcifications in a segmental or ductal distribution are highly suggestive of DCIS (ductal carcinoma in situ) and will undergo biopsy, typically with stereotactic guidance, because calcifications are not reliably visualized with US (Fig. 5.9). Calcifications that cannot be categorized definitively as benign or suspicious are reported as indeterminate and are usually also recommended for stereotactic biopsy. Calcifications that are believed to be probably benign, indicating a 2 % or less chance of malignancy, are usually recommended for observation, with follow-up mammography at 6-, 12-, and 24-month intervals if stable.


## Reporting

Radiologists use the American College of Radiology Breast Imaging Reporting and Data System (BI-RADS<sup>®</sup>) as a guide to standardize reporting, breast imaging terminology, assessment, and follow-up recommendations for mammography, US, and MRI of the breast [3]. A BI-RADS rating indicating the overall assessment of the imaging evaluation and the radiologist's recommendation for follow-up are required at the end of each mammography report. These assessment categories and recommendations are presented in Table 5.1.

The mammography report should include (and in some states is required by law) a comment on the density of the breast tissue. Breast density is affected by age, hormone replacement therapy, menstrual cycle phase, parity, body mass index, lactation, and familial predisposition. Dense tissue is more common in younger women; however, predominantly fatty or predominantly dense breast tissue can be found in women of any age. Breast density is typically subjectively categorized by the interpreting radiologist based on the amount of glandular tissue observed on mammography. Computer-assisted software programs may also be used to assess breast density. The categories are








**Fig. 5.6** (a) Right MLO and (b) left MLO views demonstrate a new asymmetry (*arrows*) in the posterior inferior aspect of the right breast. Incidentally noted is a biopsy clip from a prior benign biopsy in the right breast.

(c) Ultrasound demonstrates an irregular hypoechoic mass with indistinct margins (*arrows*). Subsequent ultrasound-guided biopsy revealed intermediate-grade ductal carcinoma in situ

**Fig. 5.5** (a) Craniocaudal and (b) mediolateral tomosynthesis images of the right breast demonstrate an area of architectural distortion (*arrows*) in the central outer right breast. (c) Ultrasound demonstrates a corresponding


irregular hypoechoic lesion (*arrows*) with acoustic shadowing. Percutaneous ultrasound-guided biopsy revealed a radial scar



**Fig. 5.7** CC view of the right breast demonstrates classic large rodlike (also called secretory) calcifications, which are dense, well defined, and often described as “cigar” shaped. They are in a ductal distribution directed toward the nipple and are usually bilateral

almost entirely fat (<25 % glandular), scattered fibroglandular (25–50 %), heterogeneously dense (50–75 %), and extremely dense (>75 % glandular) (Figs. 5.1 and 5.10). In the first two categories, 80 % of breast cancers are identified by mammography. Only 30–60 % of breast cancers may be seen on film-screen mammograms in heterogeneously and extremely dense breasts [9]. Digital mammography improves tissue contrast, increasing cancer detection in dense tissue up to 70 % compared with film-screen mammography [6].

The risk of breast cancer is increased in women with mammographically dense breasts [10]. Women with extremely dense breast tissue (>75 %) have a four- to sixfold increased risk of developing breast cancer compared to women




**Fig. 5.8** 41-year-old female with a history of right breast cancer 5 years ago, post-lumpectomy and radiation: CC mammogram of the right breast demonstrates postsurgical changes (*arrow*) from the prior lumpectomy in the right retroareolar breast with skin retraction and coarse calcifications typical of fat necrosis

with little or no dense tissue [11–14]. In addition, dense breast tissue lowers the sensitivity of mammography due to the ability of dense tissue to obscure tumors [15, 16].

## Imaging of the Male Breast

Gynecomastia is the most common reason for males to undergo breast imaging. It may present as a subareolar mass or breast enlargement that may be painful. Gynecomastia is initially evaluated with bilateral mammography to compare the breast tissue. There are three mammographic patterns of gynecomastia: early nodular, late dendritic, and diffuse glandular. The early nodular pattern occurs when the gynecomastia has been present for less than 1 year and is evident as a subareolar flame-shaped or fan-shaped den-



**Fig. 5.9** A 57-year-old asymptomatic female recalled from screening mammogram. (a) Magnified CC view of the right breast demonstrates pleomorphic calcifications (*arrows*) in the outer breast in a segmental distribution. (b, c) Stereotactic biopsy is performed utilizing paired images taken at 15° of

obliquity. (d) A specimen radiograph of the cores confirms the targeted calcifications were sampled in the biopsy. (e) CC view of the right breast demonstrates the clip (*arrow*) placed at the site of biopsy. Pathology revealed high-grade ductal carcinoma in situ

**Table 5.1** Concordance between BI-RADS® assessment categories and management recommendations

Assessment	Management	Likelihood of cancer
<i>Category 0:</i> Incomplete – need additional evaluation and/or prior mammograms for comparison	Recall for additional imaging and/or comparison with prior examination (s)	N/A
<i>Category 1:</i> Negative	Routine mammography screening	Essentially 0 % likelihood of malignancy
<i>Category 2:</i> Benign	Routine mammography screening	Essentially 0 % likelihood of malignancy
<i>Category 3:</i> Probably benign	Short-interval (6-month) follow-up or continued surveillance mammography	>0 % but ≤2 % likelihood of malignancy
<i>Category 4:</i> Suspicious	Tissue diagnosis	>2 % but <95 % likelihood of malignancy
<i>Category 4A:</i> Low suspicion for malignancy		>2 % but ≤10 % likelihood of malignancy
<i>Category 4B:</i> Moderate suspicion for malignancy		>10 to ≤50 % likelihood of malignancy
<i>Category 4C:</i> High suspicion for malignancy		>50 to <95 % likelihood of malignancy
<i>Category 5:</i> Highly suggestive of malignancy	Tissue diagnosis	≥95 % likelihood of malignancy
<i>Category 6:</i> Known biopsy-proven malignancy	Surgical excision when clinically appropriate	N/A

Reprinted with permission of the American College of Radiology (ACR). No other representation of this material is authorized without expressed, written permission from the ACR. Refer to the ACR website at [www.acr.org/Quality-Safety/Resources/BIRADS](http://www.acr.org/Quality-Safety/Resources/BIRADS) for the most current and complete version of the BIRADS® Atlas

sity (Fig. 5.11). The late dendritic appearance typically occurs after 1 year and is a flame-shaped, subareolar mass with radiating linear tissue extending into the posterior adipose tissue. The diffuse glandular pattern is due to exogenous estrogen (e.g., prostate cancer treatment, transsexuals) and is dense nodular parenchyma [17]. US is not always necessary to evaluate gynecomastia unless the mammographic appearance is atypical. Sonography of gynecomastia may reveal discrete round or hypoechoic nodular retroareolar tissue or irregular hypoechoic tissue that extends into the adjacent tissue (Fig. 5.11) [18].


Breast cancer in a man most commonly presents as a painless palpable subareolar mass. Up to 85 % of male breast cancers are invasive ductal carcinoma because the male breast lacks lobules except in states of excess estrogen, such as during treatments for prostate cancer and trans-

sexuals receiving estrogen [19–21]. DCIS occurs in up to 50 % of male breast cancers [19, 22]. The mammographic appearance of male breast cancer is that of a high-density round or oval mass with spiculated, lobulated, or indistinct margins (Fig. 5.12). Calcifications are less commonly seen than in females, occurring in only 13–30 % of cases. The calcifications are coarser and less linear than those associated with female breast cancer [19, 23]. The mass may be accompanied by skin thickening, skin or nipple retraction, and axillary adenopathy, as in female breast cancer. The differential diagnosis includes gynecomastia, fat necrosis, and metastatic disease.

The US features of invasive ductal carcinoma are typically a solid hypoechoic subareolar irregular mass with spiculated or microlobulated margins [24, 25]. Less commonly, the mass may be oval and circumscribed (Fig. 5.12).



**Fig. 5.10** A 35-year-old asymptomatic female for high-risk screening ultrasound due to strong family history and positive pathogenic BRCA-1 gene mutation. (a) Right MLO and (b) left MLO mammograms demonstrate extremely dense breasts




### High-Risk Screening

Women are considered at high risk of developing breast cancer if their estimated lifetime risk is 20 % or greater based on family history, if they have a known or suspected BRCA or other high-risk genetic mutations, or if they had mantle radiation therapy to the chest prior to age 30. Lifetime risk is calculated using one of several formulas, such as BRCAPRO, Claus, Gail, and Tyrer-Cuzick [26, 27].

The Society of Breast Imaging and American College of Radiology recommend that women with a BRCA1 or BRCA2 gene mutation or those who have not been tested but have a first-degree


relative with a known BRCA mutation have annual mammograms starting by age 30 but not before age 25. The recommendation for women with a greater than or equal to 20 % lifetime risk for breast cancer based on family history is to have annual mammography starting by age 30 (but not before age 25) or 10 years earlier than the age of diagnosis of the youngest affected relative, whichever is later. Women with a history of chest irradiation (usually as treatment for Hodgkin's disease) between the ages of 10 and 30 should have annual mammography starting 8 years after treatment (again, mammography is not recommended before age 25). Those women who have had a biopsy showing lobular carci-





**Fig. 5.11** A 50-year-old male presenting with right breast pain beneath the nipple for 1 year. (a) Right CC and (b) left CC mammograms demonstrate flame-shaped density in the right retroareolar region, consistent with

gynecomastia. (c) Ultrasound of the right retroareolar region demonstrates dispersed subareolar hypoechoic regions (*arrows*) consistent with breast tissue of gynecomastia



**Fig. 5.12** An 87-year-old male with a palpable left breast mass. (a) Left CC and (b) left magnification CC views demonstrate a multilobulated retroareolar mass with partially circumscribed and partially indistinct margins and associated pleomorphic calcifications. (c) Ultrasound of

the retroareolar left breast demonstrates a lobulated circumscribed hypoechoic mass with associated calcifications. Subsequent biopsy revealed infiltrating duct carcinoma, nuclear grade 2, and DCIS

noma in situ (LCIS), atypical lobular hyperplasia (ALH), atypical ductal hyperplasia (ADH), or DCIS should have annual mammograms from the time of diagnosis, regardless of age [27].

Although mammography has been the mainstay in breast cancer screening, other modalities, such as magnetic resonance imaging (MRI) and US, have established their relevance for women at increased risk for breast cancer.

Supplemental screening MRI has been recommended for women with a high risk of breast cancer according to a risk assessment tool that is based mainly on family history [26–28]. Several studies demonstrated significantly improved sensitivity in cancer detection in women with familial breast cancer when MRI screening was added to mammographic screening [29–31]. The cancer detection rate increased by 7–20 per 1,000 by adding screening MRI in high-risk patients [32]. The screening regimen for BRCA-positive patients, untested first-degree relatives of proven BRCA mutation carriers, and women with a >20 % lifetime risk for breast cancer consists of alternating mammography and MRI every 6 months or concurrent annual mammography and MRI starting by age 30. Women with a history of chest irradiation are recommended to have a screening MRI annually starting 8 years after the radiation therapy [27]. A recent study suggested that high-risk women who are unable to undergo MRI may benefit from screening US in addition to mammography [31].

---

## Ultrasound

US plays a pivotal role in the work-up and management of breast disease. US is a noninvasive imaging modality that uses high-frequency sound waves to produce images based on reflection from tissue interfaces. US is readily available and relatively low cost, has no associated radiation exposure, and does not require the administration of intravenous contrast. Continued technological advancements in conjunction with improved imaging skills and operator experience have

made US a critical modality in breast imaging and an indispensable adjunct to mammography. US is essential in imaging abnormalities that are not visible on mammography and is valuable in guiding needle biopsies of suspicious lesions and aspirating fluid collections. US can also be used to follow low suspicion lesions and to evaluate the response to neoadjuvant chemotherapy. In addition to greyscale imaging, elastography and Doppler sonography allow further characterization to help the radiologist appropriately categorize a lesion with regard to its stiffness and vascularity.

## Diagnostic Breast Ultrasound

A diagnostic work-up is performed when a patient has been recalled from a screening mammogram for a definite or suspected mammographic abnormality or for a symptomatic patient, with palpable abnormality being the most common symptom. Generally, mammography with additional views (i.e., spot compression, magnification, tangential view) is initially obtained in patients older than 30 years of age. If additional views alone are unable to determine whether a finding is benign or due to a summation artifact from overlapping glandular tissue, US is almost invariably the next imaging modality that is employed to better characterize the finding. All palpable abnormalities are worked up with physical exam and US. Although calcifications are generally worked up mammographically and biopsied stereotactically, US may be helpful in identifying a mammographically occult mass associated with the calcifications that could be biopsied with US guidance.

US is particularly useful in young patients (under 30 years of age) and in pregnant patients with breast complaints because of the lack of ionizing radiation and the decreased sensitivity of mammography due to dense breast tissue, which is more common in young patients. In these patients, US is the first (and typically the only) modality used to evaluate a breast concern.

Patients under 30 years of age with focal breast signs or symptoms (most commonly a palpable lump) have an incidence of malignancy less than 1 %, and the sensitivity and negative predictive value of US is extremely high [33]. Common benign findings in this age group include cysts, fibroadenomas, and normal breast tissue. US is also the first modality used in patients with a suspected breast abscess because these patients often have too much pain to tolerate the compression required for mammography (Fig. 5.13). Color Doppler ultrasound showing blood flow at the periphery rather than in the center of the lesion suggests an inflammatory process rather than a neoplasm.


US is useful in differentiating solid versus cystic breast lesions [34]. A simple cyst should be anechoic due to its homogeneous fluid content and lack of reflective interfaces. It should exhibit acoustic enhancement (hyperechogenic appearance of the tissue posterior to the structure) due to greater sound transmission through fluid than the surrounding breast tissue (Fig. 5.14). A solid mass contains internal echoes but is almost always hypoechoic relative to breast tissue. US features of benign lesions

include well-defined margins, few gentle lobulations, and a horizontal axis parallel to the chest wall (Figs. 5.3c and 5.15).


US findings of a suspicious mass include irregular or indistinct margins, a shape that is taller than it is wide, marked hypoechogenicity, and acoustic shadowing (loss of reflective interfaces posterior to the lesion due to diminished sound transmission through the lesion) (Figs. 5.16, 5.2d, and 5.6c). When a suspicious mass is identified, the remainder of the breast is evaluated, and the axilla is scanned for abnormal lymph nodes (Fig. 5.17). A lymph node is considered abnormal when it exhibits increased cortical thickness (>3 mm), eccentric cortical thickening, absence of a visible hilus, or loss of reniform shape, among other findings. However, these features are observed in both benign reactive and malignant lymphadenopathy.

## Screening Breast Ultrasound


Mammography continues to be the mainstay of breast screening, although there is a recent resurgence in the use of US as a screening tool



**Fig. 5.13** A 46-year-old female with a newly palpable mass in the periareolar right breast at 9 o'clock. Doppler color flow ultrasound image demonstrates increased peripheral flow surrounding a heterogeneous hypoechoic mass, which had mobile debris on real-time imaging. Purulent material was aspirated from this abscess




**Fig. 5.14** A 49-year-old female with right breast tenderness. Doppler color flow image of the right breast demonstrates the sonographic characteristics of a simple cyst (arrow), including anechoic contents, imperceptible wall, and enhanced posterior contrast through transmission of sound (arrow heads)



**Fig. 5.15** A 55-year-old female with a palpable lump in the left upper outer quadrant. (a) A well-defined mass is seen on sonography. (b) Ultrasound-guided core biopsy of

the mass revealed a fibroadenoma. The hyperechoic line traversing the lesion is the biopsy needle (arrows)



**Fig. 5.16** A 41-year-old asymptomatic female recalled from screening mammogram. Ultrasound of the left breast demonstrates an indistinct irregular hypoechoic mass with angulated margins that is taller than wide and demonstrates posterior acoustic shadowing (arrows); all of these findings are highly suspicious for malignancy. Ultrasound-guided core biopsy revealed intermediate-grade invasive duct carcinoma


in women with dense breast tissue, for whom mammography is less effective [6]. A recent large multicenter trial concluded that in high-risk women, the use of screening US in addition

to screening mammography increases the detection of cancer by 3–4 per 1000 over mammography alone [35]. This increased detection rate does come at the cost of increased false-positives. An average of 4.4 % of women underwent biopsy due to screening ultrasound findings, with a positive predictive value (PPV) of 9.4 % [36].

Screening whole-breast US is now being incorporated into breast-imaging practice, particularly as states implement legislation requiring that women be informed if they have dense breast tissue (>50 % glandular). Some states are now requiring that the mammography report sent to the patient include a statement that they may benefit from additional screening imaging, such as US or MRI [37].

Screening US is performed on an asymptomatic individual in conjunction with a screening mammogram in selected patient groups. The entire breast is scanned by an US technician. Images are taken of any findings and are reviewed by the radiologist. The patient is evaluated with a diagnostic US and, potentially, biopsy if suspicious lesions are found after reviewing the screening mammogram and ultrasound. Technology to enable automated screening of whole-breast ultrasound has also been developed and recently received FDA approval.





**Fig. 5.17** (a) A normal-appearing lymph node with thin cortex (*arrowheads*) and fatty hilum (*arrow*). (b) Blood flow at the hilum further characterizes this mass as a lymph node. (c) A 56-year-old female with history of breast

cancer presenting with an axillary mass. US demonstrates a markedly enlarged lymph node with a thick cortex, lack of a visible hilum, and loss of reniform shape. This mass was proven to be metastatic invasive breast cancer

## Magnetic Resonance Imaging of the Breast

MRI plays an important role in breast imaging. At present, annual screening breast MRI as an adjunct to mammography is recommended for patients with a known BRCA gene mutation, untested patients with a first-degree relative who is a BRCA carrier, patients who are considered high risk with a calculated lifetime risk of 20 % or greater, those with a history of chest irradiation

before age 30, and patients with Li-Fraumeni, Cowden, and Bannayan-Riley-Ruvalcaba syndromes and their first-degree relatives [26]. MRI has a sensitivity of 81 % in screening high-risk patients [38]. Despite its high sensitivity, MRI is not a replacement for mammography. MR images should be correlated with mammograms, US, and clinical findings for optimal interpretation. The sensitivity of MRI for detection of DCIS is less than for invasive cancers [39]. Suspicious calcifications on mammography require biopsy because



ductal carcinoma in situ lacked enhancement in 16 % of cases in a multicenter series [40]. However, not all DCIS manifests as calcifications on mammography; the sensitivity of MRI is higher than that of mammography for high-grade DCIS [41]. Although their sensitivity is high, MRI examinations suffer from a relatively lower specificity (reported at 67.4 % by Bluemke et al.), leading to the potential for an increase in follow-up studies and biopsies with negative results [42].

In addition, breast MRI is performed to evaluate the extent of disease in patients with known breast carcinoma, identify sites of residual malignancy status post-lumpectomy with positive margins, monitor the response to neoadjuvant chemotherapy, evaluate breasts for occult cancer in the presence of metastatic axillary lymphadenopathy with an unknown primary, and evaluate silicone implant integrity [43]. MRI may also be used to clarify imaging and physical findings.


The examination is performed in the prone position with simultaneous imaging of both breasts, usually in the axial plane, following the intravenous administration of gadolinium contrast. An MRI unit with a magnetic field strength of at least 1.5 T and a dedicated breast surface coil is required to obtain quality images. When MRI is performed solely to evaluate silicone implant integrity, gadolinium is not necessary.

MRI incorporates an evaluation of contrast enhancement kinetics and morphology to detect and assess breast lesions. Enhancing lesions are described as a focus/foci (an enhancing structure measuring  $\leq 4$  mm, too small to characterize), mass, or non-mass enhancement. Additional descriptors are used to explain the shape, margin, distribution, and internal enhancement pattern. However, normal fibroglandular tissue also exhibits physiological enhancement, which can make detection of malignancy more difficult and increase the chance of false-positives. Because the amount of background parenchymal enhancement is affected by hormonal status, elective MRI is performed early in the menstrual cycle, approximately days 7–10 after menses, to minimize this effect. Background parenchymal enhancement is reported as minimal, mild, moderate, or marked.

The contrast enhancement kinetics of a lesion are a reflection of the degree and type of vascularity of the lesion and are determined by obtaining 2–4 sequential scans of the breasts in 60–120 s per acquisition following the intravenous administration of gadolinium. Most MRI systems for breast imaging utilize computer software to evaluate how quickly the contrast enters and exits the mass to acquire the enhancement kinetic pattern. The percentage of increase in signal intensity as a function of time is plotted, and the program applies a designated color to the image depending on the enhancement patterns. The three categories of enhancement are persistent, plateau, and washout kinetics. Persistent kinetics (type I) indicate a continued increase in enhancement over time after the initial uptake of contrast and are suggestive of a benign lesion. Lesions with plateau kinetics (type II) have a steady enhancement pattern that does not change over time after the initial uptake phase. Washout kinetics (type III) occur when there is rapid washout of contrast from the lesion and are suggestive of malignancy. Lack of enhancement is strongly suggestive of benignity but does not necessarily exclude malignancy. Subtraction sequences are obtained during post-processing of the images and aid the detection of enhancing lesions. These sequences subtract the pre-contrast images from the post-contrast images and are also used to generate maximum intensity projection (MIP) images that provide a 3D overview of enhancing structures.

Invasive cancers will typically appear as a mass with an irregular shape with indistinct, irregular, or spiculated margins, as on mammography or US. The additional feature of internal enhancement, provided by MRI, is often heterogeneous or rim-like and usually rapid and strong with rapid washout (Fig. 5.18). Benign masses are typically oval or round in shape, with circumscribed margins and homogenous internal enhancement that progressively increases over time (persistent kinetic pattern) (Fig. 5.19).

Non-mass enhancement is described in terms of its distribution (focal, regional, linear, segmental, or diffuse) and internal features (homogenous, heterogeneous, clumped, or clustered ring) and can be due to hormonal, inflammatory, or




**Fig. 5.18** A 56-year-old female with recently diagnosed high-grade infiltrating ductal carcinoma. (a) Axial contrast-enhanced, (b) axial subtraction MR images, and (c) color MIP demonstrate an irregular mass in the left breast that exhibits enhancement with plateau and washout kinetics (color coded *green* and *red*, respectively). No

satellite lesions were identified, and the contralateral breast is unremarkable. Post-neoadjuvant chemotherapy (d) axial subtraction MR image and (e) color MIP image demonstrate complete response with no evidence of an enhancing mass within the breast. Final pathology of the segmental mastectomy demonstrated no residual invasive carcinoma

fibrocystic changes. However, it can also be a manifestation of invasive lobular cancer or DCIS. Linear or segmental clumped enhancement is suggestive of malignancy (Fig. 5.20).

MRI is valuable in the detection of breast cancer in patients presenting with carcinoma of unknown primary [44, 45], particularly if the patient presents with axillary adenopathy and a negative mammogram and US. MRI detects the primary breast cancer in two-thirds of these patients, facilitating

breast conservation instead of mastectomy (Fig. 5.21). When the suspected primary lesion is detected on MRI, a second-look US and US-guided biopsy will be performed. If the lesion cannot be found using US, MRI biopsy is performed. Targeted US performed to search for lesions detected on MRI is successful in finding a correlate between 50 and 60 % of cases; mass lesions seen on MRI are more than two times likely to be sonographically recognized than non-mass enhance-



**Fig. 5.19** A 60-year-old female with MR evaluation of the extent of disease for biopsy-proven carcinoma (not shown). (a) An axial contrast-enhanced MR of the right breast shows an oval circumscribed homogeneously


enhancing mass with (b) persistent kinetics, indicated by the *blue color* coding, consistent with benign disease. The known IDC/DCIS is not shown



**Fig. 5.20** A 50-year-old asymptomatic female for high-risk screening MR. A contrast-enhanced axial T1-weighted image demonstrates clumped linear enhancement (*arrow*) in the central upper breast. MR-guided biopsy revealed intermediate-grade invasive duct carcinoma and intermediate-grade DCIS

ment, and malignant masses are more likely than benign masses to be found with US [46, 47].


MRI is commonly used to assess the extent of disease preoperatively in patients newly diagnosed with breast cancer to aid surgical planning. MRI detects unsuspected cancer in the contralateral breast in 3 % of these patients [48]. The size of invasive carcinoma is most accurately assessed by MRI [49]. MRI can detect additional sites of malignancy in the ipsilateral breast and identify chest wall and nipple-areolar involvement. MRI detects additional disease [50–52] in 27–34 % of patients, resulting in wider surgical excision or mastectomy (Fig. 5.22). However, the use of preoperative MRI has been criticized due to its potential for overtreatment without clinical benefit. A recent meta-analysis [53] showed no reduction in local or distant recurrence rates in patients who had MRI



**Fig. 5.21** A 54-year-old female presenting with a new lump in the left axilla. **(a)** Ultrasound of the left axilla demonstrates a grossly enlarged left axillary lymph node highly suspicious for metastatic disease. US-guided core biopsy of the node revealed high-grade infiltrating duct carcinoma. **(b)** Left CC and **(c)** left MLO mammogram performed 6 weeks prior demonstrated no evidence of malignancy. MR was

subsequently performed to evaluate for an unknown primary tumor. **(d)** Contrast-enhanced axial T1-weighted images and **(e)** color MIP images demonstrate a clumped linear area of enhancement (*arrow*) with washout kinetics (color coded *red*) in the posterior left breast. **(f)** This lesion was observed on a second-look ultrasound and was biopsied with ultrasound guidance revealing infiltrating duct carcinoma





**Fig. 5.21** (continued)

preoperative staging compared with those who had not, although an earlier study [54] did show a reduction in local recurrence. The reason for this discrepancy is not clear but has been hypothesized to be due to the effectiveness of radiation and chemotherapy in treating additional sites of disease that were not removed at surgery. In addition, the COMICE (Comparative Effectiveness of MRI in Breast Cancer) trial found similar reoperation rates when comparing patients with and without preoperative MRI staging [55]. If a staging MRI for extent of disease identifies additional sites of suspected malignancy, percutaneous biopsy is performed for histological proof, either with US if found on second-look or with MRI-guided biopsy, when due to overlap in the appearance of benign and malignant lesions. False-positive lesions are common; the PPV of biopsy of additional lesions found on the extent of disease observed by MRI at Magee-Womens Hospital at the University of Pittsburgh Medical Center was 30 % after a biopsy recommendation rate of 35 % [56].


To assess the response to neoadjuvant chemotherapy (Fig. 5.18), MRI can be used to monitor

early response to treatment by revealing a change in tumor kinetics before a change in tumor size or to identify residual disease prior to surgery. MRI shows better correlation with pathological response than clinical or mammographic evaluation but still may underestimate or overestimate residual disease [57–59]. A negative MRI after neoadjuvant therapy does not exclude residual disease.

MRI can be helpful in depicting residual disease after segmental mastectomy when positive surgical margins are identified pathologically, but postoperative reactive enhancement can confound the findings. Lee et al. reported a sensitivity, specificity, and accuracy of 61.2, 69.7, and 64.6 % for MRI in identifying residual disease after excisional biopsy for breast cancer [60].

Contraindications to MRI include ferromagnetic medical devices such as intracranial aneurysm clips, cardiac pacemakers, and insulin pumps in addition to orbital metallic foreign bodies. Other contraindications include pregnancy (if gadolinium is required), claustrophobia that is not controlled by premedication, patient weight exceeding the limit of the MRI table, and an





**Fig. 5.22** A 57-year-old female recalled from screening mammogram for calcifications. **(a)** Spot compression magnification mediolateral view of the upper left breast demonstrates a small cluster of pleomorphic calcifications (*arrow*). Subsequent stereotactic biopsy revealed invasive duct carcinoma and DCIS. **(b)** An axial contrast-enhanced T1-weighted MR image demonstrates a spiculated enhancing mass in the upper left breast. **(c)** An axial image just inferior to this spiculated mass shows an area of segmental clumped non-mass enhancement (*arrows*)

suspicious for additional disease. **(d)** A sagittal contrast-enhanced T1-weighted MR image demonstrates the segmental clumped non-mass enhancement (*arrows*), suggesting a much greater extent of disease than expected based on the calcifications observed on the mammogram; the known primary carcinoma is partially visualized superior to this enhancement (*arrow head*). The segmental enhancement was biopsied with MRI guidance, revealing invasive duct carcinoma and DCIS. The identification of additional disease resulted in mastectomy

uncooperative patient or a patient who is unable to consent to the exam. Renal insufficiency is also a relative contraindication because there is an association with intravenous gadolinium and nephrogenic systemic fibrosis [61].

## Ductography


Ductography is performed to investigate the cause of unilateral clear or bloody nipple discharge and may aid the surgeon in successfully removing the lesion by delineating its location and extent. A 30-gauge sialogram or ductogram cannula is inserted into the discharging duct, which is then gently injected with radiopaque contrast material. Magnified mammography views performed with the cannula in place should reveal a filling defect in the duct if a mass is present and if the duct is adequately opacified with contrast. A papilloma usually appears as a small isolated lobulated retroareolar mass, but surgical excision is performed for histological diagnosis (Fig. 5.23). Irregularity and narrowing of the duct system suggest ductal carcinoma. Duct cannulation is not always successful and is not possible if the discharge cannot be expressed.

## New and Investigational Imaging Modalities

Although mammography, US, and MRI are effective in identifying breast malignancy, there is a role for other imaging techniques that may reduce the number of diagnostic work-ups and biopsies of benign lesions.

## Tomosynthesis

Digital breast tomosynthesis, also known as 3D mammography, involves the acquisition of 10–20 digital images of a compressed breast over a 15–50° angle using lower-dose X-rays than those used in standard digital mammography. These angled images are reconstructed as 1-mm thin sections through the breast [8]. This reconstruction allows the radiologist to scroll through the breast tissue in slices, removing the summation artifact caused by overlapping tissue. This process reduces recall rates from screening mammography and increases cancer detection [62]. A comparative study of 12,631 patients by Skaane et al. showed an improvement in the cancer detection rate from 6.1 per 1000 to 8 per



**Fig. 5.23** A 72-year-old female complaining of multiple episodes of right nipple discharge. (a) Right CC mammogram after injection of contrast material demonstrates a circumscribed lobulated filling defect in the duct (*arrow*).

(b) Ultrasound of the same region demonstrates an isoechoic circumscribed intraductal mass (calipers) within a dilated duct. Subsequent biopsy revealed an intraductal papilloma

1000 with the addition of tomosynthesis to standard screening mammography [63].

The radiation dose of a single tomosynthesis study is approximately equivalent to a two-view mammogram. Because tomosynthesis is typically performed in conjunction with a mammogram, this method doubles the radiation dose [8]. A method to synthesize a 2D mammogram from the tomosynthesis images is now FDA approved and would thus reduce the radiation dose to that of a standard mammogram while allowing acquisition of both 2D and 3D images.

## Elastography

Multiple techniques and technological approaches are used in elastography, some enabling only qualitative, and others both qualitative and quantitative assessment of lesions. The technology is incorporated with US and allows assessment of the stiffness of a lesion. Breast cancers tend to be stiffer than normal breast tissue; therefore, elastography may aid the characterization of indeterminate lesions on US and aid the determination of which lesions should be biopsied and which can be safely followed [64, 65].

## Nuclear Medicine Breast Imaging

Nuclear medicine breast imaging has been used for screening, to evaluate the response to treatment, and problem solving. It began as scintimammography with the use of a sodium iodide nuclear camera but had limited ability to detect small lesions. Breast-specific gamma imaging (BSGI) uses a single detector with a sodium iodide camera, and molecular breast imaging (MBI) uses two opposing cadmium zinc telluride (CZT) digital detectors, which allows the detection of smaller lesions and the administration of lower doses of the technetium-99m sestamibi radionuclide.

The breast is placed in mild compression to immobilize the breast and is imaged after the injection of technetium-99m sestamibi, which accumulates in the mitochondria of the cells. The

amount of cellular mitochondria is a marker of cellular proliferation. Rapidly dividing cells such as cancer cells have more mitochondria and therefore have a higher uptake of the radiotracer than normal cells [66].

The detectors create a 2D image of the breast, usually in the craniocaudal and mediolateral oblique positions, although other projections may be obtained. Each image takes approximately 10 min to acquire; therefore, standard four-view exams take 40 min (Fig. 5.24). Because this method is a physiological examination, adjunct imaging techniques must be used for lesion characterization and to guide biopsy. Technology is being developed that would incorporate guidance for biopsies.

MBI has a high degree of sensitivity, particularly for invasive cancers [67]. Studies have indicated a sensitivity of MBI of 85–100 % and a specificity of 60–90 % [66]. A disadvantage of MBI is the higher radiation dose compared to mammography. A BSGI examination provides the highest organ doses to the large intestine wall, kidneys, bladder wall, and gallbladder wall. However, the breast dose is one-tenth of this dose, with an effective dose of 2 mGy [8] when 30 mCi of technetium-99m is used. The effectiveness of imaging with smaller doses of the radionuclide is being investigated.

## Positron Emission Mammography

Positron emission mammography (PEM) was developed to overcome the limitations of whole-body PET (positron emission tomography) in detecting breast cancer and uses two opposing dedicated gamma radiation detectors placed above and below the breast while the breast is in mild compression, resulting in improved sensitivity and better spatial resolution and allowing the detection of lesions <1 cm [68]. The images are acquired after the intravenous administration of fluorine<sup>18</sup> fluorodeoxyglucose (FDG), a radionuclide that is similar in structure to glucose and is taken up by metabolically active tumor cells. The images are displayed as 12 tomographic slices in two projections of each breast, facilitating 3D

localization of abnormal FDG uptake. A disadvantage of this technology is the required fasting for 4–6 h prior to injection and the 10-min cranio-caudal and mediolateral oblique view imaging time [69]. Another disadvantage is the higher radiation dose to organs other than the breast compared with mammography. The highest organ doses associated with PEM are to the bladder, uterus, and ovaries. The breast dose is approximately 2.5 mGy [8]. A meta-analysis of 8 studies of 873 women with suspected breast malignancy reported that the pooled sensitivity and specificity values of PEM were 85 and 79 %, respectively [70]. In comparing PEM and MRI, Berg et al. found MRI to be more sensitive (14/15 vs. 3/15) for detecting contralateral cancer in women who were newly diagnosed with breast cancer. Although in a study of 388 women, Berg et al. observed similar sensitivity of the two modalities in disease detection, MRI was more sensitive in identifying women who needed additional surgery to excise all disease in the ipsilateral breast. PEM was more specific than MRI and less likely to prompt unnecessary biopsies. PEM is an alternative for women who are unable to tolerate MRI to evaluate the extent of disease [71, 72].

---

## **Image-Guided Biopsy and Localization of Non-palpable Breast Lesions**

### **Stereotactic-Guided Biopsy**

Stereotactic guidance is used for percutaneous biopsy of mammographically identified lesions, including microcalcifications, masses, asymmetries, and architectural distortions that cannot be located using US. The patient is positioned either prone or seated, and the breast is compressed between the image receptor and the compression plate. Two 15-degree angled X-ray images of the lesion allow targeting by the radiologist to produce computer-generated coordinates that are transferred to the stereotactic biopsy device. The biopsy needle is advanced into the breast, and additional images are obtained to confirm appropriate positioning.

Tissue samples are then obtained, typically utilizing an 8- to 11-gauge vacuum-assisted biopsy device (Fig. 5.9).


When the biopsy is performed for calcifications, magnification X-rays of the tissue specimen are obtained to verify that the calcifications of interest were sampled. Following biopsy, a tissue marker (“biopsy clip”) is placed at the biopsy site. The tissue marker indicates the site that was biopsied and facilitates localization if a lumpectomy or excision needs to be performed. Radiological pathological correlation should be performed to ensure that adequate sampling was performed.

### **Ultrasound-Guided Needle Aspiration and Fine-Needle Aspiration Biopsy**


Aspiration of cysts may be performed under US guidance for greatest accuracy because the needle can be visualized within the cyst; the fluid in the cyst will diminish as it is aspirated. Simple cysts do not require treatment; however, some patients desire aspiration for symptomatic relief. Complicated cysts contain homogeneous echogenicity or mobile internal echoes on US. If a mass cannot be identified as a complicated cyst or a solid mass, aspiration may be performed initially (Fig. 5.25). If the fluid is cloudy yellow, green, gray, or clear yellow, it is usually discarded. If the fluid is bloody, it should be sent for cytology. If no fluid can be aspirated, it can be assumed that the mass is solid, and a core biopsy may be performed. If the cyst contains a solid nodule, the nodule should undergo core biopsy prior to aspiration of the fluid.

Due to the increased rates of insufficient sampling and higher false-negative rates with fine-needle aspiration biopsy, core biopsy is preferred for breast masses [73, 74]. Fine-needle aspiration may still be performed in cases in which the lesion may be difficult to safely biopsy due to an adjacent vessel or breast implant. Typically, a 21- or 25-gauge needle is rapidly and manually oscillated throughout the









**Fig. 5.24** (continued)

lesion under US guidance, while suction is applied with a syringe connected to the needle with flexible tubing until there is blood or material within the needle hub. The material is immediately applied to slides, smeared, and sent for cytology.

### Ultrasound-Guided Core Biopsy

US is the most common method to guide biopsy of an indeterminate or suspicious lesion and allows visualization of the biopsy needle within the lesion in real time. US-guided core biopsy is usually performed with an 11- to 18-gauge biopsy device. Although there are a variety of devices, the most widely used are spring loaded, in which the needle is rapidly “fired” through the lesion



**Fig. 5.25** A 61-year-old female with a palpable area in the upper outer right breast. Ultrasound demonstrates an oval hypoechoic mass in the upper outer right breast with internal echoes. Aspiration was performed, yielding turbid white fluid consistent with proteinaceous debris, which was discarded. The lesion resolved upon aspiration, indicating that it was a complicated cyst

(Fig. 5.15). Vacuum-assisted core biopsy devices are also available and pull the tissue into the needle trough prior to cutting. A review of seven series including 884 cancers revealed that 840 (95 %) of the cancers were diagnosed at initial 14-gauge core biopsy [75]. In a review of 2,420 cases with long-term follow-up by Youk et al., US-guided 14-gauge core needle biopsy had a 96 % sensitivity rate and a false-negative rate of 2.4 % for malignancy (1.9 % with high-risk lesions on core were upgraded to malignancy). The malignant upgrade rate for high-risk lesions was 27 %, and the upgrade rate for DCIS at core to invasive disease was 29 % [76]. Inadequate sampling is particularly a concern with smaller

**Fig. 5.24** A 62-year-old female recalled from a screening mammogram. (a) A left spot compression magnified CC view demonstrates a spiculated mass in the inner left breast (arrow). (b) Ultrasound demonstrates a hypoechoic mass with indistinct margins and posterior shadowing.

(c) Tc 99m sestamibi left breast CC and MLO images and (d) color MBI CC images of the left breast demonstrate uptake in the inner left breast. Subsequent ultrasound-guided biopsy revealed high-grade invasive duct carcinoma

lesions. The needle may appear to be within the lesion when it is actually adjacent to it. Therefore, turning the US probe perpendicular to the lesion after a pass to visualize the needle within the lesion and acquiring more passes can be helpful in ensuring adequate sampling. If the pathology results are not concordant with the imaging finding, surgical excision or additional tissue sampling is recommended.

A biopsy marker clip should be placed into the lesion following core biopsy to facilitate removal of the lesion in cases of atypia or malignancy. A post-procedure mammogram is obtained to ensure that the clip is in the expected location.

US-guided biopsies are typically better tolerated by patients than stereotactic or MRI-guided procedures because the patient is supine and the breast is not in compression during a US-guided procedure. Therefore, a “second-look” US will often be performed to identify a sonographic correlate to a concerning MRI finding to facilitate an US-guided biopsy. Common post-procedure symptoms include bruising and pain, while hematomas are uncommon. Color Doppler should be used to ensure that blood vessels are avoided.

### **MRI-Guided Biopsy**

MRI-guided biopsies are performed when a lesion can only be identified on the MR examination. The computer system used for evaluation of enhancement patterns typically also has an adjunct system to aid in performing percutaneous biopsy. However, MRI biopsy is challenging because the lesion identified for sampling may be more difficult to identify when the patient’s breast is compressed by the grid required for immobilization and localization. The patient must remain in this position after the initial post-contrast images, during imaging to visualize the position of the introducer relative to the lesion, and during subsequent sampling with a vacuum-assisted biopsy device, clip placement, and post-biopsy imaging. After the procedure is completed, the patient is transferred from the MRI suite to the mammography suite to confirm clip placement.

### **Wire Localization**


Wire localizations may be performed preoperatively under mammographic or US guidance to direct the surgeon to the tissue that needs to be excised. This procedure is typically performed for non-palpable lesions.

When the procedure is performed using mammographic guidance, the breast is placed into compression, and a grid is projected over the skin and used to target the area to be removed. After obtaining local anesthesia, a hollow needle containing a fine wire is placed into the breast. An orthogonal mammogram is then performed to evaluate the needle depth. Once the needle is in the appropriate position, the needle is removed, and the wire is left in place. The wire has a hooked or curved end that is designed to prevent dislodgement from the breast (Fig. 5.26). When the procedure is performed using US guidance, the needle containing the wire is placed through the lesion, and the needle is removed, leaving the wire in place.

A radioactive seed may also be placed through the hollow needle instead of a wire to localize the lesion to be excised. The seed contains a small amount of I-125 within a titanium shell. The surgeon uses a gamma probe to detect the seed and guide the excision in addition to the mammogram images depicting the location of the seed.

The bracketing technique utilizes multiple wires or seeds to “bracket” the boundaries of large or complex lesions.

In all cases, craniocaudal and mediolateral mammogram views are obtained and annotated by the radiologist to delineate the lesion to be removed. A specimen radiograph should be obtained at the time of surgery to confirm that the localized lesion has been removed, along with biopsy clips, wires, or radioactive seeds. The radiologist should call the operating room to inform the surgeon that the lesion or biopsy marker and wire or seed is present in the specimen and to notify the surgeon if the excised lesion is close to the margin of the specimen so that additional tissue can be obtained immedi-



**Fig. 5.26** (a) A compression plate with a grid is placed on the breast overlying the calcifications to be localized for surgical excision. Needles (*arrows*) with coaxial wires are placed into position from a lateral approach. (b) An orthogonal view shows the position of the needles, which can be

adjusted. (c) The wires are deployed, bracketing the anterior and posterior extent of the calcifications. (d) A specimen radiograph shows the wires, the excised calcifications, and the stereotactic biopsy clip. Stereotactic biopsy and surgical excision led to a diagnosis of high-grade DCIS

ately if deemed necessary by the surgeon. It may be difficult to determine which margin is close; surgically placed markers at the margins, ortho-

gonal specimen views and careful viewing of the lesion and its relationship to the localizing markers will aid this assessment.

## References

1. Tabár L, Vitak B, Chen TH, Yen AM, Cohen A, Tot T, et al. Swedish two-county trial: impact of mammographic screening on breast cancer mortality during 3 decades. *Radiology*. 2011;260(3):658–63.
2. Coldman A, Phillips N, Warren L, Kan L. Breast cancer mortality after screening mammography in British Columbia women. *Int J Cancer*. 2007;120(5):1076–80.
3. ACR practice guidelines for the performance of screening and diagnostic mammography. Amended 2014 (Resolution 39). [www.acr.org](http://www.acr.org).
4. Sickles EA, D'Orsi CJ, Bassett LW. ACR BI-RADS® mammography. In: ACR BI-RADS® Atlas, breast imaging reporting and data system. Reston: American College of Radiology; 2013.
5. Rosenberg RD, Yankaskas BC, Abraham LA, Sickles DA, Lehman CD, Geller BM, et al. Performance benchmarks for screening mammography. *Radiology*. 2006;241(1):55–66.
6. Pisano ED, Gatsonis C, Hendrick E, Yaffe M, Baum JK, Acharyya S, Conant EF, Fajardo LL, Bassett L, D'Orsi C, Jong R, Rebner M, Digital Mammographic Imaging Screening Trial (DMIST) Investigators Group. Diagnostic performance of digital versus film mammography for breast-cancer screening. *N Engl J Med*. 2005;353(17):1773–83.
7. Hendrick RE, Pisano ED, Averbukh A, Moran C, Berns EA, Yaffe MJ, et al. Comparison of acquisition parameters and breast dose in digital mammography and screen-film mammography in the American College of Radiology Imaging Network digital mammographic imaging screening trial. *AJR Am J Roentgenol*. 2010;194(2):362–9.
8. Hendrick RE. Radiation doses and cancer risks from breast imaging studies. *Radiology*. 2010;257(1):246–53.
9. Mandelson MT, Oestreicher N, Porter PL, White D, Finder CA, Taplin SH, et al. Breast density as a predictor of mammographic detection: comparison of interval- and screen-detected cancers. *J Natl Cancer Inst*. 2000;92(13):1081–7.
10. Boyd NF, Guo H, Martin LJ, Sun L, Stone J, Fishell E, et al. Mammographic density and the risk and detection of breast cancer. *N Engl J Med*. 2007;356(3):227–36.
11. Ursin G, Ma H, Wu AH, Bernstein L, Salane M, Parisky YR, et al. Mammographic density and breast cancer in three ethnic groups. *Cancer Epidemiol Biomarkers Prev*. 2003;12:332–8.
12. Byrne C, Schairer C, Wolfe J, Parekh N, Salane M, Brinton LA, et al. Mammographic features and breast cancer risk: effects with time, age, and menopause status. *J Natl Cancer Inst*. 1995;87:670–5.
13. Harvey JA, Bovbjerg VE. Quantitative assessment of mammographic breast density: relationship with breast cancer risk. *Radiology*. 2004;230:29–41.
14. McCormack VA, dos Santos Silva I. Breast density and parenchymal patterns as markers of breast cancer risk: a meta-analysis. *Cancer Epidemiol Biomarkers Prev*. 2006;15:11159–69.
15. Sala E, Warren R, McCann J, Duffy S, Day N, Luben R. Mammographic parenchymal patterns and mode of detection: implications for the breast screening programme. *J Med Screen*. 1998;5:207–12.
16. Van Gils CH, Otten JD, Verbeek AL, Hendriks JH. Mammographic breast density and risk of breast cancer: masking bias or causality? *Eur J Epidemiol*. 1998;14:315–20.
17. Berg WA. Special topics. In: Berg WB, Yang WT, editors. *Diagnostic imaging breast*, vol. 8. 2nd ed. Manitoba: Amysis; 2014. p. 98–100.
18. Dialani V, Baum J, Mehta TS. Sonographic features of gynecomastia. *J Ultrasound Med*. 2010;29(4):539–47.
19. Günhan-Bilgen I, Bozkaya H, Ustün E, Memiş A. Male breast disease: clinical, mammographic, and ultrasonographic features. *Eur J Radiol*. 2002;43(3):246–55.
20. Goss PE, Reid C, Pintilie M, Lim R, Miller N. Male breast carcinoma: a review of 229 patients who presented to the Princess Margaret Hospital during 40 years: 1955–1996. *Cancer*. 1996;85(3):629–39.
21. Symmers WS. Carcinoma of breast in trans-sexual individuals after surgical and hormonal interference with the primary and secondary sex characteristics. *Br Med J*. 1968;2(5597):83–5.
22. Chantra PK, So GJ, Wollman JS, Bassett LW. Mammography of the male breast. *AJR Am J Roentgenol*. 1995;164(4):853–8.
23. Shi AA, Georgian-Smith D, Cornell LD, Rafferty EA, Staffa M, Hughes K, et al. Radiological reasoning: male breast mass with calcifications. *AJR Am J Roentgenol*. 2005;185(6 Suppl):S205–10.
24. Chen L, Chantra PK, Larsen LH, Barton P, Rohitopakarn M, Zhu EQ, et al. Imaging characteristics of malignant lesions of the male breast. *Radiographics*. 2006;26(4):993–1006.
25. Adibelli ZH, Oztekin O, Gunhan-Bilgen I, Postaci H, Uslu A, Ilhan E. Imaging characteristics of male breast disease. *Breast J*. 2010;16(5):510–8.
26. Breast Cancer Screening and Diagnosis. NCCN Guidelines Version 1. 2015. [www.nccn.org](http://www.nccn.org).
27. Lee CH, Dershaw DD, Kopans D, Evans P, Monsees B, Monticciolo D, et al. Breast cancer screening with imaging: recommendations from the Society of Breast Imaging and the ACR on the use of mammography, breast MRI, breast ultrasound, and other technologies for the detection of clinically occult breast cancer. *J Am Coll Radiol*. 2010;7(1):18–27.
28. American Cancer Society. Detailed guide: breast cancer. Accessed at <http://www.cancer.org/Cancer/BreastCancer/DetailedGuide/index> on 4 Feb 2013.
29. Kuhl C, Weigel S, Schrading S, Arand B, Bieling H, König R, et al. Prospective multicenter cohort study to refine management recommendations for women at elevated familial risk of breast cancer: the EVA trial. *J Clin Oncol*. 2010;28(9):1450–7.



30. Kuhl C. The current status of breast MR imaging. Part I. Choice of technique, image interpretation, diagnostic accuracy, and transfer to clinical practice. *Radiology*. 2007;244(2):356–78.
31. Berg WA, Zhang Z, Lehrer D, Jong RA, Pisano ED, Barr RG, ACRIN 6666 Investigators, et al. Detection of breast cancer with addition of annual screening ultrasound or a single screening MRI to mammography in women with elevated breast cancer risk. *JAMA*. 2012;307(13):1394–404.
32. Berg WA. Screening and management. In: Berg WA, Yang WT, editors. *Diagnostic imaging breast*. 2nd ed. Manitoba: Amirsys; 2014, 9:45.
33. Loving VA, DeMartini WB, Eby PR, Gutierrez RL, Peacock S, Lehman CD. Targeted ultrasound in women younger than 30 years with focal breast signs or symptoms: outcomes analyses and management implications. *Am J Roentgenol*. 2010;195:1472–7.
34. Harper AP, Kelly-Fry E, Noe JS, Bies JR, Jackson VP. Ultrasound in the evaluation of solid breast masses. *Radiology*. 1983;146(3):731–6.
35. Berg WA, Blume JD, Cormack JB, Mendelson EB, Lehrer D, Böhm-Vélez M, et al. Combined screening with ultrasound and mammography vs mammography alone in women at elevated risk of breast cancer. *JAMA*. 2008;299(18):2151–63.
36. Berg WA. Screening and management. In: Berg WA, Yang WT, editors. *Diagnostic imaging breast*. 2nd ed. Manitoba: Amirsys; 2014, 9:38.
37. Hooley RJ, Greenberg KL, Stackhouse RM, Geisel JL, Butler RS, Philpotts LE. Screening US in patients with mammographically dense breasts: initial experience with Connecticut Public Act 09–41. *Radiology*. 2012;265(1):59–69.
38. Berg WA. Screening and management. In: Berg WA, Yang WT, editors. *Diagnostic imaging breast*. 2nd ed. Manitoba: Amirsys; 2014, 9:44, 46.
39. Yamada T, Mori N, Watanabe M, Kimijima I, Okumoto T, Seiji K, et al. Radiologic-pathologic correlation of ductal carcinoma in situ. *Radiographics*. 2010;30(5):1183–98.
40. Schnall MD, Blume J, Bluemke DA, DeAngelis GA, DeBruhl N, Harms S, et al. Diagnostic architectural and dynamic features at breast MR imaging: multicenter study. *Radiology*. 2006;238(1):42–53.
41. Kuhl CK, Schrading S, Bieling HB, Wardelmann E, Leutner CC, Koenig R, et al. MRI for diagnosis of pure ductal carcinoma in situ: a prospective observational study. *Lancet*. 2007;370(9586):485–92.
42. Bluemke DA, Gatsonis CA, Chen MH, DeAngelis GA, DeBruhl N, Harms S, et al. Magnetic resonance imaging of the breast prior to biopsy. *JAMA*. 2004;292:2735–42.
43. American College of Radiology practice guideline for the performance of contrast-enhanced Magnetic Resonance Imaging (MRI) of the breast: 2013 [http://www.acr.org/~media/ACR/Documents/PGTS/guidelines/MRI\\_Breast.pdf](http://www.acr.org/~media/ACR/Documents/PGTS/guidelines/MRI_Breast.pdf). Accessed 20 Nov 2013.
44. de Bresser J, de Vos B, van der Ent F, Hulsewé K. Breast MRI in clinically and mammographically occult breast cancer presenting with an axillary metastasis: a systematic review. *Eur J Surg Oncol*. 2010;36(2):114–9.
45. Olson Jr JA, Morris EA, Van Zee KJ, Linehan DC, Borgen PI. Magnetic resonance imaging facilitates breast conservation for occult breast cancer. *Ann Surg Oncol*. 2000;7:411–5.
46. Abe H, Schmidt RA, Shah RN, Shimauchi A, Kulkarni K, Sennett CA, et al. MRI-directed (“Second-Look”) ultrasound examination for breast lesions detected initially on MRI: MR and sonographic findings. *Am J Roentgenol*. 2010;194(2):370–7.
47. Meissnitzer M, Dershaw DD, Lee CH, Morris EA. Targeted ultrasound of the breast in women with abnormal MRI findings for whom biopsy has been recommended. *Am J Roentgenol*. 2009;193(4):1025–9.
48. Lehman CD, Gastonis C, Kuhl CK, Hendrick RE, Pisano ED, Hanna L, et al. MRI evaluation of the contralateral breast in women with recently diagnosed breast cancer. *N Engl J Med*. 2007;356:1295–303.
49. Boetes C, Mus RD, Holland R, Barentsz JO, Strijk SP, Wobbes T, et al. Breast tumors: comparative accuracy of MR imaging relative to mammography and US for demonstrating extent. *Radiology*. 1995;197:743–7.
50. Orel SG, Schnall MD, Powell CM, Hochman MG, Solin LJ, Fowble BL, et al. Staging of suspected breast cancer: effect of MR imaging and MR-guided biopsy. *Radiology*. 1995;196:115–22.
51. Liberman L, Morris EA, Dershaw DD, Abramson AF, Tan LK. MR Imaging of the ipsilateral breast in women with percutaneously proven breast cancer. *Am J Roentgenol*. 2003;180:901–10.
52. Berg WA, Gutierrez L, NessAiver MS, Carter WB, Bhargavan M, Lewis RS, et al. Diagnostic accuracy of mammography, clinical examination, US and MR imaging in preoperative assessment of breast cancer. *Radiology*. 2004;233:830–49.
53. Houssami N, Turner R, Macaskill P, Turnbull LW, McCready DR, Tuttle T, et al. An individual person data meta-analysis of preoperative magnetic resonance imaging and breast cancer recurrence. *J Clin Oncol*. 2014;32(5):392–401.
54. Fischer U, Zachariae O, Baum F, von Heyden D, Funke M, Liersch T. The influence of preoperative MRI of the breasts on recurrence rate in patients with breast cancer. *Eur Radiol*. 2004;14(10):1725–31.
55. Turnbull L, Brown S, Harvey I, Olivier C, Drew P, Napp V, et al. Comparative effectiveness of MRI in breast cancer (COMICE) trial: a randomized controlled trial. *Lancet*. 2010;375(9714):563–71.
56. Hakim CM, Ganott MA, Vogia A, Sumkin JH. Breast imaging for pathologists. In: Dabbs DJ, editor. *Breast pathology*. Philadelphia: Saunders (Elsevier); 2012. p. 133.
57. Rosen EL, Blackwell KL, Baker JA, Soo MS, Bentley RC, Yu D, et al. Accuracy of MRI in the detection of residual breast cancer after neoadjuvant chemotherapy. *AJR Am J Roentgenol*. 2003;181(5):1275–82.



58. Yeh E, Slanetz P, Kopans DB, Rafferty E, Georgian-Smith D, Moy L, et al. Prospective comparison of mammography, sonography, and MRI in patients undergoing neoadjuvant chemotherapy for palpable breast cancer. *AJR Am J Roentgenol.* 2005;184(3):868–77.
59. Lobbes MB, Prevost R, Smidt M, Tjan-Heijnen VC, van Goethem M, Schipper R, et al. The role of magnetic resonance imaging in assessing residual disease and pathologic complete response in breast cancer patients receiving neoadjuvant chemotherapy: a systematic review. *Insights Imaging.* 2013;4(2):163–75.
60. Lee JM, Orel SG, Czerniecki BJ, Solin LJ, Schnall MD. MRI before re excision surgery in patients with breast cancer. *AJR Am J Roentgenol.* 2004;182:473–80.
61. ACR Committee on Drugs and Contrast Media. Chapter 13: Nephrogenic systemic fibrosis. *ACR Manual on Contrast Media.* Version 9. 2013. pp 81–9. [http://geiselmed.dartmouth.edu/radiology/pdf/ACR\\_manual.pdf](http://geiselmed.dartmouth.edu/radiology/pdf/ACR_manual.pdf).
62. Haas BM, Kalra V, Geisel J, Raghu M, Durand M, Philpotts LE. Comparison of tomosynthesis plus digital mammography and digital mammography alone for breast cancer screening. *Radiology.* 2013;269(3):694–700.
63. Skaane P, Bandos AI, Gullien R, Eben EB, Ekseth U, Haakenaasen U, et al. Comparison of digital mammography alone and digital mammography plus tomosynthesis in a population based screening program. *Radiology.* 2013;267:47–56.
64. Vreugdenburg TD, Willis CD, Mundy L, Hiller JE. A systematic review of elastography, electrical impedance scanning, and digital infrared thermography for breast cancer screening and diagnosis. *Breast Cancer Res Treat.* 2013;137(3):665–76.
65. Berg WA, Cosgrove DO, Doré CJ, Schäfer FK, Svensson WE, Hooley RJ, et al. Shear-wave elastography improves the specificity of breast US: the BE1 multinational study of 939 masses. *Radiology.* 2012;262(2):435–49.
66. Sun Y, Wei W, Yang HW, Liu JL. Clinical usefulness of breast-specific gamma imaging as an adjunct modality to mammography for diagnosis of breast cancer: a systemic review and meta-analysis. *Eur J Nucl Med Mol Imaging.* 2013;40(3):450–63.
67. Conners AL, Maxwell RW, Tortorelli CL, Hruska CB, Rhodes DJ, Boughey JC, et al. Gamma camera breast imaging lexicon. *AJR Am J Roentgenol.* 2012;199(6):W767–74.
68. Glass SB, Shah ZA. Clinical Utility of positron emission mammography. *Proc (Bayl Univ Med Cent).* 2013;26(3):314–9.
69. Narayanan D, Madsen KS, Kalinyak JE, Berg WA. Interpretation of positron emission mammography: feature analysis and rates of malignancy. *AJR Am J Roentgenol.* 2011;196(4):956–70.
70. Caldarella C, Treglia G, Giordano A. Diagnostic performance of dedicated positron emission mammography using fluorin-18-fluorodeoxyglucose in women with suspicious breast lesions: a meta-analysis. *Clin Breast Cancer.* 2014;14(4):241–8.
71. Berg WA, Madsen KS, Schilling K, Tartar M, Pisano ED, Hovanessian Larsen L, et al. Comparative effectiveness of positron emission mammography and MRI in the contralateral breast of women with newly diagnosed breast cancer. *AJR Am J Roentgenol.* 2012;198:219–32.
72. Berg WA, Madsen KS, Schilling K, Tartar M, Pisano ED, Hovanessian Larsen L, et al. Breast cancer: comparative effectiveness of positron emission mammography and MR imaging in presurgical planning for the ipsilateral breast. *Radiology.* 2011;258(1):59–72.
73. Pisano ED, Fajardo LL, Caudry DJ, Sneige N, Frable WJ, Berg WA, et al. Fine-needle aspiration biopsy of nonpalpable breast lesions in a multicenter clinical trial: results from the radiologic diagnostic oncology group V. *Radiology.* 2001;219(3):785–92.
74. Pisano ED, Fajardo LL, Tsimikas J, Sneige N, Frable WJ, Gatsonis CA, et al. Rate of insufficient samples for fine-needle aspiration for nonpalpable breast lesions in a multicenter clinical trial: the Radiologic Diagnostic Oncology Group 5 Study. The RDOG5 investigators. *Cancer.* 1998;82(4):679–88.
75. Berg WA. Image-guided breast biopsy and management of high-risk lesions. *Radiol Clin North Am.* 2004;42(5):935–46.
76. Youk JH, Kim EK, Kim MJ, Oh KK. Sonographically guided 14-gauge core needle biopsy of breast masses: a review of 2,420 cases with long-term follow-up. *AJR Am J Roentgenol.* 2008;190(1):202–7.

Priscilla McAuliffe

---

## Abstract

Lobular carcinoma in situ (LCIS) is a high-risk indicator lesion for and a non-obligate precursor of the development of invasive breast carcinoma. Loss of E-cadherin is the hallmark pathological feature of lobular entities. Effective clinical management of LCIS requires good communication between the radiologist, surgeon, pathologist, and medical oncologist and entails surgical excision, subsequent surveillance, and systemic and surgical strategies to reduce the risk of future invasive cancer.

---

## Keywords

Lobular carcinoma in situ • Tamoxifen • LCIS • Chemoprevention • Raloxifen • E-cadherin

---

## History

Lobular carcinoma in situ (LCIS) was first described in the 1940s [1]. LCIS was first treated similarly to invasive carcinoma—with radical mastectomy—because it was often diagnosed concurrently with invasive lobular carcinoma (ILC). It was subsequently recognized that LCIS

is a marker of risk for breast cancer that does not itself progress to malignancy, and treatment has thus evolved to close observation with early detection of subsequent malignancy. This management change was based in part on a 1978 review of 211 cases of women with LCIS treated by observation alone (without surgery). There was a 17 % incidence of subsequent invasive carcinoma, with equivalent risk in both breasts, and only six (3 %) patients died of breast cancer [2]. Close observation was associated with early breast cancer detection and high associated cure rates. However, more recently, as mammography and image-guided needle biopsies have become more widespread, the biological heterogeneity of LCIS has become more apparent, and now certain subtypes of LCIS, including the pleomorphic

---

P. McAuliffe, MD, PhD, FACS  
Department of Surgery, Division of Surgical  
Oncology, Magee-Womens Hospital,  
University of Pittsburgh Medical Center,  
300 Halket Street, Suite 2601, Pittsburgh,  
PA, 15213, USA

Magee-Women's Surgical Associates,  
Bethel Park, PA, USA  
e-mail: [mcauliffe@upmc.edu](mailto:mcauliffe@upmc.edu)

variant, are recognized as indolent precursors of ILC for which surgical resection with negative margins and often radiation therapy is indicated.

---

## Epidemiology

The incidence of LCIS is difficult to estimate because it lacks specific clinical abnormalities and is always identified incidentally [3]. LCIS is generally not detectable by palpation on physical exam, by mammogram, or by gross pathological examination [4]. LCIS is identified in 0.5–3.9 % of breast biopsy specimens [5, 6].

The mean age at diagnosis of LCIS is 10–15 years younger than that for invasive breast cancer. It has been described as being more common in premenopausal than in postmenopausal women [2, 7]. However, while LCIS is more often diagnosed in women between age 40 and 50, a review of the Surveillance, Epidemiology and End Results (SEER) program database from 1978 to 1998 revealed that LCIS increased during that time period in all age groups [5]. Interestingly, in women older than age 50, the incidence of LCIS increased concurrently with the incidence of ILC, whereas in women younger than 50, an increase in ILC was not observed as LCIS increased. In women aged 40–49 years old, rising LCIS diagnoses leveled off at approximately 1989, whereas the increase of LCIS in women aged 50–79 years old was the most profound and sustained. The reason for this increase in LCIS in postmenopausal women is likely multifactorial, including the increased availability of screening mammography, the implementation of MRI in breast cancer patient management, the use of hormone replacement therapy in postmenopausal women, and more accurate molecular diagnosis, to be discussed in the pathology section.

---

## Risk Assessment

Patients with LCIS have an 8- to 12-fold greater lifetime risk than the general population for developing invasive breast cancer in either breast

[8, 9]. Numerous studies have documented that after the diagnosis of LCIS, if diligently sought, LCIS can be found elsewhere in the index breast and also in the contralateral breast. Approximately 50 % of LCIS is multifocal, and in 30 % of patients, LCIS is found within the contralateral breast [2, 9]. However, despite the bilateral risk, cancer development is skewed toward the ipsilateral breast. Furthermore, although subsequent invasive breast cancer can be either of ductal or lobular origin, 70–89 % of invasive carcinoma after LCIS is ILC [9, 10]. The time between LCIS and invasive cancer development is approximately 15–30 years [9]. LCIS is associated with approximately 90 % of ILC cases [11].

---

## Pathology

The hallmark of LCIS is the proliferation of the epithelial cells of the terminal ductal-lobular unit, with no penetration of the basement membrane. Compared to the cells that normally line the lobular acini, LCIS cells are larger and monomorphic. There is also a loss of cellular cohesion and the presence of intra-cytoplasmic vacuoles. Mitoses and necrosis are infrequent, and nucleoli are inconspicuous, without prominent chromatin. The difference between LCIS and the high-risk lesion atypical lobular hyperplasia (ALH) is quantitative, with fewer abnormal cells and the preservation of residual lumen in the lobules with ALH compared to complete replacement of the lobular unit with LCIS. Many utilize the term “lobular neoplasia” to encompass both ALH and LCIS because they may represent early and later points on a spectrum of abnormal lobular proliferation [12]. LCIS is distinguished from ILC because it is contained by the basement membrane on hematoxylin and eosin-stained sections. Cases of mixed lobular and ductal in situ lesions have also been described, with genetic aberrations of a hybrid phenotype [12].

The pleomorphic variant of LCIS (PLCIS) is architecturally similar to LCIS. However, PLCIS has substantially larger nuclei and greater nuclear polymorphisms. In contrast to classic LCIS, PLCIS has prominent nucleoli, central necrosis,

and large, clustered calcifications. In some cases, PLCIS cells have eosinophilic cytoplasm, imparting an “apocrine appearance,” or intracytoplasmic vacuoles, imparting a “signet ring cell appearance” [12]. Her2/neu overexpression and gene amplification have been reported in PLCIS with apocrine differentiation [13]. The combination of calcifications, necrosis, and cellular features can complicate the distinction of PLCIS from high-grade DCIS. Whereas classic LCIS is generally not associated with direct clonal progression to ILC, the pleomorphic variant lesions are. These data suggest that pleomorphic LCIS may not only be a marker for increased risk of invasive breast cancer but also a direct precursor of ILC. Classic and pleomorphic LCIS can coexist in the same lesion [14].

Molecular analyses of LCIS (as well as ALH and ILC) have revealed decreased expression or the loss of the cell surface adhesion molecule E-cadherin [15]. The loss of E-cadherin is the defining molecular event of lobular breast pathology. This contrasts with ductal lesions, in which E-cadherin expression is generally maintained. Immunohistochemistry using anti-E-cadherin antibodies can be used to distinguish ductal and lobular lesions.

E-cadherin is the protein product of the *CDH1* gene (16q22.1) and is expressed on epithelial cells [12]. The cadherins are a family of adhesion proteins that span the cell membrane and, through a calcium-dependent mechanism, form dimers with cadherins on other cells and interact with the actin cytoskeleton [12]. The portion of E-cadherin that is intracytoplasmic binds to p120-catenin [16]. In normal mammary cells, p120-catenin is present at the cell membrane. However, if the E-cadherin protein is nonfunctional or lost, p120 accumulates in the cytoplasm, where it activates cytoplasmic Rho-GTPases, resulting in increased cell motility [17]. The loss of E-cadherin and the cytoplasmic accumulation of p120-catenin is pathognomonic for lobular breast pathologies [12]. This feature can be critically important when LCIS is diagnosed concurrently with lesions, such as sclerosing adenosis or radial scars, as these together can produce patterns that mimic ILC. The lack of E-cadherin staining and

cytoplasmic p120 catenin in the areas of question can differentiate LCIS and ILC [12]. Furthermore, some high-grade triple-negative DCIS may display diminished E-cadherin expression, suggesting PLCIS [12]. In addition to the loss of E-cadherin, the loss of high-molecular-weight keratins (cytokeratin 5/6, 14, and 17), which are generally present in high-grade DCIS, suggests PLCIS [12].

Some LCIS may display aberrant E-cadherin membrane expression that is not completely absent from the cell membranes, but it is fragmented, focal, or beaded. In these cases, double staining for E-cadherin and immunostaining for beta-catenin can be helpful to establish the diagnosis. The loss of beta-catenin also indicates that the E-cadherin is dysfunctional and not associated with other molecules in the cadherin-catenin complex [18, 19].

*CDH1* gene mutations, deletions, and methylation have been identified in LCIS, as well as abnormal transcriptional regulation of E-cadherin [12]. Furthermore, LCIS also exhibits a loss of heterozygosity [20]. Other target genes that have been associated with the development of LCIS include fibroblast growth factor receptor 1 (*FGFR1*) and cyclin D1 (*CCND1*) [21, 22]. Pleomorphic LCIS has also been associated with *CCND1* and the oncogenes *MYC* and *HER2* [13, 23].

---

## Diagnosis

### Clinical Presentation

The clinical presentation of patients with LCIS is highly variable. LCIS is usually not detectable by physical examination and does not have pathognomonic features on mammography. In the era of widespread mammographic screening and the shift to percutaneous breast biopsy, LCIS is most commonly diagnosed as an incidental finding on image-guided core-needle biopsy. It can also be found incidentally on surgical lumpectomy specimens removed for another indication.

Radiographically, classical LCIS is associated with small punctate calcifications in 42 % of

cases, whereas the pleomorphic variant of LCIS is more likely to have large and clustered calcifications related to the presence of comedo-type necrosis [4]. Pathological diagnosis is described above. Occasionally, even in the presence of E-cadherin, p120-catenin, beta-catenin, and cytokeratin staining, the diagnosis of LCIS is ambiguous and difficult to distinguish from DCIS. In this case, diagnosis should employ a multidisciplinary approach. However, when a definitive diagnosis cannot be rendered even after a multidisciplinary discussion or in the case of mixed LCIS and DCIS, the lesion should be managed as DCIS.

---

## Treatment

### Surgery

After an incidental diagnosis of LCIS by percutaneous image-guided core-needle biopsy, surgical excisional biopsy should be performed to rule out synchronous invasive cancer and DCIS. Percutaneous biopsy is limited by sampling error, and it can present difficulty in making a definitive histological diagnosis [24]. Upgrading to invasive cancer when the biopsy site is surgically excised can occur [25]. The goal of surgical excisional biopsy is to remove the biopsy site and any residual imaging abnormalities.

Excisional biopsy demonstrates a 0–10 % risk of synchronous invasive breast cancer and a 0–50 % risk of synchronous DCIS [6, 26, 27]. Surgical excisional biopsy is most commonly performed using a technique to localize a titanium marker clip placed radiographically during percutaneous biopsy. Two such localization techniques are wire or radioactive seed localization. To document the removal of the LCIS on excisional biopsy, mammography of the surgical specimen after excision should reveal the presence of the clip. Furthermore, the surgical pathology report should describe residual biopsy site changes due to the percutaneous core-needle biopsy. Contralateral mirror-image breast biopsy, a procedure described in the past for patients with LCIS, is no longer performed. Instead, close

observation of all remaining breast tissue is recommended.

The management of microscopic margin status in LCIS is guided by the results of several studies described below. In a study of 180 patients who underwent observation alone after margin-negative surgical excision of LCIS, the overall ipsilateral and contralateral breast cancer event rates at 12 years of follow-up were 14.4 % and 7.8 %, respectively [10]. The rate of invasive breast cancer was 5.6 %. This rate was similar whether ipsilateral or contralateral, although contralateral cancers occurred later. Nearly 85 % of subsequent ipsilateral breast tumors were detected mammographically. More than 96 % of all ipsilateral tumors occurred in the same quadrant as the original LCIS. Breast cancer-specific mortality was 1.1 % at 12 years [10]. In another study of 100 patients with LCIS in which margin status was not documented, the ipsilateral and overall breast cancer event rates were 13 % and 16 %, respectively [28]. Finally, in a retrospective analysis of 2,894 patients who underwent breast-conserving surgery for DCIS or early breast cancer between 1980 and 2007, 10 % had LCIS within the lumpectomy specimen, and of those, approximately one-third had LCIS at the margin [29]. The difference in crude local recurrence rate between the patients with LCIS within the specimen (4.5 %) and in those with no LCIS (3.8 %) was not statistically significant [29]. Furthermore, there was also no significant difference in actuarial 5- and 10-year local recurrence rates if LCIS was present at the margin (6 % and 6 %), if LCIS was present but not at the margin (1 % and 15 %), or if no LCIS was present at all (2 % and 6 %). The results of these studies suggest that re-excision to achieve negative margins for classical LCIS is not warranted. However, for the pleomorphic PLCIS subtype, re-excision to achieve negative margins is indicated. In addition, identification of LCIS in a lumpectomy specimen resected for the diagnosis of DCIS or invasive cancer should not alter surgical management of the primary breast because the presence of LCIS does not increase the rate of in-breast recurrence in patients undergoing breast conservation [29].



Once a diagnosis of LCIS has been rendered and concurrent malignancy excluded, patients with LCIS should be counseled regarding their increased lifetime risk of breast cancer development. The surgical management of LCIS is generally conservative, and only a small minority pursue bilateral risk-reducing mastectomy, although this number has recently been increasing [30]. This approach is usually reserved for patients who have additional risk factors for breast cancer development or who experience significant anxiety regarding observation and/or chemoprevention options. It is important that patients considering this option are aware that bilateral mastectomy does not completely eliminate the risk of breast cancer development [31]. Because LCIS poses no risk of regional metastasis, sentinel lymph node biopsy or axillary node dissection is not required. Immediate breast reconstruction should be offered for patients who undergo risk-reducing mastectomy for LCIS. Women should be informed about the impact of this treatment approach on quality of life, particularly body image and sexual function [32]. Nipple-areola complex-sparing mastectomy may be a viable option in carefully selected women pursuing surgical risk reduction [33].

### **Risk-Reducing Endocrine Therapy**

Risk-reducing therapy, often called “chemoprevention,” is an important treatment option for patients with LCIS. In the NSABP P-1 breast cancer prevention trial, the incidence of invasive breast cancers was reduced by 56 % in women with LCIS who received tamoxifen compared to observation alone [34]. Women with LCIS represented 6.2 % of the patients in that trial. The annual hazard rate of invasive cancer was 5.69 per 1,000 women who received tamoxifen compared with 12.99 per 1,000 women who did not. In the NSABP P-2 trial, postmenopausal women with LCIS were randomized to tamoxifen or raloxifene [35]. Women with LCIS comprised 9.2 % of the patients on the trial. There was no difference in risk reduction for invasive breast cancer between the two agents (incidence 4.30

per 1,000 vs. 4.41 per 1,000 for tamoxifen and raloxifene, respectively). Patients receiving raloxifene had a lower incidence of thromboembolic events and cataracts. There was no significant difference in the risk of other cancers, fractures, ischemic heart disease, or stroke for the two drugs. At 81 months of median follow-up, raloxifene was 78 % as effective as tamoxifen at preventing invasive disease but had fewer toxicities, with significantly fewer endometrial cancers [36]. Raloxifene may be of particular benefit to postmenopausal women with an intact uterus and a risk of osteoporosis; tamoxifen would be an appropriate choice for high-risk postmenopausal women.

### **Radiation Therapy**

Adjuvant radiation therapy is not recommended for the treatment of LCIS. If synchronous DCIS or invasive breast cancer is found in an excised LCIS specimen, the patient should be treated according to the guidelines for DCIS or invasive breast cancer and may benefit from radiation.

---

### **Surveillance**

Following excisional biopsy demonstrating LCIS, patients should undergo annual bilateral breast physical examinations and diagnostic mammography. Screening ultrasound in patients with high breast cancer risk, including LCIS, is associated with high false-positive results [37]. A recent single-institution analysis revealed that with either annual mammograms or MRI, the cancer detection rate was 13 % [38]. MRI was not associated with diagnosing breast cancer at earlier stage, smaller size, or node negativity. For this reason, the routine use of MRI for screening patients with a diagnosis of LCIS is not recommended. Patients with LCIS who undergo a bilateral mastectomy with or without reconstruction should also undergo an annual physical examination, but routine imaging is not indicated. Any suspicious lesions should be evaluated with ultrasound and biopsy analysis.

## Conclusion

LCIS is a histological finding characterized by an intact basement membrane with a loss of E-cadherin leading to a dysfunctional E-cadherin/catenin complex. LCIS confers increased long-term risk of breast cancer that may affect either breast. The pleomorphic subtype is also a non-obligate precursor to invasive cancer. Patients found to have LCIS on core-needle biopsy are evaluated with bilateral diagnostic imaging, and additional suspicious lesions are further evaluated. Marker clips should routinely be placed at the time of percutaneous image-guided biopsy. Patients diagnosed with LCIS should undergo surgical excisional biopsy with localization of the percutaneous biopsy cavity to increase accuracy. If synchronous DCIS or invasive breast cancer is diagnosed, subsequent treatment is administered according to the guidelines for these tumors. Re-excision to attain negative margins is not performed in patients with classical LCIS unless pleomorphic LCIS is identified, in which case negative margins should be achieved. Bilateral risk-reducing mastectomy is generally reserved for patients with additional risk factors for breast cancer or with extreme anxiety regarding observation and/or chemoprevention options but does not completely eradicate the risk of subsequent breast cancer development. Patients with LCIS should receive systemic risk reduction with anti-estrogen therapy, namely, tamoxifen or raloxifene. Follow-up includes clinical and imaging surveillance. All patients with LCIS should be considered for clinical trials.

## References

1. Foote FW, Stewart FW. Lobular carcinoma in situ: a rare form of mammary cancer. *Am J Pathol.* 1941;17(4):491–6.
2. Haagensen CD, Lane N, Lattes R, Bodian C. Lobular neoplasia (so-called lobular carcinoma in situ) of the breast. *Cancer.* 1978;42(2):737–69.
3. Karabakhtsian RG, Johnson R, Sumkin J, Dabbs DJ. The clinical significance of lobular neoplasia on breast core biopsy. *Am J Surg Pathol.* 2007;31(5):717–23.
4. Georgian-Smith D, Lawton TJ. Calcifications of lobular carcinoma in situ of the breast: radiologic-pathologic correlation. *AJR Am J Roentgenol.* 2001;176(5):1255–9.
5. Li CI, Daling JR, Malone KE. Age-specific incidence rates of in situ breast carcinomas by histologic type, 1980 to 2001. *Cancer Epidemiol Biomarkers Prev.* 2005;14(4):1008–11.
6. Degnim AC, King TA. Surgical management of high-risk breast lesions. *Surg Clin North Am.* 2013;93(2):329–40.
7. Beute BJ, Kalisher L, Hutter RV. Lobular carcinoma in situ of the breast: clinical, pathologic, and mammographic features. *AJR Am J Roentgenol.* 1991;157(2):257–65.
8. Frykberg ER. Lobular carcinoma in situ of the breast. *Breast J.* 1999;5(5):296–303.
9. Page DL, Kidd Jr TE, Dupont WD, Simpson JF, Rogers LW. Lobular neoplasia of the breast: higher risk for subsequent invasive cancer predicted by more extensive disease. *Hum Pathol.* 1991;22(12):1232–9.
10. Fisher ER, Land SR, Fisher B, Mamounas E, Gilarski L, Wolmark N. Pathologic findings from the National Surgical Adjuvant Breast and Bowel Project: twelve-year observations concerning lobular carcinoma in situ. *Cancer.* 2004;100(2):238–44.
11. Abdel-Fatah TM, et al. High frequency of coexistence of columnar cell lesions, lobular neoplasia, and low grade ductal carcinoma in situ with invasive tubular carcinoma and invasive lobular carcinoma. *Am J Surg Pathol.* 2007;31(3):417–26.
12. Dabbs DJ, Powe DG, Hodi Z, Lee AH, Reis-Filho JS, Ellis IO. Lobular neoplasia of the breast revisited with emphasis on the role of E-cadherin immunohistochemistry. *Am J Surg Pathol.* 2013;37(7):e1–11.
13. Chen YY, Hwang ES, Roy R, DeVries S, Anderson J, Wa C, et al. Genetic and phenotypic characteristics of pleomorphic lobular carcinoma in situ of the breast. *Am J Surg Pathol.* 2009;33(11):1683–94.
14. Sneige N, Wang J, Baker BA, Krishnamurthy S, Middleton LP. Clinical, histopathologic, and biologic features of pleomorphic lobular (ductal-lobular) carcinoma in situ of the breast: a report of 24 cases. *Mod Pathol.* 2002;15(10):1044–50.
15. Chivukula M, Haynik DM, Brufsky A, Carter G, Dabbs DJ. Pleomorphic lobular carcinoma in situ (PLCIS) on breast core needle biopsies: clinical significance and immunoprofile. *Am J Surg Pathol.* 2008;32(11):1721–6.
16. Hatzfeld M. The p120 family of cell adhesion molecules. *Eur J Cell Biol.* 2005;84(2–3):205–14.
17. Reynolds AB, Rocznik-Ferguson A. Emerging roles for p120-catenin in cell adhesion and cancer. *Oncogene.* 2004;23(48):7947–56.
18. Geyer FC, Lacroix-Triki M, Savage K, Arnedos M, Lambros MB, MacKay A, et al. beta-Catenin pathway activation in breast cancer is associated with triple-negative phenotype but not with CTNNB1 mutation. *Mod Pathol.* 2011;24(2):209–31.
19. Rakha EA, Patel A, Powe DG, Benhasouna A, Green AR, Lambros MB, et al. Clinical and biological

- significance of E-cadherin protein expression in invasive lobular carcinoma of the breast. *Am J Surg Pathol.* 2010;34(10):1472–9.
20. Lakhani SR, Collins N, Sloane JP, Stratton MR. Loss of heterozygosity in lobular carcinoma in situ of the breast. *Clin Mol Pathol.* 1995;48(2):M74–8.
  21. Natrajan R, Weigelt B, Mackay A, Geyer FC, Grigoriadis A, Tan DS, et al. An integrative genomic and transcriptomic analysis reveals molecular pathways and networks regulated by copy number aberrations in basal-like, HER2 and luminal cancers. *Breast Cancer Res Treat.* 2010;121(3):575–89.
  22. Reis-Filho JS, Simpson PT, Turner NC, Lambros MB, Jones C, Mackay A, et al. FGFR1 emerges as a potential therapeutic target for lobular breast carcinomas. *Clin Cancer Res.* 2006;12(22):6652–62.
  23. Simpson PT, Reis-Filho JS, Lambros MB, Jones C, Steele D, Mackay A, et al. Molecular profiling pleomorphic lobular carcinomas of the breast: evidence for a common molecular genetic pathway with classic lobular carcinomas. *J Pathol.* 2008;215(3):231–44.
  24. Corben AD, Edelweiss M, Brogi E. Challenges in the interpretation of breast core biopsies. *Breast J.* 2010;16 Suppl 1:S5–9.
  25. Houssami N, Ciatto S, Ellis I, Ambrogetti D. Underestimation of malignancy of breast core-needle biopsy: concepts and precise overall and category-specific estimates. *Cancer.* 2007;109(3):487–95.
  26. Arpino G, Allred DC, Mohsin SK, Weiss HL, Conrow D, Elledge RM. Lobular neoplasia on core-needle biopsy—clinical significance. *Cancer.* 2004;101(2):242–50.
  27. Hussain M, Cunnick GH. Management of lobular carcinoma in-situ and atypical lobular hyperplasia of the breast—a review. *Eur J Surg Oncol.* 2011;37(4):279–89.
  28. Ottesen GL, Graversen HP, Blichert-Toft M, Christensen IJ, Andersen JA. Carcinoma in situ of the female breast. 10 year follow-up results of a prospective nationwide study. *Breast Cancer Res Treat.* 2000;62(3):197–210.
  29. Ciocca RM, Li T, Freedman GM, Morrow M. Presence of lobular carcinoma in situ does not increase local recurrence in patients treated with breast-conserving therapy. *Ann Surg Oncol.* 2008;15(8):2263–71.
  30. Portschy PR, Marmor S, Nzara R, Virnig BA, Tuttle TM. Trends in incidence and management of lobular carcinoma in situ: a population-based analysis. *Ann Surg Oncol.* 2013;20(10):3240–6.
  31. Lostumbo L, Carbine NE, Wallace J. Prophylactic mastectomy for the prevention of breast cancer. *Cochrane Database Syst Rev.* 2010;(11):CD002748.
  32. Frost MH, Hoskin TL, Hartmann LC, Degnim AC, Johnson JL, Boughhey JC. Contralateral prophylactic mastectomy: long-term consistency of satisfaction and adverse effects and the significance of informed decision-making, quality of life, and personality traits. *Ann Surg Oncol.* 2011;18(11):3110–6.
  33. Peled AW, Duralde E, Foster RD, Fiscali AS, Esserman LJ, Hwang ES, et al. Patient-reported outcomes and satisfaction after total skin-sparing mastectomy and immediate expander-implant reconstruction. *Ann Plast Surg.* 2014;72 Suppl 1:S48–52.
  34. Fisher B, Costantino JP, Wickerham DL, Redmond CK, Kavanah M, Cronin WM, et al. Tamoxifen for prevention of breast cancer: report of the National Surgical Adjuvant Breast and Bowel Project P-1 Study. *J Natl Cancer Inst.* 1998;90(18):1371–88.
  35. Vogel VG, Costantino JP, Wickerham DL, Cronin WM, Cecchini RS, Atkins JN, et al. Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: the NSABP Study of Tamoxifen and Raloxifene (STAR) P-2 trial. *JAMA.* 2006;295(23):2727–41.
  36. Vogel VG, Costantino JP, Wickerham DL, Cronin WM, Cecchini RS, Atkins JN, et al. Update of the National Surgical Adjuvant Breast and Bowel Project Study of Tamoxifen and Raloxifene (STAR) P-2 Trial: preventing breast cancer. *Cancer Prev Res (Phila).* 2010;3(6):696–706.
  37. Berg WA, Zhang Z, Lehrer D, Jong RA, Pisano ED, Barr RG, et al. Detection of breast cancer with addition of annual screening ultrasound or a single screening MRI to mammography in women with elevated breast cancer risk. *JAMA.* 2012;307(13):1394–404.
  38. King TA, Muhsen S, Patil S, Koslow S, Oskar S, Park A, et al. Is there a role for routine screening MRI in women with LCIS? *Breast Cancer Res Treat.* 2013;142(2):445–53.

Priscilla McAuliffe

---

### Abstract

Ductal carcinoma in situ (DCIS) is defined as abnormally proliferating malignant cells confined to the breast milk ducts by the basement membrane. The prognosis of DCIS with microinvasion (DCISM) is intermediate between those of DCIS and invasive ductal cancer. DCIS is diagnosed most commonly as a mammographic abnormality but can occasionally present as a palpable breast mass. Therapy for DCIS is personalized for each patient based on tumor-to-breast size ratio, mammographic appearance, margin width, and patient preference. The benefits and risks of breast-conserving surgery versus mastectomy should be discussed in detail. Overall survival after breast-conserving therapy is equivalent to mastectomy. Re-excision is recommended for positive margins and, although debated, pathologic margins less than 2 mm. For breast conservation therapy, radiation therapy is recommended to reduce the risk of local recurrence. Mastectomy is indicated in patients with persistently positive margins after attempts at breast conservation, a contraindication to radiation therapy, or diffuse, malignant-appearing calcifications or whose anxiety about local recurrence outweighs the impact of mastectomy on quality of life. Patients undergoing mastectomy for DCIS should have sentinel lymph node biopsy. Immediate breast reconstruction should be considered for patients undergoing mastectomy. Endocrine therapy such as tamoxifen is offered for 5 years to women with estrogen receptor-positive DCIS. Patients with DCIS should have annual clinical breast exams and annual mammographic imaging of all remaining breast tissue.

---

P. McAuliffe, MD, PhD, FACS  
Department of Surgery, Division of Surgical  
Oncology, Magee-Womens Hospital,  
University of Pittsburgh Medical Center,  
300 Halket Street, Suite 2601, Pittsburgh,  
PA, 15213, USA  
e-mail: [mcauliffe@upmc.edu](mailto:mcauliffe@upmc.edu)

---

**Keywords**

Ductal carcinoma in situ • DCIS • Microinvasion

---

**History**

Before the introduction of screening mammography, most cases of ductal carcinoma in situ (DCIS) remained undetected until a palpable mass formed. However, the widespread use of screening mammography has resulted in a tenfold increase in the reported incidence of DCIS since the 1980s [1]. The Surveillance, Epidemiology, and End Results (SEER) Program database indicates that in 1975, 5.8 per 100,000 women were diagnosed with in situ breast cancer in the United States, whereas in 2007, 34.9 per 100,000 women received the same diagnosis [2].

---

**Epidemiology**

DCIS accounts for more than 25 % of all new cases of breast cancer [1]. Approximately 64,640 new diagnoses of in situ breast cancer are expected among US women in 2013, more than 85 % of which will be DCIS [1]. The median age of diagnosis of DCIS ranges from 47 to 63 years, but more than 75 % of patients will receive this diagnosis after age 50 [1]. In the United States, in women between the ages of 50 and 69, 1 case of DCIS is detected per 1,000 screening mammograms [3]. The diagnosis of DCIS peaks between ages 60 and 74, earlier than for invasive ductal carcinoma (IDC), the diagnosis of which peaks between ages 75 and 79 [4]. Many risk factors for the development of DCIS are similar to those for IDC, including female sex, older age, family history of breast cancer, BRCA 1/2 gene carriers, increased breast density, a history of previous breast biopsy, nulliparity, or older age at the time of first full-term pregnancy [4, 5]. However, studies have been inconsistent with respect to the effect of more than 5 years to estrogen hormone replacement therapy or body mass index, both

known risk factors for the development of IDC, on the risk of DCIS [4]. Oral contraceptive pills do not appear to increase the risk of DCIS [5].

**Natural History of DCIS**

Several strong pieces of evidence suggest that not all DCIS lesions are clinically significant. The prevalence of DCIS on autopsy is 1–14.3 % [6, 7]. Furthermore, in studies in which DCIS was initially misdiagnosed as benign and treated by biopsy alone, 14–53 % of DCIS progressed to IDC over a period of 10 or more years [6]. This suggests that there is a cohort of patients for whom DCIS would not have had a clinical impact on the patient's life. This underscores the critical need for research to identify markers that can aid the selection of patients for personalized treatment [5]. The development of molecular risk profiles is of particular interest [8]. Furthermore, this has also led to a discussion about the best terminology for DCIS and the possibility of omitting the term “carcinoma” [9].

---

**Pathology**

DCIS is a proliferation of malignant cells arising from the ductal epithelium in the terminal ductal-lobular unit that has not breached the ductal basement membrane. Malignant cells proliferate until the ductal lumen is obliterated. DCIS has traditionally been considered one stage in the continuum of progression from atypical ductal hyperplasia (ADH) to IDC [10]. The lesions can be heterogeneous, with variable histologic architecture, molecular and cellular characteristics, and clinical behavior. DCIS has also been associated with changes in the surrounding stroma, resulting in fibroblast proliferation, lymphocyte



infiltration, and angiogenesis [10]. Although the process is poorly understood, it is believed that most IDCs arise from DCIS, and poorly differentiated DCIS evolve from well-differentiated DCIS [11]. The genetic defects that lead to progression appear to be randomly acquired and accelerated by p53 mutation, which results in genetic instability [11]. Furthermore, genes involved in cell adhesion and signaling, motility, angiogenesis, and extracellular matrix formation that may lead to progression from DCIS to IDC have been identified [12, 13]. DCIS may represent a later stage of molecular progression because many gene mutations occur prior to invasion [14, 15].

## Classification

No single classification system for DCIS has been universally accepted. The most common subtypes based on the architectural pattern of the proliferating cells—comedo, solid, cribriform, micropapillary, and papillary—can coexist. Nuclear grade has also been used to classify lesions. DCIS with high nuclear grade and comedo necrosis is predictive of local recurrence [16]. At 8 years of follow-up, patients whose tumors had a high nuclear grade and comedo necrosis had a 20 % local recurrence rate after breast conservation surgery and irradiation, compared with 5 % for those patients whose tumors did not have necrosis and had a lower nuclear grade [16]. A classification system devised by Silverstein called the Van Nuys classification also utilized high nuclear grade and comedo-type necrosis. Their early studies suggested that this classification was strongly predictive of disease-free survival, but this utility could not be prospectively validated [17].

## Multifocality

Multifocal DCIS is defined as the presence of DCIS in two or more foci in the same breast quadrant, separated by 5 mm. Careful serial pathological subsectioning of multifocal lesions suggests that these actually represent intraductal

spread from a single focus of DCIS. In 81 of 82 mastectomy specimens, multifocal lesions that appeared to be separate using conventional pathological techniques were found to originate from the same focus [18].

## Multicentricity

Multicentric DCIS presents as separate, discontinuous foci of DCIS outside of the index breast quadrant. The incidence of multicentricity varies in the literature and likely depends on the extent of imaging and pathological review. Most local recurrences after DCIS treatment occur in the same quadrant as the index lesion, implicating residual untreated disease rather than multicentricity [19].

## Microinvasion

According to the American Joint Committee on Cancer (AJCC), microinvasion is defined as the invasion of breast cancer cells through the basement membrane at one or more foci, none of which exceeding a dimension of 1 mm. In the AJCC staging system, DCIS with microinvasion (DCISM) is classified as a “T1mic” tumor, whereas DCIS is classified as “T0.” Microinvasion upstages the AJCC cancer stage from 0 to 1. DCISM is observed in 5–10 % of cases of DCIS [20]. By definition, DCIS does not metastasize to axillary lymph nodes or distant sites, in contrast to DCISM. Axillary metastasis has been reported in 0 to 28 % of patients with DCISM [21, 22].

Microinvasion in DCIS varies according to the size and extent of the index lesion. When DCIS less than 25 mm in diameter was compared to those 25 mm and larger, the incidence of microinvasion was 2 % and 29 %, respectively [16]. The incidence of microinvasion is also higher in high-grade or comedo-type DCIS and when DCIS presents as a palpable mass or with nipple discharge [20].

Disease-specific survival is worse for DCISM than DCIS [23]. In a retrospective study of 1,248 cases of DCIS, the 10-year distant metastasis-free survival rate was significantly

higher in patients with DCIS compared to DCISM (98 % and 91 %, respectively). The overall survival rate was also higher (96.5 % vs. 88.4 %) [22]. However, when compared to IDC, metastasis-free and overall survival rates were higher in patients with DCISM. These results suggest that DCISM should be characterized as an invasive tumor with a good outcome, and the therapeutic approach for these patients should be similar to that for patients with IDC. Further study is needed to investigate the biology of microinvasion.

## Diagnosis

### Clinical Presentation

Prior to routine screening mammography, patients with DCIS presented most commonly with a palpable mass, nipple thickening or discharge, or Paget's disease of the nipple. Occasionally, DCIS was found incidentally in an otherwise benign breast biopsy specimen. With the advent of screening mammography, DCIS is more likely to be diagnosed when the tumor is still clinically occult. Patients with mammographically detected DCIS should always undergo contralateral breast imaging because patients may have synchronous occult abnormalities or cancers in the contralateral breast. To establish interval changes, current images should be compared with previous mammograms, if available. Magnification, spot compression, and other mammographic views are routinely used to further delineate the abnormality, particularly calcifications, in the index breast. Ultrasound can also be used to assess tumor size and multicentricity.

## Imaging

### Mammographic and Ultrasonographic Features

On a mammogram, DCIS can present as microcalcifications, a soft-tissue density/asymmetry, or both. Microcalcifications are

the most common mammographic manifestation of DCIS. Microcalcifications can be divided into two classes suggestive of the architectural type of DCIS: (1) linear branching type, which are more often associated with high nuclear-grade comedo-type lesions, and (2) fine, granular type, which are generally associated with micropapillary or cribriform lesions of lower nuclear grade without necrosis [18]. Mammographic findings can significantly underestimate the pathological extent of disease, particularly in cases of micropapillary DCIS [18]. Lesions were more than 2 cm larger by histological examination than by mammographic estimation in 44 % of cases of micropapillary lesions, compared with only 12 % of cases of the pure comedo subtype. Magnification views on mammography more accurately predict the extent of disease, which was underestimated in only 14 % of cases of micropapillary tumors. Hence, magnification views increase image resolution and are better able to discern the shape, number, and extent of calcifications when compared with screening mammographic films, and they should be used routinely in the evaluation of suspicious mammographic findings, particularly microcalcifications.

### Magnetic Resonance Imaging (MRI)

MRI is not routinely employed in the preoperative evaluation of patients with DCIS. Instead, mammogram remains the gold standard for radiographic evaluation of DCIS. The cost and accessibility of MRI make it less feasible as a screening method. However, it may have utility in patients at high risk for breast cancer or with extremely dense breasts. Contrast-enhanced MRI is more sensitive than mammography in detecting both DCIS and invasive cancer, but because DCIS can mimic fibrocystic change and other benign findings on MRI, it can also lead to false positives and unnecessary biopsies [24–27]. MRI is sometimes used after the initial diagnosis to identify multicentric and contralateral lesions because the presence of either of these may alter the surgical treatment strategy [28]. MRI-detected multicentric disease was detected in 4.3 % of 149 patients who presented with DCIS [29]. MRI can also detect contralateral breast

cancer in patients presenting with DCIS [28]. Of 196 patients with DCIS, MRI prompted biopsy in 18 patients, and contralateral breast cancer was detected in 5 patients (28 % of those biopsied and 2.6 % of those with DCIS). The sensitivity and specificity of detecting contralateral breast cancer are 71 % and 90 %, respectively. However, no benefit of MRI in reducing local recurrence has been observed, most likely because MRI cannot detect all clinically occult cancer in the breast, and, in addition, local failure after breast conservation is uncommon in contemporary studies [30, 31]. Finally, acting on the findings of MRI leads to the exclusion of patients from breast conservation therapy and an increase in the performance of mastectomies [26, 32]. Therefore, breast MRI likely leads to overtreatment [30].

### Diagnostic Biopsy

The preferred method for the diagnosis of DCIS is percutaneous biopsy, either with the ultrasound-guided or vacuum-assisted stereotactic core-needle biopsy technique. Patients who cannot lie prone, exceed the weight limit for the stereotactic system, or cannot cooperate during the procedure are not good candidates for stereotactic biopsy. Bleeding disorders and the concomitant use of anticoagulants are relative contraindications. Patients who are not candidates for image-guided biopsy should undergo excisional biopsy. Biopsy tissue cores are radiographed after the procedure to document the sampling of suspicious microcalcifications. Marking the biopsy site with a metallic clip is a standard of care.

Because percutaneous image-guided breast biopsy specimens represent only a sample of an abnormality observed on mammography, the results are subject to sampling error. Invasive carcinoma is detected upon lumpectomy in 10–20 % of patients in whom DCIS was diagnosed by a stereotactic core-needle biopsy, and DCIS is diagnosed upon excisional biopsy in 10–30 % of patients with atypical ductal hyperplasia or radial scar on stereotactic biopsy [33–35]. If core-needle biopsy results are discordant with the findings of imaging, repeat image-guided percu-

taneous biopsy or an excisional biopsy should be performed to clarify the diagnosis.

Excisional biopsy is performed with the assistance of a preoperative image-guided localization of the mammographic abnormality or of the previously placed metallic clip marking the biopsy site. Two localization techniques utilize either wire or radioactive seed. Specimen radiography is essential to confirm the removal of the microcalcifications of interest. The surgical pathology report should document the presence of biopsy site changes to confirm appropriate localization.

### Treatment

The treatment of DCIS is multidisciplinary. Surgical treatment can be either breast conservation (also referred to as segmental mastectomy, lumpectomy, or wide local excision) or mastectomy. Most patients who undergo breast-conserving surgery receive postoperative radiation therapy to improve local control. Postoperative systemic endocrine therapy, most often with tamoxifen, is also utilized for those patients whose tumors are estrogen receptor positive (ER+).

### Surgery: Mastectomy

Traditionally, DCIS was treated with total mastectomy. The rationale for mastectomy for DCIS was based on the high incidence of multifocality and multicentricity and the risk of occult invasion associated with the disease. Retrospective reviews indicate a mortality rate of 0–8 % after mastectomy for DCIS [36–38]. Local recurrence rates for DCIS after mastectomy are 1–3 % [36, 39, 40]. More recently, immediate reconstruction has been offered to patients with the utilization of skin-sparing or nipple-areola mastectomy in carefully chosen patients [41]. Occasionally, even after mastectomy, close or positive margins are observed on the pathological specimen, and these have been identified as an independent risk factor for locoregional recurrence. However, this rate is very low (10-year locoregional recurrence

rate of 1 %); therefore, postmastectomy radiation therapy is not warranted, except for patients with multiple positive margins that cannot be surgically excised [42].

### Breast-Conserving Surgery

The goal of breast-conserving surgery in the treatment of DCIS is to remove all suspicious calcifications and obtain negative surgical margins. Because DCIS is usually nonpalpable, breast-conserving surgery is performed with preoperative image-guided localization utilizing either a wire or a radioactive seed. In patients with extensive calcifications, bracketing of the calcifications with two or more wires or seeds may assist the excision of all suspicious calcifications. The orientation of the specimen intraoperatively with two or more marking sutures is critical for margin analysis. In addition, specimen radiography is essential to confirm removal of both the marking clip and the microcalcifications. After mammography of the surgical specimen, the specimen should be inked and then serially sectioned for pathological examination to evaluate the margin status and extent of disease.

The goal of breast-conserving surgery is to obtain tumor-free margins. The adequate margin for DCIS remains controversial. Most surgeons advocate re-excision for positive surgical margins, and many surgeons advocate re-excision for close margins, using varying thresholds of less than 1, 2, or 5 mm. Residual tumor was found on re-excision in 41 % of patients with DCIS with 0 to 1 mm margins, 31 % of patients with 1 to 2 mm margins, and 0 % of patients with greater than 2 mm margins [43]. Multifocality and an increasing number of close or involved margins have been identified as predictive of additional disease on re-excision [44]. This may be a surrogate for increased extent of disease. Recent consensus guidelines issued jointly by the Society of Surgical Oncology and the American Society for Radiation Oncology, which recommend “no ink on tumor” as the standard for an adequate margin in invasive cancer, caution that these findings cannot be extrapolated to DCIS [45].

### Role of Sentinel Lymph Node Biopsy and Axillary Staging

Because DCIS is a noninvasive disease, lymph node involvement is not expected. Thus, axillary lymph node dissection or sentinel lymph node biopsy should not be performed for DCIS treated with breast conservation [46]. If microinvasion or frank invasion is identified on final pathology after breast-conserving surgery, the patient should return to the operating room for sentinel lymph node biopsy. Sentinel lymph node biopsy was positive for metastasis among 12 % of patients with DCIS who were considered at high risk for invasion due to the presence of a large palpable mass and among 10 % of patients who had DCIS with microinvasion [47]. Patients who undergo mastectomy for DCIS should routinely undergo sentinel lymph node dissection because it is not possible to perform lymphatic mapping after a mastectomy if invasive cancer is incidentally found in the mastectomy specimen. These suggestions are reflected in the most recently issued American Society of Clinical Oncology Clinical Practice Guidelines [48]. Sentinel node dissection is associated with a small risk of lymphedema of approximately 5 % [49–52].

### Radiation Therapy

Most patients with DCIS who undergo breast-conserving surgery receive postoperative radiation therapy. Three prospective randomized studies have evaluated the role of radiation therapy following breast-conserving surgery for DCIS. In the National Surgical Adjuvant Breast and Bowel Project (NSABP) B-17 trial, 818 women with DCIS were randomized after margin-negative resection to observation or radiation therapy [53]. At 12 years of follow-up, the cumulative incidence of ipsilateral breast tumors decreased with the addition of radiation therapy from 14.6 to 8 % for noninvasive disease and from 16.8 to 7.7 % for invasive disease [54]. There was no difference in the 12-year overall survival rate with the addition of radiation: 86 % and 87 % of women were alive in the observation

versus radiation therapy group, respectively. However, 58 % of all deaths occurred before any breast cancer event. The deaths of 12 patients (3 %) in the observation group and 15 patients (3.6 %) in the radiation therapy group were attributed to invasive breast cancer [54].

A benefit of radiation therapy for DCIS was also observed in the European Organization for Research and Treatment of Cancer (EORTC) 10853 trial [55]. In this trial, 1,010 women with DCIS were randomized to breast-conserving surgery or breast-conserving surgery plus radiation therapy. At a median follow-up time of 4.25 years, radiation therapy was associated with a reduction in the incidence of non-invasive ipsilateral breast tumors from 8.8 to 5.8 % and with a reduction in the incidence of invasive ipsilateral breast tumors from 8.0 to 4.8 %. The 15-year invasive locoregional recurrence free rate was 84 % in the local excision-only group and 90 % in the local excision and radiation group [56].

A third trial conducted by the United Kingdom Coordinating Committee on Cancer Research also confirmed the benefits of radiation therapy for local control [57]. At a median follow-up of 4.4 years, the incidence of ipsilateral breast tumors decreased from 7 to 3 % for noninvasive lesions and from 6 to 3 % for invasive lesions when radiation was administered. Taken together, these three trials demonstrate that the addition of radiation therapy following breast-conserving therapy for DCIS results in an approximately 50 % relative reduction in breast cancer recurrence.

Whole-breast radiation is standard for patients undergoing breast-conserving surgery and is generally well tolerated. The most common morbidity is radiation-induced skin changes, including discoloration, fibrosis, and telangiectasias. Rare, severe side effects include damage to the heart and lungs, rib fractures, and radiation-induced secondary malignancy, angiosarcoma.

### **Partial-Breast Irradiation**

Local recurrences in the breast after a diagnosis of DCIS tend to occur in the immediate vicinity of the surgical resection cavity. Therefore, the impact of whole-breast irradiation in reducing local recurrence is most critical in the area immediately surrounding the original tumor bed. Based on this

knowledge, the goal of partial-breast irradiation is to achieve equivalent local control by focusing the treatment on the tissue surrounding the surgical resection cavity. Accelerated partial-breast irradiation is a technique in which high-dose radiation is delivered over a shorter period of time to a limited region of the breast surrounding the primary tumor site. The treatment is completed over 4–5 days, whereas conventional whole-breast external beam radiation therapy typically requires 5–6 weeks. Several methods of partial-breast irradiation have been described, including brachytherapy via multiple catheters placed in the breast parenchyma, localized conformal external beam radiation therapy, brachytherapy via bead or seed implants, single-dose intraoperative radiation therapy, and brachytherapy via a balloon catheter inserted into the cavity after breast-conserving surgery. In the American Society of Breast Surgeons' registry of accelerated partial-breast irradiation, of 194 patients with DCIS, 63 had at least 5 years of follow-up [58]. Of these, 92 % had favorable cosmetic results. The 5-year actuarial local-regional recurrence rate was 3.39 %, comparable to the rate of 7.5 % reported in the NSABP B-17 trial, which used whole-breast irradiation. The NSABP B-39/RTOG 0413 trial (ClinicalTrials.gov identifier NCT00103181), which opened in 2005, will provide additional information about the potential role of accelerated partial-breast irradiation in patients with DCIS as well as those with invasive breast cancer. In this trial, patients with DCIS of 3 cm or less or with invasive stage I or II breast cancer who undergo breast-conserving surgery with negative margins will be randomized to standard adjuvant whole-breast external beam radiation therapy or accelerated partial-breast irradiation. Patients will receive systemic therapy at the discretion of the treating physician. Local tumor control is the primary endpoint, and the secondary endpoints are disease-free and overall survival, cosmetic outcome, and treatment toxicity. Study completion is expected in 2016.

### **Omitting Radiation Therapy**

The use of radiation therapy varies depending on socioeconomic status, race, and the region of the United States in which the patient lives [59–62].



Many patients with DCIS who are candidates for breast-conserving surgery choose mastectomy because they are unable to complete 6 weeks of daily radiation therapy because of social considerations or due to concerns about postirradiation complications. An estimated 20 % of women undergoing breast-conserving surgery who would benefit from radiation therapy do not receive it as part of their treatment. Breast-conserving surgery without radiation therapy may be sufficient in selected patients with DCIS. In a study of 79 patients with DCIS who underwent margin-negative excision alone, after 124 months of follow-up, the local recurrence rate was 16 % overall: 33 % for the subgroup of patients with high-grade lesions and comedo necrosis versus only 2 % for the patients with low- or intermediate-grade lesions [16].

Margin width is an independent prognostic factor for local recurrence but is insufficient alone to predict which patients can safely forgo radiation therapy. In a retrospective analysis of 469 patients with DCIS who underwent breast conservation with margins of at least 10 mm, postoperative radiation therapy was not associated with a lower recurrence rate [63]. By contrast, on reanalysis of the NSABP B-17 data, all patient cohorts benefitted from radiation therapy, regardless of the clinical or mammographic tumor characteristics [64]. Furthermore, a prospective trial from the Dana-Farber/Harvard Cancer Center, in which radiation therapy was omitted in patients with grade 1 or 2 DCIS measuring 25 mm or less and resected with at least a 10 mm margin, was terminated early because the number of local recurrences was higher in the no-radiation group. The 10-year cumulative incidence of local recurrence was 15.6 %, and the annual local recurrence was 1.9 % per patient-year [65]. A multivariable nomogram that estimates local recurrence in women with DCIS treated with breast-conserving surgery can be used to estimate the absolute risk of ipsilateral breast tumor recurrence at 5 or 10 years, and the risk can be weighed against the use of other adjuvant treatment options. The nomogram incorporates the age at diagnosis, family history, type of patient presentation (radiological or clinical),

nuclear grade, necrosis, margins, number of excisions, and receipt of radiation and/or adjuvant endocrine therapy, all of which are factors that affect the risk of ipsilateral breast tumor recurrence [66].

Two prospective studies investigating the omission of radiation therapy after breast-conserving surgery for DCIS are the Eastern Cooperative Oncology Group E-5194 and the Radiation Therapy Oncology Group (RTOG) 9804 trials. In the E-5194 trial, all patients had breast conservation with margins measuring 3 mm or more and no radiation therapy. Patients with low- or intermediate-grade DCIS of 25 mm or less were compared to high-grade DCIS of 10 mm or less [67]. The ipsilateral breast event rate at 5 years was 6.1 % in the 565 patients in the low-/intermediate-grade group and 15.3 % in the 105 patients in the high-grade group. Long-term follow-up of this cohort is ongoing. Similarly, the RTOG 9804 trial randomized patients with low- or intermediate-grade DCIS sized 25 mm or less and excised with margins of at least 3 mm to postoperative radiation therapy versus observation. Tamoxifen use was permitted in both groups. The trial closed early. The results were presented at the American Society of Clinical Oncology (ASCO) meeting in 2012 and indicated that the addition of radiation produced a statistically significant reduction in breast cancer recurrence at 5 years to 0.4 % compared to 3.2 % in the observation group. Reports of long-term outcomes are anticipated.

## Endocrine Therapy

The NSABP B-24 trial involved 1,804 women with DCIS treated with breast-conserving surgery and radiation therapy [64]. Patients were randomized to either tamoxifen (20 mg/day) or placebo for 5 years. A total of 16 % of the patients in the study had positive margins at resection. At a follow-up of 15 years, the ipsilateral invasive breast cancer rate was lower (8.5 % after tamoxifen and 10 % after placebo), corresponding to a 32 % reduction in events. The 15-year cumulative incidence of contralateral

invasive or noninvasive breast cancer was 7.3 % and 10.8 % in the tamoxifen and placebo groups, respectively. Invasive ipsilateral recurrence was associated with an increased mortality risk, whereas the recurrence of DCIS was not. The benefits of tamoxifen persisted even in patients with positive or unknown margin status. In a subgroup analysis of patients with ER+ DCIS, those who received tamoxifen had a 59 % reduction in their relative risk of breast cancer events compared to those who received placebo. Patients with estrogen receptor-negative DCIS derived no benefit from tamoxifen [68].

In the United Kingdom Coordinating Committee on Cancer Research trial, patients who underwent breast-conserving surgery were randomized to no-adjuvant treatment, adjuvant radiation therapy, tamoxifen treatment, or adjuvant radiation therapy plus tamoxifen. Positive margins after breast conservation surgery were an exclusion criterion for the trial. Compared with 33 % of patients in the NSABP B-24 trial, only 10 % of the women were younger than 50 years of age. After a median of 4.4 years of follow-up, radiation therapy had the greatest impact on reducing ipsilateral breast cancer events, whereas tamoxifen added to radiation therapy did not portend additional benefit [57].

Adjuvant tamoxifen should only be used in ER+ DCIS. Tamoxifen is generally very well tolerated, but it has been associated with vasomotor symptoms, deep vein thrombosis, pulmonary embolism, and increased cataract formation. The risk of endometrial cancer is increased twofold to sevenfold but still remains low. Tamoxifen is also associated with an increased risk of stroke and benign ovarian cysts.

The current National Comprehensive Cancer Network guidelines recommend tamoxifen treatment for 5 years for patients with ER+ DCIS; however, they do not specifically recommend treatment with aromatase inhibitors. Aromatase inhibitors have fewer cardiovascular side effects than tamoxifen and are used in postmenopausal women. The randomized prospective clinical trial NSABP B-35 (ClinicalTrials.gov identifier NCT00053898), “Anastrozole or Tamoxifen in Treating Postmenopausal Women with Ductal

Carcinoma in Situ Who Are Undergoing Lumpectomy and Radiation Therapy”, has completed patient enrollment, and the results are expected in 2016 [69]. The International Breast Cancer Intervention Study (IBIS-II), which enrolled postmenopausal women at increased risk of breast cancer, reported a reduction in the cumulative incidence of breast cancer from 5.6 % in the placebo group to 2.8 % in the anastrozole group, with no difference in mortality [70]. A small proportion (8 %) of the 3,864 randomized women had been treated within the last 6 months for ER+ DCIS with mastectomy ( $n=326$ ). A subset analysis of this group of patients has not yet been reported.

---

## Surveillance

The National Comprehensive Cancer Network treatment guidelines recommend annual physical examination and mammogram for the follow-up of patients after breast-conserving surgery with or without radiation and/or endocrine therapy. Whether these procedures improve the detection of recurrence and outcome is not entirely clear. Both patients who undergo breast-conserving therapy and those who undergo mastectomy should be monitored for the development of new primary cancers in the contralateral breast. Annual mammograms should be performed, except in cases of mastectomy, which should be evaluated with physical exam.

---

## Predictors of Local Relapse

DCIS features associated with a greater risk of local recurrence are larger tumor size ( $>3$  cm), high nuclear grade, comedo-type necrosis, and positive margins. Involved margins are the most important prognostic variable for predicting local relapse, but as mentioned above, margins alone cannot be used independently to assess risk. Age less than 50 years and a strong family history of breast cancer are also associated with an increased risk of local recurrence. However, none of the above factors are contraindications

for breast-conserving therapy. Molecular markers, such as the overexpression of HER-2/neu, nm23, heat shock protein, and metallothionein; low expression of p21 Waf1 and Bcl2; and DNA aneuploidy, have been reported to be associated with high-grade comedo lesions, but their importance as independent prognostic variables in DCIS is not currently known [7].

## Treatment and Outcome of Local Recurrence

Patients with DCIS have excellent overall survival. In the NSABP B-17 trial, at a median follow-up of 12 years, only 27 deaths were attributed to breast cancer (3.3 %) [64]. In the B-24 trial, after 7 years of follow-up, 0.8 % died from breast cancer [64]. The management of local recurrence depends on the therapy the patient received for the primary cancer. In patients who had breast-conserving surgery without radiation therapy, re-excision to negative margins and postoperative radiation therapy are recommended. Mastectomy is the standard treatment in patients who have had breast-conserving surgery with radiation therapy. Rarely, for patients who have had mastectomy, local recurrence should be treated with wide local excision followed by postmastectomy radiation. If tissue coverage is a concern, tissue transfer techniques, such as a latissimus dorsi flap, can be employed. A reconstructive surgeon should be involved preoperatively.

In both B-17 and B-24, approximately 50 % of local recurrences were invasive. In one population-based study, younger age played a role in local recurrence after DCIS treatment and was associated with more invasive recurrence [71]. The prognosis after the treatment of local recurrence is worse when the recurrence is invasive compared to noninvasive. Disease-specific mortality after invasive recurrence is approximately 15 % [72, 73]. If a recurrence is invasive, the axilla should be staged with lymphatic mapping and sentinel lymph node dissection. Follow-up for patients with DCIS should be long term to detect not only recurrent disease but also the development of new ipsilateral or contralateral primary tumors.

## Conclusion

DCIS is most commonly diagnosed as a mammographic abnormality, which is subsequently percutaneously biopsied with image guidance. The pathological evaluation includes tumor type and grade and any evidence of microinvasion. Estrogen and progesterone receptor status should also be determined, but this is usually deferred until after surgical resection because if an invasive portion is unexpectedly identified, the receptors should be measured in that tissue, given the heterogeneity of breast lesions.

Therapy for DCIS should be personalized for each patient based on tumor size, tumor-to-breast size ratio, mammographic appearance, and margin width as well as patient preference. The benefits and risks of breast-conserving surgery and mastectomy should be discussed in detail with each patient. Most patients with DCIS are candidates for breast-conserving therapy, and this is the preferred method of local treatment because it offers equivalent overall survival compared to mastectomy. Re-excision is recommended for patients who have margins less than 2 mm on final pathological examination after breast-conserving surgery. Mastectomy is indicated in patients with persistently positive margins after attempts at breast conservation, in those with a contraindication to postoperative radiation therapy and in patients with diffuse, malignant-appearing calcifications throughout the breast. Mastectomy may also be a better option if a patient's anxiety about possible local recurrence outweighs the impact of a mastectomy on quality of life. For all patients undergoing mastectomy, immediate breast reconstruction should be considered.

Patients who have mastectomy for DCIS should also undergo sentinel lymph node biopsy. In patients who undergo breast-conserving surgery, sentinel lymph node dissection is performed if a diagnosis of invasive breast cancer is subsequently confirmed.

In patients undergoing breast-conserving surgery, the risk of local recurrence is reduced with adjuvant radiation therapy. Omitting

radiation therapy is considered in carefully selected patients with small (<1 cm in diameter), low-grade lesions that have been excised with margins of at least 5 mm and who can be observed diligently for recurrence. New genomic-based multigene assays also hold promise in identifying patients who may safely omit radiation. Partial-breast irradiation is offered most commonly in research protocols. Tamoxifen is offered for 5 years to women with estrogen receptor-positive DCIS who do not have a history of venous thromboembolism or stroke.

Posttreatment surveillance of patients treated for DCIS includes annual breast and/or chest wall physical examinations and diagnostic mammograms. All patients with DCIS should be considered for clinical trials.

## References

- Desantis C, DeSantis CE, Lin CC, Mariotto AB, Siegel RL, Stein KD, Kramer JL, et al. Breast cancer statistics, 2013. *CA Cancer J Clin*. 2014;64(1):52–62.
- Zujewski JA, Harlan LC, Morrell DM, Stevens JL. Ductal carcinoma in situ: trends in treatment over time in the US. *Breast Cancer Res Treat*. 2011;127(1):251–7.
- Lynge E, Ponti A, James T, Májek O, von Euler-Chelpin M, Anttila A, et al. Variation in detection of ductal carcinoma in situ during screening mammography: a survey within the International Cancer Screening Network. *Eur J Cancer*. 2014;50(1):185–92.
- Virnig BA, Tuttle TM, Shamlivan T, Kane RL. Ductal carcinoma in situ of the breast: a systematic review of incidence, treatment, and outcomes. *J Natl Cancer Inst*. 2010;102(3):170–8.
- Kuerer HM, Albarracín CT, Yang WT, Cardiff RD, Brewster AM, Symmans WF, et al. Ductal carcinoma in situ: state of the science and roadmap to advance the field. *J Clin Oncol*. 2009;27(2):279–88.
- Erbas B, Provenzano E, Armes J, Gertig D. The natural history of ductal carcinoma in situ of the breast: a review. *Breast Cancer Res Treat*. 2006;97(2):135–44.
- Benson JR, Wishart GC. Predictors of recurrence for ductal carcinoma in situ after breast-conserving surgery. *Lancet Oncol*. 2013;14(9):e348–57.
- Solin LJ, Gray R, Baehner FL, Butler SM, Hughes LL, Yoshizawa C, et al. A multigene expression assay to predict local recurrence risk for ductal carcinoma in situ of the breast. *J Natl Cancer Inst*. 2013;105(10):701–10.
- Omer ZB, Hwang ES, Esserman LJ, Howe R, Ozanne EM. Impact of ductal carcinoma in situ terminology on patient treatment preferences. *JAMA Intern Med*. 2013;173(19):1830–1.
- Bombonati A, Sgroi DC. The molecular pathology of breast cancer progression. *J Pathol*. 2011;223(2):307–17.
- Allred DC, Wu Y, Mao S, Nagtegaal ID, Lee S, Perou CM, et al. Ductal carcinoma in situ and the emergence of diversity during breast cancer evolution. *Clin Cancer Res*. 2008;14(2):370–8.
- Lee S, Stewart S, Nagtegaal I, Luo J, Wu Y, Colditz G, et al. Differentially expressed genes regulating the progression of ductal carcinoma in situ to invasive breast cancer. *Cancer Res*. 2012;72(17):4574–86.
- Burstein HJ, Polyak K, Wong JS, Lester SC, Kaelin CM. Ductal carcinoma in situ of the breast. *N Engl J Med*. 2004;350(14):1430–41.
- Seth A, Kitching R, Landberg G, Xu J, Zubovits J, Burger AM. Gene expression profiling of ductal carcinomas in situ and invasive breast tumors. *Anticancer Res*. 2003;23(3A):2043–51.
- Ma XJ, Salunga R, Tuggle JT, Gaudet J, Enright E, McQuary P, et al. Gene expression profiles of human breast cancer progression. *Proc Natl Acad Sci U S A*. 2003;100(10):5974–9.
- Lagios MD, Margolin FR, Westdahl PR, Rose MR. Mammographically detected duct carcinoma in situ. Frequency of local recurrence following tylectomy and prognostic effect of nuclear grade on local recurrence. *Cancer*. 1989;63(4):618–24.
- Silverstein MJ, Poller DN, Waisman JR, Colburn WJ, Barth A, Gierson ED, et al. Prognostic classification of breast ductal carcinoma-in-situ. *Lancet*. 1995;345(8958):1154–7.
- Holland R, Hendriks JH, Vebeek AL, Mravunac M, Schuurmans Stekhoven JH. Extent, distribution, and mammographic/histological correlations of breast ductal carcinoma in situ. *Lancet*. 1990;335(8688):519–22.
- Kurtz JM, Amalric R, Brandone H, Ayme Y, Jacquemier J, Pietra JC, et al. Local recurrence after breast-conserving surgery and radiotherapy. Frequency, time course, and prognosis. *Cancer*. 1989;63(10):1912–7.
- Adamovich TL, Simmons RM. Ductal carcinoma in situ with microinvasion. *Am J Surg*. 2003;186(2):112–6.
- Guth AA, Mercado C, Roses DF, Darvishian F, Singh B, Cangiarella JF. Microinvasive breast cancer and the role of sentinel node biopsy: an institutional experience and review of the literature. *Breast J*. 2008;14(4):335–9.
- de Mascarel I, MacGrogan G, Mathoulin-Pélissier S, Soubeyran I, Picot V, Coindre JM. Breast ductal carcinoma in situ with microinvasion: a definition supported by a long-term study of 1248 serially sectioned ductal carcinomas. *Cancer*. 2002;94(8):2134–42.
- Mirza NQ, Vlastos G, Meric F, Sahin AA, Singletary SE, Newman LA, et al. Ductal carcinoma-in-situ: long-term results of breast-conserving therapy. *Ann Surg Oncol*. 2000;7(9):656–64.
- Kuhl CK, Schrading S, Bieling HB, Wardelmann E, Leutner CC, Koenig R, et al. MRI for diagnosis of

- pure ductal carcinoma in situ: a prospective observational study. *Lancet*. 2007;370(9586):485–92.
25. Menell JH, Morris EA, Dershaw DD, Abramson AF, Brogi E, Liberman L. Determination of the presence and extent of pure ductal carcinoma in situ by mammography and magnetic resonance imaging. *Breast J*. 2005;11(6):382–90.
  26. Berg WA, Gutierrez L, NessAiver MS, Carter WB, Bhargavan M, Lewis RS, et al. Diagnostic accuracy of mammography, clinical examination, US, and MR imaging in preoperative assessment of breast cancer. *Radiology*. 2004;233(3):830–49.
  27. Rosen EL, Smith-Foley SA, DeMartini WB, Eby PR, Peacock S, Lehman CD. BI-RADS MRI enhancement characteristics of ductal carcinoma in situ. *Breast J*. 2007;13(6):545–50.
  28. Lehman CD, Gatsonis C, Kuhl CK, Hendrick RE, Pisano ED, Hanna L, et al. MRI evaluation of the contralateral breast in women with recently diagnosed breast cancer. *N Engl J Med*. 2007;356(13):1295–303.
  29. Hollingsworth AB, Stough RG, O'Dell CA, Brekke CE. Breast magnetic resonance imaging for preoperative locoregional staging. *Am J Surg*. 2008;196(3):389–97.
  30. Morrow M. Magnetic resonance imaging in the breast cancer patient: curb your enthusiasm. *J Clin Oncol*. 2008;26(3):352–3.
  31. Solin LJ, Orel SG, Hwang WT, Harris EE, Schnall MD. Relationship of breast magnetic resonance imaging to outcome after breast-conservation treatment with radiation for women with early-stage invasive breast carcinoma or ductal carcinoma in situ. *J Clin Oncol*. 2008;26(3):386–91.
  32. Bedrosian I, Mick R, Orel SG, Schnall M, Reynolds C, Spitz FR, et al. Changes in the surgical management of patients with breast carcinoma based on preoperative magnetic resonance imaging. *Cancer*. 2003;98(3):468–73.
  33. Darling ML, Smith DN, Lester SC, Kaelin C, Selland DL, Denison CM, et al. Atypical ductal hyperplasia and ductal carcinoma in situ as revealed by large-core needle breast biopsy: results of surgical excision. *AJR Am J Roentgenol*. 2000;175(5):1341–6.
  34. Dillon MF, McDermott EW, Quinn CM, O'Doherty A, O'Higgins N, Hill AD. Predictors of invasive disease in breast cancer when core biopsy demonstrates DCIS only. *J Surg Oncol*. 2006;93(7):559–63.
  35. Rutstein LA, Johnson RR, Poller WR, Dabbs D, Groblewski J, Rakitt T. Predictors of residual invasive disease after core needle biopsy diagnosis of ductal carcinoma in situ. *Breast J*. 2007;13(3):251–7.
  36. Balch CM, Singletary SE, Bland KI. Clinical decision-making in early breast cancer. *Ann Surg*. 1993;217(3):207–25.
  37. Fowble B. Conservative surgery and radiation in the treatment of early (stages I and II) breast cancer. Long-term results. *Helv Chir Acta*. 1989;55(6):831–6.
  38. Vezeridis MP, Bland KI. Management of ductal carcinoma in situ. *Surg Oncol*. 1994;3(6):309–25.
  39. Boyages J, Delaney G, Taylor R. Predictors of local recurrence after treatment of ductal carcinoma in situ: a meta-analysis. *Cancer*. 1999;85(3):616–28.
  40. Silverstein MJ, Cohlman BF, Gierson ED, Furmanski M, Gamagami P, Colburn WJ, et al. Duct carcinoma in situ: 227 cases without microinvasion. *Eur J Cancer*. 1992;28(2–3):630–4.
  41. Wagner JL, Fearmonti R, Hunt KK, Hwang RF, Meric-Bernstam F, Kuerer HM, et al. Prospective evaluation of the nipple-areola complex sparing mastectomy for risk reduction and for early-stage breast cancer. *Ann Surg Oncol*. 2012;19(4):1137–44.
  42. Fitzsullivan E, Lari SA, Smith B, Caudle AS, Krishnamurthy S, Lucci A, et al. Incidence and consequence of close margins in patients with ductal carcinoma-in situ treated with mastectomy: is further therapy warranted? *Ann Surg Oncol*. 2013;20(13):4103–12.
  43. Neuschatz AC, DiPetrillo T, Steinhoff M, Safaii H, Yunes M, Landa M, et al. The value of breast lumpectomy margin assessment as a predictor of residual tumor burden in ductal carcinoma in situ of the breast. *Cancer*. 2002;94(7):1917–24.
  44. Hadzikadic Gusic L, McGuire KP, Ozmen T, Soran A, Thomas CR, McAuliffe PF, et al. Margin width is not predictive of residual disease on re-excision in breast conserving therapy. *J Surg Oncol*. 2014;109(5):426–30.
  45. Moran MS, Schnitt SJ, Giuliano AE, Harris JR, Khan SA, Horton J, et al. Society of Surgical Oncology-American Society for Radiation Oncology consensus guideline on margins for breast-conserving surgery with whole-breast irradiation in stages I and II invasive breast cancer. *J Clin Oncol*. 2014;32(14):1507–15.
  46. Julian TB, Land SR, Fourchette V, Haile SR, Fisher ER, Mamounas EP, et al. Is sentinel node biopsy necessary in conservatively treated DCIS? *Ann Surg Oncol*. 2007;14(8):2202–8.
  47. Klauber-DeMore N, Tan LK, Liberman L, Kaptain S, Fey J, Borgen P, et al. Sentinel lymph node biopsy: is it indicated in patients with high-risk ductal carcinoma-in-situ and ductal carcinoma-in-situ with microinvasion? *Ann Surg Oncol*. 2000;7(9):636–42.
  48. Lyman GH, Temin S, Edge SB, Newman LA, Turner RR, Weaver DL, et al. Sentinel lymph node biopsy for patients with early-stage breast cancer: American society of clinical oncology clinical practice guideline update. *J Clin Oncol*. 2014;32(13):1365–83.
  49. DiSipio T, Rye S, Newman B, Hayes S. Incidence of unilateral arm lymphoedema after breast cancer: a systematic review and meta-analysis. *Lancet Oncol*. 2013;14(6):500–15.
  50. McLaughlin SA, Wright MJ, Morris KT, Giron GL, Sampson MR, Brockway JP, et al. Prevalence of lymphedema in women with breast cancer 5 years after sentinel lymph node biopsy or axillary dissection: objective measurements. *J Clin Oncol*. 2008;26(32):5213–9.



51. Mansel RE, Fallowfield L, Kissin M, Goyal A, Newcombe RG, Dixon JM, et al. Randomized multicenter trial of sentinel node biopsy versus standard axillary treatment in operable breast cancer: the ALMANAC Trial. *J Natl Cancer Inst.* 2006;98(9):599–609.
52. Langer I, Guller U, Berclaz G, Koechli OR, Schaar G, Fehr MK, et al. Morbidity of sentinel lymph node biopsy (SLN) alone versus SLN and completion axillary lymph node dissection after breast cancer surgery: a prospective Swiss multicenter study on 659 patients. *Ann Surg.* 2007;245(3):452–61.
53. Fisher ER, Costantino J, Fisher B, Palekar AS, Redmond C, Mamounas E. Pathologic findings from the National Surgical Adjuvant Breast Project (NSABP) Protocol B-17. Intraductal carcinoma (ductal carcinoma in situ). The National Surgical Adjuvant Breast and Bowel Project Collaborating Investigators. *Cancer.* 1995;75(6):1310–9.
54. Fisher B, Land S, Mamounas E, Dignam J, Fisher ER, Wolmark N. Prevention of invasive breast cancer in women with ductal carcinoma in situ: an update of the National Surgical Adjuvant Breast and Bowel Project experience. *Semin Oncol.* 2001;28(4):400–18.
55. Julien JP, Bijker N, Fentiman IS, Peterse JL, Delledonne V, Rouanet P, et al. Radiotherapy in breast-conserving treatment for ductal carcinoma in situ: first results of the EORTC randomised phase III trial 10853. EORTC Breast Cancer Cooperative Group and EORTC Radiotherapy Group. *Lancet.* 2000;355(9203):528–33.
56. Donker M, Litière S, Weruisky G, Julien JP, Fentiman IS, Agresti R, et al. Breast-conserving treatment with or without radiotherapy in ductal carcinoma In Situ: 15-year recurrence rates and outcome after a recurrence, from the EORTC 10853 randomized phase III trial. *J Clin Oncol.* 2013;31(32):4054–9.
57. Houghton J, George WD, Cuzick J, Duggan C, Fentiman IS, Spittle M, et al. Radiotherapy and tamoxifen in women with completely excised ductal carcinoma in situ of the breast in the UK, Australia, and New Zealand: randomised controlled trial. *Lancet.* 2003;362(9378):95–102.
58. Jeruss JS, Kuerer HM, Beitsch PD, Vicini FA, Keisch M. Update on DCIS outcomes from the American Society of Breast Surgeons accelerated partial breast irradiation registry trial. *Ann Surg Oncol.* 2011;18(1):65–71.
59. Lund MJ, Brawley OP, Ward KC, Young JL, Gabram SS, Eley JW. Parity and disparity in first course treatment of invasive breast cancer. *Breast Cancer Res Treat.* 2008;109(3):545–57.
60. Silber JH, Rosenbaum PR, Clark AS, Giantonio BJ, Ross RN, Teng Y, et al. Characteristics associated with differences in survival among black and white women with breast cancer. *JAMA.* 2013;310(4):389–97.
61. Yao N, Matthews SA, Hillemeier MM, Anderson RT. Radiation therapy resources and guideline-concordant radiotherapy for early-stage breast cancer patients in an underserved region. *Health Serv Res.* 2013;48(4):1433–49.
62. Bickell NA, Wang JJ, Oluwole S, Schrag D, Godfrey H, Hiottis K, et al. Missed opportunities: racial disparities in adjuvant breast cancer treatment. *J Clin Oncol.* 2006;24(9):1357–62.
63. Silverstein MJ, Lagios MD, Groshen S, Waisman JR, Lewinsky BS, Martino S, et al. The influence of margin width on local control of ductal carcinoma in situ of the breast. *N Engl J Med.* 1999;340(19):1455–61.
64. Wapnir IL, Dignam JJ, Fisher B, Mamounas EP, Anderson SJ, Julian TB, et al. Long-term outcomes of invasive ipsilateral breast tumor recurrences after lumpectomy in NSABP B-17 and B-24 randomized clinical trials for DCIS. *J Natl Cancer Inst.* 2011;103(6):478–88.
65. Wong JS, Chen YH, Gadd MA, Gelman R, Lester SC, Schnitt SJ, et al. Eight-year update of a prospective study of wide excision alone for small low- or intermediate-grade ductal carcinoma in situ (DCIS). *Breast Cancer Res Treat.* 2014;143(2):343–50.
66. Rudloff U, Jacks LM, Goldberg JL, Wynveen CA, Brogi E, Patil S, et al. Nomogram for predicting the risk of local recurrence after breast-conserving surgery for ductal carcinoma in situ. *J Clin Oncol.* 2010;28(23):3762–9.
67. Hughes LL, Wang M, Page DL, Gray R, Solin LJ, Davidson NE, et al. Local excision alone without irradiation for ductal carcinoma in situ of the breast: a trial of the Eastern Cooperative Oncology Group. *J Clin Oncol.* 2009;27(32):5319–24.
68. Allred DC, Anderson SJ, Paik S, Wickerham DL, Nagtegaal ID, Swain SM, et al. Adjuvant tamoxifen reduces subsequent breast cancer in women with estrogen receptor-positive ductal carcinoma in situ: a study based on NSABP protocol B-24. *J Clin Oncol.* 2012;30(12):1268–73.
69. Vogel VG, Costantino JP, Wickerham DL, Cronin WM. National surgical adjuvant breast and bowel project update: prevention trials and endocrine therapy of ductal carcinoma in situ. *Clin Cancer Res.* 2003;9(1 Pt 2):495S–501.
70. Cuzick J, Sestak I, Forbes JF, Dowsett M, Knox J, Cawthorn S, et al. Anastrozole for prevention of breast cancer in high-risk postmenopausal women (IBIS-II): an international, double-blind, randomised placebo-controlled trial. *Lancet.* 2014;383(9922):1041–8.
71. Kong I, Narod SA, Taylor C, Paszat L, Saskin R, Nofech-Moses S, et al. Age at diagnosis predicts local recurrence in women treated with breast-conserving surgery and postoperative radiation therapy for ductal carcinoma in situ: a population-based outcomes analysis. *Curr Oncol.* 2014;21(1):e96–104.
72. Silverstein MJ, Lagios MD, Martino S, Lewinsky BS, Craig PH, Beron PJ, et al. Outcome after invasive local recurrence in patients with ductal carcinoma in situ of the breast. *J Clin Oncol.* 1998;16(4):1367–73.
73. Romero L, Klein L, Ye W, Holmes D, Soni R, Silberman H, et al. Outcome after invasive recurrence in patients with ductal carcinoma in situ of the breast. *Am J Surg.* 2004;188(4):371–6.

A. Lale Dogan

## Abstract

Breast cancer is the most common cancer among women, and it has a complex genetic basis for susceptibility. Understanding the mechanisms of DNA alterations leading to carcinogenesis can provide crucial insights for resolving the development of malignant processes such as growth, invasion, and metastasis. This chapter reviews hereditary and somatic genetic alterations, epigenetic misregulations, and miRNA signatures associated with breast cancer. The chapter also emphasizes the molecular profiles of breast cancer and the critical signaling pathway alterations.

## Keywords

Breast cancer • Susceptibility genes • Somatic genetic alterations • Epigenetics • miRNA • Molecular portrait • Signal transduction pathways

## Genetics of Breast Cancer

Breast cancer is known as a multifactorial disease, with factors including advanced age, early menarche, obesity, sedentary lifestyle, late pregnancy, and menopause. All of these factors can trigger the process of carcinogenesis. In addition, family history and the accumulation of

genetic aberrations are considered the most prominent and major factors for increasing the risk of breast cancer [1]. Genetic aberrations include point mutations, deletions, amplifications, rearrangements, translocations, and duplications.

Breast cancer susceptibility increases significantly in those with a familial history of breast cancer compared with the general population. This situation can be clarified with susceptibility genes, which play key roles in breast cancer progression [2].

Breast cancer susceptibility can be classified into two forms: hereditary and sporadic [3]. Germline mutations account for 10 % of all breast cancers.

---

A.L. Dogan, MD  
Department of Basic Oncology, Hacettepe Cancer  
Institute, Hacettepe University, Sıhhiye, Ankara  
06100, Turkey  
e-mail: [laled@hacettepe.edu.tr](mailto:laled@hacettepe.edu.tr)

## Hereditary Mutations in Breast Cancer

Mutations in *BRCA1* and *BRCA2* are the primary hereditary genetic aberrations in breast cancer. These mutations account for approximately half of all hereditary breast cancers. The *BRCA1* and *BRCA2* genes encode large proteins with multiple functions. *BRCA1* is localized on the 17th chromosome and encodes an 1863 amino acid protein with a zinc finger domain. The *BRCA1* and *BRCA2* proteins are involved in many cellular functions, including the repair of double-stranded DNA breaks for protection of the genome during replication. In short, they act mainly as tumor suppressor gene products. Mutations in *BRCA1* and *BRCA2* and age of cancer onset appear to vary interdependently. *BRCA2* is localized in the 13th chromosome, and in the case of *BRCA2*, the mutation is a secondary major factor for breast cancer predisposition. The occurrence of even one of these mutations can increase the risk of breast cancer to 25 % [3–5].

It was noted that the *BRCA1/BRCA2* mutation-carrying breast cancer patients' samples differed from tumors with BRCAness (*BRCAx*), such that the *BRCA* mutation-bearing samples showed a pleomorphic structure, tubular formation, and more aggressive tumor characteristics than non-*BRCA* tumors [6]. *BRCA-1*-related breast cancers typically occur in younger women and are described as “triple negative” because of the absence of estrogen and progesterone receptors and human epidermal growth factor receptor 2 (*HER2*). When the wild-type *BRCA1* or *BRCA2* allele is lost, mutated, or silenced, defective DNA repair occasionally occurs. Consequently, additional mutations accumulate during replication, and carcinogenesis is promoted. Poly(adenosine diphosphate-ribose) polymerase-1 (*PARP1*) is an enzyme involved in single-stranded DNA repair utilizing base excision repair. *PARP1* inhibitors hold a promising therapeutic strategy in *BRCA* defective tumor cells. When *PARP* inhibition is applied, *BRCA*-deficient tumor cells are completely devoid of repair mechanisms and undergo cell cycle arrest, genomic instability, and cell death.

The *TP53* tumor suppressor gene mutation is another important mutation in breast cancer. Additionally, the *TP53* mutation is the primary symptom of Li-Fraumeni syndrome (LFS). LFS patients are particularly prone to the progression of some cancer types, notably breast cancer. The *TP53* mutation also leads to an increase in the rate of the *HER2*<sup>+</sup> breast cancer subtype because of the localization of the *TP53* and *HER2* genes on the same chromosome (17th chromosome). The loss of *p53* functions because of mutational changes may influence *HER2*-related signal transduction pathways featuring sustainable activity. Because of this phenomenon, cancer cells might display drug-resistant phenotypes [7–9].


Hereditary mutations can be divided into three categories in terms of mutation risk and the frequency of mutation. The first category includes the *BRCA1/BRCA2*, *PTEN*, and *TP53* mutations, which are classified as high-penetrance, low-frequency predisposition genes. The second category includes the *CHEK2*, *ATM*, and *PALB2* genes; these are moderate-penetrance, low-frequency predisposition genes. Finally, the third category consists of the *FGFR2*, *MAP3K1*, and *TGFBI* gene mutations, which are low-penetrance, high-frequency predisposition genes [10] (Fig. 8.1).

## Somatic Genetic Alterations in Breast Cancer

Apart from germline mutations, some gene deletions or amplifications occur as somatic alterations in breast cancer. Among these, the *HER2* (20 %), *cyclin D1* (12 %), and *WIP1* (13 %) gene amplifications and the *PTEN* and *p53* gene deletions frequently occur in breast cancer [10].

In some situations, single-nucleotide polymorphisms (SNPs) may correlate with cancer pathogenesis. For example, the *MDM2* gene, which encodes an important ubiquitin ligase that negatively regulates *p53*, has two different alleles, T-T and G-G, in the SNP390 intron region. The G-G allele leads to high binding affinity to the ER transcription factor SP-1. As a result, the G allele character on the *MDM2* gene may cause the overexpression of the *MDM2* protein, the

**Fig. 8.1** Mechanisms and related genes involved in breast cancer susceptibility



repression of p53 function, and increased ER expression [10].


One consequence of somatic mutations is copy-number alterations (CNAs). The long arm of chromosome 1 is the main region in which amplifications are detected in breast cancer. Some gene amplifications in the long arm of chromosome 16 and some gene deletions in the short arm may be observed in genome-stable tumors. Additionally, there are some gene amplifications that occur in genome-unstable tumors, concerning the 8th chromosome and the 17th chromosome in which the *MYC* and *ERBB2* genes are localized on, respectively. Furthermore, genome-unstable tumors show a poor prognosis compared to genome-stable tumors. Because of that, in luminal A, luminal B, and normal-like breast cancer subtypes, disease-free survival (DFS) rates are higher than those in basal-like and HER2<sup>+</sup> breast cancer subtypes [11] (Fig. 8.2).

### Breast Cancer Epigenetics

In higher eukaryotes, DNA methylation and chromatin modifications are essential epigenetic mechanisms that operate in association with genetic mechanisms such as replication and transcription. In many cancers, including breast cancer, tumor cells have altered patterns of methylation and histone modification, resulting in transcriptional deregulation in favor of oncogen-

esis. DNA methyltransferases catalyze methyl donor transfer from S-adenosylmethionine to cytosine bases, which undergo methylation. Cytosine-phosphate-guanine (CpG) dinucleotide sequences frequently localize at gene promoter regions, where transcription initiates. Thus, DNA methylation at the promoter proximal CpG sequences, called CpG islands, is associated with gene silencing. In tumor cells, oncogenes undergo hypomethylation, whereas tumor suppressor genes are hypermethylated. In active chromatin, unmethylated promoter regions can be transcribed by means of transcriptional machinery composed of transcription factors, co-activators with histone acetyltransferase activity (HAT), and acetylated histones H3, H4, and H2A and histone H3, which is methylated mainly at lysine residue K4 (H3-lysine 4). In heterochromatin, also called silent chromatin, transcription of the methylated regions is blocked by methylcytosine-binding protein; histone H3 is methylated at the lysine residues K9, K27, and K79 along with corepressors with histone deacetylase activity (HDAC). Furthermore, there are certain differences between epigenetic alterations and mutations; for example, mutations are irreversible changes, whereas epigenetic alterations are reversible. According to Knudson's two-hit hypothesis, the first hit is developed by epigenetic dysregulation and the second hit is the result of somatic mutations in nonhereditary tumors. Carcinogenic processes consequently evolve [12–14].

**Fig. 8.2** Copy-number alterations in breast cancer



*RASSF1A* is a tumor suppressor gene that encodes the Ras association domain-containing protein 1. In breast cancer, the deletion or altered patterns of expression of this gene appear to be related to pathogenesis. Functional studies documented decreasing levels of histone H3-lysine 4, whereas levels of histone H3-lysine 9 increased in tumor cells, resulting in *RASSF1A* silencing via epigenetic mechanisms. Previously, it was noted that the silencing of *RASSF1A* was associated with the overactivation of the *RAS* signaling pathway. As a result, tumor cells divide and proliferate in an uncontrolled fashion [15]. *BRCA1*, *E-cadherin*, *TMS1*, and *ER* are among the other genes that are hypermethylated in breast cancer [16].

Tumor cells in breast cancer subtypes that develop as a result of ER, PR, and androgen receptor hypermethylation have a more aggressive character and are inadequate for steroid hormone therapies, thus leading to a decrease in the rate of disease-free survival (DFS) [17] (Fig. 8.3).

### Molecular Portraits of Breast Cancer


*PIK3CA*, *TP53*, *GATA3*, *CDH1*, *RBI*, *MLL3*, *MAP3K1*, and *CDKN1B* gene mutations are frequently encountered in breast cancer. Although *PIK3CA* mutations are encountered at a rate of

41.3 %, *TP53* mutations are encountered at a rate of 16.1 %. *TP53* mutations are associated with the high histological grade luminal B breast cancer subtype, whereas *MAP3K1* mutations are related to luminal A breast cancer with a low histological grade. Expression of *Ki-67*, a proliferation marker, in tumor cells may result in resistance against aromatase inhibitors in advanced-stage tumors [18, 19].

Breast cancer is a heterogeneous cancer type in terms of both molecular and clinical characteristics. Even if individuals have similar clinicopathological symptoms, the disease may evolve, resulting in differing outcomes. In this respect, the classification of immunohistochemical staining may have limited utility for evaluation [20].

Investigations using pathological tissues have made it possible to develop immunohistochemical staining guidelines for classifying tumor cell characteristics. This classification is built upon the basis of ER (estrogen receptor), PR (progesterone receptor), and HER2 (human epidermal growth factor receptor) expression. Positive expression of hormone receptors occurs at a rate of 60–65 % in all breast cancer cases, and these cancers can be treated properly with hormone-based therapeutics. In breast cancer, the rate of HER2 positivity is approximately 15–20 %; this group is selectively treated with HER2-targeted





**Fig. 8.3** Chromatin modification by epigenetic mechanisms. **(a)** In active chromatin, histone acetyl transferase (*HAT*) mediates the acetylation of H2A, H3, and H4, and histone methyl transferase (*HMT*) mediates the trimethylation of lysine residue K4 at histone H3 (H3K4me3), leading to the accessibility of the unmethylated promoter

region to the co-activator protein (*CA*) and the transcription factor (*TF*). **(b)** Epigenetic gene inactivation by CpG island methylation by DNA methyl transferase (*DNMT*) is associated with histone modifications involving the trimethylation of lysine residue K27 (H3K27me3), lysine residue K9, or lysine residue K79 at histone H3

agents. The remaining 15–20 % of breast cancers lack hormone receptors and HER2 and are called triple-negative tumors [21, 22].

Microarray-based assays offer opportunity and resolution in understanding the molecular and genetic features of breast cancer. Perou and colleagues' work suggests five different breast cancer subtypes at the molecular level (molecular portraits) based on gene expression patterns. Among the subtypes they defined were the luminal A and luminal B tumor types, HER2<sup>+</sup> tumors, basal-like cancer (triple-negative breast cancer (TNBC)), and normal-like subtypes. The luminal A and luminal B subtypes are typically hormone receptor (ER or PR) positive and have a good prognosis with a 5-year survival rate of 80–85 %. Furthermore, hormone receptor-positive breast cancer patients carry *PIK3CA* mutations at a rate

of 40 %. Despite coexisting *PIK3CA* mutations and the consequent sustained PI3K signaling, these cells show low mTORC1 activity. Basal-like or triple-negative breast cancer subtypes lack ER, PR, and HER2, whereas they express *CK5/6* (cytokeratin 5/6) and *EGFR*. Generally, these subtypes are associated with sustained PI3K activity, *PTEN* deletion, and *BRCA1* and *TP53* mutations, and they ultimately have a poor prognosis. It should be noted that not all basal-like tumors are triple negative and vice versa. HER2<sup>+</sup> breast cancer is an aggressive phenotype with a poor prognosis [23, 24].

In one study, Prat and colleagues defined a new molecular-based breast cancer subtype. This subtype is called claudin-low and has low expression of luminal markers such as *ER*, *PR*, *GATA3*, *keratin 18*, and *HER2*; it also shows variable

expression of basal markers such as keratin 5, 14, and 17. Claudin-low breast cancer shows little or no expression of epithelial markers such as E-cadherin and claudin 3, making it different from basal-like breast cancer subtypes. The claudin-low tumor cell phenotype can be described as being similar to the mesenchymal phenotype; in other words, it has similar tumor-initiating cell (TIC) characteristics [25, 26].

Microarray-based gene expression profile analyses provide innovative and distinctive data that are beneficial to understand the heterogeneity of the structure of breast cancer. The Oncotype DX 21 gene analysis assay was the first approved assay for clinical use [27]. This assay is useful for determining risk factors of early-stage ER<sup>+</sup>, LN<sup>-</sup> tumors. The MammaPrint 70 gene analysis assay is another assay for clinical guidance and is useful for determining tumor proliferation and genomic grades of early-stage ER<sup>+</sup> tumors [28, 29]. Furthermore, this assay can be used to ensure prognostic disparity between LN<sup>+</sup> and LN<sup>-</sup> tumors. For other assays, such as the 8-gene recurrence score [30], the 14-gene metastatic score, and the 158-gene assay, preclinical studies are ongoing. The 50-gene analysis, which is currently called the PAM50 assay, targets all types of breast cancer and provides the advantage of intrinsic classification as indicated. Although this new assay is not yet applicable to clinics, preclinical studies are ongoing [2, 20, 31].

The implementation of the 21-gene assay data on ER<sup>+</sup> breast cancer patients treated with tamoxifen indicates that the 10-year disease-free survival rate is 96.8 % for patients with a low recurrence score, 90.9 % for patients with a mild recurrence score, and 60.5 % for patients with a high recurrence score, with a mean value of 87.8 %. Moreover, in ER<sup>+</sup> breast cancer patients treated with tamoxifen plus chemotherapy, this assay indicates that the 10-year disease-free survival rate is 95.6 % for patients with a low recurrence score, 89.1 % for patients with a mild recurrence score, and 88.1 % for patients with a high recurrence score; the mean value is 92.2 % [32].

In particular, the proliferation rate is a prognostic factor in the luminal subtype of breast cancer. Concordantly, amplifications of *FGD5*,

*METTL6*, *DTX3*, and *MRPS23* are well-known prognostic markers. For example, *FGD5* regulates the proangiogenic function of *VEGF* and increases cell proliferation. Among the others, *DTX3* promotes the Notch signaling pathway, *MRPS23* regulates proliferation and oxidative phosphorylation, and *METTL6* is related to drug sensitivity [33].

HER2<sup>+</sup> breast cancer subtypes are characterized by poor prognosis and different outcomes from systemic chemotherapy. In advanced-stage tumors, PI3K activation is usually deregulated. HER2<sup>+</sup> tumors are classified in three different clusters according to ER and LN expression and histological grades performed by 158-gene assay studies. If lymphatic infiltration occurs in early-stage HER2<sup>+</sup> tumors, it is associated with a good prognosis. In contrast, low lymphatic infiltration shows that the tumors have a more aggressive phenotype. In addition, studies have indicated that low lymphatic infiltration was associated with increasing PI3K signaling pathway activation and *IGFR1* expression. In HER2<sup>+</sup> tumors, *CXCR4*, *PLAU*, *CXCR1*, *TGFBR3*, and *STAT5A* gene expression levels are distinctive markers for evaluating tumor invasion and metastasis [34].

Recently, clarifying mechanisms of acquired resistance to HER2 targeted therapies has been of interest. One of the possible resistance mechanisms is the truncation of the HER2 receptors from the extracellular region; the resultant HER2 becomes a new variant called p95HER2. Thus, the HER2 antibody could not bind and recognize the HER2 receptors. Preclinical studies performed on cell lines show that this mechanism is based on specific HER2-mediated molecules, including *PTEN*, *PI3K*, *mTOR*, *MAPK*, and *VEGF*. However, studies performed on tissue samples show that the *AKT*, *IGFR1*, *p27 Kip1*, and *MUC4* genes are deregulated; in particular, the *PTPN11* gene shows the most alteration after HER2-targeted therapies. In short, resistance mechanisms might also be associated with changes in the expression patterns of these genes [35–37].

Basal-like tumors phenotypically resemble normal basal/myoepithelial breast cells. Basal-like breast cancer cells express basal cytokeratins

(5/6, 14, and 17), p-cadherin, and caveolin 1 and have patterns of hormone receptor and HER2 expression similar to those in normal breast cells. Additionally, basal-like breast cancer cells have *TP53* mutations and high expression levels of genes associated with proliferation. Furthermore, studies have reported that the *BRCA1* mutation is associated with this type of cancer [38].

Triple-negative breast cancer lacks hormone receptors and HER2. This type of cancer does not express basal markers and expresses basal prognostic markers less than basal-like tumors. Studies performed with the 14-gene assay show that high levels of aberration occur in the expression levels of *CLIC5*, *MATN1*, and *RPS28* in TNBC cells. Interestingly, high expression of the *IR-7*, *STAT1*, and *IFN* genes is associated with a good prognosis [39].

According to 386-gene assay studies, the TNBC subtype can be classified into seven different clusters. These clusters consist of basal-like 1 (BL1), basal-like 2 (BL2), immunomodulatory (IM), mesenchymal (M), mesenchymal-stem cell-like (MSL), luminal androgen receptor (LAR), and unstable (UNS). BL1 and BL2 clusters are characterized by genes associated with proliferation and the cell cycle. The *AURKA*, *AURKB*, *CENPA*, *CENPF*, *BUB1*, *TTK*, *CCNA2*, *PRC1*, *MYC*, *NRAS*, *PLK1*, and *BIRC5* genes are highly expressed in BL1 and BL2 tumors. Additionally, *EGFR*, *NGF*, *MET*, *Wnt/beta-catenin*, and *IGFRI* gene expression and their associated signaling pathways are activated, particularly in these clusters. The gene ontologies of the IM cluster include the overexpression/activation of TH1/TH2, NK cells, dendritic cells, B and T cell receptor immune cell signaling components, cytokine signaling, IL-12 and IL-17 cytokine signaling components, NF- $\kappa$ B, TNF, and JAK/STAT, which are standard immune response signaling transduction pathways and antigen presentation signal transduction pathways. M and MSL clusters are important in terms of motility and have distinctively aggressive characters compared to the other clusters. Signaling pathways associated with the actin regulatory protein Rho, the extracellular matrix (ECM) receptor interaction, Wnt, anaplastic lymphoma kinase (ALK),

and TGF- $\beta$  are factors that ensure cell motility. These pathways are also characteristic of the EMT marker, stem cell, and claudin-low type of breast cancer. Although the LAR cluster lacks ER, this cluster contains ample hormone regulation pathways for steroid synthesis, porphyrin metabolism, and androgen/estrogen metabolisms. Luminal character markers such as *FOXA1*, *KRT18*, and *XBPI* are the most highly expressed genes in the LAR cluster. The LAR cluster constitutes 11 % of TNBC breast cancer [40].

Although TNBC clusters show different clonal expansion patterns, somatic mutations of *TP53*, *PIK3CA*, and *PTEN* are encountered in all these clusters as a common trait [41, 42].

Immunohistochemical staining is essential for the diagnosis of breast cancer. However, because of tumor heterogeneity, immunohistochemical evaluation may not confer all the requirements for the molecular assessment of disease. It is possible to resolve copy-number alterations (CNAs) of breast cancer molecular portraits by array comparative genomic hybridization (aCGH) techniques. By this means, some subtypes can be characterized. Mutated genes can be classified using next-generation sequence technology. The results of genetic analyses in the whole genome provide single-nucleotide polymorphisms (SNPs) and copy-number variants (CNVs) for hereditary profiles and provide single-nucleotide variants (SNVs) and copy-number aberrations (CNAs) for somatic aberration profiles. In recent studies, breast cancer subtypes are classified into 10 different integrated clusters by determining the genetic profile of the somatic copy-number alterations. The only common point of 10 different integrated clusters is the high expression of mutated *PIK3CA* and *TP53* genes [20].

In the future, further analysis of next-generation sequencing data will improve the understanding of tumor heterogeneity, leading to deciding adequate therapy based on personal genomic breast cancer profiles and providing a more effective prevention of cancer [43].

Mechanisms that are involved in breast cancer susceptibility are presented in Table 8.1.

In a previous study, it was noted that in tumors in which the expression of 97 genes was

**Table 8.1** Molecular subtypes of breast cancer

Intrinsic subtypes by gene expression profiling	ER (IHC)	HER2 (IHC/ISH)	Ki-67 status (IHC)	Key molecular features [62, 25]	Predominant integrated cluster association [65, 20]
Luminal A	ER+	HER2-	Low	<i>PIK3CA</i> mutations, <i>MAP3K1</i> mutations, <i>ESR1</i> high expression, <i>GATA3</i> mutations, <i>FOXA1</i> mutations, quiet genomes: gain 1q, 8q, loss of 8p, 16q	Int cluster 2, Int cluster 3, Int cluster 4, Int cluster 7, Int cluster 8
Luminal B	ER+	HER2-	High	<i>TP53</i> mutations, <i>PIK3CA</i> mutations, <i>cyclin D1</i> amplification, <i>MDM2</i> amplification, <i>ATM</i> loss, enhanced genomic instability, focal amplifications (e.g., 8p12, 11q13)	Int cluster 1, Int cluster 2, Int cluster 6, Int cluster 9
HER2	ER+	HER2+	High	<i>HER2</i> amplification, <i>TP53</i> mutations, <i>PIK3CA</i> mutations, <i>FGFR4</i> high expression, <i>EGFR</i> high expression, <i>APOBEC</i> mutations, <i>cyclin D1</i> amplification, high genomic instability	Int cluster 5
Basal-like	ER-	HER2-	High	<i>TP53</i> mutations, <i>RB1</i> loss, <i>BRCA1</i> loss, high expression of DNA repair proteins, <i>FOXM1</i> activation, high genomic instability, focal amplifications (e.g., 8q24)	Int cluster 4, Int cluster 10
Claudin-low	ER-	HER2-	Low	<i>BRCA1</i> mutations, low expression of luminal markers, variable expression of keratin 5, 14, 17, little or no expression of <i>E-cadherin</i> and <i>claudin 3</i>	Int cluster 4

+ positive, – negative, *ER* estrogen receptor, *HER2* human epidermal growth factor receptor, *IHC* immunohistochemistry, *ISH* in situ hybridization

homogeneous and histological grade 3 was common, the overexpression of *UBEC2C*, *KPNA2*, *TPX2*, *FOXM1*, *STK6*, *CCNA2*, *BIRC5*, and *MYBL2* genes was concluded to be responsible for cell cycle progression and proliferation. However, grade 1 tumors differ from histological grade 3 samples in that they show a stable gene expression profile, whereas grade 2 tumors have heterogeneity in their gene expression profiles [44].

Driver mutations are the main mutations for carcinogenesis, and they provide a clonal selectivity advantage to cancer cells. Then, passenger mutations are triggered and help accelerate carcinogenesis. Previous studies have reported that

*AKT1*, *BRCA1*, *CDH1*, *GATA3*, *PIK3CA*, *RB1*, *PTEN*, and *TP53* mutations are driver mutations, and recent studies have added new driver gene mutations to this group, including *AKT2*, *ARID1B*, *CASP8*, *CDKN1B*, *MAP3K1*, *MAP3K13*, *NCOR1*, *SMARCD1*, and *TBX3*. *MAP3K1* and *MAP3K13* gene mutations result in deregulation of the ERK/MAPK signaling pathway. The *AKT* signaling pathway shows a sustained increase in activity in the presence of the *AKT2* mutation. Chromatin regulation is lost by means of *NCOR1*, *SMARCD1*, and *ARID1B* mutations. The *CDKN1B* mutation deregulates the cell cycle and proliferation. In the presence of the *CASP8* mutation, the apoptosis rate decreases.

The *TBX3* mutation may affect tissue morphology, but the exact mechanism remains to be clearly defined [45].

Breast cancer is divided into two different classes based on pathological form: invasive ductal carcinoma (IDC) and invasive lobular carcinoma. Analyzing the differences among these classes, the overexpression of *ERBB2*, *JAK2*, *ANKRD32*, and *NRGN* is said to be a prognostic factor for the IDC breast cancer class, and the overexpression of *VWF*, *ELN*, *DPT*, *EMCN*, *FABP4*, *CAVI*, *ADIPOG*, and *ALDH1A1* is reported as a prognostic factor for invasive lobular carcinoma [46].

The interpretation of and the lack of combined clinical and molecular data constitute a serious puzzle for molecular classification in current clustering methods. There is a new clustering method that holds more promise for combining clinical and molecular data. This methodological algorithm presents the results of combining data and is called Molecular Regularized Consensus Patient Stratification (MRCPS). Notably, the *GATA3* gene has been shown to have a major association with other deregulated genes, and mutation of *GATA3* is directly associated with the estrogen receptor [47].

### MicroRNA Signatures in Breast Cancer

MicroRNAs (miRNAs) are small RNAs of approximately 18–25 nucleotides and are endogenously expressed in cells. miRNAs can silence the expression of target genes via repressing translation or leading to the degradation of target mRNAs [48]. Thus, miRNAs modulate various cellular functions including proliferation, apoptosis, and angiogenesis.

Different tissue types have unique expression levels of specific miRNAs. Accordingly, each tumor type appears to have an miRNA signature with aberrant expression [49] (Fig. 8.4).

It is widely accepted that discrepancies among reported miRNA signatures may arise from intrinsic heterogeneity present in breast cancer tumors. Tumor stage, receptor status (HER2, ER,

PR), and vascular invasion may contribute to variability [49, 50]. The potential role of miRNAs as a tool for the diagnosis, classification, and treatment of breast cancer is currently under investigation [51–55]. miRNAs may act as oncogenes or tumor suppressors through regulating the mechanisms of proliferation, migration, invasion, metastasis, and apoptosis [56–64]. Table 8.2 presents an illustrative list of regulated miRNAs in breast cancer in association with related pathways.


---

## Biology of Breast Cancer

### Estrogen Receptors and Breast Cancer

The estrogen receptor (ER) is a steroid hormone receptor that resides in the cytoplasm and participates in cell proliferation, survival, and invasion in ER<sup>+</sup> breast cancer. The binding of estrogen is essential for translocation of the ER from the cytoplasm to the nucleus, where estrogen-bound ER dimerizes and binds to the estrogen response elements of target genes for the activation of gene expression. ER, as a transcription factor, also interacts with coregulatory proteins and other transcription factors. In addition, another form of ER, a membrane-bound or cytoplasmic protein, has non-genomic action. In short, activation of the receptors triggers phosphorylation and the activation of several receptors and signal proteins such as epidermal growth factor receptors, insulin-like growth factor-1R (IGF-1R), Src kinase, Shc adaptor protein, and phosphatidylinositol 3-kinase (PI3K). Thus, there is cross talk among growth factor receptors and estrogen receptor signaling that may contribute to resistance to antiestrogens. ER signaling may be targeted by two methods of inhibition. The common method is binding to the estrogen receptor, e.g., tamoxifen, and the other strategy involves decreasing estrogen production via aromatase inhibitors [65]. Resistance to antiestrogens is quite common, and the clinical aspects of hormone therapy will be discussed in another chapter.





**Fig. 8.4** Mechanisms of miRNA expression. MicroRNA (*miRNA*) is transcribed mainly by RNA polymerase (*pol*) II. RNA pol III is also involved in transcription. The pri-miRNA is spliced by Drosha and the DGCR8 enzymatic complex, which leads to pre-miRNA formation. Pre-miRNA is exported into the cytoplasm by the cargo protein exportin 5, and mature miRNA is obtained through

cleavage by Dicer. This mature miRNA binds to messenger RNA (*mRNA*), leading to the degradation or blockage of the translation of mRNA. The RNA-induced silencing complex (*RISC*), which is a multi-protein complex, uses miRNA as a template for recognizing complementary mRNA. *ORF* open reading frame

## Growth Factor Receptors in Breast Cancer

Growth factors and receptor tyrosine kinases (RTKs) are essential for cell proliferation and survival. The human epidermal growth factor receptor (HER) family consists of four members: EGFR/ErbB1, HER2/ErbB2, HER3/ErbB3, and HER4/ErbB4. In breast cancer, the HER family has been studied extensively. The common structure of HER proteins consists of an extracellular domain for ligand binding, a transmembrane domain, and a cytoplasmic catalytic kinase domain that drives downstream signaling pathways. The main pathways are the PI3K/Akt and

Ras/Raf/MEK/MAPK signaling pathways. The growth factor receptor pathways may be constitutively activated as a result of a few aberrations, including the overproduction of ligands, gain-of-function mutations, overexpression and/or gene amplification, and gene rearrangement. Constitutive signal activation leads to oncogenic signaling and resultant uncontrolled proliferation, survival, invasion, and metastasis processes, which are drivers of carcinogenesis. Receptor tyrosine kinases are activated by specific ligands, and activation is followed by homo- or heterodimerization of the receptors [66–69].

The HER2/neu gene is localized on chromosome 17 and encodes a transmembrane tyrosine

**Table 8.2** Some of the regulated miRNAs and their targets in breast cancer

miRNA	Tumor expression level	Validated targets	Pathways
miR-21	Up	<i>BCL2, TPM1, PDCD4, PTEN, MASPIN, RHOB, MMP3</i>	Apoptosis, invasion, metastasis
miR-125b	Down	<i>BAK, HER2, CRAF, MUC1, ERA, RTKN</i>	Proliferation, apoptosis, migration
miR-155	Up	<i>FOXO3A, SOCS1, RHOA</i>	Proliferation, TGF- $\beta$ signaling
miR-145	Down	<i>MUC1, ERA, RTKN</i>	Proliferation, apoptosis, invasion
miR-210	Up	<i>MNT, RAD52</i>	Hypoxia
miR-29c	Up	<i>B7-H3</i>	Metastasis, invasion
miR-100	Down	<i>IGF2, <math>\beta</math>-tubulin</i>	Proliferation, apoptosis
miR-10b	Down	<i>TIAM, HOXD10</i>	Migration, invasion, metastasis
let-7a-2	Down	<i>RAS, CCR7</i>	Proliferation, migration, invasion
miR-205	Down	<i>HER3, VEGFA, EMT</i>	Proliferation, invasion
miR-125b	Down	<i>ERBB2, ERBB3, EPOR, ENPEP, CK2-<math>\alpha</math></i>	Proliferation, apoptosis
miR-196a	Up	<i>ANXA1</i>	Proliferation, apoptosis
miR-497	Down	<i>BCL2, IGF-1R, cyclin E1, RAF-1</i>	Proliferation, apoptosis, invasion, metastasis
miR-181	Up	<i>ATM</i>	Proliferation, migration, invasion, TGF- $\beta$ signaling
miR-31	Down/up	<i>ITGA5, RDX, RHOA</i>	Metastasis
miR-143	Down	<i>ERK5, KRAS</i>	Proliferation, MAPK signaling
miR-191	Up	<i>BDNF, CDK6, SATB1</i>	Proliferation, migration
miR-203	Down/up	<i>SNAI2, BIRC5, LASP1</i>	Proliferation, invasion, apoptosis, migration
miR-29b	Up	<i>PI3K, CDC42, PTEN</i>	Proliferation, migration, invasion, TGF- $\beta$ signaling
miR-93	Up	<i>NRF2, LATS2, STAT3</i>	Proliferation, invasion, metastasis, migration
miR-130b	Down/up	<i>CCNG2</i>	Proliferation

kinase growth factor receptor that has no ligand-binding domain. HER2/neu gene amplification occurs in 30 % of breast cancers. Approximately 100,000 HER2 receptors exist on a normal cell surface, whereas this number reaches 2 million on a breast cancer cell [69, 70]. High levels of aneuploidy, somatic mutations such as the TP53 mutation, FGFR (fibroblast growth factor receptor), EGFR, CDK4 (cyclin-dependent kinase-4), and cyclin D1 amplifications frequently coexist in HER2<sup>+</sup> breast cancer [36]. HER2 overexpression ultimately triggers breast and over cancer progression with a poor prognosis. HER2 is the preferred partner for dimerization with HER1, HER3, and HER4. HER3 lacks a kinase domain. Although ATP binds to the kinase domain of HER3, the phospho-transfer reaction is not driven catalytically. Additionally, HER3 has six

phospho-tyrosine residues at the C-terminal that are available for PI3K binding. Thus, HER3 activation by HER2 recruits the PI3K regulatory subunit to the membrane, which initiates signaling [71]. Accordingly, the HER2-HER3 heterodimer is known to be the most potent oncogenic unit in breast cancer [72–74]. As a consequence of mutations in the extracellular domain of HER3, the oncogenic potential of the heterodimer may increase [74]. HER2 overexpression ultimately activates ligand-independent HER2/HER3/PI3K complex formation and kinase activity in tumor cells, whereas trastuzumab, a monoclonal antibody targeting HER2, destabilizes the complex. Additionally, gain-of-function mutations of PI3K and the loss of PTEN may attenuate the effectiveness of the drug. Resistance to trastuzumab can be circumvented through PI3K inhibition [8, 36].

The activation of alternate receptors, such as IGF-1R and c-met, may also lead to resistance to trastuzumab. Targeting HER2 and EGFR simultaneously may have promise for therapeutic synergy. Tyrosine kinase inhibitors are small-molecule inhibitors that target the cytoplasmic kinase domain of growth factor receptors. Lapatinib, for example, targets HER2 and EGFR simultaneously in breast cancer.

Insulin-like growth factors (IGF-1 and IGF-2) and their receptor IGF-1R expression levels have been associated with breast cancer. The upregulation of IGF-1R results in sustained activation of the PI3K/Akt pathway, thereby leading to resistance to HER2 targeted therapy and antiestrogens.

### PI3K/Akt/mTOR Signaling Pathway in Breast Cancer

The PI3K/Akt/mTOR pathway is the central signaling mechanism downstream of various RTKs (Fig. 8.5). Activating mutations of *Ras* and *PIK3CA*, the gene encoding the catalytic subunit of PI3K, may lead to aberrant signaling that may be independent of ligands binding to the growth factor receptor [8]. However, the inactivation of the PTEN tumor suppressor gene via mutation and/or deletion is widely detected in various types of tumors, including breast cancer. The loss of PTEN causes phosphatidylinositol-3,4,5-triphosphate (PIP<sub>3</sub>) accumulation, thereby activating the PI3K/Akt pathway [75].

In p110 $\alpha$ , the catalytic subunit of PI3K, mutations are observed at a rate of 20–25 % in breast cancer. E542K, E545K, and H1057R mutations are common, and PI3K gains sustained oncogenic activity with no need for RTK activation. The *PIK3CA* mutation may be associated with estrogen and progesterone receptor (ER/PR) status, lymph node metastasis, and ErbB2 overexpression in breast cancer with ultimately coexisting PTEN loss [76]. Activating mutations of genes encoding Akt kinases (*Akt1*, *Akt2*, and *Akt3*) are rare in breast cancer. The activation of Akt leads to the inhibition of FOXO transcription factors via the translocation of these factors to the cytosol. The PI3K/Akt pathway activates the downstream mammalian target of rapamycin


complex1 (mTORC1), thereby activating translation and cell growth. Additionally, the Ras/Raf/MEK pathway may be targeted in combinational therapy strategies.

The PI3K/Akt pathway plays crucial roles both in breast carcinogenesis and in targeted therapy. Thus, advances in our knowledge of agents and/or combinations targeting the pathway in the clinic hold promise for the near future [77] (Fig. 8.5).

### Hypoxia and Breast Cancer

The dysregulated growth of tumor cells leads to decreased O<sub>2</sub> availability in the tumor mass, which triggers angiogenesis. However, the blood vessels that form in tumoral tissues are abnormal because of distorted vasculature, and breast cancers ultimately contain intratumoral hypoxic regions that result in the activation of the hypoxia-inducible factors HIF-1 and HIF-2 [78, 79]. HIF-1 consists of HIF-1 $\alpha$  and HIF-1 $\beta$  subunits. HIF-1 $\alpha$  expression and activation is tightly regulated in an oxygen-dependent manner, whereas HIF-1 $\beta$  is constitutively expressed. There are also mechanisms by which HIF-1 $\alpha$  is regulated in an oxygen-independent manner. In hypoxic cells, HIF-1 $\alpha$  cannot be hydroxylated and degraded, thereby accumulating in the tumor cells. Thus, HIF-1 $\alpha$  and HIF-1 $\beta$  heterodimerize and bind to hypoxia response elements located in the target genes. HIF-2 consists of HIF-2 $\alpha$  and the HIF-1 $\beta$  heterodimer. HIF-1 and HIF-2 regulate over 1000 target genes encoding major proteins, including vascular endothelial growth factor (VEGF), which is critical for angiogenesis, and glycolytic enzymes, which are involved in the metabolic adaptation of tumor cells to hypoxia. HIF-1 $\alpha$  and HIF-2 $\alpha$  levels are found to be increased in breast cancer samples and are linked to disease progression and patient outcome [80, 81]. HIF transcriptional activity also contributes to survival, invasion, metastasis, epithelial-mesenchymal transition, stem cell maintenance, and drug resistance in tumor cells.

In hypoxic breast cancer cells, increased HIF-1 $\alpha$  transcriptional activity activates the expression of L1 cell adhesion molecule



**Fig. 8.5** Activation pathways of growth factor receptor tyrosine kinases. *ErbB1* epidermal growth factor receptor 1, *ErbB2* epidermal growth factor receptor 2, *ErbB3* epidermal growth factor receptor 3, *IGFR1* insulin-like growth factor receptor 1, *IGF1* insulin-like growth factor 1, *Tyr* tyrosine, *p* phosphate, *Grb* growth factor receptor-bound protein, *Sos* son of sevenless, *RAS* rat sarcoma, *RAF* rapidly accelerated fibrosarcoma, *MEK* mitogen-

activated and extracellular-signal-regulated kinase kinase, *ERK* extracellular-signal-regulated kinase, *PI3K* phosphatidylinositol 3-kinase, *PTEN* phosphatase and tensin homolog, *AKT* protein kinase B, *PDK1* phosphoinositide-dependent kinase-1, *mTORC 1/2* mammalian target of rapamycin complex 1/2, *FoxO* forkhead box protein O, *GSK3* glycogen synthase kinase 3, *NF-κB* nuclear factor kappa B

(L1CAM), angiopoietin-like 4 (ANGPTL4), lysyl oxidase (LOX), and LOX-like proteins 2 and 4. L1CAM increases the interaction of tumor cells with endothelial cells (ECs), leading to extravasation and lung metastasis. ANGPTL4 secretion from circulating tumor cells disrupts EC-EC interactions, thereby facilitating the extravasation of these cells into lung parenchyma. Breast cancer cells produce LOX and LOX-like proteins, which are involved in extracellular matrix modeling and the formation of pre-metastatic niches [82–86].

## Angiogenesis

Tumor growth depends on blood vessels and hypoxia acts as an activator of formation of new blood vessels, which is called neoangiogenesis.

In breast cancer, angiogenesis is a key process for invasion and metastasis. Angiogenesis is characterized by abnormal vascular architecture with abnormal function and is sustained mainly by proangiogenic factors in the tumor microenvironment, including vascular endothelial growth factor (VEGF). VEGF is involved in the expression of adhesion molecules, the proliferation of endothelial cells, and increased vascular permeability. VEGFRs also act as receptor tyrosine kinases. Tumor cell-endothelial cell interaction is mediated through the signaling unit composed of the VEGF-A-VEGFR2 interaction [87, 88]. Thus, bevacizumab, a humanized monoclonal antibody directed against VEGF-A, has been studied extensively [89].

In breast cancer, VEGF expression has been shown to be correlated with high histological grade, hormone receptor negativity, HER2 overexpres-

sion, and lymph node metastasis. Consequently, enhanced angiogenesis and metastasis may be correlated with highly proliferative tumors, such as TN/basal-like breast cancer [90, 91].

**Acknowledgment** The author gratefully acknowledges M. Emre Gedik for his valuable assistance.

## References

- McPherson K, Steel CM, Dixon JM. ABC of breast diseases. Breast cancer-epidemiology, risk factors, and genetics. *BMJ*. 2000;321(7261):624–8.
- Bydoun M, Marcato P, Dellaire G. Breast cancer genomics. In: Arceci GDNBJ, editor. *Cancer genomics*. Boston: Academic; 2014. p. 213–32.
- Gayther SA, Pharoah PD, Ponder BA. The genetics of inherited breast cancer. *J Mammary Gland Biol Neoplasia*. 1998;3(4):365–76.
- Fackenthal JD, Olopade OI. Breast cancer risk associated with BRCA1 and BRCA2 in diverse populations. *Nat Rev Cancer*. 2007;7(12):937–48.
- Nathanson KL, Wooster R, Weber BL. Breast cancer genetics: what we know and what we need. *Nat Med*. 2001;7(5):552–6.
- Lakhani SR, Gusterson BA, Jacquemier J, Sloane JP, Anderson TJ, van de Vijver MJ, et al. The pathology of familial breast cancer: histological features of cancers in families not attributable to mutations in BRCA1 or BRCA2. *Clin Can Res*. 2000;6(3):782–9.
- Melhem-Bertrandt A, Bojadziewa J, Ready KJ, Obeid E, Liu DD, Gutierrez-Barrera AM, et al. Early onset HER2-positive breast cancer is associated with germline TP53 mutations. *Cancer*. 2012;118(4):908–13.
- Hynes NE, Dey JH. PI3K inhibition overcomes trastuzumab resistance: blockade of ErbB2/ErbB3 is not always enough. *Cancer Cell*. 2009;15(5):353–5.
- Curigliano G. New drugs for breast cancer subtypes: targeting driver pathways to overcome resistance. *Cancer Treat Rev*. 2012;38(4):303–10.
- Hirshfield KM, Rebbeck TR, Levine AJ. Germline mutations and polymorphisms in the origins of cancers in women. *J Oncol*. 2010;2010:297671.
- Habermann JK, Doering J, Hautaniemi S, Roblick UJ, Bundgen NK, Nicorici D, et al. The gene expression signature of genomic instability in breast cancer is an independent predictor of clinical outcome. *Int J Cancer*. 2009;124(7):1552–64.
- Herman JG, Baylin SB. Gene silencing in cancer in association with promoter hypermethylation. *NEJM*. 2003;349(21):2042–54.
- Baylin SB, Jones PA. A decade of exploring the cancer epigenome – biological and translational implications. *Nat Rev Cancer*. 2011;11(10):726–34.
- Minning C, Mokhtar NM, Abdullah N, Muhammad R, Emran NA, Ali SA, et al. Exploring breast carcinogenesis through integrative genomics and epigenomics analyses. *Int J Oncol*. 2014;45(5):1959–68.
- Yan PS, Shi H, Rahmatpanah F, Hsiau TH, Hsiau AH, Leu YW, et al. Differential distribution of DNA methylation within the RASSF1A CpG island in breast cancer. *Cancer Res*. 2003;63(19):6178–86.
- Esteller M. Epigenetics in cancer. *NEJM*. 2008;358(11):1148–59.
- Esteller M. Relevance of DNA methylation in the management of cancer. *Lancet Oncol*. 2003;4(6):351–8.
- Ellis MJ, Ding L, Shen D, Luo J, Suman VJ, Wallis JW, et al. Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature*. 2012;486(7403):353–60.
- Elston CW, Ellis IO, Pinder SE. Pathological prognostic factors in breast cancer. *Critical Rev Oncol/Hematol*. 1999;31(3):209–23.
- Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature*. 2012;490(7418):61–70.
- Dawson SJ, Rueda OM, Aparicio S, Caldas C. A new genome-driven integrated classification of breast cancer and its implications. *EMBO J*. 2013;32(5):617–28.
- Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, et al. Repeated observation of breast tumor subtypes in independent gene expression data sets. *Proc Natl Acad Sci U S A*. 2003;100(14):8418–23.
- Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. *Nature*. 2000;406(6797):747–52.
- Sandhu R, Parker JS, Jones WD, Livasy CA, Coleman WB. Microarray-based gene expression profiling for molecular classification of breast cancer and identification of new targets for therapy. *Lab Med*. 2010;41:364–72.
- Prat A, Parker JS, Karginova O, Fan C, Livasy C, Herschkowitz JI, et al. Phenotypic and molecular characterization of the claudin-low intrinsic subtype of breast cancer. *Breast Cancer Res*. 2010;12(5):R68.
- Prat A, Perou CM. Deconstructing the molecular portraits of breast cancer. *Mol Oncol*. 2011;5(1):5–23.
- Albain KS, Barlow WE, Shak S, Hortobagyi GN, Livingston RB, Yeh IT, et al. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol*. 2010;11(1):55–65.
- Mook S, Schmidt MK, Viale G, Pruneri G, Eekhout I, Floore A, et al. The 70-gene prognosis-signature predicts disease outcome in breast cancer patients with 1–3 positive lymph nodes in an independent validation study. *Breast Cancer Res Treat*. 2009;116(2):295–302.
- Knauer M, Mook S, Rutgers EJ, Bender RA, Hauptmann M, van de Vijver MJ, et al. The predictive value of the 70-gene signature for adjuvant chemotherapy in early breast cancer. *Breast Cancer Res Treat*. 2010;120(3):655–61.



30. Sanchez-Navarro I, Gamez-Pozo A, Pinto A, Hardisson D, Madero R, Lopez R, et al. An 8-gene qRT-PCR-based gene expression score that has prognostic value in early breast cancer. *BMC Cancer*. 2010;10:336.
31. Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T, et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol*. 2009;27(8):1160–7.
32. Paik S, Tang G, Shak S, Kim C, Baker J, Kim W, et al. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol*. 2006;24(23):3726–34.
33. Gatz ML, Silva GO, Parker JS, Fan C, Perou CM. An integrated genomics approach identifies drivers of proliferation in luminal-subtype human breast cancer. *Nat Genet*. 2014;46(10):1051–9.
34. Staaf J, Ringner M, Vallon-Christersson J, Jonsson G, Bendahl PO, Holm K, et al. Identification of subtypes in human epidermal growth factor receptor 2-positive breast cancer reveals a gene signature prognostic of outcome. *J Clin Oncol*. 2010;28(11):1813–20.
35. Khoury T, Kanehira K, Wang D, Ademuyiwa F, Mojica W, Cheney R, et al. Breast carcinoma with amplified HER2: a gene expression signature specific for trastuzumab resistance and poor prognosis. *Mod Pathol*. 2010;23(10):1364–78.
36. Arteaga CL, Engelman JA. ERBB receptors: from oncogene discovery to basic science to mechanism-based cancer therapeutics. *Cancer Cell*. 2014;25(3):282–303.
37. Arribas J, Baselga J, Pedersen K, Parra-Palau JL. p95HER2 and breast cancer. *Cancer Res*. 2011;71(5):1515–9.
38. Turner NC, Reis-Filho JS. Basal-like breast cancer and the BRCA1 phenotype. *Oncogene*. 2006;25(43):5846–53.
39. Yau C, Esserman L, Moore DH, Waldman F, Sninsky J, Benz CC. A multigene predictor of metastatic outcome in early stage hormone receptor-negative and triple-negative breast cancer. *Breast Cancer Res*. 2010;12(5):R85.
40. Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, et al. Identification of human triple-negative breast cancer subtypes and pre-clinical models for selection of targeted therapies. *J Clin Invest*. 2011;121(7):2750–67.
41. Shah SP, Roth A, Goya R, Oloumi A, Ha G, Zhao Y, et al. The clonal and mutational evolution spectrum of primary triple-negative breast cancers. *Nature*. 2012;486(7403):395–9.
42. Seal MD, Chia SK. What is the difference between triple-negative and basal breast cancers? *Cancer J*. 2010;16(1):12–6.
43. Vuong D, Simpson PT, Green B, Cummings MC, Lakhani SR. Molecular classification of breast cancer. *Virchows Arch*. 2014;465(1):1–14.
44. Sotiriou C, Wirapati P, Loi S, Harris A, Fox S, Smeds J, et al. Gene expression profiling in breast cancer: understanding the molecular basis of histologic grade to improve prognosis. *J Nat Cancer Inst*. 2006;98(4):262–72.
45. Stephens PJ, Tarpey PS, Davies H, Van Loo P, Greenman C, Wedge DC, et al. The landscape of cancer genes and mutational processes in breast cancer. *Nature*. 2012;486(7403):400–4.
46. Bertucci F, Orsetti B, Negre V, Finetti P, Rouge C, Ahomadegbe JC, et al. Lobular and ductal carcinomas of the breast have distinct genomic and expression profiles. *Oncogene*. 2008;27(40):5359–72.
47. Wang C, Machiraju R, Huang K. Breast cancer patient stratification using a molecular regularized consensus clustering method. *Methods*. 2014;67(3):304–12.
48. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004;116:281–97.
49. Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, et al. MicroRNA expression profiles classify human cancers. *Nature*. 2005;435:834–8.
50. Andorfer CA, Necela BM, Thompson EA, Perez EA. MicroRNA signatures: clinical biomarkers for the diagnosis and treatment of breast cancer. *Trends Mol Med*. 2011;17(6):313–9.
51. Nygren MK, Tekle C, Ingebrigtsen VA, Makela R, Krohn M, Aure MR, et al. Identifying microRNAs regulating B7-H3 in breast cancer: the clinical impact of microRNA-29c. *Br J Cancer*. 2014;110(8):2072–80.
52. Gebeshuber CA, Martinez J. miR-100 suppresses IGF2 and inhibits breast tumorigenesis by interfering with proliferation and survival signaling. *Oncogene*. 2013;32(27):3306–10.
53. Lobert S, Jefferson B, Morris K. Regulation of beta-tubulin isoforms by micro-RNA 100 in MCF7 breast cancer cells. *Cytoskeleton*. 2011;68(6):355–62.
54. Shen L, Li J, Xu L, Ma J, Li H, Xiao X, et al. miR-497 induces apoptosis of breast cancer cells by targeting Bcl-w. *Exp Ther Med*. 2012;3(3):475–80.
55. Li D, Zhao Y, Liu C, Chen X, Qi Y, Jiang Y, et al. Analysis of MiR-195 and MiR-497 expression, regulation and role in breast cancer. *Clin Cancer Res*. 2011;17(7):1722–30.
56. Nagpal N, Ahmad HM, Molparia B, Kulshreshtha R. MicroRNA-191, an estrogen-responsive microRNA, functions as an oncogenic regulator in human breast cancer. *Carcinogenesis*. 2013;34(8):1889–99.
57. Ding X, Park SI, McCauley LK, Wang CY. Signaling between transforming growth factor beta (TGF-beta) and transcription factor SNAI2 represses expression of microRNA miR-203 to promote epithelial-mesenchymal transition and tumor metastasis. *J Biol Chem*. 2013;288(15):10241–53.
58. Zhang Z, Zhang B, Li W, Fu L, Fu L, Zhu Z, et al. Epigenetic silencing of miR-203 upregulates SNAI2 and contributes to the invasiveness of malignant breast cancer cells. *Genes Cancer*. 2011;2(8):782–91.
59. Shaker O, Maher NA, Nassar Y, Morcos G, Gad Z. Role of microRNAs -29b-2, -155, -197 and -205 as diagnostic biomarkers in serum of breast cancer females. *Gene*. 2015;560(1):77–82.

60. Zhao H, Wilkie T, Deol Y, Sneh A, Ganju A, Basree M, et al. miR-29b defines the pro-/anti-proliferative effects of S100A7 in breast cancer. *Mol Cancer*. 2015;14(1):11.
61. Chang Y-Y, Kuo W-H, Hung J-H, Lee C-Y, Lee Y-H, Chang Y-C, Lin W-C, et al. Deregulated microRNAs in triple-negative breast cancer revealed by deep sequencing. *Mol Cancer*. 2015;14:36.
62. Neel JC, Lebrun JJ. Activin and TGFbeta regulate expression of the microRNA-181 family to promote cell migration and invasion in breast cancer cells. *Cell Signal*. 2013;25(7):1556–66.
63. Ferracin M, Bassi C, Pedriali M, Pagotto S, D'Abundo L, Zagatti B, et al. miR-125b targets erythropoietin and its receptor and their expression correlates with metastatic potential and ERBB2/HER2 expression. *Mol Cancer*. 2013;12(1):130.
64. Feliciano A, Castellvi J, Artero-Castro A, Leal JA, Romagosa C, Hernandez-Losa J, et al. miR-125b acts as a tumor suppressor in breast tumorigenesis via its novel direct targets ENPEP, CK2-alpha, CCNJ, and MEGF9. *PLoS One* 2013;8(10):e76247.
65. Yager JD, Davidson NE. Estrogen carcinogenesis in breast cancer. *NEJM*. 2006;354(3):270–82.
66. Blume-Jensen P, Hunter T. Oncogenic kinase signaling. *Nature*. 2001;411(6835):355–65.
67. Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell*. 2000;103(2):211–25.
68. Hubbard SR, Miller WT. Receptor tyrosine kinases: mechanism of activation and signaling. *Curr Opin Cell Biol*. 2007;19(2):117–23.
69. Yarden Y, Pines G. The ERBB network: at last, cancer therapy meets system biology. *Nat Rev Cancer*. 2012;12(8):553–63.
70. Eccles SA. The epidermal growth factor receptor/ ErbB/HER family in normal and malignant breast biology. *Int J Develop Biol*. 2011;55(7–9):685–96.
71. Gala K, Chandralapaty S. Molecular pathways: HER3 targeted therapy. *Clin Cancer Res*. 2014;20(6):1410–6.
72. Amin DN, Campbell MR, Moasser MM. The role of HER3, the unpretentious member of the HER family, in cancer biology and cancer therapeutics. *Semin Cell Dev Biol*. 2010;21(9):944–50.
73. Roskoski R. The ErbB/HER family of protein-tyrosine kinases and cancer. *Pharmacol Res*. 2014;79:34–74.
74. Jaiswal BS, Kljavin NM, Stawiski EW, Chan E, Parikh C, Durinck S, et al. Oncogenic ERBB3 mutations in human cancers. *Cancer Cell*. 2013;23(5):603–17.
75. Carnero A. The PKB/Akt pathway in cancer. *Curr Pharm Des*. 2010;16(1):34–44.
76. Saal LH, Holm K, Maurer M, et al. PIK3CA mutations correlate with hormone receptors, node metastasis, and ErbB2, and are mutually exclusive with PTEN loss in human breast carcinoma. *Cancer Res*. 2005;65(7):2554–9.
77. Thorpe LM, Yuzugullu H, Zhao JJ. PI3K in cancer: divergent roles of isoforms, modes of activation and therapeutic targeting. *Nat Rev Cancer*. 2015;15:7–24.
78. Wang GL, Semenza GL. Purification and characterization of hypoxia-inducible factor 1. *J Biol Chem*. 1995;270:1230–7.
79. Semenza GL. Hypoxia inducible factors in physiology and medicine. *Cell*. 2012;148:399–408.
80. Yamamoto Y, Ibusuki M, Okumura Y, Kawasoe T, Kai K, et al. Hypoxia-inducible factor 1 $\alpha$  is closely linked to an aggressive phenotype in breast cancer. *Breast Cancer Res Treat*. 2008;110:465–75.
81. Helczynska K, Larsson AM, Holmquist Mengelbier L, Bridges E, Fredlund E, Borgquist S, et al. Hypoxia-inducible factor 2 $\alpha$  correlates to distant recurrence and poor outcome in invasive breast cancer. *Cancer Res*. 2008;68:9212–20.
82. Zhang H, Wong CC, Wei H, Gilkes DM, Korangath P, Chaturvedi P, et al. HIF-1-dependent expression of angiopoietin-like 4 and L1CAM mediates vascular metastasis of hypoxic breast cancer cells to the lungs. *Oncogene*. 2012;31:1757–70.
83. Erler JT, Bennewith KL, Nicolau M, Dornhöfer N, Kong C, Le QT, et al. Lysyl oxidase is essential for hypoxia-induced metastasis. *Nature*. 2006;440:1222–6.
84. Erler JT, Bennewith KL, Cox TR, Lang G, Bird D, Koong A, et al. Hypoxia-induced lysyl oxidase is a critical mediator of bone marrow cell recruitment to form the premetastatic niche. *Cancer Cell*. 2009;15:35–44.
85. Perryman L, Erler JT. Lysyl oxidase in cancer research. *Future Oncol*. 2014;10(9):1709–17.
86. Barker HE, Cox TR, Erler JT. The rationale for targeting the LOX family in cancer. *Nat Rev Cancer*. 2012;12:540–52.
87. Dvorak HF. Vascular permeability factor/vascular endothelial growth factor: a critical cytokine in tumor angiogenesis and a potential target for diagnosis and therapy. *J Clin Oncol*. 2002;20:4368–80.
88. Sitohy B, Nagy JA, Dvorak HF. Anti-VEGF/VEGFR therapy for cancer: reassessing the target. *Cancer Res*. 2012;72(8):1909–14.
89. Tredan O, Lacroix-Triki M, Guiu S, Mouret-Reynier MA, Barriere J, et al. Angiogenesis and tumor microenvironment: bevacizumab in the breast cancer model. *Target Oncol*. 2014;10(2):189–98. [Epub ahead of print]
90. Ribeiro-Silva A, Ribeiro do Vale F, Zucoloto S. Vascular endothelial growth factor expression in the basal subtype of breast carcinoma. *Am J Clin Pathol*. 2006;125:512–8.
91. Linderholm BK, Hellborg H, Johansson U, Elmberger G, Skoog L, Lehtiö J, et al. Significantly higher levels of vascular endothelial growth factor (VEGF) and shorter survival times for patients with primary operable triple-negative breast cancer. *Ann Oncol*. 2009;20:1639–46.

---

# Clinical Aspects of Estrogen and Progesterone Receptors and ERBB2 Testing

9

Ebru Sari and Suayib Yalcin

---

## Abstract

Human breast cancers depend on estrogen and/or progesterone for growth, and these effects are mediated through estrogen receptors (ERs) and progesterone receptors (PRs), respectively. The human epidermal growth factor receptor 2 (HER2) gene encodes a member of the epidermal growth factor receptor family of receptor tyrosine kinases, and its amplification with resultant overexpression plays a major role in sustaining multiple pathways in cancer growth. ERs, PRs, and HER2 status are the most important molecular markers in the standard care of all primary and recurrent/metastatic breast cancer patients and play both predictive and prognostic roles. The responsiveness of a tumor to hormone therapy is an important parameter in breast cancer management in both adjuvant and metastatic settings. Only breast cancers with HER2 amplification or overexpression respond to HER2-directed therapies. Hormonal status of the tumor is prognostic for patient outcome and site of metastasis. Hormonal receptor-positive disease represents an indolent and slowly growing tumor with longer time to disease recurrence. HER2 is a poor prognostic factor in the absence of HER2-directed therapies. Assessment of the ER/PR/HER2 status is an essential factor in the evaluation of every newly diagnosed breast cancer, and standardization of the assay methods is crucial.

---

## Keywords

Breast cancer • Estrogen • Progesterone • HER2 testing • Clinical aspects

---

E. Sari, MD  
Division of Medical Oncology, Diskapi Yildirim  
Beyazit Treatment and Research Hospital,  
İrfan Baştuğ Caddesi, Diskapi, Ankara, Turkey

S. Yalcin, MD (✉)  
Department of Medical Oncology, Hacettepe  
University Cancer Institute, Ankara 06100, Turkey  
e-mail: [syalcin@hacettepe.edu.tr](mailto:syalcin@hacettepe.edu.tr)

## Introduction

Breast cancer comprises a heterogeneous group of tumors with a wide spectrum of morphologically and molecularly different subtypes, resulting in different biological behaviors, presentation, and prognosis. The major issue in making treatment decisions is to identify the subgroup of patients who will particularly benefit from a given treatment; this concept is now being developed as precision medicine. Aside from the stage of the disease, the evidence of distant metastasis, the organ of distant metastasis, and the age, performance status, and menopausal status of the patient, the histology, grade, and molecular pattern of the tumor are also important in deciding which treatment modality is the best for an individual patient. Among the molecular alterations associated with breast cancer, estrogen receptors (ERs), progesterone receptor (PRs), and v-erb-b2 erythroblastic leukemia viral oncogene homolog 2 (ERBB2) and neuroblastoma/glioblastoma-derived oncogene homolog which is also called human epidermal growth factor receptor (HER)2 status are the most important molecular markers and are firmly established in the standard care of all primary and recurrent/metastatic breast cancer patients.

Here, we will review the clinical utility of hormonal receptor and ERBB2/HER2 testing in breast cancer, how this information is translated to treatment decision-making, the valid assays for these markers, and the guidelines for testing these important biological markers.

---

## Hormonal Receptors

### Historical Information

Breast cancer is a hormone-dependent malignancy. It arises from breast tissue that is normally responsive to endogenous hormones, and its course can often be influenced by the administration or removal of these hormones [1]. In 1896, Beatson et al. first reported dramatic regression of breast cancer after bilateral oophorectomy in a premenopausal woman with advanced disease [2].

Unfortunately, oophorectomy is not effective in all patients. In 1900, Stanley Boyd [3] reported that only one-third of patients responded to ovarian ablation and that responses lasted an average of 1–2 years. Although these data were initially disappointing, endocrine therapy became a standard of care in the treatment of breast cancer. However, because only one-third of patients responded, the question was which patients would respond to endocrine therapy and whether a marker could be used to predict this response. Identifying this marker would allow the avoidance of ineffective ablative surgery in some patients [4].

Estrogenic hormones produced in the ovary were discovered by Allen and Doisy [5]. Then, in the early 1960s, radiolabeled estrogens were first observed to be preferentially concentrated in estrogen target organs. These observations gave rise to the concept of an “estrogen receptor,” opening the door to molecular targeting in the treatment and prevention of breast cancer [4].

The finding that estrogen target tissues contained ERs and that nontarget tissues did not led to the question of whether these concepts could translate to the clinic to predict the endocrine responsiveness of breast cancer. The fact that only one in three women responds to any form of endocrine ablative therapy meant that two women were undergoing surgery and intensive medical care without any hope of a favorable outcome. Thus, if ERs are necessary for estrogen-stimulated growth, evaluating the presence or absence of ERs in a tumor specimen may be informative [4].

Jensen et al. reported in 1971 that the measurement of ER levels could predict the response to hormone therapy [6]. In general, these data were consistent with all of the clinical correlations presented at a 1974 workshop in Bethesda, Maryland, sponsored by the Breast Cancer Task Force [7]. Of the patients evaluated, only 8 % of ER-negative tumors responded to additive or ablative therapy, whereas ~60 % of patients who were ER positive had an objective response to endocrine therapy. The results of the 1974 workshop established the ER assay as a valuable predictive test for the endocrine treatment of advanced breast cancer [4].

The identification of the ER has not only proved to be a successful therapeutic target for the treatment and prevention of breast cancer but has also proved to be a selective molecular model for all subsequent efforts to design targeted therapeutics in cancer [4].

## Biology of ERs and PRs

Human breast cancers depend on estrogen and/or progesterone for growth, and this effect is mediated through ERs and PRs. ERs are ligand-regulated receptors that belong to the steroid nuclear receptor family. They function as transcription factors and transduce hormonal signals into a large variety of physiological responses in various organs [8]. The two structurally related ERs, ER $\alpha$  and ER $\beta$ , are the products of two separate genes that are differentially expressed in tissues. ER $\alpha$  is responsible for estrogen-induced mitogenic signaling in epithelial cells in breast, uterine, and ovarian tissues [9]. In the normal mammary gland, estradiol binds to both ER $\alpha$  and ER $\beta$  to control cell proliferation and differentiation [10]. The two ER isoforms are expressed at similarly low levels in the normal breast, whereas breast cancer cells express more ER $\alpha$  than ER $\beta$  [11]. The key components of ERs are the DNA-binding domain, which binds with high affinity and specificity to DNA sequences (estrogen response elements (EREs)) to regulate the transcription rates of target genes, and the ligand-binding domain, which binds estrogens [12]. The two ERs and the PR also form complexes with a number of coregulatory proteins that coordinately act to influence the transcription of estrogen-responsive genes. Mechanisms regulating ER $\alpha$  and ER $\beta$  function can occur at three levels: differential translation of exons, splicing of their messenger RNA (mRNA), and posttranslational modifications.

The effects of estrogen are mediated not only through nuclear ERs but also through cytoplasmic/membrane ERs and G protein-coupled ERs [11]. The ER proteins are generally believed to shuttle between the cytoplasm and the nucleus, and in vitro experiments have demonstrated that

ligand-free ER $\alpha$  is maintained in a non-DNA-binding form in a multichaperone complex organized around Hsp90 (heat shock protein 90) [13]. ER-mediated transcription is a highly complex process involving multiple coregulatory factors and cross talk between different signaling pathways [11].

## Genomic ER-Mediated Transcription Mechanisms

ERs and PRs function as transcription factors in the nucleus (Fig. 9.1). In the absence of hormones, histone deacetylase (HDAC) and the receptor corepressors NCoR and SMRT are bound to the receptor, forming a multichaperone complex. DNA is wrapped tightly. Estradiol binding to ER $\alpha$  activates the receptor through phosphorylation and dissociates proteins such as HSP90 [14]; during this process, the receptor undergoes conformational changes. Several kinases in the growth factor signaling networks can also activate the ER and its coregulatory proteins, and this process is termed ligand-independent activation [15]. The ER then binds to the 13-base pair ERE sequence within the promoter. ER dimers dynamically and sequentially recruit various regulatory protein complexes that contribute to chromatin remodeling, thereby strongly enhancing transcriptional activity [16]. The nuclear receptor coactivators that are associated with ERs include the general transcription factor P300/CBP. P300/CBP is ubiquitously expressed and serves as a coadaptor between nuclear receptors and DNA. P300/CBP plays a critical role in cell cycle regulation, cell differentiation, and apoptosis, and it exhibits histone acetyltransferase (HAT) activity [17, 18]. Importantly, HATs are required for the full activation of ER-mediated transcription. Methyltransferases, including CARM1 and PRMT1, are also ER $\alpha$ -associated coactivators. Members of the p160 protein family, namely, steroid receptor coactivator 1 (SRC1), SRC2, and SRC3, also play various roles [19]. After ligand activation, protein-protein interactions with other transcription factors can also result in gene regulation by indirect binding to DNA (outside EREs) [20].





**Fig. 9.1** Genomic ER-mediated transcription mechanisms. (a) Classical genomic ligand-dependent mechanisms; (b) ligand-independent genomic mechanisms.

Abbreviations: *ER* Estrogen receptor, *GFR* Growth factor receptor, *Hsp90* Heat shock protein 90, *TF* Transcription factor, *ERE* Estrogen response element, *CoA* Coactivator

Estrogen-regulated promoters are affected by dynamic and complex processes and are involved in chromatin remodeling. Consequently, these events contribute to the hormonal regulation of gene expression. Phosphorylation of coactivators can affect ER-dependent transcription, even in the absence of ligands (ligand-independent ER-mediated transcription) or in the presence of antiestrogens, by increasing their subcellular nuclear localization, their interaction with the ER, and their ability to recruit transcriptional coregulators such as the CBP/P300 coactivator to the receptor-promoter complex [11].

Estradiol-ER $\alpha$  complexes affect the transcription of genes that are involved in proliferation, differentiation, survival, stimulation of invasion, metastasis and angiogenesis. Some of these genes, like those involved in cell cycle progression (such as c-myc, cyclins D, A, and E), are activated [21].

Consequently, the growth of ER $\alpha$ -expressing (ER+) cells from breast tumors is estradiol dependent, and the removal of estradiol leads to regression. The presence of ER $\beta$  inhibits both ER $\alpha$ -mediated transcription and estradiol-induced proliferation in various types of cancer cells [22–24]. Therefore, ER $\beta$  in breast cancer lesions may be associated with more benign tumors. ER $\alpha$  and ER $\beta$  differentially regulate both the proliferation and apoptosis of normal mammary epithelial cells [25]. The ER $\alpha$ /ER $\beta$  ratio is currently hypothesized to be a key element in the regulation of estradiol activity in breast cancer cells [26].

### Posttranslational Modifications

Posttranslational modifications of ERs and PRs include phosphorylation, ubiquitylation, acetylation, and methylation. Among the multiple

kinases that can phosphorylate ER $\alpha$  are p38 mitogen-activated protein kinase (MAPK), cyclin A-CDK2, CDK7, c-Src, and pp90rsk1 [27–31]. ER $\alpha$  can also be phosphorylated by signaling molecules such as Akt (also known as protein kinase B), extracellular regulated kinase (Erk) 1/2, MAPK, protein kinase A (PKA), and p21-activated kinase 1 (PAK-1), thereby producing various responses to ligands [32–35]. Phosphorylation of the ER results in a variety of activities in different tumors. These effects involve receptor turnover, cellular localization, and transcriptional activity and are complex and interdependent.

The PR can be phosphorylated, with distinct phosphorylation sites coordinately regulated by ligands or kinases [36]. Phosphorylation of the PR, similar to ER $\alpha$ , can regulate ligand hypersensitivity [37].

### Mutations in the ER Gene

A few mutations have been reported in the ER $\alpha$  gene, including a somatic mutation that causes a single amino acid change in the ER $\alpha$  hinge domain (lysine 303 to arginine, called K303R ER $\alpha$ ). The receptor resulting from this mutation is hypersensitive to the growth-promoting effects of estrogen. Tumors with this mutation are associated with older age, larger tumor size, more lymph node-positive disease, and poor outcomes. Its incidence differs among studies [38–40]. The K303R mutation is also significantly associated with a first-degree family history of breast cancer [41].

### Nongenomic Pathways

In addition to classical ER genomic activity, which alters gene expression in the nucleus, estrogens have a more rapid class of effects (Fig. 9.2), for which plasma membrane estrogen binding sites have been described [42]. Many studies have shown that, in response to their ligands, ERs and PRs can mediate signaling cascades originating from the membrane or the cytoplasm through direct interaction with signal-transduction mediators [43]. This nongenomic ER action (also called rapid, nonnuclear, or nonclassical) occurs within seconds to a few minutes and is independent of gene transcription [44].


A small subpopulation of the classic ER $\alpha$  and  $\beta$  subtypes that are located outside the nucleus or closely related nonclassical short forms of the ER transduce rapid estrogen signaling [45–47]. Membrane receptors distinct from the classic ER, especially G protein-coupled receptor 30 (GPR30), may also contribute to the rapid effects of estrogen [48, 49]. Membrane and cytoplasmic ERs transmit their signals through kinase cascades. These cascades include growth factor receptors and cellular tyrosine kinases as well as their downstream pathways. Membrane and cytoplasmic ERs also use calcium, cyclic adenosine monophosphate (cAMP), and other second messengers for signal transduction. This signaling leads to rapid cellular responses without gene transcription, but they also regulate nuclear transcription [43, 47, 50–52]. In addition, ERs have a role in the mitochondria, where they mediate cell survival signaling [47, 53, 54].

For progesterone-induced rapid effects, distinct nonclassical and classical receptor forms are present in the membrane and the cytoplasm [55].

### ER Cross Talk with Other Signal-Transduction Pathways

In breast tumors, estrogens promote the activity of growth factor signaling pathway components, including ligands (e.g., transforming growth factor (TGF) $\alpha$ , insulin-like growth factor (IGF)-II), and receptors (EGF and IGF-I receptors) and key signal-transduction molecules (e.g., insulin receptor substrate-1) while also reducing the expression of growth-inhibitory factors (e.g., TGF $\beta$ ) and inhibiting expression of tyrosine phosphatases. Altogether, these estrogens lead to a net increase in growth factor mitogenic activity. These observations imply a positive feedback loop that augments essential signaling pathways of both the estrogen/ER and growth factors/receptor systems [50, 56, 57].

In vitro studies have shown that membrane ERs can activate various growth factor receptors, including EGFR, HER2, and IGF-IR [47, 50, 58]. This pathway involves sequential activation of G proteins, the cellular tyrosine kinase c-Src, and matrix metalloproteinases (MMPs), followed by the release of heparin-binding EGF-like growth



**Fig. 9.2** Nongenomic ER-mediated transcription mechanisms, Abbreviations: *ER* Estrogen receptor, *GPR30* G protein-coupled receptor 30, *GFR* Growth factor receptor,

*MMP* Matrix metalloproteinase, *cAMP* Cyclic adenosine monophosphate, *MAPK* Mitogen-activated protein kinase, *Erk* Extracellular regulated kinase, *TF* Transcription factor

factor (HB-EGF). HB-EGF then binds to and activates adjacent EGFR and its downstream kinase cascades (e.g., Ras/Mek/MAPK and PI3K/Akt) [50]. These cascades then activate transcription [57]. Thus, genomic and nongenomic downstream pathways of ERs potentiate each other, resulting in proliferation and cell survival.

### The Clinical Importance of ERs and PRs in Breast Cancer

Human breast cancers are dependent upon estrogen and/or progesterone for growth, and this effect is mediated through ERs and PRs. Up to two-thirds of invasive breast cancers of women younger than 50 years of age are ER or PR positive, and approximately 80 % of tumors in women older than 50 years of age are ER positive [59]. Measurement of these receptors has become a routine part of the evaluation of breast cancers.

The responsiveness of a tumor to hormone therapy is an important parameter in breast can-

cer management. However, not all patients with breast cancer benefit from hormone therapy. Tumor expression of ERs and/or PRs can best identify those women who are most likely to benefit from hormone therapy. Tumors that are negative for ERs and PRs are unlikely to respond to hormone therapy and respond better to cytotoxic chemotherapy. ERs and PRs can be utilized as both predictive and prognostic factors. A prognostic factor is any parameter available at the time of diagnosis that correlates with disease-free or overall survival. Thus, it is indicative of the inherent biological aggressiveness of a tumor and is correlated with the natural history of the disease. A predictive factor indicates the likelihood of a response to a given therapy (here, ER and PR/hormone therapy).

We also know that adjuvant hormone therapy can halve the recurrence rate of patients with ER+ breast cancer [60]. Hormone therapy is relatively nontoxic, and responses can last for many years in some patients with metastatic disease. Thus, hormone therapies offer many significant

advantages to particular subsets of breast cancer patients. The measurement of ER and PR levels in patients can identify those tumors that are most likely to benefit from hormonal agents.

ER status is strongly influenced by tumor grade and histology [59]. In one previous study, grade 1 tumors having histologies of pure tubular, colloid and classic lobular carcinoma were ER positive [61]. Patients with ER+/PR+ tumors have a better prognosis than patients with ER+/PR- tumors, who in turn have a better prognosis than patients with ER-/PR- tumors [62]. ER expression is associated with most known prognostic factors but not with nodal metastases [63]. Thus, ER status is an important marker of growth rate rather than metastatic potential.

### **ER and PR Expression as Predictive Factors for Hormone Therapy**

#### **Efficacy of ER and PR Expression in Predicting the Benefit of Hormone Therapy in the Advanced Disease Setting**

Approximately 30–40 % of patients with ER+ metastatic disease will respond to first-line hormone therapies, and another 20 % will experience disease stabilization [64–67]. The small proportion of patients who respond to hormone therapy with ER- disease may be mostly due to false-negative receptor assay results. All metastatic patients who receive hormone therapy show progression, and they often respond to a second line of hormone therapy [68]. Responses to subsequent lines of hormone therapy decline gradually but remain in the range of 20–30 %, and this response is also dependent on ER positivity. Hormone therapy provides good palliation, better quality of life, and improved survival [69, 70]. Beyond simply designating a result as “positive,” the expression level of the ER in the tumor is also very important in endocrine responsiveness. Response rates to hormone therapies are directly correlated to the level of ER expression [71, 72]. As with ERs, increasing PR levels are also associated with better response, longer time to treatment failure, and longer survival [73, 74]. Although the ER and PR expression levels are correlated, PR positivity is correlated with higher response rates independent

of ER. ER+/PR+ tumors have higher response rates than ER+/PR- tumors [73]. As a result, PR status provides important information about the responsiveness to hormone therapies.

Discordance between the hormone receptor status of the relapse/metastases and the primary tumor is an important issue. A conversion rate of 20–30 % has been reported from ER+ to ER- status, with less frequent conversion reported for ER- to ER+ status [75–79]. A significantly shorter median survival was noted with the loss of the ER in the metastasis in one of these studies [77]. Because metastases are the targets of therapy, the hormone receptor status of the metastases should be more predictive than that of the primary tumor. A previous study showed that 74 % of patients with concordant ER+ results between primary tumor and metastases responded to hormone therapy, whereas only 12 % of patients with ER+ primary tumors and ER-metastases responded [76]. A separate study that investigated biopsies from patients who developed resistance to tamoxifen reported changes in hormone receptor status and HER2 expression [80]. PR expression in metastatic lesions can also show discordance from the primary tumor. As with ERs, most of these conversions are of PR+ tumors losing PR expression [77, 78]. Sequential biopsies from patients given subsequent lines of hormone therapy have shown that loss of PR expression is associated with poor survival [81]. The metastatic tumor’s hormone receptor status is a better predictor of survival than the primary tumor’s hormone receptor status. Beyond the probable technical causes of false-negative or false-positive results, several explanations have been suggested for this discordance. Possible explanations include intratumoral heterogeneity, which can lead to clonal selection of different clones with distinct hormone receptor properties that can change over time; changes within single cells themselves as an adaptive mechanism to prior treatments and selection of more resistant clones; and tumor dedifferentiation with the development of metastasis [79, 82, 83]. Therefore, before making treatment decisions, the molecular markers of breast cancer should be retested in the metastatic lesions when possible.

### **Efficacy of ER and PR Expression in Predicting the Benefit of Adjuvant Hormonal Treatment**

The Oxford meta-analysis reviewed data from 48,000 women from randomized clinical trials with 15–20 years of follow-up. This meta-analysis showed that the benefit of 5 years of adjuvant tamoxifen treatment is dependent on the ER and PR status of the tumor. ER status was a significant predictive factor for endocrine therapy [60]. More recent large prospective trials confirmed that tamoxifen reduces the risk of distant relapse, death, and contralateral breast cancer in ER+ breast cancer [84, 85].

More recently, aromatase inhibitors have been studied in adjuvant treatment of postmenopausal patients either before, immediately after, or long after tamoxifen. For example, a TransATAC trial retrospectively collected the samples of monotherapy arms of an ATAC trial and then tested ER expression in a central laboratory. This study showed a marginally significant relationship between ER level and time to recurrence. This effect was significant in anastrozole-treated patients ( $p=0.0009$ ) but less significant in tamoxifen-treated patients ( $p=0.078$ ). There was no significant interaction between ER level and benefit of anastrozole over tamoxifen treatments [86]. ER expression of tumors in patients from the BIG1–98 trial monotherapy arms (letrozole versus tamoxifen) was also analyzed centrally. Disease-free survival (DFS) varied based on ER expression levels [87].

However, meta-analyses of early breast cancer adjuvant tamoxifen therapy have not shown any benefit of PR expression among ER+ patients. An exception was a small subgroup of patients with ER-/PR+ tumors, who also showed benefits from hormone therapy. However, ER-/PR- patients have not shown any benefit [60, 88]. Several studies have suggested that tumors in this small subset are biologically different from ER+/PR+ tumors and represent a group of tumors with worse clinicopathologic features and clinical outcome. However, whether this group truly exists or whether it represents tumors that are actually ER+, with technical challenges producing this misclassification, remains unknown [89–92]. The discrepancies in

these results may exist because relatively few studies included measurements of PR in the meta-analysis, and there may be PR measurement errors. A large retrospective study of patients receiving adjuvant tamoxifen showed that patients who had ER+/PR+ tumors experienced a 15–30 % lower risk of recurrence and death than patients who had ER+/PR- tumors [62]. Other large studies of adjuvant tamoxifen have also shown that both ER and PR expressions are predictors of the benefits of hormone therapy [90, 93, 94]. Recently, the MA.17 trial, which randomized postmenopausal women after 5 years of tamoxifen to the aromatase inhibitor letrozole versus placebo, also found that the benefit of letrozole over placebo was confined to ER+/PR+ tumors and was not seen in the ER+/PR- tumors [95].

Other than ERs, many cellular signaling networks contribute to endocrine responsiveness. These molecular pathways can modulate the ER pathway or provide alternative mitogenic and survival stimuli for the cells. Thus, resistance to hormone therapy may develop in ER+ tumors. Therefore, ER positivity alone is not sufficient for response to endocrine therapies. Some multigene predictive scores have been developed to predict hormonal responsiveness. One example is the Oncotype DX 21 gene assay, which includes several downstream ER-regulated genes and several proliferation genes in addition to ER mRNA [96].

### **Efficacy of ER and PR Expression in Predicting the Benefit of Adjuvant Chemotherapy**

Luminal A (ER/PR+, HER2-) type tumors receive less benefit from adjuvant chemotherapy, even if they have high risk of recurrence, because of lymph node positivity. The studies showing the benefit of adding taxanes to adjuvant treatment of node-positive breast cancer showed benefits irrespective of hormonal receptor status, but they did not take into account the endocrine effects of chemotherapy-induced amenorrhea in premenopausal patients [97–99]. Hormone receptor-positive breast cancer patients benefit less from adjuvant chemotherapy than hormone receptor-negative patients [100]. A study comparing adjuvant TAC



(docetaxel-adriamycin-cyclophosphamide) with FAC (fluorouracil-adriamycin-cyclophosphamide) showed a benefit of adding taxanes regardless of ER status [98]. Premenopausal patients constituted more than half of the patients in this study, and TAC was associated with a greater incidence of chemotherapy-induced amenorrhea than was FAC. Thus, the advantage of adding a taxane may have been due to its endocrine effects by inducing amenorrhea.

The meta-analysis of adjuvant chemotherapy trials supports that ER- tumors derive more benefit from chemotherapy than ER+ tumors. The ovarian ablative effects of chemotherapy are not observed in postmenopausal patients. In postmenopausal patients, the benefits of chemotherapy in reducing recurrence and mortality of ER- patients are more prominent. However, the meta-analysis still showed a significant benefit of adjuvant chemotherapy in ER+ patients, but the ER level and the PR and HER2 status were not included in the meta-analysis [60].

More recent trials (Cancer and Leukemia Group B (CALGB) and Breast Cancer Intergroup trials) also evaluated the effects of ER status on patient outcome. They compared the effect of adding taxanes, dose-dense regimens, or increasing doses of doxorubicin with standard AC (adriamycin-cyclophosphamide). Only ER- patients showed a benefit from increasing the dose of doxorubicin, adding taxanes, or using dose-dense adjuvant chemotherapy in terms of relative risk of recurrence and mortality. These aggressive adjuvant chemotherapy regimens tended to benefit ER+ patients, but the effect was not significant [101]. When both ER and HER2 status are considered, patients with HER2-, ER+ breast cancer had no benefit from the addition of paclitaxel to adjuvant AC chemotherapy. However, addition of paclitaxel was beneficial in HER2+ patients, regardless of ER status [102].

Neoadjuvant chemotherapy trials have also shown the effect of ER status in pathologic complete response (pCR) rates. pCR rates in the ER- groups are significantly higher than in ER+ patients [103].

Multigene profiles, such as the Oncotype DX 21 gene profile, provide additional information for highly endocrine-sensitive tumors and identify those patients who will not benefit from chemotherapy. Patients with ER+ tumors who also have a low risk 21-gene recurrence score receive no benefit from CMF (cyclophosphamide-methotrexate-fluorouracil) chemotherapy [104, 105]. In another study, postmenopausal patients with ER+ tumors and a low risk score with the 21-gene assay but having node positivity showed no benefit from adding chemotherapy (FAC-cyclophosphamide-doxorubicin-fluorouracil) to tamoxifen [106].

### **ER and PR Expression Levels as Prognostic Factors**

Multiple studies show a relation between the hormonal status of the tumor and patient outcome. Patients with stage I ER+ breast cancer who receive no systemic therapy after surgery have a 5–10 % lower likelihood of recurrence at 5 years in comparison with ER- patients [107, 108]. Studies with longer follow-up show that as more time passes, the difference in the rate of relapse and death significantly diminishes and eventually disappears [109–111]. ER expression is associated with a number of other well-established prognostic indicators but not with nodal metastases [63], and its prognostic significance reduces over time. ER status does not predict the metastatic potential of the tumor. However, ER expression predicts an indolent and slowly growing tumor with longer time to disease recurrence. ER+ tumors are more frequently found in older patients [59, 112], and they are well differentiated histologically [113]. They have a lower fraction of dividing cells [114], are more often diploid, and are less likely to exhibit a mutation, loss, high expression, or amplification of breast cancer-related genes such as TP53 [115], HER2 [92], or EGFR [92]. ER status is also prognostic for the site of metastasis. ER+ tumors more frequently metastasize to the bone, soft tissue, or the reproductive and genital tracts, whereas ER- tumors metastasize more often metastasize to visceral organs or the brain [116, 117].

## Methods for Measuring ER and PR Status

Assessment of ER and PR status is an essential factor in the evaluation of every newly diagnosed breast cancer. Various assay methods have been used to measure ER values in breast cancer specimens. These tests should be accurate and reliable because the results of such testing help to direct therapy. Thus, women who might benefit from endocrine therapy are provided with treatment, and those who are unlikely to derive benefit do not receive these treatments and are not exposed to unnecessary side effects [118].

The dextran-coated charcoal/ligand-binding assay (DCC/LBA) was the first method that became the standard for ER detection and measurement. Other assays, such as enzyme immunoassay and enzyme-linked immunosorbent assay (ELISA), then became available. By the late 1980s and early 1990s, immunohistochemistry (IHC) of formalin-fixed paraffin-embedded (FFPE) specimens began to replace the DCC assay because IHC has distinct advantages over DCC. These advantages include the need for smaller amounts of tissue, the ability to conduct testing on FFPE tissue (obviating the need for fresh/frozen tissue), the ability to correlate staining with histology, and the storage and retrieval of archived slides for later testing [118]. However, much data underlying the knowledge that the relationship between the response to hormone therapy and the amount of ER expression in the tumor tissue are based on studies using LBAs, and the cutoff values of positive and negative ER status were developed with reference to LBA results. IHC methods have been subject to retrospective comparisons with established methods such as LBA, and published reports have also indicated that IHC may be more predictive than LBAs in identifying patients who will derive benefit from hormone therapy [72, 119–122].

IHC is performed by first treating thin sections of tissue using a variety of antigen retrieval methods. Next, the tissue is incubated with a primary antibody directed against the ER or PR. Then, secondary detection systems (secondary antibodies) that are conjugated to an enzyme (e.g.,

horseradish peroxidase) can be used to amplify the chromogenic signal. The sections are counterstained and viewed microscopically.

ER and PR levels can also be determined by the evaluation of messenger RNA either by individual assay (northern blot analysis or reverse transcriptase polymerase chain reaction (RT-PCR)) or as part of multigene expression assays such as Oncotype DX and MammaPrint [104, 123]. However, due to the scarcity of data directly correlating these results with clinical outcomes, it is too early to recommend their routine use.

## Optimizing IHC

Optimization includes evaluating the effects of preanalytic variables (i.e., variables of testing, involving the collection, fixation, and storage of samples), analytic variables (i.e., variables associated with the method of testing itself), thresholds to define results, and postanalytic variables (i.e., variables associated with handling of results, such as reporting) [118].

The American Society of Clinical Oncology (ASCO)/Collage of American Pathologists (CAP) reported the latest guidelines for ER and PR testing in 2010 to provide clarification about testing parameters and establish mandatory proficiency testing and inspection criteria to improve the accuracy of these tests [124].

Regarding IHC assays for ERs and PRs, no gold-standard assay is available. A relevant standard would be any assay whose specific preanalytic and analytic components conformed exactly to assays whose results had been validated against clinical benefit from hormone therapy. ER and PR status should be determined in all newly diagnosed invasive breast cancers and recurrent and metastatic lesions, if appropriate. Newly diagnosed cases of ductal carcinoma in situ (DCIS) are also commonly tested for ER and PR expression because a retrospective subset analysis of the National Surgical Adjuvant Breast and Bowel Project B-24 clinical trial showed a significant reduction in subsequent breast cancer diagnosis restricted to patients with ER-positive DCIS [124].

Because the results may vary substantially between laboratories because of differences in

specimen handling, tissue fixation, antigen retrieval, and antibody type, standardization of these variables is necessary.

### Standardization of the Assays

A sample is accepted as positive for ER or PR expression if  $\geq 1\%$  of tumor cell nuclei are immunoreactive. Both the average intensity (weak, moderate, strong) and extent of staining (as a percentage) are reported. A sample is accepted as negative for ER or PR expression if  $< 1\%$  of tumor cell nuclei are immunoreactive in the presence of evidence that the sample expresses ER or PR in intrinsic controls. A sample is uninterpretable for ER or PR expression if there are no immunoreactive tumor nuclei and the internal epithelial elements (internal control) lack any nuclear staining [124].

### Optimal Preanalytic Standardization

#### Tissue Handling

Issues related to testing variation begin as soon as the breast tissue is removed from the patient. Both the warm and cold ischemic times are important variables in the analysis of labile macromolecules such as proteins, RNA, and DNA from clinical tissue samples. The time from the interruption of the blood supply to the tumor by the surgeon to the excision of the tissue specimen is defined as warm ischemia, and the time from excision of the tissue to the initiation of fixation is defined as cold ischemia. Studies have documented the progressive loss of these labile molecules after the surgical interruption of blood flow, through tissue ischemia, acidosis, and enzymatic degradation [125, 126]. Breast resection specimens should be fixed as quickly as possible in an adequate volume of fixative (optimally 10-fold greater than the volume of the specimen). The time of tissue collection (defined as the time when the tissue is handed from the surgical field) and the time the tissue is placed in fixative should be recorded. The time from tumor removal to fixation should be kept to  $\leq 1$  h [124].

After being received in the pathology laboratory, specimens should be oriented and carefully inked for surgical margin assessment, sectioned

at 5 mm intervals, and placed in 10 % neutral (phosphate) buffered formalin (NBF). If the excised specimen was obtained remotely from the grossing laboratory, the sample should be bisected through the tumor and promptly placed in NBF prior to transport. Although less optimal than immediate gross examination of the fresh sample by the pathologist, this process is preferable to storage of the sample in the refrigerator unfixed or in fixative without sectioning [124].

#### Type of Fixative

Only 10 % NBF should be used as the fixative for breast tissue specimens. Higher or lower concentrations of NBF are not acceptable [124].

#### Duration of Tissue Fixation

Breast tissue specimens must be fixed in 10 % NBF for no less than 6 h and for not more than 72 h before processing [127, 128]. Formalin penetrates tissue at a rate of approximately 1 mm/h, which is why breast excision samples must be incised in a timely fashion to initiate formalin fixation throughout the tissue. Fixation does not begin until formaldehyde has penetrated into the tissue. However, permeation of tissue by formalin is not the same as the chemical reaction of fixation, which involves protein cross-linking by formaldehyde. Underfixation of breast tissue may lead to false-negative ER results. Overfixation is likely to be less problematic than underfixation, but it can also potentially lead to false-negative results due to excessive protein cross-linking by formaldehyde [124, 127, 129].

### Analytic Standardization

#### Antibody Selection for ER and PR Testing

Testing for ER and PR expression by IHC must be performed using antibodies that have been clinically validated and show good correlation with the outcomes of patients receiving endocrine treatment. An ASCO/CAP panel provided a list of antibodies with acceptable clinical validation. Antibodies sold for research use only or investigational use only or developed by the testing facility may not be used in ER and PR testing [124]. A laboratory performing ER testing should

initially validate its proposed or existing assay against one of the clinically validated assays and demonstrate acceptable concordance. To be considered acceptable, the results of the assay must be initially 90 % concordant with those of the clinically validated assay for the ER+ and PR+ category and 95 % concordant for the ER- or PR- category [124].

#### Control Samples for IHC Assays

External and internal controls can be used to ensure that the IHC test has been performed properly. Positive and negative external controls should be included in every batch of assays. Acceptable external controls are cell lines previously defined as highly positive, intermediate and negative for the analyzed receptor to ensure that assay is working properly. Internal control defines the staining of normal epithelial elements of the analyzed tissue to ensure that handling procedures are performed properly. If an external or internal control does not produce the expected reaction, the result of patient testing must not be reported, and retesting is warranted [124].

#### Postanalytic Standardization

The percentage of cells with nuclear staining is reported by either estimation or quantitation. Quantitation may be performed either manually or by image analysis. The entire slide should be reviewed to assess the tumor-containing areas. Cytology samples with limited tumor cells and little tumor staining must have at least 100 cells counted. If cytoplasmic staining occurs, the assay should be repeated or performed on another sample. The sample should be rejected if there are obscuring artifacts such as decalcification of the sample or staining only of necrotic debris. If the test result is negative but that particular histologic type of breast cancer is unlikely to be ER- (tubular, mucinous, or lobular morphology or Nottingham score of 1), the tumor should also be subjected to confirmatory testing, such as sending the same specimen to a reference laboratory for retesting or by repeating the assay on another block or on a separate breast cancer specimen [124]. A comprehensive quality control program for ER/PR IHC analyses should be established.

This analysis should include the analysis of whether positive results in a given period could reflect the ER+ breast cancers in the patient population served by the laboratory [124].

---

## HER2 Testing

HER2 encodes a member of the epidermal growth factor receptor family of receptor tyrosine kinases, and its amplification with resultant overexpression plays a major role in sustaining multiple pathways in cancer growth. These roles include self-sufficiency in growth signals, sustained angiogenesis, increased cell division, and enhanced invasion [130, 131].

The HER2 gene encodes a 185 kDa protein that has tyrosine kinase activity. HER2 is a membrane protein that is expressed at low levels in all epithelial cells in normal fetal and adult tissues [132]. The HER2 gene is amplified in a variable percentage of human breast [133, 134], ovarian [134], bladder, endometrial [135], salivary gland [136], and gastric [137] cancers. HER2 gene amplification has been associated with increased levels of expression of HER2 mRNA and protein product [134].

## HER2 Signal-Transduction Pathways

The HER family consists of four structurally related members: the epidermal growth factor receptor (EGFR/HER1), HER2, HER3, and HER4, which are transmembrane receptor tyrosine kinases that regulate cell growth, survival, differentiation, migration, and other cellular responses. HER family receptors are activated by ligand-induced dimerization or receptor pairing [138]. They are able to homodimerize or heterodimerize with other HER family members, producing multiple receptor combinations. Dimerization leads to activation of the intrinsic tyrosine kinase domain of the receptor. Phosphorylation of tyrosine residues leads to the activation of downstream signaling pathways such as the RAS/RAF/MEK/ERK pathway and the PI3K/Akt/mTOR pathway [139]. HER2 and

HER3 are unique in that HER2 has no known ligand that stimulates its activity, and HER3 has very little enzymatic kinase activity and mainly functions as a ligand-activated dimer partner for the other family members [140]. HER heterodimers are more potent in signal transduction than are homodimers. Heterodimerization follows a strict hierarchical principle, with HER2 representing the preferred dimerization partner for all other HER receptors [141]. HER2-containing heterodimers display increased potency due to the relatively slow rate of ligand dissociation and slow rate of receptor internalization. Thus, signaling by HER2-containing dimers is prolonged and results in enhanced activation of signaling pathways. The most potent heterodimer is HER2/HER3 [142].

In addition to their function as receptors on the cell surface, HER family proteins are present in the nucleus to act as kinases and transcriptional regulators. EGFR can be transported to the nucleus and act as a tyrosine kinase. HER2 can be transported to the nucleus via endocytic vesicles. The intracellular domain of HER4 is cleaved and transported to the nucleus [143–145].

### **Clinical Significance of HER2 Amplification/Overexpression**

HER2 is amplified in approximately 15–20 % of breast cancers [146]. HER2 overexpression is associated with important clinical outcomes in breast cancer patients. HER2 overexpression is a poor prognostic factor in the absence of adjuvant treatment independent of tumor size, grade and hormone receptor status. It has been associated with shorter DFS and shorter overall survival (OS) in primary invasive node-negative breast cancer patients treated with surgery alone [147]. HER2 is also an important predictive marker for certain treatments.

### **Response to HER2-Targeted Therapies**

Much clinical evidence has shown that only breast cancers with HER2 amplification or overexpression respond to HER2-directed therapies such as monoclonal antibodies directed against

HER2 [148, 149]. Agents that target HER2 are remarkably effective in both metastatic and adjuvant settings. Trastuzumab, a humanized monoclonal antibody, improves response rates, time to progression and survival when used alone or added to chemotherapy in metastatic breast cancer [150]. Further, prospective randomized trials have shown that adjuvant trastuzumab reduces the risk of recurrence and mortality in patients with early-stage breast cancer [151–155]. Other HER2-targeted drugs, including the tyrosine kinase inhibitor lapatinib, the antibody pertuzumab, and the antibody drug conjugate ado-trastuzumab emtansine (T-DM1), improve outcomes in HER2-positive metastatic breast cancer [156–158].

### **Response to Hormone Therapies**

HER2 positivity may be associated with resistance to hormone therapies. Although this association remains controversial, some data suggest that this effect may be specific to selective estrogen modulator (SERM) therapy such as tamoxifen and perhaps not to estrogen depletion therapies such as aromatase inhibitors [86].

Preclinical studies suggest physiological cross talk between the HER2 and ER signal-transduction pathways. HER2 expression in human breast cancer cells is downregulated by estrogens [159]. Conversely, transfection and overexpression of HER2 in vitro promotes estrogen-independent growth and tamoxifen resistance in ER+ human breast cancer cells [160, 161]. Amplification and/or overexpression of HER2 may be associated with primary resistance to hormone therapy. The hypothesized mechanisms underlying the hormone independence of HER2-expressing cells include phosphorylation of the ER, ligand-independent ER activation, and regulation of hormone receptor expression [161, 162].

Some retrospective and nonrandomized clinical studies showed lower response rates of hormone therapy in HER2-overexpressing tumors. However, randomized trials in either adjuvant or metastatic settings failed to provide supporting evidence. Although some trials have concluded that patients with HER2-overexpressing tumors



are relatively resistant to adjuvant hormone therapy [163–166], a similar number could not demonstrate such a relationship [167–170].

The adverse influence of HER2 expression in response to hormone therapies is hypothesized to be limited to therapies based on a ligand-binding agent such as SERMs. Further, the response to ligand-depleting therapies such as ovarian ablation or aromatase inhibitors is not affected by HER2 overexpression. In a study of neoadjuvant endocrine therapy analyzing 250 postmenopausal women receiving either letrozole or tamoxifen for 4 months, women in the tamoxifen group with HER2+ tumors showed significantly lower response rates than those with HER2- tumors (17 % versus 40 %, respectively). However, patients receiving letrozole showed no difference in response rates between HER2+ and HER2- groups (69 % versus 53 %, respectively) [171]. Multiple studies have assessed the response rates to hormone therapy among women with HER2+ versus HER2- tumors. Although the results are mixed, most show an adverse influence of HER2 overexpression in response to treatment with tamoxifen [172–176]. However, these data may not be sufficient for using HER2 status as a factor for hormonal treatment selection (aromatase inhibitors versus tamoxifen) [177, 178].

### Response to Chemotherapy

HER2 may be associated with either sensitivity or resistance to some chemotherapeutic agents. HER2 positivity is associated with better outcomes in response to adjuvant anthracycline-containing regimens in most studies [179–181]. A meta-analysis of eight trials comparing anthracycline- versus non-anthracycline-based chemotherapy concluded that the benefits of adjuvant anthracycline use were confined to women with HER2+ tumors [182]. This effect may be secondary to coamplification of HER2 with topoisomerase II (topo2), which is the direct target of anthracyclines [183]. In a study involving patients with HER2+ metastatic breast cancer who were treated with chemotherapy with or without trastuzumab, topo2 $\alpha$  coamplification was associated with significantly improved survival when only anthracycline-containing chemotherapy was used

for treatment, compared with outcomes in HER2+ cancers lacking topo2 $\alpha$  coamplification [184]. Another meta-analysis of five randomized trials comparing adjuvant anthracycline-containing treatment with CMF (cyclophosphamide-methotrexate-fluorouracil) assessed whether HER2 expression or topo2 expression could predict response to anthracyclines [185]. Trastuzumab was not administered to patients in any of the trials included in this analysis. There was a significant improvement in both event-free survival and OS in patients with HER2 overexpression, and similar results were observed for patients with topo2 alterations. In patients without HER2 overexpression and in patients with normal topo2 levels, only improvement in event-free survival was observed, with no improvement in OS. Now, because adjuvant trastuzumab is routinely used in HER2-overexpressing breast cancer patients and because both agents are cardiotoxic, the results are more conflicting. Although individual studies of an anthracycline-based regimen plus trastuzumab versus a non-anthracycline-containing regimen plus trastuzumab have shown equivalent survival outcomes for HER2+ patients, these data are controversial [151, 184]. Many clinicians still believe that anthracyclines provide additional benefits in combination with trastuzumab for patients with HER2+ breast cancer [186]. Cardiac toxicity is a serious concern in patients with early-stage disease who are treated with both trastuzumab and anthracycline-containing chemotherapy regimens. For these reasons, accurate determination of HER2 alterations in breast carcinomas is crucially important.

HER2 amplification or overexpression has been correlated with responsiveness to paclitaxel-containing chemotherapy [102], but the data are inconsistent and contradictory. In a study of metastatic breast cancer patients who received either paclitaxel or cyclophosphamide plus epirubicin, women whose tumors were HER2+ had a significantly longer progression-free survival and OS with a taxane-containing regimen, while there were no differences between the two regimens in those whose tumors were HER2 negative [187]. In the adjuvant setting, the interaction between

HER2 status and benefit from taxanes was evaluated in a subset of 1322 women in the Cancer and Leukemia Group B (CALGB) 9344 study. This study showed that four cycles of adjuvant paclitaxel after AC significantly improved DFS and OS compared with four cycles of AC alone in women with node-positive breast cancer. There was a significant benefit from the addition of paclitaxel after AC in women with HER2+ tumors regardless of ER status and in ER- tumors regardless of HER2 status. However, there was no evidence of benefit in the subgroup of patients with ER+ and HER2- tumors [102]. Other trials conducted in the adjuvant setting have failed to note an association between HER2 overexpression and benefit from taxanes [188, 189]. Reports from neoadjuvant taxane therapy are conflicting. One suggests a worse response rate [190], and two others report no association between HER2 gene amplification or overexpression and response [191, 192].

## Methods for Measuring HER2

Because HER2 gene amplification is directly correlated with HER2 expression levels at the mRNA and protein levels, HER2 status could potentially be evaluated at any of these levels and should correspond to the HER2 status determined using any of the other measures.

Several assays have been used for HER2 determination in tissue. The studies that led to the approval of the humanized mouse 4D5 monoclonal antibody trastuzumab in HER2+ metastatic breast cancer selected the patients with an IHC assay method, known as the Clinical Trials Assay (CTA) [150, 193]. This assay used two different antibodies, 4D5 (the mouse monoclonal antibody used to produce humanized trastuzumab) and CB11 (a mouse monoclonal antibody). However, these so-called home-brew assays were not validated and are not considered appropriate for commercialization. Since the development of the CTA, a number of commercially available testing kits have received approval from FDA for the assessment of patients for whom trastuzumab may be a suitable treatment [194].

The available assays are as follows:

- Overexpression of the HER2 protein product: Western blotting, ELISA, or IHC
- Overexpression of HER2 RNA: Northern blotting or reverse transcription polymerase chain reaction (RT-PCR)
- HER2 gene amplification by in situ hybridization (ISH): Fluorescence in situ hybridization (FISH), chromogenic in situ hybridization (CISH), silver-enhanced in situ hybridization (SISH), or differential PCR

Because there is insufficient evidence to support the use of mRNA and DNA microarray assays to determine HER2 status in unselected patients, these techniques will not be discussed here.

Recommendations for tissue handling as well as preanalytic, analytic, and postanalytic factors in ER/PR testing are also suitable for HER2 testing. Therefore, these points will not be discussed again.

## Immunohistochemistry Tests

Grading of IHC assays is based on a 0, 1+, 2+, and 3+ scoring system. Tumor specimens that demonstrate complete, strong circumferential membrane staining in >10 % of tumor cells are classified as 3+ on IHC and constitute a positive result. Specimens showing strong circumferential membrane staining in <10 % and incomplete and/or weak/moderate circumferential membrane staining of >10 % of tumor cells are classified as 2+ and require confirmation of HER2 status by an alternative method, usually ISH. If there is faint or barely perceptible and incomplete membrane staining in >10 % of tumor cells, the sample is classified as IHC 1+, and this finding constitutes a negative HER2 result. If there is either no staining at all or there is faint or barely perceptible incomplete membrane staining in <10 % of tumor cells, the score is IHC 0 and is reported as HER2 negative [146].

The HercepTest, an IHC test that was approved by FDA at the same time as trastuzumab, uses a rabbit polyclonal antibody against HER2. Two other FDA-approved IHC systems are the Pathway anti-HER2/neu test (Ventana Medical Systems Inc., Tucson, AZ), which utilizes a rabbit monoclonal antibody (4B5), and the Bond

Oracle HER2 IHC System (Leica Biosystems, Newcastle upon Tyne, UK), which uses the mouse CB11 clone [195].

### In Situ Hybridization

HER2 FISH assays were initially approved by the FDA for the identification of women with node-negative disease who were at high risk of recurrence or disease-related death, as well as for selection for doxorubicin chemotherapy [196]. Later, the selection criteria were expanded to cover the assessment of patients for whom trastuzumab treatment was being considered.

When measured against an external “gold-standard” molecular characterization of HER2 status, FISH is more accurate, reproducible, and robust than IHC [196]. However, for practical and historical reasons, IHC has been more widely used as the primary test for HER2 status. IHC is comparatively quick and is viewed using a conventional bright-field microscope, and stained tissues do not degrade over time [197]. IHC testing also permits parallel viewing of tumor morphological features. In contrast, FISH is technically more demanding and requires the use of fluorescence microscopy [197].

Emerging technologies for HER2 testing include CISH and SISH. CISH uses a peroxidase enzyme-labeled probe for chromogenic detection by diaminobenzidine. SISH uses the same system with a silver-based detection system. Because these processes do not use fluorescent dye, a standard bright-field microscope can be used [197]. Automated ISH techniques may enable more rapid testing. CISH can be stored because the signal is stable [197]. An additional emerging technology is dual-color, dual-hapten, bright-field in situ hybridization (DDISH), which has all the advantages of SISH. Further, SISH requires two separate slides to detect both HER2 and CEP17, whereas DDISH uses double-stranded probes labeled with two haptens to detect both markers on a single slide [198].

When interpreting ISH results, both the average HER2 copy number signals per cell and ratio between HER2 and the chromosome 17 enumeration probe (CEP17) should be considered. When HER2 testing is performed by a validated single-

probe ISH assay, HER2 is positive if the average HER2 copy number is  $\geq 6.0$  signals/cell; HER2 is equivocal if the average HER2 copy number is  $\geq 4.0$  and  $< 6.0$  signals/cell; and HER2 expression is negative if the average HER2 copy number is  $< 4.0$  signals/cell. If equivocal, a reflex test must be ordered, either with the same specimen using dual-probe ISH or using IHC or with a new test using a new specimen if available [146].

When HER2 testing is performed by a validated dual-probe ISH assay, a positive ISH result is defined as a HER2 copy number of  $\geq 6.0$  per cell or a HER2/CEP17 ratio  $\geq 2.0$ . The equivocal result is defined as a copy number of  $\geq 4$  to  $< 6$  signals per cell and a ratio of  $< 2$ . This result should trigger an alternative test for HER2 (same specimen using IHC or an alternate ISH method or a new test using a different specimen). The ISH result is negative if the HER2/CEP17 ratio is  $< 2.0$  and the average HER2 copy number per cell is  $< 4.0$  [146].

### Key Recommendations from the 2013 ASCO/CAP Guidelines for HER2 Testing

If available, the test should be performed on a core biopsy specimen from the newly diagnosed patient. If the test result is clearly positive or clearly negative, no retesting is needed. If the test is negative and there is apparent histopathologic discordance or if specimen handling has not been consistent with guideline recommendations, a section of the tumor from the excisional specimen should be retested. If the result is positive, no further testing is needed. However, if the test is negative and there remains significant clinical concern about the result after consultation between the pathologist and the medical oncologist, it may be appropriate to repeat the test in a different block from the patient's tumor [146].

The HER2 test result must be reported as indeterminate if technical issues prevent the test from being reported as positive, negative, or equivocal. These conditions include inadequate specimen handling, artifacts (crush or edge artifacts) that make interpretation difficult, or analytical testing failure. Under these circumstances, another specimen should be requested [146].

Laboratories performing these tests should follow all accreditation requirements, one of

which is the initial testing validation. This requirement conforms to the 2010 ASCO/CAP recommendations for ER/PR testing [146].

Laboratories are responsible for ensuring the reliability and accuracy of their testing results by complying with accreditation and proficiency testing requirements for HER2 testing results. They should review and document external and internal controls with each test and each batch of tests [146].

### Histopathologic Features Suggestive of Possible HER2 Test Discordance

- A new HER2 test should be ordered if the initial HER2 test was positive in any of the following:
  1. Histologic grade 1 carcinoma of infiltrating ductal or lobular carcinoma, ER and PR positive
  2. Tubular/mucinous/cribriform (at least 90 % pure) adenoid cystic carcinoma (90 % pure), often triple negative [146]
- A new HER2 test should be ordered on the excision specimen if the initial HER2 test result in a core needle biopsy specimen of a primary breast cancer is negative in a tumor exhibiting any of the following:
  1. Grade 3 tumor.
  2. Small amount of invasive tumor in the core biopsy.
  3. The carcinoma upon definitive resection contains a high-grade carcinoma that is morphologically distinct from that in the core biopsy.
  4. The core biopsy result is equivocal for HER2 after testing by both ISH and IHC.
  5. There is doubt about the specimen handling of the core biopsy [146].

### References

1. Zumoff B, Fishman J, Bradlow HL, Hellman L. Hormone profiles in hormone-dependent cancers. *Cancer Res.* 1975;35:3365–73.
2. Stockwell S. Classics in oncology. George Thomas Beatson, M.D. (1848–1933). *CA Cancer J Clin.* 1983;33:105–21.

3. Boyd S. On oophorectomy in cancer of breast. *Br Med J.* 1900;2:1961–7.
4. Jensen EV, Jordan VC. The estrogen receptor: a model for molecular medicine. *Clin Cancer Res.* 2003;9:1980–9.
5. Allen E, Doisy EA. Landmark article Sept 8, 1923. An ovarian hormone. Preliminary report on its localization, extraction and partial purification, and action in test animals. By Edgar Allen and Edward A. Doisy. *JAMA.* 1983;250:2681–3.
6. Jensen EV, Block GE, Smith S, Kyser K, DeSombre ER. Estrogen receptors and breast cancer response to adrenalectomy. *Natl Cancer Inst Monogr.* 1971;34:55–70.
7. McGuire WLC, Carbone PP, Wollmer EP. Estrogen receptors in human breast cancer. New York: Raven; 1975.
8. Ascenzi P, Bocedi A, Marino M. Structure-function relationship of estrogen receptor alpha and beta: impact on human health. *Mol Aspects Med.* 2006;27:299–402.
9. Ali S, Coombes RC. Estrogen receptor alpha in human breast cancer: occurrence and significance. *J Mammary Gland Biol Neoplasia.* 2000;5:271–81.
10. Helguero LA, Faulds MH, Gustafsson JA, Haldosen LA. Estrogen receptors alpha (ERalpha) and beta (ERbeta) differentially regulate proliferation and apoptosis of the normal murine mammary epithelial cell line HC11. *Oncogene.* 2005;24:6605–16.
11. Renoir JM, Marsaud V, Lazennec G. Estrogen receptor signaling as a target for novel breast cancer therapeutics. *Biochem Pharmacol.* 2013;85:449–65.
12. Klinge CM. Estrogen receptor interaction with estrogen response elements. *Nucleic Acids Res.* 2001;29:2905–19.
13. Pratt WB, Toft DO. Steroid receptor interactions with heat shock protein and immunophilin chaperones. *Endocr Rev.* 1997;18:306–60.
14. Osborne CK, Schiff R. Estrogen-receptor biology: continuing progress and therapeutic implications. *J Clin Oncol.* 2005;23:1616–22.
15. Osborne CK, Schiff R, Fuqua SA, Shou J. Estrogen receptor: current understanding of its activation and modulation. *Clin Cancer Res.* 2001;7:4338s–42; discussion 411s–2s.
16. Metivier R, Penot G, Hubner MR, Reid G, Brand H, Kos M, et al. Estrogen receptor-alpha directs ordered, cyclical, and combinatorial recruitment of cofactors on a natural target promoter. *Cell.* 2003;115:751–63.
17. Lonard DM, O'Malley BW. Nuclear receptor coregulators: judges, juries, and executioners of cellular regulation. *Mol Cell.* 2007;27:691–700.
18. Ogryzko VV, Schiltz RL, Russanova V, Howard BH, Nakatani Y. The transcriptional coactivators p300 and CBP are histone acetyltransferases. *Cell.* 1996;87:953–9.
19. O'Malley BW. Coregulators: from whence came these "master genes". *Mol Endocrinol.* 2007;21:1009–13.

20. Heger Z, Guran R, Zitka O, Beklova M, Adam V, Kizek R. In vitro interactions between 17beta-estradiol and DNA result in formation of the hormone-DNA complexes. *Int J Environ Res Public Health*. 2014;11:7725–39.
21. Musgrove EA, Caldon CE, Barraclough J, Stone A, Sutherland RL. Cyclin D as a therapeutic target in cancer. *Nat Rev Cancer*. 2011;11:558–72.
22. Paruthiyil S, Parmar H, Kerekatte V, Cunha GR, Firestone GL, Leitman DC. Estrogen receptor beta inhibits human breast cancer cell proliferation and tumor formation by causing a G2 cell cycle arrest. *Cancer Res*. 2004;64:423–8.
23. Strom A, Hartman J, Foster JS, Kietz S, Wimalasena J, Gustafsson JA. Estrogen receptor beta inhibits 17beta-estradiol-stimulated proliferation of the breast cancer cell line T47D. *Proc Natl Acad Sci U S A*. 2004;101:1566–71.
24. Gougelet A, Bouclier C, Marsaud V, Maillard S, Mueller SO, Korach KS, et al. Estrogen receptor alpha and beta subtype expression and transactivation capacity are differentially affected by receptor-, hsp90- and immunophilin-ligands in human breast cancer cells. *J Steroid Biochem Mol Biol*. 2005;94:71–81.
25. Grober OM, Mutarelli M, Giurato G, Ravo M, Cicatiello L, De Filippo MR, et al. Global analysis of estrogen receptor beta binding to breast cancer cell genome reveals an extensive interplay with estrogen receptor alpha for target gene regulation. *BMC Genomics*. 2011;12:36.
26. Matthews J, Gustafsson JA. Estrogen signaling: a subtle balance between ER alpha and ER beta. *Mol Interv*. 2003;3:281–92.
27. Arnold SF, Obourn JD, Jaffe H, Notides AC. Phosphorylation of the human estrogen receptor by mitogen-activated protein kinase and casein kinase II: consequence on DNA binding. *J Steroid Biochem Mol Biol*. 1995;55:163–72.
28. Rogatsky I, Trowbridge JM, Garabedian MJ. Potentiation of human estrogen receptor alpha transcriptional activation through phosphorylation of serines 104 and 106 by the cyclin A-CDK2 complex. *J Biol Chem*. 1999;274:22296–302.
29. Chen D, Washbrook E, Sarwar N, Bates GJ, Pace PE, Thirunuvakkarasu V, et al. Phosphorylation of human estrogen receptor alpha at serine 118 by two distinct signal transduction pathways revealed by phosphorylation-specific antisera. *Oncogene*. 2002;21:4921–31.
30. Lee H, Bai W. Regulation of estrogen receptor nuclear export by ligand-induced and p38-mediated receptor phosphorylation. *Mol Cell Biol*. 2002;22:5835–45.
31. Joel PB, Smith J, Sturgill TW, Fisher TL, Blenis J, Lannigan DA. pp90rsk1 regulates estrogen receptor-mediated transcription through phosphorylation of Ser-167. *Mol Cell Biol*. 1998;18:1978–84.
32. Chen D, Pace PE, Coombes RC, Ali S. Phosphorylation of human estrogen receptor alpha by protein kinase A regulates dimerization. *Mol Cell Biol*. 1999;19:1002–15.
33. Cui Y, Zhang M, Pestell R, Curran EM, Welshons WV, Fuqua SA. Phosphorylation of estrogen receptor alpha blocks its acetylation and regulates estrogen sensitivity. *Cancer Res*. 2004;64:9199–208.
34. Wang RA, Mazumdar A, Vadlamudi RK, Kumar R. P21-activated kinase-1 phosphorylates and transactivates estrogen receptor-alpha and promotes hyperplasia in mammary epithelium. *EMBO J*. 2002;21:5437–47.
35. Likhite VS, Stossi F, Kim K, Katzenellenbogen BS, Katzenellenbogen JA. Kinase-specific phosphorylation of the estrogen receptor changes receptor interactions with ligand, deoxyribonucleic acid, and coregulators associated with alterations in estrogen and tamoxifen activity. *Mol Endocrinol*. 2006;20:3120–32.
36. Clemm DL, Sherman L, Boonyaratankornkit V, Schrader WT, Weigel NL, Edwards DP. Differential hormone-dependent phosphorylation of progesterone receptor A and B forms revealed by a phosphoserine site-specific monoclonal antibody. *Mol Endocrinol*. 2000;14:52–65.
37. Daniel AR, Qiu M, Faivre EJ, Ostrander JH, Skildum A, Lange CA. Linkage of progestin and epidermal growth factor signaling: phosphorylation of progesterone receptors mediates transcriptional hypersensitivity and increased ligand-independent breast cancer cell growth. *Steroids*. 2007;72:188–201.
38. Fuqua SA, Wiltschke C, Zhang QX, Borg A, Castles CG, Friedrichs WE, et al. A hypersensitive estrogen receptor-alpha mutation in premalignant breast lesions. *Cancer Res*. 2000;60:4026–9.
39. Herynk MH, Parra I, Cui Y, Beyer A, Wu MF, Hilsenbeck SG, et al. Association between the estrogen receptor alpha A908G mutation and outcomes in invasive breast cancer. *Clin Cancer Res*. 2007;13:3235–43.
40. Conway K, Parrish E, Edmiston SN, Tolbert D, Tse CK, Geradts J, et al. The estrogen receptor-alpha A908G (K303R) mutation occurs at a low frequency in invasive breast tumors: results from a population-based study. *Breast Cancer Res*. 2005;7:R871–80.
41. Conway K, Parrish E, Edmiston SN, Tolbert D, Tse CK, Moorman P, et al. Risk factors for breast cancer characterized by the estrogen receptor alpha A908G (K303R) mutation. *Breast Cancer Res*. 2007;9:R36.
42. Pietras RJ, Szego CM. Specific binding sites for oestrogen at the outer surfaces of isolated endometrial cells. *Nature*. 1977;265:69–72.
43. Moriarty K, Kim KH, Bender JR. Minireview: estrogen receptor-mediated rapid signaling. *Endocrinology*. 2006;147:5557–63.
44. Wehling M, Losel R. Non-genomic steroid hormone effects: membrane or intracellular receptors? *J Steroid Biochem Mol Biol*. 2006;102:180–3.
45. Li L, Haynes MP, Bender JR. Plasma membrane localization and function of the estrogen receptor alpha variant (ER46) in human endothelial cells. *Proc Natl Acad Sci U S A*. 2003;100:4807–12.



46. Wang Z, Zhang X, Shen P, Loggie BW, Chang Y, Deuel TF. A variant of estrogen receptor- $\alpha$ , hER- $\alpha$ 36: transduction of estrogen- and antiestrogen-dependent membrane-initiated mitogenic signaling. *Proc Natl Acad Sci U S A*. 2006;103:9063–8.
47. Levin ER, Pietras RJ. Estrogen receptors outside the nucleus in breast cancer. *Breast Cancer Res Treat*. 2008;108:351–61.
48. Filardo EJ. Epidermal growth factor receptor (EGFR) transactivation by estrogen via the G-protein-coupled receptor, GPR30: a novel signaling pathway with potential significance for breast cancer. *J Steroid Biochem Mol Biol*. 2002;80:231–8.
49. Filardo EJ, Quinn JA, Sabo E. Association of the membrane estrogen receptor, GPR30, with breast tumor metastasis and transactivation of the epidermal growth factor receptor. *Steroids*. 2008;73:870–3.
50. Song RX, Zhang Z, Chen Y, Bao Y, Santen RJ. Estrogen signaling via a linear pathway involving insulin-like growth factor I receptor, matrix metalloproteinases, and epidermal growth factor receptor to activate mitogen-activated protein kinase in MCF-7 breast cancer cells. *Endocrinology*. 2007;148:4091–101.
51. Song RX, McPherson RA, Adam L, Bao Y, Shupnik M, Kumar R, et al. Linkage of rapid estrogen action to MAPK activation by ER $\alpha$ -Shc association and Shc pathway activation. *Mol Endocrinol*. 2002;16:116–27.
52. Losel RM, Falkenstein E, Feuring M, Schultz A, Tillmann HC, Rossol-Haseroth K, et al. Nongenomic steroid action: controversies, questions, and answers. *Physiol Rev*. 2003;83:965–1016.
53. Chen JQ, Yager JD, Russo J. Regulation of mitochondrial respiratory chain structure and function by estrogens/estrogen receptors and potential physiological/pathophysiological implications. *Biochim Biophys Acta*. 1746;2005:1–17.
54. Pedram A, Razandi M, Wallace DC, Levin ER. Functional estrogen receptors in the mitochondria of breast cancer cells. *Mol Biol Cell*. 2006;17:2125–37.
55. Boonyaratanakornkit V, McGowan E, Sherman L, Mancini MA, Cheskis BJ, Edwards DP. The role of extranuclear signaling actions of progesterone receptor in mediating progesterone regulation of gene expression and the cell cycle. *Mol Endocrinol*. 2007;21:359–75.
56. Shou J, Massarweh S, Osborne CK, Wakeling AE, Ali S, Weiss H, et al. Mechanisms of tamoxifen resistance: increased estrogen receptor-HER2/neu cross-talk in ER/HER2-positive breast cancer. *J Natl Cancer Inst*. 2004;96:926–35.
57. Stoica GE, Franke TF, Moroni M, Mueller S, Morgan E, Iann MC, et al. Effect of estradiol on estrogen receptor- $\alpha$  gene expression and activity can be modulated by the ErbB2/PI 3-K/Akt pathway. *Oncogene*. 2003;22:7998–8011.
58. Schiff R, Massarweh SA, Shou J, Bharwani L, Mohsin SK, Osborne CK. Cross-talk between estrogen receptor and growth factor pathways as a molecular target for overcoming endocrine resistance. *Clin Cancer Res*. 2004;10:331S–6.
59. Anderson WF, Chatterjee N, Ershler WB, Brawley OW. Estrogen receptor breast cancer phenotypes in the surveillance, epidemiology, and end results database. *Breast Cancer Res Treat*. 2002;76:27–36.
60. Early Breast Cancer Trialists' Collaborative Group. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet*. 2005;365:1687–717.
61. Nadji M, Gomez-Fernandez C, Ganjei-Azar P, Morales AR. Immunohistochemistry of estrogen and progesterone receptors reconsidered: experience with 5,993 breast cancers. *Am J Clin Pathol*. 2005;123:21–7.
62. Bardou VJ, Arpino G, Elledge RM, Osborne CK, Clark GM. Progesterone receptor status significantly improves outcome prediction over estrogen receptor status alone for adjuvant endocrine therapy in two large breast cancer databases. *J Clin Oncol*. 2003;21:1973–9.
63. Arisio R, Sapino A, Cassoni P, Accinelli G, Cuccorese MC, Mano MP, et al. What modifies the relation between tumour size and lymph node metastases in T1 breast carcinomas? *J Clin Pathol*. 2000;53:846–50.
64. Manni A, Arafah B, Pearson OH. Estrogen and progesterone receptors in the prediction of response of breast cancer to endocrine therapy. *Cancer*. 1980;46:2838–41.
65. Bezwoda WR, Esser JD, Dansey R, Kessel I, Lange M. The value of estrogen and progesterone receptor determinations in advanced breast cancer. Estrogen receptor level but not progesterone receptor level correlates with response to tamoxifen. *Cancer*. 1991;68:867–72.
66. McClelland RA, Berger U, Miller LS, Powles TJ, Coombes RC. Immunocytochemical assay for estrogen receptor in patients with breast cancer: relationship to a biochemical assay and to outcome of therapy. *J Clin Oncol*. 1986;4:1171–6.
67. Mouridsen H, Gershanovich M, Sun Y, Perez-Carrion R, Boni C, Monnier A, et al. Phase III study of letrozole versus tamoxifen as first-line therapy of advanced breast cancer in postmenopausal women: analysis of survival and update of efficacy from the International Letrozole Breast Cancer Group. *J Clin Oncol*. 2003;21:2101–9.
68. Dodwell D, Wardley A, Johnston S. Postmenopausal advanced breast cancer: options for therapy after tamoxifen and aromatase inhibitors. *Breast*. 2006;15:584–94.
69. Buzdar A, Jonat W, Howell A, Jones SE, Blomqvist C, Vogel CL, et al. Anastrozole, a potent and selective aromatase inhibitor, versus megestrol acetate in postmenopausal women with advanced breast

- cancer: results of overview analysis of two phase III trials. Arimidex Study Group. *J Clin Oncol.* 1996;14:2000–11.
70. Lonning PE, Taylor PD, Anker G, Iddon J, Wie L, Jørgensen LM, et al. High-dose estrogen treatment in postmenopausal breast cancer patients heavily exposed to endocrine therapy. *Breast Cancer Res Treat.* 2001;67:111–6.
  71. Buzdar AU, Vergote I, Sainsbury R. The impact of hormone receptor status on the clinical efficacy of the new-generation aromatase inhibitors: a review of data from first-line metastatic disease trials in postmenopausal women. *Breast J.* 2004;10:211–7.
  72. Elledge RM, Green S, Pugh R, Allred DC, Clark GM, Hill J, et al. Estrogen receptor (ER) and progesterone receptor (PgR), by ligand-binding assay compared with ER, PgR and pS2, by immuno-histochemistry in predicting response to tamoxifen in metastatic breast cancer: a Southwest Oncology Group Study. *Int J Cancer.* 2000;89:111–7.
  73. Ravdin PM, Green S, Dorr TM, McGuire WL, Fabian C, Pugh RP, et al. Prognostic significance of progesterone receptor levels in estrogen receptor-positive patients with metastatic breast cancer treated with tamoxifen: results of a prospective Southwest Oncology Group study. *J Clin Oncol.* 1992;10:1284–91.
  74. Pertschuk LP, Feldman JG, Eisenberg KB, Carter AC, Thelmo WL, Cruz WP, et al. Immunocytochemical detection of progesterone receptor in breast cancer with monoclonal antibody. Relation to biochemical assay, disease-free survival, and clinical endocrine response. *Cancer.* 1988;62:342–9.
  75. Spataro V, Price K, Goldhirsch A, Cavalli F, Simoncini E, Castiglione M, et al. Sequential estrogen receptor determinations from primary breast cancer and at relapse: prognostic and therapeutic relevance. The International Breast Cancer Study Group (formerly Ludwig Group). *Ann Oncol.* 1992;3:733–40.
  76. Kuukasjarvi T, Kononen J, Helin H, Holli K, Isola J. Loss of estrogen receptor in recurrent breast cancer is associated with poor response to endocrine therapy. *J Clin Oncol.* 1996;14:2584–9.
  77. Lower EE, Glass EL, Bradley DA, Blau R, Heffelfinger S. Impact of metastatic estrogen receptor and progesterone receptor status on survival. *Breast Cancer Res Treat.* 2005;90:65–70.
  78. Sari E, Guler G, Hayran M, Gullu I, Altundag K, Ozisik Y. Comparative study of the immunohistochemical detection of hormone receptor status and HER-2 expression in primary and paired recurrent/metastatic lesions of patients with breast cancer. *Med Oncol.* 2011;28:57–63.
  79. Arslan C, Sari E, Aksoy S, Altundag K. Variation in hormone receptor and HER-2 status between primary and metastatic breast cancer: review of the literature. *Expert Opin Ther Targets.* 2011;15:21–30.
  80. Gutierrez MC, Detre S, Johnston S, Mohsin SK, Shou J, Allred DC, et al. Molecular changes in tamoxifen-resistant breast cancer: relationship between estrogen receptor, HER-2, and p38 mitogen-activated protein kinase. *J Clin Oncol.* 2005;23:2469–76.
  81. Gross GE, Clark GM, Chamness GC, McGuire WL. Multiple progesterone receptor assays in human breast cancer. *Cancer Res.* 1984;44:836–40.
  82. Bieche I, Lidereau R. Genetic alterations in breast cancer. *Genes Chromosomes Cancer.* 1995;14:227–51.
  83. Teixeira MR, Pandis N, Bardi G, Andersen JA, Mitelman F, Heim S. Clonal heterogeneity in breast cancer: karyotypic comparisons of multiple intra- and extra-tumorous samples from 3 patients. *Int J Cancer.* 1995;63:63–8.
  84. Hutchins LF, Green SJ, Ravdin PM, Lew D, Martino S, Abeloff M, et al. Randomized, controlled trial of cyclophosphamide, methotrexate, and fluorouracil versus cyclophosphamide, doxorubicin, and fluorouracil with and without tamoxifen for high-risk, node-negative breast cancer: treatment results of Intergroup Protocol INT-0102. *J Clin Oncol.* 2005;23:8313–21.
  85. Fisher B, Anderson S, Tan-Chiu E, Tan-Chiu E, Wolmark N, Wickerham DL, et al. Tamoxifen and chemotherapy for axillary node-negative, estrogen receptor-negative breast cancer: findings from National Surgical Adjuvant Breast and Bowel Project B-23. *J Clin Oncol.* 2001;19:931–42.
  86. Dowsett M, Allred C, Knox J, Quinn E, Salter J, Wale C, et al. Relationship between quantitative estrogen and progesterone receptor expression and human epidermal growth factor receptor 2 (HER-2) status with recurrence in the Arimidex, Tamoxifen, Alone or in Combination trial. *J Clin Oncol.* 2008;26:1059–65.
  87. Viale G, Regan MM, Maiorano E, Mastropasqua MG, Dell'Orto P, Rasmussen BB, et al. Prognostic and predictive value of centrally reviewed expression of estrogen and progesterone receptors in a randomized trial comparing letrozole and tamoxifen adjuvant therapy for postmenopausal early breast cancer: BIG 1–98. *J Clin Oncol.* 2007;25:3846–52.
  88. Early Breast Cancer Trialists' Collaborative Group. Tamoxifen for early breast cancer: an overview of the randomised trials. *Lancet.* 1998;351:1451–67.
  89. De Maeyer L, Van Limbergen E, De Nys K, Moerman P, Pochet N, Hendrickx W, et al. Does estrogen receptor negative/progesterone receptor positive breast carcinoma exist? *J Clin Oncol.* 2008;26:335–6; author reply 6–8.
  90. Rakha EA, El-Sayed ME, Green AR, Paish EC, Powe DG, Gee J, et al. Biologic and clinical characteristics of breast cancer with single hormone receptor positive phenotype. *J Clin Oncol.* 2007;25:4772–8.
  91. Dunnwald LK, Rossing MA, Li CI. Hormone receptor status, tumor characteristics, and prognosis: a

- prospective cohort of breast cancer patients. *Breast Cancer Res.* 2007;9:R6.
92. Dowsett M, Houghton J, Iden C, Salter J, Farnon J, A'Hern R, et al. Benefit from adjuvant tamoxifen therapy in primary breast cancer patients according oestrogen receptor, progesterone receptor, EGF receptor and HER2 status. *Ann Oncol.* 2006;17:818–26.
  93. Ferno M, Stal O, Baldetorp B, Hatschek T, Källström AC, Malmström P, et al. Results of two or five years of adjuvant tamoxifen correlated to steroid receptor and S-phase levels. South Sweden Breast Cancer Group, and South-East Sweden Breast Cancer Group. *Breast Cancer Res Treat.* 2000;59:69–76.
  94. Lamy PJ, Pujol P, Thezenas S, Kramar A, Rouanet P, Guilleux F, et al. Progesterone receptor quantification as a strong prognostic determinant in postmenopausal breast cancer women under tamoxifen therapy. *Breast Cancer Res Treat.* 2002;76:65–71.
  95. Goss PE, Ingle JN, Martino S, Robert NJ, Muss HB, Piccart MJ, et al. Efficacy of letrozole extended adjuvant therapy according to estrogen receptor and progesterone receptor status of the primary tumor: National Cancer Institute of Canada Clinical Trials Group MA.17. *J Clin Oncol.* 2007;25:2006–11.
  96. Sparano JA, Paik S. Development of the 21-gene assay and its application in clinical practice and clinical trials. *J Clin Oncol.* 2008;26:721–8.
  97. Andre F, Broglio K, Roche H, Martin M, Mackey JR, Penault-Llorca F, et al. Estrogen receptor expression and efficacy of docetaxel-containing adjuvant chemotherapy in patients with node-positive breast cancer: results from a pooled analysis. *J Clin Oncol.* 2008;26:2636–43.
  98. Martin M, Pienkowski T, Mackey J, Pawlicki M, Guastalla JP, Weaver C, et al. Adjuvant docetaxel for node-positive breast cancer. *N Engl J Med.* 2005;352:2302–13.
  99. De Laurentis M, Cancellato G, D'Agostino D, Giuliano M, Giordano A, Montagna E, et al. Taxane-based combinations as adjuvant chemotherapy of early breast cancer: a meta-analysis of randomized trials. *J Clin Oncol.* 2008;26:44–53.
  100. Henry NL, Hayes DF. Can biology trump anatomy? Do all node-positive patients with breast cancer need chemotherapy? *J Clin Oncol.* 2007;25:2501–3.
  101. Berry DA, Cirincione C, Henderson IC, Citron ML, Budman DR, Goldstein LJ, et al. Estrogen-receptor status and outcomes of modern chemotherapy for patients with node-positive breast cancer. *JAMA.* 2006;295:1658–67.
  102. Hayes DF, Thor AD, Dressler LG, Weaver D, Edgerton S, Cowan D, et al. HER2 and response to paclitaxel in node-positive breast cancer. *N Engl J Med.* 2007;357:1496–506.
  103. Mazouni C, Kau SW, Frye D, Andre F, Kuerer HM, Buchholz TA, et al. Inclusion of taxanes, particularly weekly paclitaxel, in preoperative chemotherapy improves pathologic complete response rate in estrogen receptor-positive breast cancers. *Ann Oncol.* 2007;18:874–80.
  104. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med.* 2004;351:2817–26.
  105. Paik S, Tang G, Shak S, Kim C, Baker J, Kim W, et al. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol.* 2006;24:3726–34.
  106. Albain KS, Barlow WE, Shak S, Hortobagyi GN, Livingston RB, Yeh IT, et al. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol.* 2010;11:55–65.
  107. Crowe JP, Hubay CA, Pearson OH, Marshall JS, Rosenblatt J, Mansour EG, et al. Estrogen receptor status as a prognostic indicator for stage I breast cancer patients. *Breast Cancer Res Treat.* 1982;2:171–6.
  108. Fisher B, Redmond C, Fisher ER, Caplan R. Relative worth of estrogen or progesterone receptor and pathologic characteristics of differentiation as indicators of prognosis in node negative breast cancer patients: findings from National Surgical Adjuvant Breast and Bowel Project Protocol B-06. *J Clin Oncol.* 1988;6:1076–87.
  109. Costa SD, Lange S, Klinga K, Merkle E, Kaufmann M. Factors influencing the prognostic role of oestrogen and progesterone receptor levels in breast cancer – results of the analysis of 670 patients with 11 years of follow-up. *Eur J Cancer.* 2002;38:1329–34.
  110. Hilsenbeck SG, Ravdin PM, de Moor CA, Chamness GC, Osborne CK, Clark GM. Time-dependence of hazard ratios for prognostic factors in primary breast cancer. *Breast Cancer Res Treat.* 1998;52:227–37.
  111. Schmitt M, Thomssen C, Ulm K, Seiderer A, Harbeck N, Höfler H, et al. Time-varying prognostic impact of tumour biological factors urokinase (uPA), PAI-1 and steroid hormone receptor status in primary breast cancer. *Br J Cancer.* 1997;76:306–11.
  112. Diab SG, Elledge RM, Clark GM. Tumor characteristics and clinical outcome of elderly women with breast cancer. *J Natl Cancer Inst.* 2000;92:550–6.
  113. Fisher ER, Osborne CK, McGuire WL, Redmond C, Knight 3rd WA, Fisher B, et al. Correlation of primary breast cancer histopathology and estrogen receptor content. *Breast Cancer Res Treat.* 1981;1:37–41.
  114. Wenger CR, Beardslee S, Owens MA, Pounds G, Oldaker T, Vendely P, et al. DNA ploidy, S-phase, and steroid receptors in more than 127,000 breast cancer patients. *Breast Cancer Res Treat.* 1993;28:9–20.
  115. Elledge RM, Fuqua SA, Clark GM, Pujol P, Allred DC, McGuire WL. Prognostic significance of p53 gene alterations in node-negative breast cancer. *Breast Cancer Res Treat.* 1993;26:225–35.
  116. Hess KR, Pusztai L, Buzdar AU, Hortobagyi GN. Estrogen receptors and distinct patterns of

- breast cancer relapse. *Breast Cancer Res Treat.* 2003;78:105–18.
117. Koenders PG, Beex LV, Langens R, Kloppenborg PW, Smals AG, Benraad TJ. Steroid hormone receptor activity of primary human breast cancer and pattern of first metastasis. The Breast Cancer Study Group. *Breast Cancer Res Treat.* 1991;18:27–32.
  118. Nofech-Mozes S, Vella ET, Dhesy-Thind S, Hagerty KL, Mangu PB, Temin S, et al. Systematic review on hormone receptor testing in breast cancer. *Appl Immunohistochem Mol Morphol.* 2012;20:214–63.
  119. Alberts SR, Ingle JN, Roche PR, Cha SS, Wold LE, Farr Jr GH, et al. Comparison of estrogen receptor determinations by a biochemical ligand-binding assay and immunohistochemical staining with monoclonal antibody ER1D5 in females with lymph node positive breast carcinoma entered on two prospective clinical trials. *Cancer.* 1996;78:764–72.
  120. Harvey JM, Clark GM, Osborne CK, Allred DC. Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *J Clin Oncol.* 1999;17:1474–81.
  121. Thomson CS, Twelves CJ, Mallon EA, Leake RE. Scottish Cancer Trials Breast G, Scottish Cancer Therapy N. Adjuvant ovarian ablation vs CMF chemotherapy in premenopausal breast cancer patients: trial update and impact of immunohistochemical assessment of ER status. *Breast.* 2002;11:419–29.
  122. Regan MM, Viale G, Mastropasqua MG, Maiorano E, Golouh R, Carbone A, et al. Re-evaluating adjuvant breast cancer trials: assessing hormone receptor status by immunohistochemical versus extraction assays. *J Natl Cancer Inst.* 2006;98:1571–81.
  123. Glas AM, Floore A, Delahaye LJ, Witteveen AT, Pover RC, Bakx N, et al. Converting a breast cancer microarray signature into a high-throughput diagnostic test. *BMC Genomics.* 2006;7:278.
  124. Hammond ME, Hayes DF, Dowsett M, Allred DC, Hagerty KL, Badve S, et al. American Society of Clinical Oncology/College Of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. *J Clin Oncol.* 2010;28:2784–95.
  125. Gown AM. Unmasking the mysteries of antigen or epitope retrieval and formalin fixation. *Am J Clin Pathol.* 2004;121:172–4.
  126. Diaz LK, Sneige N. Estrogen receptor analysis for breast cancer: current issues and keys to increasing testing accuracy. *Adv Anat Pathol.* 2005;12:10–9.
  127. Goldstein NS, Ferkowicz M, Odish E, Mani A, Hastah F. Minimum formalin fixation time for consistent estrogen receptor immunohistochemical staining of invasive breast carcinoma. *Am J Clin Pathol.* 2003;120:86–92.
  128. Taylor CR, Levenson RM. Quantification of immunohistochemistry – issues concerning methods, utility and semiquantitative assessment II. *Histopathology.* 2006;49:411–24.
  129. Arber DA. Effect of prolonged formalin fixation on the immunohistochemical reactivity of breast markers. *Appl Immunohistochem Mol Morphol.* 2002;10:183–6.
  130. Oh JJ, Grosshans DR, Wong SG, Slamon DJ. Identification of differentially expressed genes associated with HER-2/neu overexpression in human breast cancer cells. *Nucleic Acids Res.* 1999;27:4008–17.
  131. Kumar R, Yarmand-Bagheri R. The role of HER2 in angiogenesis. *Semin Oncol.* 2001;28:27–32.
  132. Press MF, Cordon-Cardo C, Slamon DJ. Expression of the HER-2/neu proto-oncogene in normal human adult and fetal tissues. *Oncogene.* 1990;5:953–62.
  133. Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science.* 1987;235:177–82.
  134. Slamon DJ, Godolphin W, Jones LA, Holt JA, Wong SG, Keith DE, et al. Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science.* 1989;244:707–12.
  135. Saffari B, Jones LA, el-Naggar A, Felix JC, George J, Press MF. Amplification and overexpression of HER-2/neu (c-erbB2) in endometrial cancers: correlation with overall survival. *Cancer Res.* 1995;55:5693–8.
  136. Press MF, Pike MC, Hung G, Zhou JY, Ma Y, George J, Dietz-Band J, et al. Amplification and overexpression of HER-2/neu in carcinomas of the salivary gland: correlation with poor prognosis. *Cancer Res.* 1994;54:5675–82.
  137. Park JB, Rhim JS, Park SC, Kimm SW, Kraus MH. Amplification, overexpression, and rearrangement of the erbB-2 protooncogene in primary human stomach carcinomas. *Cancer Res.* 1989;49:6605–9.
  138. Sorkin A, Goh LK. Endocytosis and intracellular trafficking of ErbBs. *Exp Cell Res.* 2008;314:3093–106.
  139. Yarden Y, Sliwkowski MX. Untangling the ErbB signalling network. *Nat Rev Mol Cell Biol.* 2001;2:127–37.
  140. Cho HS, Mason K, Ramyar KX, Stanley AM, Gabelli SB, Denney Jr DW, et al. Structure of the extracellular region of HER2 alone and in complex with the Herceptin Fab. *Nature.* 2003;421:756–60.
  141. Tzahar E, Waterman H, Chen X, Levkowitz G, Karunagaran D, Lavi S, et al. A hierarchical network of interreceptor interactions determines signal transduction by Neu differentiation factor/neuregulin and epidermal growth factor. *Mol Cell Biol.* 1996;16:5276–87.
  142. Pinkas-Kramarski R, Soussan L, Waterman H, Levkowitz G, Alroy I, Klapper L, et al. Diversification of Neu differentiation factor and epidermal growth factor signaling by combinatorial receptor interactions. *EMBO J.* 1996;15:2452–67.
  143. Giri DK, Ali-Seyed M, Li LY, Lee DF, Ling P, Bartholomeusz G, et al. Endosomal transport of ErbB-2: mechanism for nuclear entry of the cell surface receptor. *Mol Cell Biol.* 2005;25:11005–18.

144. Ni CY, Murphy MP, Golde TE, Carpenter G. gamma-Secretase cleavage and nuclear localization of ErbB-4 receptor tyrosine kinase. *Science*. 2001;294:2179–81.
145. Wang SC, Hung MC. Nuclear translocation of the epidermal growth factor receptor family membrane tyrosine kinase receptors. *Clin Cancer Res*. 2009;15:6484–9.
146. Wolff AC, Hammond ME, Hicks DG, Dowsett M, McShane LM, Allison KH, et al. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *J Clin Oncol*. 2013;31:3997–4013.
147. Press MF, Bernstein L, Thomas PA, Meisner LF, Zhou JY, Ma Y, et al. HER-2/neu gene amplification characterized by fluorescence in situ hybridization: poor prognosis in node-negative breast carcinomas. *J Clin Oncol*. 1997;15:2894–904.
148. Mass RD, Press MF, Anderson S, Cobleigh MA, Vogel CL, Dybdal N, et al. Evaluation of clinical outcomes according to HER2 detection by fluorescence in situ hybridization in women with metastatic breast cancer treated with trastuzumab. *Clin Breast Cancer*. 2005;6:240–6.
149. Seidman AD, Berry D, Cirincione C, Harris L, Muss H, Marcom PK, et al. Randomized phase III trial of weekly compared with every-3-weeks paclitaxel for metastatic breast cancer, with trastuzumab for all HER-2 overexpressors and random assignment to trastuzumab or not in HER-2 nonoverexpressors: final results of Cancer and Leukemia Group B protocol 9840. *J Clin Oncol*. 2008;26:1642–9.
150. Slamon DJ, Leyland-Jones B, Shak S, Shak S, Fuchs H, Paton V, Bajamonde A, et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med*. 2001;344:783–92.
151. Slamon D, Eiermann W, Robert N, Pienkowski T, Martin M, Press M, et al. Adjuvant trastuzumab in HER2-positive breast cancer. *N Engl J Med*. 2011;365:1273–83.
152. Romond EH, Perez EA, Bryant J, Suman VJ, Geyer Jr CE, Davidson NE, et al. Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med*. 2005;353:1673–84.
153. Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, et al. Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med*. 2005;353:1659–72.
154. Gianni L, Dafni U, Gelber RD, Azambuja E, Muehlbauer S, Goldhirsch A, et al. Treatment with trastuzumab for 1 year after adjuvant chemotherapy in patients with HER2-positive early breast cancer: a 4-year follow-up of a randomised controlled trial. *Lancet Oncol*. 2011;12:236–44.
155. Perez EA, Romond EH, Suman VJ, Jeong JH, Davidson NE, Geyer Jr CE, et al. Four-year follow-up of trastuzumab plus adjuvant chemotherapy for operable human epidermal growth factor receptor 2-positive breast cancer: joint analysis of data from NCCTG N9831 and NSABP B-31. *J Clin Oncol*. 2011;29:3366–73.
156. Geyer CE, Forster J, Lindquist D, Chan S, Romieu CG, Pienkowski T, et al. Lapatinib plus capecitabine for HER2-positive advanced breast cancer. *N Engl J Med*. 2006;355:2733–43.
157. Baselga J, Cortes J, Kim SB, Im SA, Hegg R, Im YH, Roman L, et al. Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer. *N Engl J Med*. 2012;366:109–19.
158. Verma S, Miles D, Gianni L, Krop IE, Welslau M, Baselga J, et al. Trastuzumab emtansine for HER2-positive advanced breast cancer. *N Engl J Med*. 2012;367:1783–91.
159. Dati C, Antoniotti S, Taverna D, Perroteau I, De Bortoli M. Inhibition of c-erbB-2 oncogene expression by estrogens in human breast cancer cells. *Oncogene*. 1990;5:1001–6.
160. Benz CC, Scott GK, Sarup JC, Johnson RM, Tripathy D, Coronado E, et al. Estrogen-dependent, tamoxifen-resistant tumorigenic growth of MCF-7 cells transfected with HER2/neu. *Breast Cancer Res Treat*. 1992;24:85–95.
161. Pietras RJ, Arboleda J, Reese DM, Wongvipat N, Pegram MD, Ramos L, et al. HER-2 tyrosine kinase pathway targets estrogen receptor and promotes hormone-independent growth in human breast cancer cells. *Oncogene*. 1995;10:2435–46.
162. Mueller H, Kueng W, Schoumacher F, Herzer S, Eppenberger U. Selective regulation of steroid receptor expression in MCF-7 breast cancer cells by a novel member of the heregulin family. *Biochem Biophys Res Commun*. 1995;217:1271–8.
163. Borg A, Baldetorp B, Ferno M, Killander D, Olsson H, Rydén S, et al. ERBB2 amplification is associated with tamoxifen resistance in steroid-receptor positive breast cancer. *Cancer Lett*. 1994;81:137–44.
164. Carlomagno C, Perrone F, Gallo C, De Laurentiis M, Lauria R, Morabito A, et al. c-erb B2 overexpression decreases the benefit of adjuvant tamoxifen in early-stage breast cancer without axillary lymph node metastases. *J Clin Oncol*. 1996;14:2702–8.
165. De Placido S, De Laurentiis M, Carlomagno C, Gallo C, Perrone F, Pepe S, et al. Twenty-year results of the Naples GUN randomized trial: predictive factors of adjuvant tamoxifen efficacy in early breast cancer. *Clin Cancer Res*. 2003;9:1039–46.
166. De Laurentiis M, Arpino G, Massarelli E, Ruggiero A, Carlomagno C, Ciardiello F, et al. A meta-analysis on the interaction between HER-2 expression and response to endocrine treatment in advanced breast cancer. *Clin Cancer Res*. 2005;11:4741–8.
167. Knoop AS, Bentzen SM, Nielsen MM, Rasmussen BB, Rose C. Value of epidermal growth factor receptor, HER2, p53, and steroid receptors in predicting the efficacy of tamoxifen in high-risk



- postmenopausal breast cancer patients. *J Clin Oncol.* 2001;19:3376–84.
168. Berry DA, Muss HB, Thor AD, Dressler L, Liu ET, Broadwater G, et al. HER-2/neu and p53 expression versus tamoxifen resistance in estrogen receptor-positive, node-positive breast cancer. *J Clin Oncol.* 2000;18:3471–9.
  169. Stal O, Borg A, Ferno M, Källström AC, Malmström P, Nordenskjöld B, et al. ErbB2 status and the benefit from two or five years of adjuvant tamoxifen in postmenopausal early stage breast cancer. *Ann Oncol.* 2000;11:1545–50.
  170. Love RR, Duc NB, Havighurst TC, Mohsin SK, Zhang Q, DeMets DL, et al. Her-2/neu overexpression and response to oophorectomy plus tamoxifen adjuvant therapy in estrogen receptor-positive premenopausal women with operable breast cancer. *J Clin Oncol.* 2003;21:453–7.
  171. Ellis MJ, Coop A, Singh B, Mauriac L, Llombert-Cussac A, Jänicke F, et al. Letrozole is more effective neoadjuvant endocrine therapy than tamoxifen for ErbB-1- and/or ErbB-2-positive, estrogen receptor-positive primary breast cancer: evidence from a phase III randomized trial. *J Clin Oncol.* 2001;19:3808–16.
  172. Wright C, Nicholson S, Angus B, Sainsbury JR, Farndon J, Cairns J, et al. Relationship between c-erbB-2 protein product expression and response to endocrine therapy in advanced breast cancer. *Br J Cancer.* 1992;65:118–21.
  173. Leitzel K, Teramoto Y, Konrad K, Chinchilli VM, Volas G, Grossberg H, et al. Elevated serum c-erbB-2 antigen levels and decreased response to hormone therapy of breast cancer. *J Clin Oncol.* 1995;13:1129–35.
  174. Yamauchi H, O'Neill A, Gelman R, Carney W, Tenney DY, Hösch S, et al. Prediction of response to antiestrogen therapy in advanced breast cancer patients by pretreatment circulating levels of extracellular domain of the HER-2/c-neu protein. *J Clin Oncol.* 1997;15:2518–25.
  175. Lipton A, Ali SM, Leitzel K, Demers L, Harvey HA, Chaudri-Ross HA, et al. Serum HER-2/neu and response to the aromatase inhibitor letrozole versus tamoxifen. *J Clin Oncol.* 2003;21:1967–72.
  176. Elledge RM, Green S, Ciocca D, Pugh R, Allred DC, Clark GM, et al. HER-2 expression and response to tamoxifen in estrogen receptor-positive breast cancer: a Southwest Oncology Group Study. *Clin Cancer Res.* 1998;4:7–12.
  177. Harris L, Fritsche H, Mennel R, Norton L, Ravdin P, Taube S, et al. American Society of Clinical Oncology 2007 update of recommendations for the use of tumor markers in breast cancer. *J Clin Oncol.* 2007;25:5287–312.
  178. Pritchard KI, Levine MN, Tu D. neu/erbB-2 overexpression and response to hormonal therapy in premenopausal women in the adjuvant breast cancer setting: will it play in Peoria? part II. *J Clin Oncol.* 2003;21:399–400.
  179. Paik S, Bryant J, Park C, Fisher B, Tan-Chiu E, Hyams D, et al. erbB-2 and response to doxorubicin in patients with axillary lymph node-positive, hormone receptor-negative breast cancer. *J Natl Cancer Inst.* 1998;90:1361–70.
  180. Paik S, Bryant J, Tan-Chiu E, Yothers G, Park C, Wickerham DL, et al. HER2 and choice of adjuvant chemotherapy for invasive breast cancer: National Surgical Adjuvant Breast and Bowel Project Protocol B-15. *J Natl Cancer Inst.* 2000;92:1991–8.
  181. Del Mastro L, Bruzzi P, Nicolo G, Cavazzini G, Contu A, D'Amico M, et al. HER2 expression and efficacy of dose-dense anthracycline-containing adjuvant chemotherapy in breast cancer patients. *Br J Cancer.* 2005;93:7–14.
  182. Gennari A, Sormani MP, Pronzato P, Puntoni M, Colozza M, Pfeffer U, et al. HER2 status and efficacy of adjuvant anthracyclines in early breast cancer: a pooled analysis of randomized trials. *J Natl Cancer Inst.* 2008;100:14–20.
  183. Villman K, Sjöstrom J, Heikkilä R, Hultborn R, Malmström P, Bengtsson NO, et al. TOP2A and HER2 gene amplification as predictors of response to anthracycline treatment in breast cancer. *Acta Oncol.* 2006;45:590–6.
  184. Press MF, Sauter G, Buysse M, Bernstein L, Guzman R, Santiago A, et al. Alteration of topoisomerase II-alpha gene in human breast cancer: association with responsiveness to anthracycline-based chemotherapy. *J Clin Oncol.* 2011;29:859–67.
  185. Di Leo A, Desmedt C, Bartlett JM, Piette F, Ejlertsen B, Pritchard KI, et al. HER2 and TOP2A as predictive markers for anthracycline-containing chemotherapy regimens as adjuvant treatment of breast cancer: a meta-analysis of individual patient data. *Lancet Oncol.* 2011;12:1134–42.
  186. Hayes DF. Steady progress against HER2-positive breast cancer. *N Engl J Med.* 2011;365:1336–8.
  187. Konecny GE, Thomssen C, Luck HJ, Untch M, Wang HJ, Kuhn W, et al. Her-2/neu gene amplification and response to paclitaxel in patients with metastatic breast cancer. *J Natl Cancer Inst.* 2004;96:1141–51.
  188. Martin M, Rodriguez-Lescure A, Ruiz A, Alba E, Calvo L, Ruiz-Borrego M, et al. Randomized phase 3 trial of fluorouracil, epirubicin, and cyclophosphamide alone or followed by Paclitaxel for early breast cancer. *J Natl Cancer Inst.* 2008;100:805–14.
  189. Sparano JA, Wang M, Martino S, Jones V, Perez EA, Saphner T, et al. Weekly paclitaxel in the adjuvant treatment of breast cancer. *N Engl J Med.* 2008;358:1663–71.
  190. Learn PA, Yeh IT, McNutt M, Chisholm GB, Pollock BH, Rousseau Jr DL, et al. HER-2/neu expression as a predictor of response to neoadjuvant docetaxel in patients with operable breast carcinoma. *Cancer.* 2005;103:2252–60.

191. Gonzalez-Angulo AM, Krishnamurthy S, Yamamura Y, Broglio KR, Puzstai L, Buzdar AU, et al. Lack of association between amplification of her-2 and response to preoperative taxanes in patients with breast carcinoma. *Cancer*. 2004;101:258–63.
192. Pu RT, Schott AF, Sturtz DE, Griffith KA, Kleer CG. Pathologic features of breast cancer associated with complete response to neoadjuvant chemotherapy: importance of tumor necrosis. *Am J Surg Pathol*. 2005;29:354–8.
193. Cobleigh MA, Vogel CL, Tripathy D, Robert NJ, Scholl S, Fehrenbacher L, et al. Multinational study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who have HER2-overexpressing metastatic breast cancer that has progressed after chemotherapy for metastatic disease. *J Clin Oncol*. 1999;17:2639–48.
194. Wolff AC, Hammond ME, Schwartz JN, Hagerty KL, Allred DC, Cote RJ, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol*. 2007;25:118–45.
195. Perez EA, Cortes J, Gonzalez-Angulo AM, Bartlett JM. HER2 testing: current status and future directions. *Cancer Treat Rev*. 2014;40:276–84.
196. Press MF, Slamon DJ, Flom KJ, Park J, Zhou JY, Bernstein L. Evaluation of HER-2/neu gene amplification and overexpression: comparison of frequently used assay methods in a molecularly characterized cohort of breast cancer specimens. *J Clin Oncol*. 2002;20:3095–105.
197. Penault-Llorca F, Bilous M, Dowsett M, Hanna W, Osamura RY, Rüschoff J, et al. Emerging technologies for assessing HER2 amplification. *Am J Clin Pathol*. 2009;132:539–48.
198. Kosa C, Kardos L, Kovacs J, Szollosi Z. Comparison of dual-color dual-hapten brightfield in situ hybridization (DDISH) and fluorescence in situ hybridization in breast cancer HER2 assessment. *Pathol Res Pract*. 2013;209:147–50.

Yun Wu and Aysegul A. Sahin

## Abstract

Invasive breast carcinomas comprise a heterogeneous group of lesions that differ in their molecular and pathologic features and clinical behavior. Some patients have long disease-free survival, whereas others experience the rapid development of recurrence and metastases that are fatal within a few years of the initial diagnosis. Many factors in individual tumors can be evaluated to stratify patients into subsets with varying risks of recurrence and response to different therapy modalities. This chapter describes the current standard prognostic and predictive factors of invasive breast carcinoma and discusses emerging data on molecular markers which can be considered in clinical practice.

## Keywords

Invasive breast cancer • Histopathology • Prognostic factors • Tumor biomarkers • Estrogen and progesterone receptors • Her2/neu

## Prognostic and Predictive Factors

To offer the optimal treatment plan for a patient with newly diagnosed invasive breast cancer, the clinician must determine the most likely clinical behavior of the particular tumor and assess the probable efficacies and potential side effects of

all available therapies. Unfortunately, one of the most significant features of invasive breast carcinoma is its biological and clinical diversity. Even tumors with similar clinical and histopathologic features may have significantly different clinical behaviors [1–3]. While some tumors have a long, protracted clinical course, others are very aggressive, with the rapid development of metastases that kill patients within a few years of the initial diagnosis.

A prognostic factor is any factor that can be evaluated at the time of diagnosis and that is associated with clinical outcome in the absence of systemic adjuvant therapy. By contrast, a predictive

---

Y. Wu, MD PhD • A.A. Sahin, MD (✉)  
Department of Pathology, The University of Texas  
MD Anderson Cancer Center, 1515 Holcombe  
Boulevard Unit 85 (G1.3569C), Houston, TX, 77030,  
USA  
e-mail: [asahin@mdanderson.org](mailto:asahin@mdanderson.org)

factor is a factor associated with a response or lack of response to a particular therapy [4]. In general, prognostic and predictive factors are used to forecast the natural history of a disease and its response to a particular therapy, respectively. In breast cancer, some factors are purely prognostic or predictive, whereas others have both prognostic and predictive value (Table 10.1) [4].

This chapter describes the standard prognostic and predictive factors of breast cancer that are currently used in clinical practice. Their use is based on the recommendations of multidisciplinary consensus panels of the National Institutes of Health, the College of American Pathologists (CAP), and the American Society of Clinical Oncology (ASCO) [5–8]. These prognostic and predictive factors include lymph node status, tumor size, lymphovascular invasion, tumor histologic type and grade, hormone receptor status, and Her2/neu gene overexpression or amplification (Table 10.1).

## Lymph Node Status

Axillary lymph node status at the time of diagnosis is the most powerful prognostic factor for patients with breast cancer [9–11]. The presence or absence of metastatic involvement and the number of lymph nodes involved in metastatic disease directly correlate with clinical outcome [11]. The prognosis becomes significantly poorer as the number of positive lymph nodes increases. Lymph node involvement is relatively independent of other prognostic factors, except tumor size [10, 12–17]. Histologic examination of lymph nodes is

the most accurate method for assessing the spread of disease to lymph nodes. Therefore, all dissected lymph nodes should be carefully identified and histologically evaluated for the presence of metastatic carcinoma. Lymph nodes with grossly evident metastatic tumor may be sampled by representative sections, but nodes that appear negative on gross examination should be sectioned in 2–3-mm slices and submitted for histologic evaluation in their entirety [16–20]. To optimize the accuracy of prognostic information, a minimum of ten lymph nodes should be evaluated from a full axillary lymph node dissection [13, 21–23].

Lymph nodes can be identified in the fresh state or in fixed tissue. A variety of clearing solutions (e.g., xylene, Carnoy's solution, Bouin's solution) have been used to facilitate the identification of lymph nodes but are, in general, not very practical [14, 16, 24–26]. At most institutions, a single hematoxylin- and eosin-stained section per block is routinely used for evaluation. Some researchers have advocated more extensive sampling of the lymph nodes by multiple sections and evaluation of immunohistochemical (IHC) stains for epithelial markers to detect micrometastases [16, 27, 28]. A more detailed evaluation may be justified in certain cases, such as invasive lobular carcinoma, but the routine use of exhaustive histopathologic evaluation and IHC assessment is not recommended [16–18, 27, 29].

In the past decade, the approach to axillary dissection in patients with clinically node-negative breast cancer has changed dramatically. Since several studies have demonstrated that axillary dissection does not contribute to survival, this procedure is used predominantly for staging and local control. Sentinel lymph node (SLN) biopsy is based on the concept that breast cancer metastasis follows an orderly and progressive pattern and that there is a reproducibly identifiable node (or nodes) that drains the breast and predicts the status of the remaining axillary nodes. If the SLN is negative for metastatic carcinoma, the remaining lymph nodes are presumed to be negative. Thus, the morbidity of axillary node dissection can be avoided in patients who are unlikely to benefit from the removal of lymph nodes [18, 30–32].

**Table 10.1** Summary of prognostic and predictive factors for invasive breast cancer

	Prognostic	Predictive
Nodal status	√√√	
Tumor size	√√√	
Lymphovascular invasion	√√	
Histological nuclear grade	√√	√√
Histologic type	√	√
Steroid receptors	√	√√√
Her2/neu overexpression amplification	√	√√√

Several studies, including randomized clinical trials, have demonstrated that the risk of a variety of complications in the arm (e.g., lymphedema, sensory loss, and shoulder abduction deficits) is significantly reduced in patients who undergo SLN biopsy compared to those who undergo standard axillary lymph node dissection. SLN biopsy has been adopted as the standard approach for the evaluation of regional lymph nodes and is considered the standard of care for women with a clinically node-negative T1 or T2 tumor [21, 31–35]. ASCO, the International Expert Consensus Panel on the Primary Therapy of Early Breast Cancer, and other national and international organizations have endorsed this approach [36–38].

Another very important aspect of the SLN biopsy procedure is that it identifies the lymph node or nodes most likely to harbor metastatic carcinoma. This procedure has significantly reduced the number of nodes sent for histopathologic evaluation, enabling more detailed pathologic evaluation of a smaller number of lymph nodes that are at the highest risk of harboring metastatic disease. Advances in molecular technologies have also facilitated the analysis of nodal tissue [39–44].

The American Joint Commission on Cancer (AJCC) staging system for breast cancer defines macrometastasis (>2.0 mm), micrometastasis (greater than 0.2–2 mm), and isolated tumor cells (equal or less than 0.2 mm) (Table 10.2a) [15]. Several retrospective studies have reported that the prognosis of patients with micrometastatic disease is intermediate between that of patients with node-negative disease and that of patients with lymph nodes showing indicating macrometastases. Our understanding of the clinical relevance and prognostic significance of small-volume metastases remains limited. The results of large cooperative studies evaluating the prognostic significance of small-volume disease are still pending [45–48].

## Tumor Size

After nodal status, tumor size is the most powerful prognostic factor related to the disease-free and overall survival of patients with breast cancer

**Table 10.2a** Regional lymph nodes (N)

	Lymph node status
NX	Regional lymph nodes cannot be assessed (e.g., previously removed)
N0	No regional lymph node metastases
N1	Metastases to movable ipsilateral level I and II axillary lymph node(s)
N2	Metastases in ipsilateral level I and II axillary lymph nodes that are clinically fixed or matted Or Metastases in clinically detected ipsilateral internal mammary nodes in the absence of clinically evident axillary lymph node metastases
N2a	Metastases in ipsilateral level I and II axillary lymph nodes fixed to one another (matted) or to other structures
N2b	Metastases only in clinically detected ipsilateral internal mammary nodes and in the absence of clinically evident level I and II axillary lymph node metastases
N3	Metastases in ipsilateral infraclavicular (level III axillary) lymph node(s) with or without level I and II axillary lymph node involvement Or Metastases in clinically detected ipsilateral internal mammary lymph node(s) with clinically evident level I and II axillary lymph node metastases Or Metastases in ipsilateral supraclavicular lymph node(s) with or without axillary or internal mammary lymph node involvement
N3a	Metastases in ipsilateral infraclavicular lymph node(s)
N3b	Metastases in ipsilateral internal mammary lymph node(s) and axillary lymph node(s)
N3c	Metastases in ipsilateral supraclavicular lymph node(s)

[15, 49]. Disease recurrence risk increases proportionally as tumor size increases. The pathologic assessment of tumor size is more important than the clinical assessment. Therefore, the pathology report for all newly diagnosed breast cancers should include a pathologic tumor size assessment. CAP recommends that tumor size be evaluated by gross inspection and that this evaluation be verified by microscopic evaluation.

Pathologic tumor size is the size of the invasive carcinoma component. If the tumor is associated with an in situ carcinoma component, the in situ component outside the invasive tumor is not



included in the determination of tumor size. If the in situ component is admixed with invasive carcinoma, tumor size will include the in situ carcinoma areas. The AJCC staging system for invasive breast cancer recommends that tumor size be recorded before any tumor tissue is removed for special studies [15]. If a biopsy performed prior to excision has removed a significant portion of the tumor, then the original tumor size should be reconstructed on the basis of imaging studies. If initial ultrasonographic evaluation indicates that a patient has a 1.5-cm tumor, the patient should then undergo multiple biopsies. If the excision yields a 0.2-cm invasive cancer associated with biopsy site changes, the tumor should be classified as a T1c rather than a T1a invasive cancer. When two or more distinct invasive tumors are identified, each focus should be measured individually and reported as separate tumors. The foci should not be combined as a single tumor. If there are multiple satellite nodules in association with a dominant mass, the size of the largest mass is considered for staging purposes (Table 10.2b).

**Table 10.2b** Tumor size (T)

	Tumor status
T1	Tumor $\leq 20$ mm in greatest dimension
T1mi	Tumor $\leq 1$ mm in greatest dimension
T1a	Tumor $> 1$ mm but $\leq 5$ mm in greatest dimension
T1b	Tumor $> 5$ mm but $\leq 10$ mm in greatest dimension
T1c	Tumor $> 10$ mm but $\leq 20$ mm in greatest dimension
T2	Tumor $> 20$ mm but $\leq 50$ mm in greatest dimension
T3	Tumor $> 50$ mm in greatest dimension
T4	Tumor of any size with direct extension to the chest wall and/or to the skin (ulceration or skin nodules)
T4a	Extension to the chest wall, not including only the adherence/invasion of the pectoralis muscle
T4b	Ulceration and/or ipsilateral satellite nodules and/or edema (including peau d'orange) of the skin that do not meet the criteria for inflammatory carcinoma
T4c	Both T4a and T4b
T4d	Inflammatory carcinoma

## Lymphovascular Invasion


The presence of tumor emboli in lymphovascular spaces is strongly associated with tumor size and lymph node status and is an independent prognostic indicator of both local recurrence and survival [50–52]. The invasion of the lymphovascular space is usually observed in thin-walled vessels (capillaries or lymphatics). For practical purposes, lymphatic and vascular channel invasions are considered together because their separation does not appear to have prognostic value. The considerable variation in the reported frequency of lymphovascular invasion (20–55 %) most likely represents variation in the criteria used for its identification.

The distinction of lymphovascular space invasion from retraction artifact (tissue shrinkage during processing) may be very difficult. Strict histological criteria include the identification of tumor emboli in clear spaces lined by endothelial cells at the periphery of invasive carcinoma (most commonly 0.1–0.2 cm from the edge of the invasive cancer) and differences in the shape of tumor clusters and that of the surrounding space. Furthermore, the presence of red blood cells and proximity to the neurovascular bundle are helpful in identifying the invasion of the lymphovascular space (Fig. 10.1) [53, 54]. IHC studies for endothelial markers can be used to identify this invasion, but they are not very practical, and results should be interpreted with caution.

## Tumor Histologic Type and Grade

Invasive breast carcinomas are classified on the basis of their morphological features, either the growth pattern (histologic type) or the degree of differentiation (grade). Both histologic type and grade are powerful prognostic markers, and they correlate with biological features [55, 56]. The identification of special histologic types enables further refinement of the prediction of clinical outcome. CAP recommends using the World Health Organization classification system, which recognizes 20 different histologic types (Table 10.3). The majority of breast cancers do

**Fig. 10.1** (a) Lymphovascular invasion of breast carcinoma. The tumor embolus is present within the thin-walled vessel. (b) Immunohistochemical staining for factor VIII highlights the endothelial lining of the thin-walled vessel



not exhibit specific histologic features and are classified as invasive ductal carcinoma of no special type [55, 56]. Some histologic types have distinct clinical behavior and are associated with a particularly favorable prognosis. Special histologic types that have been associated with good prognosis include tubular, invasive cribriform, mucinous, and adenoid cystic carcinomas [57–72]. Strict diagnostic criteria must be used to ensure the accuracy of diagnosis and, consequently, the prediction of outcome.

Other special types of breast cancer carry an unfavorable prognosis; these include metaplastic carcinoma and invasive micropapillary carcinoma [73–79]. Other histologic types may not have prognostic significance but are associated with a distinct biology and specific clinical behavior. Invasive lobular carcinoma, for example, is bilateral and multifocal more often than other histologic types. This type of breast cancer is more likely to occur in older patients, to be larger in size, and to be positive for steroid receptors and negative for Her2/neu overexpression.

**Table 10.3** Histological classification of special types of breast carcinomas

1	Invasive lobular carcinoma
2	Tubular carcinoma
3	Cribriform carcinoma
4	Carcinoma with medullary features
5	Metaplastic carcinoma
6	Carcinoma with apocrine differentiation
7	Salivary gland/skin adnexal-type tumors
8	Adenoid cystic carcinoma
9	Mucoepidermoid carcinoma
10	Polymorphous carcinoma
11	Mucinous carcinoma and carcinoma with signet ring cell differentiation
12	Carcinoma with neuroendocrine features
13	Invasive papillary carcinoma
14	Invasive micropapillary carcinoma
15	Secretory carcinoma
16	Oncocytic carcinoma
17	Sebaceous carcinoma
18	Lipid-rich carcinoma
19	Glycogen-rich clear cell carcinoma
20	Acinic cell carcinoma

Tumor grade has been recognized for several decades as an important component of the pathologic evaluation of breast carcinoma, and different grading systems have been used. Currently, the Nottingham combined histologic grade (the Elston-Ellis modification of the Scarff-Bloom-Richardson grading system) is widely accepted in the USA, Europe, and other parts of the world [56]. This system is also endorsed by AJCC and the European Union [15]. Although the determination of tumor grade always includes an element of subjectivity, the Nottingham grading system attempts to minimize this subjectivity by a strict definition of evaluation criteria. This grading system is based on a semiquantitative assessment of the morphologic characteristics of the tumor, including tubule formation, pleomorphism, and mitotic count. Each parameter is evaluated and assigned a value of 1 (favorable) to 3 (unfavorable), and the scores of all three categories are summed (Table 10.4a). A combined score of 3–5 points is designated grade 1 (low grade), a combined score of 6 or 7 points is grade 2 (intermediate grade), and a combined score of 8 or 9 points

is grade 3 (high grade) (Table 10.4b) (Fig. 10.2). Adherence to strict criteria is necessary for reproducibility so that grading can be used as a prognostic marker [50].

## Hormone Receptors

Estrogen and progesterone are two important endocrine steroid molecules that work together in normal breast development and play critical roles in breast differentiation and tumorigenesis [80, 81]. Estrogen and progesterone act through their nuclear receptors, the estrogen receptor (ER) and progesterone receptor (PR). ER regulates various growth factor receptor pathways, downstream oncogenes, and multiple genes, such as PR, pS2, and bcl-2. Therefore, the presence of PR indicates that ER is present and functional. ER and PR levels in breast carcinoma cells strongly correlate with patient age and inversely correlate with histologic grade, proliferation rate, and Her2/neu overexpression [82–84]. The development of drugs that target the ER pathway has significantly improved the clinical outcomes of patients with hormone receptor-positive tumors, and an evaluation of ER and PR has become a mandatory component of the pathologic evaluation of breast carcinomas [85].

Although both ER and PR are considered weak prognostic factors, the main clinical utility of these markers is in predicting the response to hormonal therapy. The absence of ER predicts a lack of response to hormonal therapy in both adjuvant and metastatic settings. Although hormone receptors were historically analyzed by ligand binding assay, IHC evaluation has been the standard of practice for the last two decades. Hormone receptors in fixed paraffin-embedded tissues are stable; therefore, receptor assays can be performed on archived tissues. Evidence suggests that the interlaboratory variance in ER and PR data is as high as 30 %. Reproducibility can be enhanced with strict adherence to the defined guidelines for ER and PR testing. ASCO and CAP have provided important guidelines on hormone receptor evaluation, emphasizing the importance of internal and external quality assurance and of preanalytical



**Table 10.4a** Breast cancer grade scoring

	1	2	3
Tubule formation	>75 % of tumor area	10–75 % of tumor area	<10 % of tumor area
Nuclear atypia/ pleomorphism	Nuclei small, slight increase in size in comparison with normal breast epithelium	Nuclei larger than normal epithelium variation in size and shape	Marked variation in size and shape occasional large bizarre forms prominent nucleoli
Mitotic count	≤4/10 HPF <sup>a</sup>	5–9/10 HPF <sup>a</sup>	≥10/10 HPF <sup>a</sup>


<sup>a</sup>HPF high-power field

**Table 10.4b** Overall grade

Grade 1 = scores 3–5

Grade 2 = scores 6 or 7

Grade 3 = scores 8 or 9



**Fig. 10.2** (a) Low-grade invasive ductal carcinoma. The majority of tumor cells form tubules that are lined by uniform tumor cells. (b) High-grade invasive ductal carcinoma characterized by solid growth of pleomorphic tumor cells. Note the numerous mitotic figures

and analytical variables and interpretation criteria (Table 10.5) [85–87]. Most guidelines recommend reporting both the proportion of positively stained nuclei and the intensity of nuclear staining (Fig. 10.3). Although different cutoff values have been used to determine positivity, clinical data indicate that ER positivity as low as 1 % can identify patients who would benefit from hormonal therapy [88–91].

ER and PR are typically evaluated at the time of initial diagnosis. Although ER and PR are generally considered static rather than dynamic tumor markers, some variance can occur depend-


ing on disease progression and hormone treatment. Suspected biological cancer progression or a second primary breast cancer may warrant the evaluation of another tumor sample for hormone receptor status.

## Her2/Neu

The human Her2/neu gene is located on chromosome 17 and encodes the transmembrane protein p185, which exhibits structural homology to the epidermal growth factor receptor [92, 93]. The

**Table 10.5** Summary of CAP guideline recommendations for ER/PR testing by IHC

1	Samples from all patients with breast carcinoma should be tested. Large, particularly multiple-core, biopsies of the tumor are preferred for testing if they are representative of the tumor (grade and type) at resection
2	The time from tissue acquisition to fixation should be as short as possible. Samples for ER and PR testing are fixed in 10 % formalin for 6–72 h. Cold ischemia time, fixative type, and the time the sample was placed in fixative must be recorded
3	Storage of slides for more than 6 weeks before analysis is not recommended
4	Validation of any test must be performed before the test is offered
5	Validation must be performed using a clinically validated ER or PR test method. Revalidation should be performed whenever there is a significant change to the test system, such as a change in the primary antibody clone or introduction of new antigen retrieval or detection systems
6	Positive for ER or PR if $\geq 1$ % of tumor cell nuclei are immunoreactive
7	Negative for ER or PR if $< 1$ % of tumor cell nuclei are immunoreactive in the presence of evidence that the sample can express ER or PR (positive intrinsic controls are observed)
8	Uninterpretable for ER or PR if no tumor nuclei are immunoreactive and internal epithelial elements present in the sample or separately submitted from the same sample lack any nuclear staining



**Fig. 10.3**

Immunohistochemical staining for estrogen receptor. This invasive ductal carcinoma shows that the majority of tumor cells are positive for estrogen receptor




epidermal growth factor receptors bind to extracellular growth factors and ligands, are activated via phosphorylation, and are internalized for signaling or degradation. All normal cells and the majority of breast cancer cells have two copies of the Her2/neu gene and produce low levels of the encoded protein p185. The Her2/neu gene is involved in several regulatory pathways, including cell motility, invasion, apoptosis, and proliferation, and affects other procarcinogenic pathways. Thus, the Her2/neu gene has diverse effects on breast cancer biology.

Extensive studies have demonstrated that amplification of the Her2/neu gene and overexpression of its associated protein are present in 15–30 % of breast cancers and are associated with high histologic grade, negative ER status, a high proliferation rate, and an aggressive clinical course with adverse clinical outcome [94–99]. Patients with a Her2/neu-positive tumor have a significantly poorer prognosis than patients with a Her2/neu-negative tumor. However, Her2/neu does not have sufficient prognostic significance to be used in clinical practice as a prognostic factor.

The importance of Her2/neu testing in clinical practice was emphasized by the development of anti-Her2/neu therapies. Trastuzumab (Herceptin) is a humanized monoclonal antibody that inter-

feres with the Her2/neu receptor. Since its approval by the US Food and Drug Administration for the treatment of invasive breast cancers with Her2/neu overexpression, trastuzumab has had a major impact on the outcomes of patients with Her2/neu-positive breast cancer. Many clinical trials have found that the combination of trastuzumab and cytotoxic chemotherapy prolongs disease-free survival and overall survival of patients with Her2/neu-positive disease. Because patients with Her2/neu-negative disease do not benefit from trastuzumab therapy, Her2/neu testing is essential for the selection of patients to receive this therapy [98].

The most commonly used methods to evaluate Her2/neu in breast cancer are IHC and fluorescent in situ hybridization (FISH) or chromogenic in situ hybridization (CISH). Using a Her2/neu-specific antibody, IHC semiquantitatively detects the Her2/neu protein at the cell membrane. IHC is a slide-based assay that can be performed on fixed, paraffin-embedded primary tumor tissues. It facilitates the microscopic visualization of the primary tumor and the semiquantitative analysis of Her2/neu protein levels in the tumor cells. FDA-approved IHC assay kits, as well as other commercial and in-house reagents, are available for IHC testing for Her2/neu (Fig. 10.4). Two-colored FISH



**Fig. 10.4**

Immunohistochemical staining for Her2/neu protein. Positive staining is defined as intense complete membranous staining


analysis and CISH analysis for Her2/neu involve a fluorescent probe or a chromogenic probe, respectively, to quantify the Her2/neu gene copy number in the tumor cells (Fig. 10.5) [94, 95]. Many laboratories use IHC as an initial screening test to identify the more than 50 % of patients whose disease is clearly Her2/neu-negative (0 or 1+). Equivocal or weakly positive cases are then retested using FISH or CISH. To minimize the subjectivity and variation of Her2/neu testing, ASCO and CAP have collaborated to provide Her2/neu testing guidelines, which are updated and refined as additional clinical data become available, to standardize the preanalytical and analytical procedures for performing Her2/neu testing and interpreting the results (Table 10.6).

Updated recommendations for Her2/neu testing in breast carcinoma are as follows. A positive result is a score of 3+ on IHC stain (intense, uniform circumferential staining) in  $\geq 10$  % of invasive carcinoma cells, in situ hybridization indicating more than six gene copies per nucleus, or a FISH gene ratio (ratio of Her2/neu gene signals to chromosome 17 signals) equal to or greater than 2. A negative result is a 0 or 1+ on IHC staining, a FISH result of less than 4 Her2/neu gene copies per nucleus, or a FISH gene ratio of less than 2.0 (Table 10.7) [95].

## Measures of Proliferation

Growth fraction or tumor proliferative capacity is one of the hallmarks of cancer and reflects an imbalance of cell proliferation and cell death.

There are many different methods for evaluating tumor proliferation: mitotic count, thymidine labeling index, S-phase fraction by flow cytometry, and IHC using antibodies against specific cell cycle antigens, such as Ki-67, proliferating cell nuclear antigen, topoisomerase II alpha, mitosin, and cyclins [100]. Mitotic count is usually expressed as the number of mitoses per light microscope high-power field. Although some technical and analytical factors, such as tissue fixation and variations in the size of high-power fields, affect mitotic counting, many studies have demonstrated that mitotic count is an independent prognostic factor. Mitotic count is the most cost-effective and easiest method to evaluate tumor proliferation, and it has been an important component of all breast carcinoma grading systems. Flow cytometric evaluation of S-phase provides important prognostic information for both patients with lymph node-negative disease and those with lymph node-positive disease. Strong correlations between high S-phase fraction and other adverse prognostic factors, such as high tumor grade, ER negativity, and Her2/neu expression, have been



**Fig. 10.5** Fluorescent in situ hybridization for the Her2/neu gene. An increased number of Her2/neu genes (*red signals*) is evident compared to 1 or 2 copies of chromosome 17 centromere sequences (*green signals*). This tumor is classified as Her2/neu gene-amplified carcinoma

**Table 10.6** Recommendations for Her2/neu testing in breast cancer: ASCO/CAP guideline update

One tumor sample from each patient with breast cancer must be tested for either Her2 protein expression or Her2 gene expression using a validated Her2 test
A Her2 test result must be reported as positive if IHC 3+ positive or ISH positive using either a single- or dual-probe ISH
A Her2 test result must be reported as equivocal and a reflex test must be ordered on the same specimen if IHC 2+ equivocal or ISH equivocal using single-probe ISH or dual-probe ISH
A Her2 test result must be reported as negative if a single test performed on a tumor specimen shows IHC 1+ negative or IHC 0 negative or ISH negative using single-probe ISH or dual-probe ISH
A Her2 test result must be reported as indeterminate if technical issues prevent one or both tests performed on a tumor specimen from being reported as positive, negative, or equivocal. This may occur if specimen handling was inadequate
The fixation process must be initiated quickly for any specimen used for Her2 testing
The laboratory must conform to the standards set for CAP accreditation or an equivalent accreditation authority, including initial test validation, ongoing internal quality assurance, ongoing external proficiency testing, and routine periodic performance monitoring
If an apparent histopathological discordance is observed in any Her2 testing situation, the pathologist should consider ordering additional Her2 testing and conferring with the oncologist and should document the decision-making process and results in the pathology report. As part of the Her2 testing process, the pathologist may pursue additional Her2 testing without conferring with the oncologist

**Table 10.7** 2013 Her2 testing guidelines update: summary of changes to the testing algorithm

Her2 result	IHC	ISH
Positive	>10 % of invasive tumor cells display strong circumferential membrane staining	Amplified ratio of Her2/CEP17 of $\geq 2.0$ or average HER2 signals $\geq 6$ signals/cell (regardless of ratio) in >10 % of tumor cells
Negative	IHC 0: no staining observed or membrane staining is incomplete, faint/barely perceptible, and in $\leq 10$ % of the invasive tumor cells IHC 1+: incomplete membrane staining that is faint/barely perceptible and in >10 % of the invasive tumor cells	Her2/CEP17 ratio $< 2$ and Her2 signals/cell $< 4$ , regardless of ratio
Equivocal reflex test must	2+ based on circumferential membrane staining, incomplete, weak, or moderate in >10 % of the invasive tumor cells; or complete and circumferential membrane intense staining in $\leq 10$ % of the invasive tumor cells	Dual-probe Her2/CEP 17 ratio $< 2.0$ with an average Her2 copy number $\geq 4.0$ and $< 6.0$ signals/cell
Indeterminate	Technical issues prevent assay from being conclusive (e.g., issues with controls, specimen handling, artifacts, or analytical failure) Assay must be repeated before diagnosis can be rendered	

reported. Widespread clinical use has been limited by the requirement for fresh tissue for flow cytometry.


The immunohistochemical evaluation of proliferation-related markers has been utilized extensively in clinical practice. Ki-67, a nonhistone nuclear protein expressed only in proliferating cells that is absent during the inactive phases

of the cell cycle, is the most commonly used marker [101, 102]. The Ki-67 labeling index, which represents the percentage of tumor cells that are positive for Ki-67 nuclear staining (Fig. 10.6), correlates well with the S-phase fraction and mitotic count and is an independent prognostic factor for breast carcinoma. A high Ki-67 index at diagnosis is usually associated



**Fig. 10.6**

Immunohistochemical staining for Ki-67. Nuclear staining in the tumor cells indicates that tumor cells are in the proliferation phase. The percentage of positive nuclei is reported as the tumor proliferation fraction. **(a)** Invasive ductal carcinoma with low tumor proliferation determined by the low number of cells with positive staining for Ki-67. **(b)** Invasive ductal carcinoma with high tumor proliferation. Note that more than 30 % of the tumor cells are positive for Ki-67



with a less favorable natural history and worse patient outcome [102].

Although the measurement of proliferation provides important prognostic information, the optimal method for use in clinical practice has not been determined. Proliferation rates are a continuum and are not bimodal. Cutoff points for low and high proliferation rates are difficult to determine. Because of the lack of standardization of evaluation methods, neither CAP nor ASCO endorses a specific test for the evaluation of tumor proliferation. In addition to its prognostic

significance, the tumor proliferation index may also provide predictive information in patients receiving neoadjuvant chemotherapy. Higher tumor response rates have been reported in breast cancer patients with a high Ki-67 index compared to those with a low Ki-67 index. However, the cutoff value for Ki-67 for clinical treatment triage remains undetermined. A change in a tumor's Ki-67 index after 2–12 weeks of endocrine therapy may also be a pharmacodynamic biomarker of treatment efficacy and a predictor of long-term outcome.

## Methods of Combining Prognostic Factors

In clinical practice, all prognostic and predictive factors described here are evaluated. Individually, each of them is clinically important. However, treatment decisions are based predominantly on the tumor stage (lymph node status and tumor size). Attempts to refine prognostication for individual patients have involved the combination of these individual factors into prognostic indices. Many different indices have been proposed; they are formulated by weighting each prognostic factor to arrive at a clinically useful stratification (good prognosis versus poor prognosis) for individual patients with invasive breast carcinoma.

1. Adjuvant! Online is a web-based statistical program that is used extensively for prognostication for patients with early-stage breast cancer ([www.adjuvantonline.com](http://www.adjuvantonline.com)). Based on the patient's age, menopausal status, tumor size, tumor grade, lymph node status, and ER status, the likelihoods of disease-free and overall survival as 10-year clinical outcomes without or with therapy (both chemotherapy and hormonal therapy) are calculated.
2. The Nottingham Prognostic Index (NPI) is the only breast carcinoma prognostic index that has been validated by both retrospective and prospective clinical studies. NPI includes lymph node status, tumor size, and tumor grade and is calculated using the following formula:  

$$\text{NPI} = \text{tumor size (0.2)} + \text{lymph node stage (1, 2, or 3)} + \text{histologic grade (1, 2, or 3)}.$$
 Lymph node stage is defined by the number of positive lymph nodes (1, all nodes negative; 2, 1–3 positive nodes; 3, 4 or more positive lymph nodes). Five different prognostic categories based on the NPI have been identified, and studies have shown that these categories are associated with significantly different 5-, 10-, and 15-year survival rates.
3. Breast Cancer Prognostic Score (BCPS) is a prognostic score calculated by evaluating the ER, PR, Her2/neu, and three components of the Nottingham histologic grade (tubule formation, mitosis, and nuclear pleomorphism).

Although initial findings have suggested that this scoring system correlates well with some other prognostic testing, such as Oncotype Dx recurrence testing, there are no clinical outcome studies to validate this finding.

## Emerging Molecular and Genomic Markers

In the past, breast cancers were simply treated based on the above described clinicopathological features (tumor size, histologic grade, lymph node status) and tumor biomarkers such as ER, PR, and HER2/neu. Systemic chemotherapy was applied nearly universally to breast cancers of high clinical stage regardless of histologic grade or biomarker profile. Even among patients with early-stage breast cancer, approximately 60 % received systemic chemotherapy [103]. However, many trials, including the National Surgical Adjuvant Breast and Bowel Project (NSABP) trial B20, demonstrated that less than 10 % of patients with node-negative, hormone receptor-positive breast cancers received significantly greater benefit from additional systemic chemotherapy compared to hormonal therapy alone, indicating that a majority of early-stage breast cancers were over-treated [104]. Therefore, identifying patients with poor prognosis that require aggressive treatment among those with the same histopathological and biomarker features remains a clinical challenge. In recent years, the application of genomic and molecular technologies to the problem of predicting breast cancer prognosis has revealed encouraging results. In 2000, Perou and his colleague were the first to examine the molecular landscape of breast cancers, and they demonstrated that the molecular signature of a breast cancer was determined by the expression of a set of genes within each tumor and that intrinsic genetic expression signatures predict the clinical outcome of the disease [105]. Sorlie et al. have categorized breast cancers into five intrinsic molecular subtypes: luminal A, luminal B, HER2 enriched, basal, and normal-like. They demonstrated that molecular subtypes were prognostic, with luminal A tumors having the best prognosis and basal and HER2-enriched tumors having the



worst prognosis [7]. Luminal types of breast cancers express high levels of ER and luminal epithelial genes compared with normal breast luminal epithelium. Compared to luminal A tumors, luminal B tumors express fewer ER-related genes and have a higher proliferation rate. Luminal B tumors also include those tumors that express ER and have HER2/neu overexpression or amplification. By contrast, HER2-enriched breast cancers exhibit HER2/neu gene amplification but do not express ER-related genes. The basal type of tumors frequently does not express ER, PR, and HER2/neu (“triple negative” tumors) but also expresses basal cytokeratins 5/6 and 17. Luminal types of breast cancers have an indolent clinical course. They have a low distant recurrence rate that persists even 10–15 years after the diagnosis. Compared to luminal A tumors, luminal B breast cancers have a more aggressive clinical course and need to be treated more aggressively, usually requiring both hormonal therapy and chemotherapy. HER2-enriched breast cancers have an aggressive natural clinical course; however, they respond very well to HER2-targeted therapy. Basal types of breast cancers have a poor prognosis; they tend to recur during the first 3 years after diagnosis, and currently there are no specific targeted therapies for them. Although the molecular classification of breast cancer has sparked major excitement among researchers and clinicians, these tests have not been fully incorporated into clinical practice. The main limitations of gene signature profiling include difficulties in reproducing the specific gene sets, the expense of testing, and standardization of reporting. Consequently, surrogates using IHC studies have been proposed by the St. Gallen International Expert Consensus Group (Table 10.8) [106]. However, there are significant discordances among the molecular classifications by IHC approximation and genetic profiling.

In the past few years, significant attempts have sought to create genomic tests that can be utilized in routine clinical practice. There are currently a few commercially available multigene prognostic/predictive tests for breast cancer. Among them, three tests have been the most widely used in clinical practice.

**Table 10.8** Immunohistochemical (IHC) surrogates for intrinsic molecular subtypes of breast cancer

Intrinsic molecular subtype	IHC surrogates
Luminal A	ER and/or PR + HER2 - Ki-67 low (<14 %)
Luminal B	HER2- ER and/or PR + Ki-67 high (≥14 %) HER2+ ER and/or PR + Any Ki-67
HER2 enriched	HER2 overexpressed or amplified ER- and PR-
Basal-like	ER-, PR-, and HER- CK5/6+ and/or CK17+ and/or EGFR+

The *Oncotype DX* test (Genomic Health, Redwood, CA, USA) is an RT-PCR-based assay to measure gene expression in formalin-fixed paraffin-embedded (FFPE) samples. It measures a panel of 21 genes, including 16 cancer-related genes (ER, PR, Bcl2, SCUBE2, HER2, GRB7, Ki-67, STK15, survivin, cyclin B1, MYBL2, stromelysin 3, cathepsin L2, GSTM1, CD68, and BAG1) and 5 housekeeping control genes (beta-actin, GAPDH, RPLPO, GUS, and TFRC). Based on the expression levels of these 21 genes, a recurrence score (RS) is generated, which stratifies breast cancer patients into three risk groups: low risk (RS <18), intermediate risk (RS 18–30), and high risk (RS ≥31), with 10-year distant recurrence rates of 6.8 %, 14.3 %, and 30.5 %, respectively [107]. The test was originally designed to predict distant recurrence within 10 years in hormone receptor-positive and node-negative breast cancers [107]. Its clinical application was later expanded to hormone receptor-positive breast cancer patients with up to three positive lymph nodes [108, 109].

The *Oncotype DX* test is the most widely used molecular test in the clinical setting for making treatment decisions. It is recommended by both the National Comprehensive Cancer Network (NCCN) and the St. Gallen Consensus [110, 111]. The test is specifically applied to hormone receptor-positive breast cancers with 0–3 positive nodes that are to be treated with hormonal

therapy. The general consensus is that hormonal therapy without systemic chemotherapy is sufficient for patients with a low RS. However, for patients with a high RS, systemic chemotherapy is warranted in addition to hormonal therapy to further reduce the future distant recurrence by 50 %. An ongoing prospective Trial Assigning Individualized Options for Treatment (Rx) (TAILORx, NCT00310180) has been designed to address whether breast cancer patients with intermediate RS need systemic chemotherapy in addition to hormonal therapy.

The *MammaPrint* assay is the second most commonly ordered molecular test approved by the US Food and Drug Administration (FDA). It is a 70-gene expression assay developed by The Netherlands Cancer Institute based on 78 treatment-naïve node-negative breast cancers that were 5 cm or less from women younger than 55 years old [112]. This 70-gene expression assay was further validated in several hundreds of patients with both node-positive and node-negative diseases [113–115]. It is prognostic for early distant recurrence within the first 5 year after diagnosis and predictive for chemoresponse in poor prognostic patients. The MammaPrint assay can be applied to both hormone-positive and hormone-negative breast cancers. This 70-gene test is a better prognostic predictor than clinicopathological risk assessments, including the NIH and St. Gallen risk assessments and Adjuvant! Online [114, 115]. The general consensus is that no systemic chemotherapy is required in patients with low-risk cancers determined by both clinical evaluation and the MammaPrint test, whereas systemic chemotherapy is recommended in patients with high-risk cancers determined by both assessments. In practice, however, some breast cancers have conflicting risks as evaluated by clinical assessments such as Adjuvant! Online and molecular assessments such as the MammaPrint assay. An ongoing prospective trial “Microarray in Node-Negative and 1–3 Positive Lymph Node Disease May Avoid Chemotherapy” (MINDACT, NCT00433589) is designed to address whether chemotherapy can be safely avoided in patients who are predicted to be at low risk by the MammaPrint test but at high risk by clinical

assessment using Adjuvant! Online. Preliminary results presented at the recent European Cancer Congress meeting are promising [116].

The *Prosigna* test is a PAM50-based assay offered by NanoString Technologies (Seattle, WA) that has been recently cleared by the US FDA. It measures the expression of 50 genes using a proprietary technology called the “nCounter Dx Analysis System.” These 50 genes were selected from the original gene set used for molecular classification by Perou et al. [7, 105]. Based on the expression levels of these 50 genes and clinical variables, a risk of recurrence (ROR) score is generated that correlates to one of the five molecular subtypes, i.e., luminal A, luminal B, HER-enriched, basal, or normal-like subtypes [117]. The ROR score is also used to assess the risk of distant recurrence at 10 years for hormone receptor-positive stage I–III breast cancers to be treated with adjuvant endocrine therapy [117]. A study has demonstrated that the PAM50 assay is superior to IHC profiling for both prognosis and the prediction of benefit from adjuvant tamoxifen [118]. Recent studies have also demonstrated that the Prosigna assay is superior to the Oncotype DX test in predicting late distant recurrence after 5–10 years [119, 120].

The key features of the abovementioned multigene prognostic/predictive tests and their clinical applications are summarized in Table 10.9.

## Conclusion

The accurate diagnosis of breast cancer is a critical prerequisite to the therapy decision-making process. Most of the prognostic factors currently used in clinical practice are based on meticulous pathologic evaluation of the primary tumor and lymph nodes. In addition to lymph node status and the size of the primary tumor, tumor grade and the presence of lymphovascular space invasion are independent prognostic factors associated with clinical outcome and should be carefully evaluated and reported for all newly diagnosed breast cancers. ER/PR and Her2/neu status should be evaluated by standard methods and reported as recommended by national guidelines to determine the likelihood of response to endocrine and anti-Her2/neu therapy,

**Table 10.9** Emerging molecular and genomic tests for invasive breast cancers

	Assay	Provider	Tissue type	Clinical indications	Prognostic/predictive value
Oncotype DX	qRT-PCR	Genomic Health (Redwood City, CA, USA)	FFPE	ER+ 0–3 node positive To be treated with tamoxifen or aromatase inhibitors (AI)	Prognostic for distant recurrence in 10 years Predictive for chemoresponse in high recurrence score group
MammaPrint	Microarray	Agendia (Amsterdam, Netherlands)	Fresh/frozen or FFPE	0–3 node positive <5 cm All ages ER+ or ER-	Prognostic for early distant recurrence within first 5 years after diagnosis Predictive for chemoresponse in poor prognosis group
Prosigna (PAM 50)	nCounter	NanoString Technologies (Seattle, WA, USA)	FFPE	Stage I–III All ages ER+ or ER-	Prognostic based on assigned intrinsic molecular subtypes Predictive for tamoxifen benefit in luminal A cancers

respectively. New technologies and genome-wide approaches have the potential to identify additional prognostic and predictive markers for invasive breast cancer. Future research should focus on the identification of these markers and the correlation of their significance with those of established factors described in this chapter.

## References

- Hortobagyi GN. Treatment of breast cancer. *N Engl J Med.* 1998;339:974–98.
- Fisher B. Biological and clinical considerations regarding the use of surgery and chemotherapy in the treatment of primary breast cancer. *Cancer.* 1977;40(1 Suppl):574–87.
- Tabar L, Fagerberg G, Day NE, Duffy SW, Kitchin RM. Breast cancer treatment and natural history: new insights from results of screening. *Lancet.* 1992;339:412–4.
- Eifel P, Axelson JA, Costa J, Crowley J, Curran WJ Jr, Deshler A, et al. National Institutes of Health Consensus Development Conference Statement: adjuvant therapy for breast cancer, November 1–3, 2000. *J Natl Cancer Inst.* 2001;93(13):979–89.
- Mittendorf EA, Ballman KV, McCall LM, Yi M, Sahin AA, Bedrosian I, et al. Evaluation of the stage IB designation of the American Joint Committee on Cancer staging system in breast cancer. *J Clin Oncol.* 2014;32:1–10.
- Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, et al. Prognostic factors in breast cancer. College of American Pathologists Consensus Statement 1999. *Arch Pathol Lab Med.* 1999;124(7):966–78.
- Goldhirsch A, Wood WC, Coates AS, Gelber RD, Thürlimann B, Senn HJ, et al. Strategies for subtypes – dealing with the diversity of breast cancer: highlights of the St. Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2011. *Ann Oncol.* 2011;22(8):1736–47.
- National Institutes of Health NIH consensus conference. Treatment of early-stage breast cancer. *JAMA.* 1991;265:391–5.
- Wilking N, Rutqvist LE, Carstensen J, Mattsson A, Skoog L. Prognostic significance of axillary nodal status in primary breast cancer in relation to the number of resected nodes. Stockholm Breast Cancer Study Group. *Acta Oncol.* 1992;31:29–35.
- Carter CL, Allen C, Henson DE. Relation of tumor size, lymph node status, and survival in 24,740 breast cancer cases. *Cancer.* 1989;63:181–7.
- Fisher B, Bauer M, Wickerham DL, Redmond CK, Fisher ER, Cruz AB, et al. Relation of number of positive axillary nodes to the prognosis of patients with primary breast cancer. An NSABP Update. *Cancer.* 1983;52(9):1551–7.
- Nemoto T, Vana J, Bedwani RN, Baker HW, McGregor FH, Murphy GP. Management and survival of female breast cancer: results of a national survey by the American College of Surgeons. *Cancer.* 1980;45(12):2917–24.

13. Whitworth P, McMasters KM, Tafra L, Edwards MJ. State-of-the-art lymph node staging for breast cancer in the year 2000. *Am J Surg*. 2000;180(4):262–7.
14. Tavassoli F, Eusebi V, editors. Tumors of the mammary gland, AFIP atlas of tumor pathology, vol. 4. Maryland: ARP Press; 2009.
15. In Edge SB, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti A, editors. AJCC cancer staging manual. 7th ed. New York: Springer; 2010.
16. Badve S. Sentinel lymph node biopsy. In: Dabbs DJ, editor. Breast pathology. 1st ed. Elsevier/Saunders: Philadelphia; 2012. p. 93–116.
17. Hoda SA. Invasive ductal carcinoma: assessment of prognosis with morphologic and biologic markers. In: Rosen's breast pathology. 4th ed. Philadelphia: Lippincott Williams & Wilkins, a Wolters Kluwer business; 2014. p. 413–68.
18. Weaver DL. Sentinel node biopsy and lymph node classification in the 7th edition staging manual. In: O'Malley FP, Pinder SE, Mulligan AM, editors. Breast pathology. 2nd ed. Philadelphia: Elsevier; 2011. p. 295–305.
19. Hartveit F, Samsonsen G, Tangen M, Halvorsen JF. Routine histological investigation of the axillary nodes in breast cancer. *Clin Oncol*. 1982;8(2):121–6.
20. Hartveit F. The routine histological investigation of axillary lymph nodes for metastatic breast cancer. *J Pathol*. 1984;143(3):187–91.
21. Ho A, Morrow M. The evolution of the locoregional therapy of breast cancer. *Oncologist*. 2011;16(10):1367–79.
22. Kingsley WB, Peters GN, Cheek JH. What constitutes adequate study of axillary lymph nodes in breast cancer. *Ann Surg*. 1985;201:311–4.
23. Fisher B, Wolmark N, Bauer M, Redmond C, Gebhardt M. The accuracy of clinical nodal staging and of limited axillary dissection as a determinant of histologic nodal status in carcinoma of the breast. *Surg Gynecol Obstet*. 1981;152:765–72.
24. Luz DA, Ribeiro Jr U, Chassot C, Collet E, Silva Fde S, Ceconello I, et al. Carnoy's solution enhances lymph node detection: an anatomical dissection study in cadavers. *Histopathology*. 2008;53(6):740–2.
25. Gregurek SF, Wu HH. Can GEWF solution improve the retrieval of lymph nodes from colorectal cancer. *Arch Pathol Lab Med*. 2009;133(1):83–6.
26. Morrow M, Evans J, Rosen PP, Kinne DW. Does clearing of axillary lymph nodes contribute to accurate staging of breast carcinoma? *Cancer*. 1984;53:1329–32.
27. Lester SC, Bose S, Chen YY, Connolly JL, de Baca ME, Fitzgibbons PL, et al. Protocol for the examination of specimens from patients with invasive carcinoma of the breast. *Arch Pathol Lab Med*. 2009;133(10):1515–38.
28. Dowlatshahi K, Fan M, Anderson JM, Bloom KJ. Occult metastases in sentinel nodes of 200 patients with operable breast cancer. *Ann Surg Oncol*. 2001;8(8):675–81.
29. Lawrence WD. Association of Directors of Anatomic and Surgical Pathology. ADASP recommendations for processing and reporting lymph node specimens submitted for evaluation of metastatic disease. *Virchows Arch*. 2001;439(5):601–3.
30. Neuman HB. Management of the axilla. *Surg Clin North Am*. 2013;93(2):429–44.
31. Rao R, Euhus D, Mayo HG, Balch C. Axillary node interventions in breast cancer: a systematic review. *JAMA*. 2013;310(13):1385–94.
32. Thompson AM. New standards of care in the management of the axilla. *Curr Opin Oncol*. 2012;24(6):605–11.
33. Zarebczan Dull B, Neuman HB. Management of the axilla. *Surg Clin North Am*. 2013;93(2):429–44.
34. Noguchi M, Morioka E, Ohno Y, Noguchi M, Nakano Y, Kosaka T. The changing role of axillary lymph node dissection for breast cancer. *Breast Cancer*. 2013;20(1):41–6.
35. Giuliano AE, Hunt KK, Ballman KV, Beitsch PD, Whitworth PW, Blumencranz PW, et al. Axillary dissection vs no axillary dissection in women with invasive breast cancer and sentinel node metastasis: a randomized clinical trial. *JAMA*. 2011;305(6):569–75.
36. Lyman GH, Giuliano AE, Somerfield MR, Benson 3rd AB, Bodurka DC, Burstein HJ, et al. American Society of Clinical Oncology guideline recommendations for sentinel lymph node biopsy in early-stage breast cancer. *American Society of Clinical Oncology. J Clin Oncol*. 2005;23(30):7703–20.
37. Mansel RE, Fallowfield L, Kissin M, Goyal A, Newcombe RG, Dixon JM, et al. Randomized multicenter trial of sentinel node biopsy versus standard axillary treatment in operable breast cancer: the ALMANAC Trial. *J Natl Cancer Inst*. 2006;98(9):599–609. Erratum in: *J Natl Cancer Inst*. 2006;98(12):876.
38. Krag DN, Anderson SJ, Julian TB, Brown AM, Harlow SP, Costantino JP, et al. Sentinel-lymph-node resection compared with conventional axillary-lymph-node dissection in clinically node-negative patients with breast cancer: overall survival findings from the NSABP B-32 randomized phase 3 trial. *Lancet Oncol*. 2010;11(10):927–33.
39. Weaver DL. Pathology evaluation of sentinel lymph nodes in breast cancer: protocol recommendations and rationale. *Mod Pathol*. 2010;23 Suppl 2:S26–32.
40. Turner RR, Giuliano AE, Hoon DS, Glass EC, Krasne DL. Pathologic examination of sentinel lymph node for breast carcinoma. *World J Surg*. 2001;25(6):798–805.
41. Cserni G. Evaluation of sentinel lymph nodes in breast cancer. *Histopathology*. 2005;46(6):697–702.
42. Cserni G. Surgical pathological staging of breast cancer by sentinel lymph node biopsy with special emphasis on the histological work-up of axillary sentinel lymph nodes. *Breast Cancer*. 2004;11(3):242–9; discussion 264–6.
43. Fritzsche FR, Reineke T, Morawietz L, Kristiansen G, Dietel M, Fink D, et al. Pathological processing techniques and final diagnosis of breast cancer sentinel lymph nodes. *Ann Surg Oncol*. 2010;17:2892–8.

44. Weaver DL, Le UP, Dupuis SL, Weaver KA, Harlow SP, Ashikaga T, et al. Metastasis detection in sentinel lymph nodes: comparison of a limited widely spaced (NSABP protocol B-32) and a comprehensive narrowly spaced paraffin block sectioning strategy. *Am J Surg Pathol.* 2009;33:1583–9.
45. Chen SL, Hoehne FM, Giuliano AE. The prognostic significance of micrometastases in breast cancer: a SEER population-based analysis. *Ann Surg Oncol.* 2007;14:3378–84.
46. Wu Y, Mittendorf EA, Kelten C, Tucker SL, Wei W, Middleton LP, et al. Occult axillary lymph node metastases do not have prognostic significance in early stage breast cancer. *Cancer.* 2012;118(6):1507–14.
47. Sahin AA, Guray M, Hunt KK. Identification and biologic significance of micrometastases in axillary lymph nodes in patients with invasive breast cancer. *Arch Pathol Lab Med.* 2009;133(6):869–78.
48. Galimberti V, Cole BF, Zurrida S, Viale G, Luini A, Veronesi P, et al. Axillary dissection versus no axillary dissection in patients with sentinel-node micrometastases (IBCSG 23–01): a phase 3 randomized controlled trial. *Lancet Oncol.* 2013;14(4):297–305.
49. Blamey RW, Hornmark-Stenstam B, Ball G, Blichert-Toft M, Cataliotti L, Fourquet A, et al. ONCOPOOL: a European database for 16,944 cases of breast cancer. *Eur J Cancer.* 2010;46:56–71.
50. Shatat L, Gloyeske N, Madan R, O'Neil M, Tawfik O, Fan F. Microinvasive breast carcinoma carries an excellent prognosis regardless of the tumor characteristics. *Hum Pathol.* 2013;44(12):2684–9.
51. Lee AH, Pinder SE, Macmillan RD, Mitchell M, Ellis IO, Elston CW, et al. Prognostic value of lymphovascular invasion in women with lymph node negative invasive breast carcinoma. *Eur J Cancer.* 2006;42:357–62.
52. Lee A, DeLellis R, Silverman M, Heatley GJ, Wolfe H. Prognostic significance of peritumoral lymphatic and blood-vessel invasion in node-negative carcinoma of the breast. *J Clin Oncol.* 1990;8:1457–65.
53. Rosen PP. Tumor emboli in intramammary lymphatics in breast carcinoma. Pathologic criteria for diagnosis and clinical significance. *Pathol Annu.* 1983;18:215–32.
54. Van den Eynden GG, Van der Auwera I, Van Laere SJ, Colpaert CG, van Dam P, Dirix LY, et al. Distinguishing blood and lymph vessel invasion in breast cancer: a prospective immunohistochemical study. *Br J Cancer.* 2006;94:1643–9.
55. Lakhani SR, Ellis IO, Schnitt SJ, Tan PH, vande Vijver MJ, editors. *World Health Organization classification of tumors. Pathology and genetics of tumors of the breast and female genital organs.* 4th ed. Lyon: IARC Press; 2012.
56. Ellis IO, Galea M, Broughton N, Locker A, Blamey RW, Elston CW. Pathological prognostic factors in breast cancer. II. Histological type. Relationship with survival in a large study with long-term follow-up. *Histopathology.* 1992;20:479–89.
57. Pusztai L, Sotiriou C, Buchholz TA, Meric F, Symmans WF, Esteva FJ, et al. Molecular profiles of invasive mucinous and ductal carcinomas of the breast: a molecular case study. *Cancer Genet Cytogenet.* 2003;141:148–53.
58. Kulkarni N, Pezzi CM, Greif JM, Suzanne Klimberg V, Bailey L, et al. Rare breast cancer: 933 adenoid cystic carcinomas from the National Cancer Data Base. *Ann Surg Oncol.* 2013;20(7):2236–41.
59. Thompson K, Grabowski J, Saltzstein SL, Sadler GR, Blair SL. Adenoid cystic breast carcinoma: is axillary staging necessary in all cases? Results from the California Cancer Registry. *Breast J.* 2011;17(5):485–9.
60. Parl FF, Richardson LD. The histologic and biologic spectrum of tubular carcinoma of the breast. *Hum Pathol.* 1983;14:694–8.
61. Page DL, Dixon JM, Anderson TJ, Lee D, Stewart HJ. Invasive cribriform carcinoma of the breast. *Histopathology.* 1983;7:525–36.
62. Venable JG, Schwartz AM, Silverberg SG. Infiltrating cribriform carcinoma of the breast: a distinctive clinicopathologic entity. *Hum Pathol.* 1990;21:333–8.
63. Cabral AH, Recine M, Paramo JC, McPhee MM, Poppiti R, Mesko TW. Tubular carcinoma of the breast: an institutional experience and review of the literature. *Breast J.* 2003;4:298–301.
64. Anan K, Mitsuyama S, Tamae K, Nishihara K, Iwashita T, Abe Y, et al. Pathological features of mucinous carcinoma of the breast are favourable for breast-conserving therapy. *Eur J Surg Oncol.* 2001;27:459–63.
65. Scopsi L, Andreola S, Pilotti S, Bufalino R, Baldini MT, Testori A, et al. Mucinous carcinoma of the breast. A clinicopathologic, histochemical and immunocytochemical study with special reference to neuroendocrine differentiation. *Am J Surg Pathol.* 1994;18:702–11.
66. Winchester DJ, Chang HR, Graves TA, Menck HR, Bland KI, Winchester DP. A comparative analysis of lobular and ductal carcinoma of the breast: presentation, treatment and outcomes. *J Am Coll Surg.* 1998;186:416–22.
67. Cao AY, Huang L, Wu J, Lu JS, Liu GY, Shen ZZ, et al. Tumor characteristics and the clinical outcome of invasive lobular carcinoma compared to infiltrating ductal carcinoma in a Chinese population. *World J Surg Oncol.* 2012;10:152.
68. Dixon JM, Anderson TJ, Page DL, Lee D, Duffy SW. Infiltrating lobular carcinoma of the breast. *Histopathology.* 1982;6:149–61.
69. Mersin H, Yildirim E, Gulben K, Berberoglu U. Is invasive lobular carcinoma different from invasive ductal carcinoma? *Eur J Surg Oncol.* 2003;29:390–5.
70. Du Toit RS, Locker AP, Ellis IO, Elston CW, Nicholson RI, Blamey RW. Invasive lobular carcinomas of the breast: the prognosis of histopathological subtypes. *Br J Cancer.* 1989;60:605–9.



71. Tse GMK, Ma TKF, Chu WCW, Lam WWM, Poon CSP, Chan W-C. Neuroendocrine differentiation in pure type mammary mucinous carcinoma is associated with favourable histologic and immunohistochemical parameters. *Mod Pathol.* 2004;17:568–72.
72. Tang F, Wei B, Tian Z, Gilcrease MZ, Huo L, Albarracin CT, et al. Invasive mammary carcinoma with neuroendocrine differentiation: histological features and diagnostic challenges. *Histopathology.* 2011;59(1):106–15.
73. Vingiani A, Maisonneuve P, Dell’orto P, Farante G, Rotmensz N, Lissidini G, et al. The clinical relevance of micropapillary carcinoma of the breast: a case-control study. *Histopathology.* 2013;63(2):217–24.
74. Walsh MM, Bleiweiss IJ. Invasive micropapillary carcinoma of the breast: eighty cases of an under-recognized entity. *Hum Pathol.* 2001;32:583–9.
75. Zekioglu O, Erhan Y, Ciris M, Bayramoglu H, Ozdemir N. Invasive micropapillary carcinoma of the breast: high incidence of lymph node metastasis with extranodal extension and its immunohistochemical profile compared with invasive ductal carcinoma. *Histopathology.* 2004;44:18–23.
76. Nassar H, Wallis T, Andea A, Dey J, Adsay V, Visscher D. Clinicopathologic analysis of invasive micropapillary differentiation in breast carcinoma. *Mod Pathol.* 2001;14:836–41.
77. DownsKelly E, Nayeemuddin KM, Albarracin C, Wu Y, Hunt KK, Gilcrease MZ. Matrix-producing carcinoma of the breast: an aggressive subtype of metaplastic carcinoma. *Am J Surg Pathol.* 2009;33:534–41.
78. Lee H, Jung SY, Ro JY, Kwon Y, Sohn JH, Park IH, et al. Metaplastic breast cancer: clinicopathological features and its prognosis. *J Clin Pathol.* 2012;65(5):441–6.
79. Tse GM, Tan PH, Putti TC, Lui PC, Chaiwun B, Law BK. Metaplastic carcinoma of the breast: a clinicopathological review. *J Clin Pathol.* 2006;59:1079–83.
80. Anderson E. The role of oestrogen and progesterone receptors in human mammary development and tumorigenesis. *Breast Cancer Res.* 2002;4:197–201.
81. Allred CD. Commentary: hormone receptor testing in breast cancer: a distress signal from Canada. *Oncologist.* 2008;13:1134–6.
82. Blamey RW. Guidelines on endocrine therapy of breast cancer EUSOMA. *Eur J Cancer.* 2002;38:615–34.
83. Harvey JM, Clark GM, Osborne CK, Allred DC. Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *J Clin Oncol.* 1999;17:1474–81.
84. Cui X, Schiff R, Arpino G, Osborne CK, Lee AV. Biology of progesterone receptor loss in breast cancer and its implications for endocrine therapy. *J Clin Oncol.* 2005;23:7721–35.
85. Hammond MEH, Hayes DF, Dowsett M, Allred DC, Hagerty KL, Badve S, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. *J Clin Oncol.* 2010;28:2784–95. Erratum in *Clin Oncol.* 2010;28:3543.
86. Goldstein NS, Ferkowicz M, Odish E, Mani A, Hastah F. Minimum formalin fixation time for consistent estrogen receptor immunohistochemical staining of invasive breast carcinoma. *Am J Clin Pathol.* 2003;120:86–92.
87. Harris L, Fritsche H, Mennel R, Norton L, Ravdin P, Taube S, et al. American Society of Clinical Oncology 2007 update of recommendations for the use of tumor markers in breast cancer. *J Clin Oncol.* 2007;25:5287–312.
88. Harvey JM, Clark GM, Osborne CK, Allred DC. Predicting response to adjuvant endocrine therapy in breast cancer. *J Clin Oncol.* 2000;17:1474–81.
89. Rhodes A, Jasani B, Barnes DM, Bobrow LG, Miller KD. Reliability of immunohistochemical demonstration of oestrogen receptors in routine practice: interlaboratory variance in the sensitivity of detection and evaluation of scoring systems. *J Clin Pathol.* 2000;53:125–30.
90. Umemura S, Itoh J, Itoh H, Serizawa A, Saito Y, Suzuki Y, et al. Immunohistochemical evaluation of hormone receptors in breast cancer: which scoring system is suitable for highly sensitive procedures? *Appl Immunohistochem Mol Morphol.* 2004;12:8–13.
91. Leake R, Barnes D, Pinder S, Ellis I, Anderson L, Anderson T, et al. Immunohistochemical detection of steroid receptors in breast cancer: a working protocol on behalf of the UK Receptor Group, UK NEQAS, the Scottish Breast Cancer Pathology Group, and the Receptor and Biomarker Study Group of the EORTC. *J Clin Pathol.* 2000;53:634–5.
92. Ross JS, Slodkowska EA, Symmans WF, Pusztai L, Ravdin PM, Hortobagyi GN. The HER-2 receptor and breast cancer: ten years of targeted anti-HER-2 therapy and personalized medicine. *Oncologist.* 2009;14(4):320–68.
93. Sauter G, Lee J, Bartlett JMS, Slamon DJ, Press MF. Guidelines for human epidermal growth factor receptor 2 testing: biologic and methodologic considerations. *J Clin Oncol.* 2009;27:1323–33.
94. Wolff AC, Hammond ME, Schwartz JN, Hagerty KL, Allred DC, Cote RJ, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol.* 2007;25:1–28.
95. Wolff AC, Hammond ME, Hicks DG, Dowsett M, McShane LM, Allison KH, et al. Recommendations for Human Epidermal Growth Factor Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Update. *J Clin Oncol.* 2013;31(31):3997–4013.

96. Hayes DF, Thor AD, Dressler LG, Weaver D, Edgerton S, Cowan D, et al. HER2 and response to paclitaxel in node-positive breast cancer. *N Engl J Med*. 2007;357:1496–506.
97. Muss HB, Thor AD, Berry DA, Kute T, Liu ET, Koerner F, et al. c-erbB-2 expression and response to adjuvant therapy in women with node-positive early breast cancer. *N Engl J Med*. 1994;330:1260–6.
98. Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, et al. Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med*. 2005;353:1659–72.
99. Dent S, Oyan B, Honig A, Mano M, Howell S. HER2-targeted therapy in breast cancer: a systematic review of neoadjuvant trials. *Cancer Treat Rev*. 2013;39(6):622–31.
100. Luporsi E, André F, Spyrtos F, Martin PM, Jacquemier J, Penault-Llorca F, et al. Ki-67: level of evidence and methodological considerations for its role in the clinical management of breast cancer: analytical and critical review. *Breast Cancer Res Treat*. 2012;132(3):895–915.
101. Polley MY, Leung SC, Gao D, Mastropasqua MG, Zabaglo LA, Bartlett JM, et al. An international study to increase concordance in Ki67 scoring. *Mod Pathol*. 2015;38:1–9.
102. Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomized trials. *Lancet*. 2005;365(9472):1687–717.
103. Fisher B, Dignam J, Wolmark N, DeCillis A, Emir B, Wickerham DL, et al. Tamoxifen and chemotherapy for lymph node-negative, estrogen receptor-positive breast cancer. *J Natl Cancer Inst*. 1997;89(22):1673–82.
104. Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. *Nature*. 2000;406(6797):747–52.
105. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A*. 2001;98(19):10869–74.
106. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med*. 2004;351(27):2817–26.
107. Goldstein LJ, Gray R, Badve S, Childs BH, Yoshizawa C, Rowley S, et al. Prognostic utility of the 21-gene assay in hormone receptor-positive operable breast cancer compared with classical clinicopathologic features. *J Clin Oncol*. 2008;26(25):4063–71.
108. Albain KS, Barlow WE, Shak S, Hortobagyi GN, Livingston RB, Yeh IT, et al. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomized trial. *Lancet Oncol*. 2010;11(1):55–65.
109. [http://www.nccn.org/professionals/physician\\_gls/pdf/breast.pdf](http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf). Accessed on 1 Apr 2014.
110. Gnant M, Harbeck N, Thomssen C. St. Gallen 2011: summary of the consensus discussion. *Breast Care (Basel)*. 2011;6:136–41.
111. van't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, et al. Gene expression profiling predicts clinical outcome of breast cancer. *Nature*. 2002;415(6871):530–6.
112. Ma XJ, Wang Z, Ryan PD, Isakoff SJ, Barmettler A, Fuller A, et al. A two-gene expression ratio predicts clinical outcome in breast cancer patients treated with tamoxifen. *Cancer Cell*. 2004;5(6):607–16.
113. van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, et al. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med*. 2002;347(25):1999–2009.
114. Buyse M, Loi S, van't Veer L, Viale G, Delorenzi M, Glas AM, et al. Validation and clinical utility of a 70-gene prognostic signature for women with node-negative breast cancer. *J Natl Cancer Inst*. 2006;98(17):1183–92.
115. Rutgers E, Piccart-Gebhart MJ, Bogaerts J, et al. Baseline results of the EORTC 10041/MINDACT TRIAL (Microarray In Node 0–3 positive Disease may Avoid ChemoTherapy). On behalf of the MINDACT TRANSBIG Studygroup. Elsevier. Abstract#2062. European Congress of Cancer Meeting; 2013.
116. Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T, et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol*. 2009;27(8):1160–7.
117. Chia SK, Bramwell VH, Tu D, Shepherd LE, Jiang S, Vickery T, et al. A 50-gene intrinsic subtype classifier for prognosis and prediction of benefit from adjuvant tamoxifen. *Clin Cancer Res*. 2012;18(16):4465–72.
118. Dowsett M, Sestak I, Lopez-Knowles E, Sidhu K, Dunbier AK, Cowens JW, et al. Comparison of PAM50 risk of recurrence score with oncotype DX and IHC4 for predicting risk of distant recurrence after endocrine therapy. *J Clin Oncol*. 2013;31(22):2783–90.
119. Naoi Y, Noguchi S. Multi-gene classifiers for prediction of recurrence in breast cancer patients. *Breast Cancer*. 2015. Published online.
120. Martín M, González-Rivera M, Morales S, de la Haba-Rodríguez J, González-Cortijo L, Manso L, et al. Prospective study of the impact of the Prosigna assay on adjuvant clinical decision-making in unselected patients with estrogen receptor positive, human epidermal growth factor receptor negative, node negative early-stage breast cancer. *Curr Med Res Opin*. 2015;6:1129–37.

Cagatay Arslan, M. Kadri Altundag,  
and Y. Yavuz Ozisik

---

## Abstract

Among women, breast cancer accounts for one-third of cancer cases and is the second most frequent cause of death. Improvements in treatment agents and screening procedures have increased the diagnosis of early breast cancer and survival rates. Adjuvant chemotherapy and endocrine treatment decrease the mortality of early breast cancer by approximately 50 %. However, not all early breast cancer patients benefit equally from adjuvant endocrine treatment and/or chemotherapy. Patients at high risk are classically identified based on clinicopathological factors, such as age, tumor size, histopathological grade, nodal status, hormone and HER2 receptor positivity, and menopausal status. However, for patients with early breast cancer, using these standard clinicopathological factors might not thoroughly show the individual risk of disease recurrence and the benefits from adjuvant systemic chemotherapy. Many patients with early breast cancer do not derive benefit from adjuvant systemic chemotherapy. Quality-of-life issues, acute and long-term side effects of systemic chemotherapy, and the cost of unnecessary treatments are the main factors of concern for this group of patients. Quantitative approaches for defining prognoses and for individualizing treatments are required. In recent years, molecular signatures of gene expression have been correlated with breast cancer recurrence risk. Several tests for genomic expression have been developed and validated on specimens from previous phase III studies to improve the prognostication of early breast cancer patients and/or the prediction of adjuvant systemic treatment. The most commonly used genomic expression-based tests used for prognostic information and for the prediction of chemotherapy benefits in early breast cancer are summarized below.

---

C. Arslan, MD (✉)  
Department of Medical Oncology, Izmir University  
Faculty of Medicine, Medical Park Hospital,  
Karsiyaka, Izmir, Turkey  
e-mail: [arslancagatay@yahoo.com](mailto:arslancagatay@yahoo.com)

M.K. Altundag, MD • Y.Y. Ozisik, MD  
Department of Medical Oncology, Hacettepe  
University Cancer Institute, Ankara 06100, Turkey  
e-mail: [altundag66@yahoo.com](mailto:altundag66@yahoo.com); [yozisik@hacettepe.edu.tr](mailto:yozisik@hacettepe.edu.tr)

**Keywords**

Early breast cancer • Gene arrays • Prognosis • Prediction of treatment

**Introduction**

Breast cancer accounts for one-third of cancer cases among women and is the second most frequent cause of death. [1] It consists of heterogeneous subtypes that differ in clinical presentation and disease course. Improvements in treatment agents and screening procedures have increased the diagnosis of early breast cancer and survival rates. Chemotherapy, endocrine treatment (ET), and trastuzumab comprise the main armamentarium for the adjuvant treatment of breast cancer. Adjuvant chemotherapy and ET decrease the mortality of early breast cancer by approximately 50 % [2]. However, early breast cancer patients do not benefit equally from adjuvant ET and/or chemotherapy. The recurrence risk of disease for hormone receptor-positive early breast cancer with tamoxifen after surgery is 15 % in 10 years, and the survival benefit of adjuvant chemotherapy in the same group of patients is 3–10 % [3, 4]. Patients at high risk are classically identified based on clinicopathological factors, such as age, tumor size, histopathological grade, nodal status, hormone and HER2 receptor expression, and menopausal status. However, using these standard clinicopathological factors might not thoroughly show the individual risk of disease recurrence and the benefit from adjuvant systemic chemotherapy for early breast cancer patients. Many early breast cancer patients do not benefit from adjuvant systemic chemotherapy [5]. Quality-of-life issues, acute and long-term side effects of systemic chemotherapy, and cost of the unnecessary treatments are the main concerns for this group of patients. During the past two decades, the level of knowledge regarding the molecular pathways and underlying genetic changes in breast cancer has improved. However, treatment decisions still often rely on classical histopathological and immunohistochemical techniques. Numerous biomarkers have been

studied to define the residual risk of recurrence, but none of them have been recommended for routine use [6–11].

Quantitative approaches for defining prognosis and individualization of treatments are thus required. In recent years, molecular signatures of gene expression have been correlated with the risk of breast cancer recurrence [12–15]. Questions of reproducibility and the need for fresh or fresh-frozen tissue have limited their clinical application. A number of genomic expression tests were developed regarding this unmet medical need. Several genomic expression tests have been developed and validated on specimens of previous phase III studies to improve the prognostication of early breast cancer patients and/or the prediction of the utility of adjuvant systemic treatment (Table 11.1) [16]. In this chapter, using satisfactory data from the literature, we review the most commonly used genomic expression-based tests for predicting the prognosis of and chemotherapy benefit for early breast cancer.

**Gene Array Tests****Oncotype DX**

The Oncotype DX (Genomic Health, Inc., Redwood City, CA) uses real-time reverse-transcriptase chain reaction (RT-PCR) to quantify the expression levels of 21 genes in formalin-fixed, paraffin-embedded (FFPE) breast cancer tissue samples [17]. The test, which is regulated by Clinical Laboratory Improvement Amendments (CLIA) and the College of American Pathologists (CAP), is performed in a central laboratory in the USA. This assay uses a calculation model to generate a risk score ranging between 0 and 100 based on 5 reference and 16 cancer-related genes. The reference genes,

**Table 11.1** Summary of selected approved genomic tests for breast cancer [16]

	Oncotype DX™	MammaPrint™	Prosigna™
Generic name	21-gene signature	70-gene signature	PAM50
Company	Genomic Health	Agendia	NanoString
Method	qRT-PCR	Microarray	qRT-PCR nCounter™ (for Prosigna)
Target genes	16 genes 5 controls	70 genes	50 genes 5 controls
Specimen	FFPE	Fresh/Frozen FFPE (2011)	FFPE
Analyses	Central Lab.	Central Lab.	Localized
Prognostic index	Risk score (RS)	MammaPrint index	Risk of recurrence (ROR)
Indication	Prognostic Predictive	Prognostic Predictive	Prognostic Intrinsic subtype classifier
Population studied	N0-1, ER positive	N0-1, <61 years old	N0-1, ER positive, postmenopausal
Evidence	NSABP B14 NSABP B20 ECOG 2197 TransATAC NSABP B28 SWOG 8814 JBCRG Northern California	TRANSBIG	NCIC MA.12 NCIC MA.5 ABCSG-8 TransATAC

*GAPDH*, *ACTB* ( $\beta$ -actin), *RPLPO*, *GUS*, and *TFRC*, are used for normalization. The cancer-related genes used in the assay include genes related to proliferation (*Ki-67*, *Survivin*, *MYBL2*, *CCNB1* [cyclin B1], and *STK15*), invasion (*CTSL2* [cathepsin L2] and *MMP-11* [stromolysin 3]), and hormone receptors (*ER*, *PR*, *BCL2*, and *SCUBE2*) as well as *HER2* (*GRB7* and *HER2*), *GSTM1*, *BAG1*, and *CD68*.

A multistep approach was adopted for developing the assay for expression levels of tumor-related genes by the Genomic Health researchers. Routinely used tumor blocks were used for this purpose and to validate the assay. High-throughput RT-PCR was used to quantify gene expression levels in FFPE tumor tissue sections [18]. In the second phase, they chose 250 candidate genes from the published literature, genomic databases, and experiments based on DNA arrays performed on fresh-frozen tissue [12–14, 19]. In the third phase, data from 447 breast cancer patients from 3 different studies, including patients from the tamoxifen-only arm of the NSABP B-20 trial, were used to test the correla-

tion of 250 candidate genes with the recurrence of breast cancer [20–22]. In the fourth phase, a panel of 16 cancer-related genes and 5 reference genes were selected from the results of three studies based on the strength of their performance in the previous studies and the consistency of the primer and probe performance in the assay. An algorithm was designed based on the expression of these 21 genes for computing a recurrence score (RS) for each tumor sample [17]. The possible RSs ranged between 0 and 100, where a higher recurrence score indicated a higher likelihood of recurrence. The RS was derived from the reference-normalized expression measurements of the 16 cancer-related genes. Reproducibility within and between blocks was also assessed. Based on their RS, patients were divided into 3 risk categories, including low (<18), intermediate (18–30), and high ( $\geq 31$ ). The RS was prognostic for estrogen receptor (ER)-positive early breast cancer patients with positive (1–3 lymph nodes involved) and negative lymph node involvement who were treated with tamoxifen. A low RS predicted no likelihood of recurrence in 10 years and



little benefit from chemotherapy. Adjuvant chemotherapy showed benefits in high-RS patients but not in patients with a low RS.

Oncotype DX was tested in a community-based population from Northern California [23]. The 21-gene assay was prognostic in ER-positive patients with and without tamoxifen treatment ( $p=0.003$  and  $p=0.03$ , respectively). There were 220 patients and 570 controls in this study. Archived tumor tissues were tested. Nearly 50 % of the patients were in the low-risk group. The risk for death from breast cancer within 10 years in tamoxifen-treated patients was 2.8 (95 % confidence interval (CI): 1.7–3.9), 10.7 (95 % CI: 6.3–14.9), and 15.5 % (95 % CI: 7.6–22.8) in the low-, intermediate-, and high-risk patients, respectively. The risk of death within 10 years in patients with no tamoxifen treatment was 6.2 (95 % CI: 4.5–7.9), 17.8 (95 % CI: 11.8–23.3), and 19.9 (95 % CI: 14.2–25.2) in low-, intermediate-, and high-risk patients, respectively.

The Oncotype DX assay was further tested in hormone receptor-positive early breast cancer patients from Japan [24]. All patients were treated with tamoxifen. Among the total patients, 280 patients had tumor tissues that were adequate for the assay. Of these patients, 48 % were in the low-risk group, 20 % were in the intermediate-risk group, and 33 % were in the high-risk group. Distant recurrence risks in 10 years were 3.3 (95 % CI: 1.1–10), 0, and 24.8 % (95 % CI: 15.7–37.8) in low-, intermediate-, and high-risk lymph node-negative patients, respectively. Differences between the low-risk and high-risk groups for distant recurrence were significant (log rank  $p<0.001$ ). There was also a significant difference in overall survival (OS) between the low-risk and high-risk groups ( $p=0.008$ ).

## Pivotal Trials of Oncotype DX

### NSABP B14 Trial

The data obtained from the large, multicenter NSABP B14 trial and the FFPE tumor tissues were used to validate this 21-gene RT-PCR assay for RS detection in early stage, node-negative, ER-positive breast cancer patients,

who had been treated with tamoxifen [17]. The RS was calculated as low, intermediate, or high for each patient, as previously defined. The data from the prospective NSABP B14 trial were used for validation. Cutoff points were determined based on the results of the NSABP B14 study. According to the study results, the rates of recurrence in 10 years were 6.8 (95 % CI: 4–9.6), 14.3 (95 % CI: 8.3–20.3), and 30.5 % (95 % CI: 23.6–38.4) for the low-, intermediate-, and high-risk groups, respectively. The risk of recurrence in the high-risk group was similar to that for the lymph node-positive patients [25]. In this study, 51 % of the patients were in the low-risk group, and 27 % of the patients were in the high-risk group. Age and tumor size are standard factors used for predicting recurrence. However, when RS was added to the multivariate Cox model, recurrence could no longer be predicted based on age or tumor size. Moreover, all patients with tumors smaller than 1 cm ( $N=109$ ) were not in the low-risk group. Forty-four of these patients were in the intermediate- or high-risk groups, which means a risk of 15–20 % for recurrence in 10 years. A subgroup of the patients with low-grade tumors also showed high RS and high rates of recurrence in this study. Additionally, concordance of pathologists was moderate for poorly differentiated tumors and low for the well- and intermediate-differentiated tumors. HER2 amplification was detected in 55 of 668 patients (8 %). The 10-year recurrence was 75 % (95 % CI: 63.2–86.9) in HER2-amplified and 86 % (95 % CI: 83.1–88.9) in HER2-nonamplified breast tumors ( $p=0.08$ ). In the Cox model including RS and classical factors (estrogen, progesterone receptors, HER2 DNA amplification), any RS was a significant predictor of distant recurrence. The RS provided significant predictive power independent of age or tumor size ( $p<0.001$ ). The RS was also prognostic ( $p<0.001$ ) and could be used as a continuous function to predict the recurrence in each patient [17]. It is important to note that all patients were treated with tamoxifen; thus, outcomes must be evaluated considering the effects of both tamoxifen and the natural disease course.

### TransATAC Trial

ATAC is a phase III trial that included 9366 postmenopausal early breast cancer patients, including both ER-positive and ER-negative patients. This study compared 5 years of adjuvant treatment with tamoxifen alone, anastrozole alone, and tamoxifen in combination with anastrozole [26]. In the translational arm of the ATAC study (TransATAC), the risk of recurrence was evaluated using the Oncotype DX assay in axillary node-negative or node-positive, hormone receptor-positive postmenopausal breast cancer patients who were treated with tamoxifen and anastrozole. RNA was extracted from 1372 tumor blocks from patients in the monotherapy arms of this study. Available scores were obtained from 1231 patients (node positive,  $N=306$ ; node negative,  $N=872$ ; node status unknown,  $N=52$ ). Multivariate analysis showed that RS was significantly associated with time to disease recurrence ( $p<0.001$  and  $p=0.002$  for node-negative and node-positive patients, respectively). There was a poor correlation between RS and “Adjuvant! Online” for estimating prognosis ( $p<0.001$ ). In node-negative patients, the disease recurrence rate within 9 years was 4, 12, and 25 % for the low-, intermediate-, and high-risk groups, respectively. In node-positive patients, the 9-year disease recurrence rates were 17, 28, and 49 % for the low-, intermediate-, and high-risk groups, respectively. The hazard ratio (HR) for disease recurrence was 2.7 for high-RS and 1.8 for low-RS lymph node-positive patients. Similar results were obtained for OS. For any RS, the risk of distant recurrence was higher in node-positive than node-negative patients and in patients with 4 or more positive nodes than in patients with 1–3 positive nodes. Prognostic value was similar in the tamoxifen and anastrozole groups. In the original study, anastrozole showed a 16 % risk reduction for distant recurrence compared with tamoxifen [27]. However, in this study, HRs for distant recurrence were similar in the tamoxifen and anastrozole treatment arms, and RS did not interact with any treatment arm. Relative risk reductions for anastrozole compared with tamoxifen were similar in all RS groups. A higher risk reduction with anastrozole in patients with high

RS might be expected. However, the number of cases was too small to allow for such an analysis.

### Studies of Prediction for Chemotherapy

#### NSABP B20 Trial

In the original NSABP B20 trial, there were 2363 ER-positive, axillary lymph node-negative early breast cancer patients [28]. The aim was to examine the benefit of CMF (cyclophosphamide, methotrexate, 5-fluorouracil) or MF (methotrexate, 5-fluorouracil) chemotherapy followed by 5 years of tamoxifen treatment. The Oncotype DX assay was studied in these tumor blocks, and prospective clinical outcomes were investigated in that study group to examine whether the Oncotype DX assay could predict the benefits of chemotherapy. Tumor blocks with sufficient tumor tissue for the RS assay were obtained in 670 patients and adequate samples were obtained from 651 tumor blocks. Of the total patients, 227 (29.5 %) were treated with tamoxifen, and 424 (27.7 %) were treated with chemotherapy plus tamoxifen. Among the 651 assessable patients, the proportions of patients without distant recurrence in 10 years were 92.2 % in the chemotherapy plus tamoxifen group and 87.8 % in the tamoxifen group. Disease recurrence (locoregional or distant) was observed in 90.1 % of the chemotherapy plus tamoxifen patients and in 83.5 % of the tamoxifen patients. The 10-year survival estimate was 89.5 % in patients treated with chemotherapy plus tamoxifen and 86.4 % in patients treated with only tamoxifen. There were 353 (54.2 %) patients with low RS, 134 (20.6 %) with intermediate RS, and 164 (25.2 %) with high RS.

This study showed that the benefit of chemotherapy was not equivalent across all ER-positive and axillary lymph node-negative early breast cancer patients in the NSABP B20 study. The Oncotype DX assay was shown to predict the chemotherapy (CMF or MF) benefit in this group of patients. The magnitude of the benefit of chemotherapy for distant recurrence was greater for the high-RS group than for the intermediate- and low-RS groups. A Kaplan-Meier estimate indicated

that 10 years of freedom from the disease recurrence improved from 60 to 88 % in patients in the high-RS group. There was no demonstrable risk reduction with chemotherapy regarding the 10-year disease recurrence rates in the low-RS group (relative risk, 1.31; CI, 0.46–3.78). A significant risk reduction (27.6 % reduction in absolute risk) was shown with chemotherapy in the high-RS group (relative risk, 0.26; CI, 0.13–0.53). The benefit of chemotherapy was not clear in the intermediate-RS group (relative risk, 0.61; CI, 0.24–1.59). In a multivariate analysis of Cox models containing chemotherapy treatment and RS, the interaction between chemotherapy treatment and RS was significant ( $p=0.038$ ).

### NSABP B28 Trial

The current standard adjuvant treatment of ER-positive, axillary lymph node-positive breast cancer in pre- or postmenopausal patients is ET plus chemotherapy (ET+CT) [2]. Nevertheless, exploratory analyses show that breast cancer patients with high levels of ER positivity and a lack of HER2 overexpression may not derive substantial benefit, even if those patients show positive axillary nodes [29, 30].

The original phase III NSABP B28 trial included 3060 pre- or postmenopausal ER-positive ( $N=2019$ ) and ER-negative/borderline ( $N=1041$ ) axillary lymph node-positive, early breast cancer patients [31].

The treatment groups were tamoxifen plus AC (doxorubicin plus cyclophosphamide) and AC plus paclitaxel. In this trial, 2687 patients received concurrent ET. RS analysis was performed in 1065 tumor blocks of ER-positive patients treated with endocrine therapy [32]. In the univariate analysis, RS predicted disease-free survival (DFS), disease recurrence-free interval, and OS ( $p<0.001$  for each). In a multivariate analysis, RS was identified as an independent prognostic factor for DFS, disease recurrence-free interval, and OS ( $p<0.001$ ). DFS was 76 (95 % CI: 71–80), 57 (95 % CI: 52–62), and 48 % (95 % CI: 42–53) for the low-, intermediate-, and high-RS groups, respectively. The disease recurrence-free interval was 81 (95 % CI: 76–85), 65 (95 % CI: 60–70), and 56 % (95 % CI: 50–61) for the low-, interme-

diate-, and high-RS groups, respectively. OS was 90 % (95 % CI: 86–93), 75 (95 % CI: 70–79), and 63 % (95 % CI: 57–68) for the low-, intermediate-, and high-RS groups, respectively. In patients with low RS, adding paclitaxel to anthracycline-based adjuvant treatment did not provide any additional benefit to the final outcome. Based on these results, aggressive chemotherapy may not be warranted for patients with low RSs [32].

A subgroup of ER-positive breast cancer patients shows a low risk of disease recurrence even with positive axillary nodes. These patients are unlikely to benefit from adjuvant chemotherapy. Oncotype DX RS was also shown to be prognostic in axillary lymph node-positive, concordant with axillary lymph node-negative tamoxifen-treated breast cancer patients. Oncotype DX RS may also predict patients who will not benefit from the addition of chemotherapy (i.e., patients with low RS).

### SWOG 8814 Trial

Recent data from the SWOG 8814 trial showed Oncotype DX to be prognostic and predictive in ER-positive, axillary lymph node-positive breast cancer patients [33]. The SWOG 8814 study, a parent trial, was a phase III trial and showed that adjuvant CAF (cyclophosphamide, doxorubicin, 5-fluorouracil) chemotherapy plus tamoxifen was superior to tamoxifen alone for DFS and OS in postmenopausal, estrogen and/or progesterone receptor (ER/PR)-positive, axillary lymph node-positive breast cancer patients. The Oncotype DX assay was applied to specimens from a tumor bank for RS analysis. That study was performed by Genomic Health, Inc., and the investigators were blinded to patients' clinical data and outcome. Clinical data and outcome were combined with the Oncotype DX RS results after all assays were completed. Tumor samples were available for 664 of 1477 patients (45 %). The parent trial included three arms: tamoxifen only, concomitant CAF and tamoxifen, and sequential CAF and tamoxifen. In this translational study, the concomitant CAF and tamoxifen arm was omitted due to inferior efficacy. Sufficient RNA was obtained from 149 patients in the tamoxifen arm and 219 patients in the CAF plus tamoxifen arm (367 total

patients) for RT-PCR analysis. The 21-gene RS was prognostic in tamoxifen-treated, ER-positive, axillary lymph node-positive patients and predictive for adjuvant chemotherapy with CAF regimen in patients with high RS [33]. The RS was prognostic for DFS in tamoxifen-treated patients ( $p=0.006$ ). There was a significant benefit of chemotherapy in the high-RS group (log rank  $p=0.03$ ; HR, 0.59; 95 % CI, 0.35–1.01), but no benefit was detected in patients with low RS (log rank  $p=0.97$ ; HR, 1.02; 95 % CI, 0.54–1.93) regarding DFS. A low RS identifies patients who may not benefit from adjuvant chemotherapy despite positive nodes. The benefit of adjuvant chemotherapy in the high-RS group was independent from the number of positive nodes. The benefit of chemotherapy on DFS was significant for the first 5 years of follow-up. However, there was no additional prediction of chemotherapy benefit for DFS beyond 5 years, despite the continued presence of the cumulative benefit after 10 years. Similar results were obtained for the prognostic value of the 21-gene RS. Ten-year OS estimates were 77, 68, and 51 % for patients with high, intermediate, and low RS, respectively. Breast cancer-specific survival (BCSS) and OS were significantly better in patients with high RS but not low and intermediate RS when treated with CAF and tamoxifen compared to tamoxifen only. In the exploratory analysis after adjustment for classical risk factors, including age, histopathological grade, race, tumor size, PR status, and HER2 status, both treatment and RS remained significant. The limited number of samples obtained from patients included in the original study was an important caveat of this study. The probability of a chemotherapy effect in low-RS patients cannot be ruled out completely due to the small sample size and broad CI ranges in this study. This study challenged the standard adjuvant chemotherapy treatment of patients with axillary lymph node-positive, ER-positive breast cancer.

### ECOG 2197 Trial

In the original ECOG 2197 trial, there were 2185 pre- or postmenopausal ER-positive and lymph node-negative and node-positive (1–3 nodes) breast cancer patients treated with doxorubicin-

containing chemotherapy or docetaxel plus ET [34]. The Oncotype DX assay was tested in the doxorubicin-containing chemotherapy arm ( $N=465$ ) and was a significant prognostic marker of disease recurrence in this study group. There was a significant correlation between RS and 5-year disease recurrence rates in both lymph node-positive and node-negative patient groups ( $p<0.001$ ). Of the lymph node-positive patients, 46 % were in the low-RS group. Patients with zero or one positive node had a 5-year disease recurrence rate under 3 %, and patients with two or three positive nodes had a 5-year disease recurrence rate of 8 %. However, good outcomes may be attributed to low RS, chemotherapy, or both. Ten-year follow-up analysis also showed that RS was still a prognostic marker for disease recurrence for lymph node-positive and lymph node-negative breast cancer patients [35].

Based on all of these studies, Oncotype DX has shown a high level of utility for estimating the risk of distant recurrence in 10 years and the benefit of chemotherapy in ER-positive, HER2-negative early breast cancer patients with up to three metastatic lymph nodes. The Oncotype DX assay has been adopted by the international guidelines of ESMO, ASCO, NCCN, and St. Gallen [36–39].

### Impact on Treatment Decision

The Oncotype DX assay changed treatment decisions made on classical clinicopathological factors in several studies. Its impact on decision-making seems to increase with the results of the prospective trials. In a survey study in the USA, oncologists were asked about the treatment recommendation for their most recent ER-positive, lymph node-positive breast cancer patient after getting the Oncotype DX assay result [40]. The vast majority of the patients had one to three positive lymph nodes (92.5 %), and most (96.5 %) had a tumor size smaller than 5 cm. Of the 160 physicians who responded (16 % of the original sample), 86 % made decisions before obtaining the RS. However, 51 % of those oncologists changed their decision after receiving the RS result. Treatment decisions were changed to ET alone from ET+CT in one-third of those cases. Chemotherapy was eliminated in

49 % of the intermediate RS group and in 21 % of the low-RS group of patients. Additional chemotherapy decisions were made in 9 % of the cases.

In a study from Israel, 951 patients with one to three positive nodes received ET with or without chemotherapy [41]. Treatment decisions for 282 patients were made according to Oncotype DX, and there were 669 controls. In the Oncotype DX group, chemotherapy was given to all patients with high RS (100 %), 37 % of patients with intermediate RS, and 7 % of patients with low RS. Chemotherapy was given to 24.5 % of all Oncotype DX patients and 70.4 % of controls ( $p < 0.001$ ). However, the patients' clinicopathological features were not balanced between the groups, and patients in the control group had bigger tumor sizes, higher grade, and more positive lymph nodes. After adjustment according to these factors, Oncotype DX was associated with a 65 % decrease in chemotherapy usage.

A study of 50 lymph node-positive breast cancer patients from Australia showed a 26 % change in the treatment decision with Oncotype DX, and the majority of the treatment decisions were to omit chemotherapy [42]. A study of 42 axillary lymph node-positive ER-positive breast cancer patients (22 with macrometastasis) from Spain showed that 73 % of the patients had low RS [43], and the recommendation of chemotherapy decreased from 55 to 17 % with the use of Oncotype DX ( $p = 0.021$ ).

In a recent meta-analysis of 8 studies (1437 patients) on the impact of Oncotype DX on treatment decisions, the adjuvant therapy recommendation changed in 33.4 % of patients due to Oncotype DX RS versus the decision recommended based on clinicopathological factors [44]. After Oncotype DX, the adjuvant chemotherapy recommendation was 83.4 % in patients with high RS, 37.4 % with intermediate RS, and 5.8 % with low RS. The overall chemotherapy recommendation was 28.2 % after using the Oncotype DX assay.

### Prospective Clinical Trials

The phase III SWOG 1007 (RxPONDER, NCT01272037) trial is designed to determine the effect of chemotherapy with adjuvant endocrine therapy in patients with ER-positive, 1–3 axillary

lymph node-positive breast cancer and low or intermediate RS (less than or equal to 25) [45]. This study will provide important information about the patients in whom chemotherapy can be omitted in the low- and intermediate-RS groups. It will also address issues of quality of life and long-term side effects such as premature menopause and weight gain.

The prospective TAILORx trial was designed mainly to investigate the benefits of chemotherapy in the intermediate RS group (scores of 11–25) of ER-positive, axillary lymph node-negative early breast cancer patients [46]. The Oncotype DX assay was applied prior to treatment, and patients were divided into three RS groups: low (RS <11), intermediate (RS 11–25), and high (RS >25). Patients in the low-RS group were treated with adjuvant ET, and those in the high-RS group were treated with ET+CT. Enrollment was completed in 2010, and a total of 11,000 patients were recruited in the study. The results will be compiled in 2015.

WSG PLAN B is a phase III trial run by the West German Study Group. In this study, two different chemotherapy regimens were compared in 2549 patients with early breast cancer [47]. Oncotype DX, uPA/PAI-1, Ki67, and histopathological grade were used for comparison of chemotherapy regimens in patients with ER-positive axillary lymph node-positive or high-risk node-negative breast cancer.

WSG ADAPT is another phase II/III trial in pre- and postmenopausal early breast cancer patients [47]. It aims to individualize the adjuvant treatment decision in early breast cancer patients and is using the Oncotype DX assay with conventional prognostic factors (nodal status). Dynamic changes in proliferation rates and apoptosis were checked after a short course of treatment. The aim was to establish early predictive surrogate molecular markers for outcome by assessing the response to a repeated biopsy after 3 weeks of induction treatment.

### MammaPrint

MammaPrint is based on DNA microarray technology. Using gene-expression profiling, the



Netherlands Cancer Institute™ and its spinoff company Agendia™ developed a 70-gene prognostic signature called MammaPrint™ for axillary lymph node-negative early breast cancer patients. These 70 genes are involved in the cell cycle, invasion, proliferation, angiogenesis, metastasis, and signal transduction. None of these genes are assessed by Oncotype DX. The 70-gene signature was developed for a dichotomous risk classification as low or high risk of disease recurrence in 5 years in a cohort of node-negative breast cancer patients, who were not treated systemically [14]. This gene signature defined the low- and high-risk groups for 5-year disease recurrence [14]. The low-risk group was identified as having a 13 % risk of distant metastasis in 10 years, and the high-risk group was identified as having a 56 % risk of distant metastasis in 10 years without adjuvant treatment. The first validation study was performed with the samples from the tumor bank of the Netherlands Cancer Institute. A total of 295 tumor samples were obtained from breast cancer patients (ER negative and ER positive, lymph node negative and node positive, untreated, treated with chemotherapy and/or endocrine therapy) [15]. Metastasis-free survival and OS were higher in the low-risk group than in the high-risk group in lymph node-negative and node-positive patients and all patients were evaluated using the MammaPrint assay. Of the patients with low-risk profiles, 85 % were disease recurrence-free, and 50.6 % of the patients with high-risk profiles were disease recurrence-free in 10 years. In the multivariate analysis, independent prognostic markers for disease recurrence were high-risk profile with MammaPrint™, tumor size, and absence of chemotherapy. This 70-gene expression profile was developed on microarrays containing 25,000 60-mer oligonucleotides that are not designed for routine clinical practice. This 70-gene prognosis profile, which was translated to a customized microarray (MammaPrint™), contains a reduced set of 1900 probes suitable for high-throughput processing [48]. It allows the use of less RNA and a short processing time of 5 days. To validate its prognostic value, the RNA of 162 patients from two previous studies was used

for hybridization with this custom array. Custom microarray results were compared with the original analysis, and they showed an extremely high correlation of prognosis prediction ( $p < 0.0001$ ). This first study showed that this small, custom microarray can be a reliable diagnostic tool for predicting the outcome of disease in breast cancer patients. MammaPrint™ was intended for use in women under 61 years of age with stage I and II, ER-positive or ER-negative early breast cancer. It was approved by the US FDA in 2007 for determining the risk of distant recurrence at 5 and 10 years but not for predicting the benefit of chemotherapy. Based on the evidence supporting the prognostic value of MammaPrint™, it has been incorporated into the ESMO and St. Gallen guidelines as a prognostic tool [37, 38]. The test was initially developed and validated for fresh or fresh-frozen tumor tissue [15, 49]. A recent study of 211 matched samples showed 91.5 % consistency between using FFPE and using fresh or frozen specimens [50].

Another MammaPrint™ validation study was performed by TRANSBIG researchers in 307 patients with early, axillary lymph node-negative breast cancer, with a median follow-up of 13.6 years, in 5 different European countries [51]. MammaPrint™ results were compared with other clinicopathological risk classification systems. Patients were defined as low risk if their 5-year probability of distant metastasis-free survival was above 90 %. The clinicopathological low-risk group was defined as a 10-year OS probability greater than 88 % (for ER-positive patients) or 92 % (for ER-negative patients). MammaPrint was identified as an independent prognostic marker for distant metastasis-free survival and OS. In the univariate analysis of the 70-gene signature in high- versus low-risk patients, the HR for time to distant metastasis was 2.32 (95 % CI, 1.35–4;  $p = 0.002$ ), HR for OS was 2.79 (95 % CI, 1.6–4.87;  $p < 0.001$ ), and the HR for DFS was 1.56 (95 % CI, 1.04–2.16;  $p = 0.032$ ). MammaPrint was a more powerful prognostic tool for distant metastasis-free survival and OS in comparison with clinicopathological factors defined by Nottingham Prognostic Index, St. Gallen Criteria or Adjuvant! Online.

When adjusted for the clinical risk groups, the 70-gene risk score results showed HRs of 2.13 (95 % CI: 1.19–3.82), 2.63 (95 % CI: 1.45–4.79), and 1.36 (95 % CI: 0.91–2.03) for time to distant metastasis, OS, and DFS, respectively.

### Data for Patients with Positive Lymph Nodes

The 70-gene assay result was also prognostic in lymph node-positive breast cancer patients and predicted the outcome better than the other classical clinicopathological factors. A validation study was conducted with 241 patients from two European institutes, with one to three positive lymph nodes and operable T1–T3 early breast cancer. Adjuvant treatment decisions were made according to national guidelines from the two different countries (the Netherlands and Italy). MammaPrint™ predicted that patients in the low-risk group would have 91 and 96 % distant metastasis-free survival and BCSS, respectively, in 10 years [52]. For the poor prognostic group, the 10-year distant metastasis-free survival and BCSS were 76 and 76 %. The 70-gene risk score was superior to other clinicopathological factors for predicting BCSS. In the multivariate analysis, the 70-gene risk score HR was 7.19 (95 % CI: 1.8–28.43;  $p=0.005$ ) for BCSS.

The MammaPrint™ assay was studied using frozen tumors from 173 N2 (with 4–9 axillary lymph node metastatic) breast cancer patients from the Netherlands and Italy [53]. Seventy patients (40 %) were classified as genomic high risk, and 103 (60 %) were classified as genomic low risk. Patients in the genomic high-risk group were more often grade 3 (60 %), hormone receptor negative and HER2 positive (25 %). The 5-year OS was 97 and 76 % in the low- and high-risk groups, respectively ( $p<0.01$ ). Distant metastasis-free survival in 5 years was 87 % for low-risk patients and 63 % for high-risk patients ( $p<0.01$ ). In the luminal A subgroup, the 70-gene assay result was the only independent risk factor for distant metastasis and breast cancer-specific death in breast cancer patients with 4–9 positive lymph nodes. MammaPrint can be integrated into the selection of treatment strategy in this group of patients.


### Data on Adjuvant Treatment Decisions

The performance of the 70-gene signature was assessed in a prospective observational community-based study that included 427 early breast cancer (cT1-3N0M0) patients [54]. Adjuvant systemic treatment decisions were given considering the 70-gene signature, the Dutch CBO 2004 guidelines, and the preferences of the physicians and patients. The median follow-up duration was 62 months. In the 70-gene signature group, 15 % (33/219) of low-risk patients and 81 % (169/208) of high-risk patients received adjuvant chemotherapy. The 5-year probability of disease recurrence-free survival according to the 70-gene score was 97.0 % for low-risk and 91.7 % for high-risk patients. The 5-year probabilities of disease recurrence-free survival for the Adjuvant! Online low- and high-risk groups were 96.7 and 93.4 %, respectively. There were 124 patients in the 70-gene signature low-risk and Adjuvant! Online high-risk groups. Of these patients, 94 (76 %) did not receive adjuvant chemotherapy, and the 5-year disease recurrence-free interval was 93.4 %. The adjuvant chemotherapy decision would decrease by 32 % (94/295) if the 70-gene signature had been used in the Adjuvant! Online high-risk group of patients. The prognostic value of the 70-gene signature was again shown for the 5-year probability of disease recurrence-free survival, showing that in the low-risk 70-gene signature group, omission of chemotherapy did not compromise outcome.

The 70-gene assay was studied for the predictive value of chemotherapy in 541 patients whose tumor samples were obtained from 7 previously reported studies conducted between 1984 and 2006 with known adjuvant treatment data [55]. There were 315 patients in the ET group and 226 patients in the ET+CT group. According to the 70-gene assay, 252 (47 %) patients were in the low-risk group, and 289 (53 %) patients were in the high-risk group. BCSS at 5 years was 97 % in patients treated with ET and 99 % in patients treated with ET+CT in the low-risk group (HR, 0.58; 95 % CI, 0.07–4.98,  $p=0.20$ ). In the high-risk group, BCSS was 81 % in patients treated with ET and

94 % in patients treated with ET+CT (HR, 0.21; 95 % CI 0.07–0.59,  $p < 0.01$ ). Distant disease-free survival in the low-risk group was 93 % in patients treated with ET and 93 % in patients treated with ET+CT (HR, 0.26; 95 % CI, 0.03–2.02,  $p = 0.2$ ). In the high-risk group, distant disease-free survival was 76 % and 88 % in the patients treated with ET and ET+CT, respectively (HR, 0.35; 95 % CI, 0.17–0.71,

$p < 0.01$ ). There was a significant benefit of adding CT to ET for the group identified as high risk according to the 70-gene signature (Fig. 11.1). This benefit was not significant in patients with low risk according to the 70-gene signature. High-risk ER-positive breast cancer patients may be treated with more intensive treatment (i.e., chemotherapy) strategies in addition to adjuvant endocrine therapies.



**Fig. 11.1** Panel (a), five-year breast cancer-specific survival by treatment within the 70-gene signature groups (70-gene low risk on the left, high risk on the right). Panel (b), five-year distant disease-free survival by treatment within the 70-gene signature groups (70-gene; low risk on the left, high risk on the right). Abbreviations: BCSS

breast cancer-specific survival, DDFS distant disease-free survival,  $n$  number, ET endocrine therapy, ET+CT endocrine+chemotherapy, HR univariate hazard ratio (Reproduced from Reference [55] with kind permission of Springer Science+Business Media)

MammaPrint can affect the adjuvant chemotherapy recommendation. Using the MammaPrint™ assay may decrease the variability of adjuvant treatment decisions. A cohort of 194 patients from 4 different countries in Europe was used to measure the impact of MammaPrint™ on adjuvant treatment decisions [56]. Patients' clinicopathological data were sent to different multidisciplinary teams with and without the MammaPrint™ assay result, and adjuvant treatment decisions were provided for each patient. Treatment decisions in ER-positive and HER2-negative patients were changed in 37, 24, 28, and 35 % of cases by the Dutch, Belgian, Italian, and Spanish teams, respectively. MammaPrint™ increased the interinstitutional agreement in treatment advice (i.e., whether to utilize chemotherapy) from 51 to 75 %.

### Prospective Trials

The MINDACT (Microarray In Node-negative and 1–3 node-positive Disease may Avoid ChemoTherapy) trial was designed to investigate the clinical utility of MammaPrint™ (70-gene profile) with clinicopathological criteria for the selection of ER-positive early breast cancer patients with one to three positive nodes for adjuvant chemotherapy. The clinicopathological risk definition was made using a modified version of Adjuvant! Online. Patients were also defined as low or high risk by MammaPrint™. The MINDACT trial hypothesizes that using the MammaPrint™ assay will outperform the clinicopathological classification through the better classification of patients as high or low risk and by reducing chemotherapy usage by 10–20 % without impairing the outcome. Patients who were defined as low risk by both Adjuvant! Online and MammaPrint™ were given ET, and adjuvant ET+CT were given to patients characterized as high risk by both Adjuvant! Online and MammaPrint™. Patients were randomized to the ET or chemotherapy arms if the risk groups differed between Adjuvant! Online and MammaPrint™. The study was designed to address whether a non-anthracycline-containing regimen (docetaxel plus capecitabine) may be used instead of an anthracycline-containing regimen. There were

two arms for ET: letrozole only and tamoxifen followed by letrozole.

The pilot phase of the study included 800 patients [57]. Of those patients, 386 (48 %) were in the low-risk group based on both Adjuvant! Online and MammaPrint, and 198 (24.8 %) patients were in the high-risk group based on both Adjuvant! Online and MammaPrint™. There were 216 (27 %) discordant patients, of whom 75 (9.4 %) were in the low-risk group from Adjuvant! Online and the high-risk group from MammaPrint™ and 141 were in the high-risk group from Adjuvant! Online and the low-risk group from MammaPrint™. There was an 8.25 % difference (95 % CI: 4.7–11.8 %;  $p < 0.001$ ) between the high-risk groups defined by Adjuvant! Online (42 %) and MammaPrint™ (34 %). There was a high consistency among the treatments in both groups (>92 %).

### Prosigna

The NanoString Prosigna assay uses the expression of 50 target genes and eight constitutively expressed normalization genes. The test can be performed using qRT-PCR, but Prosigna™ relies on the NanoString nCounter Analysis System, which delivers direct, multiplexed measurement of gene expression. The assay is highly sensitive and precise. It uses 250 ng of RNA from FFPE tumor tissue. Assay controls are included along with test samples, and the process meets the pre-defined quality criteria. The PAM50 test was generated as a second-generation multigene expression assay and was developed to define the intrinsic breast cancer subtypes as luminal A/B, HER2 enriched, and basal-like from FFPE tissue [58]. PAM50 is more effective than classical immunohistochemical methods and clinicopathological factors for subtyping breast cancer. It uses a set of 50 genes and 5 control genes for analysis. In addition to classifying the tumor's intrinsic subtype, PAM50 gives a numeric score for the patient's distant recurrence probability by calculating the molecular subtype correlations, a subset of proliferation genes and pathological tumor size. The PAM50 test was adopted for performance by nCounter

Analysis in a local molecular pathology lab (Prosigna™ Breast Cancer Gene Signature Assay, NanoString Technologies, Seattle). Training is required for the operators to demonstrate proficiency in studying this locally used test. The embedded software automatically applies all of the quality thresholds to the data. A clinically validated algorithm is used for expressing the risk of recurrence and the intrinsic subtype, both of which are prognostic indicators of the risk of disease recurrence for breast cancer. This test works with either frozen or FFPE samples and uses multiplexed gene-specific fluorescently labeled probe pairs to measure gene expression.

PAM50 reflects the underlying biology associated with the ER and HER2 pathways, and it also includes proliferation genes and markers of basal phenotype. Luminal A and B breast cancers are the most frequent subtypes. Luminal A tumors are characterized by lower expression levels of proliferation genes and ERBB2 and by lower recurrence rates compared with luminal B tumors, and these characteristics can be shown by the PAM50 risk of recurrence score.

RNA is extracted from tumor tissue, and the samples are hybridized without reverse transcription or amplification for both capture and reporter probes for the measured genes and assay controls. After hybridization, the target-probe complexes are processed on the nCounter Analysis System. A minimum threshold of expression for normalization genes must be met by the test sample data to ensure that the assay signal is high enough to produce precise results.

The test was prognostic in untreated (no systemic treatment) or tamoxifen-treated early breast cancer patients [58, 59]. It was validated in 786 stage I and II ER- and/or PR-positive postmenopausal patients (in this validation study, only 40 of 789 patients were premenopausal) and provided accurate information for subtyping and prognosis by risk of recurrence [59]. This risk of recurrence score gave the estimated 10-year recurrence probability in postmenopausal early breast cancer patients with ET. Despite clinical ER positivity, 10 of the cases were assigned to nonluminal subtypes by PAM50 in this study. PAM50 expression for

intrinsic subtyping provided more prognostic information than standard clinical factors and IHC.

PAM50 was validated on over 2400 patients from two large retrospective studies [60, 61]. Using the mRNA of 1017 ER-positive early breast cancer patients from the ATAC trial, the PAM50 risk of recurrence score was studied and compared with Oncotype DX and IHC4 results [60]. IHC4 is a distant recurrence index derived from immunohistochemical analysis of estrogen and progesterone receptors, HER2 and Ki67. Additional prognostic information beyond that in the clinical treatment score, which integrates the prognostic information from nodal status, tumor size, histopathological grade, age, and treatment, was greater in the PAM50 risk of recurrence than the Oncotype DX RS for the overall population and for each subgroup (lymph node-negative and node-positive and HER2-negative/lymph node-negative subgroups). In the node-negative/HER2-negative subgroup, prognostic information obtained using the PAM50 risk of recurrence score was more accurate than the Oncotype DX RS. The correlation between risk of recurrence and clinical treatment score was similar to what was previously found in the TransATAC study between Oncotype DX RS and Adjuvant! Online. The risk of recurrence includes more clinicopathological information than the other factors. The PAM50 risk of recurrence score can be applied in the context of clinicopathological factors involved in the clinical treatment score.


One of the other large validation studies was on the patients from the ABCSG-8 trial. PAM50 was evaluated for obtaining the risk of recurrence and defining subtypes from the FFPE tumor tissues of 1478 patients from the ABCSG-8 trial [61]. The original study was a phase III prospective design of 3901 patients to test tamoxifen versus tamoxifen followed by anastrozole in the adjuvant treatment of ER-positive early breast cancer patients. Both risk of recurrence and intrinsic subtypes (luminal A/B, basal-like, and HER2 enriched) were defined using PAM50. There were 1004 (67.9 %) patients in the luminal A, 418 (28.3 %) patients in the luminal B, 48 (3.3 %) patients in the HER2-enriched, and 8 (0.5 %) patients in the



basal-like subgroups according to the PAM50 test. The aim was to test whether the risk of recurrence score adds prognostic value in predicting disease recurrence beyond standard clinical factors. For 10-year disease recurrence risk, lower than 10 % was defined as low risk, and higher than 20 % was defined as high risk (Fig. 11.2). The risk of recurrence score added prognostic information to the clinical predictors in all subgroups ( $p < 0.0001$ ). The luminal A subgroup had a lower risk of recurrence at 10 years compared with the luminal B subgroup ( $p < 0.0001$ ). Ten-year distant recurrence-free survival rates were higher in the luminal A subgroup than in the luminal B (HR, 2.85; 95 % CI, 2.04–4;  $p < 0.0001$ ) (Fig. 11.3). Low-risk and high-risk groups were discriminated using the risk of recurrence score in all subgroups of patients. As a result, the PAM50 test was validated in this study for predicting disease recurrence in ER-positive postmenopausal early breast cancer patients. A 10-year disease recurrence under 3.5 % in the risk of recurrence low category makes it unlikely that additional chemotherapy would improve the outcome.

The combined analysis of the TransATAC and ABCSG-8 trials with 2137 patients showed that


the risk of recurrence adds significant prognostic information for late recurrences (>5 years) in women with hormone receptor-positive early-stage breast cancer [62]. Median follow-up time was 10 years for that analysis. Predefined risk stratification showed significant differences between risk groups based on the risk of recurrence for 10-year distant recurrence rates. Patients with recurrence in the first 5 years were excluded from this analysis. The risk of recurrence score for the patients with recurrence ( $53.7 \pm 20.4$ ) in the first 5 years was higher than the patients with no recurrence ( $41.89 \pm 19.5$ ) ( $p < 0.001$ ). There were more late recurrences and more patients with poor differentiation, larger tumor size, and >3 positive lymph nodes in the TransATAC study than the ABCSG-8 trial. Of the 2137 patients analyzed, 1530 (73.8 %) women had luminal A and 542 (26.2 %) women had luminal B breast cancer. Patients in the luminal B subgroup had a 2.9-fold higher risk of distant recurrence (HR, 2.9; 95 % CI, 2.07–4.02;  $p = 0.001$ ). Patients were divided according to the 10-year recurrence probability as having a low, intermediate, and high risk of recurrence (lower than 10, 10–20 %, and higher than 20 %,



**Fig. 11.2** Prognosis based on PAM50 classifier for DRFS (Reproduced from Reference [61] with permission of Oxford University Press on behalf of the European Society for Medical Oncology. © The author (Michael Gnant) 2013)

	No. at risk:	0	1	2	3	4	5	6	7	8	9	10
Low	502	497	488	479	469	460	447	439	412	331	250	
Int	478	466	454	437	423	416	400	387	370	289	220	
High	498	484	465	428	399	379	347	330	301	238	198	

**Fig. 11.3** Distant recurrence-free survival: Kaplan–Meier plot of luminal (A, B) subtypes with 95 % CI (Reproduced from Reference [61] with permission of Oxford University Press on behalf of the European Society for Medical Oncology. © The author (Michael Gnant) 2013)



respectively). The risk of distant recurrence in 5–10 years was 2.4 % (95 % CI: 1.6–3.5), 8.3 % (95 % CI: 6.1–11.2), and 16.6 % (95 % CI: 13.1–20.9) in patients in the low, intermediate, and high risk of recurrence groups, respectively. The risk of late recurrence was 6.9 (HR, 6.9; 95 % CI, 4.54–10.47) times higher in the high-risk group than in the intermediate-risk group and 3.3 (HR, 3.26; 95 % CI, 2.07–5.13) times higher in the intermediate-risk group than the low-risk group. Based on the risk of recurrence score, patients in the high-risk group may be separated for extended therapy.

Tissue samples from the MA.12 NCIC CTG (National Cancer Institute of Canada Clinical Trials Group) prospective trial, which compared tamoxifen and placebo in early premenopausal breast cancer patients, were used for PAM50 risk of recurrence evaluation. The aim was to evaluate intrinsic breast cancer subtypes and prognosis with PAM50 and IHC [63], and 395 patients were evaluated. PAM50 gave significant prognostic information for both DFS and OS, but IHC analysis could not. The 5-year DFS and OS rates were higher in the luminal A group and lower in the HER2-enriched group according to PAM50 (*p*=0.0003). Classification of the patients into intrinsic subtypes using PAM50 was also more

effective than using IHC. This study also concluded that PAM50 was predictive for adjuvant tamoxifen treatment in node-negative and node-positive premenopausal breast cancer patients. There was significant interaction with 10-year DFS probability and the risk of recurrence score in lymph node-negative and node-positive early breast cancer patients treated with tamoxifen.

The phase III GEICAM/9906 study population (*N*=814) was evaluated for PAM50 breast cancer subtyping and clinical standard markers [64]. The standard IHC panel for breast cancer (ER, PR, and HER2) could not adequately define the PAM50 expression subtypes in this study. There was high agreement between biomarker scoring by protein IHC and gene expression, but gene expression determinants for ESR1 and ERBB2 status were more prognostic.

In a population-based study, 1319 women with breast cancer from the LACE and Pathways cohort were tested for intrinsic subtyping [65]. According to PAM50 subtyping, 53.1 % of the patients were luminal A, 20.5 % were luminal B, 13 % were HER2 enriched, 9.8 % were basal-like, and 3.6 % were normal-like. Among low-risk hormone receptor-positive patients with classical clinicopathological tests, only 76.5 % were categorized as luminal A by PAM50. In this

population-based cohort, African-American women were more likely to have basal-like tumors (OR: 4.4; 2.3–8.4), and Asian and Pacific Islander women had reduced odds of the basal-like subtype (OR: 0.5; 0.3–0.9). In another population-based cohort of 1691 patients, early (<5 years) and late (>5 years) risks of recurrence were determined based on the PAM50 risk of recurrence score according to intrinsic subtypes of breast cancer [66]. The risks of disease recurrence and death were lower in patients with luminal A tumors compared with luminal B, HER2-enriched, and basal-like subgroups of breast cancer at 2, 5, and 10 years. In that study, PAM50 better defined the patients with lower risk from the higher risk of recurrence tumors than the standard immunohistochemistry or tumor grade.

### Prediction of Response to Treatment

In a study of 104 postmenopausal ER-positive breast cancer patients, tumor biopsies were taken before and 2 weeks after the beginning of treatment with anastrozole [67]. The risk of recurrence was calculated by PAM50 for luminal A and B tumors. Among the pretreatment samples, all intrinsic subtypes were present, but the luminal subgroups were most highly represented. The decrease in Ki67 levels was evaluated between subgroups according to PAM50, and there was a similar proportionate decrease in Ki67 levels in the luminal A and B subgroups (mean suppression: 75 % for both), which suggests that patients in the luminal A and B subgroups derive similar benefit from anastrozole treatment. However, in the subgroups of basal-like (15 %) and HER2-enriched (50 %) cancers, Ki67 reductions were low under treatment with anastrozole. The normal breast-like subgroup showed the greatest reduction of Ki67 (83 %) with anastrozole treatment. Residual Ki67 staining remained high after 2 weeks of anastrozole treatment in the luminal B subgroup. The PAM50 risk of recurrence score was significantly associated with clinical outcome ( $p=0.03$ ) and antiproliferative response to anastrozole treatment ( $p=0.0019$ ). These data show that the short-term response to anastrozole treatment may be similar between luminal sub-

groups and that higher residual Ki67 levels might show poor response to anastrozole treatment in the luminal B subgroup.

Another validation study was the MA.5 NCIC CTG trial, which randomized early premenopausal, lymph node-positive breast cancer patients into CMF versus CEF (cyclophosphamide, epirubicin, and 5-fluorouracil) chemotherapy. PAM50 classified 467 patients into intrinsic subtypes, and the HER2-enriched subtype strongly predicted anthracycline sensitivity [68].

A subset of patients from the GEICAM/9906 phase III trial who were identified by PAM50 as having low proliferation status derived a larger benefit from weekly paclitaxel [49]. The original GEICAM/9916 study tested FEC versus FEC followed by weekly paclitaxel in node-positive early breast cancer patients. The PAM50 risk of recurrence was studied in 820 patients, and the median follow-up was 8.7 years. The median OS was higher in the FEC plus weekly paclitaxel arm compared with the FEC arm (HR, 0.693;  $p=0.013$ ). A benefit from weekly paclitaxel treatment was achieved only in patients with a low PAM50 risk of recurrence score (HR, 0.23;  $p<0.001$ ).

The NCIC CTG M1.21 trial was a phase III study of 2104 patients who were  $\leq 61$  years old and had high-risk node-negative or node-positive disease [69]. It tested different anthracycline and taxane combinations for the adjuvant treatment of breast cancer. Randomization was performed among doxorubicin plus cyclophosphamide and paclitaxel (AC/T), dose-dense CEF, and dose-dense, intense epirubicin, cyclophosphamide, and paclitaxel (EC/T) groups. Intrinsic subtyping was performed in 1094 available patients with the PAM50 assay [70]. Of these patients, 27 % were in the luminal A subgroup, 23 % were in the luminal B subgroup, 18 % were in the HER2-enriched subgroup, and 32 % were in the basal-like subgroup. Dose-dense CEF and dose-dense, intense EC/T treatments were superior to AC/T treatment ( $p=0.01$ ). In the multivariate analysis, a high risk of recurrence was associated with worse RFS ( $p=0.03$ ). However, categorical risk of recurrence was neither prognostic nor predictive for any treatment. In the multivariate analysis, intrinsic

subtyping with PAM50 had a significant prognostic effect on RFS ( $p=0.002$ ). Compared with luminal A, the hazard ratios were 1.48 (95 % CI: 0.92–2.37) for luminal B, 2.68 (95 % CI: 1.6–4.48) for HER2 enriched, and 1.97 (95 % CI: 1.1–3.53) for basal-like. Intrinsic subtypes were not predictive of treatment benefit (AC/T vs. EC/T-CEF). However, subgroup analysis showed that the nonluminal subtype was predictive for taxane benefit compared with the luminal subtype ( $p=0.05$ ).

Based on the results of these trials, PAM50 provides valuable information for intrinsic subtyping and also distant relapse-free survival and likelihood of recurrence at 10 years, at least for ER-positive and tamoxifen-treated breast cancer patients. PAM50 subtype classification is superior to IHC for both prognosis and predicting the benefit of tamoxifen. The risk of recurrence score gives an individual risk assessment in early stage, hormone receptor-positive, pre- and postmenopausal breast cancer patients and allows subtype classification. The risk of recurrence score estimates the probability of disease recurrence in pre- and postmenopausal, hormone receptor-positive early breast cancer patients treated with ET. The HER2-enriched subtype, as defined by PAM50, was a strong predictor of adjuvant anthracycline treatment. In light of these data, PAM50 was approved in Europe in 2012 and by the US FDA in 2013.

### Other Genomic Tests

Blueprint® is an 80-gene microarray that was specifically designed for molecular subclassification in early-stage (stage I and II), lymph node-negative or node-positive and ER-positive or ER-negative breast cancer patients [71]. In a neoadjuvant chemotherapy study of 426 breast cancer patients, the Blueprint® assay reassigned 22 % of the previously classified patients with IHC/FISH [72]. Of 211 patients classified as hormone receptor-positive/HER2-negative according to IHC/FISH, 37 were reclassified into the basal ( $N=35$ ) and HER2 ( $N=2$ ) subtypes by the Blueprint® assay. The pathological complete response rate was higher in the HER2 subgroup defined by Blueprint®

(53 %) than in the HER2 subgroup defined by IHC/FISH (38 %) ( $p=0.047$ ).

Endopredict® (Sividon Diagnostics GmbH, Germany) is an 11-gene assay that was recently introduced to predict disease recurrence risk in ER-positive and HER2-negative early breast cancers treated with adjuvant ET alone [73]. Eight selected genes and three additional control genes were used for quantification of mRNA levels by RT-PCR. FFPE can be used for Endopredict® analysis. The assay result is expressed as the Endopredict® (EP) score and was used in combination with nodal status and tumor size (EPclin) for a comprehensive risk estimation. EPclin was used in two validation studies of 378 and 1324 patients from the ABCSG-6 and ABCSG-8 studies, respectively [73, 74]. The 11-gene EP risk score was an independent predictor of disease recurrence in a multivariate analysis, and the EPclin score outperformed conventional clinicopathological risk factors. This test is used in laboratories in several centers in Germany, Austria, and Switzerland.

MapQuantDx® (Genomic Grade, Ipsogen, France) is a 97-gene histologic grade predictor that was developed in recent years. It is a microarray-based assay that calculates the Genomic Grade Index (GGI). It provides prognostic information in addition to standard clinicopathological variables and had significant impact on treatment decisions in several retrospective studies [75, 76]. Fresh or frozen tissue is needed for this 97-gene signature. However, a 6-gene PCR genomic grade was developed from the initial 97 genes for FFPE tissue. A high correlation was shown between the microarray and RT-PCR assays studied in frozen and FFPE tissues. The prognostic value of PCR-GGI was confirmed on FFPE samples [77].

The Rotterdam 76-gene signature was developed at the Erasmus University Cancer Center in Rotterdam (the Netherlands) [78, 79]. It was introduced to the market by Veridex (Raritan, USA). This 76-gene assay includes proliferation genes, none of which overlap with MammaPrint® or Oncotype DX®. Fresh or frozen tissue is needed for this assay. Validation was performed in patients with node-negative breast cancer that

was either positive or negative for hormone receptors. The 76-gene assay was validated in a group of 198 node-negative, systemically untreated breast cancer patients, and a high-risk group was defined for early disease recurrence in this study [80]. A very low-risk subgroup was also defined in another validation study in breast cancer patients treated with adjuvant ET [81].

### Conclusion

Numerous multigene assays have been developed based on multigene profiling assays to avoid over- and undertreatment and to better define the prognosis and predictive markers in early-stage breast cancer. Some of these assays provide subclassifications based on gene expression profiling. The main limitation for many gene profiling assays is the lack of “level of evidence I” due to the need for crucial prospective data and sufficient numbers of robust retrospective studies. Very few assays are suitable for use in common clinical practice due to technical and clinical validity. Clinicians must carefully consider the indications of these gene array-based assays regarding differences in technical prerequisites, reproducibility, clinical validity, underlying evidence, and clinical impact on the special patient populations. Among the most widely used multigene assays are Oncotype DX™, MammaPrint™, and PAM50™. Less expensive and more feasible assays are needed for decision-making about adjuvant or neoadjuvant treatment of breast cancer. Avoiding chemotherapy for a group of patients who would not benefit is very important for acute and long-term toxicities, quality-of-life issues, and cost. Based on numerous retrospective studies, ongoing prospective studies will provide important information about the treatment prediction in early breast cancer. Technical ease is a major concern for the general implementation of these tests in clinics. The availability of the test in a local lab is an advantage. However, the need for special equipment and trained personnel is critical for the local facility. Several other assays are undergoing validation studies or are otherwise in the process of development.

### References

1. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin.* 2013;63:11–30.
2. Early Breast Cancer Trialists' Collaborative G. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet.* 2005;365:1687–717.
3. Polychemotherapy for early breast cancer: an overview of the randomised trials. *Early Breast Cancer Trialists' Collaborative Group. Lancet.* 1998;352:930–42.
4. Tamoxifen for early breast cancer: an overview of the randomised trials. *Early Breast Cancer Trialists' Collaborative Group. Lancet.* 1998;351:1451–67.
5. Bedard PL, Cardoso F. Can some patients avoid adjuvant chemotherapy for early-stage breast cancer? *Nature reviews. Clin Oncol.* 2011;8:272–9.
6. Bryant J, Fisher B, Gunduz N, Costantino JP, Emir B. S-phase fraction combined with other patient and tumor characteristics for the prognosis of node-negative, estrogen-receptor-positive breast cancer. *Breast Cancer Res Treat.* 1998;51:239–53.
7. Henderson IC, Patek AJ. The relationship between prognostic and predictive factors in the management of breast cancer. *Breast Cancer Res Treat.* 1998;52:261–88.
8. Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, et al. Prognostic factors in breast cancer. *College of American Pathologists Consensus Statement 1999. Arch Pathol Lab Med.* 2000;124:966–78.
9. Bast Jr RC, Ravdin P, Hayes DF, Bates S, Fritsche Jr H, Jessup JM, et al. 2000 update of recommendations for the use of tumor markers in breast and colorectal cancer: clinical practice guidelines of the American Society of Clinical Oncology. *J Clin Oncol.* 2001;19:1865–78.
10. Goldhirsch A, Glick JH, Gelber RD, Coates AS, Senn HJ. Meeting highlights: International Consensus Panel on the Treatment of Primary Breast Cancer. *Seventh International Conference on Adjuvant Therapy of Primary Breast Cancer. J Clin Oncol.* 2001;19:3817–27.
11. Eifel P, Axelson JA, Costa J, Crowley J, Curran Jr WJ, Deshler A, et al. National Institutes of Health Consensus Development Conference Statement: adjuvant therapy for breast cancer, November 1–3, 2000. *J Natl Cancer Inst.* 2001;93:979–89.
12. Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. *Nature.* 2000;406:747–52.
13. Golub TR, Slonim DK, Tamayo P, Huard C, Gaasenbeek M, Mesirov JP, et al. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science.* 1999;286:531–7.
14. van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, et al. Gene expression profiling predicts clinical outcome of breast cancer. *Nature.* 2002;415:530–6.



15. van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, et al. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med*. 2002;347:1999–2009.
16. Harbeck N, Sotlar K, Wuerstlein R, Doisneau-Sixou S. Molecular and protein markers for clinical decision making in breast cancer: today and tomorrow. *Cancer Treat Rev*. 2014;40:434–44.
17. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med*. 2004;351:2817–26.
18. Cronin M, Pho M, Dutta D, Stephans JC, Shak S, Kiefer MC, et al. Measurement of gene expression in archival paraffin-embedded tissues: development and performance of a 92-gene reverse transcriptase-polymerase chain reaction assay. *Am J Pathol*. 2004;164:35–42.
19. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A*. 2001;98:10869–74.
20. Esteva FJ, Sahin AA, Coombes K, Baker J, Cronin M, Walker M, Watson D, Cristofanilli M, Shak S, Hortobagyi GN. Multi-gene RT-PCR assay for predicting recurrence in node negative breast cancer patients – M.D. Anderson Clinical Validation Study [abstract]. *Breast Cancer Res Treat*. 2003;82:A17. doi:10.1023/B:BREA.0000003871.38587.8b. <http://www.sabcs.org>.
21. Cobleigh MA, Bitterman P, Baker J, Cronin M, Liu M-L, Borchik R, Tabesh B, Mosquera J-M, Walker MG, Shak S. Tumor gene expression predicts distant disease-free survival (DDFS) in breast cancer patients with 10 or more positive nodes: high throughput RT-PCR assay of paraffin-embedded tumor tissues [abstract]. *Proc Am Soc Clin Oncol*. 2003;22:A3415.
22. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, Baehner R, Walker M, Watson D, Park T, et al. Multi-gene RT-PCR assay for predicting recurrence in node negative breast cancer patients – NSABP studies B-20 and B-14 [abstract]. *Breast Cancer Res Treat*. 2003;82:A16. <http://www.sabcs.org>.
23. Habel LA, Shak S, Jacobs MK, et al. A population-based study of tumor gene expression and risk of breast cancer death among lymph node-negative patients. *Breast Cancer Res*. 2006;8(3):R25. doi:10.1186/bcr1412.
24. Toi M, Iwata H, Yamanaka T, Masuda N, Ohno S, Nakamura S, et al. Clinical significance of the 21-gene signature (Oncotype DX) in hormone receptor-positive early stage primary breast cancer in the Japanese population. *Cancer*. 2010;116:3112–8.
25. Fisher B, Redmond C, Legault-Poisson S, Dimitrov NV, Brown AM, Wickerham DL, et al. Postoperative chemotherapy and tamoxifen compared with tamoxifen alone in the treatment of positive-node breast cancer patients aged 50 years and older with tumors responsive to tamoxifen: results from the National Surgical Adjuvant Breast and Bowel Project B-16. *J Clin Oncol*. 1990;8:1005–18.
26. Baum M, Budzar AU, Cuzick J, Forbes J, Houghton JH, Klijn JG, et al. Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early breast cancer: first results of the ATAC randomised trial. *Lancet*. 2002;359:2131–9.
27. Arimidex TAOiCTG, Forbes JF, Cuzick J, Buzdar A, Howell A, Tobias JS, et al. Effect of anastrozole and tamoxifen as adjuvant treatment for early-stage breast cancer: 100-month analysis of the ATAC trial. *Lancet Oncol*. 2008;9:45–53.
28. Fisher B, Dignam J, Wolmark N, DeCillis A, Emir B, Wickerham DL, et al. Tamoxifen and chemotherapy for lymph node-negative, estrogen receptor-positive breast cancer. *J Natl Cancer Inst*. 1997;89:1673–82.
29. Berry DA, Cirincione C, Henderson IC, Citron ML, Budman DR, Goldstein LJ, et al. Estrogen-receptor status and outcomes of modern chemotherapy for patients with node-positive breast cancer. *JAMA*. 2006;295:1658–67.
30. Hayes DF, Thor AD, Dressler LG, Weaver D, Edgerton S, Cowan D, et al. HER2 and response to paclitaxel in node-positive breast cancer. *N Engl J Med*. 2007;357:1496–506.
31. Mamounas EP, Bryant J, Lembersky B, Fehrenbacher L, Sedlacek SM, Fisher B, et al. Paclitaxel after doxorubicin plus cyclophosphamide as adjuvant chemotherapy for node-positive breast cancer: results from NSABP B-28. *J Clin Oncol*. 2005;23:3686–96.
32. Mamounas E, Tang G, Paik S, et al. Association between the 21-gene recurrence score (RS) and benefit from adjuvant paclitaxel (Pac) in node-positive (N+), ER-positive breast cancer patients (pts): results from NSABP B-28 [Abstract]. *Cancer Res*. 2012;72(24 Suppl):S1–10.
33. Albain KS, Barlow WE, Shak S, Hortobagyi GN, Livingston RB, Yeh IT, et al. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol*. 2010;11:55–65.
34. Goldstein LJ, Gray R, Badve S, Childs BH, Yoshizawa C, Rowley S, et al. Prognostic utility of the 21-gene assay in hormone receptor-positive operable breast cancer compared with classical clinicopathologic features. *J Clin Oncol*. 2008;26:4063–71.
35. Sparano JA, O'Neill A, Gray RJ, et al. 10-year update of E2197: phase III doxorubicin/docetaxel (AT) versus doxorubicin/cyclophosphamide (AC) adjuvant treatment of LN+ and high-risk LN- breast cancer and the comparison of the prognostic utility of the 21-gene Recurrence Score (RS) with clinicopathologic features. *J Clin Oncol*. 2012;30(Suppl 15) [abstract 1021].
36. Senkus E, Kyriakides S, Penault-Llorca F, Poortmans P, Thompson A, Zackrisson S, et al. Primary breast

- cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2013;24 Suppl 6:vi7–23.
37. Goldhirsch A, Winer EP, Coates AS, Gelber RD, Piccart-Gebhart M, Thurlimann B, et al. Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. *Ann Oncol.* 2013;24:2206–23.
  38. Harris L, Fritsche H, Mennel R, Norton L, Ravdin P, Taube S, et al. American Society of Clinical Oncology 2007 update of recommendations for the use of tumor markers in breast cancer. *J Clin Oncol.* 2007;25:5287–312.
  39. [http://www.nccn.org/professionals/physician\\_gls/pdf/breast.pdf](http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf). Accessed: 1 Jan 2015.
  40. Oratz R, Kim B, Chao C, Skrzypczak S, Ory C, Bugarini R, et al. Physician survey of the effect of the 21-gene recurrence score assay results on treatment recommendations for patients with lymph node-positive, estrogen receptor-positive breast cancer. *J Oncol Pract.* 2011;7:94–9.
  41. Klang S, Liebermann N, Rizel S, et al. The recurrence score and chemotherapy treatment in node-positive, ER+ early-stage breast cancer patients in Israel. *J Clin Oncol.* 2010;28(15 Suppl):[Abstract] 6075.
  42. de Boer RH, Baker C, Speakman D, Chao CY, Yoshizawa C, Mann GB. The impact of a genomic assay (Oncotype DX) on adjuvant treatment recommendations in early breast cancer. *Med J Aust.* 2013;199:205–8.
  43. Estevez LG, Calvo I, Abad MF, et al. A retrospective study in the Spanish population with Oncotype dx recurrence score (RS) in breast cancer patients with positive and negative-lymph nodes. *J Clin Oncol.* 2013;31(Suppl; [abstract] e11531).
  44. Carlson JJ, Roth JA. The impact of the Oncotype Dx breast cancer assay in clinical practice: a systematic review and meta-analysis. *Breast Cancer Res Treat.* 2013;141:13–22.
  45. Goncalves R, Bose R. Using multigene tests to select treatment for early-stage breast cancer. *J Natl Compr Cancer Network.* 2013;11:174–82; quiz 82.
  46. Sparano JA. TAILORx: trial assigning individualized options for treatment (Rx). *Clin Breast Cancer.* 2006;7:347–50.
  47. Gluz O, Kreipe H, Dehnhardt T, Christgen M, Kates R, Liedtke C, et al. Prospective comparison of risk assessment tools in early breast cancer (recurrence score, uPA/PAI-1, central grade, and luminal subtypes): final correlation analysis from the phase III WSG plan B trial. In: San Antonio Breast Cancer symposium; 2011. [Abstract] S4–3.
  48. Glas AM, Floore A, Delahaye LJ, Witteveen AT, Pover RC, Bakx N, et al. Converting a breast cancer microarray signature into a high-throughput diagnostic test. *BMC Genomics.* 2006;7:278.
  49. Martin M, Prat A, Rodriguez-Lescure A, Caballero R, Ebbert MT, Munarriz B, et al. PAM50 proliferation score as a predictor of weekly paclitaxel benefit in breast cancer. *Breast Cancer Res Treat.* 2013;138:457–66.
  50. Sapino A, Roepman P, Linn SC, Snel MH, Delahaye LJ, van den Akker J, et al. MammaPrint molecular diagnostics on formalin-fixed, paraffin-embedded tissue. *J Mol Diagn.* 2014;16:190–7.
  51. Buyse M, Loi S, van't Veer L, Viale G, Delorenzi M, Glas AM, et al. Validation and clinical utility of a 70-gene prognostic signature for women with node-negative breast cancer. *J Natl Cancer Inst.* 2006;98:1183–92.
  52. Mook S, Schmidt MK, Viale G, Pruneri G, Eekhout I, Floore A, et al. The 70-gene prognosis-signature predicts disease outcome in breast cancer patients with 1–3 positive lymph nodes in an independent validation study. *Breast Cancer Res Treat.* 2009;116:295–302.
  53. Saghatchian M, Mook S, Pruneri G, Viale G, Glas AM, Guerin S, et al. Additional prognostic value of the 70-gene signature (MammaPrint((R))) among breast cancer patients with 4–9 positive lymph nodes. *Breast.* 2013;22:682–90.
  54. Drukker CA, Bueno-de-Mesquita JM, Retel VP, van Harten WH, van Tinteren H, Wesseling J, et al. A prospective evaluation of a breast cancer prognosis signature in the observational RASTER study. *Int J Cancer.* 2013;133:929–36.
  55. Knauer M, Mook S, Rutgers EJ, Bender RA, Hauptmann M, van de Vijver MJ, et al. The predictive value of the 70-gene signature for adjuvant chemotherapy in early breast cancer. *Breast Cancer Res Treat.* 2010;120:655–61.
  56. Cusumano PG, Generali D, Ciruelos E, Manso L, Ghanem I, Lifrange E, et al. European inter-institutional impact study of MammaPrint. *Breast.* 2014;23:423–8.
  57. Rutgers E, Piccart-Gebhart MJ, Bogaerts J, Delaloge S, Veer LV, Rubio IT, et al. The EORTC 10041/BIG 03–04 MINDACT trial is feasible: results of the pilot phase. *Eur J Cancer.* 2011;47:2742–9.
  58. Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T, et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol.* 2009;27:1160–7.
  59. Nielsen TO, Parker JS, Leung S, Voduc D, Ebbert M, Vickery T, et al. A comparison of PAM50 intrinsic subtyping with immunohistochemistry and clinical prognostic factors in tamoxifen-treated estrogen receptor-positive breast cancer. *Clin Cancer Res.* 2010;16:5222–32.
  60. Dowsett M, Sestak I, Lopez-Knowles E, Sidhu K, Dumbier AK, Cowens JW, et al. Comparison of PAM50 risk of recurrence score with oncotype DX and IHC4 for predicting risk of distant recurrence after endocrine therapy. *J Clin Oncol.* 2013;31:2783–90.
  61. Gnani M, Filipits M, Greil R, Stoeger H, Rudas M, Bago-Horvath Z, et al. Predicting distant recurrence

- in receptor-positive breast cancer patients with limited clinicopathological risk: using the PAM50 Risk of Recurrence score in 1478 postmenopausal patients of the ABCSG-8 trial treated with adjuvant endocrine therapy alone. *Ann Oncol.* 2014;25:339–45.
62. Sestak I, Cuzick J, Dowsett M, Lopez-Knowles E, Filipits M, Dubsy P, et al. Prediction of late distant recurrence after 5 years of endocrine treatment: a combined analysis of patients from the Austrian breast and colorectal cancer study group 8 and arimidex, tamoxifen alone or in combination randomized trials using the PAM50 risk of recurrence score. *J Clin Oncol.* 2014.
63. Chia SK, Bramwell VH, Tu D, Shepherd LE, Jiang S, Vickery T, et al. A 50-gene intrinsic subtype classifier for prognosis and prediction of benefit from adjuvant tamoxifen. *Clin Cancer Res.* 2012;18:4465–72.
64. Bastien RR, Rodriguez-Lescure A, Ebbert MT, Prat A, Munarriz B, Rowe L, et al. PAM50 breast cancer subtyping by RT-qPCR and concordance with standard clinical molecular markers. *BMC Med Genomics.* 2012;5:44.
65. Sweeney C, Bernard PS, Factor RE, Kwan ML, Habel LA, Quesenberry Jr CP, et al. Intrinsic subtypes from PAM50 gene expression assay in a population-based breast cancer cohort: differences by age, race, and tumor characteristics. *Cancer Epidemiol Biomarkers Prev.* 2014;23:714–24.
66. Caan BJ, Sweeney C, Habel LA, Kwan ML, Kroenke CH, Weltzien EK, et al. Intrinsic subtypes from the PAM50 gene expression assay in a population-based breast cancer survivor cohort: prognostication of short- and long-term outcomes. *Cancer Epidemiol Biomarkers Prev.* 2014;23:725–34.
67. Dumbier AK, Anderson H, Ghazoui Z, Salter J, Parker JS, Perou CM, et al. Association between breast cancer subtypes and response to neoadjuvant anastrozole. *Steroids.* 2011;76:736–40.
68. Cheang MC, Voduc KD, Tu D, Jiang S, Leung S, Chia SK, et al. Responsiveness of intrinsic subtypes to adjuvant anthracycline substitution in the NCIC.CTG MA.5 randomized trial. *Clin Cancer Res.* 2012;18:2402–12.
69. Burnell M, Levine MN, Chapman JA, Bramwell V, Gelmon K, Walley B, et al. Cyclophosphamide, epirubicin, and Fluorouracil versus dose-dense epirubicin and cyclophosphamide followed by Paclitaxel versus Doxorubicin and cyclophosphamide followed by Paclitaxel in node-positive or high-risk node-negative breast cancer. *J Clin Oncol.* 2010;28:77–82.
70. Liu S, Chapman JA, Burnell MJ, Levine MN, Pritchard KI, Whelan TJ, et al. Prognostic and predictive investigation of PAM50 intrinsic subtypes in the NCIC CTG MA.21 phase III chemotherapy trial. *Breast Cancer Res Treat.* 2015;149:439–48.
71. Krijgsman O, Roepman P, Zwart W, Carroll JS, Tian S, de Snoo FA, et al. A diagnostic gene profile for molecular subtyping of breast cancer associated with treatment response. *Breast Cancer Res Treat.* 2012;133:37–47.
72. Whitworth P, Stork-Sloots L, de Snoo FA, Richards P, Rotkis M, Beatty J, et al. Chemosensitivity predicted by BluePrint 80-gene functional subtype and MammaPrint in the Prospective Neoadjuvant Breast Registry Symphony Trial (NBRST). *Ann Surg Oncol.* 2014;21:3261–7.
73. Filipits M, Rudas M, Jakesz R, Dubsy P, Fitzal F, Singer CF, et al. A new molecular predictor of distant recurrence in ER-positive, HER2-negative breast cancer adds independent information to conventional clinical risk factors. *Clin Cancer Res.* 2011;17:6012–20.
74. Dubsy P, Filipits M, Jakesz R, Rudas M, Singer CF, Greil R, et al. EndoPredict improves the prognostic classification derived from common clinical guidelines in ER-positive, HER2-negative early breast cancer. *Ann Oncol.* 2013;24:640–7.
75. Reyat F, Bollet MA, Caly M, Gentien D, Carpentier S, Peyro-Saint-Paul H, et al. Respective prognostic value of genomic grade and histological proliferation markers in early stage (pN0) breast carcinoma. *PLoS One.* 2012;7:e35184.
76. Metzger-Filho O, Michiels S, Bertucci F, Cateau A, Salgado R, Galant C, et al. Genomic grade adds prognostic value in invasive lobular carcinoma. *Ann Oncol.* 2013;24:377–84.
77. Toussaint J, Sieuwerts AM, Haibe-Kains B, Desmedt C, Rouas G, Harris AL, et al. Improvement of the clinical applicability of the Genomic Grade Index through a qRT-PCR test performed on frozen and formalin-fixed paraffin-embedded tissues. *BMC Genomics.* 2009;10:424.
78. Wang Y, Klijn JG, Zhang Y, Sieuwerts AM, Look MP, Yang F, et al. Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet.* 2005;365:671–9.
79. Foekens JA, Atkins D, Zhang Y, Sweep FC, Harbeck N, Paradiso A, et al. Multicenter validation of a gene expression-based prognostic signature in lymph node-negative primary breast cancer. *J Clin Oncol.* 2006;24:1665–71.
80. Desmedt C, Piette F, Loi S, Wang Y, Lallemand F, Haibe-Kains B, et al. Strong time dependence of the 76-gene prognostic signature for node-negative breast cancer patients in the TRANSBIG multicenter independent validation series. *Clin Cancer Res.* 2007;13:3207–14.
81. Zhang Y, Sieuwerts AM, McGreevy M, Casey G, Cufer T, Paradiso A, et al. The 76-gene signature defines high-risk patients that benefit from adjuvant tamoxifen therapy. *Breast Cancer Res Treat.* 2009;116:303–9.

Saadettin Kilickap, M. Kadri Altundag,  
and Y. Yavuz Ozisik

---

## Abstract

In clinical practice, although local recurrence or distant metastasis develops in some individuals who have been assessed as low risk despite treatment, some individuals with high-risk disease do not relapse despite systemic and local therapy. Therefore, oncologists need to determine objective prognostic factors to identify early recurrence and metastasis in patients with breast cancer. Based on the presumption of residual disease, clinicians have recently tried to identify micrometastases by using disseminated tumor cells (DTCs) in the bone marrow and circulating tumor cells (CTCs) from the peripheral blood. However, identifying the presence of DTCs and CTCs has required experienced staff and special laboratory.

DTCs are known as epithelial cells in the bone marrow, and they are also considered micrometastases in the bone marrow. The morphological characteristics of DTCs include large cells with a large nucleus, nuclear granulation or stippling, strong or irregular staining for cytokeratin, and cytokeratin filaments. DTCs are observed in approximately 30 % of early-stage breast cancer patients. Tumor cells circulating in the peripheral blood of patients with cancer are called CTCs. CTCs are cells that have entered the peripheral blood circulation after having detached from an existing primary tumor or its metastases. DTCs and CTCs can be used to predict progression-free and overall survival as well as response to treatment.

---

## Keywords

Breast cancer • Bone marrow • Micrometastases • Circulating tumor cells  
• Prognosis • Survival

---

S. Kilickap, MD, MSc (✉) • M.K. Altundag, MD  
Y.Y. Ozisik, MD  
Department of Medical Oncology, Hacettepe  
University Cancer Institute, Ankara, Turkey  
e-mail: [skilickap@yahoo.com](mailto:skilickap@yahoo.com);  
[kaltunda@hacettepe.edu.tr](mailto:kaltunda@hacettepe.edu.tr); [yozisik@hacettepe.edu.tr](mailto:yozisik@hacettepe.edu.tr)

## Background

Breast cancer is the worldwide leading cause of cancer death in women. Breast cancer-related mortality is associated with distant metastases. The 5-year relative survival rate is 22 % for stage IV breast cancer [1]. The response to treatment and survival outcomes of patients with breast cancer differ among individuals. The aim of treatment in patients with advanced stage breast cancer is palliative. By contrast, surgery is an essential treatment for early-stage breast cancer, and local or distant recurrence is the most important issue at follow-up in these patients. Among patients with favorable prognoses, approximately 20 % develop recurrence during the follow-up period, whereas other patients have long-term recurrence-free survival despite unfavorable prognoses [2, 3]. The aim of treatment in early-stage breast cancer is to achieve a cure or longer disease-free and overall survival. Adjuvant therapy with chemotherapy and/or radiotherapy following curative surgery aims to eradicate residual microscopic tumors. The traditional approach in the treatment of breast cancer is based on preventing and identifying the development of metastatic disease in early-stage breast cancer. At this stage, the presence of minimal residual disease or metastatic tumor cells is presumed but not certain. Clinical and pathological features associated with distant metastasis, such as high tumor grade, large tumor size, lymph node involvement, lack of hormone receptor expression, and overexpression of HER2, determine the treatment strategies. In clinical practice, while local recurrence or distant metastasis develops in some individuals who were assessed as low risk despite treatment, some individuals with high-risk disease do not relapse after systemic and local therapy. Therefore, oncologists must establish objective prognostic factors for identifying early recurrence and metastasis in patients with breast cancer.

The risk of recurrence after curative surgery in early-stage breast cancer is estimated with the 21-gene Oncotype DX and the 70-gene MammaPrint in patients with hormone receptor-positive cancers [4, 5]. After risk estimation,

patients with a high recurrence score are treated with chemotherapy and hormone therapy, whereas women with a low recurrence score are treated only with endocrine therapy. Despite classification as “high risk” for recurrence, some patients do not develop relapses within their lifetimes. Because of the presumption of residual disease, clinicians have recently tried to identify micrometastases based on DTCs in the bone marrow and CTCs from the peripheral blood. These cells can be detected with immunohistochemistry and molecular techniques. However, the detection of DTCs and CTCs requires experienced staff and specialized laboratories.

---

## Detection of Bone Marrow Micrometastases

In the bone marrow, DTCs can be identified as epithelial cells and are also recognized as micrometastases. Immunohistochemistry is the most common method used to detect micrometastases in bone marrow aspirates. The aspirates obtained from bone marrow are stained for epithelial cells. DTCs are observed in approximately 30 % of early-stage breast cancer patients [2]. The morphological characteristics of DTCs include large cells with a large nucleus, nuclear granulation or stippling, strong or irregular staining for cytokeratin, and cytokeratin filaments. The presence of at least one CK+ cell in the bone marrow is defined as DTC [6]. DTC may be established using immunohistochemistry and molecular techniques. The terms DTC or micrometastasis have been recently used in guidelines such as those of the American Society of Clinical Oncology and the San Gallen Consensus [7, 8].

The prognostic value of bone marrow micrometastases in breast cancer is illustrated in Table 12.1. The value of prognostic DTCs detected in the bone marrow after adjuvant chemotherapy is associated with poor prognosis [13]. In a study evaluating whether DTCs represented a prognostic factor in patients with breast cancer, the authors reported DTCs in 15 % of these patients [13]. The presence of DTCs in the bone marrow of breast cancer patients was nega-



**Table 12.1** Prognostic significance of bone marrow micrometastases in selected important breast cancer trials

	Year	N	Method	Positivity rate (%)	Prognostic value	Outcome
Braun et al. [2]	2005	4703	IHC	31	DFS, DDFS, OS	Prognostic
Bidard et al. [9]	2008	621	IHC	15	DMFS, OS	Prognostic
Molino et al. [10]	2008	125	IHC	31	RFS, OS	Non-prognostic
Falck et al. [11]	2012	401	IF, IHC	38	DDFS, OS	Non-prognostic
Hartkopf et al. [12]	2014	3141	IHC	26	DFS, OS	Prognostic

*IHC* immunohistochemistry, *IF* immunofluorescence, *DFS* disease-free survival, *DDFS* distant disease-free survival, *OS* overall survival, *DMFS* distant metastasis-free survival, *RFS* relapse-free survival

tively correlated with survival. Those results have been supported by studies using adjuvant and neoadjuvant chemotherapy [14, 15].

The detection of DTCs before surgery for early-stage breast cancer is an independent prognostic factor and is associated with shorter survival [2, 9, 16–18]. Most of these studies have shown a correlation between DTCs and other prognostic factors, including tumor size, histologic grade, and lymph node status.

In a meta-analysis that included individual data from 4703 patients, the frequency of DTCs was approximately 31 % [2]. These patients tended to have larger tumors, hormone receptor-negative cancers, and lymph node metastases. The presence of micrometastases in the bone marrow of patients with early-stage breast cancer before the initiation of primary therapy was an independent unfavorable prognostic factor for disease-free, distant-free, and overall survival in breast cancer. The mortality rate in patients with micrometastases is 2.4 times higher than in those without micrometastases. Additionally, persistent DTCs at follow-up after therapy in these patients predicted the risk of relapse and death and were associated with shortened overall survival [19]. Moreover, the results were consistent with those of several small studies in which the presence of DTCs in the bone marrow was a prognostic factor for survival [17, 18].

Molino et al. investigated the prognostic value of epithelial tumor cells in bone marrow in patients with breast cancer [10]. The authors found that the 10-year probability of relapse-free and overall survival rates in the patients in whom

epithelial cells were observed in the first bone marrow sample was similar with those without epithelial cells in the bone marrow. In Cox's model, the rates of relapse and death were different between patients with and without micrometastases in the first bone marrow samples. In a prospective observational study from Sweden, DTCs in the bone marrow were found in 38 % of 401 patients with breast cancer [11]. Tumor characteristics were similar between patients with and without detected DTCs. Detection of DTCs was not a prognostic factor for distant disease-free survival (DFS) or overall survival. These results did not support the results of previous published studies. Recently, a single-center study from Germany addressed whether the presence of DTCs had prognostic value in early-stage breast cancer patients [12]. In this study, which had a large sample size, data from 3141 patients were analyzed. DTCs were detected in 26 % of these patients. Patients who were DTC positive in the bone marrow tended to have larger tumor sizes as well as lymph node-positive and HER2-positive cancer. The presence of DTCs before surgery was associated with shortened disease-free and overall survival. The detection of DTCs in early-stage breast cancer is an independent predictor of disease-free and overall survival. However, DTCs are not detected in approximately half of patients with early breast cancer, and relapse can occur in patients without detectable DTCs [2].

The association between subtypes of breast cancer and the presence of DTCs in the bone marrow was examined [20]. In the luminal A

group, DTC detection was associated with a significantly poorer prognosis for relapse and death compared with the other groups. The lowest rate was observed in the basal-like group. However, for all subtypes, the relapse risk was significantly higher in DTC-positive patients than in DTC-negative patients.

Despite data suggesting that micrometastases in the bone marrow are an independent prognostic factor for survival, there are concerns regarding its validation. Thus, the clinical utility of DTC detection remains unclear.

## DTCS and CTCs

Bone marrow aspiration is an invasive procedure and is often painful, and standardizing its quality is particularly difficult. Therefore, effort has been focused on CTCs in the peripheral blood. However, it is unclear whether DTCs are the same cells as CTCs. To date, the accuracy of this hypothesis has been evaluated in clinical studies [21, 22] showing a positive correlation between DTCs and CTCs. DTCs were detected at a higher rate than CTCs.

Furthermore, these results were confirmed by a recent study performed using AdnaTest for the detection of CTCs [23]. Although these results support the possibility that DTC and CTC may be the same cells, the evidence remains insufficient.

## CTCS and Breast Cancer

Tumor cells circulating in the peripheral blood of patients with cancer are called CTCs. CTCs are cells that have entered the peripheral blood circulation after having detached from an existing primary tumor or its metastases. The presence of CTCs in the peripheral blood of patients with cancer was first described by Thomas Ashworth in 1869 [24]. However, cancer investigators have more recently focused on the molecular characterization and prognostic value of CTCs in many tumors, including breast, prostate, colorectal, ovarian, and pancreatic cancers [25]. Additionally, detecting CTCs can help to better understand the

biology of tumors and their metastasis in cancer patients [26, 27]. In many solid tumors, baseline CTC detection can provide useful information for estimating the prognosis and efficacy of treatment. The potential benefits of CTC detection in cancer patients are shown in Table 12.2. However, many unanswered questions remain, including the optimal method for quantifying and characterizing CTCs [28]. Another problem is the high heterogeneity of CTCs.

The CellSearch (Veridex LLC, Raritan, NJ, USA) system, a semi-automated methodology to detect and count CTCs in breast, colorectal, and prostate cancers, has been approved by the US Food and Drug Administration [29–31]. To detect CTCs, a 7.5 ml sample of peripheral blood is often sufficient. CTCs are analyzed using antibodies targeting three proteins: cytokeratin (CK), epithelial cell adhesion molecule (EpCAM), and CD45. Additionally, cell nuclei are fluorescently labeled with the DAPI (4',6-diamidino-2-phenylindole) nuclear dye. CTCs are defined as positive for nuclei and for epithelial cell adhesion molecule (EpCAM) and CK expression but negative for expression of the common leukocyte antigen CD45. The detection of at least one CTC per 7.5 ml of blood sample is considered positive. The advantages and disadvantages of CTC analysis are summarized in Table 12.3.

**Table 12.2** The potential benefits of CTC detection

Estimation of the risk of metastasis or tumor progression
Monitoring of treatment efficacy
Identification of resistance mechanisms
Understanding of the biology of metastasis
Prognosis

**Table 12.3** Advantages and disadvantages of CTC analysis

Advantages	Disadvantages
Minimally invasive	Expensive
Monitoring of treatment response	Isolation can be difficult
May be a prognostic marker	Requires experienced staff and specialized equipment
	Cutoff value uncertain

Recently, several technologies have been used to isolate CTCs, but they have not been approved by the FDA. These technologies include AdnaTest® (AdnaGen AG, Langerhagen, Germany), MACS (magnetic activated cell sorting system), and MagSweeper [32, 33]. Using AdnaTest, CTCs are isolated with antibodies against EpCAM and MUC1 [34]. The results of studies comparing CellSearch with other methods in patients with advanced cancer are conflicting. Muller et al. found that the CellSearch system is superior to the AdnaTest in metastatic breast cancer [35]. In contrast, another study demonstrated that CellSearch® and CTC-Chip systems were similarly effective [36]. However, at present, only the CellSearch system has been approved by the FDA.

Although CTCs can provide prognostic information in breast cancer, their utility as predictive markers is less certain. Studies about the prog-

nostic significance of CTCs in breast cancer have focused on both early-stage and metastatic disease. CTCs can be detected in the blood of many patients with breast cancer, even those without established metastasis. The prognostic significance of the detection of CTCs in breast cancer trials is presented in Table 12.4.

### Early-Stage Breast Cancer

CTCs can be detected in early-stage breast cancer and are associated with a high risk of relapse [39, 52, 53]. Many clinical studies have evaluated the prognostic value of CTCs in early-stage breast cancer. These studies include patients treated with both neoadjuvant and adjuvant chemotherapy. In these studies, the presence of CTCs in the peripheral blood has been associated with an increased risk of relapse and reduced

**Table 12.4** Prognostic significance of the detection of circulating tumor cells in selected important breast cancer trials

	Year	N	Method	Positivity rate (%)	Prognostic value	Outcome
<i>Non-metastatic breast cancer</i>						
Xenidis et al. [37]	2006	167	RT-PCR	22	DFS, OS	Predictive and prognostic
Ignatiadis et al. [38]	2007	444	RT-PCR	41	DFS, OS	Predictive
Pierga et al. [14]	2008	118	CellSearch	23	DFS	Prognostic for early relapse
Rack et al. [39]	2010	2026	CellSearch	22	DFS, OS	Prognostic
Bidard et al. [40]	2010	115	CellSearch	23	DFS, OS	Predict overall survival
Franken et al. [41]	2012	404	CellSearch	19	DFS	Prognostic
Lucci et al. [42]	2012	302	CellSearch	24	PFS, OS	Prognostic
<i>Metastatic breast cancer</i>						
Cristofanilli et al. [29]	2004	177	CellSearch	49	PFS, OS	Predictive
Hayes et al. [43]	2006	177	CellSearch	54	PFS, OS	Predictive for PFS but not OS
Giuliano et al. [44]	2011	235	CellSearch	40	PFS, OS	Prognostic
Müller et al. [35]	2012	254	CellSearch	50	PFS, OS	Predictive
			AdnaTest	40		
Giordano et al. [45]	2012	517	CellSearch	40	PFS, OS	Predictive
Pierga et al. [46]	2012	267	CellSearch	44	PFS, OS	Predictive
Wallwiener et al. [47]	2013	468	CellSearch	42	PFS, OS	Prognostic
Jiang et al. [48]	2013	294	CellSearch	77	PFS, OS	Prognostic
Smerage et al. [49]	2014	595	CellSearch	54	PFS, OS	Prognostic
Wallwiener et al. [50]	2014	393	CellSearch	34	PFS, OS	Predictive
Giuliano et al. [51]	2014	492	CellSearch	62	OS	Predictive

RT-PCR real-time polymerase chain reaction, DFS disease-free survival, PFS progression-free survival, OS overall survival

disease-free and overall survival [37, 54–56]. In a phase II trial including patients who were treated with neoadjuvant chemotherapy,  $\geq 1$  CTC was a predictive surrogate marker in predicting overall survival [14]. In the adjuvant setting, the presence of  $\geq 5$  CTCs/7.5 ml of peripheral blood was predictive of decreased survival in patients with HER2-positive but hormone receptor-negative cancer [38]. Furthermore, the presence of one or more CTC was associated with early recurrence and decreased overall survival in chemo-naïve operable breast cancer patients [42].

Rack et al. investigated the prognostic value of CTCs in early breast cancer [15]. In a large, multi-center, prospective, randomized trial, called the SUCCESS study, they used the CellSearch system to analyze CTCs at baseline and after the completion of chemotherapy. At baseline, CTCs were found in 21.5 % of 2026 patients who were diagnosed with early-stage breast cancer, and they were detected in 22.1 % of 1492 patients after the chemotherapy. The authors concluded that there was no association between the presence of CTCs and tumor characteristics, including size, grade, and hormone receptor status. However, they showed that the presence of CTCs before systemic chemotherapy was an independent prognostic factor for DFS and overall survival and that CTCs were associated with poor prognosis. In patients who were positive for CTCs, DFS at 36 months was 88.1 %, whereas it was 93.7 % in CTC-negative patients. Similarly, the breast cancer-related death rate was significantly higher in CTC-positive patients than in CTC-negative patients (40.9 % vs. 20.8 %, respectively). Additionally, these authors also analyzed the prognostic value of CTCs in early breast cancer patients. The DFS and overall survival after the completion of chemotherapy were significantly lower in persistently CTC-positive patients than in persistently CTC-negative patients (85.9 % vs. 93.9 % for DFS and 92.8 % vs. 97.6 % for OS, respectively).

Moreover, the prognostic impact of the presence of CTCs in early-stage breast cancer patients was preoperatively evaluated in two different studies [41, 42]. In one study including 404 patients with stage I–III breast cancer, CTCs were detected in approximately 20 % of cases at

the time of primary surgery [41]. In the other study, Lucci et al. found that CTCs were detected preoperatively in 24 % of patients [42]. Both studies suggested that the presence of CTCs before primary surgery was associated with early recurrence and an increased risk of breast cancer-related death, and these results have been confirmed in many small studies. CTCs were detected in 18–30 % of cases with early-stage breast cancer [40, 53, 57, 58]. The frequency of CTCs in early-stage breast cancer is lower than in metastatic patients [15, 29, 59].

The presence of CTCs and its prognostic impact were also evaluated in patients with locally advanced breast cancer in the neoadjuvant setting (Table 12.3). Two phase III trials, GeparQuattro and GeparQuinto, examined the relationship between tumor response and the presence of CTCs in breast cancer patients in the neoadjuvant setting [59, 60]. In both of these trials, CTCs were present in approximately 22 % of patients. CTC detection was similar across different tumor characteristics. There was no association between CTC detection and pathologic complete response in these neoadjuvant chemotherapy studies. In a French study, REMAGUS02, the presence of CTCs at baseline was inversely correlated with survival in patients treated with neoadjuvant chemotherapy [61].

Finally, in a meta-analysis including locally advanced stage breast cancer patients treated with neoadjuvant chemotherapy, the change in the number of CTCs was not associated with the pathologic response rate [62]. However, the presence of CTCs was associated with shorter DFS and overall survival.

## Metastatic Breast Cancer

Metastatic disease is the most common cause of breast cancer-related mortality. The traditional prognostic factors affect treatment strategies and prognosis in breast cancer. However, new treatment strategies and prognostic factors are needed in patients with asymptomatic visceral metastases and in those with the disease confined to non-visceral organs.

The prognostic role of CTCs in metastatic breast cancer has been investigated in many trials. Most of these studies have shown that the presence of CTCs is a prognostic factor for both disease-free and overall survival in the metastatic setting. Cristofanilli et al. showed that CTCs with a cutoff of 5 CTCs/7.5 ml of peripheral blood in patients with breast cancer represented an independent prognostic factor for progression-free and overall survival in the metastatic setting [29]. Elevated CTC counts were associated with significantly shorter progression-free and overall survival compared with the rates in patients with <5 CTCs per 7.5 ml of blood. The outcomes were highly predictive. Despite debates about the appropriate cutoff value for CTCs, the same value was supported in additional studies [43, 63, 64]. Thus, whether CTCs act as a prognostic and predictive surrogate marker in many solid tumors has been evaluated by multiple investigators [35, 45–47, 65]. Additionally, the CTC count may predict the response to treatment in breast cancer patients. Lack of CTC response after chemotherapy is associated with a worse prognosis and shorter progression-free survival (PFS) and overall survival [66, 67].

The CTC count at baseline is an independent prognostic factor for PFS and overall survival in metastatic breast cancer [48]. The effect is very prominent in the HER2-positive histologic subtype of breast cancer, regardless of treatment.

The presence of CTCs in peripheral blood may also be a potential marker of micrometastatic disease in patients with breast cancer. The prognostic value of elevated CTCs may not only correlate with higher metastatic tumor burden and metastatic sites but also predict micrometastatic disease [44, 49–51, 65, 68]. The results of a retrospective study including 492 advanced stage breast cancer patients suggested that the number of metastatic sites increased with a pretreatment level of  $\geq 5$  CTCs/7.5 ml compared with <5 CTCs/7.5 ml [44]. The authors showed that the development of new metastatic lesions and metastatic sites increased in patients with higher baseline CTC counts and that CTCs could be used as an indicator of metastatic potential in patients with limited metastatic dissemination.

However, whether the detection of CTCs using only the baseline measurement correctly identifies the prognosis remains unclear. Some clinical studies indicate that serial enumeration of CTCs should be performed to predict prognosis rather than a single measurement [43, 46, 50, 51, 69, 70]. Wallwiener et al. showed that serial enumeration of CTCs was more effective as a prognostic indicator and was useful for therapeutic monitoring of metastatic breast cancer [50]. The authors found that PFS and overall survival were significantly higher in patients with negative CTC status compared with CTC-positive patients after one cycle of treatment. They recommended that monitoring of CTCs should be performed at baseline and after one cycle of treatment. They also suggested that changes in CTC status from baseline to completion of one treatment cycle were predictive for PFS and overall survival.

In the SWOG S0500 trial, the presence of CTCs before first-line chemotherapy in metastatic breast cancer patients was strongly prognostic and was associated with decreased survival [49]. The worst survival was prominent in patients who did not show reduced numbers of CTCs after chemotherapy. In a phase II study, the authors evaluated the prognostic role of CTCs in HER2-positive metastatic breast cancer patients with brain metastasis; CTCs were detected at baseline and at the third week after chemotherapy with lapatinib and capecitabine [71]. The proportion of patients with detectable CTCs decreased from 49 to 18 % after only one cycle of chemotherapy. The response was significantly higher in patients who did not show CTCs at the third week, and the 1-year overall survival rate in these patients (84 %) was significantly higher than for those with  $\geq 1$  CTCs (43 %).

### CTCS and Breast Cancer Subtypes

The authors of the SWOG S0500 trial analyzed all patients with metastatic breast cancer according to three biologic subtypes [49]. The first group consisted of patients who were negative for CTCs at baseline (group A), and group B was



defined as patients who were initially positive for CTCs but who showed decreased CTCs after only one cycle of chemotherapy. In contrast, group C consisted of patients who showed increased CTCs after the first cycle of chemotherapy. In subtype analysis, the median overall survival was significantly lower in group C than in the other groups. The median overall survival rates were 35 months, 23 months, and 13 months in groups A, B, and C, respectively. Additionally, within each subgroup, the median overall survival was higher in patients with HER2-negative but hormone receptor-positive cancers than in those with hormone receptor-negative cancer.

Although within group A, the worst prognosis was found in triple-negative patients, and the median overall survival was still higher than for HER2-negative but hormone receptor-positive patients (22 vs. 15 months, respectively). Furthermore, the lowest median overall survival was observed in patients who showed increases in CTCs from the baseline (9.5 months), regardless of biologic subtype. In this group, 75 % of patients died within approximately 15 months. This study also showed that the quantification of CTCs after chemotherapy provided additional information regarding survival.

In their study, Rack et al. showed no association between the presence of CTCs and histological subtypes, including luminal, basal-like, and HER2-positive cancers [70]. However, the presence of CTCs at baseline was an independent prognostic factor for reduced DFS. In all subtypes, DFS was significantly reduced in node-positive patients compared with those with lymph node-negative cancer.

Lymph node positivity at the time of initial diagnosis is also a prognostic marker for survival. In the SUCCESS and EUDRA-CT trials, CTC detection was higher in patients with lymph node-positive cancer [72]. These trials showed that lymph node metastasis is an independent prognostic factor for survival in multivariate analysis.

### Conclusion

Whether the presence of tumor cells in the bone marrow and peripheral blood is suffi-

cient to detect micrometastases and to predict survival remains unclear. In the absence of overt metastases, the detection of DTCs in the bone marrow or CTCs in the peripheral blood is a comparable factor to predictors such as tumor size, differentiation, lymphatic involvement, and HER2 and hormonal status. Evidence suggests that CTCs represent residual disease and are associated with disease-free, recurrence-free, and overall survival. Further large studies and scientific evidence are needed to support whether DTCs and/or CTCs reflect patient outcomes.

### References

1. <http://www.cancer.org/cancer/breastcancer/detailedguide/breast-cancer-survival-by-stage>. Accessed date: 21.2.2015.
2. Braun S, Vogl FD, Naume B. A pooled analysis of bone marrow micrometastasis in breast cancer. *N Engl J Med*. 2005;353:793–802.
3. Vincent-Salomon A, Bidard FC, Pierga JY. Bone marrow micrometastasis in breast cancer: review of detection methods, prognostic impact and biological issues. *J Clin Pathol*. 2008;61:570–6.
4. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, et al. Expression of the 21 genes in the Recurrence Score assay and tamoxifen clinical benefit in the NSABP study B-14 of node negative, estrogen receptor positive breast cancer. *J Clin Oncol*. 2005;23:abstract 510.
5. van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, et al. A gene expression signature as a predictor of survival in breast cancer. *N Engl J Med*. 2002;347:1999–2009.
6. Fehm T, Braun S, Muller V, Janni W, Gebauer G, Marth C, et al. A concept for the standardized detection of disseminated tumor cells in bone marrow from patients with primary breast cancer and its clinical implementation. *Cancer*. 2006;107:885–92.
7. Harris L, Fritsche H, Mennel R, Norton L, Ravdin P, Taube S, et al. American Society of Clinical Oncology 2007 update of recommendations for the use of tumor markers in breast cancer. *J Clin Oncol*. 2007;25:5287–312.
8. Goldhirsch A, Ingel JN, Gelber RD, Coates AS, Thürlimann B, Senn HJ, et al. Thresholds for therapies: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2009. *Ann Oncol*. 2009;20:1319–29.
9. Bidard FC, Vincent-Salomon A, Gomme S, Nos C, de Rycke Y, Thiery JP, et al. Disseminated tumor cells of breast cancer patients: a strong prognostic factor for

- distant and local relapse. *Clin Cancer Res.* 2008;14:3306–11.
10. Molino A, Giovannini M, Micciolo R, Auriemma A, Fiorio E, Santo A, et al. Bone marrow micrometastases in breast cancer patients: a long-term follow-up study. *Clin Med Oncol.* 2008;2:487–90.
  11. Falck AK, Bendahl PO, Ingvar C, Isola J, Jönsson PE, Lindblom P, et al. Analysis of and prognostic information from disseminated tumour cells in bone marrow in primary breast cancer: a prospective observational study. *BMC Cancer.* 2012;12:403.
  12. Hartkopf AD, Taran FA, Wallwiener M, Hahn M, Becker S, Solomayer EF, et al. Prognostic relevance of disseminated tumour cells from the bone marrow of early stage breast cancer patients – results from a large single-centre analysis. *Eur J Cancer.* 2014;50:2550–9.
  13. Naume B, Fehm T, Wiedswang G, Juckstock J, Borgen E, Rack B, et al. Persistence of isolated tumor cells in the bone marrow of breast cancer patients predicts increased risk for relapse – a European pooled analysis. *Cancer Res.* 2009;69:abstr. 304.
  14. Pierga JY, Bidard FC, Mathiot C, Brain E, Delaloge S, Giachetti S, et al. Circulating tumor cell detection predicts early metastatic relapse after neoadjuvant chemotherapy in large operable and locally advanced breast cancer in a phase II randomized trial. *Clin Cancer Res.* 2008;14:7004–10.
  15. Rack BK, Schindlbeck C, Schneeweiss A, Hilfrich J, Lorenz R, Beckmann MW, et al. Prognostic relevance of circulating tumor cells (CTCs) in peripheral blood of breast cancer patients before and after adjuvant chemotherapy. *J Clin Oncol.* 2008;26:abstr. 503.
  16. Braun S, Pantel K, Muller P, Janni W, Hepp F, Kutenich CR, et al. Cytokeratin-positive cells in the bone marrow and survival of patients with stage I, II, or III breast cancer. *N Engl J Med.* 2000;342:525–33.
  17. Diel IJ, Kaufmann M, Costa SD, Holle R, von Minckwitz G, Solomayer EF, et al. Micrometastatic breast cancer cells in bone marrow at primary surgery: prognostic value in comparison with nodal status. *J Natl Cancer Inst.* 1996;88:1652–8.
  18. Braun S, Pantel K. Micrometastatic bone marrow involvement: detection and prognostic significance. *Med Oncol.* 1999;16:154–65.
  19. Janni W, Vogl FD, Wiedswang G, Synnestvedt M, Fehm T, Juckstock J, et al. Persistence of disseminated tumor cells in the bone marrow of breast cancer patients predicts increased risk for relapse—a European pooled analysis. *Clin Cancer Res.* 2011;17:2967–76.
  20. Naume B, Zhao X, Synnestvedt M, Borgen E, Russnes HG, Lingjaerde OC, et al. Presence of bone marrow micrometastasis is associated with different recurrence risk within molecular subtypes of breast cancer. *Mol Oncol.* 2007;1:160–71.
  21. Müller V, Stahmann N, Riethdorf S, Rau T, Zabel T, Goetz A, et al. Circulating tumor cells in breast cancer: correlation to bone marrow micrometastases, heterogeneous response to systemic therapy and low proliferative activity. *Clin Cancer Res.* 2005;11:3678–85.
  22. Pierga JY, Bonneton C, Vincent-Salomon A, de Cremoux P, Nos C, Blin N, et al. Clinical significance of immunocytochemical detection of tumor cells using digital microscopy in peripheral blood and bone marrow of breast cancer patients. *Clin Cancer Res.* 2004;10:1392–400.
  23. Fehm T, Becker S, Becker-Pergola G, Sotlar K, Gebauer G, Dürr-Störzer S, et al. Presence of apoptotic and nonapoptotic disseminated tumor cells reflects the response to neoadjuvant systemic therapy in breast cancer. *Breast Cancer Res.* 2006;8:R60.
  24. Ashworth TR. A case of cancer in which cells similar to those in the tumours were seen in the blood after death. *Med J Australia.* 1869;14:146–7.
  25. Lianidou ES, Strati A, Markou A. Circulating tumor cells as promising novel biomarkers in solid cancers. *Crit Rev Clin Lab Sci.* 2014;51:160–71.
  26. Lianidou ES, Mavroudis D, Georgoulis V. Clinical challenges in the molecular characterization of circulating tumour cells in breast cancer. *Br J Cancer.* 2013;108:2426–32.
  27. Lianidou ES, Markou A, Strati A. Molecular characterization of circulating tumor cells in breast cancer: challenges and promises for individualized cancer treatment. *Cancer Metastasis Rev.* 2012;31:663–71.
  28. Parkinson DR, Dracopoli N, Petty BG, Compton C, Cristofanilli M, Deisseroth A, et al. Considerations in the development of circulating tumor cell technology for clinical use. *J Transl Med.* 2012;10:138.
  29. Cristofanilli M, Budd GT, Ellis MJ, Stopeck A, Matera J, Miller MC, et al. Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med.* 2004;351:781–91.
  30. Cohen SJ, Alpaugh RK, Gross S, O'Hara SM, Smirnov DA, Terstappen LW, et al. Isolation and characterization of circulating tumor cells in patients with metastatic colorectal cancer. *Clin Colorectal Cancer.* 2006;6:125–32.
  31. Okegawa T, Nutahara K, Higashihara E. Prognostic significance of circulating tumor cells in patients with hormone refractory prostate cancer. *J Urol.* 2009;181:1091–907.
  32. Mostert B, Sleijfer S, Foekens JA, Gratama JW. Circulating tumor cells (CTCs): detection methods and their clinical relevance in breast cancer. *Cancer Treat Rev.* 2009;35:463–74.
  33. Talasz AH, Powell AA, Huber DE, Berbee JG, Roh KH, Yu W, et al. Isolating highly enriched populations of circulating epithelial cells and other rare cells from blood using a magnetic sweeper device. *Proc Natl Acad Sci U S A.* 2009;106:3970–5.
  34. Allan AL, Keeney M. Circulating tumor cell analysis: technical and statistical considerations for application to the clinic. *J Oncol.* 2010;2010:426218.
  35. Müller V, Riethdorf S, Rack B, Janni W, Fasching PA, Solomayer E, et al. Prognostic impact of circulating tumor cells assessed with the Cell Search System™

- and AdnaTest Breast™ in cancer patients: the DETECT study. *Breast Cancer Res.* 2012;14:R118.
36. Punnoose EA, Atwal SK, Spoerke JM, Savage H, Pandita A, Yeh RF, et al. Molecular biomarker analyses using circulating tumor cells. *PLoS One.* 2010;5:e12517.
  37. Xenidis N, Perraki M, Kafousi M, Apostolaki S, Bolonaki I, Stathopoulou A, et al. Predictive and prognostic value of peripheral blood cytokeratin-19 mRNA-positive cells detected by real-time polymerase chain reaction in node-negative breast cancer patients. *J Clin Oncol.* 2006;24:3756–62.
  38. Ignatiadis M, Xenidis N, Perraki M, Apostolaki S, Politaki E, Kafousi M, et al. Different prognostic value of cytokeratin-19 mRNA positive circulating tumor cells according to estrogen receptor and HER2 status in early-stage breast cancer. *J Clin Oncol.* 2007;25:5194–202.
  39. Rack BK, Schindlbeck C, Andergassen U, Schneeweiss A, Zwingers T, Lichtenegger W, et al. Use of circulating tumor cells (CTC) in peripheral blood of breast cancer patients before and after adjuvant chemotherapy to predict risk for relapse: the SUCCESS trial. *J Clin Oncol.* 2010;28 Suppl 15:1003.
  40. Bidard FC, Mathiot C, Delaloge S, Brain E, Giachetti S, de Cremoux P, et al. Single circulating tumor cell detection and overall survival in nonmetastatic breast cancer. *Ann Oncol.* 2010;21:729–33.
  41. Franken B, de Groot MR, Mastboom WJ, Vermees I, van der Palen J, Tibbe AG, et al. Circulating tumor cells, disease recurrence and survival in newly diagnosed breast cancer. *Breast Cancer Res.* 2012;14:R133.
  42. Lucci A, Hall CS, Lodhi AK, Bhattacharyya A, Anderson AE, Xiao L, et al. Circulating tumour cells in non-metastatic breast cancer: a prospective study. *Lancet Oncol.* 2012;13:688–95.
  43. Hayes DF, Cristofanilli M, Budd GT, Ellis MJ, Stopeck A, Miller MC, et al. Circulating tumor cells at each follow-up time point during therapy of metastatic breast cancer patients predict progression-free and overall survival. *Clin Cancer Res.* 2006;12:4218–24.
  44. Giuliano M, Giordano A, Jackson S, Hess KR, De Giorgi U, Mego M, et al. Circulating tumor cells as prognostic and predictive markers in metastatic breast cancer patients receiving first-line systemic treatment. *Breast Cancer Res.* 2011;13:R67.
  45. Giordano A, Cristofanilli M. CTCs in metastatic breast cancer. *Recent Results Cancer Res.* 2012;195:193–201.
  46. Pierga JY, Hajage D, Bachelot T, Delaloge S, Brain E, Campone M, et al. High independent prognostic and predictive value of circulating tumor cells compared with serum tumor markers in a large prospective trial in first-line chemotherapy for metastatic breast cancer patients. *Ann Oncol.* 2012;23:618–24.
  47. Wallwiener M, Hartkopf AD, Baccelli I, Riethdorf S, Schott S, Pantel K, et al. The prognostic impact of circulating tumor cells in subtypes of metastatic breast cancer. *Breast Cancer Res Treat.* 2013;137:503–10.
  48. Jiang ZF, Cristofanilli M, Shao ZM, Tong ZS, Song EW, Wang XJ, et al. Circulating tumor cells predict progression-free and overall survival in Chinese patients with metastatic breast cancer, HER2-positive or triple-negative (CBCSG004): a multicenter, double-blind, prospective trial. *Ann Oncol.* 2013;24:2766–72.
  49. Smerage JB, Barlow WE, Hortobagyi GN, Winer EP, Leyland-Jones B, Srkalovic G, et al. Circulating tumor cells and response to chemotherapy in metastatic breast cancer: SWOG S0500. *J Clin Oncol.* 2014;32:3483–9.
  50. Wallwiener M, Riethdorf S, Hartkopf AD, Modugno C, Nees J, Madhavan D. Serial enumeration of circulating tumor cells predicts treatment response and prognosis in metastatic breast cancer: a prospective study in 393 patients. *BMC Cancer.* 2014;14:512.
  51. Giuliano M, Giordano A, Jackson S, De Giorgi U, Mego M, Cohen EN, et al. Circulating tumor cells as early predictors of metastatic spread in breast cancer patients with limited metastatic dissemination. *Breast Cancer Res.* 2014;16:440.
  52. Krishnamurthy S, Cristofanilli M, Singh B, Reuben J, Gao H, Cohen EN, et al. Detection of minimal residual disease in blood and bone marrow in early stage breast cancer. *Cancer.* 2010;116:3330–7.
  53. Riethdorf S, Muller V, Zhang L, Zhang L, Rau T, Loibl S, et al. Detection and HER2 expression of circulating tumor cells: prospective monitoring in breast cancer patients treated in the neoadjuvant GeparQuattro trial. *Clin Cancer Res.* 2010;16:2634–45.
  54. Xenidis N, Markos V, Apostolaki S, Perraki M, Pallis A, Sfakiotaki G, et al. Clinical relevance of circulating CK-19 mRNA-positive cells detected during the adjuvant tamoxifen treatment in patients with early breast cancer. *Ann Oncol.* 2007;18:1623–31.
  55. Xenidis N, Ignatiadis M, Apostolaki S, Perraki M, Kalbakis K, Agelaki S, et al. Cytokeratin-19 mRNA-positive circulating tumor cells after adjuvant chemotherapy in patients with early breast cancer. *J Clin Oncol.* 2009;27:2177–84.
  56. Saloustros E, Perraki M, Apostolaki S, Kallergi G, Xyrafas A, Kalbakis K, et al. Cytokeratin-19 mRNA-positive circulating tumor cells during follow-up of patients with operable breast cancer: prognostic relevance for late relapse. *Breast Cancer Res.* 2011;13:R60.
  57. Sandri MT, Zorzino L, Cassatella MC, Bassi F, Luini A, Casadio C, et al. Changes in circulating tumor cell detection in patients with localized breast cancer before and after surgery. *Ann Surg Oncol.* 2010;17:1539–45.
  58. Biggers B, Knox S, Grant M, Kuhn J, Nemunaitis J, Fisher T, et al. Circulating tumor cells in patients undergoing surgery for primary breast cancer: preliminary results of a pilot study. *Ann Surg Oncol.* 2009;16:969–71.
  59. Fehm T, Muller V, Aktas B, Janni W, Schneeweiss A, Stickeler E, et al. HER2 status of circulating tumor

- cells in patients with metastatic breast cancer: a prospective, multicenter trial. *Breast Cancer Res Treat.* 2010;124:403–12.
60. Untch M, Loibl S, Bischoff J, Eidtmann H, Kaufmann M, Blohmer JU, et al. Lapatinib versus trastuzumab in combination with neoadjuvant anthracycline-taxane-based chemotherapy (GeparQuinto, GBG 44): a randomised phase 3 trial. *Lancet Oncol.* 2012;13:135–44.
  61. Bidard FC, Belin L, Delalogue S, Lerebours F, Ngo C, Reyal F, et al. Time-dependent prognostic impact of circulating tumor cells detection in non-metastatic breast cancer: 70-month analysis of the REMAGUS02 study. *Int J Breast Cancer.* 2013;2013:130470.
  62. Fei F, Du Y, Di G, Wu J, Shao Z. Are changes in circulating tumor cell (CTC) count associated with the response to neoadjuvant chemotherapy in local advanced breast cancer? A meta-analysis. *Oncol Res Treat.* 2014;37(5):250–4.
  63. Dawood S, Broglio K, Valero V, Reuben J, Handy B, Islam R, et al. Circulating tumor cells in metastatic breast cancer: from prognostic stratification to modification of the staging system? *Cancer.* 2008;113:2422–30.
  64. Zhang L, Riethdorf S, Wu G, Wang T, Yang K, Peng G, et al. Meta-analysis of the prognostic value of circulating tumor cells in breast cancer. *Clin Cancer Res.* 2012;18:5701–10.
  65. Giordano A, Egleston BL, Hajage D, Bland J, Hortobagyi GN, Reuben JM, et al. Establishment and validation of circulating tumor cell-based prognostic nomograms in first-line metastatic breast cancer patients. *Clin Cancer Res.* 2013;19:1596–602.
  66. Martin M, Custodio S, de Las Casas ML, Garcia-Saenz JA, de la Torre JC, Bellon-Cano JM, et al. Circulating tumor cells following first chemotherapy cycle: an early and strong predictor of outcome in patients with metastatic breast cancer. *Oncologist.* 2013;18:917–23.
  67. Budd GT, Cristofanilli M, Ellis MJ, Stopeck A, Borden E, Miller MC, et al. Circulating tumor cells versus imaging-predicting overall survival in metastatic breast cancer. *Clin Cancer Res.* 2006;12:6403–9.
  68. Giordano A, Giuliano M, De Laurentiis M, Arpino G, Jackson S, Handy BC, et al. Circulating tumor cells in immunohistochemical subtypes of metastatic breast cancer: lack of prediction in HER2-positive disease treated with targeted therapy. *Ann Oncol.* 2012;23:1144–50.
  69. Giordano A, Giuliano M, Handy BC, Ueno NT, Alvarez RH, Cohen EN, et al. Clinical value of circulating tumor cells (CTC) in first-line metastatic breast cancer according to type of treatment and immunohistochemical molecular subtype. San Antonio Breast Cancer symposium 2010: Abstracts, PD04-02.
  70. Liu MC, Shields PG, Warren RD, Cohen P, Wilkinson M, Ottaviano YL, et al. Circulating tumor cells: a useful predictor of treatment efficacy in metastatic breast cancer. *J Clin Oncol.* 2009;27:5153–9.
  71. Pierga JY, Bidard FC, Cropet C, Tresca P, Dalenc F, Romieu G, et al. Circulating tumor cells and brain metastasis outcome in patients with HER2-positive breast cancer: the LANDSCAPE trial. *Ann Oncol.* 2013;24(12):2999–3004.
  72. Rack B, Schindlbeck C, Jückstock J, Andergassen U, Hepp P, Zwingers T, et al. Circulating tumor cells predict survival in early average-to-high risk breast cancer patients. *J Natl Cancer Inst.* 2014;106(9). doi:10.1093/jnci/dju273.

Sitki Tuzlali

---

## Abstract

In this chapter, pathological features of breast carcinoma are discussed. Classification is based on the recent WHO classification of breast carcinoma, and specific gross and microscopic features of in situ and invasive breast carcinomas are explained. Morphological groups, grading of DCIS, and the necessary information that should be in a surgical pathology report are discussed. Recent information about columnar cell lesions and flat epithelial atypia of the breast are discussed, along with their clinical importance. Atypical lobular hyperplasia and lobular carcinoma in situ (LCIS) are discussed under the heading “lobular neoplasia.” Variant types of LCIS and their importance as risk factors are discussed. Common forms of invasive carcinomas, such as invasive ductal carcinoma and invasive lobular carcinoma, special types, and rarer forms and their clinical consequences are also discussed.

---

## Keywords

Breast • Breast cancer • Breast carcinoma • Ductal carcinoma in situ • Invasive ductal carcinoma • Lobular carcinoma • Tubular carcinoma • Cribriform carcinoma • Mucinous carcinoma • Medullary carcinoma • Papillary carcinoma

---

## Pathology of Breast Cancer

Histopathologically, breast carcinoma is simply divided into two major categories with respect to its confinement to the ductal-lobular system of the breast or not:

1. In situ
2. Invasive

---

S. Tuzlali, MD  
Department of Pathology, Istanbul Medical Faculty,  
Istanbul University, Istanbul, 34390, Turkey  
e-mail: [stuzlali@hotmail.com](mailto:stuzlali@hotmail.com)



## Carcinoma In Situ

Ductal carcinoma in situ  
Lobular carcinoma in situ

### Ductal Carcinoma In Situ (DCIS)

DCIS is characterized by neoplastic proliferation of epithelial cells that is confined to the ductal-lobular system of the breast without evidence of invasion through the basement membrane into the surrounding stroma. DCIS encompasses a heterogeneous group of lesions that differ with regard to their presentation, histopathological features, biological markers, and risk for progression to invasive cancer [1]. In areas where mammographic breast screening is not performed, DCIS constitutes approximately 5 % of breast cancers, but within screening programs, it comprises approximately 20–25 % of these tumors [2]. Approximately 10–20 % of DCIS cases are bilateral.


These tumors are traditionally classified according to their architecture and are divided into comedo and non-comedo subtypes. The non-comedo subtype is further subdivided into solid, cribriform, micropapillary, and papillary types. Recent grading systems use the nuclear grade alone or in combination with necrosis [3].

DCIS is generally divided into three grades according to the nuclear features [3, 4]:

*High nuclear grade DCIS:* The tumor is composed of large, pleomorphic cells, often with prominent nucleoli. The nuclei are more than 2.5 times the diameter of red blood cells. Chromatin is coarse and clumped, and its distribution is irregular. Comedo necrosis with or without microcalcification is frequent but not necessary. Polarization toward the luminal surface is usually lost. Mitoses may be frequent (Fig. 13.1).


*Low nuclear grade DCIS:* The cells are small, monotonous cells that form arcades, micropapillae, and cribriform and solid patterns. Their nuclei are uniform and 1.5–2.5 times the size of normal red blood cells. Nuclei are usually but not invariably small [1]. The chromatin is finely dispersed. Nucleoli are inconspicuous. Mitoses are sparse, and the cells are polarized toward the luminal spaces (Fig. 13.2).

*Intermediate grade DCIS:* When the lesion cannot be assigned easily to the high- or low-grade DCIS categories, it is diagnosed as intermediate grade. The features mentioned above are usually intermediate between low and high nuclear grade DCIS.



**Fig. 13.1** DCIS of high nuclear grade. Proliferation of pleomorphic cells in two spaces in the center, and comedo necrosis is apparent

**Fig. 13.2** DCIS of low nuclear grade. Proliferation of monotonous, uniform cells forming a micropapillary and cribriform architecture



In the presence of foci of different grades, the case should be graded according to the highest grade.

*Pathology reporting for DCIS:*

A pathology report for DCIS should include the following [3–6]:

- *Size/extent of the lesion:* Precisely measuring the extent of DCIS is often not possible. The volume of the breast tissue that is involved by DCIS is estimated by the pathologist based on the preferred sampling method. Mammographic correlation is also necessary, and this information should be provided by the clinician.
- *Nuclear grade*
- *The presence or absence of necrosis and its type:* The type of necrosis can be classified as punctate or comedo. Comedo necrosis is the classical central necrosis in the duct lumina with karyorrhectic debris. This form of necrosis is associated with mammographic microcalcifications. Punctate necrosis present small foci or single-cell necrosis that is indistinct at low magnification.
- *Architectural pattern(s):* The comedo, solid, cribriform, micropapillary, papillary patterns are considered in the traditional classification schemes.

- *Cell polarization:* The presence or loss of polarization toward the luminal surfaces is considered in some grading schemes [7].
- *Location of microcalcifications:* When microcalcifications are present, their localization should be reported (in DCIS alone, in benign breast tissue, or in both). This information provides the correlation with mammographic findings.
- *Surgical margin status:* The surgeon provides the orientation using sutures or clips. In the presence of microcalcifications, specimen mammography should be provided. The specimen should be linked by the pathologist, and sampling is performed in any of several methods, depending upon the pathologist's choice.

Necrosis and polarization appear to have secondary importance compared with nuclear grade.

In our pathology practice, sampling the whole lesion is mandatory to exclude any minute foci of invasion before giving the diagnosis of DCIS.

**Differential Diagnosis**

- *Lobular carcinoma in situ (LCIS) versus DCIS:* DCIS with a solid pattern must sometimes be distinguished from LCIS. This distinction may be difficult on a morphological

basis. Immunohistochemically, the presence of E-cadherin may be helpful in categorizing the individual case in favor of DCIS.


- *Usual ductal hyperplasia (UDH) and atypical ductal hyperplasia (ADH) versus low-grade DCIS:* The difference between ADH and low-grade DCIS lies in the extent of the involvement of the duct system. In ADH, there is partial involvement of multiple spaces, whereas low-grade DCIS involves the entire duct space. It is proposed by Page and Tavassoli that in order to describe a lesion as low-grade DCIS, complete involvement should include at least two spaces or be larger than 2 mm [8, 9]. Lesions occupying fewer than two spaces or a total area smaller than 2 mm are called ADH (Fig. 13.3). This distinction is imperfect, and the level of concordance and consistency in their diagnosis is low [3].
- *Foci of microinvasion:* DCIS extending into a terminal ductal-lobular unit (TDLU) or an adjacent benign proliferative lesion such as sclerosing adenosis (SA) or radial scar may create the impression of microinvasion. The absence of invasive foci can be confirmed by immunohistochemically demonstrating the presence of myoepithelial cells (using antibodies against smooth muscle actin, p63, CD10, calponin, etc.) or basement membrane

(using antibodies against collagen type IV or laminin).

- *Invasive cribriform carcinoma (ICC):* This unusual invasive carcinoma can sometimes be mistaken as DCIS of cribriform type. The diagnosis in ICC is based on the recognition of the infiltrative pattern and the absence of myoepithelial markers by immunohistochemistry.
- *Receptor status:* Most cases of DCIS are positive for estrogen receptor (ER). Positivity (defined as  $\geq 1\%$  of tumor cells) is observed in 70–85 % of the cases [3, 4]. Expression correlates with the grade of DCIS. Almost all cases of ER-negative DCIS are of high nuclear grade. Progesterone receptor (PR) expression is lower than ER expression.

### Columnar Cell Lesions and Flat Epithelial Atypia

Lesions lacking intraluminal proliferation have long been recognized, and they have been given a variety of names with regard to cell morphology and the presence or absence of atypia. In 2003, Schnitt et al. classified these lesions as follows [10]:



**Fig. 13.3** Atypical ductal hyperplasia (ADH). Architectural and cellular features of a low-grade DCIS but covering less than 2 mm, in a terminal ductal-lobular unit (TDLU)



Columnar cell change (CCC)  
 Columnar cell hyperplasia (CCH)  
 CCC with atypia  
 CCH with atypia

Now, a simplified terminology combining the latter two under the term flat epithelial atypia (FEA) has become widely used [3, 11]:

CCC  
 CCH  
 FEA


Here, CCC and CCH with atypia are classified under the category FEA.

*CCC and CCH* are lesions in the TDLU that are characterized by enlarged, variably dilated acini lined by columnar epithelial cells [3]. These lesions are microscopic in size and are increasingly detected because of mammographic microcalcifications. The cells have ovoid nuclei that are oriented perpendicularly to the basement membrane. They have evenly dispersed, fine chromatin and inconspicuous nucleoli. The lesions are frequently associated with intraluminal secretion and microcalcification. Lesions in which the epithelial lining is composed of one or two cell layers are categorized as CCC. If there is cellular stratification of more than two layers and

piling up of several layers, the term CCH is used. Columnar cell lesions are associated with very low risk for subsequent development of invasive breast cancer, and these lesions do not increase this risk independent of the concurrent proliferative changes [12].

*FEA*: Lesions exhibiting cellular atypia in addition to the architectural patterns described for CCC and CCH are categorized as FEA. FEA is characterized by the replacement of native epithelial cells with one to several layers of monotonous, cuboidal to columnar cells with low-grade cytologic atypia. The cells often have apical snouts. Well-developed bridges or arcades are absent (Fig. 13.4).

FEA corresponds to Azzopardi's "clinging carcinoma, monomorphic type." Other flat proliferations corresponding to the "clinging carcinoma, high-grade polymorphous type" should be categorized as DCIS. If a lesion with low-grade nuclear features has well-developed bridges, arcades, or bulbous micropapillae, it should be diagnosed ADH or low-grade DCIS depending on the quantity of the lesion (see above). The risk of subsequent invasive breast cancer in FEA is low and is substantially lower than the risk associated with established forms of ADH [3]. FEA is often associated with ADH, low-grade DCIS, lobular neoplasia (LN), and tubular carcinoma (TC).



**Fig. 13.4** Flat epithelial atypia. The duct is lined by one to three layers of cells with low-grade atypia and apical snouts. There is no bridge or arcade formation

Unlike the normal breast and UDH, where ER and PR immunostaining is heterogeneous and limited to approximately 10–15 % of the cells, CCL and FEA reveal diffuse and homogenous staining in all lesional cells. Most of the cells show immunostaining for low-molecular-weight cytokeratins and are negative for CK5/6.

### Lobular Neoplasia: Lobular Carcinoma In Situ (LCIS)

The entire spectrum of atypical epithelial lesions originating in the TDLU of the breast, characterized by the proliferation of generally small, non-cohesive cells, is called LN. The terms atypical lobular hyperplasia (ALH) and LCIS reflect the extent of the lesion. In both types of lesions, proliferating cells are cuboidal or polygonal monotonous and poorly cohesive cells with clear or light cytoplasm. Pagetoid spread of these cells between the surface epithelial cells and basement membrane is a common finding. When more than half of the acini of a lobular unit are distended and distorted, the lesion is called LCIS (Fig. 13.5). Lesser involvement with cells showing the same characteristics is called ALH [8] (Fig. 13.6). The differentiation of these two is sometimes subjective. Thus, the term LN, which does not differentiate between LCIS and ALH, was introduced [3].

**Pleomorphic LCIS** The cells are markedly pleomorphic with large nuclei. Central necrosis and microcalcifications may be present [13] (Fig. 13.7). These are usually mammographically detected lesions.


**LCIS with Comedo Necrosis** In addition to the classical, small, monotonous cells of LCIS, there is comedo-type necrosis in the central portion of spaces [14].

The morphological distinction from solid-type DCIS is discussed above.

LN and LCIS are almost uniformly positive for ER and PR and negative for E-cadherin. Classical LCIS and LCIS with comedo necrosis are negative for H2 and p53 and have a low Ki-67 index. However, pleomorphic LCIS may have H2 and p53 overexpression and moderate to high Ki-67 [15].

LN is classically accepted as a risk indicator of breast cancer development for both breasts; however, recent, carefully conducted cohort studies suggest that the risk is higher in the ipsilateral breast (68 % versus 24 %) [16]. The clinical and molecular evidence available suggests that ALH and LCIS are clonal and neoplastic and that these lesions are both risk indicators and non-obligate precursors of breast cancer [17].


The florid form of LCIS is more frequently associated with an invasive component than the nonflorid form of it (87 % versus 73 %, respectively).




**Fig. 13.5** Lobular carcinoma in situ (LCIS). Proliferation of small, uniform, discohesive cells that completely fill and distend the TDLUs



**Fig. 13.6** Atypical lobular hyperplasia (ALH). Proliferation of small, uniform cells that slightly distend the acini of the lobule



**Fig. 13.7** LCIS of pleomorphic type with necrosis. The cells were E-cadherin negative. There is necrosis in the center of one of the spaces



respectively). The invasive component is lobular in 100 % of florid LCIS lesions but only in 82 % of nonflorid LCIS lesions [18]. Recent evidence also suggests that the florid form of LCIS is genetically more advanced compared with the indolent phenotype of classic LCIS [19]. This difference may explain the greater frequency of concurrent invasive carcinoma in florid LCIS compared with classic LCIS [19].

### Microinvasive Carcinoma

This lesion is characterized by one or more clearly separate microscopic foci of infiltration of tumor cells into the mammary stroma, each less than or equal to 1 mm in size, and most commonly seen in a background of high-grade DCIS [3, 20].

Microinvasive carcinoma is most commonly observed in a background of extensive high-grade

DCIS with prominent inflammatory infiltration [3] accompanied by stromal edema and desmoplasia. This entity is commonly overdiagnosed. Central consultation usually downgrades the lesion. Even in cases initially suspected or diagnosed as microinvasion, subsequent review downgrades the diagnosis in 80 % of cases [21]. Differentiation from DCIS is described above. These foci are commonly lost in serial sections performed for immunostaining.

The prognosis is not clearly different from patients with DCIS of equivalent grade.

## Invasive Carcinomas

Invasive carcinomas can broadly be divided into two categories, invasive carcinoma of no special type (NST) and special subtypes [3]. Invasive carcinoma NST and invasive lobular carcinoma (ILC) constitute the major types of breast carcinoma. Cytoarchitectural and spread patterns of some carcinomas are distinctive enough to be recognized as special subtypes especially when associated with a particular behavior [22].

According to the recent WHO classification, invasive breast carcinomas are classified as indicated in Table 13.1 [3].

### Invasive Carcinoma of No Special Type (NST)

This carcinoma is the most common type of invasive breast cancer and represents up to 75 % of the cases in published series. Terms such as infiltrating ductal carcinoma and invasive ductal carcinoma, not otherwise specified (NOS) are also used. There is a wide range in the frequency of invasive carcinoma NST because there are no strict criteria for inclusion of these tumors in the special types and because different centers have varying attitudes in categorizing cases with different amounts of NST and special types. A tumor should be called invasive ductal carcinoma (IDC) NST if it cannot be categorized as one of

**Table 13.1** WHO classification of breast cancer

Invasive carcinoma of no special type
Pleomorphic carcinoma
Carcinoma with osteoclast-like stromal giant cells
Carcinoma with choriocarcinomatous features
Carcinoma with melanocytic features
Special types:
Invasive lobular carcinoma
Classical lobular carcinoma
Solid lobular carcinoma
Alveolar lobular carcinoma
Pleomorphic lobular carcinoma
Tubulolobular carcinoma
Mixed lobular carcinoma
Tubular carcinoma
Cribriform carcinoma
Mucinous carcinoma
Carcinoma with medullary features
Medullary carcinoma
Atypical medullary carcinoma
Invasive carcinoma NST with medullary features
Carcinoma with apocrine differentiation
Carcinoma with signet ring cell differentiation
Invasive micropapillary carcinoma
Metaplastic carcinoma of no special type
Low-grade adenosquamous carcinoma
Fibromatosis-like metaplastic carcinoma
Squamous cell carcinoma
Spindle cell carcinoma
Metaplastic carcinoma with mesenchymal differentiation
Chondroid differentiation
Osseous differentiation
Other types of mesenchymal differentiation
Mixed metaplastic carcinoma
Myoepithelial carcinoma
<i>Rare types</i>
Carcinoma with neuroendocrine features
Neuroendocrine tumor, well differentiated
Neuroendocrine carcinoma, poorly differentiated (small-cell carcinoma)
Secretory carcinoma
Invasive papillary carcinoma
Acinic cell carcinoma
Mucoepidermoid carcinoma
Polymorphous carcinoma
Oncocytic carcinoma
Lipid-rich carcinoma

**Table 13.1** (continued)

Glycogen-rich clear cell carcinoma
Sebaceous carcinoma
Salivary gland/skin adnexal-type tumors
Cylindroma
Clear cell hidradenoma
Epithelial-myoepithelial tumors
Pleomorphic adenoma
Adenomyoepithelioma
Adenomyoepithelioma with carcinoma
Adenoid cystic carcinoma

Modified from Lakhani et al. [3]

the special or rare types. There is great variation in their appearance.

**Gross Features** IDC NST has no specific gross features. It also shows a great variation in size, ranging from a few millimeters to huge masses. In typical cases, these tumors have irregular, stellate borders (Fig. 13.8). They have a firm consistency, and their cut surface is generally gray-white with a gritty sensation. Less frequently, the tumor may have a nodular configuration with circumscribed margins and relatively softer consistency.

**Microscopic Features** The tumor cells are arranged in sheets, clusters, cords, trabeculae, and glands/tubules or sometimes in a solid pattern with no or little intervening stroma. Cellular features also show great variability. Nuclei may be uniform and regular or highly pleomorphic with very prominent and multiple nucleoli. Mitotic activity is also highly variable (Fig. 13.9). IDC NST may have histopathological characteristics of special types, but less than 50 % of the tumor will fall into this category. In other words, in IDC NST, at least 50 % of the tumor should be composed of a nonspecialized type. The tumor stroma may be abundant.

When a proportion of specialized histopathological forms accompany the IDC NST, these carcinomas are described as “mixed type” [3].

Pleomorphic carcinoma, carcinoma with osteoclast-like stromal giant cells, carcinoma with choriocarcinomatous features, and carcinoma with melanocytic features are not recognized as

distinct special types but as variants of IDC NST [3]. The latter two are exceptionally rare.


## Pleomorphic Carcinoma

*Pleomorphic carcinoma* is characterized by the proliferation of bizarre, highly anaplastic, and sometimes multinucleated cells. Approximately one-third of the cases have a metaplastic spindle cell component [23, 24]. Spindle cell components and tumor size (>5 cm) are associated with poor clinical outcome [24].

This prognostically unfavorable tumor represents the extreme end of the morphological spectrum of grade III infiltrating ductal carcinoma [23].

## Carcinoma with Osteoclast-Like Stromal Giant Cells


The distinctive feature is the presence of osteoclastic giant cells (OGCs). Grossly, they have a striking red-brown cut section with a hemorrhagic appearance, especially in cases with plenty of



**Fig. 13.8** Typical gross appearance of an invasive carcinoma located in the center of the specimen. Grayish-white tumor with ill-defined borders. Fibrocystic changes below the tumor and fatty appearance in the rest of the excision



**Fig. 13.9** Invasive ductal carcinoma NOS. Grade 3 carcinoma with sheets of cells with pleomorphic nuclei and frequent mitosis



OGCs. These cells are generally associated with a fibroblastic and hypervascular stroma which contains inflammatory cells, erythrocytes, and hemosiderin as the evidence of recent and past hemorrhages. OGCs are usually close to the edges of carcinomatous glands or in the intervening stroma, and they are sometimes present in glandular lumens. They may appear as though they are fused with the glandular component and may be difficult to discern (Fig. 13.10). Associated carcinomas are mostly well to moderately differentiated, showing a relatively more common cribriform pattern. OGCs are positive for CD 68, acid phosphatase, and lysozyme but negative for cytokeratin and alkaline phosphatase. They are negative for ER, PR, cytokeratin, epithelial membrane antigen (EMA), actin, and S-100 protein [25–27]. This immunohistochemical profile, along with the absence of any epithelial features in ultrastructural examination, supports the histiocytic origin of these cells [27]. They also express the osteoclast markers MMP-9, TRAP, and cathepsin K, and they appear to form in response to the specific hypervascular stroma which secretes cytokines such as VEGF and of MMP-12 [28, 29].

Axillary lymph node involvement has been reported in one-third of the cases [26, 27]. Distant metastasis to a variety of sites has also been reported [26, 27]. The 5-year survival rate is approximately 70 %, which is similar to or

slightly better than in patients with ordinary invasive ductal carcinoma [3].

### Invasive Lobular Carcinoma

ILC is a carcinoma composed of non-cohesive cells that are individually dispersed or arranged in a single-file linear pattern in a fibrous stroma [3]. ILC represents the 5–15 % of invasive breast carcinomas [3]. In most series, its incidence is approximately 10 % [27].


ILC frequently presents as a mass with irregular borders that sometimes cannot be defined macroscopically, and the breast tissue appears normal with only a firm consistency by palpation [26]. The size ranges from occult, microscopic lesions to tumors that diffusely involve the entire breast [26]. Sometimes ILC forms numerous, fine, hard nodules that grossly and microscopically mimic.

The incidence of synchronous or metachronous bilateral carcinoma in ILCs is almost twice that observed in IDCs [30, 31].

### Classical ILC

ILC is characterized by the proliferation of small, uniform cells that lack cohesion and are dispersed individually in a fibrous stroma or arranged in

**Fig. 13.10** Carcinoma with osteoclast-like giant cells. The cribriform architecture is formed by the tumor cells. Multinucleated giant cells resemble osteoclasts in the fibroblastic stroma



linear cords. These cords usually present a concentric pattern around nonneoplastic ducts, forming the “targetoid pattern” (Fig. 13.11). The tumor cells are bland or monotonous and have round to ovoid nuclei. Mitoses are uncommon.

ILC has some histologic variants that differ from the classical type in terms of their histological growth or cytological patterns but still lack cellular cohesion.

### Solid Variant

Uniform, small, and non-cohesive cells are arranged in sheets. There is little stroma. These tumors are often more pleomorphic and have a higher mitotic rate compared with the classical ILC [30, 32].

### Alveolar Variant

*Alveolar variant* exhibits small, globular aggregates of 20 or more cells [33].

### Tubulolobular Variant

*Tubulolobular variant ILC* exhibits small tubules and cords of neoplastic cells in a lobular configuration reminiscent of ILC [34]. It has

many features that are intermediate between tubular carcinoma and classical ILC [34].

This tumor is morphologically different from tumors showing a mixture of tubular carcinoma and classical ILC, which should be categorized in the mixed category.

### Pleomorphic Variant

*Pleomorphic ILC* exhibits the growth pattern of classical ILC but a greater degree of cellular atypia and pleomorphism and a higher mitotic rate than classical ILC. These cells retain their lobular characteristics with their single-file and/or targetoid arrangement and non-cohesive appearance. LCIS is present in 45–60 % of the cases [13, 35, 36] and is frequently of the pleomorphic type [36]. Pleomorphic ILC may show apocrine [13] or histiocytoid [37, 38] differentiation and may be composed of signet ring cells [3].

### Mixed Type

These cases exhibit mixtures of the abovementioned variants and were described by Dixon [30] as “none of these patterns are prominent.” Lobular differentiation accompanying IDC NST



is observed in approximately 5 % of invasive breast cancers [3].

ILC is almost invariably ER positive. PR positivity is present in approximately 70–80 % of cases. Her-2 positivity with immunohistochemistry or in situ hybridization is very rare and is generally limited to pleomorphic ILC. Immunohistochemically, E-cadherin is absent or reduced in ILC compared with IDC. However, a subset of ILCs express E-cadherin, ranging from 10 to 16 % of ILCs [39, 40], and this subset is described as being aberrant without any significance or any correlation with the known prognostic parameters [40, 41].

Most ILCs also show loss of membrane-specific catenin immunoreactivity in parallel to E-cadherin loss [39] and mislocalization of catenin p120 in the cytoplasm [42].

In general, ILCs have more favorable prognostic features than IDC NST. A higher frequency of ILC was placed in the good Nottingham Prognostic Index group (40 % compared with 21 % for IDC) [43], and it has a better or similar outcome in the short-term period (first 6–10 years).

However, the long-term outcome for ILC is worse than for IDC NST [43, 44].

A more favorable outcome is reported for the classical type than the pleomorphic type [26]. The differences in outcome between variant forms and classical ILC have not been statistically significant


[26]. Rakha et al. found that survival in patients with pleomorphic lobular carcinoma was associated with mitotic score but not with nuclear pleomorphism [45].

Distinctive patterns of metastases are associated with ILC. ILC shows a higher frequency of metastases in the intra-abdominal serosal surfaces and retroperitoneum, leptomeninges, gastrointestinal tract, and gynecologic organs and a lower frequency of pulmonary metastases [3, 25–27].

### Tubular Carcinoma

Tubular carcinoma (TC) is a low-grade (grade I) carcinoma with a particularly favorable prognosis. It is composed of well-differentiated tubular structures lined by a single layer of cells and has open lumina. Pure TC accounts for approximately 2 % of invasive breast cancers. Its frequency is higher in populations where screening mammography is used. TC is more likely to be smaller lesions with less frequent nodal involvement and a better outcome than IDC NST.

TC often presents as an ill-defined, gray-white, firm to hard, stellate mass with an average size of 1.3 cm (0.2–5 cm). The cut surfaces frequently show elastotic, yellow streaks.



**Fig. 13.11** Invasive lobular carcinoma. Small, discohesive tumor cells form the single-file pattern around the intact duct


Microscopically, the tubules are haphazardly arranged in a typical desmoplastic stroma. The lumina of the tubules are oval or rounded with angulated ends. The single cells lining these tubules have little nuclear pleomorphism with inconspicuous nucleoli, and they exhibit very few mitoses. The cells may have apical snouts, but this characteristic has no diagnostic significance (Fig. 13.12). Myoepithelial cell layer and basal membrane are lacking in contrast to the nonneoplastic proliferations. TC occurs in association with FEA and low-grade DCIS.

There is a lack of consensus regarding the proportion of tubules necessary to establish a diagnosis of TC. In several studies, the threshold for tubule formation was set between 75 and 100 %. However, a cutoff of 90 % is more widely accepted [46]. Patients diagnosed with TC with this cutoff and small lesions have the same overall survival as the age-matched general population [46, 47].

Tubule formation in less than 90 % of the tumor should be regarded as mixed type. One exception that should be considered is the cribriform pattern. In the presence of invasive cribriform carcinoma (ICC) intermingled with TC, these areas are also regarded as tubule formation.

### Differential Diagnosis

- *Microglandular adenosis (MGA)*: The TC is sometimes composed of small, round tubules of relatively uniform caliber that are irregularly dispersed in a fibrofatty stroma with a resemblance to MGA. Glands in MGA are more rounded and regular and contain secretory material [3, 27]. The myoepithelium is lacking in both types of lesions, and immunostaining reveals no staining for calponin, p63, CD10, or cytokeratin 5. The basement membrane is lacking in TC, which can be demonstrated around the glands of MGA by periodic acid-Schiff (PAS) staining and immunostaining for collagen IV and laminin [3, 25–27]. Epithelial membrane antigen (EMA), which is present in TC, is absent in MGA [27].
- *Sclerosing adenosis (SA)*: SA is a lobulocentric proliferation with a compressed and distorted appearance of the tubular structures. Myoepithelial cells and basement membrane are always present in SA and can be highlighted with the immunostaining described above. TC does not have a lobulocentric growth pattern and does not contain myoepithelial cells or a basement membrane.
- *Complex sclerosing lesion (radial scar)*: The central fibroelastotic core of this lesion may have



**Fig. 13.12** Tubular carcinoma. Tumor cells form tubules, some of which have microcalcifications in their lumens

a few, distorted, entrapped, pseudoinfiltrative glands, creating diagnostic difficulty through its resemblance to TC. The glands at the periphery of the core are hyperplastic and dilated. This zoning phenomenon is lacking in TC. The glands in CSL also contain myoepithelial cells and a basement membrane.

Women with “pure” TC have an excellent prognosis. The frequency of axillary lymph node metastasis is approximately 10 %. TC has a better prognosis than grade I IDC or tubular mixed carcinomas, independent of other prognostic factors [46, 47]. In a follow-up of 127 months (4–217 months), recurrent disease was found in 13.2 % of patients with TCs, with no cancer-specific deaths, compared with patients with grade I IDCs, in which the recurrent disease rate was 29.4 % and the cancer-specific death rate was 9 % [46].

---

## Invasive Cribriform Carcinoma

ICC is a low-grade carcinoma with excellent prognosis in which the majority of the invasive component shows a cribriform pattern of growth, similar to intraductal cribriform carcinoma. Pure ICC consists of an invasive cribriform pattern in more than 90 % of the tumor [48, 49]. The tumor cells, which have mild to moderate pleomorphism, are arranged to form cribriform spaces (sievelike pattern). Mitoses are rare. There are no specific gross features for this tumor.

## Differential Diagnosis

*Adenoid cystic carcinoma (ACC)*: ICC most closely resembles ACC. ICC is composed of one cell type and lacks the basal-myoepithelial type. There is mucinous material in the cribriform spaces, and tumor cells are diffusely positive for ER. In ACC, there are two cell types, basal-myoepithelial and luminal, and secretory and basement membrane-like material is present in the glandular spaces; ACC also shows a triple-negative immunoprofile.

*Cribriform DCIS*: ICC has a more irregular and angular cribriform pattern with more haphazard distribution compared with cribriform DCIS. Cribriform DCIS has a myoepithelial cell layer around the cribriform structures.

In cribriform DCIS, 100 % of cases are ER positive, 69 % of cases are PR positive [50], and HER2 expression is absent [27].

The prognosis of ICC is favorable [49] and similar to TC [48]. The 10-year overall survival is 90–100 % [49, 50].

---

## Carcinoma with Medullary Features

These tumors exhibit some or all of the following features: a circumscribed or pushing border, syncytial growth pattern, cells with high-grade nuclei, and prominent lymphoid infiltration. According to the 2003 WHO classification [51], tumors that fulfill all these criteria are called medullary carcinoma (MC), and tumors that fulfill most but not all of these criteria are called atypical MC. In the more recent classification, there has been an attempt to categorize these tumors into three groups, under the heading “Carcinomas with medullary features,” as follows [3]:

- MC
- Atypical MC
- IDC NST with medullary features

The criteria that distinguish these groups are vague and have poor interobserver reproducibility. Distinction between the latter two groups is particularly difficult. In our institutional practice, we prefer reserving the term MC for tumors exhibiting all of the features described above, using very strict criteria, and calling the tumors exhibiting some of these features atypical MC.

Despite poor clinicopathologic features, patients with medullary histology demonstrate favorable long-term distant relapse-free survival compared with patients with IDC NST. Local control rates of MC and IDC are comparable [52]. In a retrospective study of 165 cases of basal-like

carcinomas, the Nottingham group has found that prominent inflammation and anastomosing sheets in at least 30 % of the tumor were associated with a better prognosis in a univariate analysis [53]. The combination of these two features was present in 17 % of tumors and was an independent prognostic factor in a multivariate analysis. They also proposed a simplified definition of medullary-like type based on these two features [53].

## Mucinous Carcinoma


Mucinous carcinoma is characterized by the production of extracellular and/or intracellular mucinous material. Clusters of generally small and uniform cells are seen as floating in large amounts of mucin. A lesion is called pure mucinous carcinoma if the mucinous component constitutes more than 90 % of the lesion [54]. Mucinous carcinoma is also observed as part of a mixed carcinoma with IDC NST. The axillary lymph nodes are rarely involved.

Gross examination of mucinous carcinomas reveals a circumscribed, gelatinous mass with pushing margins and soft consistency. The cut surface has a glistening appearance. Confluent hemorrhagic areas are frequent [27] (Fig. 13.13). The tumor size ranges from 0.5 to 20 cm. Despite having large diameters, axillary nodal involvement is infrequent.

Microscopically, there are clusters of tumor cells floating in mucin lakes separated by delicate fibrovascular septae. The clusters are variable in size. Nuclear atypia is generally low. Mucinous carcinoma can be divided into two categories, types A and B [55]:

*Type A mucinous carcinoma:* This is the classical or non-endocrine variant and is characterized by larger quantities of mucin. Mucin is always extracellular [27] (Fig. 13.14).

*Type B mucinous carcinoma:* This type is more cellular with large clusters and has frequent neuroendocrine differentiation. Intracytoplasmic mucin is abundant in type B lesions (Fig. 13.15).



**Fig. 13.13** Mucinous carcinoma. The tumor has well-defined borders and a lobulated appearance with a glistening and partially hemorrhagic cut surface

Mucinous carcinoma is usually positive for ER and PR and negative for HER2.

Type AB mucinous carcinoma constitutes 20 % of the cases and is an intermediate lesion that has features of both types.

The most important entity in the differential diagnosis is the “mucocele-like lesion” [54]. Mucinous carcinoma should also be distinguished from myxoid fibroadenomas, especially in fine-needle aspiration biopsies.


Mucinous carcinomas have a favorable outcome [47]. In a follow-up series of 11,400 cases of pure mucinous carcinoma, the 5-, 10-, 15-, and 20-year survival rates were 94, 89, 85, and 81 %, respectively [56]. Nodal involvement was associated with significant disease-free survival and overall survival [47]. The separation of cases as types A and B has no clinical significance.

## Carcinomas with Signet Ring Cell Differentiation


Cells with signet ring cell differentiation have abundant mucin in their cytoplasm, which pushes the nucleus to one side, creating the typical signet ring cell appearance (Fig. 13.16). Carcinomas with extensive signet ring cell differentiation are



**Fig. 13.14** Mucinous carcinoma type A with a huge amount of extracellular mucin and low-grade cellular atypia



**Fig. 13.15** Mucinous carcinoma type B with a more cellular appearance and a higher grade of cellular atypia



rare. Focal signet ring cell differentiation is more commonly observed.

Prominent signet ring cell differentiation is most common in ILC.

Pathologists should sometimes distinguish these cases from gastrointestinal metastasis. The presence of an in situ component is a sign in favor of primary breast cancer. In difficult cases, steroid receptor expression and antibodies specific to breast carcinoma such as GCDFP or mammaglobin are helpful.

The prognostic importance of signet ring cell differentiation is uncertain [3].

### **Carcinoma with Apocrine Differentiation**

This class includes any invasive carcinoma having cells with cytological features of apocrine differentiation. These cells have abundant, eosinophilic, granular cytoplasm and large nuclei with



prominent nucleoli. There is a transition to cells with foamy cytoplasm resembling sebaceous cells, which may sometimes dominate the histology. Focal apocrine differentiation is not very rare. However, a tumor is called “pure” apocrine carcinoma if 90 % of the lesion is composed of these cells. ER and PR expression is usually negative. Androgen receptor (AR) positivity is encountered in more than 70 % of apocrine carcinomas. GCDFP-15 is characteristic but not specific for apocrine cells [27, 57]. From a practical perspective, we do not call tumors pure apocrine carcinoma if there is ER or PR expression. AR expression in ER-/PR-/HER2+ tumors, which commonly show apocrine differentiation, and a subset of triple-negative apocrine tumors suggests that these tumors together form a molecular apocrine group [58].

A recent study with long-term follow-up revealed that patients with pure apocrine carcinomas (negative for ER and PR and positive for AR) have a shorter disease-free survival than patients with IDC NST and apocrine-like IDC (ER or PR positive and AR negative) [59].

### Invasive Micropapillary Carcinoma (IMPC)

This type of tumor which was first described by Tavassoli and her colleagues [60] is now regarded as a distinct entity. It accounts for 0.9–1.7 % of


invasive breast carcinomas when occurring in pure form and up to 7.6 % when admixed with other types of mammary carcinoma [61]. Most of the patients present with a palpable mass [61].

The tumor is composed of small, hollow, or morula-like clusters of tumor cells that lack fibrovascular cores and surrounded by clear stromal spaces (Fig. 13.17). The “reverse polarity” of cancer cells is typical and can be facilitated by immunohistochemical demonstration of MUC1 in the stroma-facing surface. This reaction may also be used in differentiating these spaces from lymphovascular invasion or retraction artifacts [3, 61]. The presence of an in situ component is helpful in excluding the rare cases of metastatic ovarian serous papillary carcinoma to the breast.

Most cases are grade 2 or 3 carcinomas, and the majority of them are ER and PR positive. HER2 overexpression is present in less than 10–35 % of cases [3].


High-resolution microarray comparative genomic hybridization has revealed that high cyclin D1 expression, high proliferation rates, and MYC (8q24) amplification are significantly associated with IMPCs [62].

IMPCs present more frequently with lymphovascular invasion and lymph node metastasis when compared with the IDC NST [63]. The Ki-67 proliferative index is significantly higher in IMP carcinomas with p63 expression (nuclear or cytoplasmic)



**Fig. 13.16** Carcinoma with signet ring cell differentiation. Nuclei of most of the cells are pushed to one side by mucous, creating the “signet ring” appearance

**Fig. 13.17** Invasive micropapillary carcinoma. Lobular configuration of the invasive tumor. The clusters of tumor cells are surrounded by clear spaces. DCIS is evident on the upper right, and microcalcification is visible in the lower left



than in those without and is also higher in cases with lymph node metastasis than in cases without [64]. However, the association of this histology with survival remains unclear. In a recent series of 49 patients, IMPC histology did not add any independent information to the risk of locoregional or distant relapse or to overall survival [65].

### Metaplastic Carcinoma

Metaplastic carcinoma encompasses a group of neoplasms that are characterized by the differentiation of the neoplastic epithelium into squamous and/or mesenchymal-looking elements, including but not restricted to spindle, chondroid, osseous, and rhabdomyoid cells [3]. The tumor may be entirely composed of metaplastic elements or may include a mixture of carcinoma and metaplastic elements. Its incidence is less than 1 % [66, 67]. It is usually diagnosed as T2 disease, and the mean size is 3.4–4.4 cm [66].

These tumors can present either as a circumscribed nodule or as a mass with indistinct borders. Cystic changes can occur, especially in cases that are accompanied by squamous cell carcinoma (SCC).

The recent WHO classification [3] categorizes metaplastic carcinomas in a descriptive manner.

### Low-Grade Adenosquamous Carcinoma (LGASC)


This tumor is similar to the infiltrating syringomatous tumors of the salivary glands and microcystic adnexal carcinomas of the skin of the lip [68]. The patients present with a palpable mass [69], and grossly, the tumors are smaller than other forms of metaplastic carcinoma [26]. They have a hard consistency and ill-defined borders [69]. Microscopically, there are well-defined tubules and glands in a spindle cell background. Squamous differentiation is seen as solid nests, syringoma-like areas, and isolated inconspicuous foci in glandular structures and solid cords. Squamous differentiation may be extensive with large keratinizing cyst formations. In our experience, this rare tumor is an underdiagnosed entity and therefore may be left untreated; during their long evolution, they recur and metastasize.

### Fibromatosis-Like Metaplastic Carcinoma

This tumor is characterized by bland spindle cells having slender nuclei with tapered ends. Nuclear atypia is mild or absent. These cells are arranged in wavy, interlacing fascicles. Focal squamous differentiation is observed. Because of the bland appearance of tumor cells, this tumor may be



**Fig. 13.18** Metaplastic carcinoma. The tumor has squamous (*left*) and chondromyxoid components



underdiagnosed as benign [70]. The tumor is always positive for keratins [70] and p63 [3].

### Squamous Cell Carcinoma

Grossly, it is often a cystic lesion [3]. The cavity is lined by squamous cells, often with bland nuclear features. The infiltrating squamous cells form sheets and nests with varying degrees of differentiation. Combination of patterns with transition to spindle cells or to less differentiated forms may occur. A rare variant is the acantholytic type of SCC, which may be confused with angiosarcoma. The irregular spaces lined by atypical squamous cells create a pseudoglandular and/or pseudovascular appearance. SCC can be easily discriminated by the positive staining of these cells for keratin and the absence of FVIII and CD34 [71].

An origin from the overlying skin should also be excluded.

SCC may be mixed with an invasive ductal carcinoma NST. Focal squamous differentiation can also be found in IDC NST and may accompany carcinomas with medullary features.

### Spindle Cell Carcinoma

This tumor is characterized by the pseudosarcomatous growth pattern of its neoplastic spindle

cells. The distinction between spindle cell carcinoma and primary sarcomas of the breast, including fibrosarcoma and malignant fibrous histiocytoma, may be problematic. The presence of focal squamous differentiation and small clusters of spindle cells with more epithelioid histology is the clues for differentiating these lesions. Epithelial differentiation can be demonstrated by immunohistochemistry using a panel of antibodies (high-molecular-weight cytokeratins). P63 staining is also very common [72].

### Metaplastic Carcinoma with Mesenchymal Differentiation

These tumors display an admixture of carcinomatous and mesenchymal elements. Mesenchymal components include chondroid, osseous, and rhabdomyoid elements with varying degrees of differentiation.

Metaplastic carcinomas often contain a mixture of different elements (Fig. 13.18).

**Matrix-Producing Carcinoma** This is a subgroup of metaplastic carcinomas that show an abrupt transition from epithelial to mesenchymal elements without intervening spindle cells.

Over 90 % of metaplastic carcinomas are triple-negative cancers and express keratin 5/6, 14 and EGFR [3]. Immunohistochemically, they

show a basal-like phenotype, regardless of the types of metaplastic elements. They also overexpress EGFR in over half of the cases [73, 74]. Microarray-based gene expression profiling has demonstrated that metaplastic carcinomas are part of the spectrum of basal-like breast carcinomas and display a myoepithelial and epithelial-to-mesenchymal transition-like molecular composition [75].

Lymph node metastases are less frequent in metaplastic carcinomas than in IDC NST. However, distant metastasis can occur in the absence of lymph node metastasis, as seen in other triple-negative breast cancers [3].

---

### Carcinomas with Neuroendocrine Features

These carcinomas exhibit morphological and immunohistochemical features of endocrine tumors, similar to those observed in the GI tract and lung, with the formation of solid, trabecular, glandular, and organoid structures. In the recent WHO classification [3], neuroendocrine breast carcinomas are categorized as follows:

- Neuroendocrine tumor, well differentiated
- Neuroendocrine carcinoma, poorly differentiated/small-cell carcinoma
- Invasive breast carcinoma with neuroendocrine differentiation

Invasive cancers of NST and other special types may show endocrine differentiation.

These tumors do not have any specific clinical presentation.

### Well-Differentiated Neuroendocrine Tumor

The tumor consists of densely cellular, solid nests and trabeculae of cells separated by a thin fibrovascular stroma [76]. These tumors are of low or intermediate grade (WHO). There is chromogranin positivity in more than 50 % of cases [77]. Other endocrine markers, such as synaptophysin and CD

56, are also positive. These tumors are typically positive for ER and PR and negative for HER2.

### Neuroendocrine Carcinoma

The tumor is composed of highly atypical cells with hyperchromatic nuclei and scant cytoplasm. Mitotic figures are frequent, and necrosis may accompany the lesion. The tumor should be distinguished from metastatic small-cell carcinoma of the lung; this distinction cannot be made on the sole basis of morphology. The presence of an in situ component supports the diagnosis of the breast as the primary cancer. Monoclonal NSE is positive in all cases of small-cell carcinomas and other neuroendocrine markers in approximately 50 % of the cases [3]. ER and PR expression may also be observed in over 50 % of cases and is generally correlated to the degree of differentiation. SCC is negative for HER2 expression [77, 78].

**Invasive Breast Carcinoma with Neuroendocrine Differentiation** Mucinous carcinoma of type B and solid papillary carcinoma (SPC) are the two tumors representing the most frequent examples of this category [3, 76].

---

### Secretory Carcinoma

Secretory carcinoma presents as a well-circumscribed mobile mass. The median age of presentation is 25 years. Microscopically, tumors show microcystic, tubular, and solid patterns, frequently in combination with each other. The characteristic finding is the presence of intracellular and extracellular secretory material showing positive staining with PAS. ER, PR, and HER2 are absent. EMA, alpha-lactalbumin, and S-100 protein are frequently present. There is a high expression rate of basal-like markers (CK5/6 or epidermal growth factor receptor) in secretory carcinomas [79]. Tognon et al. [80] showed that 12 of 13 of their cases of secretory breast carcinoma expressed the *ETV6-NTRK3* gene fusion. Laé et al. [81] demonstrated that secretory breast carcinoma with the *ETV6-NTRK3* fusion gene belongs to the pheno-

typic spectrum of basal-like breast carcinomas and that immunohistochemical and genetic features of secretory breast carcinomas distinguished them from other basal-like breast cancers. Secretory carcinoma has an indolent clinical behavior, especially in children and young adults [79].

---

## Papillary Lesions

These lesions, especially from the clinical perspective, are often confused with each other. For this reason, all of them will be discussed consecutively under the title “papillary lesions.”

---

### Intraductal Papillary Carcinoma (IDPC)

IDPC is a malignant, noninvasive neoplastic epithelial proliferation with papillary architectural features that occurs in the lumen of the ductal-lobular system [3]. Two types of IDPC exist:

- Central, solitary: Presentation may include nipple discharge.
- Peripheral, multifocal: Presentation may be as a mass.

Microscopically, ducts or TDLU are filled with slender, branching fibrovascular stalks, lined by a single layer or several layers of monomorphic epithelial cell population. Tumor cells have a bland appearance. High-grade nuclear features are rare. Solid, cribriform, and micropapillary patterns also exist. There is complete or near-complete (90%) absence of myoepithelial cells in the fibrovascular cores. However, there are myoepithelial cells at the periphery of the involved duct [82, 83].

---

### Encapsulated Papillary Carcinoma (EPC)

This lesion has a fibrous capsule, and its size ranges between 0.5 and 8 cm [84]. It is also called intracystic papillary carcinoma. All papillary intraductal carcinomas arise in a background of a

variably cystically dilated duct. The main difference in EPC is the loss of myoepithelial cells at the periphery of the lesion. EPC lacks these cells both in the fibrovascular cores and at the periphery [83, 84]. The absence of these cells and the reported cases of metastatic cases raises the possibility that these tumors represent low-grade invasive carcinomas with an expansile growth pattern [85]. However, the presence of continuous and intense collagen IV expression at the periphery is regarded as highly suggestive of a noninvasive carcinoma that is confined within an intact basement membrane [85].

EPC without an adjacent DCIS or any invasive component has a very favorable prognosis with adequate local therapy. The presence of associated DCIS confers a higher risk of local recurrence and meticulous radiologic examination before surgery is necessary.

---

### Solid Papillary Carcinoma

SPC is a variant of papillary carcinoma that is characterized by compact cellular growth within multiple nodules representing dilated ducts [61]. It presents in older women [82]. The neoplastic cells are ovoid or spindle cells of low to intermediate grade, and they have a streaming pattern. They are homogenous and do not form papillary or cribriform patterns. The most important indicator of the papillary nature of the lesion is the presence of thin, fibrovascular cores that are inconspicuous at low magnification. Neuroendocrine differentiation is frequent. Mucin production is common, and invasive mucinous carcinoma may coexist. Other types of invasive carcinoma may also be observed [86]. The distinction between in situ and invasive disease in SPC is difficult. Some authors regard this entity as an expansile variant of invasive carcinoma [86, 87]. SPC has an indolent clinical course even in cases with obvious invasion [86].

In the papillary lesions mentioned above, the lesion is called in situ if there is any doubt about the invasion. If there is obvious invasion, the staging should be conducted according to the measurement of the invasive component.



## Invasive Papillary Carcinoma

Invasive papillary carcinoma (IPC) is a carcinoma with a predominantly papillary morphology in its “invasive” component. This is a rare lesion, and there are no specific clinical and macroscopic features of this tumor. It should be distinguished from invasive carcinomas arising from EPC and SPC. Many cases in older series may have included such cases in this category [61].

## Adenoid Cystic Carcinoma

ACC is a carcinoma of low-grade malignant potential that is histologically similar to its counterpart in the salivary gland. This is a rare tumor. Approximately half of the cases arise from the subareolar region [88]. ACC is usually a circumscribed tumor.

Histologically, the tumor has the following basic patterns: tubular, cribriform, trabecular, and solid [89]. The neoplastic cells, which are epithelial and myoepithelial (basal), are arranged to form glandular spaces and pseudolumina [27]. Pseudolumina contain a myxoid acidic substance that is surrounded by myoepithelial cells. In smaller spaces, small spherules or cylinders of hyaline material are formed. True glandular spaces are surrounded by luminal cells and contain neutral mucosubstances. With occasional exceptions, ACC is triple negative [27].

Breast ACC rarely involves the axillary lymph nodes, and survival is excellent [88, 90]. A solid variant with basaloid features has a higher frequency of axillary lymph node metastasis [91].

## Glycogen-Rich Clear Cell Carcinoma

This is a carcinoma in which 90 % or more of the tumor cells have abundant clear cytoplasm containing glycogen [3]. It accounts for 1–3 % of breast carcinomas.

Histologically, the tumor cells have polygonal, sharply defined contours. The clear or finely granular cytoplasm contains PAS-positive diastase-labile glycogen. ER is present in 50 % of the cases, and PR is absent [26].

This tumor should be distinguished from lipid-rich carcinoma, histiocytoid carcinoma, and metastatic renal cell carcinoma [27].

There are conflicting reports regarding the prognosis of these tumors [92, 93].

## Inflammatory Carcinoma

Inflammatory carcinoma (IC) is an aggressive form of breast carcinoma with distinct clinical features. Clinically, there is rapid breast enlargement with edema and erythema of the skin (orange peel skin). Currently, there are no definitive molecular or pathological diagnostic criteria for IC. Therefore, the diagnosis is based on the clinical findings described above [94]. Signs and symptoms required for a diagnosis of IC include erythema occupation of at least one-third of the breast, edema and/or peau d’orange of the breast, and/or a warm breast, without an underlying palpable mass in the majority of cases [94, 95]. The onset of these signs and symptoms should be rapid; the duration of signs and symptoms at initial presentation should be  $\leq 3$  months [94].

IC is not considered a specific histological subtype of breast carcinoma, and there are no special pathological diagnostic criteria for IC [94, 95]. The underlying carcinoma is most often IDC NST of high grade; there may or may not be a distinct mass.

The pathognomonic histopathologic finding in IC is the presence of many lymphovascular tumor emboli in the papillary and reticular dermis overlying the breast. Although skin emboli are sometimes noted in the skin of patients with non-IC, emboli in patients with non-IC are usually less numerous and smaller than the skin emboli in patients with IBC [94]. The absence of tumor emboli in skin punch biopsies should not negate the diagnosis of IC because dermal tumor emboli can be detected in no more than 75 % of the cases despite meticulous sampling and sectioning [95]. Approximately 55 % of the cases are negative for ER and PR, 45 % is HER2 positive, and 33 % is triple negative [95].

Survival is worse than in patients with locally advanced breast cancer without IC [3].

Mucoepidermoid carcinoma, polymorphous carcinoma, oncocytic carcinoma, sebaceous carcinoma, lipid-rich carcinoma, and acinic cell carcinoma are very rare tumors and beyond the scope of this chapter.

## References

- Ellis IO. Intraductal proliferative lesions of the breast: morphology, associated risk and molecular biology. *Mod Pathol.* 2010;23 Suppl 2:S1–7.
- Pinder SE. Ductal carcinoma in situ (DCIS): pathological features, differential diagnosis, prognostic factors and specimen evaluation. *Mod Pathol.* 2010;23:S8–13.
- Lakhani SR, Ellis IO, Schnitt SJ, Tan PH, van de Vijver MJ. World health organization classification of tumors, WHO classification of tumors of the breast. 2nd ed. Lyon: IARC; 2012.
- Lester SC, Bose S, Chen YY, Connolly JL, de Baca ME, Fitzgibbons PL, et al. Protocol for the examination of specimens from patients with ductal carcinoma in situ of the breast. *Arch Pathol Lab Med.* 2009;133(1):15–25.
- Consensus conference on the classification of ductal carcinoma in situ. *Hum Pathol.* 1997;28:1221–5.
- Schwartz GF, Solin LJ, Olivetto IA, Ernster VL, Pressman P. The consensus conference on the treatment of in situ ductal carcinoma of the breast, 22–25 April 1999. *Breast.* 2000;9:177–86.
- Holland R, Peterse JL, Millis RR, Eusebi V, Faverly D, van de Vijver MJ, et al. Ductal carcinoma in situ: a proposal for a new classification. *Semin Diagn Pathol.* 1994;11(3):167–80.
- Page DL, Dupont WD, Rogers LW, Rados MS. Atypical hyperplastic lesions of the female breast. A long-term follow-up study. *Cancer.* 1985;55:2698–708.
- Tavassoli FA, Norris HJ. A comparison of the results of long-term follow-up for atypical intraductal hyperplasia and intraductal hyperplasia of the breast. *Cancer.* 1990;65(3):518–29.
- Schnitt SJ, Vincent-Salomon A. Columnar cell lesions of the breast. *Adv Anat Pathol.* 2003;10:113e2.
- Schnitt SJ. Clinging carcinoma: an American perspective. *Semin Diagn Pathol.* 2010;27(1):31–6.
- Aroner SA, Collins LC, Schnitt SJ, Connolly JL, Colditz GA, Tamimi RM. Columnar cell lesions and subsequent breast cancer risk: a nested case–control study. *Breast Cancer Res.* 2010;12(4):R61. doi:10.1186/bcr2624.
- Eusebi V, Magalhaes F, Azzopardi JG. Pleomorphic lobular carcinoma of the breast: an aggressive tumor showing apocrine differentiation. *Hum Pathol.* 1992;23:655–62.
- Fadare O, Dadmanesh F, Varado-Cabrero I, et al. Lobular intraepithelial neoplasia [lobular carcinoma in situ] with comedo-type necrosis: a clinicopathologic study of 18 cases. *Am J Surg Pathol.* 2006;30:1445–53.
- O'Malley FP. Lobular neoplasia: morphology, biological potential and management in core biopsies. *Mod Pathol.* 2010;23:S14–25. doi:10.1038/modpathol.2010.35.
- Page DL, Schuyler PA, Dupont WD, Jensen RA, Plummer Jr WD, Simpson JF. Atypical lobular hyperplasia as a unilateral predictor of breast cancer risk: a retrospective cohort study. *Lancet.* 2003;361:125–9.
- Lopez-Garcia MA, Geyer FC, Lacroix-Triki M, Marchió C, Reis-Filho JS. Breast cancer precursors revisited: molecular features and progression pathways. *Histopathology.* 2010;57(2):171–92. doi:10.1111/j.1365-2559.2010.03568.x.
- Bagaria SP, Shamonki J, Kinnaird M, Ray PS, Giuliano AE. The florid subtype of lobular carcinoma in situ: marker or precursor for invasive lobular carcinoma? *Ann Surg Oncol.* 2011;18(7):1845–51. doi:10.1245/s10434-011-1563-0.
- Shin SJ, Lal A, De Vries S, Suzuki J, Roy R, Hwang ES, et al. Florid lobular carcinoma in situ: molecular profiling and comparison to classic lobular carcinoma in situ and pleomorphic lobular carcinoma in situ. *Hum Pathol.* 2013;44(10):1998–2009.
- Edge SB, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti III A. American Joint Committee on Cancer (AJCC) cancer staging manual. 7th ed. New York: Springer; 2010.
- Prasad ML, Osborne MP, Giri DD, Hoda SA. Microinvasive carcinoma (T1mic) of the breast: clinicopathologic profile of 21 cases. *Am J Surg Pathol.* 2000;24(3):422–8.
- Rosai J. Breast. In: Rosai and Ackerman's surgical pathology. 9th ed. New York: Mosby; 2004. p. 1763–876.
- Silver SA, Tavassoli FA. Pleomorphic carcinoma of the breast: clinicopathological analysis of 26 cases of an unusual high-grade phenotype of ductal carcinoma. *Histopathology.* 2000;36(6):505–14.
- Nguyen CV, Falcón-Escobedo R, Hunt KK, Nayeemuddin KM, Lester TR, Harrell RK, et al. Pleomorphic ductal carcinoma of the breast: predictors of decreased overall survival. *Am J Surg Pathol.* 2010;34(4):486–93. doi:10.1097/PAS.0b013e3181ce97bf.
- Tavassoli FA, Norris HJ. Breast carcinoma with osteoclastlike giant cells. *Arch Pathol Lab Med.* 1986;110(7):636–9.
- Rosen PP. Rosen's breast pathology. 3rd ed. Philadelphia: Lippincott Williams & Wolters Kluwer business; 2009.
- Tavassoli FA, Eusebi V. Tumors of the mammary gland. AFIP atlas of tumor pathology. Fourth series; fasc 10. Washington, DC: American Registry of Pathology in collaboration with the Armed Institute of Pathology; 2009.
- Athanasou NA, Wells CA, Quinn J, Ferguson DP, Heryet A, McGee JO. The origin and nature of stromal osteoclast-like multinucleated giant cells in breast

- carcinoma: implications for tumour osteolysis and macrophage biology. *Br J Cancer*. 1989;59(4):491–8.
29. Shishido-Hara Y, Kurata A, Fujiwara M, Itoh H, Imoto S, Kamma H. Two cases of breast carcinoma with osteoclastic giant cells: are the osteoclastic giant cells pro-tumoural differentiation of macrophages? *Diagn Pathol*. 2010;5:55. doi:10.1186/1746-1596-5-55.
  30. Dixon JM, Anderson TJ, Page DL, Lee D, Duffy SW. Infiltrating lobular carcinoma of the breast. *Histopathology*. 1982;6(2):149–61.
  31. Tavassoli FA. *Pathology of the breast*. 2nd ed. New York: Appleton & Lange; 1999.
  32. Fechner RE. Histologic variants of infiltrating lobular carcinoma of the breast. *Hum Pathol*. 1975;6(3):373–8.
  33. Shousha S, Backhous CM, Alaghaband-Zadeh J, Burn I. Alveolar variant of invasive lobular carcinoma of the breast. A tumor rich in estrogen receptors. *Am J Clin Pathol*. 1986;85(1):1–5.
  34. Fisher ER, Gregorio RM, Redmond C, Fisher B. Tubulolobular invasive breast cancer: a variant of lobular invasive cancer. *Hum Pathol*. 1977;8(6):679–83.
  35. Weidner N, Semple JP. Pleomorphic variant of invasive lobular carcinoma of the breast. *Hum Pathol*. 1992;23(10):1167–71.
  36. Middleton LP, Palacios DM, Bryant BR, Krebs P, Otis CN, Merino MJ. Pleomorphic lobular carcinoma: morphology, immunohistochemistry, and molecular analysis. *Am J Surg Pathol*. 2000;24(12):1650–6.
  37. Walford N, ten Velden J. Histiocytoid breast carcinoma: an apocrine variant of lobular carcinoma. *Histopathology*. 1989;14(5):515–22.
  38. Eusebi V, Foschini MP, Bussolati G, Rosen PP. Myoblastomatoid (histiocytoid) carcinoma of the breast. A type of apocrine carcinoma. *Am J Surg Pathol*. 1995;19(5):553–62.
  39. Rakha EA, Patel A, Powe DG, Benhasouna A, Green AR, Lambros MB, et al. Clinical and biological significance of E-cadherin protein expression in invasive lobular carcinoma of the breast. *Am J Surg Pathol*. 2010;34(10):1472–9.
  40. Qureshi HS, Linden MD, Divine G, Raju UB. E-cadherin status in breast cancer correlates with histologic type but does not correlate with established prognostic parameters. *Am J Clin Pathol*. 2006;125(3):377–85.
  41. Da Silva L, Parry S, Reid L, Keith P, Waddell N, Kossai M, et al. Aberrant expression of E-cadherin in lobular carcinomas of the breast. *Am J Surg Pathol*. 2008;32(5):773–83.
  42. Sarrío D, Pérez-Mies B, Hardisson D, Moreno-Bueno G, Suárez A, Cano A, et al. Cytoplasmic localization of p120ctn and E-cadherin loss characterize lobular breast carcinoma from preinvasive to metastatic lesions. *Oncogene*. 2004;23(19):3272–83.
  43. Rakha EA, El-Sayed ME, Powe DG, Green AR, Habashy H, Grainge MJ, et al. Invasive lobular carcinoma of the breast: response to hormonal therapy and outcomes. *Eur J Cancer*. 2008;44(1):73–83.
  44. Pestalozzi BC, Zahrieh D, Mallon E, Gusterson BA, Price KN, Gelber RD, et al. International Breast Cancer Study Group: Distinct clinical and prognostic features of infiltrating lobular carcinoma of the breast: combined results of 15 International Breast Cancer Study Group clinical trials. *Eur J Cancer*. 2008;44(1):73–83.
  45. Rakha EA, van Deurzen CH, Paish EC, Macmillan RD, Ellis IO, Lee AH. Pleomorphic lobular carcinoma of the breast: is it a prognostically significant pathological subtype independent of histological grade? *Mod Pathol*. 2013;26(4):496–501.
  46. Rakha EA, Lee AHS, Evans AJ, Menon S, Assad NY, Hodi Z, et al. Tubular carcinoma of the breast: further evidence to support its excellent prognosis. *J Clin Oncol*. 2010;28(1):99–104.
  47. Diab SG, Clark GM, Osborne CK, Libby A, Allred DC, Elledge RM. Tumor characteristics and clinical outcome of tubular and mucinous breast carcinomas. *J Clin Oncol*. 1999;17(5):1442–8.
  48. Page DL, Dixon JM, Anderson TJ, Lee D, Stewart HJ. Invasive cribriform carcinoma of the breast. *Histopathology*. 1983;7(4):525–36.
  49. Ellis IO, Galea M, Broughton N, Locker A, Blamey RW, Elston CW. Pathological prognostic factors in breast cancer. II. Histological type. Relationship with survival in a large study with long term of follow up. *Histopathology*. 1992;20:479–89.
  50. Venable JG, Schwartz AM, Silverberg SG. Infiltrating cribriform carcinoma of the breast: a distinctive clinicopathologic entity. *Hum Pathol*. 1990;21(3):333–8.
  51. Tavassoli FA, Devilee P. *World health organization classification of tumors, tumors of the breast and female genital organs*. 2nd ed. Lyon: IARC Press; 2003.
  52. Vu-Nishino H, Tavassoli FA, Ahrens WA, Haffty BG. Clinicopathologic features and long-term outcome of patients with medullary breast carcinoma managed with breast-conserving therapy (BCT). *Int J Radiat Oncol Biol Phys*. 2005;62(4):1040–7.
  53. Marginean F, Rakha EA, Ho BC, Ellis IO, Lee AH. Histological features of medullary carcinoma and prognosis in triple-negative basal-like carcinomas of the breast. *Mod Pathol*. 2010;23(10):1357–63. doi:10.1038/modpathol.2010.123.
  54. Tan PH, Tse GM, Bay BH. Mucinous breast lesions: diagnostic challenges. *J Clin Pathol*. 2008;61(1):11–9.
  55. Capella C, Eusebi V, Mann B, Azzopardi JG. Endocrine differentiation in mucoid carcinoma of the breast. *Histopathology*. 1980;4(6):613–30.
  56. Di Saverio S, Gutierrez J, Avisar E. A retrospective review with long term follow up of 11,400 cases of pure mucinous breast carcinoma. *Breast Cancer Res Treat*. 2008;111(3):541–7.
  57. Tavassoli FA, Purcell CA, Bratthauer GL, Man Y. Androgen receptor positivity along with loss of

- bcl-2, ER, and PR expression in benign and malignant apocrine lesions of the breast. Implications for therapy. *Breast J.* 1996;2:1–10.
58. Niemeier LA, Dabbs DJ, Beriwal S, Striebel JM, Bhargava R. Androgen receptor in breast cancer: expression in estrogen receptor-positive tumors and in estrogen receptor-negative tumors with apocrine differentiation. *Mod Pathol.* 2010;23(2):205–12.
  59. Dellapasqua S, Maisonneuve P, Viale G, Pruneri G, Mazzarol G, Ghisini R, Mazza M, Iorfida M, Rotmensz N, Veronesi P, Luini A, Goldhirsch A, Colleoni M. Immunohistochemically defined subtypes and outcome of apocrine breast cancer. *Clin Breast Cancer.* 2013;13(2):95–102. doi:10.1016/j.clbc.2012.11.004.
  60. Siriaunkgul S, Tavassoli FA. Invasive micropapillary carcinoma of the breast. *Mod Pathol.* 1993;6(6):660–2.
  61. Ueng SH, Mezzetti T, Tavassoli FA. Papillary neoplasms of the breast: a review. *Arch Pathol Lab Med.* 2009;133:893–907.
  62. Marchiò C, Irvani M, Natrajan R, Lambros MB, Savage K, Tamber N, et al. Genomic and immunophenotypic characterization of pure micropapillary carcinomas of the breast. *J Pathol.* 2008;215(4):398–410.
  63. Zekioglu O, Erhan Y, Ciris M, Bayramoglu H, Ozdemir N. Invasive micropapillary carcinoma of the breast: high incidence of lymph node metastasis with extranodal extension and its immunohistochemical profile compared with invasive ductal carcinoma. *Histopathology.* 2004;44(1):18–23.
  64. Yamaguchi R, Tanaka M, Kondo K, Yokoyama T, Kaneko Y, Yamaguchi M, et al. Characteristic morphology of invasive micropapillary carcinoma of the breast: an immunohistochemical analysis. *Jpn J Clin Oncol.* 2010;40(8):781–7.
  65. Vingiani A, Maisonneuve P, Dell'orto P, Farante G, Rotmensz N, Lissidini G, et al. The clinical relevance of micropapillary carcinoma of the breast: a case-control study. *Histopathology.* 2013;63(2):217–24.
  66. Luini A, Aguilar M, Gatti G, Fasani R, Botteri E, Brito J, et al. Metaplastic carcinoma of the breast, an unusual disease with worse prognosis: the experience of the European Institute Of Oncology and review of the literature. *Breast Cancer Res Treat.* 2007;101(3):349–53.
  67. Tavassoli FA. Classification of metaplastic carcinomas of the breast. *Pathol Annu.* 1992;27(Pt 2):89–119.
  68. Foschini MP, Eusebi V. Carcinomas of the breast showing myoepithelial cell differentiation. A review of the literature. *Virchows Arch.* 1998;432(4):303–10.
  69. Van Hoeven KH, Drudis T, Cranor ML, Erlandson RA, Rosen PP. Low-grade adenosquamous carcinoma of the breast. A clinicopathologic study of 32 cases with ultrastructural analysis. *Am J Surg Pathol.* 1993;17(3):248–58.
  70. Sneige N, Yaziji H, Mandavilli SR, Perez ER, Ordonez NG, Gown AM, et al. Low-grade (fibromatosis-like) spindle cell carcinoma of the breast. *Am J Surg Pathol.* 2001;25(8):1009–16.
  71. Eusebi V, Lamovec J, Cattani MG, Fedeli F, Millis RR. Acantholytic variant of squamous-cell carcinoma of the breast. *Am J Surg Pathol.* 1986;10(12):855–61.
  72. Reis-Filho JS, Milanezi F, Paredes J, Silva P, Pereira EM, Maeda SA, et al. Novel and classic myoepithelial/stem cell markers in metaplastic carcinomas of the breast. *Appl Immunohistochem Mol Morphol.* 2003;11(1):1–8.
  73. Reis-Filho JS, Schmitt FC. p63 expression in sarcomatoid/metaplastic carcinomas of the breast. *Histopathology.* 2003;42(1):94–5.
  74. Reis-Filho JS, Milanezi F, Steele D, Savage K, Simpson PT, Nesland JM, et al. Metaplastic breast carcinomas are basal-like tumours. *Histopathology.* 2006;49(1):10–21.
  75. Weigelt B, Kreike B, Reis-Filho JS. Metaplastic breast carcinomas are basal-like breast cancers: a genomic profiling analysis. *Breast Cancer Res Treat.* 2009;117(2):273–80.
  76. Righi L, Sapino A, Marchiò C, Papotti M, Bussolati G. Neuroendocrine differentiation in breast cancer: established facts and unresolved problems. *Semin Diagn Pathol.* 2010;27(1):69–76.
  77. Sapino A, Righi L, Cassoni P, Papotti M, Pietribiasi F, Bussolati G. Expression of the neuroendocrine phenotype in carcinomas of the breast. *Semin Diagn Pathol.* 2000;17(2):127–37.
  78. Shin SJ, DeLellis RA, Ying L, Rosen PP. Small cell carcinoma of the breast: a clinicopathologic and immunohistochemical study of nine patients. *Am J Surg Pathol.* 2000;24(9):1231–8.
  79. Li D, Xiao X, Yang W, Shui R, Tu X, Lu H, et al. Secretory breast carcinoma: a clinicopathological and immunophenotypic study of 15 cases with a review of the literature. *Mod Pathol.* 2012;25(4):567–75.
  80. Tognon C, Knezevich SR, Huntsman D, Roskelley CD, Melnyk N, Mathers JA, et al. Expression of the ETV6-NTRK3 gene fusion as a primary event in human secretory breast carcinoma. *Cancer Cell.* 2002;2:367–76.
  81. Laé M, Fréneaux P, Sastre-Garau X, Chouchane O, Sigal-Zafrani B, Vincent-Salomon A. Secretory breast carcinomas with ETV6-NTRK3 fusion gene belong to the basal-like carcinoma spectrum. *Mod Pathol.* 2009;22:291–8.
  82. Collins LC, Schnitt SJ. Papillary lesions of the breast: selected diagnostic and management issues. *Histopathology.* 2008;52(1):20–9.
  83. Collins LC, Carlo VP, Hwang H, Barry TS, Gown AM, Schnitt SJ. Intracystic papillary carcinomas of the breast: a reevaluation using a panel of myoepithelial cell markers. *Am J Surg Pathol.* 2006;30(8):1002–7.
  84. Esposito NN, Dabbs DJ, Bhargava R. Are encapsulated papillary carcinomas of the breast in situ or

- invasive? A basement membrane study of 27 cases. *Am J Clin Pathol.* 2009;131(2):228–42.
85. Mulligan AM, O'Malley FP. Metastatic potential of encapsulated (intracystic) papillary carcinoma of the breast: a report of 2 cases with axillary lymph node micrometastases. *Int J Surg Pathol.* 2007;15:143–7.
  86. Nassar H, Qureshi H, Adsay NV, Visscher D. Clinicopathologic analysis of solid papillary carcinoma of the breast and associated invasive carcinomas. *Am J Surg Pathol.* 2006;30(4):501–7.
  87. Nicolas MM, Wu Y, Middleton LP, Gilcrease MZ. Loss of myoepithelium is variable in solid papillary carcinoma of the breast. *Histopathology.* 2007;51(5):657–65.
  88. Ghabach B, Anderson WF, Curtis RE, Huycke MM, Lavigne JA, Dores GM. Adenoid cystic carcinoma of the breast in the United States (1977 to 2006): a population-based cohort study. *Breast Cancer Res.* 2010;12(4):R54.
  89. Marchiò C, Weigelt B, Reis-Filho JS. Adenoid cystic carcinomas of the breast and salivary glands (or 'The strange case of Dr Jekyll and Mr Hyde' of exocrine gland carcinomas). *J Clin Pathol.* 2010;63(3):220–8.
  90. Kulkarni N, Pezzi CM, Greif JM, Suzanne Klimberg V, Bailey L, Korourian S, et al. Rare breast cancer: 933 adenoid cystic carcinomas from the National Cancer Data Base. *Ann Surg Oncol.* 2013;20(7):2236–41.
  91. Shin SJ, Rosen PP. Solid variant of mammary adenoid cystic carcinoma with basaloid features: a study of nine cases. *Am J Surg Pathol.* 2002;26(4):413–20.
  92. Hayes MM, Seidman JD, Ashton MA. Glycogen-rich clear cell carcinoma of the breast. A clinicopathologic study of 21 cases. *Am J Surg Pathol.* 1995;19(8):904–11.
  93. Kuroda H, Sakamoto G, Ohnisi K, Itoyama S. Clinical and pathological features of glycogen-rich clear cell carcinoma of the breast. *Breast Cancer.* 2005;12(3):189–95.
  94. Yamauchi H, Woodward WA, Valero V, Alvarez RH, Lucci A, Buchholz TA, et al. Inflammatory breast cancer: what we know and what we need to learn. *Oncologist.* 2012;17(7):891–9.
  95. Robertson FM, Bondy M, Yang W, Yamauchi H, Wiggins S, Kamrudin S, et al. Inflammatory breast cancer: the disease, the biology, the treatment. *CA Cancer J Clin.* 2010;60(6):351–75.



Ekrem Yavuz

---

## Abstract

This chapter describes the intraoperative pathological examination of breast lesions and the pathological examination of sentinel lymph nodes in breast cancer cases. Intraoperative pathological examination may be performed for the rapid diagnosis of breast malignancy, the assessment of the surgical margins of breast-conserving excision specimens, and the pathological analysis of sentinel lymph nodes. The most commonly used methods for intraoperative pathological examination of breast lesions are cytological and frozen section examinations in addition to gross analysis. The pathological examinations of sentinel lymph nodes necessitate careful gross examination and serial and/or step sectioning. Immunostaining using antibodies against pancytokeratin can also be performed. Sentinel lymph node metastases should be clearly defined as macro- or micrometastasis or isolated tumor cells. The differential diagnosis of subtypes of metastasis and mimickers is detailed.

---

## Keywords

Breast • Breast cancer • Frozen sections • Cytology • Differential diagnosis • Breast-conserving surgery • Sentinel lymph node biopsy • Immunohistochemistry • Micrometastasis • Extramedullary hematopoiesis

---

E. Yavuz, MD  
Department of Pathology,  
Istanbul Medical Faculty,  
Istanbul University, Millet Street, Fatih,  
Istanbul 34390, Turkey  
e-mail: [yavuze@istanbul.edu.tr](mailto:yavuze@istanbul.edu.tr)

The intraoperative pathological examination of resected breast tissue is performed for a number of reasons, including microscopic diagnosis of the breast lesion, assessment of surgical margins and sentinel lymph node status, and, rarely, determination of tissue adequacy for subsequent paraffin-block examination. The type of method used for the intraoperative

pathological examination depends on the experience and circumstances of the pathologist and varies from a simple gross examination to complex molecular techniques. However, the most commonly used methods are cytological and frozen section (FS) examinations in addition to gross analysis. Regardless of the intraoperative pathological method used, indefinite results may be obtained.

---

### **Intraoperative Pathological Diagnosis of Breast Lesions**

Surgeons prefer a definite preoperative diagnosis to develop a better strategy for the operation. Thus, the majority of breast lesions are diagnosed preoperatively by core biopsy or fine-needle aspiration cytology. Consequently, FS examination of a primary breast tumor is requested only when these preoperative procedures are not diagnostic, have failed to determine the presence of an invasion, or have been omitted. Routine FS examination is not recommended for grossly indistinct and “possibly benign” breast lesions, although a small percentage of these specimens may contain grossly undetectable in situ or invasive breast carcinoma. Many pathologists are reluctant to perform FS analysis on breast lesions with a diameter of less than 1 cm because the frozen tissue remaining from the intraoperative examination is not ideal for determining prognostic and predictive parameters of the breast carcinoma. However, in our opinion, careful gross slicing of a breast lesion even with a diameter of 5 mm provides a sufficient amount of tissue for subsequent paraffin-block examination [1–3].

There is also a debate on the use of FS for mammographically detected non-palpable breast lesions. Some authors have reported little difficulty in using FS in this setting, unless the lesion was only a mammographic calcification [4–6]. However, we agree with the recommendation that FS not be routinely performed for breast lesions without a grossly detectable mass because of the difficulty of subsequent gross and microscopic examinations. The tissue may be distorted by freezing artifacts. Furthermore, some portion


of the tissue may be lost, and tissue orientation may be very difficult [7].

The FS diagnosis of a grossly detectable breast lump is straightforward and has high specificity, sensitivity, and accuracy rates close to 100 % [8, 9]. However, because urgent intraoperative diagnosis of a primary breast lump is rarely requested, young pathologists should be aware of the possible diagnostic pitfalls in this setting. Intraoperative diagnosis of a primary breast lump by touch print cytology can be performed accurately and can be used as an adjunct method to FS but requires experience in the cytological features of breast lesions [10–12]. Furthermore, because rapid preoperative cytological analysis of a breast lump can be performed in more appropriate circumstances, pathologists deserve the right to refuse to employ only cytological methods to diagnose a malignancy during the operation.


### **Intraoperative Pathological Assessment of Surgical Margins**

The intraoperative pathological assessment of surgical margins plays a critical role in breast-conserving surgery (BCS) because surgical margin negativity is the most important relevant parameter for local recurrence after BCS for breast cancer. However, there are many issues limiting the success rates of this procedure that a pathologist must address. Due to the lipid-rich characteristics of breast tissue, the surface of the excised tissue is frequently irregular and contains crevasses and defects. Tissue flattening, which is referred to as the “pancake phenomenon,” after the removal of the breast tissue also contributes to inaccurate results [13–15].

It is the surgeon’s responsibility to submit an intact specimen and include appropriate orientation markings. The pathologist can easily be oriented to the specimen with a short suture at the superior margin and a long suture at the lateral margin placed by the surgeon (Fig. 14.1). Upon receipt, the pathologist should approach the specimen as if it contains a malignant tumor. The surface of the specimen should be marked using




**Fig. 14.1** Gross appearance of a resected right breast lump with orientation sutures. A short suture at the superior margin, a long stitch at the lateral margin, and long and short stitches at the anterior margin were placed by the surgeon



**Fig. 14.2** Distance from the irregular breast lump to the superior, inferior, anterior, and posterior margins can be observed after gross slicing of an inked specimen

India ink or another dye that is resistant to processing solutions and remains visible at the edge of the sample during the microscopic examination. If a cotton applicator is used and the ink is rapidly fixed in Bouin's or a similar solution, the ink will not permeate into the tissue crevasses. A breast excision specimen sliced by the surgeon may be reassembled but will not be as accurate as an intact specimen. The most important part of the surgical margin assessment is the gross examination. The pathologist must be informed about the number and size of the lumps and the radiological findings. For this reason, it is preferable to receive the patient's file during the intraoperative pathological examination. After carefully slicing the specimen, the relationship of an apparent tumor to the six margins can be observed and reported grossly (Figs. 14.2 and 14.3). However, the gross impressions may fail to estimate the





**Fig. 14.3** Surgical margin assessment in a breast lesion containing comedo-type necrotic areas and grossly suggesting a ductal carcinoma in situ

involvement of a surgical margin adjacent to the microscopic focus of an invasive or in situ carcinoma. Hence, a FS examination should be performed if a margin is grossly close to the tumor. However, the success of surgical margin assessment depends on the pathologist's experience and the methods used. A random sectioning of grossly normal surgical margins is not recommended because it has limited sensitivity [16].

In our opinion, samples for FS should be perpendicular to the inked surface because the true distance between the tumor and the margin can be determined (Figs. 14.4 and 14.5). Samples taken parallel to the surface (enface samples) can also be frozen sectioned for surgical margin assessment. In this setting, any tumor in the section indicates margin positivity. However, if no tumor is detected in the section, we can estimate a tumor-free distance of at least 2 mm between the tumor and the margin because a shaved sample is usually at least 2 mm in thickness. The sampling method, whether perpendicular to the

**Fig. 14.4** Frozen section appearance of an invasive ductal carcinoma sectioned perpendicular to the margin. The fatty part of the breast tissue has been lost due to difficulties with frozen sectioning (H&E  $\times 4$  original magnification)



**Fig. 14.5** Gross appearance of frozen-sectioned tissue may also facilitate surgical margin assessment because freezing usually highlights neoplastic tissue

inked surface or a shaved sample, should be clearly stated on the report.

Some surgeons prefer to perform re-excisions by shaving the excision cavity at the initial operation. These specimens should also be marked by suture or ink for orientation by the operator. These re-excisions are helpful in the decision to reoperate if there is a “close” or positive margin in the excision [17–19].

Surgical shavings from the excision cavity have revealed that negative excision margins during the intraoperative pathological examination do not absolutely assure that all the carcinomatous tissue in the region has been successfully removed. The likelihood of the presence of a

carcinoma in a shave biopsy of the tumor bed after a negative lumpectomy margin varies from 9 to 39 % [19–21]. This variation may reflect the characteristics of either the primary tumor or the surgical team. Invasive lobular carcinomas and similar tumors showing microscopic multifocal growth patterns, an extensive intraductal component and extensive lymphovascular invasion, are more likely to result in a false-negative intraoperative surgical margin assessment. Furthermore, the type and extent of the radiological methods used and the experience of the surgeon and pathologist have a great impact on the success rates of BCS.

Retrospective analyses of the impact of FS on BCS have revealed that during the first operation, additional excisions are performed in 24–27 % of cases based on FS results, while 5–9 % necessitated a second re-excision due to definitive histopathological examination.[22–24].

Cytological methods can also be used for margin assessment. Intraoperative touch preparation cytology (IOTPC) or “imprint cytology” is based on the ability of malignant cells but not benign mammary fat tissue to adhere to glass slides. To assess margin status, glass slides are first brought against the borders of the excised specimen, and then the slides are rapidly fixed and stained. Some studies have reported that IOTPC is



inexpensive, accurate, and rapid and conserves tissue for permanent sectioning and histopathological examination [25–27]. Klimberg et al. [25] reported a diagnostic sensitivity and specificity of 96 and 100 %, respectively, in a study of 428 patients. Weinberg et al. [28] stated that IOTPC significantly reduced local recurrence (LR) rates compared to other methods. However, there are limitations regarding the use of IOTPC. It necessitates experience in breast cytology, and some pathologists are very reluctant to rely only on cytological methods intraoperatively. In our opinion, a proportion of indefinite results are inevitable when using cytology for margin assessment, particularly for tumors with low nuclear atypia, such as invasive lobular carcinoma. Another disadvantage of IOTPC is that close margins cannot be observed because only superficial tumor cells are detected with this technique. Therefore, no information regarding margin distance, tumor multifocality, or the presence of either in situ or invasive carcinoma can be provided by this method. Cox et al. [16] reported three false-positive interpretations with cytology and no false-positive and five false-negative interpretations with FS. Cytospin preparations can be prepared from BCS specimens by scraping the surface. Veronesi et al. [29] used the monoclonal antibody B72.3 to detect carcinomatous cells in cytospin preparations obtained by scraping. However, they detected immunoreactive cells in 33 % of the cytospin specimens, whereas only 12 % had definitive margin positivity. We suggest that the addition of rapid immunohistochemistry to the cytological method can increase the sensitivity of diagnosing positive margins but will not resolve any of the outlined limitations of using cytological methods for margin assessment.

Reporting the pathological assessment of excisions is also problematic. The existence of carcinomatous cells on the inked surface should be reported as positive margins, and the extent of the positivity should be detailed if possible. However, the definition of a close margin is debated. The definition “close to the margin” is conventionally used for tumor foci that are less than one high-power field away from the surgical

margin, whereas some pathologists use this term for tumors that are less than 3 mm from the surgical margin. We only use the term “positive margin”; otherwise, we give the distance to the inked surface.

---

## **Pathological Examination of Sentinel Lymph Nodes**

### **Intraoperative Pathological Diagnosis of Sentinel Lymph Nodes**


Intraoperative pathological examination of sentinel lymph nodes (IPESLN) should be performed if the result will influence the type of surgical approach. According to this definition, an axillary lymph node dissection (ALND) should be performed if the intraoperative sentinel lymph node (SLN) diagnosis is metastatic and omitted if the SLN is metastasis-free. However, clinicians should be aware of the limitations of IPESLN. In a number of patients, metastatic foci will not be detected because of inadequate sampling and the difficulty of detecting small metastatic foci in SLNs. Therefore, patients must be informed of the possibility or risk of a second operation for completion of ALND. A recent study by the American College of Surgeons Clinical Oncology Group (ACOSOG) demonstrated that the completion of ALND after a pathological diagnosis of SLN metastasis was not superior to SLN biopsy alone in terms of disease-free and overall survival for early-stage clinically N0 breast cancer patients [30]. This result will lead to a decrease in the use of IPESLNs. However, this study included only patients treated with BCS and radiation therapy. Therefore, pathologists will continue to perform IPESLNs.

IPESLNs can be performed either using FS or imprint cytology or a combination of both techniques. Another cytological method for intraoperative SLN analysis is the evaluation of cells scraped from the cut surface of the node.

Cytological methods have some advantages compared to frozen sectioning. Cytology conserves tissue for subsequent permanent histopathological examination. In addition,



**Fig. 14.6** Imprint cytology of the sentinel lymph node showing metastatic breast cancer cells within the lymphoid cellular background (H&E  $\times 60$  original magnification)




cytological analysis is not time-consuming, and it permits the analysis of specimens prepared from multiple cut surfaces. Cytological preparations can be rapidly stained with H&E (Fig. 14.6), Giemsa, or other stains depending on the pathologist's preference. However, many pathologists are not skilled at cytological diagnosis and are reluctant to use cytology for IPESLN because observing sparse carcinomatous cells in the highly cellular background of the lymph node imprint may be very difficult. Another disadvantage of cytology is that the detected metastasis cannot be measured, although sparse metastatic cells on imprint preparations usually correspond to either micrometastasis or isolated tumor cells (ITCs). Several studies have reported controversial results with regard to imprint cytology for IPESLNs. In a meta-analysis of 32 studies, Tew et al. [31] concluded that intraoperative cytology (IC) is simple, rapid, and sensitive for macrometastases (MAM) but not micrometastases (MIM). The pooled sensitivity of IC was 63 % (81 % for MAM and 22 % for MIM), and its specificity was 99 %. These success rates are slightly worse than those for frozen sectioning reported in another meta-analysis of the use of frozen sectioning for IPESLNs [32].

The FS method for IPESLN (Fig. 14.7) has been evaluated in several studies, and satisfactory results have usually been reported. In a meta-analysis of the use of frozen sectioning for IPESLN, Liu et al. reported that the mean sensitivity was 73 % (94 % for MAM and 40 % for MIM/ITC) and the mean specificity was 100 % [32]. In a study directly comparing IC and FS methods for IPESLN, van Diest et al. observed false-negative rates of 13 % for FS and 38 % for IC, with 88 % concordance between FS and IC [33]. The sensitivity and overall accuracy (87 % and 95 %, respectively) of FS were substantially greater than those of IC (62 and 83 %, respectively). In a study by Turner et al. [34], combining IC and FS methods resulted in an overall accuracy of 93.2 %. However, the combined intraoperative method detected 87 % of MAMs but only 28 % of MIMs. At our institution, we perform IC for IPESLNs because we are concerned about tissue loss during IPESLN, and there is a consensus in the multidisciplinary team that the main objective of IPESLNs is to detect all MAMs. As mentioned above, we do not consider either FS or IC reliable for ruling out MIMs and ITCs.

Molecular biology techniques, such as reverse transcriptase polymerase chain reaction,

**Fig. 14.7** Frozen section of a breast carcinoma metastasis in the sentinel lymph node. A sharp contrast between the carcinoma and lymphoid parenchyma can be observed (H&E  $\times 20$  original magnification)




are under investigation for their potential applicability for the evaluation of SLNs. Although these techniques are highly sensitive and permit the evaluation of large amounts of tissue, the tissues are destroyed during the procedure, and consequently, it is not possible to determine the cell from which the signal originated. Viale et al. compared molecular techniques with FS for IPESLNs using a commercially available molecular assay in a series of 293 SLNs [35]. Using the molecular technique, they correctly detected 51 of 52 MAMS and 5 of 20 MIMs. They concluded that the sensitivity of the molecular assay was comparable to that of histopathological examination of the entire SLN by serial sectioning at 1.5–2 mm. Similarly, many studies using molecular techniques have reported satisfactory results in detecting even small metastases during IPESLN [36–38]. We believe that molecular techniques should be evaluated as research studies and are not ready to use in routine IPESLNs. The American Society of Clinical Oncology (ASCO) has recommended that molecular approaches should remain investigational and tissue potentially required for histological diagnosis should not be utilized for investigational purposes until the diagnosis is secure [39].

## Permanent Pathological Analysis of Sentinel Lymph Nodes

### Gross Examination

The submitted tissue may be a single or a few SLNs or axillary fatty tissue containing one or more lymph nodes. Fatty tissue should be carefully dissected to identify all lymph nodes. All lymph nodes should be measured and sliced into 2-mm-thick sections. We prefer producing slices parallel to the longitudinal axis of the SLN (Fig. 14.8). If isosulfan blue dye or methylene blue is used, the afferent lymphatic vessel can be observed, and the section can be made where the afferent lymphatic vessel is connected to the SLN. Such sectioning facilitates the detection of tiny metastases, which are typically located at the marginal sinus into which the lymph fluid first drains. Some pathologists prefer transverse slicing if the SLN is larger than 0.5 cm. All slices should be inspected for changes in color and consistency. The cut surface of a positive lymph node ranges from tan to gray-white in color and from normal to hard in consistency. In cases of partial involvement, a sharp contrast between the metastatic focus and normal lymphoid tissue can easily be observed. However, permeative metastases of lobular carcinomas usually exhibit no or subtle



**Fig. 14.8** Gross appearance of a sentinel lymph node detected using blue dye and showing fatty changes. Slicing was performed parallel to the longitudinal axis

gross findings. Each SLN should be submitted in a separate cassette, and all nodes should be submitted for microscopic examination.

### Sectioning

The majority of MAMs and MIMs are easily detected during standard H&E-stained section examination, an approach endorsed by leading surgical pathology organizations [40, 41]. Although superficial serial sectioning limits sampling to the upper limits of the paraffin block, virtually all MAMs can be diagnosed if the gross slicing is performed as recommended [42–44].

However, if multiple step sections from different levels of the block are examined, the majority of MIMs can be detected. Various protocols for the pathological examination of SLNs have been reported, but there is no consensus as to which is the most cost-effective [34, 45–47]. Step sectioning of the block completely at 0.2-mm or smaller intervals enables the detection of all MIMs but will result in too many sections. Decreasing the step-sectioning interval will obviously result in an increase in the detection of MIMs and ITCs [48]. Interestingly, a recent study reported that 0.02-mm and 0.150-mm interval sectioning produced comparable results for the detection of metastatic deposits in SLNs [49]. Weaver suggested that “the size of the largest acceptable missed metastasis must be determined prospectively and then the

sectioning strategy that will systematically detect metastases larger than the threshold must be employed” [50].

### Use of Immunohistochemistry


The use of antibodies to cytokeratin facilitates the detection of small clusters of metastatic carcinomatous cells, particularly in lobular carcinoma metastases that may manifest as discohesive cells throughout the sinuses (Fig. 14.9a, b) [51]. Nevertheless, the clinical significance of these small clusters of metastatic cells, which are usually detected by immunohistochemistry, is unknown and the subject of ongoing clinical studies.

Guidelines published by the College of American Pathologists [40] and others [52, 53] state that immunohistochemical analysis is not required for the evaluation of SLN. However, in the latest recommendation paper, a liberal approach was allowed by The American Society of Clinical Oncology [39]. They stated that the decision to utilize immunohistochemical analysis and act on the results remains, for now, a matter of discussion among individual surgeons, oncologists, and pathologists, based on a determination of the best course for their patients as assessed from their own experience and a review of the available literature. A survey of European pathology laboratories identified 123 different protocols described by 240 respondents, 71 % of which used immunohistochemistry to evaluate SLNs found to be negative in H&E-stained sections [54]. We prefer using pancytokeratin immunostaining in our routine practice because it accurately determines the extent of metastatic foci. Moreover, most pathologists dealing with SLNs have encountered undetected MAMS identified by immunohistochemistry. Immunohistochemistry is most useful for the discrimination of isolated tumor cells from MIMs.

### Histopathology

Breast carcinoma metastases in SLNs usually recapitulate the architectural and cytological patterns of the primary tumor. MAMs usually disrupt the lymph node architecture and can easily be detected at scanning power. Lobular





**Fig. 14.9** (a) Microscopic appearance of an invasive lobular carcinoma in the sentinel lymph node. Observing the neoplastic cells (*arrows*) may be difficult due to the dyshesive nature and the small size of the neoplastic cells (H&E  $\times 40$  original magnification).


(b) Immunohistochemical examination using anti-cytokeratin antibody highlights the metastatic cells dispersed within the parenchyma of the sentinel lymph node (anti-pancytokeratin, Mayer's hematoxylin counterstaining  $\times 20$  original magnification)

carcinomas may diffusely infiltrate the lymph node parenchyma as single cells or small clusters. A MAM is defined as a lymph node metastasis of breast carcinoma measuring greater than 2 mm in size [55]. A differential diagnosis of MAMs includes other malignancies, such as malignant melanoma, lung carcinoma, and malignant lymphomas.


A MIM is defined as a lymph node metastasis measuring greater than 0.2 mm and/or more than 200 cells in a cross section of a single SLN but not greater than 2.0 mm in size [55]. In patients with MIMs in the SLN, a MAM may exist in the non-sentinel lymph node of the completion ALND in approximately 15 % of patients [56]. The majority of MIMs are cohesive malignant cell aggregates without specific patterns and without desmoplasia. Because MIMs are usually found within the subcapsular sinus, careful examination of these areas is critical, and pancytokeratin immunostaining is helpful as previously stated. Differential diagnosis of MIMs includes MAMs and ITCs and also some benign lesions, such as nevus cell aggregates, heterotopic glands, and mechanical transport of benign breast epithelium. Multiple MIMs may exist in a SLN, and these should not be diagnosed as MAMs even when the aggregate diameter of the metastatic deposits exceeds 2 mm [57]. Nevus

cell aggregates can be found in the lymph node capsule and, occasionally, in capsular invaginations within the lymph node parenchyma. The cells of nevus cell aggregates resemble the cells of cutaneous nevi. They have round to ovoid nuclei with fine chromatin, pale-staining cytoplasm, and indistinct cell borders (Fig. 14.10). Melanin pigment is variably present. The distinction can usually be made by routine microscopic analysis; however, in some lesions, immunohistochemical examination using both pancytokeratin and a melanocytic marker may be necessary. Multiple nevus cell aggregates are possible, and both MIM and nevus cell aggregates may be present in the same SLN [58]. There may be heterotopic glandular epithelium in axillary lymph nodes, which may exhibit ductal, squamous, or salivary gland-type differentiation and even proliferative and metaplastic changes [59]. However, the diagnosis of heterotopic epithelium during the microscopic examination of a SLN should be made with great caution because some metastatic carcinomas with low nuclear atypia, such as tubular and tubulolobular carcinoma, may resemble nonneoplastic glandular inclusions (Fig. 14.11). For this discrimination, an observation of the myoepithelial cell layer or the low proliferation capacity of the cells using immunohistochemistry may help.

**Fig. 14.10** Microscopic appearance of a nevus cell aggregate (*arrow*) at the capsule of the sentinel lymph node. Nevus cells lack nuclear atypia, and their cytoplasmic borders are indistinct, as with dermal nevi (*asterisk*: nevus cells within the capsule invagination through the lymphoid parenchyma) (H&E  $\times 20$  original magnification)



**Fig. 14.11** Glandular structure (*asterisk*) simulating a benign heterotopic epithelium within the capsule of a sentinel lymph node that was diagnosed as “isolated tumor cells” resulting from a tubular carcinoma (H&E  $\times 20$  original magnification)




ITCs are defined as small tumor-cell clusters with a diameter of less than or equal to 0.2 mm (Fig. 14.12). Another definition of ITCs is the existence of less than 200 tumor cells in a single cross section of a single SLN. The latter definition is particularly useful for differentiating MIMs and ITCs in invasive lobular carcinoma cases [55]. Microscopically, ITCs are usually located within the subcapsular sinus, although they may occasionally be observed within the medullary sinus


and lymph node parenchyma. In the case of a few malignant cells within the afferent lymphatics, one should carefully look for a metastatic focus inside the lymph node. Although the metastatic cells within the afferent lymphatics are traditionally not used for staging purposes, pathologists should note their existence in the report. The differential diagnosis of ITCs includes MIMs, nevus cell aggregates, heterotopic glandular structures, extramedullary hematopoiesis, and benign mechanical



**Fig. 14.12** Positive staining for anti-cytokeratin in a small cluster composed of a few metastatic cells from an invasive ductal carcinoma located in the subcapsular sinus of the sentinel lymph node. According to size criteria (less than 0.2 mm), it should have been diagnosed as “isolated tumor cells.” Note the slight immune positivity in reticulum cells (anti-pancytokeratin, Mayer’s hematoxylin counterstaining  $\times 40$  original magnification)




**Fig. 14.13** Metastatic cells from an invasive lobular carcinoma within the medullary sinus highlighted with anti-cytokeratin immunostaining. Neoplastic cells are either isolated or in two- to three-cell groups. This case should have been diagnosed as “isolated tumor cells” if less than 200 neoplastic cells were counted in the single cross section (anti-pancytokeratin, Mayer’s hematoxylin counterstaining  $\times 40$  original magnification)



transport (BMT) of displaced epithelium. The distinction between ITCs and MIMs in invasive ductal carcinomas is relatively easy and depends on the measurement of the largest focus. However, this is a problematic issue in invasive lobular carcinoma cases because the metastasis is frequently dispersed within the sinuses and lymph node parenchyma as isolated single cells or two- to three-cell groups. In this setting, the alternative size criterion of less than 200 cells in a single cross

section of a single lymph node is very useful (Fig. 14.13). Nevertheless, in some cases, the distinction may be extremely difficult, and the pathologist must make a final judgment according to his or her impression. Extramedullary hematopoiesis also occurs in SLNs, as in other organs. The composition of hematopoietic elements may vary. Megakaryocytes and their immature forms are large cells with multiple nuclei that can be misdiagnosed as ITCs (Fig. 14.14). Immunohistochemical

**Fig. 14.14** Microscopic appearance of extramedullary hematopoiesis in a sentinel lymph node. A megakaryocyte with nuclear multilobation (*arrow*) and granulocytic precursors with dense eosinophilic cytoplasm (*arrow heads*) are shown (H&E  $\times 40$  original magnification)




stains are very useful for this discrimination. Both benign and malignant breast tissue may be traumatically displaced by either a biopsy procedure or vigorous breast massage after the injection of radiocolloid or dye tracer for the detection of SLNs [60]. The displaced epithelial cells may also enter the lymphatics and drain into the SLN. Interestingly, fine-needle aspirations produce mechanically transported epithelium within the SLN more commonly than core and excisional biopsy procedures [61]. The mechanical transport of epithelial cells to the SLN is more common in papillary breast lesions because they are usually friable lesions [62]. Some morphological and immunohistochemical details facilitate the identification of BMT. Accompanying hemosiderin-laden histiocytes around epithelial cells, a lack of nuclear atypia (Fig. 14.15 a, b) and proliferative activity in epithelial cells, a myoepithelial cell layer at the periphery of the epithelial cell cluster (Fig. 14.15c), a mixture of high- and low-molecular-weight cytokeratin expression (Fig. 14.15d), and entrapped epithelial cells within the granulation tissue at the biopsy site in the breast are helpful morphological and immunohistochemical features for the differential diagnosis of BMT and ITC/MIM. Nevertheless, some pathologists favor the diagnosis of ITC in patients with an invasive carcinoma to avoid understaging because it is usually very difficult to diagnose BMT to the SLN.

### Role of the Pathological Examination of SLNs in Special Circumstances

Although pathologists usually do not have an active role in selecting patients in which an SLN biopsy (SLNB) will be performed, they should be aware of controversial circumstances. SLNB is currently used in clinically early-stage invasive breast cancer. ASCO recommends against the use of SLNB in patients with T3–T4 breast cancers, inflammatory breast cancer, pregnancy, male breast cancer, prior non-oncological breast surgery, preoperative systemic chemotherapy, and prior axillary surgery [39]. In patients with ductal carcinoma in situ (DCIS), the frequency of SLNB positivity in previous studies has varied from 1.9 to 9 % [63–67]. SLNB should be performed in patients with DCIS when a mastectomy is indicated because performing an SLNB is impossible after finding an invasive tumor during a permanent pathological examination of the mastectomy specimen. SLNB may also be performed in patients with large, high-grade, or palpable DCIS cases to avoid a secondary operation because it is not infrequent to find an invasive focus in these types of DCIS cases [39]. The use of SLNB in breast cancer patients with preoperative chemotherapy (Fig. 14.16) is controversial. In studies with small institutional case series, the false-negative rate ranged from 0 to 33 % [68–75]. In cases






**Fig. 14.15** Case of benign mechanical epithelial transport in the sentinel lymph node. The primary breast lesion was misdiagnosed as invasive carcinoma, and the paraffin-block consultation resulted in a diagnosis of sclerosing papilloma. (a) Epithelial cell clusters within the lymphoid parenchyma (H&E  $\times 10$  original magnification). (b) High magnification of the rectangle in “a” showing the lack of

obvious nuclear atypia (H&E  $\times 60$  original magnification). (c) The myoepithelial cell layer at the periphery of the cluster highlighted with anti-p63 immunostaining (arrows) (anti-p63, Mayer’s hematoxylin counterstaining  $\times 40$  original magnification). (d) (anti-cytokeratin 14, Mayer’s hematoxylin counterstaining  $\times 4$  original magnification)

**Fig. 14.16** Imprint cytology of a breast carcinoma metastasis in a sentinel lymph node excised after neoadjuvant chemotherapy. Note the cellular enlargement caused by chemotherapy (H&E  $\times 20$  original magnification)



with neoadjuvant chemotherapy, the success rate of identifying the SLN improves with experience, but the false-negative rate does not change [76]. In the NSABP B-27 trial, the false-negative rate of SLNs was 10.7 % [77]. The ASCO panel concluded that there are insufficient data to recommend SLNBs in breast cancer with neoadjuvant chemotherapy and that an SLNB should be performed only in clinically negative axillary lymph nodes, whether in the preoperative or postoperative settings [39].

## References

- Rosen PP, Senie R, Schottenfeld D, Ashikari R. Noninvasive breast carcinoma: frequency of unsuspected invasion and implication for treatment. *Ann Surg.* 1979;89:98–103.
- Speights Jr VO. Evaluation of frozen sections in grossly benign breast biopsies. *Mod Pathol.* 1994;7:762–5.
- Niemann TH, Lucas JG, Marsh Jr WL. To freeze or not to freeze. A comparison of methods for the handling of breast biopsies with no palpable abnormality. *Am J Clin Pathol.* 1996;106:225–8.
- Bianchi S, Palli D, Ciatto S, Galli M, Giorgi D, Vezzosi V, et al. Accuracy and reliability of frozen section diagnosis in a series of 672 nonpalpable breast lesions. *Am J Clin Pathol.* 1995;103:199–205.
- Ferreiro JA, Gisvold JJ, Bostwick DG. Accuracy of frozen-section diagnosis of mammographically directed breast biopsies. Results of 1,490 consecutive cases. *Am J Surg Pathol.* 1995;19:1267–71.
- Tinnemans JG, Wobbes T, Holland R, Hendriks JH, van der Sluis RF, Lubbers EJ, et al. Mammographic and histopathologic correlation of nonpalpable lesions of the breast and the reliability of frozen section diagnosis. *Surg Gynecol Obstet.* 1987;165:523–9.
- Association of Directors of Anatomic and Surgical Pathology. Immediate management of mammographically detected breast lesions. *Am J Surg Pathol.* 1993;12:850–1.
- Fessia L, Ghiringhello B, Arisio R, Botta G, Aimone V. Accuracy of frozen section diagnosis in breast cancer detection. A review of 4436 biopsies and comparison with cytodiagnosis. *Pathol Res Pract.* 1984;179:61–6.
- Kaufman Z, Lew S, Griffel B, Dinbar A. Frozen-section diagnosis in surgical pathology. A prospective analysis of 526 frozen sections. *Cancer.* 1986;57:377–9.
- Esteban JM, Zaloudek C, Silverberg SG. Intraoperative diagnosis of breast lesions. Comparison of cytologic with frozen section technics. *Am J Clin Pathol.* 1987;88:681–8.
- De Rosa G, Boschi R, Boscaino A, Petrella G, Vetrani A, Palombini L, et al. Intraoperative cytology in breast cancer diagnosis: comparison between cytologic and frozen section techniques. *Diagn Cytopathol.* 1993;9:623–31.
- Veneti S, Ioannidou-Mouzaka L, Toufexi H, Xenitides J, Anastasiadis P. Imprint cytology. A rapid, reliable method of diagnosing breast malignancy. *Acta Cytol.* 1996;40:649–52.
- Spivack B, Khanna MM, Tafra L, Juillard G, Giuliano AE. Margin status and local recurrence after breast-conserving surgery. *Arch Surg.* 1994;138:1371–4.
- Schnitt SJ, Abner A, Gelman R, Connolly JL, Recht A, Duda RB, et al. The relationship between microscopic margins of resection and the risk of local recurrence in patients with breast cancer treated with breast conserving surgery and radiation therapy. *Cancer.* 1994;74:1746–2751.
- Graham RA, Homer MJ, Katz J, Rothschild J, Safaii H, Supran S. The pancake phenomenon contributes to the inaccuracy of margin assessment in patients with breast cancer. *Am J Surg.* 2002;184:89–93.
- Cox CE, Ku NN, Reintgen D, Greenberg HM, Nicosia SV, Wangenstein S. Touch preparation cytology of breast lumpectomy margins with histologic correlation. *Arch Surg.* 1991;126:490–3.
- Barthelmes L, Al Awa A, Crawford DJ. Effects of cavity margin shavings to ensure completeness of excision on local recurrence rates following breast conserving therapy. *Eur J Surg Oncol.* 2003;29:644–8.
- Malik HZ, George WD, Mallon EA, Harnett AN, Macmillan RD, Purushotham AD. Margin assessment by cavity shaving breast-conserving surgery: analysis and follow-up of 543 patients. *Eur J Surg Oncol.* 1999;25:464–9.
- Cao D, Lin C, Woo SH, Vang R, Tsangaris TN, Argani P. Separate cavity margin sampling at the time of initial breast lumpectomy significantly reduces the need for reexcisions. *Am J Surg Pathol.* 2005;29:1625–32.
- Guidi AJ, Connolly JL, Harris JR, Shmitt SJ. The relationship between shaved margin and inked margin status in breast excision specimens. *Cancer.* 1997;70:1568–73.
- Rubin P, O'Honlon D, Browell D, Callanan K, Shrimankar J, Scott D, et al. Tumour bed biopsy detects the presence of multifocal disease in patients undergoing breast conservation therapy for primary breast carcinoma. *Eur J Surg Oncol.* 1996;22:23–6.
- Riedl O, Fitzal F, Mader N, Dubsy P, Rudas M, Mittlboeck M, et al. Intraoperative frozen section analysis for breast-conserving therapy in 1016 patients with breast cancer. *Eur J Surg Oncol.* 2009;35:264–70.
- Olson TP, Harter J, Munoz A, Mahvi DM, Breslin T. Frozen section analysis for intraoperative margin assessment during breast-conserving surgery results in low rates of re-excision and local recurrence. *Ann Surg Oncol.* 2007;14:2953–60.

24. Camp ER, McAuliffe PF, Gilroy JS, Morris CG, Lind DS, Mendenhall NP, et al. Minimizing local recurrence after breast conserving therapy using intraoperative shaved margins to determine pathologic tumor clearance. *J Am Coll Surg*. 2005;201:855–61.
25. Klimberg VS, Westbrook KC, Korourian S. Use of touch preps for diagnosis and evaluation of surgical margins in breast cancer. *Ann Surg Oncol*. 1998;5:220–6.
26. Valdes EK, Boolbol SK, Ali I, Feldman SM, Cohen JM. Intraoperative touch preparation cytology for margin assessment in breast-conservation surgery: does it work for lobular carcinoma? *Ann Surg Oncol*. 2007;14:2940–5.
27. Bakhshandeh M, Tutuncuoglu SO, Fischer G, Masood S. Use of imprint cytology for assessment of surgical margins in lumpectomy specimens of breast cancer patients. *Diagn Cytopathol*. 2007;35:656–9.
28. Weinberg E, Cox C, Dupont E, White L, Ebert M, Greenberg H, et al. Local recurrence in lumpectomy patients after imprint cytology margin evaluation. *Am J Surg*. 2004;188:349–54.
29. Veronesi U, Frante G, Galimberti V, Greco M, Luini A, Sacchini V, et al. Evaluation of resection margins after breast conservative surgery with monoclonal antibodies. *Eur J Surg Oncol*. 1991;17:338–41.
30. Giuliano AE, Hunt KK, Ballman KV, Beitsch PD, Whitworth PW, Blumencranz PW, et al. Axillary dissection vs no axillary dissection in women with invasive breast cancer and sentinel node metastasis. A randomized clinical trial. *JAMA*. 2011;305(6):569–75.
31. Tew K, Irwig L, Matthews A, Crowe P, Macaskill P. Metaanalysis of sentinel node imprint cytology in breast cancer. *Br J Surg*. 2005;92:1068–80.
32. Liu LC, Lang JE, Lu Y, Roe D, Hwang SE, Ewing CA, et al. Intraoperative frozen section analysis of sentinel lymph nodes in breast cancer patients. A meta-analysis and single-institution experience. *Cancer*. 2011;117:250–8.
33. Van Diest PJ, Torrenza H, Borgstein PJ, Pijpers R, Bleichrodt RP, Rahusen FD, et al. Reliability of intraoperative frozen section and imprint cytological investigation of sentinel lymph nodes in breast cancer. *Histopathology*. 1999;35:14–8.
34. Turner RR, Ollila DW, Krasne DL, Giuliano AE. Histopathologic validation of sentinel lymph node hypothesis for breast carcinoma. *Ann Surg*. 1997;226:271–6.
35. Viale G, Dell'orto P, Biasi MO, Stufano V, De Brito Lima LN, Paganelli G, et al. Comparative evaluation of an extensive histopathologic examination and a realtime reverse-transcription-polymerase chain reaction assay for mammaglobin and cytokeratin 19 on axillary sentinel lymph nodes of breast carcinoma patients. *Ann Surg*. 2008;247:136–42.
36. Klingler S, Marchal F, Rauch P, Kenouchi O, Chretien AS, Genin P, et al. Using one-step nucleic acid amplification (OSNA) for intraoperative detection of lymph node metastasis in breast cancer patients avoids second surgery and accelerates initiation of adjuvant therapy. *Ann Oncol*. 2013;24:2305–9.
37. Hughes SJ, Xi L, Raja S, Gooding W, Cole DJ, Gillanders WE, et al. A rapid, fully automated, molecular-based assay accurately analyzes sentinel lymph nodes for the presence of metastatic breast cancer. *Ann Surg*. 2006;243:389–98.
38. Tamaki Y, Akiyama F, Iwase T, Kaneko T, Tsuda H, Sato K, et al. Molecular detection of lymph node metastases in breast cancer patients: results of a multicenter trial using the one-step nucleic acid amplification assay. *Clin Cancer Res*. 2009;15:2879–84.
39. Lyman GH, Giuliano AE, Somerfield MR, Benson 3rd AB, Bodurka DC, Burstein HJ, et al. American Society of Clinical Oncology guideline recommendations for sentinel lymph node biopsy in early-stage breast cancer. *J Clin Oncol*. 2005;23:7703–20.
40. Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, et al. Prognostic factors in breast cancer: College of American Pathologists Consensus Statement 1999. *Arch Pathol Lab Med*. 2000;124:966–78.
41. Association of Directors of Anatomic and Surgical Pathology. Recommendations for processing and reporting of lymph node specimens submitted for evaluation of metastatic disease. *Am J Clin Pathol*. 2001;115:799–801.
42. Viale G, Bosari S, Mazzarol G, Galimberti V, Luini A, Veronesi P, et al. Intraoperative examination of axillary sentinel lymph nodes in breast carcinoma patients. *Cancer*. 1999;85:2433–8.
43. Turner RR. Histopathologic processing of the sentinel lymph node. *Semin Breast Dis*. 2002;5:35–40.
44. Weaver DL, Krag DN, Ashikaga T, Harlow SP, O'Connell M. Pathologic analysis of sentinel and nonsentinel lymph nodes in breast carcinoma: a multicenter study. *Cancer*. 2000;88:1099–107.
45. Cserni G. Metastases in axillary sentinel lymph nodes in breast cancer as detected by intensive histopathological work up. *J Clin Pathol*. 1999;52:922–4.
46. Turner RR, Ollila DW, Stern S, Giuliano AE. Optimal histopathologic examination of the sentinel lymph node for breast carcinoma staging. *Am J Surg Pathol*. 1999;23:263–7.
47. Zhang PJ, Reisner RM, Nangia R, Edge SB, Brooks JJ. Effectiveness of multiple-level sectioning in detecting axillary nodal micrometastasis in breast cancer: a retrospective study with immunohistochemical analysis. *Arch Pathol Lab Med*. 1998;122:687–90.
48. Weaver DL, Le UP, Dupuis SL, Weaver KA, Harlow SP, Ashikaga T, et al. Metastasis detection in sentinel lymph nodes: comparison of a limited widely spaced (NSABP protocol B-32) and a comprehensive narrowly spaced paraffin block sectioning strategy. *Am J Surg Pathol*. 2009;33:1583–9.
49. Kannurn S, Chewatanakornkul S, Premprapha T, Thongsuksai P. Comparability of different pathologic protocols in sentinel lymph node evaluation an analysis of two step-sectioning methods for the same patients with breast cancer. *Arch Pathol Lab Med*. 2009;133:1437–40.



50. Weaver DL. Pathologic evaluation of sentinel lymph nodes in breast cancer: a practical academic perspective from America. *Histopathology*. 2005;46:697–706.
51. Wells CA, Heryet A, Brochier J, Gatter KC, Mason DY. The immunocytochemical detection of axillary micrometastases in breast cancer. *Br J Cancer*. 1984;50:193–7.
52. Kollias J, Gil PG, Chatterton B, Raymond W, Collins PJ. Sentinel node biopsy in breast cancer: recommendations for surgeons, pathologists, nuclear physicians and radiologists in Australia and New Zealand. *Aust N Z J Surg*. 2000;70:132–6.
53. Cserni G. Histopathologic examination of the sentinel lymph nodes. *Breast J*. 2006;12(5 Suppl 2):S152–6.
54. Cserni G, Amendoeira I, Apostolikas N, Bellocq JP, Bianchi S, Boecker W, et al. Discrepancies in current practice of pathological evaluation of sentinel lymph nodes in breast cancer. Results of a questionnaire-based survey by the European Working Group for Breast Screening Pathology. *J Clin Pathol*. 2004;57:695–701.
55. Green FL, Page DL, Fleming ID, Fritz AG, Balch CM, Haller DG, et al. *AJCC cancer staging manual*. 6th ed. New York: Springer; 2002.
56. Van Rijk MC, Peterse JL, Nieweg OE, Oldenburg HS, Rutgers EJ, Kroon BB. Additional axillary metastases and stage migration in breast cancer patients with micrometastasis or submicrometastasis in sentinel lymph nodes. *Cancer*. 2006;107:467–71.
57. Rivera M, Merlin S, Hoda RS, Gopalan A, Hoda SA. Minimal involvement of sentinel lymph node in breast carcinoma: prevailing concepts and challenging problems. *Int J Surg Pathol*. 2004;12:301–6.
58. Biddle DA, Evans HL, Kemp BL, El-Naggar AK, Harvell JD, White WL, et al. Intraparenchymal nevus cell aggregates in lymph nodes: a possible diagnostic pitfall with malignant melanoma and carcinoma. *Am J Surg Pathol*. 2003;27:673–81.
59. Ichihara S, Ikeda T, Kimura K, Hanatate F, Yamada F, Hasegawa M, et al. Coincidence of mammary and sentinel lymph node papilloma. *Am J Surg Pathol*. 2008;32:784–92.
60. Diaz NM, Cox CE, Ebert M, Clark JD, Vrcel V, Stowell N, et al. Benign mechanical transport of breast epithelial cells to sentinel lymph nodes. *Am J Surg Pathol*. 2004;28(12):1641–5.
61. Moore KH, Thaler HT, Tan LK, Borgen BJ, Cody 3rd HS. Immunohistochemically detected tumor cells in the sentinel lymph nodes of patients with breast carcinoma: biologic metastasis or procedural artifact? *Cancer*. 2004;100:929–34.
62. Nagi C, Bleiweiss I, Jaffer S. Epithelial displacement in breast lesions: a papillary phenomenon. *Arch Path Lab Med*. 2005;129:1465–9.
63. Cox CE, Haddad F, Bass S, Cox JM, Ku NN, Berman C, et al. Lymphatic mapping in the treatment of breast cancer. *Oncology*. 1998;12:1283–92.
64. Bass SS, Lyman GH, McCann CR, Ku NN, Berman C, Durand K, et al. Lymphatic mapping and sentinel lymph node biopsy. *Breast J*. 1999;5:288–95.
65. Pendas S, Dauway E, Giuliano R, Ku N, Cox CE, Reintgen DS. Sentinel node biopsy in ductal carcinoma in situ patients. *Ann Surg Oncol*. 2000;7:15–20.
66. Yi M, Krishnamurthy S, Kuerer HM, Meric-Bernstam F, Bedrosian I, Ross MI, et al. Role of primary tumor characteristics in predicting positive sentinel lymph nodes in patients with ductal carcinoma in situ or microinvasive breast cancer. *Am J Surg*. 2008;196:81–7.
67. Ballehaninna UK, Chamberlain RS. Utility of intraoperative frozen section examination of sentinel lymph nodes in ductal carcinoma in situ of the breast. *Clin Breast Cancer*. 2013;13:350–8.
68. Miller AR, Thomason VE, Yeh IT, Alrahan A, Sharkey FE, Stauffer J, et al. Analysis of sentinel lymph node mapping with immediate pathologic review in patients receiving preoperative chemotherapy for breast carcinoma. *Ann Surg Oncol*. 2002;9:243–7.
69. Nason KS, Anderson BO, Byrd DR, Dunnwald LK, Eary JF, Mankoff DA, et al. Increased false negative sentinel node biopsy rates after preoperative chemotherapy for invasive breast carcinoma. *Cancer*. 2000;89:2187–94.
70. Tafta L, Verbanac KM, Lannin DR. Preoperative chemotherapy and sentinel lymphadenectomy for breast cancer. *Am J Surg*. 2001;182:312–5.
71. Julian TB, Patel N, Dusi D, Olson P, Nathan G, Jasnosz K, et al. Sentinel lymph node biopsy after neoadjuvant chemotherapy for breast cancer. *Am J Surg*. 2001;182:407–10.
72. Haid A, Tausch C, Lang A, Lutz J, Fritzsche H, Peschina W, et al. Is sentinel lymph node biopsy reliable and indicated after preoperative chemotherapy in patients with breast carcinoma? *Cancer*. 2001;92:1080–4.
73. Fernandez A, Cortes M, Benito E, Azpeitia D, Prieto L, Moreno A, et al. Gamma probe sentinel node localization and biopsy in breast cancer patients treated with a neoadjuvant chemotherapy scheme. *Nucl Med Commun*. 2001;22:361–6.
74. Balch GC, Mithani SK, Richards KR, Beauchamp RD, Kelley MC, et al. Lymphatic mapping and sentinel lymphadenectomy after preoperative therapy for stage II and III breast cancer. *Ann Surg Oncol*. 2003;10:616–21.
75. Ozmen V, Unal ES, Muslumanoglu ME, Igci A, Canbay E, Ozcinar B, et al. Axillary sentinel node biopsy after neoadjuvant chemotherapy. *Eur J Surg Oncol*. 2010;36:23–9.
76. Stearns V, Ewing CA, Slack R, Penannen MF, Hayes DF, Tsangaris TN. Sentinel lymphadenectomy after neoadjuvant chemotherapy for breast cancer may reliably represent the axilla except for inflammatory breast cancer. *Ann Surg Oncol*. 2002;9:235–42.
77. Mamounas EP, Brown A, Anderson S, Smith R, Julian T, Miller B, et al. Sentinel node biopsy after neoadjuvant chemotherapy in breast cancer: results from National Surgical Adjuvant Breast and Bowel Project Protocol B-27. *J Clin Oncol*. 2005;23:2694–702.

Sennur Ilvan

## Abstract

Fibroepithelial tumors of the breast represent a heterogeneous group of biphasic tumors composed of a proliferation of epithelial and stromal components. Fibroadenomas and phyllodes tumors constitute the major entities.

These tumors are one of the most challenging diagnostic lesions for pathologists. It can be difficult to make a clear distinction microscopically between fibroadenomas and benign phyllodes tumors. There are no reliable morphological features or immunohistochemical markers predicting phyllodes tumors.

## Keywords

Breast • Biphasic tumor • Phyllodes tumor • Fibroadenoma • Tubular adenoma • Hamartoma • Pathology

## Fibroepithelial Tumors

Fibroepithelial tumors are characterized by a simultaneous proliferation of both epithelial and mesenchymal elements. Fibroadenomas and phyllodes tumors constitute the major entities. Several breast lesions, such as hamartoma, fibroadenomatoid hyperplasia, and tubular adenoma, are also included in this category.

---

S. Ilvan, MD  
Department of Pathology,  
Cerrahpasa School of Medicine, Istanbul University,  
Fatih, Istanbul, Turkey  
e-mail: [sennurilvan@hotmail.com](mailto:sennurilvan@hotmail.com)

## Fibroadenoma

Fibroadenomas are the most common benign breast tumors in women 20–30 years but may be present at any age. They originate in the terminal duct lobular unit. The initial processes of epithelial and stromal proliferation in multiple lobules either occur spontaneously or are precipitated by hyperestrogenic states. A gradual confluence of the hyperplastic lobules is followed by the formation of fibroadenomatoid nodules. Finally, the fibroadenomatous nodules coalesce to form the fibroadenoma [1].

Fibroadenomas typically present as palpable, slow-growing, firm, mobile masses and are

generally less than 3 cm in size. Less frequently, some of these tumors, particularly the juvenile variant, may reach very large dimensions. If rapid growth occurs, the lesion may undergo infarction, particularly in pregnant and lactating women, and may be mistaken for malignancy. Fibroadenomas are usually solitary, but in some patients, bilateral and multiple lesions can be observed [2]. Because fibroadenomas originate in the terminal duct lobular unit, which is absent in the normal male mammary duct system, fibroadenomas are rarely observed in the male breast [3].

Most fibroadenomas are clinically identifiable, but in 25 % of cases, they are nonpalpable and can be diagnosed by mammography and ultrasound. Small-sized impalpable fibroadenomas are detected as mammographic nodular densities or as calcified lesions. Ultrasonography shows a well-defined homogeneous hypoechoic mass [2, 4].

Some chromosomal abnormalities have been described in fibroadenomas, such as the deletion of 17p and numerical changes in chromosomes 16, 17, and 21 [5]. Clonality studies have demonstrated the polyclonality of both stroma and epithelium, although monoclonality has been demonstrated in stromal areas of phyllodes tumors [1]. Because the stroma and epithelium of fibroadenomas exhibit estrogen and progesterone receptor expression, it is believed that fibroadenomas develop as a result of unopposed estrogenic influences [1, 6].


Patients who receive cyclosporin A for immunosuppression after organ transplantation are reported to be predisposed to fibroadenoma development [7]. The risk of developing breast carcinoma is dependent on the presence of proliferative changes in the fibroadenoma itself or in the surrounding breast tissue, as well as a family history of breast carcinoma [4].

Grossly, fibroadenomas are ovoid and well circumscribed. The cut surface is gray or white, solid, rubbery, and often somewhat lobulated. Slit-like spaces corresponding to epithelial structures upon microscopic examination may also be recognized. Rarely, the cut surface may appear myxoid. In older patients, the lesion may be fibrotic and calcified [2].

Microscopically, fibroadenomas consist of an admixture of stromal and epithelial elements. Two growth patterns are recognized. The intracanalicular pattern is characterized by the proliferation of stromal cells around stretched and compressed ducts (Fig. 15.1). In the pericanalicular pattern, stroma surrounds glandular structures with open lumina. These patterns often coexist and are thought to have no clinical significance. The glands have an inner epithelial layer and an outer myoepithelial layer, which are morphologically similar to normal breast ducts. The epithelial component may exhibit typical ductal hyperplasia, sclerosing adenosis, or benign metaplastic changes, such as apocrine or squamous metaplasia. Rarely, lobular or ductal carcinoma in situ and invasive carcinoma may be present within a fibroadenoma [2, 8]. The stroma in fibroadenomas is variable in both appearance and the degree of cellularity; it may be sclerotic, cellular, or myxoid. Usual fibroadenomas exhibit collagenized stroma with low cellularity. Spindle-shaped stromal cells have bland ovoid to elongated nuclei and do not display cytological atypia. Mitotic figures are rare but may be present, particularly in young and pregnant women. A hyalinized, sclerotic stroma is observed more frequently in fibroadenomas in older women. Chondroid, osseous, and smooth muscle metaplasia can very rarely occur in the stroma. Infrequently, scattered enlarged multinucleated stromal giant cells may be present. These cells are benign and should not be misinterpreted as evidence of malignancy [9].

Fibroadenomas with increased stromal cellularity and without leaflike processes are referred to as cellular fibroadenomas. The stroma may display prominent myxoid changes, particularly in younger women. Such myxoid fibroadenomas have been reported in association with Carney syndrome [2]. Juvenile fibroadenoma is observed mostly in patients younger than 20 years of age and is characterized by rapid growth, massive size, cellular stroma, a pericanalicular epithelial growth pattern, and marked epithelial hyperplasia with irregular tufts similar to that observed in gynecomastia [10]. Fibroadenomas that contain cysts greater than 3 mm, sclerosing adenosis, epithelial calcifications, or papillary apocrine

**Fig. 15.1** Fibroadenoma, intracanalicular pattern (HE, 40×)



hyperplasia are considered complex fibroadenomas. Complex fibroadenomas are associated with a slightly increased risk of subsequent development of breast cancer compared with simple fibroadenomas [11].

An intracanalicular fibroadenoma with increased stromal cellularity can be misinterpreted as a benign phyllodes tumor. The distinction is based on the degree of stromal cellularity and the presence of frond-like projections into cystic spaces (leaflike appearance). The distinction is particularly difficult in core-needle biopsy. In this situation, a descriptive diagnosis that leads to excision is appropriate.

### **Fibroadenomatous Change (Fibroadenomatoid Hyperplasia)**

Fibroadenomatous change is a term used to describe histological changes in the breast similar to those observed in fibroadenomas but in which no discrete mass is formed.

### **Tubular Adenoma**

Many authorities consider tubular adenoma to be a variant of fibroadenoma. These relatively rare lesions usually occur in young women. On gross

examination, tubular adenomas are softer and typically tan-brown in color. They are circumscribed nodular lesions composed of closely packed, small, round acinar structures surrounded by scant, loose cellular stroma [2].

### **Mammary Hamartoma**

These lesions occur at any age but are most commonly observed in premenopausal and perimenopausal women. The clinical features are similar to those of fibroadenoma. Hamartomas are composed of mammary ducts, lobules, collagenous stroma, and adipose tissue. In contrast to normal breast tissue, these structures are disorganized and irregularly dispersed. Because of their similarity to normal breast tissue, histological diagnosis may be impossible. In these situations, mammographic findings are required for correct diagnosis. Smooth muscle may be present, and lesions with marked smooth muscle fibers have been termed myoid hamartomas [9].

### **Phyllodes Tumor**

Phyllodes tumors are uncommon fibroepithelial breast neoplasms that account for 0.3–1.0 % of all primary breast tumors. Johannes Mueller, who

first described phyllodes tumors in 1838, called them cystosarcoma phyllodes, “sarcoma” for the fleshy nature of the lesion and “phyllodes” for the leaflike appearance [1]. This term is currently considered inappropriate because the tumors are rarely cystic and most of them follow a benign course. The World Health Organization (WHO) currently supports the term “phyllodes tumor” in its classification of breast tumors. These tumors are classified as benign, borderline, and malignant based on a variety of histopathological variables [9]. Whether phyllodes tumors arise from preexisting fibroadenomas or develop de novo is controversial. It has been speculated that phyllodes tumors begin as fibroadenomas, and subsequently a single stromal cell undergoes mutation and develops into a phyllodes tumor [12].

While phyllodes tumors can be observed at any age, including adolescence, they occur most frequently in women 40–50 years of age. Few cases have been reported in the male breast [3]. Phyllodes tumors usually present as unilateral, painless, lobulated, and freely movable breast masses and rarely cause skin changes, such as ulceration, nipple retraction, or nipple discharge. It is generally difficult to clinically differentiate them from fibroadenoma. Patients with phyllodes tumors usually have a history of large, rapidly growing masses [13].

Mammography reveals a rounded or lobulated, sharply defined opaque mass in most cases. It is not possible to reliably distinguish between benign and malignant phyllodes tumors by mammography or ultrasonography [13].

Macroscopically, phyllodes tumors vary in size from a few centimeters to 20 cm in diameter, with an average size of 4–5 cm. Although malignant variants tend to be larger than benign lesions, there is no consistent relationship between tumor size and malignancy. Similar to fibroadenomas, phyllodes tumors have well-delineated borders but appear fleshier, with bulging cut surfaces. Clefts and cystic spaces may be observed, particularly in larger tumors. Malignant lesions may be less well defined, and the clefts may be less obvious. Hemorrhage and necrosis are more common in borderline and malignant tumors [2, 13].

Microscopically, phyllodes tumors are composed of epithelium and cellular stroma arranged


in an intracanalicular pattern. Projections of this cellular stroma into cystic spaces form the leaflike pattern characteristic of phyllodes tumors (Fig. 15.2). The morphology of the stromal component varies from only slightly more cellular than that of a fibroadenoma to frankly sarcomatous. In addition, heterogeneity in stromal cellularity within a phyllodes tumor is common. Stromal cellularity and mitotic activity are often increased in periglandular areas. The epithelial component may exhibit usual ductal hyperplasia or metaplastic changes, such as squamous or apocrine [1, 2].

Based on the WHO classification, phyllodes tumors are classified as benign, borderline, and malignant according to stromal cellularity, cellular atypia and pleomorphism, mitotic index, tumor margin, stromal overgrowth, and the presence or absence of heterologous differentiation (Table 15.1) [9].

Benign phyllodes tumors typically have pushing borders and mild stromal cellularity, with little or no atypia (Fig. 15.2). The stroma is usually condensed around the epithelial component. Mitotic figures are infrequent; they rarely exceed 1–2 per 10 high-power fields and are generally located in the periductal areas. Stromal overgrowth, defined as at least one microscopic field observed at 4× magnification that contains only stroma without epithelial components, is not present. Benign heterologous stromal elements, such as skeletal muscle, cartilage, and bone, may be present. The presence of bizarre stromal giant cells should not be taken as a mark of malignancy. Benign phyllodes tumors can be difficult to distinguish from cellular fibroadenoma. Compared to fibroadenoma, phyllodes tumors are characterized in most cases by expansion and the increased cellularity of the stromal component. In addition, numerous leaflike projections into cystic spaces and stromal mitosis favor a diagnosis of phyllodes tumor. When there is histological ambiguity, the WHO Working Group recommends a diagnosis of fibroadenoma instead of phyllodes tumor. Some authors advocate using the term “benign fibroepithelial neoplasm” with an explanation of the diagnostic difficulty [9].



**Fig. 15.2** Benign phyllodes tumor with a leaflike structure (HE, 100×)



**Table 15.1** Histological features used in the classification of phyllodes tumor subtypes [9]

	Benign	Borderline	Malignant
Tumor border	Well defined	Well defined, focally infiltrative	Infiltrative
Stromal cellularity	Mild	Moderate	Marked
Stromal cell atypia	None to mild	Mild to moderate	Marked
Mitotic activity	<5/10 HPF	5–9/10 HPF	≥10/10 HPF
Stromal overgrowth	Absent	Absent or very focal	Often present
Malignant heterologous elements	Absent	Absent	May be present

*HPF* high-power fields

Borderline phyllodes tumors exhibit a relatively expanded stromal component and pronounced intracanalicular growth pattern compared with benign forms. Stromal atypia is mild to moderate, and mitotic figures range from 5 to 9 per 10 high-power fields. Tumor margins may be pushing or focally infiltrative. Stromal overgrowth is absent or very focal.


In malignant phyllodes tumors, the stroma is more expanded and highly cellular, and the tumor borders are infiltrative. The stromal cells exhibit marked atypia and brisk mitotic activity (≥10 mitoses/10 high-power fields). Stromal overgrowth is common and may be so prominent that the appearance mimics a primary breast sarcoma. In such cases, extensive sampling is required for correct diagnosis. Although the stroma most often resembles a fibrosarcoma, heterologous stromal differentiation, such as liposarcomatous,

chondrosarcomatous, or osteosarcomatous elements, can be observed (Fig. 15.3).

The local recurrence rates are 10–17 %, 14–25 %, and 23–30 % for benign, borderline, and malignant phyllodes tumors, respectively. Recurrence rates are strongly correlated with the amount of normal breast tissue present at the resection margins. Histological parameters predicting recurrence are stromal overgrowth, atypia, and mitotic activity. Axillary node involvement is rare, but hematogenous spread occurs in the lung, bone, and liver. Metastases generally consist of stromal component alone. Histological type is the most important predictor of metastatic spread, and most distant metastases develop from malignant tumors [9, 13].

The expression levels of many biological markers, such as p53, c-kit, CD10, β-catenin, vascular endothelial growth factor, and CD34,

**Fig. 15.3** Malignant phyllodes tumor showing liposarcomatous differentiation (HE, 100×)



have been explored in phyllodes tumors. Most of them are variably expressed depending on the grade of the tumor. However, the clinical value of these markers is limited in predicting the behavior and classification of phyllodes tumors [14, 15].

## References

1. Tavassoli FA, Eusebi V. Tumors of the mammary gland. Silver Spring: ARP Press; 2009. p. 315–40.
2. Pinder SE, Mulligan AM, O'Malley FP. Fibroepithelial lesions, including fibroadenoma and phyllodes tumors. In: O'Malley FP, Pinder SE, Mulligan AM, editors. Breast pathology. 2nd ed. Philadelphia: Elsevier Saunders; 2011. p. 121–38.
3. Ansah-Boateng Y, Tavassoli FA. Fibroadenoma and cystosarcoma phyllodes of the male breast. *Mod Pathol.* 1992;5:114–6.
4. El-Wakeel H, Umpleby HC. Systematic review of fibroadenoma as risk factor for breast cancer. *Breast.* 2003;12:302–7.
5. Barbosa ML, Ribeiro EM, Silva GF, Maciel ME, Lima RS, Cavalli LR, et al. Cytogenetic findings in phyllodes tumor and fibroadenomas of the breast. *Cancer Genet Cytogenet.* 2004;154:156–9.
6. Sapino A, Bosco M, Cassoni P, Castellano I, Arisio R, Cserni G, et al. Estrogen receptor-beta is expressed in stromal cells of fibroadenoma and phyllodes tumors of the breast. *Mod Pathol.* 2006;19:599–606.
7. Iaria G, Pisani F, De Luca L, Sforza D, Manuelli M, Perone L, et al. Prospective study of switch from cyclosporine to tacrolimus for fibroadenomas of the breast in kidney transplantation. *Transplant Proc.* 2010;42:1169–70.
8. Limite G, Esposito E, Sollazzo V, Ciancia G, Formisano C, Di Micco R, et al. Lobular intraepithelial neoplasia arising within breast fibroadenoma. *BMC Res Notes.* 2013;6:267.
9. Tan PH, Tse G, Lee A. Fibroepithelial tumors. In: Lakhani SR, Ellis IO, Schnitt SJ, Tan PH, van de Vijver MJ, editors. WHO classification of tumours of the breast. 4th ed. Lyon: IARC Press; 2012. p. 141–7.
10. Ng WK, Mrad MA, Brown MH. Juvenile fibroadenoma of the breast: treatment and literature review. *Can J Plast Surg.* 2011;19:105–7.
11. Sklair-Levy M, Sella T, Alweiss T, Craciun I, Libson E, Mally B. Incidence and management of complex fibroadenomas. *Am J Roentgenol.* 2008;190:214–8.
12. Abe M, Miyata S, Nishimura S, Iijima K, Makita M, Akiyama F, et al. Malignant transformation of breast fibroadenoma to malignant phyllodes tumor: long-term outcome of 36 malignant phyllodes tumors. *Breast Cancer.* 2011;18:268–72.
13. Parker SJ, Harries SA. Phyllodes tumours. *Postgrad Med J.* 2001;77:428–35.
14. Ho SK, Thike AA, Cheok PY, Tse GMK, Tan PH. Phyllodes tumours of the breast: the role of CD34, vascular endothelial growth factor and  $\beta$ -catenin in histological grading and clinical outcome. *Histopathology.* 2013;63:393–406.
15. Korcheva VB, Levine J, Beadling C, Warrick A, Countryman G, Olson NR, et al. Immunohistochemical and molecular markers in breast phyllodes tumors. *Appl Immunohistochem Mol Morphol.* 2011;19:119–25.

Zerrin Calay

**Abstract**

A variety of reactive and neoplastic lesions of the breast are characterized by a proliferation of spindle cells. The pathologist must be aware of the clinical, radiological, and morphological overlap between reactive and neoplastic spindle cell lesions of the breast. In addition, metaplastic (spindle cell) carcinoma is far more common than spindle cell sarcoma in the breast.

Among the vascular lesions of the breast, angiosarcoma is more common and may appear very bland, simulating hemangioma.

Core biopsy samples must be evaluated very carefully to interpret spindle and vascular lesions. In general, excision is recommended because of morphological overlap, and clinicopathological correlation is necessary for correct diagnosis.

**Keywords**

Breast • Mesenchymal tumor • Pseudoangiomatous stromal hyperplasia • Myofibroblastoma • Fibromatosis • Nodular fasciitis • Vascular lesions

**Mesenchymal Tumors of the Breast**

Many neoplastic and nonneoplastic mesenchymal lesions have been reported in the breast. In this chapter, only lesions that are observed more frequently or are important for differential diagnosis will be discussed.

Z. Calay, MD  
Department of Pathology, Cerrahpaşa School of  
Medicine, Istanbul University, Istanbul, Turkey  
e-mail: [zcalay@gmail.com](mailto:zcalay@gmail.com)

**Pseudoangiomatous Stromal  
Hyperplasia (PASH)**

PASH is a benign myofibroblastic proliferation that simulates a vascular lesion. This proliferation is thought to represent an exaggerated myofibroblastic response to endogenous or exogenous hormonal stimuli, particularly progesterone [1].

PASH is commonly observed as an incidental microscopic finding in breast biopsies or in association with other breast lesions, such as fibroadenomas, phyllodes tumors, and gynecomastia.

Rarely, this proliferation can be florid and form a clinically evident mass mimicking fibroadenoma. In these cases, the terms “nodular or tumorous PASH” have been used [2]. Gross examination of tumorous PASH reveals a round-oval, well-circumscribed, rubbery mass with a homogenous cut surface that is tan to white in color. Typical PASH is characterized by thin, slit-like spaces set in a densely hyalinized stroma. The spaces are often lined by bland spindled myofibroblasts that may mimic the endothelial cells of blood vessels. Occasionally, PASH is more cellular with a fascicular growth pattern that is similar to myofibroblastoma. Multinucleated giant cells have been reported in some examples of PASH [3]. Cytologic atypia and mitotic activity of the myofibroblasts may rarely be observed [4]. The myofibroblasts exhibit staining for CD34 and variable staining for SMA (smooth muscle actin) and desmin.

PASH is a benign lesion that is adequately treated by local excision, although recurrences have been reported [1]. The major importance of PASH is that it must be distinguished histologically from a true vascular lesion, specifically, low-grade angiosarcoma.

---

## Myofibroblastoma

Myofibroblastoma is an uncommon benign tumor of the breast composed of fibroblasts and myofibroblasts. Clinically, myofibroblastomas are slow-growing, circumscribed, mobile masses that are usually mistaken for fibroadenomas on physical examination and mammography. Myofibroblastoma is now considered part of the same spectrum as spindle cell lipoma [1, 5], with which it shares the same chromosomal rearrangements affecting region 13q [6].

Gross examination reveals a rubbery-firm, oval-round nodule with a whorled cut surface. A variety of patterns may be observed histologically. Myofibroblastoma is usually composed of uniform, oval-spindle cells arranged as short fascicles admixed with hyalinized bands of collagen. Morphological variants are reported as epitheloid, cellular, infiltrative, decidual, colla-

genized, lipomatous, and myxoid types [7]. This morphological variation may cause diagnostic difficulties.

Differential diagnosis includes a variety of reactive and benign spindle cell lesions, spindle cell sarcoma, and carcinoma. Myofibroblastoma cells typically exhibit CD34 and desmin expression, with variable positivity for actin, bcl-2, and CD99. The expression of ER, PR, and androgen receptors is common. Therefore, an epitheloid variant of myofibroblastoma may be confused with invasive lobular carcinoma because both tumors express hormone receptors [8].

Myofibroblastomas are benign and are adequately treated by local excision.

---

## Desmoid-Type Fibromatosis

Mammary fibromatosis (desmoid type) is an infiltrative, locally aggressive proliferation of fibroblasts and myofibroblasts that typically presents as a firm mass that may be mistaken for carcinoma clinically and mammographically. It can occur within the breast parenchyma but frequently arises from the pectoral fascia and extends into the breast. There is an association with previous trauma, particularly surgery and including implants [9]. Desmoid-type fibromatosis occasionally occurs in patients with familial adenomatous polyposis (FAP) [1].

The margins of the lesion are infiltrative and difficult to determine on gross examination. The cut surface reveals a firm consistency with a gray-white to tan appearance. Fibromatosis is characterized by broad sheets of spindle cells producing a storiform or herringbone pattern. The lesion irregularly infiltrates the adjacent breast parenchyma. Nuclear expression of beta-catenin is observed in approximately 80 % of cases [1]; however, beta-catenin can also be expressed in spindle cell carcinoma, phyllodes tumors, and fibrosarcoma. CD34 expression is absent.

Scarring from prior trauma, such as surgery, and fibromatosis, such as metaplastic carcinoma (spindle cell carcinoma), are the most important lesions for the differential diagnosis of fibroma-

tosis. An immunohistochemistry panel should be performed using a broad panel of antibodies to keratins, including those of high molecular weight [1].

The major clinical concern in patients with fibromatosis is local recurrence, which is observed in 20–30 % of cases. Wide local excision is required to prevent local recurrence.

---

## Nodular Fasciitis

Nodular fasciitis is a self-limited, mass-forming fibroblastic-myofibroblastic proliferation that is clonal [7]. Nodular fasciitis is very uncommon in the breast but is important to recognize because it may clinically, radiographically and histologically mimic a malignant tumor. Typically, it grows rapidly and may be tender and painful. These lesions regress spontaneously within a few months [1].

Nodular fasciitis arises in the subcutis or, less often, the mammary parenchyma and is a well-circumscribed grayish nodule that may display central cystic changes. It is composed of short, spindle-shaped cells loosely arranged in a “tissue culture” fashion. Mitoses are typically frequent. They form solid masses and do not infiltrate around ducts and lobules. The cells are negative for CD34 and beta-catenin but positive for actin and focally positive for desmin.

The major differential diagnostic considerations are malignant spindle cell tumors (including spindle cell carcinomas and sarcomas) and fibromatosis.

Although nodular fasciitis will spontaneously regress, in general, local excision is performed for definitive diagnosis. Local excision is curative in nearly all cases [1, 7].

---

## Inflammatory Myofibroblastic Tumor

Inflammatory myofibroblastic tumor (IMT) is a low-grade neoplasm consisting of myofibroblastic spindle cells mixed with prominent inflamma-

tory cells, usually plasmacytes. IMT is very rare in the breast [10].

IMT in the breast presents as a painless, circumscribed firm mass. The majority of lesions are benign with a local recurrence rate of 10–25 % [1].

---

## Vascular Lesions of the Breast

Vascular lesions of the breast are a heterogeneous group. They can be either benign or malignant; however, vascular lesions with atypical but not frankly malignant features have also been described.

### Benign Vascular Lesions

Benign vascular lesions of the mammary parenchyma are relatively uncommon. These include perilobular hemangioma, hemangioma (capillary, cavernous and venous), and angiomatosis. In addition, there are a variety of benign vascular lesions that involve the subcutaneous tissue [4].

The major clinical importance of these lesions is that they must be distinguished from angiosarcoma. Some angiosarcomas are very well differentiated and simulate benign blood vessels; conversely, some benign vascular lesions may exhibit atypia. Occasionally, differential diagnosis is impossible, particularly if it is a core biopsy; surgical excision should be performed for exclusion.

### Atypical Vascular Lesions

Atypical vascular lesions of the breast are angioformative proliferations that can develop in the skin after BCS and radiation therapy for breast carcinoma [11]. These lesions typically present as small papules or plaques with pink, red, or brown discoloration. They are characterized by dilated vessels in the dermis that often have complex branching and anastomosing areas. Although they have a benign clinical course, some may be precursors of angiosarcoma [12].



## Angiosarcoma

Angiosarcoma is the most frequent primary sarcoma of the breast, with an incidence of approximately 0.05 % of breast malignancies. It may arise spontaneously (primary angiosarcoma) or following radiation therapy for breast cancer (secondary angiosarcoma).

Angiosarcoma of the mammary parenchyma presents as a painless mass, and those that involve the skin may appear as areas of bluish-red discoloration (Fig. 16.1).

Histologically, these lesions are characterized by anastomosing vascular channels that dissect the adipose tissue and mammary stroma (Fig. 16.2). The endothelial cells have atypical hyperchromatic nuclei, but endothelial multilayering and mitoses are often absent. Poorly differentiated angiosarcoma has a more solid, cellular

growth pattern, typically with spindled morphology and reduced formation of vascular channels (Fig. 16.3). Angiosarcomas are classified as low, intermediate, and high grades based on a combination of histological features, although grading has no prognostic value [13].

Differential diagnoses for well-differentiated angiosarcoma include PASH, angioliipoma, and benign vascular lesions. Poorly differentiated angiosarcoma must be differentiated from spindle cell carcinoma and other sarcomas.

Angiosarcomas require complete excision and are usually treated by mastectomy. The prognosis is generally poor, particularly for secondary angiosarcomas [4].

## Other Mesenchymal Lesions

Benign mesenchymal tumors that occur elsewhere in the body have been described in the breast, including lipoma, angioliipoma, leiomyoma, neurofibroma, schwannoma, and granular cell tumor of the breast. Granular cell tumor is a benign tumor derived from Schwann cells of peripheral nerves; however, they may appear malignant clinically (irregular and firm mass), radiologically (ill-defined spiculated lesion), and pathologically (infiltrative growth pattern) [1].


Malignant mesenchymal tumors of the breast other than angiosarcomas are extremely rare. After angiosarcoma, the second-most common type of primary sarcoma is liposarcoma. However, any type of sarcoma may occur in the breast as a primary lesion, including fibrosarcoma, leiomyosarcoma, rhabdomyosarcoma, malignant peripheral nerve sheath tumor, and osteosarcoma.

A sarcomatous-appearing tumor in the breast is far more likely to be a metaplastic carcinoma (spindle cell carcinoma) or a malignant phyllodes tumor than a primary sarcoma. Therefore, extensive sampling of the tumor and immunostaining using multiple anti-cytokeratin antibodies are essential for correct diagnosis.




**Fig. 16.1** Angiosarcoma. In this mastectomy specimen, the skin has areas of bluish-red discoloration

**Fig. 16.2** Angiosarcoma, low grade. (Well-formed, interanastomosing vascular spaces are present. The endothelial cells lining the vascular channels show mild nuclear hyperchromasia, but there is no evidence of endothelial cell tufting.) HE, 100×



**Fig. 16.3** Angiosarcoma, high grade. (This tumor has a more solid, cellular growth pattern, typically with spindled morphology and reduced formation of vascular channels.) HE, 200×



## References

1. Lakhani SR, Ellis IO, Schnitt SJ, Tan PH, Vijver MJ. Mesenchymal tumours. In: WHO classification of tumours of the breast. 4th ed. Lyon: IARC; 2012. p. 125–39.
2. Wieman SM, Landercasper J, Johnson JM, Ellis RL, Wester SM, Lambert PJ, et al. Tumoral pseudoangiomatous stromal hyperplasia of the breast. *Am Surg*. 2008;74:1211–4.
3. Comunoğlu N, Comunoğlu C, Ilvan S, Calay Z, Muslumanoglu M. Mammary pseudoangiomatous stromal hyperplasia composed of predominantly giant cells: an unusual variant. *Breast J*. 2007;13:568–70.
4. Schnitt JS, Collins LC. Biopsy interpretation of the breast. 2nd ed. Philadelphia: Wolters Kluwer/Lippincott Williams & Wilkins; 2013. p. 387–408.
5. Magro G, Bisceglia M, Michal M, Eusebi V. Spindle cell lipoma-like tumor, solitary fibrous tumor and myofibroblastoma of the breast: a clinico-pathological analysis of 13 cases in favor of a unifying histogenetic concept. *Virhows Arch*. 2002;440:249–60.
6. Pauwels P, Sciort R, Croiset F, Rutten H, Van den Berghe H, Dal CP. Myofibroblastoma of the breast: genetic link with spindle cell lipoma. *J Pathol*. 2000;191:282–5.

7. Shin SJ, Rabban JT. Mesenchymal neoplasms of the breast. In: Dabbs DJ, editor. *Breast pathology*. 1st ed. Philadelphia: Elsevier-Saunders; 2012. p. 596–642.
8. Magro G. Epitheloid-cell myofibroblastoma of the breast: expanding the morphologic spectrum. *Am J Surg Pathol*. 2009;33:1085–92.
9. Jamshed S, Farhan MI, Marchal MB, Nahabedian MY, Liu MC. Fibromatosis of the breast after mammary prosthesis implantation. *Clin Adv Hematol Oncol*. 2008;6:687–94.
10. Haj M, Weiss M, Loberant N, Cohen I. Inflammatory pseudotumor of the breast: case report and literature review. *Breast J*. 2003;9:423–5.
11. Weaver J, Billings SD. Postradiation cutaneous vascular tumors of the breast: a review. *Semin Diagn Pathol*. 2009;26:141–9.
12. Patton KT, Deyrup AT, Weiss SW. Atypical vascular lesions after surgery and radiation of the breast: a clinicopathologic study of 32 cases analyzing histologic heterogeneity and association with angiosarcoma. *Am J Surg Pathol*. 2008;32:943–50.
13. Nascimento AF, Raut CP, Fletcher CD. Primary angiosarcoma of the breast: clinicopathologic analysis of 49 cases, suggesting that grade is not prognostic. *Am J Surg Pathol*. 2008;32:1896–904.

---

# Index

## A

Abrikosov, A.I., 38  
Adejolu, M., 19  
Allen, E., 162  
Alltundag, M.K., 207–224, 229–236  
Anatomy, 1–12  
Andacoglu, O., 48  
Arslan, C., 207–224  
Arun, B.K., 57–78  
Ashworth, T, 232  
Aspiration, 18, 19, 21, 38, 61, 103, 104, 115–117, 232, 252, 268, 278  
Axilla, 1–7, 11, 12

## B

Bayraktar, S., 57–78  
BCS. *See* Breast-conserving surgery (BCS)  
Beatson, M.D., 162  
Benign breast disease, 19–20  
Berg, J.C., 48  
Berg, W.A., 115  
Bianchi, S., 48  
Bidard, F.C., 231, 233  
Biphasic tumor, 20  
BI-RADS. *See* Breast imaging reporting and data system (BI-RADS)  
Bone marrow, 229–236  
Boyd, S., 162  
Bracketing, 118, 119, 136  
Braun, S., 231  
BRCA, 61, 62, 64–72, 75, 77, 78, 101, 103, 106, 132, 146, 148, 149, 151, 152  
Breast  
  imaging, 16, 18–21, 28, 33, 36, 37, 40, 89–120, 134  
  inflammation, 17–18  
Breast-conserving surgery (BCS), 126, 135–140, 268, 270, 271, 291  
Breast imaging reporting and data system (BI-RADS), 16, 95, 100  
Breast-specific gamma imaging (BSGI), 114

Brennan, S.B., 34  
Brenner, R.J., 48  
Bunting, D.M., 48  
Byrski, T., 70

## C

Calay, Z., 289–293  
Cawson, J.N., 48  
Chemoprevention, 68, 70, 72–78, 127, 128  
Circulating tumor cells (CTCs), 157, 229–236  
Clinical aspects, 153, 161–177  
Computed tomography (CT), 18, 94, 212–214, 216–217, 236  
Cox, C.E., 271  
Cribriform carcinoma, 192, 248  
Cristofanilli, M., 233, 235  
CT. *See* Computed tomography (CT)  
CTCs. *See* Circulating tumor cells (CTCs)  
Cytology, 17, 20, 61, 115, 117, 172, 245, 251, 256, 268, 270–272, 274, 279, 284

## D

Dean, P.B., 44  
Development, 1–3, 7–8, 127, 128, 132, 192, 246  
Diabetic mastopathy, 17, 20  
Diagnostic, 33, 35, 90–95, 103, 113, 116, 128, 135, 141, 195, 223, 254, 262, 268, 271, 286  
Diego, E.J.B., 23–53  
Differential diagnosis, 20, 26–27, 31, 35–36, 47–48, 101, 243–244, 253–255, 275, 276, 278, 289–292  
Dixon, J.M., 251  
3-D Mammogram, 113–115  
Doisy, E.A., 162  
Domchek, S.M., 69  
Ductal carcinoma in situ (DCIS), 20, 30, 60, 61, 65, 66, 76, 77, 95, 97, 99, 101–103, 107–109, 112, 118, 119, 125–128, 131–141, 170, 242–248, 253, 254, 258, 261, 278, 284

**E**

- Early breast cancer, 123, 126, 189, 208–216, 218–224, 231, 234  
 E-cadherin, 64, 125, 126, 128, 148, 150, 152, 244, 246, 247, 252  
 Elastography, 103, 114  
 Embryology, 1–3  
 Epigenetics, 64, 147–149  
 Estrogen, 7, 48, 61, 98, 128, 132, 146, 162, 192, 209, 244, 284  
 Estrogen and progesterone receptors and Her2/neu, 8, 61, 140, 146, 156, 161–177, 192, 218, 284  
 Extramedullary hematopoiesis

**F**

- Falck, A.K., 231  
 Fat necrosis, 16, 34–36, 47, 98, 101  
 Fibroadenoma, 19, 21, 23–27, 36, 93, 95, 104, 105, 255, 283–286, 289, 290  
 Fibromatosis, 248, 258–259, 290–291  
 Franken, B., 233  
 Frozen sections, 268–273  
 Fu, C.Y., 34

**G**

- Ganott, M., 89–120  
 Gene arrays, 207–224  
 Giordano, A., 233  
 Giuliano, M., 233  
 Granular cell tumor, 38–43, 292  
 Griffith, B., 89–120  
 Gynecomastia, 8, 34, 48–53, 98, 101, 102, 282, 289

**H**

- Hamartoma, 36–38, 283, 285  
 Hamperl, H., 43  
 Hartkopf, A.D., 231  
 Hayes, D.F., 233  
 Hereditary breast cancer, 64–66, 146  
 HER-2 testing, 140, 252  
 High-risk screening, 90, 101, 109  
 Histopathology, 25, 176, 177, 187–189, 197, 199, 208, 213, 214, 219, 241, 242, 249, 262, 270, 271, 273–277  
 Holley, S.O., 34

**I**

- IDC. *See* Invasive ductal carcinoma (IDC)  
 Ignatiadis, M., 233  
 IHC. *See* Immunohistochemistry (IHC)  
 Ilvan, S., 283–288  
 Immunohistochemistry (IHC), 25, 65, 125, 152, 171–172, 175–176, 188, 190, 199, 222, 230, 231, 244, 252, 255, 267, 274, 275, 279, 291  
 Intraductal papilloma, 20, 27–34, 60, 113  
 Invasive breast cancer, 6, 58, 65–67, 72, 73, 75, 106, 124–128, 138, 140, 187–202

- Invasive ductal carcinoma (IDC), 65, 101, 132, 153, 191, 193, 194, 198, 248, 250, 259, 270, 277  
 Involution, 9, 19

**J**

- Jaffer, S., 34  
 Jensen, E.V., 162  
 Jiang, Z.F., 233

**K**

- Kibil, W., 34  
 Kilickap, S., 229–236  
 Klimberg, V.S., 271

**L**

- Lactation, 8, 17–19, 25, 28, 29, 95, 284  
 Laé, M., 260  
 Lakhani, S.R., 249  
 Lale Dogan, A., 145–159  
 LCIS. *See* Lobular carcinoma in situ (LCIS)  
 Lee, C.H., 111  
 Lee, E.H., 66  
 Lee, K.A., 48  
 Linda, A., 48  
 Lipoma, 3, 31–36, 40, 290, 292  
 Liu, L.C., 272  
 Lobular carcinoma in situ (LCIS), 20, 61, 67, 69, 73, 74, 76, 123–128, 153, 167, 177, 191, 192, 243, 246–248, 252, 270, 273–277  
 Lopez-Medina, A., 48  
 Lucci, A., 234

**M**

- Magnetic resonance imaging (MRI), 19, 37, 40, 41, 44, 45, 47, 66–68, 72, 95, 103, 106–113, 115, 118, 122, 124, 127, 134, 135  
 Mammogram, 8, 20, 21, 24, 25, 31, 33–35, 37, 44, 45, 50, 51, 63, 72, 74, 90–95, 98–103, 105–108, 110, 112–114, 117, 118, 124, 127, 132, 134, 139, 141  
 Manfrin, E., 48  
 Mastectomy, 7, 10–12, 65, 66, 68, 69, 76, 109, 112–113, 123, 127, 128, 133, 135–140, 292  
 MBI. *See* Molecular breast imaging (MBI)  
 MC. *See* Medullary carcinoma (MC)  
 McAuliffe, P., 123–128, 131–141  
 McGuire, K.P., 1–12  
 Medullary carcinoma (MC), 65, 248, 254  
 Mesenchymal tumor, 289–293  
 Microinvasion, 133–134, 136, 140, 244  
 Micrometastases, 188, 189, 229–236, 272  
 MicroRNAs (miRNA), 153–155  
 miRNA. *See* MicroRNAs (miRNA)  
 Molecular breast imaging (MBI), 114, 117  
 Molecular portrait, 148–153  
 Molino, A., 231  
 Morgan, C., 48



Mucinous carcinoma, 198, 248, 255, 256, 260, 261  
 Müller, V., 233  
 Myofibroblastoma, 289–291

## N

Nipple discharge, 16, 17, 19, 28, 29, 50, 113, 133, 261, 286  
 Nodular fasciitis, 291  
 Nuclear medicine breast imaging, 114

## O

Ozisik, Y.Y., 229–236

## P

Page, D.L., 244  
 Papillary carcinoma, 31, 192, 248, 257, 261  
 PASH. *See* Pseudoangiomatous stromal hyperplasia (PASH)  
 Pastor-Barriuso, R., 67  
 Pathology, 19–20, 24–25, 30, 34–35, 37, 41–42, 45–46, 52–53, 60–61, 65, 99, 108, 118, 124–125, 132–136, 171, 189, 241–263  
 Patterson, J.A., 48  
 PEM. *See* Positron emission mammography (PEM)  
 Perou, C.M., 149, 199, 201  
 Pfeiffer, R.M., 67  
 Phyllodes tumors (PTs), 20, 283–290, 292  
 Physiology, 1–12, 17, 18, 48–49, 107, 114, 163, 173  
 Pierga, J.Y., 233  
 Poland's syndrome, 2  
 Positron emission mammography (PEM), 114–115  
 Prat, A., 149  
 Prediction of treatment, 208, 213, 222–223  
 Prevention, 52, 57–78, 118, 127, 162, 163, 197, 291  
 Progesterone, 8, 9, 18, 65, 81, 146, 148, 156, 161–175, 192, 212, 219, 284, 289  
 Prognosis, 138, 150, 153, 166, 167, 169, 173, 187–189, 192, 195–197, 199, 212, 214, 230–232, 234–236, 254, 255  
 Prognostic factors, 138, 150, 153, 166, 167, 173, 187, 189, 192, 195–197, 201, 202, 230–232, 234, 235, 255  
 Prophylactic mastectomy, 65, 68  
 Prophylactic oophorectomy, 69, 72  
 Pseudoangiomatous stromal hyperplasia (PASH), 20, 38, 289–290  
 PTs. *See* Phyllodes tumors (PTs)

## R

Rack, B.K., 233, 234, 236  
 Radial scar, 43–53, 95, 97, 135, 244, 253  
 Radioactive Seed, 12, 118, 119, 126, 135, 136  
 Rahka, E.A., 48  
 Rakha, E.A., 252  
 Raloxifen, 69, 72, 74–75, 77, 127, 128  
 Ready, K.J., 67

Resetskova, E., 48  
 Risk factors, 16, 18, 20, 25, 48, 57–78, 127, 128, 132, 135, 150, 213, 216  
 Risk models, 67  
 Rizzo, M., 34  
 Rudzinski, S.M., 89–120

## S

Sahin, A.A., 187–202  
 Sanchez, M.R., 32  
 Sari, E., 161–177  
 Sauter, E.R., 15–21  
 Schnitt, S.J., 244  
 Sentinel lymph node biopsy, 11, 12, 127, 136–140  
 Sentinel node biopsy, 11–12  
 Signal transduction pathways, 146, 151, 165–166, 172–175  
 Singletary, S.E., 61  
 Skaane, P., 114  
 Smerage, J.B., 233  
 Somatic genetic alterations, 146–147  
 Sorlie, T., 199  
 Stereotactic, 95, 99, 104, 112, 115, 118, 119, 135  
 Survival, 16, 19, 60, 66, 68–71, 133, 134, 136, 140, 149–150, 153, 154, 156, 164–168, 173–174, 188–190, 195, 199, 208, 210, 235, 236, 250, 252–255, 262, 271  
 Susceptibility genes, 65, 145

## T

Tabar, L., 44  
 Tamoxifen, 52, 68, 69, 71, 73–78, 127, 128, 135, 138, 139, 150, 153, 168, 169, 173, 202, 208–213, 218, 219, 221, 223  
 Tavassoli, F.A., 244, 257  
 TC. *See* Tubular carcinoma (TC)  
 Tew, K., 272  
 Tognon, C., 260  
 Tomosynthesis, 2, 94, 95, 97, 113–114  
 Toth, D., 48  
 Townsend, C.M., 3, 7  
 Tubular adenoma, 25, 27, 28, 283, 285  
 Tubular carcinoma (TC), 65, 192, 245, 248, 251–254, 276  
 Tumor biomarkers, 187  
 Turner, R.R., 272  
 Tuzlali, S., 241–263

## U

Ultrasound (US), 19, 24, 25, 29, 30, 33, 34, 37, 38, 41, 44, 72, 92–94, 97, 101–106, 110, 113, 115–118, 127, 134, 135, 284

## V

Van Diest, P.J., 272  
 Vascular lesions, 28–292

Veronesi, U., 271  
Viale, G., 273

**W**

Wallwiener, M., 233, 235  
Weinberg, E., 271  
Wire localization, 25, 118–120  
Wu, Y., 187–202

**X**

Xenidis, N., 233

**Y**

Yalcin, S., 161–177  
Yavuz, E., 267–280  
Youk, J.H., 118