
Acquired Demyelinating Diseases

Alex Rovira and Kelly K. Koeller

Introduction

Demyelinating disorders of the central nervous system (CNS) have a variety of etiologies and can be separated into primary (e.g., multiple sclerosis (MS) and other idiopathic inflammatory-demyelinating diseases) and secondary (e.g., infectious, ischemic, metabolic, or toxic) diseases. Brain and spinal cord magnetic resonance imaging (MRI) is the imaging modality of choice to assess demyelinating disorders and, together with the clinical and laboratory findings, can accurately classify them in most cases [1–3]. This review will highlight the important imaging manifestations of some acquired demyelinating diseases that allow more specific diagnosis.

Idiopathic Inflammatory-Demyelinating Diseases of the CNS

The term idiopathic inflammatory-demyelinating disease (IIDD) encompasses a broad spectrum of CNS disorders that can be differentiated according to their severity, clinical course, and lesion distribution, as well as their imaging, laboratory, and pathological findings. The spectrum includes monophasic, multiphasic, and progressive disorders, ranging from highly localized forms to multifocal or diffuse variants [2]. Relapsing-remitting and secondary progressive MS are the two most common forms of IIDD. MS can also have a progressive course from onset (primary progressive and progressive-relapsing MS). Fulminant forms of IIDD include

a variety of disorders that have in common the severity of the clinical symptoms, an acute clinical course, and atypical findings on MRI. The classic fulminant IIDD is Marburg disease. Balo concentric sclerosis, Schilder disease, and acute disseminated encephalomyelitis (ADEM) can also present with severe, acute attacks.

Some IIDDs have a restricted topographic distribution, as is the case with neuromyelitis optica spectrum disorders (NMOSDs), which can have a monophasic or, more often, a relapsing course.

Multiple Sclerosis

MS is a progressive inflammatory, demyelinating, and neurodegenerative autoimmune disease characterized pathologically by perivascular infiltrates of mononuclear inflammatory cells, demyelination, and axonal loss and gliosis, with the formation of focal and diffuse abnormalities in the brain and spinal cord, mainly affecting the optic nerves, brainstem, spinal cord, and cerebellar and periventricular white matter, although cortical and subcortical gray matter damage is also prominent, resulting in chronic progressive disability for the majority of people with the disorder.

The high sensitivity of MRI in depicting brain and spinal cord demyelinating plaques has made this technique the most important paraclinical tool in current use, not only for the early and accurate diagnosis of MS but also for understanding the natural history of the disease and monitoring and predicting the efficacy of disease-modifying treatments.

MRI is the most sensitive imaging technique for detecting MS plaques throughout the brain and spinal cord. Proton density (PD) or T2-weighted MR images (especially those of fluid-attenuated inversion recovery (FLAIR) sequence) show areas of high signal intensity in the periventricular white matter in 98 % of MS patients. MS plaques are generally round to ovoid in shape and range from a few millimeters to more than one centimeter in diameter. They are typically discrete and focal at the early stages of the disease but become

À. Rovira
Section of Neuroradiology and MR Unit (IDI), Department of Radiology, Hospital Universitari Vall d'Hebron, Autonomous University of Barcelona, Barcelona, Spain
e-mail: Alex.rovira@idi.gencat.cat

K.K. Koeller (✉)
Section of Neuroradiology, Department of Radiology, Mayo Clinic, Rochester, MN, USA
e-mail: Koeller.Kelly@mayo.edu

confluent as the disease progresses, particularly in the posterior hemispheric periventricular white matter. MS plaques tend to affect the deep white matter rather than the subcortical white matter, whereas small vessel ischemic lesions tend to involve the subcortical white matter more than the periventricular white matter [4, 5]. The total T2 lesion volume of the brain increases by approximately 5 %–10 % each year in the relapsing forms of MS.

Both acute and chronic MS plaques appear hyperintense on T2-weighted sequences, reflecting their increased tissue water content. The signal increase indicates edema, inflammation, demyelination, reactive gliosis, and/or axonal loss in proportions that differ from lesion to lesion. The vast majority of MS patients have at least one ovoid periventricular lesion, whose major axis is oriented perpendicular to the outer surface of the lateral ventricles. The ovoid shape and perpendicular orientation are derived from the perivenular location of the demyelinating plaques noted on histopathology (Dawson's fingers).

MS lesions tend to affect specific regions of the brain, including the periventricular white matter situated superolateral to the lateral angles of the ventricles, the calloso-septal interface along the inferior surface of the corpus callosum, the cortico-juxtacortical regions, and the infratentorial regions. Focal involvement of the periventricular white matter in the anterior temporal lobes is typical for MS and rarely seen in other white matter disorders. The lesions commonly found at the calloso-septal interface are best depicted by sagittal T2-FLAIR images; so this sequence is highly recommended for diagnostic MRI studies.

Histopathological studies have shown that a substantial portion of the total brain lesion load in MS is located within the cerebral cortex. Presently available MRI techniques are not optimal for detecting cortical lesions because of poor contrast resolution between normal-appearing gray matter (NAGM) and the plaques in question and because of the partial volume effects of the subarachnoid spaces and CSF surrounding the cortex. Cortical lesions are better visualized by 2D or 3D T2-FLAIR sequences and newer MR techniques such as 3D double inversion recovery (DIR) MR sequences which selectively suppress the signal from white matter and cerebrospinal fluid (CSF). Juxtacortical lesions that involve the "U" fibers are seen in two-thirds of patients with MS. They are a rather characteristic finding in early stages of the disease and are best detected by T2-FLAIR sequences.

Posterior fossa lesions preferentially involve the floor of the fourth ventricle, the middle cerebellar peduncles, and the brainstem. Most brainstem lesions are contiguous with the cisternal or ventricular CSF spaces and range from large confluent patches to solitary, well-delineated paramedian lesions or discrete "linings" of the CSF border zones. Predilection for these areas is a key feature that helps to identify MS plaques and to differentiate them from focal areas of ischemic

demyelination and infarction that preferentially involve the central pontine white matter. Because of their short acquisition time and greater sensitivity, T2-weighted sequences are preferred over T2-FLAIR sequences for detecting posterior fossa lesions [6].

Approximately 10–20 % of T2 hyperintensities are also visible on T1-weighted images as areas of low signal intensity compared with normal-appearing white matter. These so-called T1 black holes have a different pathological substrate that depends, in part, on the lesion age. The hypointensity is present in up to 80 % of recently formed lesions and probably represents marked edema, with or without myelin destruction or axonal loss. In most cases, the acute (or wet) "black holes" become isointense within a few months as inflammatory activity abates, edema resolves, and reparative mechanisms like remyelination become active. Less than 40 % evolve into persisting or chronic black holes [7], which correlate pathologically with the most severe demyelination and axonal loss, indicating areas of irreversible tissue damage. Chronic black holes are more frequent in patients with progressive disease than in those with relapsing-remitting disease and more frequent in the supratentorial white matter as compared with the infratentorial white matter. They are rarely found in the spinal cord and optic nerves.

MS lesions of the spinal cord resemble those in the brain. The lesions can be focal (single or multiple) or diffuse and mainly affect the cervical cord segment. On sagittal scans, the lesions characteristically have a cigar shape and rarely exceed two vertebral segments in length. On cross section, they typically occupy the lateral and posterior white matter columns, extend to involve the central gray matter, and rarely occupy more than one half the cross-sectional area of the spinal cord [8].

Acute spinal cord lesions can produce a mild-to-moderate mass effect with spinal cord swelling and may show contrast enhancement. Active lesions are rarer in the spinal cord than the brain and are almost always associated with new clinical symptoms. The prevalence of spinal cord abnormalities is as high as 74–92 % in established MS and depends on the clinical phenotype of MS. In a clinically isolated syndrome (CIS), the prevalence of spinal cord lesions is lower, particularly if there are no referable symptoms such as myelopathy. Nevertheless, asymptomatic spinal cord lesions are found in 30–40 % of patients with a CIS. In relapsing-remitting MS, the spinal cord lesions are typically multifocal. In secondary progressive MS, the abnormalities are more extensive and diffuse and are commonly associated with spinal cord atrophy. In primary progressive MS, spinal cord abnormalities are quite extensive as compared with brain abnormalities. This discrepancy may help to diagnose primary progressive MS in patients with few or no brain abnormalities.

Longitudinal and cross-sectional MR studies have shown that the formation of new MS plaques is often associated

with contrast enhancement, mainly in the acute and relapsing stages of the disease [9]. The gadolinium enhancement varies in size and shape but usually lasts from a few days to weeks, although steroid treatment shortens this period. Incomplete ring enhancement on T1-weighted gadolinium-enhanced images, with the open border facing the gray matter of the cortex or basal ganglia, is a common finding in active MS plaques and is a helpful feature for distinguishing between inflammatory-demyelinating lesions and other focal lesions such as tumors or abscesses [10].

Focal enhancement can be detected before abnormalities appear on unenhanced T2-weighted scans and can reappear in chronic lesions with or without a concomitant increase in size. Although enhancing lesions also occur in clinically stable MS patients, their number is much greater when there is concomitant clinical activity. Contrast enhancement is a relatively good predictor of further enhancement and of subsequent accumulation of T2 lesions but shows no (or weak) correlation with progression of disability and development of brain atrophy. In relapsing-remitting and secondary progressive MS, enhancement is more frequent during relapses and correlates well with clinical activity. For patients with primary progressive MS, serial T2-weighted studies show few new lesions and little or no enhancement with conventional doses of gadolinium, despite steady clinical deterioration. Contrast-enhanced T1-weighted images are routinely used in the study of MS to provide a measure of inflammatory activity in vivo. The technique detects disease activity 5–10 times more frequently than clinical evaluation of relapses, suggesting that most of the enhancing lesions are clinically silent. Subclinical disease activity with contrast-enhancing lesions is four to ten times less frequent in the spinal cord than the brain, a fact that may be partially explained by the large volume of brain as compared with the spinal cord. High doses of gadolinium and a long postinjection delay can increase the detection of active spinal cord lesions.

Multiple Sclerosis Variants

Marburg Disease

Marburg disease (also termed malignant MS) is a rare, acute MS variant that occurs predominantly in young adults. It is characterized by a confusional state, headache, vomiting, gait unsteadiness, and hemiparesis. This entity has a rapidly progressive course with frequent, severe relapses leading to death or severe disability within weeks to months after onset of the clinical signs, mainly due to brainstem involvement or mass effect with herniation. Most patients who survive subsequently develop a relapsing form of MS. Because Marburg disease is often preceded by a febrile illness, it can also be

considered a fulminant form of ADEM when it has a monophasic course. At the time of the clinical presentation, it may be difficult to find good predictors of whether the patient will have a fulminant course, develop mild or severe MS, or even develop MS at all. Pathologically, Marburg lesions are more destructive than those of typical MS or ADEM and are characterized by massive macrophage infiltration, demyelination (not restricted to the perivascular areas), hypertrophic astrocytes, and severe axonal injury, with features compatible with the histomorphologic diagnosis of a severe, acute demyelinating disease. Despite the destructive nature of these lesions, areas of remyelination are often observed [11]. In Marburg disease, MRI typically shows multiple focal T2 lesions of varying sizes, which may coalesce to form large white matter plaques disseminated throughout the hemispheric white matter and brainstem. Perilesional edema is often present, and enhancement is commonly seen. A similar imaging pattern is seen in ADEM.

Schilder Disease

Schilder disease is a rare acute or subacute disorder that can be defined as a specific clinical-radiologic presentation of MS. It commonly affects children and young adults. The clinical spectrum of Schilder disease includes psychiatric predominance, acute intracranial hypertension, intermittent exacerbations, and progressive deterioration. Imaging studies show large ring-enhancing lesions involving both hemispheres, sometimes symmetrically, and located preferentially in the parieto-occipital regions. These large, focal demyelinating lesions can resemble a brain tumor, an abscess, or even an adrenoleukodystrophy. MRI features that suggest possible Schilder disease include large and relatively symmetric involvement of the brain hemispheres, incomplete ring enhancement, minimal mass effect, restricted diffusivity, and sparing of the brainstem [2, 12]. Histopathologically, Schilder disease consistently shows well-demarcated demyelination and reactive gliosis with relative sparing of the axons. Microcystic changes and even frank cavitation can occur. The clinical and imaging findings usually show a dramatic response to steroids.

Balo Concentric Sclerosis

Balo concentric sclerosis is a rare IIDD subtype, considered a variant of MS, with characteristic radiologic and pathologic features. The disease was formerly considered an aggressive MS variant, leading to death in weeks to months after onset, and in which the diagnosis was made on histopathologic findings at postmortem examination. However, with the widespread use of MRI, this MS variant is often

identified in patients who later have a complete or almost complete clinical recovery [13]. The pathologic hallmarks of the disease are large demyelinated lesions showing a peculiar pattern of alternating layers of preserved and destroyed myelin. One possible explanation for this pattern is that sublethal tissue injury is induced at the edge of the expanding lesion, which would then stimulate expression of neuroprotective proteins to protect the rim of periplaque tissue from damage, thereby resulting in alternating layers of preserved and nonpreserved myelinated tissue [14]. These alternating bands are best identified with T2-weighted sequences, which typically show thick concentric hyperintense bands corresponding to areas of demyelination and gliosis, alternating with thin isointense bands corresponding to normal myelinated white matter. This pattern can be also identified on T1-weighted images as alternating isointense (preserve myelin) and hypointense (demyelinated) concentric rings. These bands, which may eventually disappear over time, can appear as multiple concentric layers (onion skin lesion), as a mosaic, or as a “floral” configuration. The center of the lesion usually shows no layering because of massive demyelination (“storm center”). Contrast enhancement and decreased diffusivity are common in the outer rings (inflammatory edge) of the lesion [13].

The Balo pattern can be isolated, multiple, or combined with typical MS-like lesions, and the lesion structure can vary from one or two to several alternating bands, with a total size from one to several centimeters. Lesions occur predominantly in the cerebral white matter, although brainstem, cerebellum, and spinal cord involvement has also been reported.

Neuromyelitis Optica Spectrum Disorders

Neuromyelitis optica (NMO) is an autoimmune inflammatory disorder of the CNS with a predilection for the optic nerves and spinal cord. The discovery of NMO-IgG and an NMO-specific autoantibody directed against aquaporin-4 (AQP4-Ab), the major water channel in the CNS, clearly identified NMO as a disease separate from MS.

This uncommon and topographically restricted form of IIDD is characterized by severe unilateral or bilateral optic neuritis and complete transverse myelitis, which occur simultaneously or sequentially over a varying period of time (weeks or years). AQP4-Ab has been also demonstrated in patients with conditions other than classic NMO, including isolated longitudinally extensive transverse myelitis, defined by lesions spanning over more than three segments; monophasic or recurrent isolated optic neuritis; and certain types of brainstem encephalitis (particularly if the diencephalon or medulla oblongata is affected). As most of these patients later develop NMO, various groups have suggested classifying these symptoms as high-risk syndromes for NMO when

they occur in AQP4-Ab-seropositive patients. In addition, it has been proposed that AQP4-Ab-positive classic NMO and AQP4-Ab-positive high-risk syndromes be referred to as NMO-spectrum disorder (NMOSD) or autoimmune AQP4 channelopathy [15]. The incidence and prevalence of NMO are unknown, and there are important differences in its regional distribution worldwide.

The index events of new-onset NMO are severe unilateral or bilateral optic neuritis, acute myelitis, or a combination of these symptoms. The myelitis attacks appear as complete transverse myelitis with severe bilateral motor deficits, sensory-level, bowel and bladder dysfunction, pain and significant residual neurologic injury. Optic neuritis attacks are generally more severe than those typically seen in MS. Approximately 85 % of patients have a relapsing course with severe acute exacerbations and poor recovery, which leads to increasing neurologic impairment and a high risk of respiratory failure and death due to cervical myelitis. Patients who experience acute optic neuritis and transverse myelitis simultaneously or within days of each other are much more likely to have a monophasic course. On the other hand, a relapsing course correlates with NMO-IgG seropositivity, a longer interval between attacks, older age at onset, female gender, and less severe motor impairment after the myelitic onset. Although the initial attacks are more severe in patients proven to have monophasic NMO, the long-term neurologic prognosis is somewhat better in this group because patients do not accumulate disability from recurrent attacks.

Clinical features alone are insufficient to diagnose NMO; CSF analysis and MRI are usually required to confidently exclude other disorders. CSF pleocytosis (>50 leukocytes/mm³) is often present, while oligoclonal bands are seen less frequently (20–40 %) than in MS patients (80–90 %). AQP4-Ab detection is reported to have a sensitivity of 73 % and a specificity of 91 % for NMO. AQP4-Ab may be helpful to distinguish this form of IIDD from MS and it can predict relapse and conversion to NMO in patients presenting with a single attack of longitudinally extensive myelitis. NMO-IgG is positive in 52 % of patients with relapsing transverse myelitis and in 25 % of patients with recurrent idiopathic optic neuritis [16].

Wingerchuk et al. [17] proposed a revised set of criteria for diagnosing NMO. These criteria remove the absolute restriction on CNS involvement beyond the optic nerves and spinal cord, allow any interval between the first events of optic neuritis and transverse myelitis, and emphasize the specificity of longitudinally extensive spinal cord lesions on MRI and NMO-IgG-seropositive status. More recently the International Panel for NMO diagnosis developed new diagnostic criteria that define the unifying term NMOSD, which is stratified by serologic testing (with or without AQP4-IgG) [18]. These new criteria require, in patients with AQP4-IgG, core clinical and MRI findings related to optic nerve, spinal

cord, area postrema, other brainstem, diencephalic, or cerebral presentations. However, more stringent clinical and MRI criteria are required for diagnosis of NMO-spectrum disorders without AQP4-IgG or when serologic testing is unavailable.

MRI of the affected optic nerve demonstrates swelling and loss of blood–brain barrier integrity with gadolinium enhancement that can extend into the optic chiasm. The spinal cord lesions in NMO typically extend over three or more contiguous vertebral segments and occasionally the entire spinal cord (longitudinally extensive spinal cord lesions); they are centrally located (preferential central gray matter involvement) and affect much of the cross-section on axial images. During the acute and subacute phase, the lesions are tumefactive and show contrast uptake. In some cases, the spinal cord lesions are small at the onset of symptoms, mimicking those of MS, and then progress in extent over time. The presence of very hyperintense spotty lesions on T2-weighted images (“bright spotty sign”) is a specific feature that helps differentiate NMO from MS, particularly in patients without longitudinally extensive spinal cord lesions [19], and likely reflects the highly destructive component of the inflammatory lesion. Spinal cord lesions can progress to atrophy and necrosis, and may lead to syrinx-like cavities on T1-weighted images.

NMO was long considered a disease without brain involvement, and a negative brain at disease onset was considered a major supportive criterion for the diagnosis of NMO. However, various studies have shown that brain MRI abnormalities exist in a significant proportion (50–85 %) of patients [20]. Brain MRI lesions are often asymptomatic but sometimes are associated with symptoms even at disease onset. The brain lesions are commonly nonspecific. They can be dot-like or patchy, <3 cm in diameter, and located in the deep white matter, brainstem, or cerebellum. Nonetheless, some brain MRI features appear to be quite characteristic and distinct from MS lesions. These abnormalities may have parallel sites with high AQP4 expression adjacent to the ventricular system at any level, such as the hypothalamus, periventricular areas surrounding the third and lateral ventricles, cerebral aqueduct, corpus callosum, and dorsal brainstem adjacent to the fourth ventricle. The appearance of periventricular lesions in NMO is quite characteristic. In contrast to MS, where periventricular lesions are discrete, oval-shaped, and perpendicular to the ependymal lining due to their periventricular distribution (Dawson’s fingers), NMO lesions are not oval-shaped, located immediately adjacent to the lateral ventricles following the ependymal lining in a disseminated pattern, and are often edematous and heterogeneous [20]. As opposed to what occurs in MS, NMO lesions do not affect the cortical gray matter.

Involvement of the corpus callosum has been described in 18 % of AQP4-seropositive NMO patients. The lesions are

multiple, large, and edematous, show heterogeneous signal intensity on T2-weighted images, and sometimes affect the entire thickness of the corpus callosum.

Lesions may also affect areas where AQP4 expression is not particularly high, such as the corticospinal tracts. These lesions, which can be unilateral or bilateral and may affect the posterior limb of the internal capsule and cerebral peduncle of the midbrain, are contiguous and often longitudinally extensive [20].

Other brain MRI findings described in NMO include extensive and confluent hemispheric white matter lesions and radial hemispheric lesions (sometimes corresponding to an extension of periventricular lesions), which are likely related to vasogenic edema involving the white matter tracts. These lesions usually do not show mass effect or contrast enhancement but there may be a “cloud-like” pattern of enhancement, defined as multiple patches of enhancing lesions with blurred margins [20].

Some of the typical brain MRI findings may be specific to clinical presentations, such as intractable vomiting and hiccup (linear dorsal medullary lesions involving the area postrema and nucleus tractus solitarius) or a syndrome of inappropriate antidiuretic hormone secretion (hypothalamic and periaqueductal lesions) [21].

Distinguishing NMO from MS is critical, particularly in the early stages, since the treatment and prognosis of these disorders differ. In fact, some evidence suggests that MS-modifying treatments such as interferon- β , natalizumab, and laquinimod exacerbate NMO. By contrast, several immunosuppressants (e.g., azathioprine, rituximab, mitoxantrone) seem to help in preventing NMO relapses.

NMOSD can be associated with systemic autoimmune diseases such as systemic lupus erythematosus and Sjögren syndrome). Whether the neurologic manifestations are solely due to NMO-spectrum disorder or are a manifestation of these diseases is controversial, although optic neuritis and transverse myelitis are rare presentations of them, and several studies have shown that patients with systemic autoimmune diseases and NMO-IgG-positive antibodies always have optic neuritis, myelitis, or NMO.

Acute Disseminated Encephalomyelitis

ADEM is a severe, immune-mediated inflammatory disorder of the CNS that predominantly affects the white matter of the brain and spinal cord. In the absence of specific biologic markers, the diagnosis of ADEM is based on clinical and radiologic features. This disorder affects children more commonly than adults, and, in contrast to MS, shows no gender preponderance. The estimated incidence is 0.8 per 100,000 population per year. In most cases, the clinical onset of disease is preceded by viral or bacterial infections, usually

nonspecific upper respiratory tract infections. ADEM may also develop following a vaccination (postimmunization encephalomyelitis). Patients commonly present with non-specific multifocal symptoms, which developed subacutely over a period of days, frequently associated with encephalopathy (relatively uncommon in MS), defined as an alteration in consciousness (e.g., stupor, lethargy) or a behavioral change unexplained by fever, systemic illness, or postictal symptoms. Although ataxia, encephalopathy, and brainstem symptoms are frequently present in both pediatric and adult cases, certain signs and symptoms appear to be age related. In childhood ADEM, long-lasting fever and headaches occur more frequently, while in adult cases, motor and sensory deficits predominate. In general, the disease is self-limiting and the prognostic outcome favorable.

Although ADEM usually has a monophasic course, multiphasic forms have been reported, raising diagnostic difficulties in distinguishing these cases from MS. This multiphasic form, which accounts for less than 4 % of ADEM cases, is defined as a new encephalopathic event consistent with ADEM, separated by 3 months after the initial illness but not followed by any further events. The second ADEM event can involve either new or re-emergent neurologic symptoms, signs, and MRI findings. Relapsing disease following ADEM that occurs beyond a second encephalopathic event is no longer consistent with multiphasic ADEM but rather indicates a chronic disorder, most often leading to the diagnosis of MS or NMO [22].

An ADEM event as the first manifestation of the classic relapsing form of MS occurs in 2–10 %. According to the International Pediatric Multiple Sclerosis Study Group, the diagnosis of MS is met if, after the initial ADEM, a second clinical event meets the following three requirements: [1] it is nonencephalopathic; [2] it occurs 3 months or more after the incident neurologic illness; and [3] it is associated with new MRI findings consistent with the 2010 McDonald criteria for dissemination in space [22]. The presence of hypointense lesions and two or more periventricular lesions is an MRI feature that supports an MS diagnosis in children with acute CNS demyelination [23]. Unlike the lesions in MS, ADEM lesions are often large, patchy, and poorly marginated on MRI. There is usually asymmetric involvement of the subcortical and central white matter and cortical gray-white junction of the cerebral hemispheres, cerebellum, brainstem, and spinal cord. Lesions confined to the periventricular white matter and corpus callosum are less common than in MS. The gray matter of the thalamus and basal ganglia is often affected, particularly in children, and typically in a symmetric pattern [40]. However, the frequency of thalamic involvement in adult ADEM does not differ from that of adult MS. This can be explained by the fact that involvement of this structure is less common in adult ADEM than in childhood ADEM. Four patterns of cerebral involvement

have been proposed to describe the MRI findings in ADEM [24]: (1) ADEM with small lesions (less than 5 mm); (2) ADEM with large, confluent, or tumefactive lesions and frequent extensive perilesional edema and mass effect; (3) ADEM with additional symmetric deep gray matter involvement; and (4) acute hemorrhagic encephalomyelitis. Gadolinium enhancement of one or more lesions occurs in 14–30 % of cases [22]. The pattern of enhancement varies and can be complete or incomplete ring shaped, nodular, gyral, or spotty. Although ADEM is usually a monophasic disease, new lesions may be seen on follow-up MRI within the 1st month of the initial attack.

Most MRI lesions appear early in the course of the disease, supporting the clinical diagnosis of ADEM. Nonetheless, in some cases, there may be a delay of more than 1 month between the onset of symptoms and the appearance of lesions on MRI. Therefore, a normal brain MRI obtained within the first days after the onset of neurologic symptoms suggestive of ADEM does not exclude this diagnosis.

The spinal cord is affected in less than 30 % of ADEM patients [24], predominantly in the thoracic region. The spinal cord lesion is typically large, causes swelling, and shows variable enhancement. In most ADEM patients, partial or complete resolution of the MRI abnormalities occurs within a few months of treatment. This course is positively associated with a final diagnosis of ADEM.

Infectious Inflammatory-Demyelinating Disorders

Progressive Multifocal Leukoencephalopathy

Progressive multifocal leukoencephalopathy (PML) is overwhelmingly a disease of the immunocompromised patient, and most (55–85 %) cases are related to acquired immunodeficiency syndrome (AIDS). There is a wide age range of involvement, with the peak age of presentation in the sixth decade. The disease is caused by reactivation of a papovavirus (JC virus) that selectively attacks the oligodendrocyte, leading to demyelination. Treatment with monoclonal antibody therapy (natalizumab, rituximab, efalizumab) or other immunomodulatory drugs, commonly used in patients with MS and other disorders, has also been linked with PML [25]. Unlike patients with ADEM, untreated patients with PML have an extremely poor prognosis, with death common in the first 6 months following establishment of the diagnosis. Although there is no specific treatment, combination antiretroviral therapy (cART) has not only resulted in a lower incidence of PML in AIDS patients but also substantially improved survival times of afflicted patients, now at 50 % 1-year survival [25]. Unfortunately, about 20 % of PML cases are linked to a robust inflammatory response to

pathogens associated with recovery of the immune system after a period of immunosuppression. This condition, known as PML–IRIS (immune reconstitution inflammatory syndrome), has been associated with intracranial masses with generous amounts of surrounding vasogenic edema on MRI. Enhancement may also occur [41].

Like ADEM, the lesions of PML are characterized by little mass effect or enhancement. Most lesions involve the subcortical white matter and deep cortical layers of the parieto-occipital or frontal white matter, although gray matter and posterior fossa lesions are also common, occurring in up to 50 % of cases. PML lesions tend to be more confluent in their appearance than ADEM lesions, and scalloping of the lateral margin of the lesion at the gray matter–white matter junction is common. Subtle signal intensity changes in the white matter may precede clinical suspicion of the PML–IRIS. While development of mass effect and temporary enhancement in the early phase of cART has been linked to better survival, similar imaging manifestations may also be seen in monoclonal antibody-treated PML patients [25, 26].

Human Immunodeficiency Virus Encephalopathy

Human immunodeficiency virus (HIV) encephalopathy results from direct infection of the brain by the virus itself. Since the advent of cART, the prevalence of the disease has markedly decreased, and the temporal progression has been slowed. Most patients are severely immunocompromised at the time of onset and exhibit psychomotor slowing, impaired mental status, and memory difficulties. Histologically, demyelination and vacuolation with axonal loss are noted, along with occasional microglial nodules. Mild cerebral atrophy is the first and sometimes only imaging feature of the disease, which is also known as AIDS dementia complex, HIV dementia, HIV-associated dementia complex, and HIV-associated neurocognitive disorder (HAND). Involvement of the central white matter, basal ganglia, and thalamus is characteristic. Typically, bilaterally symmetric abnormal T2 hyperintensity in the basal ganglia and small focal areas in the periventricular regions are noted on T2-weighted MR images [27]. Regression of these findings has been seen following institution of cART.

White Matter Disease from Toxic Imbalance

Chronic Alcohol Ingestion and Its Consequences

Brain abnormalities in alcoholics include atrophy, Marchiafava-Bignami disease, Wernicke encephalopathy,

osmotic myelinolysis, and consequences of liver cirrhosis such as hepatic encephalopathy and coagulopathy [28]. All the reported entities are not specific of alcohol and can be found in many other toxic or metabolic conditions. Ethanol direct brain toxicity is caused by under-regulation of receptors of N-methyl-D-aspartate and abnormal catabolism of homocysteine, resulting in an increased susceptibility to glutamate excitatory and toxic effects. Moreover, immune response occurs mediated by lipid peroxidation products that bind to neurons resulting in neurotoxicity. Neuroimaging studies show a characteristic distribution of loss of volume: initially there is infratentorial predominance with atrophy of the cerebellar vermis and hemispheres. Frontal and temporal atrophy is subsequently evident, followed by diffuse atrophy of the brain. The possibility of partial reversibility of these alterations is also observed. In pregnancy, ethanol determines inhibition of maturation of Bergmann's fibers of cerebellum, with consequential marked cerebellar atrophy.

Hepatic Encephalopathy

The term *hepatic encephalopathy* (HE) includes a wide spectrum of neuropsychiatric abnormalities occurring in patients with liver dysfunction. Most cases are associated with cirrhosis and portal hypertension or portal-systemic shunts, but the condition can also be seen in patients with acute liver failure and rarely those with portal-systemic bypass and no associated intrinsic hepatocellular disease. Although HE is a clinical condition, several neuroimaging techniques, particularly MRI, may eventually be useful for the diagnosis because they can identify and measure the consequences of CNS increase in substances, which, under normal circumstances, are efficiently metabolized by the liver. Classical MR abnormalities in chronic HE include high signal intensity in the globus pallidum on T1-weighted images, likely a reflection of increased tissue concentrations of manganese and an elevated glutamine/glutamate peak coupled with decreased myoinositol and choline signals on proton MR spectroscopy, representing disturbances in cell volume homeostasis secondary to brain hyperammonemia [29]. Recent data have shown that white matter abnormalities, also related to increased CNS ammonia concentration, can also be detected with several MRI techniques: magnetization transfer ratio measurements show significantly low values in otherwise normal-appearing brain white matter, T2-Flair sequences reveal diffuse and focal high-signal intensity lesions in the hemispheric white matter, and diffusion-weighted imaging (DWI) discloses increased white matter diffusivity. All these MR abnormalities, which return to normal with restoration of liver function, probably reflect the presence of mild diffuse interstitial brain edema, which seems to play an essential role in the pathogenesis of HE.

In acute HE, bilateral symmetric signal intensity abnormalities on T2-weighted images, often with associated restricted diffusion involving the cortical gray matter, are commonly identified. Involvement of the subcortical white matter and the basal ganglia, thalami, and midbrain may also be seen. These abnormalities that can lead to intracranial hypertension and severe brain injury reflect the development of cytotoxic edema secondary to the acute increase of brain hyperammonemia.

Marchiafava-Bignami Disease

Marchiafava-Bignami disease is a rare complication of chronic alcoholism, characterized by demyelination and necrosis of the corpus callosum, with rare involvement of extracallosal regions. Etiology remains unknown but is believed to be caused by toxic agents in red wine of low quality and lack of group B vitamins. Rarely, the disease has also been reported in nonalcoholic patients. Symptoms are mainly represented by cognitive deficits, psychosis, hypertonia, and interhemispheric disconnection, until coma and death. Typical MRI features in the acute phase are corpus callosum hyperintensity on T2-weighted sequences and FLAIR, without significant mass effect, with peripheral enhancement. Diffusion is restricted due to cytotoxic edema. In chronic forms, necrosis of the genu and splenium can be detected [30].

Wernicke Encephalopathy

Wernicke encephalopathy (WE) is an acute condition first described by the French ophthalmologist Gayet in 1875 and later by the German neurologist Wernicke in 1881, caused by a deficiency of vitamin B1 (thiamine). WE develops frequently but not exclusively in alcoholics. Other potential causes include extended fasting, malabsorption, digitalis poisoning, and massive infusion of glucose without vitamin B1 in weak patients. The autopsy incidence is reported to be 0.8–2 % in random autopsies and 20 % in chronic alcoholics. The classic clinical triad of ocular dysfunctions (nystagmus, conjugate gaze palsy, ophthalmoplegia), ataxia, and confusion is observed in only 30 % of cases. Treatment consists of thiamine infusion and avoids irreversible consequences including Korsakoff dementia or death. Memory impairment and dementia are related to damage of the mammillary bodies, anterior thalamic nuclei, and interruption of the diencephalic–hippocampal circuits. Depletion of thiamine leads to failure of conversion of pyruvate to acetyl CoA and α -ketoglutarate to succinate, altered pentose monophosphate shunt, and the lack of Krebs cycle, with cerebral lactic acidosis, intra- and extracellular edema,

swelling of astrocytes, oligodendrocytes, myelin fibers, and neuronal dendrites. Neuropathological aspects include neuronal degeneration, demyelination, hemorrhagic petechiae, proliferation of capillaries and astrocytes in periaqueductal gray substance, mammillary bodies, thalami, pulvinar, III cranial nerves nuclei, and cerebellum. On MRI, bilateral and symmetric hyperintensities on T2-weighted sequences and FLAIR are evident at the level of the already mentioned structures, mainly mammillary bodies and thalami [31, 32]. DWI shows areas of high signal with apparent diffusion coefficient (ADC) signal reduced due to cytotoxic edema, although the ADC can sometimes be high due to the presence of vasogenic component. Rarely, T1-weighted images show bleeding ecchymotic hemorrhages in the thalami and mammillary bodies, a sign considered clinically unfavorable. In 50 % of cases, contrast enhancement is present in periaqueductal regions. Marked contrast enhancement of mammillary bodies is evident in 80 % of cases, even prior to the development of visible changes in T2-weighted sequences, and is considered highly specific for WE. In chronic forms, change of signal in T2-weighted sequences becomes less prominent due to the diffuse brain atrophy, more pronounced at the level of mesencephalon and mammillary bodies.

Osmotic Demyelination Syndrome

Osmotic demyelination syndrome (ODS) usually occurs in the setting of osmotic changes, typically with the rapid correction of hyponatremia. This causes destruction of the blood–brain barrier with hypertonic fluid accumulation in extracellular space, resulting in a noninflammatory demyelination. The most common damage is in the pontine fibers [33]. ODS is observed in alcoholics with nutritional deficiency. The most common symptoms include paralysis, dysphagia, dysarthria, and pseudobulbar palsy. Death is frequent. Rarely, ODS affects other regions, especially the basal ganglia, thalami, and deep white matter. MRI usually shows an area of high signal on T2-weighted sequences in the central part of the pons, sparing ventrolateral portions and corticospinal tracts. The lesion is moderately hypointense in T1 and may show positive contrast enhancement. If the patient survives, the acute phase can evolve into a cavitated pontine lesion.

White Matter Disease Associated with Radiation Therapy and Chemotherapy

Treatment strategies designed to target cancer cells are commonly associated with deleterious effects to multiple organ systems, including the CNS. As both radiation and

chemotherapy alone can be associated with significant toxicity, the combination of radiation and chemotherapy may be particularly harmful to the CNS. With advanced treatment regimens and prolonged survival, neurologic complications are likely to be observed with increasing frequency.

Neurotoxicity can result from direct toxic effects of the drug or radiation on the cells of the CNS or indirectly through metabolic abnormalities, inflammatory processes, or vascular adverse effects.

Recognition of treatment-related neurologic complications is critically important, because symptoms may be confused with metastatic disease, tumor progression, paraneoplastic disorders, or opportunistic infections, and discontinuation of the offending drug may prevent irreversible CNS injury.

Radiation Injury and Necrosis

It is widely accepted that the white matter of the CNS is prone to radiation-induced injury, compared with gray matter. Radiation encephalopathy has been classically divided into three stages according to its timing after radiotherapy: early, early-delayed, and late-delayed reactions [34]. Within the first several weeks of therapy, patients may experience acute declines with focal neurologic deficits. These effects are possibly related to increased edema, which has been supported by the observation that steroid treatment often results in clinical improvement. Early- delayed adverse effects usually occur within 1–6 months of treatment and are thought to be a result of demyelination. This syndrome is characterized by somnolence, fatigue, and cognitive dysfunction, consistent with dysfunction of the frontal network systems. Late-delayed side effects occur months to years after cessation of treatment and are largely irreversible and progressive. These late changes are commonly associated with progressive cognitive deficits. In more severe cases of late-delayed radiation injury, imaging and histopathological studies may demonstrate leukoencephalopathy and/or focal necrosis.

In all types of white matter radiation-induced damage, CT and mainly MRI may demonstrate variable degrees of white matter signal changes related to an increase in free tissue water in the involved areas. This may result from endothelial damage, causing increased capillary permeability and vasogenic edema, or from demyelination. However, the degree of these white matter changes correlates poorly with the functional deterioration observed in the patients. MRI findings in early reactions occurring during course of treatment are non-specific. MRI may be normal or demonstrate poorly defined multifocal lesions in both hemispheres that usually disappear spontaneously. MRI in early-delayed reactions also may show signal changes involving not only the hemispheric white matter but also the basal ganglia and the cerebral

peduncles, which resolve completely without treatment. These early-delayed changes have been reported in children with acute lymphatic leukemia who have been treated with both whole brain irradiation and chemotherapy. These changes have no correlation with clinical manifestations and have no clear prognostic significance. Late-delayed reactions can be subdivided into diffuse and focal radiation necrosis injury. Diffuse radiation injury is characterized by white matter changes that are “geographic” in nature, i.e., the areas of abnormal signal intensity or attenuation are limited to the regions of the brain that conform to the radiation portal. This can produce striking differences between the involved zones and the spared surrounding white matter. The involved territories are often symmetric and do not enhance on post-contrast studies. While originally reported in children with leukemia, diffuse necrotizing leukoencephalopathy has also been observed following treatment for many other malignancies in both children and adults. The disease may occur following chemotherapy alone, but the incidence of disease is highest when chemotherapy is combined with radiation therapy. Both the histologic findings and imaging features bear resemblance to radiation necrosis. Axonal swelling, demyelination, coagulation necrosis, and gliosis dominate the histologic picture.

Diffuse white matter changes, with hypoattenuation on CT and T1 and T2 prolongation on MRI, are common and often involve an entire hemisphere. Microbleeds can occur as a sign of vasculopathy. Radiation-induced leukoencephalopathy may be associated with progressive brain atrophy, and patients may present with cognitive decline, gait abnormalities, and urinary incontinence. However, the more common mild-to-moderate cognitive impairment is inconsistently associated with radiological findings and frequently occurs in patients with normal appearing. Recently, more sensitive tools to quantify the damage to brain tissue, such as diffusion tensor imaging (DTI), have been applied in patients treated with brain radiotherapy and have shown early and progressive diffuse changes in normal-appearing white matter indicating radiation-induced demyelination and mild structural degradation of axonal fibers, which are undetectable with conventional MRI.

Focal radiation necrosis usually manifests as a ring-like or irregular enhancing mass lesion located in the white matter, which may become hemorrhagic. The MRI features commonly seen in radiation necrosis include a soap bubble-like interior and Swiss cheese-like interior. Compared with lesions with the soap bubble pattern, Swiss cheese lesions are larger, more variable in size, and more diffuse. This pattern can be visualized as a result of diffuse necrosis affecting the white matter and cortex with diffuse enhancement at the margins with intermixed foci of necrosis. The rim of enhancement is often thinner, more uniform, and more aligned to the gray matter–white matter junction than in

malignant tumors. As radiation necrosis progresses, it can lead to severe shrinkage of the white matter and cortex and result in focal brain atrophy with hydrocephalus.

Quite frequently is impossible to distinguish radiation necrosis from recurrent malignant brain tumor, such as glioblastoma multiforme, using conventional MRI. Metabolic imaging (e.g., positron emission tomography) may facilitate differentiating between the two diseases as radiation necrosis is iso- to hypometabolic, while recurrent high-grade tumors are typically hypermetabolic. MR spectroscopy (MRS) may also be useful as radiation necrosis frequently shows a characteristic lactic acid peak and near-normal peaks for N-acetyl-aspartate (NAA) and choline, while recurrent high-grade gliomas typically show elevated choline levels compared to NAA without or with elevated lactic acid levels. Perfusion imaging can identify the areas of increased blood flow associated with tumor recurrence whereas radiation necrosis is not expected to contain any increased blood flow [35].

Chemotherapy-Associated Neurotoxicity

Neurotoxicity has been observed with virtually all categories of chemotherapeutic agents. Neurologic complications may range from acute encephalopathy, headache, seizures, visual loss, cerebellar toxicity, and stroke to chronic side effects, including chronic encephalopathy, cognitive decline, and dementia.

Among the most puzzling aspects of cancer therapy-related toxicity is the occurrence of delayed and progressive neurologic decline, even after cessation of treatment. Anticancer agents affect brain function through both direct and indirect pathways. It is also conceivable that additional variables play important roles, including the timing of treatment, combination of different treatment modalities, patient age, integrity of the blood–brain barrier, and cognitive function prior to treatment initiation.

Imaging studies have provided evidence that structural and functional CNS changes occur in a significant number of patients treated with chemotherapy. Some agents, such as methotrexate or carmustine, are well known to cause a leukoencephalopathy syndrome, especially when administered at a high dose, intrathecally, or in combination with cranial radiotherapy. Nonenhancing, confluent, periventricular white matter lesions, necrosis, ventriculomegaly, and cortical atrophy characterize this syndrome. White matter abnormalities following high-dose chemotherapy have been detected in up to 70 % of treated individuals and usually have a delayed onset of several months [36].

A delayed leukoencephalopathy syndrome with distinct DWI abnormalities on MRI indicative of cytotoxic edema within cerebral white matter has been recently described.

This syndrome appeared to mimic a stroke-like syndrome and has been seen mainly in patients receiving methotrexate, 5-fluorouracil (5-FU), capecitabine, and carmofofur [37].

Mineralizing Microangiopathy

In children with cancer who have been treated with either chemotherapy alone or in combination with radiation therapy, deposits of calcium in and around small penetrating blood vessels of the deep brain lead to local areas of necrosis. This process is termed mineralizing microangiopathy and is the most common neuroradiologic abnormality noted in this group of patients. The disease has a predilection for the basal ganglia, especially the putamen, and, more rarely, the cerebral cortex. On CT and MRI, evidence of cortical atrophy and abnormal attenuation/signal intensity changes within the white matter are commonly noted. Of all chemotherapeutic agents, methotrexate is the one classically associated with mineralizing microangiopathy. Patients younger than 5 years of age who have meningeal leukemia and have received high-dose methotrexate therapy are at greatest risk of developing this complication of therapy [38].

Vascular Causes of White Matter Disease

Reversible Encephalopathy Syndrome

Although not a truly demyelinating condition, reversible encephalopathy syndrome is noteworthy because of its affinity for the posterior cerebral white matter territories and is important in the differential diagnosis of demyelinating disease. Under normal circumstances, cerebral perfusion pressure is maintained at a relatively constant level by autoregulation, a physiologic mechanism that compensates for wide changes in systemic blood pressure. Hypertensive encephalopathy is believed to result from loss of normal autoregulation, with competing regions of vasodilatation and vasoconstriction and endothelial dysfunction. The vessels of the posterior cerebral circulation, lacking less sympathetic innervation compared to those of the anterior circulation, are unable to vasoconstrict in a normal manner and bear the brunt of these vascular changes. Reversible vasogenic edema is the result and is associated with visual field deficits, as well as headaches, somnolence, and an overall impaired mental status. The terms posterior reversible encephalopathy syndrome (PRES) and reversible posterior leukoencephalopathy syndrome (RPLS) have been popularized in the literature to describe this scenario that is most commonly seen in hypertensive states and/or the presence of immunosuppression (particularly cyclosporine A and tacrolimus), chemotherapy, eclampsia, and renal failure. While it commonly

involves the posterior cerebral white matter, other sites may also be affected including unilateral cerebral hemispheric or isolated brainstem involvement in patients following aortic valve surgery [39, 40]. Accordingly, it has been suggested that perhaps the terminology should be changed to simply “reversible encephalopathy” [40].

On MR studies, bilaterally symmetric abnormal T2 hyperintensity, representing vasogenic edema, is most commonly seen in the distribution of the posterior circulation, although other sites including the frontal lobes and corpus callosum may be noted as well. Cortical and subcortical lesions may be better detected on FLAIR sequences [41]. Diffusion-weighted imaging (DWI) may be normal or show restricted water diffusion in regions of infarction that correlate with poorer prognosis [42]. Susceptibility-weighted imaging may show areas of hemorrhage within involved territories. With early treatment and limited involvement of the brain, many of these imaging abnormalities will completely resolve, and most patients recover within 2 weeks [43]. However, when larger areas or regions of infarction are involved, permanent neurologic deficits or even death are possible [42]. Vascular narrowing has been observed on angiographic studies. Perfusion studies reported in the literature indicate normal to increased perfusion in these zones. When biopsies of these regions have been performed, white matter edema is seen histologically.

Aging and Ischemic Demyelinating Disorders

Small focal lesions on T2-weighted images are quite common in the white matter of adult subjects [44]. They are not associated with mass effect, do not enhance, and are typically isointense compared to normal white matter on T1-weighted images. When these lesions have been biopsied, histologic examination reveals a spectrum of findings including gliosis (partial), loss of myelination, and vasculopathy. They tend to be located in the periventricular white matter, centrum semiovale, and optic radiations. In contrast to MS, they do not involve the corpus callosum or the juxtacortical U-fibers, important distinguishing features [3]. Since the lesions are so ubiquitous and appear to be a part of “normal” aging, various terms have been proposed: senescent white matter changes or disease, deep white matter ischemia, leukoaraiosis, etc. In general, the more lesions present, the more likely it is that the patient will have cognitive problems or difficulties with neuropsychologic testing. However, it is not possible to predict a particular patient’s status simply based on the imaging appearance alone.

In adult patients between 30 and 50 years of age, the presence of periventricular and subcortical lesions in a patient with a family history of similarly affected relatives should raise the possibility of cerebral autosomal dominant

arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL). A defect in the notch3 gene on the long arm of chromosome 19 has been identified and apparently evokes an angiopathy affecting small- and medium-sized vessels. Most lesions occur in the frontal and temporal lobes and less commonly in the thalamus, basal ganglia, internal and external capsules, and brainstem [45].

Conclusions

Several neuroradiologic imaging techniques have been extensively applied to the study of white matter disorders. These examinations play an important role in narrowing the differential diagnosis of this heterogeneous group of disorders. Due to the lack of specificity of imaging findings in many of these conditions, the imaging features should be correlated with clinical manifestations and laboratory results to improve their diagnostic value.

References

1. Smith AB, Smirniotopoulos JG (2010) Imaging evaluation of demyelinating processes of the central nervous system. *Postgrad Med J* 86:218–229
2. Cañellas AR, Gols AR, Izquierdo JR et al (2007) Idiopathic inflammatory-demyelinating diseases of the central nervous system. *Neuroradiology* 49:393–409
3. Aliaga ES, Barkhof F (2014) MRI mimics of multiple sclerosis. *Handb Clin Neurol* 122:291–316
4. Rovira A, León A (2008) MR in the diagnosis and monitoring of multiple sclerosis: an overview. *Eur J Radiol* 67:409–414
5. Filippi M, Rocca MA (2011) MR imaging of multiple sclerosis. *Radiology* 259:659–681
6. Rovira À, Wattjes MP, Tintoré M et al (2015) Evidence-based guidelines: MAGNIMS consensus guidelines on the use of MRI in multiple sclerosis-clinical implementation in the diagnostic process. *Nat Rev Neurol* 11:471–482
7. Bagnato F, Jeffries N, Richert ND et al (2003) Evolution of T1 black holes in patients with multiple sclerosis imaged monthly for 4 years. *Brain* 126:1782–1789
8. Lycklama G, Thompson A, Filippi M et al (2003) Spinal-cord MRI in multiple sclerosis. *Lancet Neurol* 2:555–562
9. Cotton F, Weiner HL, Jolesz FA et al (2003) MRI contrast uptake in new lesions in relapsing–remitting MS followed at weekly intervals. *Neurology* 60:640–646
10. Masdeu JC, Quinto C, Olivera C et al (2000) Open-ring imaging sign: highly specific for atypical brain demyelination. *Neurology* 54:1427–1433
11. Popescu BF, Lucchinetti CF (2012) Pathology of demyelinating diseases. *Annu Rev Pathol* 7:185–217
12. Mehler MF, Rabinowich L (1989) Inflammatory myelinoclastic diffuse sclerosis (Schilder’s disease): neuroradiologic findings. *AJNR Am J Neuroradiol* 10:176–180
13. Hardy TA, Miller DH (2014) Baló’s concentric sclerosis. *Lancet Neurol* 13:740–746
14. Stadelmann C, Ludwin S, Tabira T et al (2005) Tissue preconditioning may explain concentric lesions in Baló’s type of multiple sclerosis. *Brain* 128:979–987
15. Trebst C, Jarius S, Berthele A et al (2014) Update on the diagnosis and treatment of neuromyelitis optica: recommendations of the Neuromyelitis Optica Study Group (NEMOS). *J Neurol* 261:1–16

16. Lennon VA, Wingerchuk DM, Kryzer TJ et al (2004) A serum auto-antibody marker of neuromyelitis optica: distinction from multiple sclerosis. *Lancet* 364:2106–2112
17. Wingerchuk DM, Lennon VA, Pittock SJ et al (2006) Revised diagnostic criteria for neuromyelitis optica. *Neurology* 66:1485–1489
18. Wingerchuk DM, Banwell B, Bennett JL et al (2015) International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* 85:177–189
19. Yonezu T, Ito S, Mori M et al (2014) “Bright spotty lesions” on spinal magnetic resonance imaging differentiate neuromyelitis optica from multiple sclerosis. *Mult Scler* 20:331–337
20. Kim HJ, Paul F, Lana-Peixoto MA et al (2015) MRI characteristics of neuromyelitis optica spectrum disorder: an international update. *Neurology* 84:1165–1173
21. Mitsu T, Fujihara K, Nakashima I et al (2005) Intractable hiccup and nausea with periaqueductal lesions in neuromyelitis optica. *Neurology* 65:1479–1482
22. Krupp LB, Tardieu M, Amato MP et al (2013) International Pediatric Multiple Sclerosis Study Group criteria for pediatric multiple sclerosis and immune-mediated central nervous system demyelinating disorders: revisions to the 2007 definitions. *Mult Scler* 19:1261–1267
23. Verhey LH, Branson HM, Shroff MM et al (2011) MRI parameters for prediction of multiple sclerosis diagnosis in children with acute CNS demyelination: a prospective national cohort study. *Lancet Neurol* 10:1065–1073
24. Tenenbaum S, Chamoles N, Fejerman N (2002) Acute disseminated encephalomyelitis: a long-term follow-up study of 84 pediatric patients. *Neurology* 59:1224–1231
25. Tan CS, Koralknik IJ (2010) Progressive multifocal leukoencephalopathy and other disorders caused by JC virus: clinical features and pathogenesis. *Lancet Neurol* 9:425–437
26. Wattjes MP, Barkhof F (2014) Diagnosis of natalizumab-associated progressive multifocal leukoencephalopathy using MRI. *Curr Opin Neurol* 27:260–270
27. Gottumukkala RV, Romero JM, Riascos RF et al (2014) Imaging of the brain in patients with human immunodeficiency virus infection. *Top Magn Reson Imaging* 23:275–291
28. Geibprasert S, Gallucci M, Krings T (2010) Alcohol-induced changes in the brain as assessed by MRI and CT. *Eur Radiol* 20:1492–1501
29. Rovira A, Alonso J, Cordoba J (2008) MR imaging findings in hepatic encephalopathy. *AJNR Am J Neuroradiol* 29:1612–1621
30. Arbelaez A, Pajon A, Castillo M (2003) Acute Marchiafava-Bignami disease: MR findings in two patients. *AJNR Am J Neuroradiol* 24:1955–1957
31. Gallucci M, Bozzao A, Splendiani A et al (1990) Wernicke encephalopathy: MR findings in five patients. *AJR Am J Roentgenol* 155:1309–1314
32. Zuccoli G, Santa Cruz D, Bertolini M et al (2009) MR imaging findings in 56 patients with Wernicke encephalopathy: nonalcoholics may differ from alcoholics. *AJNR Am J Neuroradiol* 30:171–176
33. Ruzek KA, Campeau NG, Miller GM (2004) Early diagnosis of central pontine myelinolysis with diffusion-weighted imaging. *AJNR Am J Neuroradiol* 25:210–213
34. Soussain C, Ricard D, Fike JR et al (2009) CNS complications of radiotherapy and chemotherapy. *Lancet* 374:1639–1651
35. Cha S (2006) Update on brain tumor imaging: from anatomy to physiology. *AJNR Am J Neuroradiol* 27:475–487
36. Rimkus Cde M, Andrade CS, Leite Cda C et al (2014) Toxic leukoencephalopathies, including drug, medication, environmental, and radiation-induced encephalopathic syndromes. *Semin Ultrasound CT MR* 35:97–117
37. Baehring JM, Fulbright RK (2008) Delayed leukoencephalopathy with stroke-like presentation in chemotherapy recipients. *J Neurol Neurosurg Psychiatry* 79:535–539
38. Davis P, Hoffman JJ, Pearl G et al (1986) CT evaluation of effects of cranial radiation therapy in children. *AJR Am J Roentgenol* 147:587–592
39. Wijdicks EF, Campeau N, Sundt T (2008) Reversible unilateral brain edema presenting with major neurologic deficit after valve repair. *Ann Thorac Surg* 86:634–637
40. McKinney AM, Short J, Truwit CL et al (2007) Posterior reversible encephalopathy syndrome: incidence of atypical regions of involvement and imaging findings. *AJR Am J Roentgenol* 189:904–912
41. Casey SO, Sampaio RC, Michel E, Truwit CL (2000) Posterior reversible encephalopathy syndrome: utility of fluid-attenuated inversion recovery MR imaging in the detection of cortical and subcortical lesions. *AJNR Am J Neuroradiol* 21:1199–1206
42. Covarrubias D, Luetmer P, Campeau N (2002) Posterior reversible encephalopathy syndrome: prognostic utility of quantitative diffusion-weighted MR images. *AJNR Am J Neuroradiol* 23:1038–1048
43. Post JD, Beauchamp NJ (1998) Reversible intracerebral pathologic entities mediated by vascular autoregulatory dysfunction. *Radiographics* 18:353–367
44. Kloppenborg RP, Nederkoorn PJ, Geerlings MI, van den Berg E (2014) Presence and progression of white matter hyperintensities and cognition: a meta-analysis. *Neurology* 82:2127–2138
45. van dem Boom R, Lesnick Oberstein S et al (2006) Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy: MR imaging changes and apolipoprotein E genotype. *AJNR Am J Neuroradiol* 27:359–362