

# Chapter 14

## Role of Biochar in Remediating Heavy Metals in Soil

Amina Hayyat, Maryam Javed, Iqra Rasheed, Shafaqat Ali,  
Munazzam Jawad Shahid, Muhammad Rizwan, Muhammad Tariq Javed,  
and Qasim Ali

### 14.1 Introduction

Soil contamination in these days has become a worldwide problem of greatest magnitude. Despite comprehensive regulations, a huge increase in concentration of different elements and compounds from threshold level is observed in recent years [1]. The main reasons of contamination are natural processes like volcanic eruption and weathering of rocks, as well as anthropogenic activities such as smelting, mining, and overapplication of agrochemicals such as fertilizers and pesticide [1, 2]. Industrial and technological advancement increased pollutant intake into the environment. Heavy metals among these pollutants have great influence on fertility of agricultural soils [3]. Urbanization leads to promote the farmer's concern to use contaminated soil for the production of food crops [4]. Urban agricultural soils are contaminated because of waste water irrigation to increase the crop yield [5]. The wastewater used for irrigation is rich in toxic heavy metals which are major contributor to heavy metal pollution in the soils irrigated are amended with waste water and material [6, 7]. Heavy metals are toxic and stored in the environment because they cannot be broken down [8, 9]. Due to its persistent nature, heavy metals become part of the food chain and pose risks to human health and degrade soil quality. Generally soil chemistry and the chemical form of heavy metal are responsible for the existence of these metals in crops and plants. In soil, commonly the accessibility of all the metals surges due to acidic environments; the transfer factor of a particular

---

A. Hayyat • M. Javed • I. Rasheed • S. Ali (✉) • M.J. Shahid • M. Rizwan  
Department of Environmental Sciences and Engineering, Government College University,  
Allama Iqbal Road, 38000 Faisalabad, Pakistan  
e-mail: [shafaqataligill@yahoo.com](mailto:shafaqataligill@yahoo.com)

M.T. Javed • Q. Ali  
Department of Botany, Government College University,  
Allama Iqbal Road, 38000 Faisalabad, Pakistan

element to the plants is also affected by the crop's type, such as Cd which is mainly considered a toxic element in soil and fertilizers because of its highest transfer factor for any crop [10]. The transfer of heavy metals from soil to plant is one of the basic mechanisms of human exposure to heavy metals through the food chain [11].

Here the overview of heavy metal pollution due to Pb, Cd, Cr, As Ni, Zn, and Cu is described. The concentration of these heavy metals above threshold level is lethal to human health. Some elements such as Cd and Pb, without microbial degradation above optimum level of concentration, put the plant growth at risk due to permanent adulteration in soil [12]. The continuous use of inorganic and biological fertilizers increased heavy metal pollution in soil [1]. It is very necessary to minimize the accessibility and phyto-availability of heavy metals to plants along with the restoration of contaminated soil for safe and healthy food production [13]. In situ remediation techniques have been emphasized in various studies about remediation of heavy metals. In recent years researchers much investigated the use of biochar as in situ soil amendment, and it was found to be effective in reducing the mobility of heavy metals in soils [9, 14]. Biochar method is known as Terra Preta de Indio, introduced from the dark soil of the Amazon basin and is of high utility. Biochar have high chemical stability in contaminated soils and is a C-rich material. A lot of researches are initiated to explore the distinctive use of biochar for continuing C sequestration [14]. Biochar which is porous and has high C content is prepared by pyrolysis of organic waste [14]. Carter et al. [15] defined biochar as follows: "it is a porous carbonaceous solid material manufactured by the process of thermo-chemical decomposition under little supply of oxygen appropriate for the benign and continuing storage of carbon." The International Biochar Initiative (IBI) described biochar as a charcoal which is used as a tool for agricultural and environmental management [9].

Biochar has proved to be a very effective tool for treatment of contaminated soils due to these reasons: it effectively adsorbs heavy metals and decreases bioavailability and toxin-induced stress to plants and microorganisms [13, 16]. Biochar compounds are a good source of organic material and mineral nutrients for microbes. It promotes the beneficial microbes that promote remediation and protect them from predators [17]. Biochar improves the soil fertility and plant growth by improving physical and chemical properties of soil and also increases the availability of useful nutrients [14]. In soils the use of biochar has proved to raise the stable C pool and minimize the increasing concentration of atmospheric CO<sub>2</sub> [18].

## 14.2 Biochar Production and Properties

### 14.2.1 Biomass Pyrolysis

Biochar is a fine-grained porous and carbonaceous solid material synthesized from waste biomass residues under limited oxygen condition and low to medium temperatures (450–650 °C) by the slow pyrolysis [19, 20]. Biochar is manufactured from renewable resources such as green waste and chicken manure [21]. With recent

advancements, biochar can be produced by thermal decomposition of various kinds of organic feedstocks such as crop biomass, wood, agricultural residues (cereal straw, hazelnut and peanut shell, wheat straw, etc.), and industrial organic waste (sewage sludge and de-inking paper sludge [1]. Forest-remain biochar has low metal content, so the frequent use of this biochar has no negative effect on the agricultural soils [22].

### ***14.2.2 Properties of Biochar***

Biochar, a carbon-rich material, is now well known because of its agronomic benefits and ability to moderate climate change by carbon sequestration potential [23–25]. The biochar proved to have an effective role as a soil conditioner and fertilizer [26]. The properties of biochar include highly porous structure, high surface area, pH, cation exchange capacity (CEC), adsorptive capacity, carbon content, organic matter content, and high water-holding capacity. It reduces CO<sub>2</sub> emission, retains nutrients and, pesticide [3, 12, 26, 27], making it a perfect soil amendment to remediate heavy metals and to recover the fertility. Activation of biochar is very effective to improve the adsorption capacity of biochar. The nutrient retention and uptake by plants are enhanced due to activation of biochar as compared to non-activated biochar [28]. Therefore, steam activation is an exciting opportunity for prospective biochar applications because it revealed to almost double the constructive effects of biochars in all illustrations.

### ***14.2.3 Factors Affecting Biomass Properties***

The properties of biochar are dependent upon the type of feedstock and the production procedure. Depending upon these two main parameters, the composition of organic and inorganic contaminants in the biochar also changed, and application of this biochar may cause adulteration in the soil [22]. The source material of biochar may affect the carbon sequestration and conditioning capacity of soil [20]. Other factors such as the type of soil, the type of metal, the nature of biomass, the thermal decomposition conditions (pyrolysis), and the quantity of biochar used [13] may also have prominent effect on properties of biochars.

## **14.3 Heavy Metal-Contaminated Soils**

Mostly the heavy metals such as Cu(II), Cd(II), and Ni(II) are found together in contaminated soils. Oxidation-reduction and acid-base properties of heavy metal ions affect the mobility of these heavy metals. Zn(II) and Pb(II) retain in soil, while Cr(II), Cu(II), and Cd(II) move through soil pore water. Cd(II) and Cu(II) form a

complex with available natural organic material in the soil [26]. Mobility and bioavailability of Cu and Pb are a worldwide matter of concern especially in polluted soils of mining, shooting, and industrialized areas [29]. The frequent use of sewage, municipal composts, manure, mining wastes, and copper-containing fungicides is the main cause of Cu contamination in soil [30]. Higher concentration of Cu has detrimental effect on soil and reduces the population of bacteria, fungi, earthworms, and plant organic content in soil. It also disrupts the nutrient cycle and activity of enzymes such as arylsulfatase, phosphatase, dehydrogenase, and  $\beta$ -glucosidase [16]. Higher concentrations of Cu in fruits and vegetables cause gastrointestinal cancer [31]. In areas rich in Pb and Zn rocks, Cd is obtained from lithogenic sources. Recently, the increased level of Cd in contaminated soils is due to the frequent application of fertilizers and sewage sludge on agricultural soils.

However, plants have high tolerance level for Cd as compared to animals and humans due to this reason: Cd is only toxic to plants at significantly higher concentrations. Disease itai-itai caused by high ingestion of Cd may also cause cancer and damage the kidney [2]. Municipal waste incineration; coal combustion; Pb, Cu, or Zn smelter; electroplating; nickel-cadmium batteries; and pigment production are the major anthropogenic sources of Cd in the environment (World Health Organization, <http://www.euro.who.int/en/home>). Consequently, Cd content is increased due to the usage of phosphatic fertilizers and sewage sludge [32]. Smelting processes are the main cause of Zn production. Among other sources, cosmetics, galvanized products, television, coating of metals, rubber and tire industries, and Zn alloys are prominent. However, as compared to Cd, Zn is less toxic. The presence of Zn in soil is affected by pH, organic content, and structure and nature of parent material. Zn is a transition metal and is an essential micronutrient for many biological processes, but it is toxic at higher concentrations; Zn is the most abundant trace heavy metal existing in agroecosystems [33]. Zn is also entering in the environment from sources such as municipal waste treatment plants and burning of coal and waste.

Municipal waste treatment plants and burning of coal and waste are also sources of zinc. The bioavailability of zinc in alkaline soil increases due to its solubility. Zn(II) also has the mobility in acidic soils [34]. The adverse effects of Zn on plant physiology are widely reported [35]. The solubility and bioavailability of Pb increase in soil due to weather and oxidation processes which modify the metallic Pb in soil. After its bioavailability, Pb promptly adsorbed on the sediment and soil particles, and it also tends to accumulate in plants and animal bodies [29]. Arsenic (As)-contaminated soils deteriorate the water quality because it leaches down from the soil and contaminated water sources such as the rivers and canals. The process of oxidative phosphorylation and synthesis of ATP in cells is also disturbed from high concentration of arsenic [3].

Arsenic also has detrimental effect on the activity of microbial population, soil biota, and nutrient cycles [36]. Ni originated in the environment both from the natural and anthropogenic process. The weathering of rocks and human activities such as smelting, plating, and mining are the main sources of Ni contamination of soil. For agricultural soils, application of organic waste material such as sewage sludge and fertilizer application are the main causes of contamination [1]. Chromium exists

in the contaminated soil as Cr(III) and Cr(VI) ions. Chromium oxidation states are responsible for toxicity toward plants and animals, such as Cr(III) which is an essential nutrient and has less solubility in acidic and alkaline soils, whereas Cr(VI) is highly soluble in acidic and alkaline soils considered as carcinogen. Cr(VI) has a harmful effect and disturbs the biological activity of the soil. The soil properties play the main role in the availability of heavy metals to plants for uptake [10, 37]. In soil, the various enzymatic activities of bacteria are disrupted due to modification of soil environment by chromium (Cr).

### ***14.3.1 Heavy Metal Remediation by Biochar***

Namgay et al. [38] documented a decrease in the accessibility of heavy metals after the contaminated soil was amended with biochar, due to which plant absorption of the heavy metals is reduced. Unlike many other biological amendments, biochar having the ability to increase soil pH [39] might have improved sorption of these metals, consequently decreasing their bioavailability for plant uptake.

### ***14.3.2 Heavy Metals Found in Soil***

Generally Cu, Zn, As, Cr, Co, Ni, Sb, Hg, Th, Pb, Se, Si, and Cd are heavy metals that originate in soil which may be extremely harmful to human and plant life by contamination of soil and water. Heavy metals do not have the ability to biodegrade, so they can persist in polluted soils for a longer time [40]. There is a prerequisite to remove these metals, and the best convenient way is by environmental friendly techniques, i.e., biochar. From a long period, biochar is being applied to overcome the problem of heavy metal contamination and to improve soil fertility. Depending on the soil type, diverse types of biochar are used for different types of heavy metals, as demonstrated in Table 14.1.


Biochar has high pH and organic carbon content; higher concentrations of phosphorous, calcium, and magnesium; and low particular surface area than activated carbon [41]. That's why the addition of biochar brought a notable proliferation in soil cation exchange capability [42]. Biochar is considerably more active in restraining soil Pb than AC [41]. Biochar has excellent adsorption capacity due to its asymmetrical plates and porous structure (Fig. 14.1) [42].

The biochar produced from different sources showed similar results [43]. The maximum falloff of transferable Pb was achieved at 10% application rate of biochars with steady reduction of 68 and 30% for sugar cane biogases and orange peel, respectively [3]. Biochar is prosperous in nutrients, i.e., nitrogen, calcium, magnesium, and phosphorous in addition to carbon. Concentration of carbon and nitrogen are reduced with the increase in temperature, although Mg, Ca, and P were augmented by rise in temperature [43]. 93% of lead was absorbed by biochar at 100,

**Table 14.1** Heavy metal removal by different types of biochar

Contaminants	Biochar type	Matrix	Effects	References
As and Cu	Hardwood	Soil	Mobilization due to enhanced pH and DOC	Beesley et al. [67]
As, Cr, Cd, Cu, Ni, Pb, and Zn	Sewage sludge (500–550 °C)	Soil	Immobilization of As, Cr, Ni, and Pb due to rise in soil pH. Mobilization of Cu, Zn, and Cd due to highly available concentrations in biochar	Khan et al. [68]
Cd and Zn	Hardwood	Soil	Immobilization due to enhanced pH	Beesley et al. [67]
Cd, Cu, and Pb	Chicken manure and green waste (550 °C)	Soil	Immobilization due to partitioning of metals from exchangeable phase to less bioavailable organic-bond fraction	Park et al. [20]
Cu	Broiler litter (700 °C)	Soil	Cation exchange; electrostatic interaction; sorption on mineral ash content; complexation by surface functional groups	Uchimiya et al. [69]
Cu and Pb	Oakwood	Soil	Complexation with phosphorus and organic matter	Karami et al. [70]
Pb	Dairy manure (450 °C)	Soil	Immobilization by hydroxypyromorphite formation	Cao et al. [41]
Pb	Oakwood (400 °C)	Soil	Immobilization by rise in soil pH and adsorption on biochar	Ahmad et al. [29]
Pb	Rice straw	Soil	Nonelectrostatic adsorption	Jiang et al. [42]
Pb, Cu and Zn	Broiler litter (300 and 600 °C)	Soil	Stabilization of Pd and Cu	Uchimiya et al. [48]
Ni, Cu, Pb, and Cd	Cottonseed hulls (200–800 °C)	Soil	Surface functional groups of biochar-controlled metal sequestration	Uchimiya et al. [71]

while at 200 and 350, approximately the entire lead was removed from the soil solution [43]. The pH is the main parameter for disturbing adsorption and/or desorption of heavy metals in acidic soils. The amplified system pH by application of biochar increases the adsorption of Pb(II) in the soils. Biochar results in pH increase that flourishes the negative surface charges in the soil and enhanced the attraction for cations [44]. pH rise is beneficial for heavy metal control in bulk soils. The increasing amount of biochar reduced the acid soluble Pb(II) and Cu(II) by 18.8–77.0% and 19.7–100.0%, respectively [42]. The uses of biochar, mussel shell, and cow bone reduced the lead phyto-availability by 55.50%, 71.22%, and 70.47%, respectively, in army firing soil [45].



**Fig. 14.1** SEM photographs of rice straw biochar at different magnifications: (a)  $\times 800$ , (b)  $\times 50,000$ , and (c) TEM photograph of the biochar

When the application rate of biochar produced from cow manure increased, the nutrient uptake, maize production, and photochemical properties of a dry land sandy soil considerably improved [46]. Sewage sludge-derived biochar successfully eradicates  $Pb^{2+}$  from acidic soil at early pH 5, 4, 3, and 2 with the capabilities of 30.88, 24.80, 20.11, and 16.11  $mg\ g^{-1}$ , respectively [47]. Biochar with phosphorus-rich manure serves as a fertilizer and is also used to remove heavy metal, specifically lead [48]. Historically, phosphorus in the biochar encouraged modification of less constant  $PbCO_3$  to more stable  $Pb_5(PO_4)_3OH$ , liable for soil Pb restriction [41]. The application of biochar produced from cotton sticks put a positive impact on the cadmium-stressed soils by increasing the plant growth. It is due to the unique capacity of biochar that the metal ion is separated and the cadmium ion movement to the aerial tissue of plants is controlled [49].

The biochar synthesized from swine manure at  $450\ ^\circ C$  could contribute as a possible amendment for the control of heavy metals ( $Cd^{2+}$ ) in sandy soil [9]. Biochar produced from green waste restrained lead, copper, and cadmium by 36.8%, 22.9%,

and 30.3 %, respectively, for pointed soil and by 72.9, 0.901, and 42.7 % for naturally polluted soils [20]. Bamboo-derived biochar can adsorb nickel, chromium, copper, and mercury, from both water and soils, and cadmium from polluted soils [40].

Table 14.2 shows the effect of biochar application on the mobility of heavy metals in soils.

In basic soils, carbonaceous tools, irrespective of biomass and pyrolysis, improved the Cu(II) restriction than Ni(II) [26] while the Cu(II) adsorption amplified with pH rise (3.5–6.0). Cu(II) has an adsorption capability in the following order: canola straw char < soybean straw char < peanut straw char [50]. Organic segments of biochars and natural organic matter can stimulate Cu(II) extraction by basic soil because of more carboxyl contents [26]. The unsaturated biochar detached about 70 % of Cr(III) at equilibrium time, although only 30 % of As(V) in batch kinetic trials, suggesting that biochar is highly effective in eliminating cations than anions [51]. Biochar derived from chicken manure is more active in controlling metals as well as plant growth than biochar derived from green waste. So, biochar derived from chicken manure can be used to improve phyto-stabilization of metal tainted soils [20].

Jiang et al. [42] explored that biochar derived from rice straw influences the bio-availability and mobility of Pb(II), Cu(II), and Cd(II) in an Ultisol. When the amendment dosage of biochar increased, the acid removable Pb(II) and Cu(II) reduced by 18.8e77.0 % and 19.7e100 %, respectively. With the addition of 5 mmol kg<sup>-1</sup> of these heavy metals, for treatments with 3 and 5 % biochar, the reducible Pb(II) was 2.0 and 3.0 times greater than the samples deprived of biochar. Pore structure of the biochars produced from crop straws (i.e., rice, corn, wheat, and cotton) is more developed as compared to wood char because wood char has greater lignin content. Biochars with

**Table 14.2** Effect of biochar application on the mobility of heavy metals in soils

Feedstock	Production temperature (°C)	Contaminants	Effects	References
Bamboo	Not available	Cd	Combined effect of electrokinetics, removal of extractable Cd by 76.9 % within 12 days	Ma et al. [72]
Hardwood	450	As, Cu, Cd, and Zn	Reduction in Cd in soil pore water by tenfolds; Zn concentrations reduced 300- and 45-folds, respectively, in column leaching test	Beesley et al. [67]; Beesley and Marmiroli [73]
Hardwood	450	As, Cd, Cu, Pb, and Zn	Biochar surface mulch enhanced As and Cu mobility in the soil profile; little effect on Pb and Cd	Beesley and Dickinson [74]
Wood	200 and 400	Cd and Zn	Reduction in Cd and Zn leaching loss by >90 %	Debela et al. [75]



lower lignin content have larger surface areas and more developed pore volumes. In case of Cd the order of corn straw > cotton straw > wheat straw > rice straw > poplar shaving is for the sorption capacity of biochar that was not stringently reliable to the surface area of biochars [52]. Uchimiya et al. [53] reported that a biochar adsorbs Cd, Cu, Pb, and Ni and perceived that the tendency of the elimination order was Ni < Cd < Cu < Pb. The affinity for metal immobilization upsurges in the following order: Pb(II) > Cu(II) > Zn(II) > Cd(II) [26].

Table 14.3 shows the effect of biochar application on the bioavailability of heavy metals in soils.

**Table 14.3** Effect of biochar application on the bioavailability of heavy metals in soils

Feedstock	Production temperature (°C)	Contaminant	Effects	References
Cotton stacks	450	Cd	Reduction of the bioavailability of Cd in soil by adsorption or Co precipitation	Zhou et al. [76]
Hardwood-derived biochar	450	As	Significant reduction of As in the foliage of <i>Miscanthus</i>	Hartley et al. [77]
Eucalyptus	550	As, Cd, Cu, Pb, and Zn	Decrease in As, Cd, Cu, and Pb in maize shoots	Namgay et al. [38]
Orchard prune residue	500	Cd, Cr, Cu, Ni, Pb, and Zn	Significant reduction of the bioavailable Cd, Pb, and Zn with Cd showing the greatest reduction; an increase in the pH, CEC, and water-holding capacity	Fellet et al. [78]
Chicken manure and green waste	550	Cd, Cu, and Pb	Significant reduction of Cd, Cu, and Pb accumulation by Indian mustard	Park et al. [20]
Chicken manure	550	Cr	Enhanced soil Cr(VI) reduction to Cr(III)	Choppala et al. [79]
Sewage sludge	500	Cu, Ni, Zn, Cd, Pb	Significant reduction in plant availability of the metals studied	Méndez et al. [80]
Rice straw	Not clear	Cd, Cu, and Pb	Significant reduction in concentrations of free Cu, Pb, and Cd in contaminated soils Identification of functional groups on biochar with high adsorption affinity to Cu	Jiang et al. [42]
Quail litter	500	Cd	Reduction of the concentration of Cd in physic nut; greater reduction with the higher application rates	Suppadit et al. [81]
Oakwood	400	Pb	Bioavailability reduction by 75.8%; bioaccessibility reduction by 12.5%	Ahmad et al. [29]

## **14.4 Effects of Biochar on Soil**

The increasing population of human and their activities put huge pressure on agriculture land to fulfill need of food. This overburden on cultivated land raised the problems of soil erosion and degradation and depletion of organic matter and vital nutrients from the soil. To overcome these problems, biochar can be used as an effective tool [54]. Biochar can be utilized to get agricultural, environmental, and economic benefits although the nature of these benefits can vary by type of biochar and nature of soil [55].

### ***14.4.1 Agricultural Benefits***

Application of biochar in soil results in the following positive impacts on the soil of agriculture lands.

#### **14.4.1.1 Soil Fertility**

The extreme use of fertilizers exacerbates the leaching of macronutrients from the agriculture lands leading to decrease in soil fertility, amplification of acidity, and increased requirements of fertilizers that resulted in ultimate low crop yield [56]. Biochar fulfills the nutrient requirement of the soil and improves the soil fertility and productivity that resulted in optimum yield of crops [57].

#### **14.4.1.2 Crop Productivity**

Biochar offers large surface area for soil microbes resulting in increase in microbial growth. It improves the degradation of organic matter and improves retention time of microbes in soil and increases availability of nutrients which are favorable for soil growth for better crop production [27, 49, 54]. The productivity of soil also increases due to the addition of biochar in soil because it increases the availability of C compound and minimizes the requirement of artificial fertilizers [58].

#### **14.4.1.3 Water Holding Capacity**

Biochar enhances the water-holding capacity of soil by improving the soil quality in terms of physical, biological, and chemical fertility and better plant root density [59].

#### **14.4.1.4 Cation Exchange Capacity**

The application of biochar causes the stabilization of heavy metal, increases CEC, and increases the availability of higher mineral essential for plant growth [48, 60].

#### **14.4.1.5 pH of soil**

The specific pH properties of biochar make the soil acidic for evaporation of metal ion in soil. The soils with low pH and CEC will also have the low adsorption capacity for metals [61].

#### **14.4.1.6 Uptake of Heavy Metals**

The addition of biochar reduces the heavy metal uptake by plant roots and minimizes the chance of low productivity due to toxicity in plant body [59].

#### **14.4.1.7 Waste Reduction**

The production of biochar is also an innovative technique to make beneficial reuse of waste and to minimize the waste volume [41].

#### **14.4.1.8 Moisture Holding Capacity**

The moisture holding capacity is improved due to application of biochar due to retention of pollution for shorter or longer time duration [62].

#### **14.4.1.9 Adsorbent Capacity**

Biochar has high adsorbent capacity for adsorption of heavy metals and minerals in soil. Due to its high competency to adsorb organic contaminants for purification of soil from contaminants, it improves the soil fertility [63, 64].

#### **14.4.1.10 Soil Respiration Rate**

The respiration of soil improves by decomposition of biochar with the help of microbes (bacteria) resulting in C production which is used by microbes for microbial activity in soil [1].

#### **14.4.1.11 Residing Capacity of Soil-Living Organisms**

Biochar also improves the resistance capacity of living organisms residing in soil, i.e., earthworms (nematodes), insects, fungus, etc. [58].

#### **14.4.1.12 Nutrient Availability**

High concentration of ash in biochar increases the availability of nutrients which satisfies the need of soil for nutrients. In case of low ash concentration in biochar, compost or manure can be added to maintain the ratio of nutrients [65].

#### **14.4.1.13 Rate of Germination**

The rate of seed germination increased due to direct interaction of biochar with soil [58].

### ***14.4.2 Environmental Benefits***

Interaction of soil with biochar results in subsequent effects on environment:

#### **14.4.2.1 Carbon Sequestration**

Biochar has great resistance for biotic and biotic degradation in soil which emphasizes its importance in carbon cycle as carbon sink [66]. Carbon sequestration resulting in reduction of CO<sub>2</sub> in the atmosphere is due to the long-term availability of biochar in soil. It also has the ability to minimize the effect of climate change, decrease in GHG (greenhouse gas), and NO<sub>x</sub> emissions by causing decline in C emission as output from burning fossil fuels. Methane and nitrous oxide produced from carbon cycle and nitrification/denitrification process, respectively, can be reduced by application of biochar, and thus biochar can play an important role to solve issues of global warming [54, 56, 57, 62].

### ***14.4.3 Economic Benefits***

Application of biochar is also beneficial for the economy.

#### **14.4.3.1 Economically Beneficial**

Biochar is desirable nowadays because of its low energy demand, ease of use, low cost, and no pretreatment for use in soil.

### 14.4.3.2 Environment Friendly

The eco-friendly nature of biochar made it desirable, and nowadays it is being widely used in various applications because it (1) can be reuseable, (2) can recycle the organic waste, and (3) can reduce waste quantity [64].

## 14.5 Conclusion

Global industrialization leads pollutants to the environment. Among these pollutants, heavy metals have the profound effect on fertility of soils. Biochar amendments remediate heavy metal toxicity in agricultural soils through different processes. Biochar is an effective tool for contaminated soil due to these different processes: (1) adsorption of heavy metals; (2) highly porous structure and high surface area; (3) CEC; (4) reduced CO<sub>2</sub> emission; (5) high water-holding capacity; (6) retention of pesticides, PAHs, and PSBs; (7) reduction of bioavailability and toxin-induced stress to microorganisms and plants; (8) C-rich material having high chemical stability in the contaminated soils and mineral nutrients for microbes; (9) reduction of the mobility in soil; (10) protection of microbes from predators and introduction of beneficial microbes that promote remediation; and (11) improvement of soil fertility and plant growth by improving physical and biological properties of soil by provision and maintenance of nutrients.

However, these processes vary with biochar type, nature of soil, type of plants, type of metal toxicity, conditions of thermal decomposition (pyrolysis), and the quantity of biochar used. Therefore, we should use biochar according to soil contamination. However, the various facts of opinions founded on comprehensive point of views should not be snubbed. Variant consequences recommend that recent biochar application to soil is not a standard example, as an alternative extensive concern of the properties related to each specific biochar material and how those properties could cure a particular soil scarcity are mandatory [55]. Biochar use in soil has been suggested to increase the stable C pool and limit the growing concentration of atmospheric CO<sub>2</sub>. In conclusion much more investigations are required to check the long-term environmental and economic feasibility of biochar application to remediate heavy metal-contaminated soils.

## References

1. Mendez A, Ferreiro JP, Gasco G (2014) Biochar from pyrolysis of drinking paper sludge and its use in the treatment of a nickel polluted soil. *J Anal Appl Pyrolysis* 107:46–52
2. Lu H, Li Z, Fu S, Mendez A, Gasco G, Ferreiro JP (2014) Can biochar and phytoextractors be jointly used for cadmium remediation? *PLoS One* 9:1–7

3. Abdelhafez AA, Li J, Abbas MHH (2014) Feasibility of biochar manufactured from organic wastes on the stabilization of heavy metals in a metal smelter contaminated soil. *Chemosphere* 117:66–71
4. Nabulo G (2009) Assessing risks to human health from peri-urban agriculture in Uganda. Ph.D. thesis, University of Nottingham
5. Mahmood A, Malik RN (2014) Human health risk assessment of heavy metals via consumption of contaminated vegetables collected from different irrigation sources in Lahore, Pakistan. *Arab J Chem* 7:91–99
6. Singh KP, Mohon D, Sinha S, Dalwani R (2004) Impact assessment of treated/untreated wastewater toxicants discharge by sewage treatment plants on health, agricultural and environmental quality in wastewater disposal area. *Chemosphere* 55:227–255
7. Mapanda F, Mangwayana EN, Nyamangara J, Giller KE (2005) The effect of long-term irrigation using wastewater on heavy metal contents of soils under vegetables in Harare, Zimbabwe. *Agr Ecosyst Environ* 107:151–165
8. Anwar J, Shafique U, Zaman WU, Salman M, Dar A, Anwar S (2010) Removal of Pb(II) and Cd(II) from water by adsorption on peels of banana. *Bioresour Technol* 101:1752–1755
9. Xu D, Zhao Y, Sun K, Gao B, Wang Z, Jin J, Zhang Z, Wang S, Yan Y LX, Wu F (2014) Cadmium adsorption on plant- and manure-derived biochar and biochar-amended sandy soils: Impact of bulk and surface properties. *Chemosphere* 113:320–326
10. Javied S, Mehmood T, Choodhry MM, Tufail M, Irfan N (2009) Heavy metal pollution from phosphate rock used for the production of fertilizer in Pakistan. *Microchem J* 91:94–99
11. Khan S, Rehman S, Khan AZ, Khan MA, Shah MT (2010) Soil and vegetables enrichment with heavy metals from geological sources in Gilgit, northern Pakistan. *Ecotoxicol Environ Saf* 73:1820–1827
12. Kloss S, Zehetner F, Obergur E, Buecker J, Kitzler B, Wenzel WW, Wimmer B, Soja G (2014) Trace element concentrations in leachates and mustard plant tissue (*Sinapis alba* L.) after biochar application to temperate soils. *Sci Total Environ* 481:498–508
13. Al-Wabel MI, Usman ARA, El-Naggar AH, Aly AA, Ibrahim HM, Elmaghraby S, Al-Omran A (2014) Conocarpus biochar as a soil amendment for reducing heavy metal availability and uptake by maize plants. *Saudi J Biol Sci* 22:503–511
14. Houben D, Evrard L, Sonnet P (2013) Mobility, bioavailability and pH-dependent leaching of cadmium, zinc and lead in a contaminated soil amended with biochar. *Chemosphere* 92:1450–1457
15. Carter S, Shackley S, Sohi S, Suy TB, Haefele S (2013) The impact of biochar application on soil properties and plant growth of pot grown lettuce (*Lactuca sativa*) and cabbage (*Brassica chinensis*). *Agronomy* 3:404–418
16. Mackie KA, Marhan S, Ditterich F, Schmidt HP, Kandeler E (2015) The effects of biochar and compost amendments on copper immobilization and soil microorganisms in a temperate vineyard. *Agric Ecosyst Environ* 201:58–69
17. Rutigliano FA, Romano M, Marzaioli R, Baglivo I, Baronti S, Miglietta F, Castaldi S (2014) Effect of biochar addition on soil microbial community in a wheat crop. *Eur J Soil Biol* 60:9–15
18. Fang Y, Singh B, Singh BP (2015) Effect of temperature on biochar priming effects and its stability in soils. *Soil Biol Biochem* 80:136–145
19. Lehmann J, Joseph S (2009) Biochar for environmental management: an introduction. In: Lehmann J, Joseph S (eds) *Biochar for environmental management-science and technology*. Earthscan, London, pp 1–9
20. Park JH, Choppala GK, Bolan NS, Chung JW, Chuasavathi T (2011) Biochar reduces the bio-availability and phytotoxicity of heavy metals. *Plant Soil* 348:439–451
21. Hass A, Gonzalez JM, Lima IM, Godwin HW, Halvorson JJ, Boyer DG (2011) Chicken manure biochar as liming and nutrient source for acid Appalachian soil. *J Environ Qual* 41:1096–1106

22. Lucchini P, Quilliam RS, DeLuca TH, Vamerali T, Jones DL (2014) Does biochar application alter heavy metal dynamics in agricultural soil? *Agric Ecosyst Environ* 184:149–157
23. Marris E (2006) Putting the carbon back: black is the new green. *Nature* 442:624–626
24. Lehmann J (2007) A handful of carbon. *Nature* 447:143–144
25. Khan S, Reid BJ, Li G, Zhu YG (2014) Application of biochar to soil reduces cancer risk via rice consumption: A case study in Miaoqian village, Longyan, China. *Environ Int* 68:154–161
26. Uchimiya M, Lima IM, Klasson KT, Wartelle LH (2010) Contaminant immobilization and nutrient release by biochar soil amendment: Roles of natural organic matter. *Chemosphere* 80:935–940
27. Fellet G, Marmiroli M, Marchiol L (2014) Elements uptake by metal accumulator species grown on mine tailings amended with three types of biochar. *Sci Total Environ* 468–469:598–608
28. Borchard N, Wolf A, Laabs V, Aeckersberg R, Scherer HW, Moeller A, Amelung W (2012) Physical activation of biochar and its meaning for soil fertility and nutrient leaching—a greenhouse experiment. *Soil Use Manage* 28:177–184
29. Ahmad M, Lee SS, Yang JE, Ro HM, Lee YH, Ok YS (2012) Effects of soil dilution and amendments (mussel shell, cow bone, and biochar) on Pb availability and phytotoxicity in military shooting range soil. *Ecotoxicol Environ Saf* 79:225–231
30. Uchimiya M, Klasson KT, Wartelle LH, Lima IM (2011) Influence of soil properties on heavy metal sequestration by biochar amendment: 2. Copper desorption isotherms. *Chemosphere* 82:1438–1447
31. Turkdogan MK, Fevzi K, Kazim K, Ilyas T, Ismail U (2003) Heavy metals in soil, vegetables and fruits in the endemic upper gastrointestinal cancer region of Turkey. *Environ Toxicol Pharm* 13:175–179
32. Lu H, Li Z, Fu S, Mendez A, Gasco G, Ferriero JP (2015) Combining phytoextraction and biochar addition improves soil biochemical properties in a soil contaminated with Cd. *Chemosphere* 119:209–216
33. Baccio D, Kopriva S, Sebastiani L, Rennenberg H (2005) Does glutathione metabolism have a role in the defence of poplar against zinc excess. *New Phytol* 167:73–80
34. Cameron RE (1992) Guide to site and soil description for hazardous waste site characterization, EPA/600/4-91/029, vol 1, Metals. US Environmental Protection Agency, Washington, DC
35. Manivasagaperumal R, Balamurugan S, Thiagarajan G, Sekar J (2011) Effect of zinc on germination, seedling growth and biochemical content of cluster bean (*Cyamopsis tetragonoloba* (L.) Taub). *Curr Bot* 2:11–15
36. Gregory SJ, Anderson CWN, Arbestain MC, McManus MT (2014) Response of plant and soil microbes to biochar amendment of an arsenic-contaminated soil. *Agric Ecosyst Environ* 191:133–141
37. Wyszowska J (2002) Soil contamination by chromium and its enzymatic activity and yielding. *Pollut Environ Stud* 11:79–84
38. Namgay T, Singh B, Singh BP (2010) Influence of biochar application to soil on the availability of As, Cd, Cu, Pb, and Zn to maize (*Zea mays* L.). *Soil Res* 48:638–647
39. Novak JM, Busscher WJ, Laird DL, Ahmedna M, Watts DW, Niandou MAS (2009) Impact of biochar amendment on fertility of a south eastern coastal plain soil. *Soil Sci* 174:105–112
40. Zhang X, Wang H, He L, Lu K, Sarmah A, Li J, Bolan NS, Pei J, Huang H (2013) Using biochar for remediation of soils contaminated with heavy metals and organic pollutants. *Environ Sci Pollut Res* 20:8472–8483
41. Cao X, Ma L, Liang Y, Gao B, Harris W (2011) Simultaneous immobilization of lead and atrazine in contaminated soils using dairy-manure biochar. *Environ Sci Technol* 45:4884–4889
42. Jianga J, Xua R, Jianga T, Li Z (2012) Immobilization of Cu(II), Pb(II) and Cd(II) by the addition of rice straw derived biochar to a simulated polluted Ultisol. *J Hazard Mater* 229–230:145–150

43. Cao X, Harris W (2010) Properties of dairy-manure-derived biochar pertinent to its potential use in remediation. *Bioresour Technol* 101:5222–5228
44. Jiang T, Jiang J, Xu R, Li Z (2012) Adsorption of Pb (II) on variable charge soils amended with rice-straw derived biochar. *Chemosphere* 89:249–256
45. Ahmad M, Lee SS, Lim JE, Lee S, Cho JS, Moon DH, Hashimoto Y, Ok YS (2014) Speciation and phytoavailability of lead and antimony in a small arms range soil amended with mussel shell, cow bone and biochar: EXAFS spectroscopy and chemical extractions. *Chemosphere* 95:433–441
46. Uzoma KC, Inoue M, Andry H, Fujimaki H, Nishihara E, Zahoor A (2011) Effect of cow manure biochar on maize productivity under sandy soil condition. *Soil Use Manage* 27:205–212
47. Lu H, Zhang W, Yang Y HX, Wang S, Qiu R (2012) Relative distribution of Pb 2D sorption mechanisms by sludge-derived biochar. *Water Res* 46:854–862
48. Uchimiya M, Bannon DI, Wartell LH, Lima IM, Klasson KT (2012) Lead retention by broiler litter biochars in small arms range soil: impact of pyrolysis temperature. *Agric Food Chem* 60:5035–5044
49. Younis U, Shah MHR, Danish S, Malik SA, Ameer A (2014) Biochar role in improving biometric and growth attributes of *S. oleracea* and *T. corniculata* under cadmium stress. *Int J Biosci* 5:84–90
50. Tong X, Li J, Yuan J, Xu R (2011) Adsorption of Cu (II) by biochars generated from three crop straws. *Chem Eng J* 172:828–834
51. Agrafioti E, Bouras G, Kalderis D, Diamadopoulos E (2013) Biochar production by sewage sludge pyrolysis. *J Anal Appl Pyrolysis* 101:72–78
52. Sun J, Lian F, Liu Z, Zhu L, Song Z (2014) Biochars derived from various crop straws: characterization and Cd(II) removal potential. *Ecotoxicol Environ Saf* 106:226–231
53. Uchimiya M, Lima IM, Klasson KT, Chang SC, Wartelle LH, Rodgers JE (2010) Immobilization of heavy metals ions (CuII, CdII, NiII, and PbII) by broiler litter-derived biochar in water and soil. *J Agric Food Chem* 58:5538–5544
54. Fisher D, Glaser B (2012) Synergism between compost & biochar for sustainable soil amelioration. *Intech Open Sci/Open Mind* 32:167–198
55. Novak JM, Busscher WJ (2012) Selection and use of designer biochars to improve characteristics of southeastern USA Coastal Plain degraded soils. In: Lee JE (ed) *Advanced biofuels and bioproducts*. Springer, New York
56. Yao Y, Gao B, Zhang M, Inyang M, Zimmerman AR (2012) Effect of biochar amendment on sorption & leaching of nitrate, ammonium & phosphate in a sandy soil. *Chemosphere* 89:1467–1471
57. Jones DL, Murphy DV, Khalid M, Ahmed W, Jones GE, Deluca TH (2011) Short term biochar-induced increase in soil CO<sub>2</sub> release in both biotically & abiotically mediated. *Soil Biol Biochem* 43:1723–1731
58. Ahmad M, Rajapaksha AU, Lim JE, Zhang M, Bolan N, Mohan D, Vithanage M, Lee SS, Ok YS (2014) Biochar as a sorbent for contaminant management in soil and water: a review. *Chemosphere* 99:19–33
59. Younis U, Qayyum MF, Shahi MHR, Malik SA (2014) Nutrient shift modeling in Fenugreek (*Trigonella corniculata* L.) under biochar and cadmium treatment. *Int J Biosci* 5:64–74
60. Bolan NS, Kunhikrishnan A, Choppala GK, Thangarajan R, Chung JW (2012) Stabilization of carbon in composts and biochars in relation to carbon sequestration and soil fertility. *Sci Total Environ* 424:264–270
61. Jiang J, Xu RK (2013) Evaluating role of alkali and organic functional group in Cu(II) immobilization. *Bioresour Technol* 133:537–545
62. Kolodyriska D, Wnetrzak R, Leahy JJ, Hayes MHB, Kwapisirski W, Hubicki Z (2012) Kinetic and adsorption characterization of biochar in metal ions removal. *Chem Eng J* 197:295–305
63. Xu X, Cao X, Zhao L, Wang H, Yu H, Gao B (2013) Removal of Cu, Zn and Cd from aqueous solutions by the dairy manure-derived biochar. *Environ Sci Pollut Res* 20:358–368



64. Ahmad M, Lee SS, Rajapaksha AU VM, Zhang M, Cho JS, Lee SE, Ok YS (2013) Trichloroethylene adsorption by pine needle biochars produced at various pyrolysis temperatures. *Bioresour Technol* 143:615–622
65. Ska DK, Trzak RW, Leahy JJ, Hayes MHB, Ski WK, Hubicki Z (2012) Kinetic and adsorption characterization of biochar in metal ions removal. *Chem Eng J* 197:295–305
66. Harvey OR, Kuo LJ, Zimmerman AR, Louchouart P, Amonette JE, Herbert BE (2012) An index based approach to assessing recalcitrance and soil carbon sequestration potential of engineered black carbons (biochar). *Environ Sci Technol* 46:1415–1421
67. Beesley L, Jiménez EM, Eyles JLG (2010) Effects of biochar and green waste compost amendments on mobility, bioavailability and toxicity of inorganic and organic contaminants in a multi-element polluted soil. *Environ Pollut* 158:2282–2287
68. Khan S, Chao C, Waqas M, Arp HPH, Zhu YG (2013) Sewage sludge biochar influence upon rice (*Oryza sativa* L.) yield, metal bioaccumulation and greenhouse gas emissions from acidic paddy soil. *Environ Sci Technol* 47:8624–8632
69. Uchimiya M, Wartelle LH, Klasson T, Fortier CA, Lima IM (2011) Influence of pyrolysis temperature on biochar property and function as a heavy metal sorbent in soil. *J Agric Food Chem* 59:2501–2510
70. Karami N, Clemente R, Moreno-Jiménez E, Lepp NW, Beesley L (2011) Efficiency of green waste compost and biochar soil amendments for reducing lead and copper mobility and uptake to ryegrass. *J Hazard Mater* 191:41–48
71. Uchimiya M, Klasson KT, Wartelle LH, Lima IM (2011) Influence of soil properties on heavy metal sequestration by biochar amendment: 1. Copper sorption isotherms and the release of cations. *Chemosphere* 82:1431–1437
72. Ma JW, Wang H, Luo QS (2007) Movement-adsorption and its mechanism of Cd in soil under combining effect of electrokinetics and a new type of bamboo charcoal. *Environ Sci* 28:1829–1834
73. Beesley L, Marmiroli M (2011) The immobilisation and retention of soluble arsenic, cadmium and zinc by biochar. *Environ Pollut* 159:474–480
74. Beesley L, Dickinson N (2011) Carbon and trace element fluxes in the pore water of an urban soil following greenwaste compost, woody and biochar amendments, inoculated with the earthworm *Lumbricus terrestris*. *Soil Biol Biochem* 43:188–196
75. Debela F, Thring RW, Arocena JM (2012) Immobilization of heavy metals by Co-pyrolysis of contaminated soil with woody biomass. *Water Air Soil Pollut* 223:1161–1170
76. Zhou JB, Deng CJ, Chen JL, Zhang QS (2008) Remediation effects of cotton stalk carbon on cadmium (Cd) contaminated soil. *Ecol Environ* 17:1857–1860
77. Hartley W, Dickinson NM, Riby P, Lepp NW (2009) Arsenic mobility in brownfield soils amended with green waste compost or biochar and planted with *Miscanthus*. *Environ Pollut* 157:2654–2662
78. Fellet G, Marchiolli DVG, Peressotti A (2011) Application of biochar on mine tailings: effects and perspectives for land reclamation. *Chemosphere* 83:1262–1297
79. Choppala GK, Bolan NS, Mallavarapu M, Chen Z, Naidu R (2012) The influence of biochar and black carbon on reduction and bioavailability of chromate in soils. *J Environ Qual* 41:1–10
80. Mendez A, Gomez A, Ferreiro JP, Gasco G (2012) Effects of sewage sludge biochar on plant metal availability after application to a Mediterranean soil. *Chemosphere* 89:1354–1359
81. Suppadit T, Kitikoon V, Phubphol A, Neumnoi P (2012) Effect of quail litter biochar on productivity of four new physic nut varieties planted in Cadmium-contaminated soil. *Chilean J Agric Res* 72:125–132