

# Chapter 6

## Vertebrate Axial Patterning: From Egg to Asymmetry

Douglas W. Houston

**Abstract** The emergence of the bilateral embryonic body axis from a symmetrical egg has been a long-standing question in developmental biology. Historical and modern experiments point to an initial symmetry-breaking event leading to localized Wnt and Nodal growth factor signaling and subsequent induction and formation of a self-regulating dorsal “organizer.” This organizer forms at the site of notochord cell internalization and expresses primarily Bone Morphogenetic Protein (BMP) growth factor antagonists that establish a spatiotemporal gradient of BMP signaling across the embryo, directing initial cell differentiation and morphogenesis. Although the basics of this model have been known for some time, many of the molecular and cellular details have only recently been elucidated and the extent that these events remain conserved throughout vertebrate evolution remains unclear. This chapter summarizes historical perspectives as well as recent molecular and genetic advances regarding: (1) the mechanisms that regulate symmetry-breaking in the vertebrate egg and early embryo, (2) the pathways that are activated by these events, in particular the Wnt pathway, and the role of these pathways in the formation and function of the organizer, and (3) how these pathways also mediate antero-posterior patterning and axial morphogenesis. Emphasis is placed on comparative aspects of the egg-to-embryo transition across vertebrates and their evolution. The future prospects for work regarding self-organization and gene regulatory networks in the context of early axis formation are also discussed.

**Keywords** Vertebrate embryology • Axis formation • Cortical rotation • Spemann organizer • Dorsoventral patterning • Anteroposterior patterning • Embryonic induction • Nieuwkoop center • Anterior visceral endoderm • Gastrulation

---

D.W. Houston (✉)


Department of Biology, The University of Iowa, 257 BB, Iowa City, IA 52242, USA  
e-mail: [douglas-houston@uiowa.edu](mailto:douglas-houston@uiowa.edu)

## Abbreviations

AP	Anteroposterior, Anterior-to-posterior
AVE	Anterior visceral endoderm
BMP	Bone morphogenetic protein
CRD	Cysteine-rich domain
DEP domain	Dishevelled, Eg110, Pleckstrin
DFC	Dorsal forerunner cell
DIX domain	Dishevelled, Axin
DV	Dorsoventral, Dorsal-to-ventral
dYSL	Dorsal yolk syncytial layer
EMT	Epithelial-to-mesenchymal transition
EpiSC	Epiblast stem cells
ES	Embryonic stem
EVL	Enveloping layer
FGF	Fibroblast growth factor
GPCR	G protein-coupled receptor
HMG	High mobility group
ICM	Inner cell mass
MAPK	Mitogen-activated protein kinase
MBT	Mid-blastula transition
MPF	Maturation promoting factor
PCP	Planar cell polarity
PDZ domain	Postsynaptic density protein (PSD95), Disc large tumor suppressor (Dlg1), and zonula occludens1 protein (ZO-1)
PMZ	Posterior marginal zone
TALEN	TAL-effector nuclease
TE	Trophectoderm
Tgfb	Transforming growth factor beta
UV	Ultraviolet irradiation

## 6.1 Introduction

Bilaterality is a central feature of animal body organization. In certain invertebrates, such as some insects and cephalopods, this feature is determined by the structure of the egg itself (Wilson 1928), but vertebrates and many other animals define the plane of bilateral symmetry *de novo* in each embryo. In vertebrates, this plane is ultimately defined by the formation of the generalized vertebrate tissues, the dorsal neural tube, notochord and somites. The vertebrate body axis fully forms during gastrulation, following the internalization of axial mesendoderm at the future dorsal midline of the embryo (Fig. 6.1). This event initiates at the dorsal (upper) lip of the forming blastopore, the importance of which was first clearly recognized by



**Fig. 6.1** Vertebrate axial organization. (a) Diagram of a sagittal section through a *Xenopus* gastrula, showing the involution of the dorsal mesoderm (d.m., dark red) at the dorsal lip. The neural plate (n.p., blue) overlies the dorsal mesoderm. bl blastocoel, v.l.m. ventrolateral mesoderm (orange), e endoderm (yellow). (b) Sagittal (left panel) and coronal (right panel) diagrams of a tailbud-stage *Xenopus* embryo showing the elongated anterior-to-posterior axis and organization of tissues within. The neural tube is located dorsally and will form the entire central nervous system (c.n.s.). The dorsal mesoderm gives rise to the notochord and somites, ventrolateral mesoderm (v.l.m.) will form the kidneys, body wall muscles and vascular system. The endoderm forms the gut and its derivative organs. The cement gland (c.g.), a larval amphibian anchoring structure, is shown at the anterior end. After Hausen and Riebesell (1991)

Spemann in amphibians, and was defined as the “organizer” of axis formation (Spemann 1918; Spemann and Mangold 1924). Spemann performed transplantation experiments in salamanders, demonstrating that the dorsal lip could induce and organize a normally patterned second body axis when grafted to the opposite (ventral) side of a host gastrula. In this “secondary embryo,” the organizer cells contributed mostly to notochord themselves, whereas host cells populated the bulk of the induced axis, which included neural tube, somites, intermediate mesoderm and gut endoderm. Additional experiments showed that organizer might also contribute to anteroposterior patterning of the embryo, demonstrating a central role for the organizer tissue in controlling cell interactions during development.

Although these main findings were firmly established by the 1930s, it was not until the 1990s that the cellular and molecular mechanisms underlying the action of the organizer were revisited, resulting in the identification of conserved growth factor antagonists and transcription factors. The background and history of this work has been written about exhaustively by Spemann and his contemporaries and later by modern authors (Spemann 1938; Waddington 1940; Hamburger 1988; Grunz 2004). As outlined later in this chapter, the conservation of the organizer extends to the cellular and genetic levels and largely defines the core mechanisms of early vertebrate body plan formation.

In contrast to the conservation of the organizer and its components, the ultimate origins of axial bilateral symmetry in vertebrates are seemingly more diverse. Axis formation was first extensively studied using amphibians and was linked to cytoplasmic localizations in the egg. This was evident in the formation of a natural marker of the future dorsal side, what came to be called the “gray crescent” (Roux 1888). Early mechanistic studies suggested the crescent formed by rotation of the outer cortex over the yolky inner cytoplasm (reviewed in Clavert 1962; Ancel and Vintemberger 1948). This “cortical rotation” was verified by later authors and found to involve the

organization and polarization of microtubules dorsally and the transport of dorsalizing determinants (Gerhart et al. 1989). Similar overall patterns are seen in primitive fish (Clavert 1962), suggesting that axis specification through cortical rotation in the fertilized egg is an ancestral condition in vertebrates.

By contrast, sauropsids (birds and reptiles) and more derived fish (teleost and selachians/dogfish) lack an obvious physical marker of dorsoventral polarity. These eggs contain abundant yolk and undergo discoidal cleavage, and axis formation occurs after significant cleavage in the blastoderm. In birds and reptiles, evidence suggests that rotation of the egg during passage through the oviduct affects axis formation in the blastoderm. Similar gravitational mechanisms were originally thought to exist in dogfish and teleosts (Clavert 1962), although recently, mechanisms involving cytoskeletal polarization in the cortex, analogous to the amphibian cortical rotation have been found in teleosts (zebrafish and medaka).

With the exception of the egg-laying monotremes, which undergo discoidal cleavage and are likely similar to reptiles with regard to axial patterning, mammals represent a significant divergence from this broad trend. The eggs of therian mammals have lost yolk, reverted to holoblastic cleavage (secondary holoblastic cleavage) and evolved the blastocyst structure to facilitate implantation. Consequently, the first cell fate decisions are centered on distinguishing the embryo proper from extraembryonic lineages rather than on establishing bilateral symmetry. Axial patterning is thus rather late, only becoming apparent after implantation, about a week into development. Early blastomeres retain pluripotency for an extended time and axis formation requires multiple reciprocal interactions with extraembryonic tissues.

Although there was evidence that formation of the organizer depended on polarization of the egg, the mechanisms connecting the two were totally unknown to early embryologists. Studies in amphibians unexpectedly found that the organizer was itself formed through induction, rather than by inheriting gray crescent material. This organizer-inducing activity was predominantly found in dorsovegetal cells of the blastula, later termed the “Nieuwkoop center” after its discoverer, and its formation depended on cortical rotation (Gerhart et al. 1989). These experiments were a critical link in the chain of causality from egg to organizer and were represented in various three- and four-signal models familiar to developmental biologists (Slack 1991). The cortical rotation → Nieuwkoop center → organizer model has been a useful conceptual tool and has directed much of the research into the molecular basis for these processes and their conservation across vertebrates. It is now appreciated that cortical rotation results in dorsal Wnt/beta-catenin signaling, activating Transforming growth factor beta (Tgfb)/Nodal signaling in the vegetal cells, which induce and pattern the organizer in the overlying equatorial cells. Analogous mechanisms have been found acting in the teleost dorsal yolk syncytial layer (dYSL) of the egg and in the avian posterior marginal zone (PMZ) epiblast, based on molecular and functional data, suggesting deep conservation of these processes in the early vertebrate lineage.

Recent cellular and molecular characterization of axis formation and patterning has produced a wealth of examples of such deeply conserved vertebrate developmental mechanisms. Vertebrate embryology has historically been a comparative science, with investigations encompassing a wide range of diverse organisms. More recently,

the use of specialized molecular and genetic approaches has largely limited the study of axis formation to a few tractable vertebrates, notably the mouse, chicken, *Xenopus*, and zebrafish. However, these species are all fairly evolutionarily derived representatives of their respective groups, making inference of primitive vertebrate characters problematic. With the growing ease of high-throughput genome analyses, stem cell technology and programmable genome editing, the barriers to performing comparative molecular and genetic studies are becoming increasingly reduced, potentially heralding a return to a broad comparative approach to vertebrate development.


In the context of the egg-to-embryo transition, the formation of the body axis is perhaps a defining process, since early developmental processes become organized into a unit comprising an individual. Indeed the idea of individuality in twinned embryos was an inspiration for Spemann to begin studying the embryological mechanisms of axis formation (Hamburger 1988), and remains relevant in current bioethics arguments regarding human embryos. This chapter shall review the core concepts relating to the origins and patterning of the axis, focusing on recent advances in understanding intracellular reorganizations, intercellular signaling events and cellular migrations. Emphasis has been given to recent molecular advances in the context of first discoveries and initial functional studies. Many of the ideas in this chapter have been extensively reviewed separately in the context of certain organisms, molecules or individual processes, but this chapter will attempt to tie these together to generate a more unified picture of axial development throughout the vertebrates.

## 6.2 Origins of Axial Polarity in the Egg and Early Embryo

The process of determining the initial plane of bilaterality and axis formation was first examined closely in the amphibians, where the gray crescent served as marker of the future axis (Fig. 6.2). Like most vertebrate eggs, amphibian eggs are initially symmetrical about the animal–vegetal axis (axisymmetrical), with the animal pole being the site of polar body extrusion, by definition, and more darkly pigmented. The vegetal pole is less pigmented and more concentrated with yolk. Using localized fertilization of frog eggs (*Rana (Lithobates) spp.*), Newport and Roux showed that the meridian of sperm entry coincided with the embryonic midline (and often the first cleavage plane), with the dorsoanterior axis forming opposite the sperm entry side (Newport 1851, 1854; Roux 1885, 1887). Importantly, Roux noted an apparent shifting of the animal–vegetal axis toward the sperm entry point, along with the appearance of a lightened area in the pigment on the opposing side. This feature formed before first cleavage and indicated the axial plane of the embryo, irrespective of the cleavage plane, which could be highly variable relative to sperm entry, depending on species (Roux 1887, 1888). Later studies confirmed these observations, further showing that the position of this “gray crescent”<sup>1</sup> strongly

---

<sup>1</sup>The nomenclature of the gray crescent has been quite variable. Roux (1888) referred to this feature as a “crescent-shaped gray seam” (*halbmondförmigen grauen Saumes*). Morgan and Tsuda



**Fig. 6.2** Gray crescent formation in amphibians. *Top panel*, diagram of an amphibian egg (e.g., *Rana*) before (*left*) and after fertilization (*right*). The heavily pigmented animal pole (an) and the paler vegetal pole (veg) are indicated. After fertilization, corticocyttoplasmic movements opposite to the sperm entry point (s.e.p.) result in the appearance of the gray crescent (g.c.) on the prospective dorsal side. *Bottom panel*, images of a *Rana* egg at fertilization (**a**), and at 20 min post-fertilization, showing the gray crescent (**b**; dorsal view, *arrow*). *Bottom panel* reproduced from Rugh (1951)

predicts the area in which the future dorsal lip of the blastopore would form (Morgan and Tsuda 1894; Schultze 1899; Roux 1903; Morgan and Boring 1903; Brachet 1904). These classical embryological observations established that the dorsal axis and bilateral symmetry were determined upon fertilization of the egg and likely occurred prior to the first division, thus disputing the long-held idea that the cleavage itself was determinative.

Despite the important nature of the connection between the gray crescent and the dorsal axis, the mechanisms of gray crescent formation and function remained elusive for many years. In the first comprehensive effort to understand the cellular changes underlying gray crescent formation, Ancel and Vintemberger (1948; reviewed in Clavert 1962) examined the movement of electrocautery wounds that were made either in the deep yolk or on the surface of frog eggs. The motion of these markers revealed that outer egg cortex moves dorsally relative to the stationary deeper yolk cytoplasm. Although other models were also considered, such as asymmetric cortical contraction, a later series of marking and egg manipulation studies in *Xenopus* largely substantiated and clearly documented the cortical rotation model (Vincent et al. 1986; Vincent and Gerhart 1987).

---

(1894) used the term “white crescent,” whereas Morgan and Boring (1903) used “grey crescent” [sic], translated as “*graue Feld*” (gray field). Later this became universally referred to as the gray crescent/*grauer Halbmond/croissant gris*.

Measurements of relative cortical displacement using superficially or deeply placed fluorescent dyes found that cortical rotation begins about halfway through the first cell cycle, eventually covering an average 30° of arc over the dense yolky cytoplasm. Also, cortical movement progressed along an animal–vegetal meridian, and toward the future dorsal side of the egg (generally away from the sperm entry point), with the region of greatest movement correlating with the position of the axial midline (Vincent et al. 1986; Gerhart et al. 1989). It is thought that changes in fluid dynamics in the fertilized egg result in a low viscosity/high elasticity shear zone in the subcortical region as well as an increase in firmness in the deep cytoplasm, allowing this differential movement between two parts of the egg (Elinson 1983; Gerhart et al. 1989).


In an extensive comparison of axis forming mechanisms, Clavert (1962) indicates that, in addition to amphibians, primitive fish including lampreys, lungfish, and chondrosteian fish (sturgeons and paddlefish) likely form gray crescents, suggesting that these organisms also likely undergo cortical rotation. With these older comparative studies and more recent molecular data taken together, it is apparent that the basics of the amphibian cortical rotation model are conserved throughout the anamniotes (ichthyosaurs). And vestiges may exist even in some amniotes. It is also now generally appreciated that cortical rotation establishes a self-organizing, transient microtubule polarity in the zygote that is critical for the transport of cortical cytoplasmic dorsal determinants and activation of Wnt/beta-catenin signaling (reviewed in Gerhart 2004; Weaver and Kimelman 2004; Houston 2012). Wnt/beta-catenin signaling (see Sect. 6.3.5) also plays a key role in bird and mammal axis formation, but this is likely controlled by mechanisms other than cytoplasmic localization.

## 6.2.1 Mechanisms of Amphibian Cortical Rotation

### 6.2.1.1 Cortical Rotation in Anurans

A number of studies have shown that cortical rotation is primarily controlled by the assembly of parallel microtubule arrays in the vegetal cortex (Fig. 6.3). Treatment of fertilized eggs during the middle of the first cell cycle with microtubule-depolymerizing agents such as ultraviolet irradiation (UV), exposure to anti-microtubule drugs, cold and high pressure, can inhibit gray crescent formation and/or block axis formation in both frog and salamander eggs (Malacinski et al. 1975; Manes et al. 1978; Manes and Elinson 1980; Scharf and Gerhart 1983; Vincent and Gerhart 1987). Correspondingly, impressive arrays of parallel microtubules are assembled in the vegetal cortical region during the period of cortical rotation in *Xenopus* and *Rana* (*Lithobates*) (Elinson and Rowing 1988).

Microtubules are completely disassembled in the egg at fertilization, but progressively repolymerize over the first 35 min, approximately when relative cortical movement begins. Microtubules in the cortical region initially form a disorganized network that gradually becomes organized as cortical rotation progresses. By mid-cortical rotation, microtubules are predominantly oriented with the plus ends



**Fig. 6.3** Events of cortical rotation in *Xenopus*. Microtubules are disassembled during oocyte maturation, and are absent from the egg cortex (*left panels*). Certain RNAs are localized to the vegetal cortex during oogenesis (blue) and encode proteins critical for cortical rotation and dorsalization (e.g., *trim36*, *wnt11b*). After fertilization, the incoming sperm pronucleus and associated centrosome initiate astral microtubule assembly. Cortical microtubule assembly also begins, forming a network by 40 min post-fertilization. A shear zone forms and microtubules associate with the yolk cytoplasmic core (not shown) and cortical rotation begins, under the action of kinesin-like proteins (kinesins). Relative cortical movement occurs dorsally, possibly the result of nudging by ventrally positioned astral microtubules, and rapidly orients microtubule plus ends dorsally (Olson et al. 2015) (*middle panel*). Microtubule assembly and organization becomes robust by 60 min post-fertilization and full cortical rotation commences, continuing until first cleavage. Rapid transport of dorsalizing activity occurs along parallel microtubule arrays using kinesin-like motors (*right panel*). The corresponding bottom panels show live images of microtubules labeled with Enconsin microtubule-binding domain tagged GFP (EMTB-GFP), showing progressive assembly and alignment during cortical rotation (Olson et al. 2015)

towards the dorsal side (Houliston and Elinson 1991; Olson et al. 2015). Live imaging studies indicate that the arrays associate and move with the deeper cytoplasm (Houliston 1994; Larabell et al. 1996; Olson et al. 2015). These microtubule arrays are transient and are progressively depolymerized upon first cleavage under the control of MPF activation (Marrari et al. 2003), thus terminating cortical rotation.

Surprisingly little is known about the regulation of microtubule activity during cortical rotation. Generalized kinesin-related protein activity in the cortex proper is thought to tether the microtubule array to the cortex and facilitate movement (Marrari et al. 2003). This has been assessed using function-blocking antibodies but specific roles for individual kinesins have not been identified. Kinesin1/Kif5b appears dispensable in *Xenopus* (Marrari et al. 2000, 2003), and Dynein has been shown to act early in rotation, as shown by injection of the antagonistic Dctn2 (Dynamitin/p50; Marrari et al. 2004). Recently, a suite of mRNAs localized to the vegetal pole in oocytes has also been implicated in regulating microtubule assembly. Maternal mRNA depletion experiments show that reductions in *perilipin2* (*plin2*; Chan et al. 2007), *tripartite motif containing 36* (*trim36*; Cuykendall and Houston 2009), and *dead end homolog 1* (*dnd1*; Mei et al. 2013) lead to abnormal



microtubule array formation and failure of cortical rotation. Trim36 can function as a single RING-finger-type ubiquitin ligase, and this activity is essential for its role in microtubule assembly (Cuykendall and Houston 2009). Dnd1 is an RNA-binding protein required to tether *trim36* mRNA to the cortex, facilitating locally enriched Trim36 protein levels (Mei et al. 2013). Dnd1 is typically associated with germline specification (Weidinger et al. 2003), and it is not known whether these functions are related. The role of Plin2 is unclear. The protein is associated with lipid droplets (Chan et al. 2007), but a structural role for the *plin2* RNA has also been suggested (Kloc 2009). A different set of localized mRNAs are involved in vegetal microtubule organization and transport in the zebrafish zygote (Nojima et al. 2010; Ge et al. 2014), although with slightly different functions (see Sect. 6.2.2). It remains to be determined how these localized molecules interact with microtubule regulatory proteins and motor proteins to control microtubule assembly in cortical rotation.

The initial cue for the direction of cortical rotation in normal development is thought to be sperm entry, as this site is generally opposite the direction of movement. The central model for orientation of the array is a reciprocal positive feedback loop, during which random asymmetry in microtubule growth is refined and amplified by rotation of the cortex (Gerhart et al. 1989; Gerhart 2004). Microtubules growing into the cortex, originating from the sperm aster and within the cortex may provide the initial movement cue (Houlston and Elinson 1991; Schroeder and Gard 1992). Cortical movement then serves to progressively stabilize microtubule growth and formation in the same direction. High-resolution live imaging of microtubule assembly and orientation has verified that cortical rotation begins before there is visible bias in plus end directionality or microtubule alignment (Olson et al. 2015), an observation that was suggested from earlier studies but never directly shown (Larabell et al. 1996). Additionally, plus end orientation occurs almost as soon as cortical rotation begins, indicating that directionality is determined in a punctuated manner rather than progressively (Olson et al. 2015).

In vivo, sperm entry or slight asymmetry with respect to gravity could be sufficient to initiate cortical movement, although a “vector summation” of microtubule polymerization forces, as initially proposed (Gerhart et al. 1989) cannot be ruled out. The shear-induced alignment of organelles (endoplasmic reticulum) may also play a role in perpetuating alignment, since ER and microtubules are often interdependent (Terasaki et al. 1986). Because cortical movement can have a role in determining the orientation of microtubules, the overall role of cortical rotation may be thought of as twofold; first to generate relative displacement of the cortex, and second to align the microtubule array facilitating the faster and longer range transport of determinants.

Evidence for these determinants came from 90° egg tipping experiments, which cause the axis to form in the uppermost part of a tipped egg (Ancel and Vintemberger 1948; Kirschner et al. 1980; Gerhart et al. 1981). Also, tipping can rescue axial development following UV-irradiation (Scharf and Gerhart 1980; Chung and Malacinski 1980). In amphibian eggs, denser yolk accumulates in the vegetal pole, which when tipped off axis, results in a tendency to “fall” downward against the cortex, which is immobilized in these experiments, creating relative displacement. Tipping does not restore microtubules (Zisckind and Elinson 1990), further suggesting that the relative

displacement of cortical dorsal determinants is essential, whether achieved normally by microtubule motive force or experimentally by gravitational force.

Other studies indicated the existence of an essential, transplantable dorsalizing activity associated with the cortex/subcortical cytoplasm (Yuge et al. 1990; Hainski and Moody 1992; Holowacz and Elinson 1993; Kikkawa et al. 1996; Kageura 1997). And, live imaging studies have shown various substances moving dorsally within the shear zone during cortical rotation. These include a subset of pigment granules and organelles, fluorescent beads, and certain GFP fusion proteins (Miller et al. 1999; Weaver et al. 2003). Their movement is rapid ( $\sim 50 \mu\text{m}/\text{min}$ ) and saltatory, consistent with generalized kinesin-based transport along microtubules. Transport can be measured from  $30^\circ$ – $120^\circ$  of arc from the vegetal pole, equal to and greater than the overall relative cortical displacement (Rowning et al. 1997; Miller et al. 1999; Weaver et al. 2003). Interestingly, this distribution matches that of dorsalizing cytoplasm taken from the egg (Fujisue et al. 1993; Holowacz and Elinson 1993). Additionally, stimulation of microtubule assembly with deuterium can hyperdorsalize embryos, potentially through wide-spread distribution of this dorsalizing material along many egg meridians (Scharf et al. 1989; Miller et al. 1999). The identity of the molecules responsible for the activity of this cytoplasm in vivo is unclear but are likely related to Wnt/beta-catenin signaling (see Sect. 6.3.2).

Cortical rotation can thus be considered a robust self-organizing symmetry-breaking process that integrates cytoskeletal and physical forces to generate a single direction for the short-range relative displacement of the cortex and for the long-range distribution of molecules and putative determinants towards the presumptive dorsal side.

### 6.2.1.2 Cortical Rotation in Urodeles

Although much of the recent cell and molecular characterization of cortical rotation has been done in anurans (*Xenopus* and *Rana* (*Lithobates*)), urodeles *Triturus* and *Ambystoma* are known to form gray crescents (Bánki 1927; Clavert 1962). However, urodele eggs are normally polyspermic and the relationships between the site of sperm entry or male pronucleus formation and the site of the gray crescent are unclear. Recently, relative cortical displacements analogous to those in *Xenopus* have been observed in *Cynops* (Fujisue et al. 1991), which also exhibits vegetal microtubule array assembly during the period of cortical rotation (Iwao et al. 1997). Curiously, although some species are ventralized by UV-irradiation (see above), irradiation of *Cynops* eggs dorsalizes embryos (Doi et al. 2000), suggesting that putative dorsal determinants are more widely dispersed in these eggs and would remain so in the absence of microtubule assembly and cortical rotation. This situation may possibly mimic the random dispersion of determinants occurring in deuterium-treated *Xenopus* eggs. Thus, the basic mechanisms of microtubule-dependent cortical rotation and dorsal determinant transport are conserved in amphibians.

Urodeles are thought to lack vegetal cortical localization of RNAs (Elinson and del Pino 2011; Houston 2013), which is interesting given the connection between


localized RNAs and cortical rotation in *Xenopus* and zebrafish. However, it has not been specifically demonstrated whether the exact RNAs implicated, including *trim36* and *syntabulin* (*sybu*), are in fact unlocalized in urodeles. Since these RNAs are partially associated with the germ plasm, which is not found in urodeles (Nieuwkoop and Sutasurya 1979), one might expect an absence of localization. Urodeles may however localize important components posttranslationally. Nevertheless the basic mechanisms of polarizing the egg and distributing dorsal determinants appear conserved, but are not well understood in either case.

### 6.2.2 Cortical Rotation and Dorsal Determinant Transport in Zebrafish

Axis formation in zebrafish similarly relies on asymmetric localization of dorsal determinants and activation of Wnt/beta-catenin signaling. The polarizing mechanisms and the similarity of these to classical amphibian cortical rotation are only now becoming apparent, however. It has been traditionally thought that typical cortical rotation does not occur in teleost fish. The origin of this assumption is mysterious, but likely can be attributed to initial observations on the importance of a polarized dYSL in teleost axial patterning rather than formation of a gray crescent-like clear crescent, which does occur in non-teleosts (Long 1983; Ho 1992). There do not appear to have been any classical embryological studies directly addressing either relative displacement of cytoplasm and cortex or the existence of transplantable vegetal cortical cytoplasm.

However, parallel microtubule arrays have been noted at the vegetal pole cortex of the early post-fertilization (~20 min) medaka and zebrafish egg (Jesuthasan and Stähle 1997). By 30 min post-fertilization, this array is offset from the vegetal pole, giving bilateral symmetry to the egg. During cleavage of the blastoderm, a second set of microtubule arrays forms along the animal–vegetal axis (Strähle and Jesuthasan 1993; Solnica-Krezel and Driever 1994), and polarized transport of fluorescent beads has been observed to move animally into the YSL and marginal blastomeres (Jesuthasan and Stähle 1997). Disruption of either set of microtubules with UV, cold, or nocadazole disrupts axis formation as well as epiboly (Strähle and Jesuthasan 1993; Solnica-Krezel and Driever 1994; Jesuthasan and Stähle 1997).

Thus a two-step transport pathway is proposed (Fig. 6.4). Asymmetry is initially established by the localization of determinants to the future dorsal vegetal side of the egg, followed by generalized upward movement of material into the YSL and marginal blastoderm cells (the dorsal determinants being carried along only on the dorsal side). In support of this idea, a recent live imaging study has demonstrated for the first time, as was tacitly assumed, that the plus ends of zebrafish vegetal cortical microtubules are oriented dorsally as in frogs (Tran et al. 2012). Given the relationship between movement of the cytoplasmic core and microtubule orientation in *Xenopus* eggs (Olson et al. 2015), it is likely that relative cortical movement, at least locally, is involved in orienting these microtubules in teleost eggs as well.



**Fig. 6.4** Dorsal determinant transport in zebrafish. (a) Sequence of events in wildtype embryos. RNAs and other dorsal determinants are localized vegetally during oogenesis (blue). After fertilization, cytoplasm streams to the animal pole, forming the future blastoderm. Microtubule assembly initiates ~20 min post-fertilization at the vegetal pole of the yolk cell; localized RNAs and Syntabulin protein (Sybu) are shifted toward the future dorsal side. Microtubule networks in the lateral cortex facilitate global transport animal-ward, which on the dorsal side would contain axis determinants. (b) In *hecate* (*hec*) mutants lacking Grip2a, maternal vegetal localization occurs, but cortical rotation and microtubule assembly are deficient post-fertilization. This image is reproduced and modified from, Ge X, Grotjahn D, Welch E, Lyman-Gingerich J, Holguin C, Dimitrova E, et al. (2014) Hecate/Grip2a acts to reorganize the cytoskeleton in the symmetry-breaking event of embryonic axis induction. PLoS Genet 10(6): e1004422. doi:10.1371/journal.pgen.1004422, under the terms of the Creative Commons Attribution License (CC BY 4.0)

Additional insight into microtubule assembly in zebrafish and further support for the dual-range model of transport in the yolk cell has recently come from analysis of maternal-effect mutants in zebrafish (Nojima et al. 2004, 2010; Lyman-Gingerich et al. 2005; Ge et al. 2014). The mutants *hecate* (*hec*; Ge et al. 2014) and *tokkaebi* (*tkk*; Nojima et al. 2010) are ventralized with near complete penetrance and harbor disruptions in *syntabulin* (*sybu*) and *glutamate receptor interacting protein 2a* (*grip2a*) loci, respectively. Parallel microtubule assembly at the vegetal pole is normal in *tkk* mutants but disrupted in *hec* mutants. Upward movement to the YSL/

blastoderm margin is normal in *hec* eggs, underscoring the independence of these two transport systems. *Sybu* encodes a potential cargo linking protein for Kif5b, suggesting a role in microtubule transport of dorsal determinants vegetally. *Grip2a* encodes a scaffolding protein important for subcellular localization in mammalian neurons.

Both *sybu* and *grip2a* mRNAs localize to the vegetal cortex, along with *wnt8a* mRNA (see below), and these RNAs all undergo an off-center “shift,” mirroring that of the microtubule array (Nojima et al. 2010; Lu et al. 2011; Ge et al. 2014). *Sybu* protein fails to localize to the prospective dorsal side in nocodazole-treated eggs, suggesting it is trafficked by microtubules. The exact role of *Grip2a* is not known but it may recruit protein complexes to vesicles or organelles that attach to and help align microtubules. Interestingly, *grip2* mRNA is localized in *Xenopus*, but follows a germ plasm-like pattern and is not thought to play a role in axis formation (Tarbashevich et al. 2007). Similarly, *sybu* is localized to the germ plasm in *Xenopus* and may play an undefined role in axis formation, possibly in transport or in Wnt activation (Colozza and De Robertis 2014).


In zebrafish, maternal loss-of-function mutants have implicated *kif5ba* in vegetal microtubule formation and axis formation (Campbell et al. 2015), although its role is complex. Organized vegetal microtubules fail to form and *wnt8a* does not shift dorsally. However, *grip2a* asymmetric translocation still occurs and *sybu* RNA is not maintained vegetally (Campbell et al. 2015). It is unclear to what extent these phenotypes reflect roles for Kif5ba in localizing components vegetally during oogenesis or more acute roles during microtubule organization and transport.

### 6.2.3 *Asymmetry in Early Amniote Embryos*

Initial axis formation in fish and frogs occurs in the fertilized egg; the dorsal determinants are either inherited directly by dorsal cells (frogs, primitive fish) or transmitted from the uncleaved yolk cell to peripheral dYSL and overlying dorsal marginal blastomeres (teleost fish). The axis in amniotes (birds, reptiles and mammals) relies on mainly on reciprocal interactions between upper embryonic and lower extraembryonic tissue layers (epiblast and hypoblast, respectively) and asymmetric cell movements, with only hints that early asymmetry in the egg or early embryo are involved. Additionally, the links to localized activation of growth factor signaling pathways are much less clear.

#### 6.2.3.1 *The Role of Gravity in Axis Formation in Sauropsids*

Classic experiments in the chick have suggested that axis specification occurs in response to gravity as the egg rotates as it passes through the oviduct (Fig. 6.5). Axis formation in reptile embryos is thought to occur in a similar fashion, although is less thoroughly studied in this regard. Bird and reptile embryos are both highly polyspermic, making it unlikely that sperm entry plays a role in axis formation



**Fig. 6.5** Model for establishment of asymmetry in bird eggs. *Left*, sectional view of a uterine chicken egg viewed from the sharp end. The direction of rotation is indicated; because of this rotation, the lighter blastoderm cytoplasm is maintained off angle as it continually floats to the highest point. The blastoderm is exposed to the subgerminal cytoplasm, which is hypothesized to contain axis determinants (*blue*). At this stage, the blastoderm is several thousand cells and has not formed the area pellucida epiblast. *Right*, top view of 2–3 day embryo showing anterior-to-posterior axial polarity. This embryo would conform to von Baer’s rule, with head oriented away with the blunt end positioned left. *ant.* anterior, *post.* posterior

(Waddington 1956). The long-noted von Baer’s rule of thumb suggested a relationship between the axis of the egg and that of the embryonic axis (von Baer 1828). This axis is most often (~70%) situated perpendicular to the long axis of the egg, with the left side of the embryo oriented towards the blunt end. Eggs rotate clockwise about the long axis as they travel through the oviduct (~0.2 rpm) and continue rotating in the uterus, where they acquire the shell membranes and shell. As the egg rotates, the blastoderm is maintained at an angle of ~45°, balancing the inertia of the rotation with the tendency of the buoyant blastoderm to float on the dense yolk (Clavert 1962). In this arrangement, the anterior of the embryo forms at the lower end. Most eggs enter the uterus and are laid sharp-end first and thus end up following von Baer’s rule. The minority of cases where the blunt end enters first are thus truly exceptions that prove the rule, since it is only the embryo’s orientation with respect to the external egg shape that is changed; the posterior of the embryo still forms at the upper end of the blastodisc.

Both in utero and in vitro experiments have defined a critical period for axis establishment in the uterus (Vintemberger and Clavert 1959; Clavert 1961; reviewed in Clavert 1962). Egg orientation was manipulated at different times prior to egg-laying and the orientation of the embryonic axis was altered if the presentation of the egg was changed at least six hours prior to laying. Additionally, eggs could be removed and incubated in vitro in a rotating cylinder and the orientation of the embryo was determined by the relative direction of rotation. Similar to the experiments in frogs, the effects of earlier rotations could be overridden by later manipulations up to the critical period of axis formation. This period correlates with the time

when the chick blastoderm thins to a single layer, forming the *area pellucida* epiblast. Another set of studies showed that uterine eggs could be incubated in a variety of orientations without rotation, or even hung from a chalaza without shells, and the axis always formed along the gravitational axis with the posterior end uppermost (Kochav and Eyal-Giladi 1971; Eyal-Giladi and Fabian 1980). Thus, it is the response to gravity that is critical, not the stress of movement or effect of rotation per se.

Axial polarity and bilateral symmetry in the blastoderm is evident both morphologically and molecularly prior to primitive streak formation. The future posterior half of the embryo, in which the primitive streak forms, can first be distinguished by the formation of a ridge of cells in the deep layer of the posterior *area opaca*, Koller's/Rauber's sickle (Callebaut and Van Nueten 1994). Additionally, the hypoblast layer (analogous to the anterior visceral endoderm, see below) begins to form in this region, coalescing from isolated hypoblast islands delaminating from the epiblast in a posterior-to-anterior progression. The hypoblast is then replaced by the endoblast (posterior visceral endoderm), derived from Koller's sickle, in the same progression (Stern 1990; Stern and Downs 2012). Molecular analyses have also identified early differential gene expression in this region, including *Gdf1* (*alias cVg1*; Seleiro et al. 1996; Shah et al. 1997) in the PMZ of the epiblast and *Goosecoid* (*Gsc*) in Koller's sickle (Izpisúa Belmonte et al. 1993). Transplantation experiments showed non-cell/tissue autonomous induction of ectopic axes, sparking parallels between the PMZ and the amphibian Nieuwkoop center (see Sect. 6.3.5).

The mechanisms leading to these developmental events in the posterior are unknown. The prevailing hypothesis for the initiation of this posterior polarity is the differential exposure of presumptive areas to maternal cytoplasm during cleavage. In the chicken egg, a particular cytoplasmic layer, the subgerminal ooplasm (gamma- and delta-ooplasm) is contained in a central region below the blastodisc and overlying the latebra and Nucleus of Pander (Callebaut 2005). With the blastoderm offset from the animal pole by the inertia of the rotating egg (see above), this cytoplasm would have more prolonged contact with cells arising in the upper (future posterior end) of the embryo. The subgerminal cytoplasm may also be differentially inherited by primordial germ cells, which form in response to cytoplasmic determinants (germ plasm) in birds (Tsunekawa et al. 2000). It is thus possible that unknown axial determinants might also be localized to this region, as in amphibians.

### 6.2.3.2 Early Polarization of the Mammalian Embryo

The evolution of implantation in therian mammals resulted in many changes in the structure of the egg and early embryo, including a loss of yolk, the reemergence of holoblastic cleavage and the early segregation of embryonic and extra embryonic tissues, forming a preimplantation blastocyst (see Chap. 4). In light of these major alterations in life history, it has long been of interest to determine the extent that axes in the mammalian embryo are determined by cytoplasmic asymmetries as in other vertebrates.


Early authors concluded that this must be the case, although these conclusions were admittedly based on a few cases of poorly characterized abnormal embryos generated following blastomere perturbations (Waddington 1956). However, in contrast to amphibians, the separation of early mammalian (rodent and rabbit) blastomeres does not result in complementary embryos either having or lacking dorsal structures (Seidel 1956; Tarkowski 1959; Tarkowski and Wróblewska 1967). Additionally, early mammalian blastomeres demonstrate a high degree of developmental plasticity, with each of the four-to-eight cell blastomeres contributing to all cell lineages in chimeric embryos (Tarkowski 1961; Mintz 1964; Kelly 1977). Furthermore, cell fate specification with respect to epiblast/primitive endoderm/trophectoderm is largely dependent on cell polarity related to inside or outside cell position within the morula, as well as on the timing of asymmetric cell division in generating inside cells (Hillman et al. 1972; Ziomek and Johnson 1980; Pedersen et al. 1986; Morris et al. 2010) (see also Chap. 4). Ablation experiments have found that removal of the animal or vegetal poles from fertilized eggs and early blastomeres is fully compatible with normal development (Zernicka-Goetz 1998; Ciemerych et al. 2000), unlike the case in amphibians. It is therefore generally concluded that segregation of maternal determinants in the egg is unlikely to direct axis formation or cell fate patterning in mammals, or that if such a bias exists it can be readily overridden by other cellular interactions.

Nevertheless, axis specification requires that symmetry breaking occur at some point prior to gastrulation. When this asymmetry is established and to what extent it depends on earlier developmental bias or is more or less random has been a recurring debate. There have been various attempts to correlate cleavage patterns in the early embryo with asymmetries in the blastocyst and conceptus and with the eventual anteroposterior axis of the embryo. A preponderance of the evidence however suggests that much of this observed “bilateral symmetry” likely results from physical constraints imposed by the *zona pellucida* (vitelline membrane) or other external constraints and is not connected to the orientation of the anteroposterior axis (for detailed reviews of this literature, please see Takaoka and Hamada 2012; Zernicka-Goetz 2013; Bedzhov et al. 2014 and references therein).

The most compelling evidence for an early cell fate bias is the observation that the order and orientation of rotational cleavages in the mouse embryo can influence blastomere fate in the blastocyst (Fig. 6.6). In particular, the vegetal blastomere (distal to the polar body) that arises from a particular tetrahedral cleavage pattern (which occurs in a subset of cases), will disproportionately contribute to the trophectoderm in normal embryos (Piotrowska-Nitsche et al. 2005; Torres-Padilla et al. 2007a). Furthermore, chimeras composed exclusively of vegetal blastomeres fail to survive (Piotrowska-Nitsche et al. 2005), likely because these cells cannot generate sufficient numbers of pluripotent epiblast cells to support development (Morris et al. 2012b).

Lineage labeling studies of individual or all four cells have found a similarly biased contribution of four cell-stage blastomeres to either inner cell mass (ICM) or trophectoderm (TE) fates in a subset of embryos (Fujimori et al. 2003; Tabansky et al. 2013). Importantly, this developmental bias was reflected in cell lineage but not in relative cell positioning toward the embryonic or abembryonic poles of the





**Fig. 6.6** Early bias of mouse blastomeres towards lineage fate but not axial polarity. Two-cell blastomeres undergo rotational cleavage (*dotted white lines* indicate cleavage planes), generating a fraction of embryos with a tetrahedral cell arrangement. In this formation, vegetal blastomeres are biased towards contributing to the trophoblast (*dark gray*) in the blastocyst. The corresponding animal blastomeres are biased towards contributing to the inner cell mass (*blue*). After Zernicka-Goetz et al. (2009)

blastocyst. For technical reasons however, the specific contribution of vegetal blastomeres could not be assessed in these studies. Mechanisms underlying this bias may include epigenetic regulation of cell polarity (Parfitt and Zernicka-Goetz 2010), decreased pluripotency transcription factor occupancy at target genes (Plachta et al. 2011) or a combination of factors. How this differential regulation is initiated is unknown, but the lack of maternal influences and a lack of differential gene expression in two- and three-cell blastomeres (VerMilyea et al. 2011) suggest that this bias is either an emergent property or a positioning effect in the four-cell stage embryo.

It is unclear at present whether any bias in early blastomere fate can be connected to axis specification in the mammalian embryo. Although this will be discussed further in Sect. 6.5.1 and similar to the bird embryo, the proximal events in mammalian axis formation involve the asymmetric migration of cells in the extraembryonic anterior visceral endoderm/hypoblast.

### 6.3 Initiation of Axis Induction by Dorsal Determinant Signaling

Numerous models have been suggested for how early asymmetries in the egg and embryo can lead to the specification of the organizer and ultimately to axis formation. Classical views, perhaps influenced by the importance of cytoplasmic localizations in invertebrates, suggested that the amphibian gray crescent contained precursors or determinants of the organizer (Wilson 1928). Another influential idea was that of a dorsal “cortical field” intersecting with a vegetal yolk gradient to determine the position of the organizer (Dalcq and Pasteels 1937). Later experiments showed that mesoderm in general, and the organizer in particular, required inductive cell–cell signaling by the vegetal prospective endoderm (see Chap. 7; Boterenbrood and Nieuwkoop 1973; Gimlich and Gerhart 1984; Dale et al. 1985), suggesting that cell-autonomous inheritance of organizer determinants was subordinate

to mesoderm induction. Importantly with respect to axis formation, Nieuwkoop and colleagues showed that the blastula vegetal mass is dorsoventrally patterned, with only the dorsovegetal cells being able to induce dorsal mesoderm/organizer. This dorsal signaling center, or “Nieuwkoop center” as it became known (Gerhart et al. 1989), was also demonstrated by transplantation of dorsal vegetal cells into UV-ventralized hosts (or ventrally into normal hosts), resulting in largely non-cell-autonomous organizer and axis induction (Gimlich and Gerhart 1984; Gimlich 1986; Kageura 1990).

Cortical rotation emerged as the candidate upstream event leading to Nieuwkoop center formation in dorsovegetal cells, as embryos ventralized by UV-irradiation lack both Nieuwkoop center and organizer activity (Smith et al. 1985; Gerhart et al. 1989). Also, because the extent of mesoderm induction is unchanged in ventralized embryos (Cooke and Smith 1987), a hypothesis was formed that the Nieuwkoop center generates a distinct dorsalizing signal or a competence modifying signal, which acts along with a general mesoderm inducer to induce the organizer. This idea became enshrined in the influential three-signal models of axis formation (Smith et al. 1985; Smith 1989; Heasman 1997). It is now recognized, owing to the work of many labs over many years, that this “dorsal signal” is not a unique signal at all, but represents an early and elevated wave of Nodal-related Tgfb signaling that is regulated by dorsally enriched Wnt/beta-catenin signaling and other maternal factors (see below).

Although many of these studies were conducted using *Xenopus* embryos, transplantation experiments have shown that localized regions in the blastula-equivalent stages of the zebrafish and chicken embryo can induce axes non-cell autonomously (dYSL, Mizuno et al. 1999; PMZ epiblast, Bachvarova et al. 1998). These regions also ultimately act through elevated Nodal signaling, either downstream of or in concert with Wnt/beta-catenin signaling, suggesting that the mechanisms of axis induction are widely conserved vertebrate development. In mammals however, Nodal signaling likely precedes obvious Wnt asymmetry and is the main determinant of axis formation, albeit in conjunction with Wnt signaling. In this section, the roles of early Wnt signaling in establishing dorsal fates in amphibians and fish are reviewed, along with the conserved but divergent roles of Wnt and Nodal signaling in regulating organizer formation across vertebrates.

### 6.3.1 Basic Wnt Signaling Mechanisms

Since its initial discovery as a mammalian oncogene (Nusse and Varmus 1982), signaling by the deeply conserved *Wnt1* (*int-1/wingless* (*wg*)) family of growth factors has emerged as a central feature of many aspects of animal development and disease. The reception of Wnt signals and intracellular signal transduction mechanisms has been extensively studied *in vivo* in both vertebrate and invertebrate organisms as well as in tissue culture cells. Although there are many variations that are important in specific tissues and disease states, three main arms of the pathway are widely implicated in vertebrate axis formation. These are: (1) the regulation of Ctnnb1

protein stability (beta-catenin protein hereafter), nuclear localization and transcriptional activity, (2) the regulation of cytoskeletal organization and cell polarity, and (3) the release of calcium from intracellular stores. With the caveats in mind that much of Wnt signaling entails complex, context-dependent and networked interactions, it remains useful to understand the basic features of Wnt pathways involved in early axis development. There are numerous comprehensive reviews on different aspects of Wnt signaling (MacDonald et al. 2009; Hikasa and Sokol 2013); here the key evidence of Wnt signaling in axis formation and the core signal transduction mechanisms most involved in axis formation and patterning will be briefly reviewed.


### 6.3.1.1 Wnt/Beta-Catenin Signal Transduction Mechanisms

In Wnt-unstimulated cells, beta-catenin protein is constitutively turned over by the activity of a multiprotein “destruction” complex (Fig. 6.7). This complex contains Axin1, Adenomatous polyposis coli (Apc), Glycogen synthase kinase 3 beta (Gsk3b), and Casein kinase 1 alpha (Ck1a/Csnk1a1), and serves to regulate the phosphorylation of cytoplasmic beta-catenin by Gsk3b and Ck1a (reviewed in MacDonald et al. 2009; Clevers and Nusse 2012; Hikasa and Sokol 2013). These phosphorylations in the beta-catenin N-terminus allow recognition by members of the beta-transducin repeat containing E3 ubiquitin protein ligase family (Btrc, Jiang and Struhl 1998; Liu et al. 1999a) and target phospho-beta-catenin for ubiquitylation, resulting in degradation by proteasomes (Aberle et al. 1997). Axin is thought to be a key limiting component of this complex, regulating the assembly of the destruction complex (Lee et al. 2003) and recruiting the beta-catenin kinases (Ck1a and Gsk3b) for the priming and processive phosphorylation, respectively, of beta-catenin (Liu et al. 2002; Amit et al. 2002).


Wnt stimulation inactivates the destruction complex through a little understood mechanism involving recruitment of the cytoskeletal adaptor protein Dishevelled (Dvl homologs 1–3). This blocks the activity of Gsk3b (Siegfried et al. 1992, 1994; Cook et al. 1996) and prevents beta-catenin phosphorylation, thereby allowing cytoplasmic accumulation and subsequent nuclear localization of beta-catenin (MacDonald et al. 2009; Clevers and Nusse 2012; Hikasa and Sokol 2013). Nuclear beta-catenin interacts with the LEF/TCF family of High Mobility Group (HMG) domain transcription factors (Brunner et al. 1997, see below) either to activate or to “derepress” transcription of Wnt-responsive genes (Fig. 6.7).

Signaling is initiated by binding of Wnts to one of seven-transmembrane domain Frizzled (Fzd) receptors plus a coreceptor, lipoprotein receptor-related protein Lrp5 or Lrp6 (Lrp5/6; reviewed in He et al. 2004). Fzds are heterotrimeric G protein-coupled receptors that are activated in response to Wnts (Slusarski et al. 1997b; Katanaev and Buestorf 2009). Wnts bind to Fzd through the receptor’s extracellular cysteine-rich domain (CRD), with key contacts being made between the Wnt lipid moiety and a separate hydrophobic “index finger” interacting with grooves in the CRD (Janda et al. 2012). Wnts also interact with the Lrp6 extracellular domain, leading to clustering of the receptors and coreceptors (Tamai et al. 2000; Mao et al. 2001; Kato et al. 2002;

- Wnt



+ Wnt



**Fig. 6.7** Generalized Wnt signaling networks. In the absence of activating Wnt ligands (*top panel*, -Wnt), beta-catenin protein (Ctnnb1) is phosphorylated by destruction complex components and tagged for proteasomal degradation. In the nucleus, Tcf711/Tcf3 represses Wnt target promoter activity through recruitment of Groucho. Upon stimulation with Wnt ligand, a variety of pathways are activated (see text for details). Predominantly positive-acting components with respect to beta-catenin regulation are shown in *green*, negative components in *red*, beta-catenin-independent components are *light blue*. Beta-catenin is shown in *yellow*. *Circles* indicate component nodes, *lines* indicate edges, or interacting components. This arrangement is not meant to convey specific exact binding relationships or stoichiometry. Wnt1 is shown as a beta-catenin-activating ligand, whereas Wnt5 is shown as a Wnt/PCP and Wnt/Calcium-stimulating ligand. Plot was generated with iGraph in R (Csardi and Nepusz 2014). *txn* transcription

Liu et al. 2003; Itasaki et al. 2003). The induced proximity of the intracellular domains of Fzd and Lrp6 is necessary and sufficient to initiate downstream signaling and inhibition of beta-catenin phosphorylation (Tolwinski et al. 2003).

The inhibition of beta-catenin degradation following Wnt receptor activation remains incompletely understood. Recent observations together with extensive data on biochemical interactions have suggested that Wnt-Fzd binding activates Dvl, possibly through GPCR activation, to recruit Axin/Gsk3b/Ck1a complexes to the Lrp6 intracellular domain, resulting in phosphorylation by Gsk3b and Ck1a (Mao et al. 2001; Tolwinski et al. 2003; Cliffe et al. 2003; Tamai et al. 2004; Davidson et al. 2005; Zeng et al. 2005, 2008; Egger-Adam and Katanaev 2010; Jernigan et al. 2010). Lrp6 phosphorylation occurs at PPPSPxS motifs, which serve as sites for additional Axin complex recruitment and are thought to directly inhibit destruction complex Gsk3b activity (Piao et al. 2008; Cselenyi et al. 2008; Wu et al. 2009). Furthermore, Axin itself is a Gsk3b substrate (Yamamoto 1999) and Gsk3b inhibition results in Axin dephosphorylation and its dissociation from phospho-Lrp6 and beta-catenin (Kim et al. 2013). Axin is then free to be either phosphorylated again to reconstitute beta-catenin destruction complexes or degraded. These data are consistent with a kinetic analysis of beta-catenin regulation, which suggests that Wnt signaling results in partial inhibition of both Gsk3b and Ck1a activities (Hernández et al. 2012). It is possible that this effect could be explained by the inactivation of a subset of limiting and finite destruction complexes through Axin dephosphorylation, which would then depress overall beta-catenin phosphorylation at the population level in a distributed manner.

In a separate but not necessarily mutually exclusive model, Dvl recruitment leads to multimerization of phospho-Lrp6-Fzd complexes, followed by accumulation of Dvl aggregates, leading to positive feedback recruitment and inactivation of destruction complexes (Metcalf and Bienz 2011; Dobrowolski and De Robertis 2012). There is also evidence that these receptor complexes are incorporated into signaling endosomes (Lrp6 signalosomes) to stabilize and amplify signaling (Bilic et al. 2007). Other data suggest that these signalosomes are eventually sequestered into multivesicular bodies, leading to the longer-term removal of Gsk3b activity and the inability to phosphorylate newly synthesized beta-catenin (Taelman et al. 2010).

### 6.3.1.2 Transcriptional Regulation by Wnt/Beta-Catenin Signaling

Transcriptional responses in response to beta-catenin are mediated by binding to Lymphoid Enhancer-binding Factor 1 (Lef1)/Transcription factor 7 (T-cell specific, HMG-box; Tcf7) proteins. These proteins are constitutively nuclear and typically repress target genes by recruiting Groucho family repressors (Roose et al. 1998; Barker et al. 2000). Beta-catenin accumulation can lead to displacement of Groucho and activation of target genes, through a combination of derepression and transcriptional activation, mediated by distinct Lef1/Tcf7 proteins. Tcf7l1 (Tcf3 hereafter) likely exclusively acts as a transcriptional repressor during early development, with Lef1 and Tcf1 proteins serving as activators, and Tcf7l2 (Tcf4) exhibiting

spliceform-dependent activator and repressive functions (Pukrop et al. 2001; Gradl et al. 2002; Wöhrle et al. 2007; Weise et al. 2010).

This protein family has diverged in function, with Tcf3 primarily performing a repressive role during early development and others acting as beta-catenin-dependent co-activators later in development (see Sect. 6.5.3). Tcf3 constructs lacking the ability to interact with beta-catenin, by deletion of the N-terminal beta-catenin-binding domain (deltaNTcf3), have been used to inhibit Wnt/beta-catenin-regulated transcription, as these cannot be derepressed or activated by beta-catenin. Expression of deltaNTcf3 during the cleavage stages efficiently ventralizes embryos (Molenaar et al. 1996; Pelegri and Maischein 1998), but fails to inhibit ventrolateral development or to block a late Wnt overexpression effect in *Xenopus* (i.e., anterior truncations; Hamilton et al. 2001). Additionally, experimental depletion of Tcf3 is sufficient to activate Wnt target gene expression during vertebrate axis development (Kim et al. 2000; Houston et al. 2002; Dorsky et al. 2003; Merrill et al. 2004) and in embryonic stem cells (Yi et al. 2008). Recent data from mouse studies in which the mutant deltaNTcf3 was knocked into the endogenous *Tcf711* locus have substantiated the idea that Tcf3-mediated repression is critical for its role in early development (Wu et al. 2012a). Gastrulation proceeded normally in these mice, suggesting that the proper amount of transcriptional derepression of Tcf3 targets can occur in the absence of beta-catenin-Tcf3 interactions during axis formation. However, beta-catenin interactions with Tcf3 and with other Lef1/Tcf7 proteins are required later in development and in cancer cells (Wu et al. 2012a; Shy et al. 2013).

Derepression of Tcf3 is sufficient for Wnt target gene activation, although co-activators are also required for normal development, suggesting both likely operate in vivo. Beta-catenin recruits a number of co-activators including p300 and the conserved nuclear complex containing Pygopus and Bcl9 proteins (Kramps et al. 2002; Parker et al. 2002; Belenkaya et al. 2002). Pygo/Bcl9 are thought to be dedicated to Wnt signaling and may regulate the extent that Tcfs and beta-catenin associate with chromatin (Hoffmans et al. 2005; Fiedler et al. 2008; Mieszczynek et al. 2008). Also, beta-catenin has also been implicated in establishing poised chromatin architecture prior to major zygotic gene activation. Evidence in *Xenopus* suggests that beta-catenin recruits Histone H3 Arginine 8 Methyltransferase (Prmt2; Blythe et al. 2010) to modify chromatin at target loci prior to the onset of target gene expression. Thus, Wnt target genes are regulated both by direct transcriptional activation following beta-catenin recruitment and by beta-catenin-regulated changes to chromatin, modes of regulation that may be temporally uncoupled. It is unclear, however to what extent the Pygo-regulated mechanisms and Prmt2 chromatin modifications are interrelated or instead exhibit overlapping or redundant regulation of Wnt targets genes.

### 6.3.1.3 Wnt/Planar Cell Polarity (PCP) Signaling

The first studies on Wnt signaling focused on the regulation of beta-catenin in development and cancer. Subsequent work found additional roles for a subset of Wnt ligands other components in controlling cell movements during axis organization

and/or antagonism of the Wnt/beta-catenin pathway (Rauch et al. 1997; Rothbächer et al. 2000; Wallingford et al. 2000; Veeman et al. 2003a) (Fig. 6.7). In vertebrates, these beta-catenin-independent Wnt pathways (often referred to, malapropos, as the “noncanonical” Wnt pathways) were shown to act through conserved *Drosophila* planar cell polarity (PCP) homologs and/or through release of intracellular calcium (Wnt/PCP and Wnt/Calcium pathways). Although these different pathway designations are convenient conventions, there is likely a large degree of overlap and interaction among them in vivo, particularly in the case of the Wnt/PCP and Wnt/Calcium pathways, and the ultimate outcome of signaling is likely dependent on the complement of Fzd receptors and coreceptors present on a given cell. This important point was exemplified early on by experiments showing that Wnt5a, traditionally considered a beta-catenin-independent Wnt ligand, could induce second axes in *Xenopus* when co-expressed with its cognate receptor Frizzled 5 (He et al. 1997), and recently by studies demonstrating Wnt5a-mediated regulation of beta-catenin-dependent and -independent Wnt signaling in mammals (Mikels and Nusse 2006; van Amerongen et al. 2012).

PCP signaling in vertebrates involves a set of components largely homologous to those mediating planar cell polarity signaling during imaginal disc development in insects (Vinson and Adler 1987; Krasnow and Adler 1994). This core set of proteins controls asymmetric Fzd1 localization (Strutt 2001) independently of Wnt ligands (Lawrence et al. 2002) and have been characterized genetically and biochemically in *Drosophila* (reviewed in Maung and Jenny 2011; Jose Maria Carvajal-Gonzalez 2014). These proteins include Fzd, Dishevelled, Flamingo (Fmi, a seven transmembrane pass cadherin), Prickle (Pk, a LIM and PET domain protein), strabismus/Van Gogh (Stbm/Vang, a four transmembrane protein with a PDZ motif), and Diego (Dgo, an ankyrin repeat protein). Homologous proteins also control epithelial cell and tissue polarity in vertebrates, notably in the inner ear (reviewed in Veeman et al. 2003a; Bayly and Axelrod 2011). Additionally, vertebrate PCP proteins are critical for controlling cell shape and cell migration in mesenchymal-type cells. Cell intercalation and cell migration during vertebrate gastrulation and neurulation in particular are dependent on Wnt/PCP signaling (reviewed in Solnica-Krezel and Sepich 2012).

The mechanisms of signal transduction during Wnt/PCP signaling in vertebrates are more varied and less well characterized than those of the beta-catenin-dependent pathway. Activation of the Wnt/PCP pathway in vertebrates is dependent on certain Wnt-Fzd combinations and a different set of coreceptors instead of Lrp5/6, including Ryk (Kim et al. 2008; Yoshikawa et al. 2003), Ror2 (Schambony and Wedlich 2007; Gao et al. 2011), and various Glypican proteoglycans (Topczewski et al. 2001; Ohkawara et al. 2003). Additionally, the transmembrane protein encoding Protein tyrosine kinase 7 (Ptk7) has been characterized as a novel regulator of PCP signaling (Lu et al. 2004; Yen et al. 2009). The role of Ptk7 is unclear, but it may represent an additional Wnt coreceptor modulating beta-catenin inhibition and activation with PCP signaling (Peradziryi et al. 2011; Hayes et al. 2013; Bin-Nun et al. 2014; Linnemannstöns et al. 2014). Dvl involvement is also critical for beta-catenin-independent Wnt signaling, although different domains are important for each function by controlling protein complex assembly and subcellular localization (Axelrod

et al. 1998; Boutros and Mlodzik 1999; Rothbächer et al. 2000). The N-terminal DIX domain (Dishevelled, Axin) is critical for beta-catenin regulation whereas the C-terminal DEP (Dvl, Egl-10, Pleckstrin) domain regulates PCP and calcium signaling. In *Drosophila*, Dvl associates with Fzd1 and localizes to the distal cell margin. This complex inhibits the distal accumulation of Vang/Pk complexes, which are restricted proximally.

In vertebrates, similar complexes are implicated but the assembly and asymmetry of these is less understood. Fzd/Dvl association likely occurs following GPCR activation, and Dvl and Fzds accumulate in asymmetric puncta in cells in various vertebrate tissues undergoing PCP. In the zebrafish gastrula, Dvl-GFP is localized to the posterior membrane of cells whereas injected *Drosophila* Prickle-GFP localizes to the opposite, anterior edge (Ciruna et al. 2006; Yin et al. 2008). Additionally, in the mouse posterior notochord/node, Prickle2 and Vangl1 colocalize at the anterior edge of cells (Antic et al. 2010) and Dvl-GFP localizes posteriorly (Hashimoto et al. 2010). However, in other tissues such as the cochlea, Vangl2 and Fzd3 colocalize (Wang and Nathans 2007), but Prickle2 and Fzd6 localize to opposite sides (Deans et al. 2007). Celsr1 (a vertebrate Fmi homolog; cadherin, EGF LAG seven-pass G-type receptor 1) may also play a role in recruiting Dvl/Fz complexes to adherens junctions in the neural plate and mediating subsequent signaling (Nishimura et al. 2012). Thus in vertebrates, the roles of the different core PCP components may have diverged following gene duplication and the acquisition of Wnt ligand dependence and may have taken on tissue- or cell type-specific roles.

Dvl recruitment in the context of Wnt/PCP signaling is implicated in the control of cytoskeletal dynamics through the activation of small GTPases. Dvl can recruit the Formin-related Daam1 protein to activate Rho in an Wnt-dependent manner and regulate actin dynamics (Habas et al. 2001). Additionally, Rho activation can lead to Rho kinase (Rok2) activation to control cell shape (Marlow et al. 2002; Tahinci and Symes 2003). In a separate and parallel pathway, Dvl can directly activate Rac downstream of Wnt, leading the stimulation of filopodial extensions and Mapk8 (Jun N-terminal kinase, JNK) activation (Habas et al. 2003; Tahinci and Symes 2003). The coordinate activity of Rho and Rac, and potentially other small GTPases, is required for cell intercalation and convergent extension morphogenesis in many developing tissues.

#### 6.3.1.4 Wnt/Calcium Release Signaling

Certain Wnt-Fzd combinations can stimulate the release of intracellular calcium stores (reviewed in Veeman et al. 2003a; Kohn and Moon 2005) and signal independently of beta-catenin. The regulation of this pathway also begins with Fzd-mediated heterotrimeric G protein activation and involves well-characterized GPCR responses, namely phosphoinositide turnover (Slusarski et al. 1997b), activation of cGMP-phosphodiesterase (Ahumada et al. 2002), as well as Calmodulin-dependent protein kinase 2 (Camk2) and Protein kinase C (Prkca) activation (Sheldahl et al. 1999; Kuhl et al. 2000). Many of the same coreceptors involved in Wnt/PCP



signaling are also critical for the Wnt/calcium pathway, suggesting that these pathways overlap considerably (Fig. 6.7). In line with this idea, overexpression of Dvl can initiate calcium flux and activate Camk2 and Prkca in fish and frog embryos (Sheldahl et al. 2003). Similarly, overexpression of Prickle1 indirectly regulates calcium dynamics (Veeman et al. 2003b). Also, recruitment of Dvl to the membrane during PCP signaling requires a calcium-regulated PKC isoform, Prkcd (Kinoshita et al. 2003). Wnt/PCP and Wnt/Calcium are likely to be tightly integrated, owing to shared components and shared roles in regulating morphogenesis during gastrulation and beta-catenin antagonism.

Evidence suggests that Wnt/Calcium signaling is essential for inhibiting beta-catenin activation during axis formation. Loss of maternal Wnt5b in zebrafish eliminates calcium flux in the blastula and triggers ectopic beta-catenin activity, resulting in dorsalized embryos (Westfall et al. 2003). This effect was partially rescued by Camk2, suggesting that calcium-mediated activation of this pathway is sufficient to suppress beta-catenin activity. Wnt/Calcium is also implicated in activating Nemo-like kinase (Nlk) (Ishitani et al. 1999, 2003; Meneghini et al. 1999) and Nfat nuclear translocation (Saneyoshi et al. 2002) to antagonize beta-catenin activity.

### 6.3.1.5 Wnt Secretion and Extracellular Regulation

Wnts are secreted and are modified by glycosylation (Brown et al. 1987; Papkoff et al. 1987) and lipidation (Willert et al. 2003). Efficient secretion of Wnts requires glycosylation and palmitoleoylation, the latter of which is mediated by the Porcupine (Porcn) family of acyl transferases (van den Heuvel et al. 1993; Kadowaki et al. 1996; Hofmann 2000; Tanaka et al. 2000). Tyrosine sulfation has also been observed and may be necessary for activity in some cases (Cha et al. 2009). Wnt secretion also requires trafficking of Wnt from the Golgi apparatus to the plasma membrane by the Wntless Wnt ligand secretion mediator (Wls; *alias* Evi/Gpr177/Wingful) as well as efficient recycling of Wls through the endosome-retromer system (Bartscherer et al. 2006; Coudreuse 2006). Interestingly, *Wls* is a direct Wnt/beta-catenin target gene in mouse and is required for extracellular Wnt signaling during mouse axis formation (Fu et al. 2009), indicating that Wnt activity potentiates its own signaling. Additional evidence suggests Wnt proteins may also be packaged into lipoprotein particles and/or exosome vesicles (Panáková et al. 2005; Gross et al. 2012). Wnts can act as both long- and short-range signaling molecules in the extracellular space, acting as developmental morphogens (Zecca et al. 1996). Wnt signaling gradients can also interact with those of a Wnt antagonist, Dkk1, to establish hair follicle spacing through a Turing-like reaction-diffusion mechanism (Sick et al. 2006), illustrating one of the complex ways this pathway can be used to establish tissue patterns in development.

Wnt signaling can be tightly regulated in the extracellular space by a host of different Wnt antagonists. Many of these proteins belong to large protein families and have redundant and tissue-specific functions throughout development (Cruciat and Niehrs 2013). The main secreted Wnt antagonists involved in axial patterning are the

Secreted Frizzled-related proteins (Sfrps), which bind directly to Wnts and antagonize different Wnt ligands, and Dickkopf1 (Dkk1), which acts at the level of the Wnt/Lrp6 receptor complex. In addition, the secreted Notum pectinacetyltransferase homolog was identified as a Wnt antagonist in *Drosophila* (Giraldez et al. 2002; Gerlitz and Basler 2002) and is thought to act by promoting membrane shedding of Glypican Wnt coreceptors (Kreuger et al. 2004). Recent data from flies and vertebrates also suggests that Notum acts as a Wnt deacylase, cleaving the Wnt palmitoleate moiety, resulting in Wnt ligand oxidation and inactivation (Kakugawa et al. 2015; Zhang et al. 2015). Notum is conserved and is involved in feedback regulation of Wnt signaling body axis patterning in Planaria (Petersen and Reddien 2011), and recent data suggest a role in dorsoventral neural tube patterning in zebrafish (Flowers et al. 2012).

Transmembrane antagonists have recently been identified as well. Those with roles in axis formation include the leucine-rich repeat protein Trogoblast glycoprotein (Tbgb/Waif1) and Tiki1 (Trabd2a). Tbgb is thought to act as a feedback Wnt inhibitor, acting in Wnt-receiving cells to alter Lrp6 subcellular localization (Kagermeier-Schenk et al. 2011). Trabd2a/Tiki1 is a transmembrane metalloproteinase enriched in the organizer that can cleave a subset of Wnt ligands, causing their abnormal oxidation and oligomerization and reduced receptor binding (Zhang et al. 2012).

### 6.3.2 *Wnt/Beta-Catenin Signaling in Early Axis Formation*

The central role of Wnt/beta-catenin signaling in axis formation was initially demonstrated largely through simple overexpression experiments in *Xenopus* and zebrafish embryos. The first of these was the induction of axis duplications in *Xenopus* by injected mouse *Wnt1* mRNA (McMahon and Moon 1989). *Xenopus wnt8a* (*Xwnt-8*; Christian et al. 1991; Sokol et al. 1991, Smith and Harland 1991), and several other Wnt ligands (Wolda et al. 1993; Du et al. 1995; Kelly et al. 1995a) can also induce secondary axes and rescue UV ventralization. Importantly, beta-catenin also exhibits axis inducing activity (Funayama et al. 1995; Guger and Gumbiner 1995), and both Wnt and beta-catenin can induce axial structures non-cell autonomously when expressed in vegetal blastomeres, suggesting that this activity acts analogously to a Nieuwkoop center (Smith and Harland 1991; Guger and Gumbiner 1995).

Interestingly, later overexpression of *wnt8a* during gastrulation (from injected plasmid DNA as opposed to mRNA) causes a loss of anterior structures, indicating roles for Wnts in patterning of the axis as well as its induction (Christian and Moon 1993). Other Wnts, including Wnts 4, 5a and 11b, do not elicit axis duplications but disrupt gastrulation movements and cell adhesion when overexpressed (Moon et al. 1993; Ku and Melton 1993; Du et al. 1995). Additionally, these Wnts can be antagonistic to the axis-inducing Wnts in some cases (Torres et al. 1996). The same ligands were also shown to trigger intracellular calcium release when expressed in the early zebrafish embryo (Slusarski et al. 1997a, b).

Loss-of-function experiments have established that Wnt/beta-catenin signaling is essential for axis formation in vertebrates. The first evidence for this came from

antisense oligonucleotide mRNA depletion of maternal *ctnnb1* mRNA in *Xenopus* oocytes, leading to embryos lacking axial structures and dorsal-specific gene expression (Heasman et al. 1994). Additionally, in Nieuwkoop conjugate experiments, late blastula vegetal masses from *ctnnb1*-depleted embryos fail to induce dorsal mesoderm in animal caps, suggesting that maternal Wnt/beta-catenin is essential for the generation of the Nieuwkoop signal and acts upstream of other axis-inducing molecules (Wylie et al. 1996). Analysis of heterochronic Nieuwkoop conjugates pre- and post-midblastula transition also showed that dorsal and general mesoderm induction are primarily zygotic events (Wylie et al. 1996). Beta-catenin is present in dorsal nuclei prior to major zygotic genome activation in the *Xenopus* morula and blastula as well as in the zebrafish dYSL and dorsal marginal blastomeres (Schneider et al. 1996; Jesuthasan and Stähle 1997; Kelly et al. 2000; Dougan et al. 2003) demonstrating that Wnt/beta-catenin signaling is active in the relevant region of the embryo.

In addition to these data, genetic studies in zebrafish identified a requirement for a maternally expressed beta-catenin in normal axis formation (*ctnnb2/ichabod*; Kelly et al. 2000). Furthermore, in the mouse, genetic deletion of *Ctnnb1* results in embryos lacking axial structures and anteroposterior polarity, resulting from lack of all mesoderm and failure to form the anterior visceral endoderm (AVE) (Haegel et al. 1995; Huelsken et al. 2000; Morkel et al. 2003). Mouse beta-catenin is predominantly required in the epiblast, as shown in chimeric embryo experiments (Huelsken et al. 2000) and is not required maternally (De Vries et al. 2004), reflecting a different mode of activation in mammals. The main Wnt ligand expressed at this time, *Wnt3*, is primarily expressed in the posterior epiblast and is required for embryonic axis and mesoderm formation (Liu et al. 1999b). Interestingly, the formation of the AVE is normal in *Wnt3* null mice, indicating a differential role for beta-catenin in the development of this tissue. A Wnt ligand-independent role for beta-catenin in anteroposterior patterning has been proposed (Morkel et al. 2003), possibly through regulation of *TdGF1* (*teratocarcinoma-derived growth factor 1*, *alias Cripto/fr11*) expression and subsequent effects on Nodal activity (see Sect. 6.5). Although genetic manipulations are less tractable in the chicken, studies using extracellular Wnt inhibitors suggested that Wnt signaling is required for experimental axis induction (Skromne and Stern 2001).

### 6.3.3 *Asymmetric Activation of Wnt/Beta-Catenin Signaling in Early Amphibian and Fish Embryos*

#### 6.3.3.1 *Xenopus*

Despite the well-documented roles for beta-catenin in axis formation in *Xenopus*, and more recently in zebrafish, it remains relatively unclear how Wnt/beta-catenin signaling is initiated in early embryos, as well as the extent that these activating mechanisms are conserved. Cytoplasmic transplantation studies in *Xenopus* identified the presence of a cytoplasmic, transplantable dorsalizing activity in the vegetal

cortical region (Darras et al. 1997; Marikawa et al. 1997; Marikawa and Elinson 1999). By correlating its activity with various axis-inducing molecules, this vegetal cortical cytoplasm was found to mimic intracellular activation of the Wnt/beta-catenin signaling pathway (Marikawa and Elinson 1999). Curiously, UV-irradiation experiments in *Xenopus* oocytes indicated that this cytoplasm showed cell cycle-dependent sensitivity to UV Irradiation of the egg disrupts microtubule assembly and cortical rotation, although the activity of the vegetal cortical cytoplasm itself is not affected. By contrast, UV-irradiation of full-grown oocytes effectively does eliminate the dorsalizing ability of vegetal cytoplasm and ventralizes embryos (Holwill et al. 1987; Elinson and Pasceri 1989). Eggs irradiated as oocytes undergo normal cortical rotation and are not rescued by tipping (Elinson and Pasceri 1989), suggesting that a critical component of axis induction is absent.

The target of UV irradiation in the oocyte is not known, but either its action is completed by oocyte maturation or it is subsequently sequestered and no longer susceptible to irradiation. These features may be useful in identifying potential candidate molecules. In apparent support of a direct cytoplasmic beta-catenin activation model, particles of exogenous Wnt-activating proteins, Dvl2-GFP (Miller et al. 1999) and Frat1-GFP (Weaver et al. 2003) were shown to undergo dorsal translocation during cortical rotation, suggesting that the dorsalizing activity might be composed of beta-catenin stabilizing agents. These molecules might then directly stabilize beta-catenin, or act by sensitizing dorsal cells to Wnt signals. Further indications of potential Wnt ligand-independent dorsalizing mechanisms came from observations that overexpression of secreted Wnt antagonists were unable to suppress endogenous axis formation (Hoppler et al. 1996; Leyns et al. 1997; Wang et al. 1997).

More recent studies in *Xenopus* suggest a more typical Wnt signaling model, with maternal Wnt11b acting to induce beta-catenin activity. Maternal *wnt11b* is localized to the vegetal cortex during oogenesis via a mitochondrial cloud-dependent pathway (see Chap. 8) and was initially considered a prime candidate for the vegetal dorsally activity, based on its activity in UV-rescue experiments. In light of the evidence favoring the direct activation model (see above), and other experiments showing that Wnt11b can regulate beta-catenin-independent signaling, a role for *wnt11b* in axis formation was later discounted. However, a reinvestigation using antisense oligo mediated maternal mRNA depletion showed that maternal *wnt11b* is indeed required for axis formation (Tao et al. 2005).

Wnt11b is thought to act in concert with uniformly expressed Wnt5a (Cha et al. 2008), forming extracellular complexes with each other and with other proteins, including heparin sulfate proteoglycans and the Nodal coreceptor Tdgf1 (Tao et al. 2005; Cha et al. 2008, 2009). The activity of this Wnt complex can be antagonized by maternal Dkk1, suggesting a model in which cortical rotation tips the balance of Wnt activity to overcome generalized Wnt antagonism (Cha et al. 2008). The mechanism of Wnt11b enrichment dorsally following cortical rotation is unclear, as both enrichment of total *wnt11b* RNA (Tao et al. 2005) or enhanced polyadenylation (Schroeder et al. 1999) have been proposed. A similar mechanism, albeit with different Wnts and antagonists, has been proposed in zebrafish (see below). However, Wnt11b can also regulate beta-catenin levels in an autocrine fashion in fully grown


oocytes (Kofron et al. 2007), providing the possibility that Wnt activation may wholly or partially occur before fertilization. Possible mechanisms for Wnt activation in *Xenopus* are shown Fig. 6.8 (top panels).

Although there is good evidence for secreted Wnt11b activity, it is unclear when Wnt11b is required. Also, this role for Wnt11b has not been reconciled with the cytoplasmic activation model. One potential unifying model has been proposed; that ongoing Wnt signaling in the oocyte generates activated Lrp6 signaling endosomes that are transported dorsally (Dobrowolski and De Robertis 2012). However, it is not known to what extent these Lrp6-containing endosomes are formed in the oocyte. Lrp6 is phosphorylated in eggs but becomes dephosphorylated following egg activation (Davidson et al. 2009), suggesting that stable signaling complexes may actually be inactivated prior to cortical rotation. Furthermore, analyses of the relative activity of vegetal cortical cytoplasm suggests it acts at the level of the destruction complex and does not mimic the activity of activated receptors (Marikawa and Elinson 1999).

### 6.3.3.2 Zebrafish

The regulation of Wnt signaling in zebrafish also has intriguing parallels and differences to the situation in *Xenopus* (Fig. 6.8, bottom panels). Maternal/zygotic mutants of *wnt11* form the axis normally in zebrafish (Heisenberg et al. 2000), suggesting that fish use other mechanisms or other Wnts. Maternal *wnt8a* has been proposed to act as a dorsal determinant in zebrafish (Kelly et al. 1995a; Lu et al. 2011). *Wnt8a* is the only vegetally localized *wnt* transcript in the yolk cell, and is shifted asymmetrically following cortical rotation. Vegetal localization also occurs through a mitochondrial cloud dependent mechanism, similar to frog *wnt11b* (Lu et al. 2011). In the fish however, injection of a dominant-negative Wnt8a construct was able to reduce the expression of *chrd* and *dharma*, the latter being a direct Wnt target gene. Full-length Wnt8a also rescues to some extent embryos ventralized by nocodazole (Lu et al. 2011). Also, depletion of Sfrp1 or Frzb hyperdorsalizes embryos, indicating dorsal enrichment of a maternal Wnt, Wnt8a in the case of fish, may be a trigger to overcome generalized Wnt antagonism in the embryo. While this and the similar mechanism proposed to act in frog are intriguing (Cha et al. 2008), it remains to be seen whether relatively small changes in Wnt levels involved are responsible for dorsalization.

Evidence in fish also suggests that other mechanisms of Wnt antagonism are critical in suppressing beta-catenin activation. Notably, maternal/zygotic mutants for *wnt5b* are hyperdorsalized, owing to defective Wnt/calcium signaling and failure to repress Wnt/beta-catenin signaling (Westfall et al. 2003). A similar but not well-characterized pathway may exist in frogs (Saneyoshi et al. 2002). In *Xenopus*, the maternal role of Wnt5b has not been assessed, however Wnt5a is proposed to activate instead of repress Wnt/beta-catenin signaling in frogs (Cha et al. 2008). Axis formation is largely normal however in *wnt5a*, *wnt5b* double mutant mice (Agalliu et al. 2009; spinal cord formation), suggesting that the roles of Wnt5 paralogues may be



**Fig. 6.8** Models for Wnt/beta-catenin activation in *Xenopus* and zebrafish. **(a)** During cortical rotation in *Xenopus* (top) and zebrafish (bottom), beta-catenin stabilizing dorsalizing activity is transported into the equatorial region of the embryo by microtubule-mediated rotation of the cortex and through transport along microtubule arrays. Candidates for this activity include *wnt11b* and Lrp6/Dvl particles in *Xenopus* and *wnt8a* in zebrafish. **(b)** By the cleavage stages (16–128-cell stage), beta-catenin becomes activated and enriched in dorsal vegetal and marginal nuclei until MBT. In *Xenopus*, priming of Wnt target genes occurs through dimethylation of Histone3 at arginine 8 (H3R8). In zebrafish, beta-catenin accumulates in dorsal marginal and dYSL nuclei, and is antagonized by multiple antagonists and calcium signaling mediators. **(c)** During the peri-MBT stages, beta-catenin activates direct Wnt targets and cooperates with maternal T-domain proteins (Vegt, Eomes) to activate nodal initially on the dorsal side. The combination of nodal and BMP antagonism induced by beta-catenin induce the formation of the organizer (*gsc, chrd*)

species-specific. Recent data suggest that calcium transients downstream of the chemokine receptor *Ccr7* GPCR signaling are also involved in suppressing beta-catenin activity (Wu et al. 2012b). The prominence of calcium regulation mechanisms suggests tight control over beta-catenin stabilization in the zebrafish blastula.

### 6.3.4 Beta-Catenin Activity Dorsalizes the Primary Germ Layers

Beta-catenin stabilization in the blastula following cortical rotation is the central mechanism for establishing early dorsal fates across all three primary germ layers (Fig. 6.8). In the vegetal prospective endodermal cells, beta-catenin is critical for the dynamic regulation of Nodal expression and signaling, which is directly involved in mesendoderm induction and patterning and likely constitutes what is referred to as the Nieuwkoop center (see Chap. 7). In *Xenopus*, the typical *nodal* homolog genes (e.g., *nodal* homologs 1, 2, 4, 5, and 6) are expressed in a temporally and spatially

graded fashion in the late blastula/early gastrula. These genes depend on vegetally localized maternal *vegt* activity, showing high dorsovegetal expression at the onset of gastrulation, followed by a shift ventrolaterally by late gastrulation, mirrored by Smad2 activity (Agius et al. 2000; Faure et al. 2000; Lee et al. 2001). Maternal beta-catenin directly patterns this activity first by contributing to early *nodal* paralogue expression (*nodal5*, *nodal6*) in dorsovegetal blastomeres prior to and immediately after the onset of zygotic transcription at the mid-blastula transition (MBT; see Chap. 9) (Takahashi et al. 2000; Rex et al. 2002; Xanthos et al. 2002; Hilton et al. 2003; Blythe et al. 2010). This role of beta-catenin may be related to its function in recruiting Prmt2 to prime gene expression during the cleavage stages (Blythe et al. 2010). Beta-catenin also synergizes with Nodal activity to generate higher *nodal1* expression dorsally in the early gastrula, although mid-late gastrula expression becomes uniform and is independent of beta-catenin (Agius et al. 2000; Lee et al. 2001).

In addition to overlapping with Vegt, beta-catenin might also functionally interact with Gdf1 (*alias* Vg1), which is encoded by another maternally localized mRNA (Rebagliati et al. 1985; Melton 1987; Weeks and Melton 1987). Gdf1 is a Nodal-related Tgfb family ligand and activates Smad2 signaling through Nodal receptors/coreceptors. Wnt/beta-catenin activity synergizes with *gdf1* overexpression to induce axial structures in *Xenopus* (Cui et al. 1996). Maternally depleted *gdf1* embryos lack anterior structures and are deficient in the expression of certain organizer markers, including secreted antagonists *nog*, *chrd*, *dkk1*, and *cer* (Birsoy et al. 2006). Early Smad2 activation is also compromised, indicating that Gdf1 contributes to overall Nodal signaling on the dorsal side of the late blastula (Birsoy et al. 2006). Thus, beta-catenin activity can enrich Nodal expression and activity dorsally at multiple regulatory levels.

This temporal control of Nodal activity is functionally significant. In Nieuwkoop conjugate experiments, late blastula vegetal explants from *ctnmb1*-depleted *Xenopus* embryos fail to induce dorsal fates in early equatorial or animal cap tissue (Xanthos et al. 2002; Wylie et al. 1996). However, if *ctnmb1*-depleted vegetal explants from late gastrula embryos are used, which now express *nodal* genes, dorsal mesoderm is induced (Xanthos et al. 2002). Additionally, pre-MBT Nodal activity regulated by beta-catenin is essential for normal axis formation, possibly for perpetuating *nodal* expression through autoinduction (Skirkanich et al. 2011). Beta-catenin also likely contributes to Nodal ongoing autoregulation, both directly and indirectly. Dorsal beta-catenin activity is required to repress *microRNAs 15* and *16* (*miR15/16*) through an unknown mechanism (Martello et al. 2007). These miRs target and downregulate an essential component of the Nodal receptor complex, *activin A receptor, type IIA* (*acrv2a*) (Martello et al. 2007), which contributes to the early dorsal bias in Nodal activity.

In zebrafish, maternal beta-catenin signaling similarly regulates early *nodal* homolog gene expression. Two paralogues are expressed in the early dYSL (*nodal1/squint* and *nodal2/cyclops*); *nodal1* is likely a direct beta-catenin target (Kelly et al. 2000) and *nodal2* is autoregulated by Nodal signaling itself. Genetic and other loss-of-function experiments support a role for the early Nieuwkoop center *nodal* genes in axis formation. Single and double mutants for *nodal1*, *nodal2* (*squint*, *cyclops*) exhibit axial patterning and mesendoderm defects (Rebagliati et al. 1998; Feldman

et al. 1998; Sampath et al. 1998; Erter et al. 1998), and these abnormalities can be rescued by YSL-specific expression of Nodal ligands. Additionally, *nodal1* is maternally expressed in zebrafish and its mRNA localized to the presumptive dorsal side at the 4-cell stage, through an uncharacterized mechanism involving microtubules (Gore et al. 2005). The role of this maternal transcript is not fully clear; it may lead to enhanced Nodal activity dorsally, although there are hints that the 3'UTR may help promote dorsal nuclear beta-catenin accumulation by acting as a noncoding RNA (Lim et al. 2012).

In addition to stimulating higher levels of Nodal activity, dorsal beta-catenin directly initiates a set of gene regulatory interactions that integrate with Nodal signaling in the prospective mesendoderm (Fig. 6.8). In *Xenopus*, Wnt/beta-catenin signaling directly activates the paired homeobox paralogues *siamois1* (*sia1*) and *siamois2* (*twin/sia2*) in dorsovegetal cells in the blastula (Lemaire et al. 1995; Brannon and Kimelman 1996; Brannon et al. 1997). *Sia1/2* integrate with Nodal/Tgfb signaling in activating additional genes in the anterior endoderm and organizer, including *cerberus*, *hhex*, and *gooseoid*, whose promoters are bound by *Sia1* (Ishibashi et al. 2008; Rankin et al. 2011; Sudou et al. 2012). Maternal beta-catenin signaling is also required to repress early *bmp4* expression dorsally in *Xenopus* (Baker et al. 1999), and this is also mediated through *Sia1/2*. These proteins are required to repress Bone Morphogenetic Protein (BMP) expression and activity dorsally, which is essential for dorsal mesoderm formation (see Sect. 6.4.1). *Sia1/2* activate the expression of several BMP antagonists, including *noggin* (*nog*) and *chordin* (*chrd*) (Ishibashi et al. 2008), and also indirectly repress *bmp4* expression, through an unknown intermediate. *Nog* and *chrd* expression are maintained by Nodal signaling, but initial expression depends only on maternal beta-catenin (Wessely et al. 2001). Although *Sia1/2* are predominantly expressed in mesendodermal precursors, their expression can extend into some equatorial precursors that escape mesoderm induction (Kuroda et al. 2004), where they specify early neural fate and the prospective anterior neuroectoderm (Ishibashi et al. 2008; Klein and Moody 2015).

Similarly, in zebrafish maternal beta-catenin activates expression of *dharma* in dorsal marginal blastomeres and dYSL. *Dharma* is a paired homeobox gene somewhat functionally analogous to *sia1*, but unrelated by descent (Fekany et al. 1999; Kelly et al. 2000). *Dharma* has numerous roles in dorsoventral patterning, including direct repression of *bmp2b* (Koos and Ho 1999; Leung et al. 2003a). Interestingly, both *sia1* and *dharma* are expressed prior to major zygotic gene activation in frogs and fish (Yang et al. 2002; Leung et al. 2003b; Blythe et al. 2010) suggesting very early and direct roles for these proteins in dividing the embryo into prospective BMP-expressing or -absent territories. Additionally in fish, the accumulation of nuclear beta-catenin occurs concomitantly with a change in cell division patterns relative to the body axis as well as slower cell division in the presumptive dorsal shield region, but a causal relationship between these events has not been established (Keller et al. 2008).

Although the activation of Nodal and *Sia1/2* appears to mediate many of the functions of maternal beta-catenin signaling, other less characterized pathways are likely required as well. One other beta-catenin target gene, *nodal homolog 3.1*



(*nodal3.1*; Smith et al. 1995), likely functions in controlling morphogenesis during organizer formation. Nodal 3.1 is an atypical member of the Nodal protein family present only in anuran amphibians (Smith et al. 1995). This protein has BMP antagonist activity (Hansen et al. 1997) and, during gastrulation, is restricted to the superficial epithelial layer of the organizer (Glinka et al. 1996), a region with discrete morphogenetic regulatory ability (Shih and Keller 1992a). Loss-of-function experiments suggest that Nodal3.1 regulates convergent extension, acting as a Fibroblast growth factor (FGF) receptor ligand (Yokota et al. 2003), although this mechanism is not well understood.


In summary, beta-catenin function is critical at many different regulatory levels to specify dorsal fate across all prospective germ layers in the early blastula in *Xenopus* and zebrafish. These general functions would include generating dorsal-inducing signals, mediating competence to respond to those signals, reducing the ability of dorsal cells to respond to ventralizing BMP signals, and regulating dorsal morphogenesis. In this context, the strong expression of *nodals* in the dorsovegetal blastomeres is likely responsible for the Nieuwkoop center phenomenon, although beta-catenin activity is required in prospective organizer mesoderm and ectoderm as well. In amniotes, however, axis formation is also governed by the dynamic regulation of Nodal signaling in concert with Wnt/beta-catenin activity. In these cases, the initial establishment of restricted Nodal signaling occurs without early maternal Wnt signaling and the role of a Nieuwkoop center analog in inducing the organizer is less clear.

### 6.3.5 *Wnt/Beta-Catenin and Nodal Signaling During Axis Formation in Amniotes*

#### 6.3.5.1 Chick

The role of concerted Wnt and Tgfb signaling in the dynamic regulation of Nodal expression and activity is conserved during amniote axis formation. In contrast however to the case in fish and frogs, axis formation in birds and mammals is driven by spatially restricted Nodal activity overlapping with more generalized Wnt/beta-catenin signaling in the marginal zone of the epiblast (Fig. 6.9a). In the chicken, there is no evidence for localized Wnt expression in the egg or early cleavage stages. *Wnt8a* (alias, *CWnt8C*) is the predominant Wnt expressed prior to gastrulation and is enriched in the PMZ epiblast, but is also found in a decreasing gradient around the marginal zone from posterior-to-anterior (Hume and Dodd 1993; Skromne and Stern 2001). By contrast, the Nodal-related gene *Gdf1* is specific to the PMZ at the same stage (Shah et al. 1997; Skromne and Stern 2001) and represents one of the earliest asymmetrically expressed genes in the chick blastoderm (and the first to be discovered).

Grafting of *Gdf1*-expressing cells induces an ectopic axis anywhere in the marginal zone (Shah et al. 1997; Skromne and Stern 2001). Interestingly, ectopic expression of Wnt alone is not a robust axis inducer in chicken (Joubin and Stern 1999). *Gdf1* and *Wnt8a* can synergize in axis induction, and the axis-inducing abil-



**Fig. 6.9** Models for axis induction signaling in chick and mouse. **(a)** In the chick blastoderm (left panel, top/dorsal view, ~stage X, Eyal-Giladi and Kochav 1976) the outer marginal zone of the epiblast expresses *Wnt8a* in a posterior-to-anterior gradient (purple shading). In the PMZ, *Pitx2* (yellow) activates *Gdf1* expression (green). Subsequently (middle panel), the newly formed hypoblast (below the plane of the page) begins anterior migration and *Gdf1* + *Wnt8a* cooperate to induce *Lef1* in the PMZ and *Nodal* in the adjacent epiblast. *Gata2* is expressed in the anterior marginal zone and antagonizes *Gdf1* long-range. *Nodal* (magenta) is antagonized by *Cerberus* (*Cer*), which is expressed in the hypoblast. By the initial primitive streak stage (right panel, stage 2+, Hamburger and Hamilton 1951), the anterior migration of the hypoblast and migration of the endoblast beneath the posterior epiblast removes the inhibition of *Nodal* and allows feed-forward signaling leading to primitive streak formation. The same signaling molecules are expressed in the primitive streak and induce organizer genes in Hensen's node (dotted circle, *Gsc*, *Chrd*) at the anterior tip of the streak. **(b)** In the mouse, the earliest asymmetries are the expression of *Lefty1* and *Cerl* (red-brown) at the tip of the postimplantation AVE (left panel, ~E5.0). These genes are regulated by *Nodal* (green) and *Tdgf1*, and *Tdgf1* is regulated by beta-catenin in the absence of secreted *Wnt* ligand activity (stop symbol). *Lefty1* and *Cerl* antagonize *Nodal* and feedback regulation drives AVE migration towards the proximal egg cylinder on one side (right panel, ~E5.5). *Nodal* activity is restricted to the posterior epiblast and is responsible for *Wnt3* expression (blue), which in turn maintains *Nodal*. These signals cooperate to induce the primitive streak, which induces Hensen's node toward the distal tip later in gastrulation. *a* anterior, *p* posterior, *ExE* extraembryonic ectoderm, *VE* visceral endoderm, *ParE* parietal endoderm

ity of *Gdf1* is blocked by co-expression with *Wnt* antagonists (Skromne and Stern 2001). These *Gdf1* and *Wnt8a* signals cooperate to induce *Lef1*, a *Wnt* signaling transcriptional activator, in the PMZ (Skromne and Stern 2001) coincident with the activation of *Nodal* in the adjacent area pellucida epiblast (Skromne and Stern 2002). *Nodal* itself is required for primitive streak and axis formation (Bertocchini and Stern 2002). Based on shared embryological and molecular characteristics, it has been speculated that the PMZ epiblast in chick represents the equivalent of the Nieuwkoop center (Bachvarova et al. 1998). However, in birds and likely in mammals

as well (see below), the middle regions of the primitive streak also express *Gdf1*, *Wnt8a*, and *Nodal* and may be more relevant for inducing the anterior streak and Hensen's node/organizer (Joubin and Stern 1999; Bertocchini and Stern 2002). The Nieuwkoop center in chicks may thus be considered a primitive streak-inducer, as opposed to its more traditional organizer-inducing amphibian role (also see Sect. 6.7). As compared to mammalian axis formation, which depends initially on Nodal asymmetry (see below), the chick may represent an intermediate condition, where Wnt signaling is widely active prior to gastrulation but the role of Nodal-related proteins in inducing the axis is becoming predominant.

The mechanisms controlling localized *Gdf1* expression in the PMZ remain largely unknown. In contrast to the strict requirement for maternal factors in fish and amphibians, there is only a presumptive early influence of maternal determinants in birds, and axis formation is highly self-organizing (i.e., regulative). Classic embryological studies showed that partitioning of the blastoderm results in axis initiation in each individual fragment (Lutz 1949; Spratt and Haas 1960; Callebaut and Van Nueten 1995), suggesting that axis determinants are not uniquely restricted to one region. In recent reinvestigations of this phenomenon, *Gdf1* expression reinitiates stochastically at the "new posterior" pole of these blastoderm explants and regulates formation of the new primitive streak (Bertocchini et al. 2004). The *Wnt8a* gradient presumably reforms around the new site of *Gdf1*, suggesting that *Wnt8a* is likely downstream or independent of *Gdf1*. There is no evidence to suggest whether this regulation is direct or indirect. Data also show that the transcription factor *Gata2* is expressed earlier than *Gdf1*, and in a roughly complementary anteriorly biased gradient (Bertocchini and Stern 2012). Both *Gata2* and *Gdf1* control each other's expression indirectly through a global signaling gradient, likely mediated by BMP signaling downstream of *Gata2*. Additionally, recent microarray screening and functional analyses identified the Pitx2 homeodomain protein as an essential upstream activator of *Gdf1* expression in the PMZ (Torlopp et al. 2014). This Pitx/*Gdf1*/*Nodal* regulatory relationship is deeply conserved in several well-known developmental processes, including sea urchin axis patterning, amphibian mesendoderm induction and left-right patterning (Torlopp et al. 2014 and references therein), indicating these genes form a robust module can be readily redeployed for novel functions. However, there are as yet no clues to any mechanisms that would potentially activate Pitx2 or connect this pathway to any maternal asymmetries. Interestingly, recent data suggest that chick embryos form a yolk syncytial layer (Nagai et al. 2015), similar to that of teleosts. This is likely the result of independent convergent evolution and it will be exciting to learn to what extent this structure might function similarly in axis induction.

### 6.3.5.2 Mouse

In contrast to the case in chick, in which the graded expression of *Wnt8a* in the marginal zone still plays some role in normal axis development, several lines of evidence in the mouse suggest that active Wnt/beta-catenin signaling is not part of the early axial polarization mechanism (Fig. 6.9b). Many Wnt genes are expressed in the preimplantation blastocyst (Kemp et al. 2005), although studies in Wnt reporter

mice initially suggested that Wnt signaling is not active in the blastocyst prior to implantation (Mohamed et al. 2004; Na et al. 2007). Recent studies with more sensitive reporter constructs have detected some beta-catenin-dependent activity in the ICM/epiblast of peri-implantation blastocysts (E3.0–4.5; Granier et al. 2011; ten Berge et al. 2011). Similarly, transient nuclear beta-catenin has been observed by immunolocalization at the same stage (Chazaud and Rossant 2006). However, the extent of activity is unclear, since overexpression of stabilized beta-catenin does not ectopically induce Wnt target genes before gastrulation (Kemler et al. 2004). Thus, although there are indications of early but non-polarized beta-catenin activity in the preimplantation mammalian embryo, the role of this signaling appears to be dispensable or easily compensated for, at least with regard to axis formation.

Correspondingly, mutants for the major Wnts, Wnt/beta-catenin coreceptors, as well as regulators of Wnt secretion and activity all undergo normal preimplantation development, including anteroposterior patterning in the AVE. These mutants also fail to form mesoderm and do not gastrulate. These Wnt/beta-catenin pathway mutants include (but are not limited to) *Wnt3* (Liu et al. 1999b), *Lrp5/6* (Kelly et al. 2004), *Mesdc1* (Hsieh et al. 2003), *Wntless* (Fu et al. 2009), and *Porcn* (Biechele et al. 2011, 2013; Barrott et al. 2011). Since beta-catenin mutants exhibit these same mesoderm defects but additionally fail to form the AVE, a Wnt ligand-independent role for beta-catenin in anteroposterior patterning has been proposed (Morkel et al. 2003), possibly mediated by activation of Tdgl expression and subsequent effects on Nodal activity (see Sect. 6.5.2). These results suggest possible similarities to the potential role of Wnt ligand independent signaling in *Xenopus* axis formation (see Sect. 6.3.3), but too little is known about either mechanism to draw meaningful comparisons. There are other agonist ligands of Wnt/beta-catenin signaling, such as Norrin and R-spondin (Cruciat and Niehrs 2013), but these also rely on Lrp5/6. Other potential mechanisms of regulating beta-catenin activity may exist, such as regulation through Hippo signaling (Varelas et al. 2010) or through Seven in absentia homologs (Siah) (Topol et al. 2003), although these have not been extensively investigated in the context of axis formation.

Rather than relying on polarized Wnt activity, the earliest asymmetries in the mouse blastocyst are the expression of Nodal antagonists *Cer1* and *Lefty1* in the primitive endoderm of the peri-implantation blastocyst (~E4.0; Torres-Padilla et al. 2007b). Expression of these genes likely depends on FoxH1-mediated Nodal signaling (Takaoka et al. 2006; Torres-Padilla et al. 2007b), initiating a cascade of both positive and negative feedback regulation of Nodal signaling in the embryo. *Cer1/Lefty1* asymmetry is likely to arise stochastically at this early stage, as the blastocyst contains a rather small number of cells and it is unlikely that the few *Cer1/Lefty1* expressing cells would arise precisely in the center. These cells are then thought to lead and propagate AVE migration toward the future anterior side, i.e., towards the side of initial asymmetry (Takaoka et al. 2011; Morris et al. 2012a), restricting Nodal activity to the prospective posterior. This aspect will be discussed further in Sect. 6.5. Additionally, *Nodal* is expressed at low levels in the ICM prior to implantation, but there is no evidence that Wnt/beta-catenin signaling activates this expression. Later in development, prior to and during gastrulation, *Nodal* expression is maintained by *Wnt3* signals in the posterior epiblast and in the forming primitive

streak (Brennan et al. 2001; Ben-Haim et al. 2006), indicating that Wnt regulation of *Nodal* remains conserved, but is deployed modularly in development.

Interestingly, although Wnt may not activate early *Nodal* expression in the blastocyst, data suggest that Tcf3 repression of *Nodal* is a conserved feature of vertebrate development. Deletion of mouse *Tcf3* (Merrill et al. 2004) results in the formation of secondary axes (Hensen's nodes and notochord) and in upregulation of *Nodal* target genes (e.g., *Foxa2*; Hoodless et al. 2001; Yamamoto et al. 2001) in the pregastrula embryo. And, *Nodal* is upregulated in *Tcf3* null embryonic stem (ES) cells (Cole et al. 2008). In fish and frogs, early-localized Wnt signals provide an initial signal to relieve repression by Tcf3 (Kim et al. 2000; Houston et al. 2002; Dorsky et al. 2003). Mammals must have evolved different mechanisms to overcome this repression in the absence of pervasive early Wnt activity. Recent data suggest that *Nodal* initiation and potentiation may be controlled by a distinct enhancer bound by pluripotency transcription factors Oct4 and Klf4, in addition to ongoing Smad2-dependent signaling (Papanayotou et al. 2014). Interestingly, *Tcf3* null ES cells also exhibit derepression of other genes, in addition to *Nodal*, that are co-regulated by pro-pluripotency factors (Cole et al. 2008), suggesting that in normal ES cells, these proteins may help overcome Tcf3-mediated repression in the absence of strong activation of Wnt/beta-catenin signaling.

Is there a mammalian Nieuwkoop center? In line with a lack of asymmetry in the mammalian egg, clonal analyses and transplant experiments performed in the pregastrula mouse embryo have not identified a region with the requisite Nieuwkoop properties; indeed transplanted epiblast cells typically change fate according to their new position (Lawson et al. 1991; Lawson and Pedersen 1992; Parameswaran and Tam 1995). Also, there is a high degree of cell mixing in the epiblast during normal development (Gardner and Cockcroft 1998), which would be inconsistent with a lineage-restricted Nieuwkoop center. However, there is overlap of *Nodal* and *Wnt3* in the posterior proximal epiblast and subsequently throughout the primitive streak (Tam et al. 2006) and high levels of *Nodal* signaling are required for organizer formation (Vincent et al. 2003). The early proximal epiblast may have some hallmarks of a Nieuwkoop center, although the region of highest and most complete axis-inducing activity is the anterior primitive streak itself, prior to Hensen's node formation (Kinder et al. 2001), similar to the case in chicken. Interestingly, these cells do not significantly contribute to the organizer/Hensen's node or notochord, instead making contributions to the anterior mesendoderm (Kinder et al. 2001). Thus, organizer induction in mammals may still require non-cell-autonomous induction by Wnt and *Nodal* signals, with the signals being generated from the nascent mesendoderm, as opposed to strictly extra-mesodermal Nieuwkoop center-like mechanisms.

## 6.4 The Organizer and Dorsoventral Patterning

Axis formation and patterning are largely determined by a discrete zone of midline mesoderm in the prospective dorsal region of the gastrula. The collective cell signaling and morphogenetic properties of these cells and their descendants during

normal development and under experimental conditions led to their designation as the “organization center” or “organizer” (*Organisationszentrum/Organisator*) (Spemann 1921; Spemann and Mangold 1924). During experiments to test the extent of cell fate determination in the gastrula, Spemann and Mangold showed that the dorsal (upper) lip of the newt gastrula could retain its fate when transplanted to the ventral side, typically forming a normally patterned “secondary embryo” that matched the axial organization of the host. By grafting dorsal lips from albino embryos into pigmented hosts (Fig. 6.10), they assessed the contribution of donor cells, and strikingly demonstrated that the secondary dorsal tissues including the neural tube were formed from induced host cells, with donor cells forming notochord and somites (Spemann 1921; Spemann and Mangold 1924). The term organizer was thus coined by Spemann to reflect the ability of the dorsal (upper) blastopore lip to direct the development of the characteristic embryonic structures in vertebrates (notochord, somites, dorsal neural tube).


Spemann’s characterization of the organizer was highly influential and rapidly led to the identification of homologous regions in other vertebrates: birds (chick and duck, Hunt 1929; Waddington 1930), mammals (chick into rabbit, Waddington 1934; rabbit into rabbit/chick, Waddington 1936, 1937; mouse into frog, Blum et al. 1992; and mouse into mouse, Beddington 1994), lamprey (Bytinski-Salz 1937; Yamada 1938), and teleost fish (minnow, perch, and trout, Oppenheimer 1934a, b; Luther 1935). Similar dorsalizing and neuralizing activities have also been found for the prospective notochordal cells of invertebrate cephalochordates (Tung et al. 1962), suggesting the organizer is a basal feature of the chordates.

In bird and mammalian embryos, organizer activity is limited to the anterior primitive streak, including the discrete anterior tip (Hensen’s node, “*Knoten*,” Viebahn 2001; Blum et al. 2007).<sup>2</sup> In teleosts, the organizer corresponded to a thickened area of the blastoderm on the dorsal side, termed the embryonic shield. The initial characterization of the organizer identified several embryological properties of the organizer that are central to its functions and are still studied in the context of axial development. These are: the tissue-autonomous involution and convergent extension of the dorsal lip and the differentiation into axial mesoderm (notochord), the induction and regionalization of the neural plate, and self-organization/developmental plasticity (i.e., regulation) (Spemann 1938).

As intriguing as these observations were, the organizer remained largely a phenomenon until the application of molecular biological methods to developmental biology in the 1980s–1990s. These were first used to characterize the neural inducing activity of the organizer. This aspect was the historical readout of organizer

---

<sup>2</sup>In this chapter, Hensen’s node is used to refer to the anterior tip of the primitive streak in all birds and mammals, and is considered equivalent to the dorsal lip/organizer. Often the mouse organizer is referred to as the “node” without the eponym. However, the node can also refer to the posterior notochord “node” involved in left-right patterning, which lacks organizer activity. This terminology can cause confusion since the latter structure is embryologically distinct from Hensen’s node. In human embryology, Hensen’s node is referred to as the primitive node/knot or Hensens’ knot (Gray 1918; Larsen et al. 2009).




**Fig. 6.10** The organizer experiment of Spemann and Mangold. (a) Diagrammatic model of Spemann and Mangold's dorsal lip transplantation from a lightly pigmented species (*light gray*) to the ventral region of a darker species (*dark gray*). (b) Image of a *Xenopus* tadpole following the successful grafting of an early gastrula dorsal lip, showing the endogenous axis (1° axis) and the induced partial axis (2° axis). The dark pigment in the head of the 2° axis is related to abnormal head development in the induced axis. (c) Diagram of a cross section through an embryo resulting from a dorsal lip transplant as in (a). The typical lineage contribution of the donor lip is lightly shaded reflecting the species origin, indicating contributions to the notochord, floor plate and medial somite (c is after Spemann and Mangold 1924). v ventral, d dorsal

function prior to the molecular era, owing to ease of identifying induced neural plates and nervous system tissue in sectioned material. Beginning with the observations that dissociation and delayed reaggregation of amphibian animal cap ectoderm could cause neural differentiation (Grunz and Tacke 1989, 1990; Saint-Jeannet et al. 1990; Godsave and Slack 1991), and the cloning of organizer-specific transcription factors (Cho et al. 1991; Dirksen and Jamrich 1992; Taira et al. 1992), a series of experiments from many labs identified antagonism of Bone Morphogenetic Protein (BMP) signaling activity as a key activity of the organizer in vertebrate axial development (Smith and Harland 1992; Sasai et al. 1994, 1995; Wilson and Hemmati-Brivanlou 1995; Zimmerman et al. 1996). The early history of the molecular characterization of the organizer and its impact on the fields of developmental biology and the evolution of development has been extensively reviewed

over the years by many of the key participants (Harland 1994, 2008; Elinson and Holowacz 1995; De Robertis and Sasai 1996; Harland and Gerhart 1997; Chang and Hemmati-Brivanlou 1998; De Robertis et al. 2000; De Robertis 2006, 2009; Harland and Grainger 2011).

### 6.4.1 The Differentiation of Axial Mesoderm and the Role of BMP Antagonism

It is clear from numerous studies that the formation of a ventral-to-dorsal BMP activity gradient through extracellular antagonism underlies the main functions of the organizer in dorsoventral axial patterning (Fig. 6.11a). BMPs are members of the Tgfb superfamily of secreted growth factors. These function as dimers and activate heteromeric cell surface serine/threonine kinase receptors, resulting in the phosphorylation of BMP-specific Smad transcription factors (Smad 1/5/8) (Chap. 7; reviewed in Little and Mullins 2006; Ozair et al. 2012). BMPs are expressed in non-organizer tissue, namely the ventrolateral regions in amphibians and posterior primitive streak of the epiblast in chick and mouse. These molecules function in a



**Fig. 6.11** Dorsoventral patterning of the gastrula. (a) Image of an early gastrula *Xenopus* embryo (left image) showing *gsc* mRNA expression in the organizer region (purple). This area expresses BMP antagonists Chrd, Fst, and Nog in a dorsoventral gradient (blue shading), and is complementary to a gradient of BMP signaling (red shading), resulting in a graded pattern of phospho-Smad1/5 (p-Smad1/5). On the right is a late gastrula showing continued ventroposterior patterning by BMP signaling, specifying in progressively later fashion (differing line thicknesses) the anterior neural crest (a.n.c.), posterior neural crest (p.n.c.) and epidermis (ep.), in addition to the underlying germ layers (not shown). (b) Simplified network model of secreted protein interactions acting in dorsoventral patterning. Chrd secreted by the organizer antagonizes Bmp activity, mediated by Bmp4/7 ventrally and Admp within the organizer. Bmp activity inhibits *chrd* expression. Tld/Bmp1 acts as a Chrd inhibitor via proteolysis; Szl is a Tld inhibitor, indirectly promoting Chrd activity in the extreme ventral region (dotted arrow). The critical reciprocal control interactions responsible for self-organization are indicated with blue lines; Bmp4/7 positively controls its own expression but inhibits *admp* expression. Bmps also upregulate *cv2/bmper*, a ventral Bmp antagonist, and inhibit *chrd* dorsally. Model after De Robertis (2009)



dose-dependent fashion to promote ventrolateral mesendoderm and epidermis, and can prevent neural development of dissociated animal caps and inhibit dorsal mesendoderm formation in explant assays (Jones et al. 1992; Dale et al. 1992; Fainsod et al. 1994; Suzuki et al. 1994; Graff et al. 1994; Wilson and Hemmati-Brivanlou 1995; Hammerschmidt et al. 1996b; Streit et al. 1998).

The first insights into the molecular biology of the organizer came from the isolation of conserved homeobox and forkhead transcription factors expressed in the organizer: *gooseoid* (*gsc*), *lim homeobox 1* (*lhx1*), and *forkhead box a4* (*foxa4*) (Cho et al. 1991; Taira et al. 1992; Dirksen and Jamrich 1992). Importantly, ectopic expression of *gsc* in ventral cells caused second axes with notochordal tissue, indicating that *gsc* was not merely a marker of, but a critical functional component of the organizer (Cho et al. 1991). Studies in other organisms subsequently found a deeply conserved role for these proteins in regulating axial development in both protostomes and deuterostomes and even to some extent in diploblasts (Martindale 2005).

Although it was clear that overexpression of these proteins, notably Gsc, could induce most aspects of organizer function and regulate cell–cell signaling during axis formation (Niehrs et al. 1993), the nature of these signals was not immediately apparent. In a series of seminal experiments, expression cloning and differential cDNA screening in *Xenopus* identified secreted molecules encoded by *noggin* (*nog*; Smith and Harland 1992; Lamb et al. 1993), *chordin* (*chrd*; Sasai et al. 1994), and *follistatin* (*fst*; Hemmati-Brivanlou et al. 1994) as potent dorsalizing and neuralizing molecules specifically expressed in the organizer. These genes turned out to encode extracellular BMP antagonists that bind to secreted BMPs and inhibit the activation of BMP receptors (Zimmerman et al. 1996; Piccolo et al. 1996). Importantly, overexpression of these BMP antagonists mimicked the action of the organizer in nearly every respect.

Functional interference and genetic manipulations confirmed the requirements for BMP signaling and BMP antagonism in dorsoventral patterning. Of particular importance, genetic mutations in zebrafish *bmp7* and *bmp2b*, BMP receptor *acvr1* and *smad5* (*snailhouse*, *swirl*, *lost-a-fin*, and *somitabun* mutants, respectively) result in dorsalized embryos (Kishimoto et al. 1997; Nguyen et al. 1998; Hild et al. 1999; Dick et al. 2000; Schmid et al. 2000; Mintzer et al. 2001), whereas embryos lacking *chrd* function (*chordino*; Hammerschmidt et al. 1996a, b) are ventralized. Results using antisense oligo-based loss-of-function in *Xenopus* similarly showed dorsalization upon reduction of BMPs and ventralization following inhibition of Nog, Chrd, Fst (Oelgeschläger et al. 2003; Khokha et al. 2005; Reversade et al. 2005). In the mouse and the chick, the roles of BMPs are more complex owing to their functions in extra embryonic tissues and in controlling cell proliferation (Hogan 1996). Nonetheless, *Nog* and *Chrd* are expressed in the anterior primitive streak and Hensen's node (Connolly et al. 1997; Streit et al. 1998; Streit and Stern 1999; Bachiller et al. 2000), and *Nog*, *Chrd* double mouse mutants exhibit dorsoanterior truncations (Bachiller et al. 2000). Additionally, genetic deletion of *Bmp4* in mouse results in ventral mesoderm defects (Winnier et al. 1995).

A host of other BMP antagonists as well as antagonists of other ligands were also discovered and a commonly accepted model emerged; namely that dorsoventral patterning across germ layers is mediated by a gradient of BMP activity originating

ventrally and antagonized dorsally by organizer molecules. Additional support for this model came from characterization of patterns of BMP activity, finding that levels of phosphorylated active Smad1 were enriched ventrally and absent from the organizer (Faure et al. 2000; Schohl and Fagotto 2002).

BMP antagonism derived from the organizer also patterns the convergent extension movements of the axial mesoderm (also see Sect. 6.6). High levels of BMP inhibit convergent extension behavior in *Xenopus* (Graff et al. 1994), and in fish there is evidence that BMP signaling inhibits expression of convergent extension-promoting Wnt/PCP ligands, *wnt11* and *wnt5b* (Myers et al. 2002). BMP signaling may also inhibit dorsal mesoderm involution but this has not been extensively studied (Nakayama et al. 1998). And, BMPs can promote epithelial-mesenchymal transitions and ingression behavior, which occurs in more ventrolateral (non-organizer) mesoderm. Thus, the production of secreted BMP antagonists is essential and largely sufficient for the self-differentiation and morphogenesis of the organizer into notochord mesoderm and for the patterning of the surrounding germ layers.

#### **6.4.2 The Role of the Organizer and BMP Antagonism in Neural Induction**

Because of the historical link to early organizer studies and a general interest in the development of the central nervous system, the role of the organizer in neural induction has received considerable attention. Also, conceptually, neural induction is thought of differently from axis and mesendoderm induction. Mesoderm and endoderm are induced from ectoderm, whereas ectoderm develops if inducers are absent, and can thus be considered the “default” germ layer. Surprisingly and perhaps counter-intuitively, experiments in *Xenopus* suggested that neural, rather than epidermal, was the default state of the ectoderm (Hemmati-Brivanlou and Melton 1997; Chang and Hemmati-Brivanlou 1998; Stern 2006; Ozair et al. 2012). Inhibition of mesendoderm-inducing Tgfb signaling using dominant-negative receptors or endogenous antagonists (Hemmati-Brivanlou and Melton 1994; Chang and Harland 2007) or depletion of maternal *vegt* mRNA (Zhang et al. 1998) results in neuroectoderm formation in all presumptive germ layers. Critical evidence for the neural default model includes experiments indicating that BMPs induce epidermis in dissociated cells that would otherwise become neuralized, and that Nog, Chrd and Fst act as direct neural inducers in ectoderm tissue by blocking BMP signaling. Interestingly, these antagonists elicited only anterior neural fates, which hinted at possible mechanisms of anteroposterior patterning by the organizer (discussion of this aspect will be deferred to the following section).

Whereas the central requirement for BMP antagonism is clear, the extent that BMP inhibition alone is sufficient for neural induction is unclear. Studies in chick and *Xenopus* have indicated that neural induction also requires Wnt antagonism (Wilson et al. 2001; Pera et al. 2003; Fuentealba et al. 2007) as well as ongoing FGF signaling (Linker and Stern 2004; Delaune et al. 2005; Marchal et al. 2009). Subsequent experiments using different methodologies have suggested that FGF

signaling is particularly involved in establishing “pre-neural” ectodermal fate in the chick and is not sufficient for neural induction, whereas in *Xenopus*, BMP antagonism is largely sufficient in vivo (Wills et al. 2010; Pinho et al. 2011). Experiments in mammalian embryonic stem cells indicate that culture under low density or Tgfb-inhibited conditions can lead to neural development (Tropepe et al. 2001; Chambers et al. 2009), largely supporting the default model in mammalian cells.

The relative roles of FGF signaling and/or Wnt antagonism may reflect species level differences in the function of the organizer. The *Xenopus* organizer expresses *shisa2*, an antagonist of Fzds and FGF receptors (Yamamoto et al. 2005). Shisa homologues are not expressed in the organizer in chicken and mouse, although they are expressed in anterior regions (Furushima et al. 2007; Hedge and Mason 2008). Thus, FGF signaling may be less critical in *Xenopus* neural induction because it is normally inhibited in the presumptive neural region. FGF signaling may contribute to neural induction by inhibiting Smad1, through MAPK-mediated linker domain phosphorylation (Pera et al. 2003), or by inducing the expression of various pre-neural genes (Sheng et al. 2003; Pinho et al. 2011). In chick, non-organizer Wnt signaling can promote BMP activity, both indirectly by inhibiting the ability of FGF activity to repress *Bmp* expression and by acting more directly to promote BMP activity (Wilson et al. 2001). This latter activity may occur at the level of C-terminally phosphorylated active Smad1. FGF/MAPK and Gsk3b activity (i.e., absence of Wnt signaling) can phosphorylate the linker region of active Smad1 leading to its turnover, whereas limiting MAPK activity of Wnt treatment can prolong active Smad1 signaling (Fuentealba et al. 2007).

In general, FGF may be more critical in cases where neural development occurs over a longer time course and many rounds of cell divisions, as in the chick and mouse. Additionally, FGFs may have different roles in the context of an epiblast epithelial architecture (i.e., pseudostratified, interdigitating columnar epithelia). A similar idea has come from experiments in embryonic stem cells, where FGF signaling and low Wnt activity are thought to promote a pluripotent stem cell state resembling the postimplantation epiblast (EpiSC) (reviewed in Ozair et al. 2012). Thus FGFs may not just promote a “pre-neural” state, but a “pre-germ layer differentiation” state, which would encompass neural fate.

### 6.4.3 *Self-Organization and Developmental Plasticity in the Organizer*

In addition to the neural inducing and mesoderm patterning functions, the developmental plasticity of the organizer can be largely explained by BMP regulation. Early experiments by Spemann showed that partial constriction of the gastrula along the midline led to double-headed tadpoles, each with a normally formed anterior axis (Spemann 1903). Thus, half an organizer can restore normal bilateral symmetry locally and scale its effects accordingly. Similarly, rather normal embryos develop from dorsal half explants as opposed to hyperdorsalized embryos, which would be the expected result in the absence of self-organization (Reversade and De Robertis 2005).

Recent experiments in *Xenopus* have shown that this self-organization ability relies on the feedback regulation of Chrd flux in the embryo (De Robertis 2009) (Fig. 6.11b). The stability and activity of Chrd is regulated in the extracellular space by proteolytic degradation, mediated by Bmp1, a member of the Tolloid family of metallopeptidases. The activity of Bmp1 can be inhibited by the Sizzled (Szl) protein (Lee et al. 2006), a secreted Frizzled related protein. Szl is expressed ventrally in *Xenopus* as well as zebrafish, and becomes restricted to an extreme ventral domain in the ventral mesoderm (Salic et al. 1997; Yabe et al. 2003; Collavin and Kirschner 2003; Martyn and Schulte-Merker 2003). Szl therefore promotes Chrd activity, creating a relatively shallow gradient of dorsalizing activity, which can explain the apparent paradox that *szl* loss-of-function embryos are ventralized despite its expression ventrally (Yabe et al. 2003; Collavin and Kirschner 2003; Martyn and Schulte-Merker 2003; Lee et al. 2006). Chrd and BMPs proteins can form long-range gradients within Brachet's cleft, the narrow intraembryonic space in the gastrula separating the ectoderm from the mesendoderm, possibly using the fibronectin-rich extracellular matrix to facilitate distribution (Plouhinec et al. 2013).

Self-organization becomes possible because key dorsal and ventral genes encode proteins with opposing molecular activities, but are under differential transcriptional control. Although the organizer expresses many growth factor antagonists, this region also contains several BMP receptor agonists, *admp* (*anti-dorsalizing morphogenetic protein*) and *bmp2* (Moos et al. 1995; Joubin and Stern 1999; Lele et al. 2001; Inomata et al. 2008). A subset of BMP antagonists are expressed in a ventral domain, including *bmpcr/crossveinless2* (Ambrosio et al. 2008) and *bambi* (Onichtchouk et al. 1999). A central element of this regulatory network is the positive feedback control of the dorsal BMP agonists (De Robertis 2009): The prominent ventrally expressed BMPs (*bmp4/7*) are positively regulated by BMP signaling itself, whereas *admp/bmp2* agonists are inhibited by BMP activity. Similarly, the ventrally expressed BMP antagonists and *szl* are induced by BMPs, whereas *chrd* is repressed. Additionally, all BMPs are inhibited by binding to Chrd and can be released for action by Bmp1-mediated cleavage of Chrd.

Thus, in a dorsal half explant, loss of BMPs and Szl would lead to enhanced Chrd cleavage and inhibition resulting in higher *admp/bmp2* expression. These ligands would then accumulate in the area of highest Bmp1/lowest Chrd activity, activating *bmp4/7* expression. Increasing levels of these "new" ventral BMPs would limit the extent of *chrd* and *admp* expression as well as establish a new *szl* domain, resetting the Chrd gradient to the size of the new "embryo" (Reversade and De Robertis 2005). Analogous regulatory networks have been uncovered in zebrafish (Lele et al. 2001; Xue et al. 2014) and are likely to exist in chick, as orthologues of the main genes are present. Mammals appear to lack *Admp* and *Szl* (although *Bmp2* is present), making it unclear whether similar axial self-organization occurs or to what extent mechanistically similar processes have evolved. As the appreciation of network effects in biology is growing, it will be interesting to discover whether these principles of embedded antagonistic proteins under complementary transcriptional control are general features of self-organization in development.

#### 6.4.4 Molecular Interpretation of BMP Gradient Signaling

Accumulated data has provided insight into the mechanisms through which cells interpret different BMP signaling levels to achieve different dorsoventral cell fates. Typical models invoke a positional information mechanism, in which the graded concentration of a morphogen determines cell fate. However, it is becoming clear that there are spatiotemporal aspects to growth factor signaling gradients in general, and to BMP gradients in particular. Recent experiments have shown that dorsalized maternal-zygotic BMP receptor (*acvr1*) mutant fish (*lost-a-fin*) are rescued by inducible *acvr1* expression only if expression is initiated prior to the midgastrula stage (Tucker et al. 2008). Similarly, inducible expression of *chrd* after the middle of gastrulation fails to dorsalize wildtype embryos. Analogous results were seen in *Xenopus* and in fish using timed application of an anti-BMPR drug (Wawersik et al. 2005; Kwon et al. 2010), further supporting the idea that early but not late exposure to BMP signals is critical for inhibiting dorsal fates and promoting ventral fates.

Temporal regulation was particularly evident in timed *Chrd* induction experiments. Dorsoanterior markers become progressively refractory to BMP inhibition, whereas more lateral-posterior tissues (neural crest and pronephros) became progressively more sensitive (Tucker et al. 2008). These studies thus suggest that BMP signaling is active later and longer in ventroposterior tissues and never active in prospective organizer. Additionally, dorsoventral and anteroposterior patterning occurred along a similar time course, indicating these processes are coordinated (see also Sect. 6.5.2). Interestingly in these studies, embryos do not seem to sense cumulative BMP exposure, as ventroposterior fates could still be induced with a late burst of BMP signals (Tucker et al. 2008). As discussed above, the BMP gradient is self-organizing, therefore a small pulse of BMP activity could potentially regenerate the activity gradient to regulate different cell fates. Threshold dose and temporal aspects of exposure to BMP signals thus cooperate to specify axial patterning.

The nature of interpretation of the BMP gradient at the transcriptional level may also partly underlie this temporally progressive dorsoventral patterning. In *Xenopus* ectoderm cells, competence to respond to BMP is maintained throughout gastrulation (Simeoni and Gurdon 2007). And transcription can be triggered rapidly and requires the continuous receptor activation to maintain appropriate steady-state levels of Smad1 in the nucleus (Simeoni and Gurdon 2007). These responses are distinct from those elicited during Activin/Tgfb-mediated mesoderm induction (Chap. 7), which are characterized by a more limited period of competence, extending signaling from active endocytic complexes, delayed onset of transcription, and cumulative sensing of ligand levels (Dyson and Gurdon 1998; Shimizu and Gurdon 1999; Bourillot et al. 2002; Piepenburg et al. 2004; Jullien and Gurdon 2005). The molecular bases for the differences between Tgfb and BMP signaling, despite similar pathway architecture, is not known but may underlie the general tendency of ventral signals to limit dorsal ones. These differences may also help explain the temporal aspects of BMP responses, as discussed above, as well as the importance of Smad1 as a center of signaling integration.

BMP-mediated transcriptional regulation is controlled by the formation of phosphorylated Smad1/Smad4 complexes and subsequent nuclear translocation and association with target promoters. In general, Smad-responsive cis-regulatory elements are thought to mediate low affinity interactions and Smad-associated cofactors are therefore required for specific target promoter interactions. Smad1-binding transcriptional cofactors Znf423 (Oaz; Seoane et al. 2000 and Hivep1 (Human immunodeficiency virus type I enhancer binding protein 1/Schnurri1; Yao et al. 2006) have been characterized and facilitate binding of BMP-activated Smads to BMP-responsive enhancer elements. These cofactors likely recruit general transcriptional activators or repressors depending on the cellular or epigenetic context (Blitz and Cho 2009).

#### **6.4.5 Complexity of Cross-Regulation Between BMP Signaling and the Organizer**

BMP signaling activates a conserved cascade of gene regulation involving several immediate response genes, including the homeobox *ventx* genes, as well as *msx1* and *wnt8a* (Gawantka et al. 1995; Schmidt et al. 1996; Ault et al. 1996; Ladher et al. 1996; Onichtchouk et al. 1996; Suzuki et al. 1997; Hoppler and Moon 1998), and a secondary target, *even-skipped homolog 1 (evx1/xhox3)* (Ruiz i Altaba et al. 1991). Additionally, Ventx and Gsc proteins cross-repress each other's expression, mediating part of the negative feedback regulation of BMP and organizer gene expression (Fainsod et al. 1994; Gawantka et al. 1995; Onichtchouk et al. 1996; Trindade et al. 1999; Sander et al. 2007). In the mesoderm, Brachyury homolog (T) provides an essential input to *ventx* expression (and hence *gsc* repression) through its interaction with Smad1 (Messenger et al. 2005). Interestingly, double inhibition of Ventx and Gsc results in normal embryos (Sander et al. 2007) indicating that these proteins are only strictly required to regulate each other, and that redundant mechanisms pattern the axis in their absence. Multiple such independent and redundant cross-regulatory interactions between Gsc-Ventx and other protein pairs likely would provide robustness to dorsoventral patterning mechanisms.


Similar cross-regulatory loops exist in zebrafish (Nikaido et al. 1997; Melby et al. 2000; Kawahara et al. 2000; Imai et al. 2001), although the Ventx genes have undergone a unique evolutionary trajectory in other organisms (Scerbo et al. 2014). These genes appear to have been lost in rodents and are only expressed in a limited set of hematopoietic cells in humans (Rawat et al. 2010). The Ventxs bear ancient homology to Nanog, a gene implicated in pluripotency and primitive endoderm formation (Kozmik et al. 2001). Nanog may have been co-opted for some of the functions of Ventx in mammals, including possibly Gsc repression (Vallier et al. 2009), whereas Ventx may have substituted for Nanog function in fish and amphibians (Camp et al. 2009; Scerbo et al. 2012). This case is particularly likely in the anuran lineage, which appears to have secondarily lost Nanog (Scerbo et al. 2014). In spite of this reduced dependency on Ventx in mammals, cross-antagonism between BMP signaling and Gsc in patterning the axis/primitive streak has been

evolutionarily maintained, as this relationship exists as reciprocal repression of *EVX1* (activated by BMP signaling) and *GSC* (Kalisz et al. 2012) in human ES cells. Thus, although the involvement of the well-characterized *Ventx* proteins is variable throughout evolution, possibly related to the timing and requirement for ventral mesoderm in hematopoiesis in early development (Kozmik et al. 2001), there remains an ancient and conserved network of reciprocal repression between BMPs and organizer genes.

## 6.5 Anteroposterior Axis Patterning

In general, within the dorsal blastopore/anterior primitive streak, the sequence of internalizing mesoderm determines anteroposterior character, with anterior mesendoderm (anterior definitive endoderm and prechordal plate) being followed by the chordamesoderm (notochord). Signals from the organizer have long been implicated in establishing anteroposterior fates along the body axis. This effect was seen in Spemann and Mangold's early experiments in which second axes induced by organizer transplantation often showed varying degrees of completeness at the anterior end (Spemann and Mangold 1924). These results were roughly correlated with the stage of gastrula from which the donor dorsal lip was taken; earlier lips induced more complete axes with well-formed heads and later lips induced truncated axes or second tails (Spemann 1931). These experiments were interpreted to suggest that the organizer comprised a spatially distinct "head organizer" that would induce forebrain in the early gastrula, and a "trunk organizer" that would induce hindbrain and spinal cord in more posterior ectoderm during later gastrulation. The remaining region, the midbrain, is formed largely by interactions between forebrain and hindbrain.

Work by Nieuwkoop and others later suggested a different model. Implantation of folded strips of competent animal cap ectoderm at various anteroposterior levels in the prospective neuroectoderm generated a graded pattern of neural fates in the implant, with anterior neural/neural plate border tissue forming distally in the implant and more posterior neural forming proximally (Nieuwkoop 1952, 1999) (Fig. 6.12a). These observations were interpreted to suggest a neural "activation" step that initiates a tendency toward forebrain differentiation, followed by a "transforming" event that converts activated anterior neural tissue to more posterior fates (Nieuwkoop 1952, 1999). Earlier experiments had shown that artificial neural induction in urodele animal caps using nonspecific chemical and physical methods invariably generated forebrain (Holtfreter 1944). Also, explants taken from early prospective neuroectoderm showed progressive waves of activation followed by transformation, occurring from posterior to anterior and coincident with mesoderm internalization (Eyal-Giladi 1954). These different activities were found to depend on the nature of the underlying axial mesoderm, with activation predominating in the prechordal mesoderm and anterior notochord and transformation predominating in the posterior notochord (Sala 1955). Experimental data have at various times favored one model or the other,



**Fig. 6.12** Models for anteroposterior axis patterning in vertebrates. **(a)** Depiction of Nieuwkoop's ectodermal fold implantation experiments (Nieuwkoop 1952; Nieuwkoop and Nigtevecht 1954). Dorsal posterior view of a neurula stage amphibian embryo; neural fate is represented as a gradient from light-to-dark with darker color indicating more posterior fates; the epidermis is yellow. The implanted folds are shown as boxes, divided to show the approximately position of induced neural fates. Each fold is characterized by a distal epidermal portion (ep.) bounded by general neural/neural plate border (activated tissue); this is followed proximally by graded neural fates, reflecting the hypothesized influence of a transforming gradient (as opposed to a distinct inducer at each AP level). **(b)** Molecular interpretation of Nieuwkoop's model. In the gastrula, neural induction is accomplished by BMP antagonism, which induces neural tissue with forebrain character. **(c)** During later gastrulation, the expression of Wnts directly induces posterior fates in anterior neural-fated tissue in a dose-dependent fashion. FGF signaling is required in a permissive role. Wnt antagonists expressed in the anterior mesendoderm limit the extent of Wnt signaling and the anterior remains forebrain

but accumulating molecular evidence supports a modified version of the Nieuwkoop model (Fig. 6.12b and c). This support is based on three major findings:

1. that BMP antagonists, Noggin, Chordin and Follistatin act as endogenous neural inducers, and induce exclusively anterior neural fate in competent ectoderm (see Sect. 6.4, Smith and Harland 1992; Hemmati-Brivanlou et al. 1994; Sasai et al. 1994),
2. the identification of "transforming" molecules, such as FGFs, Wnts, Nodals, and retinoids, which posteriorize anterior tissue (reviewed in Stern 2005) but do not directly neuralize ectoderm per se, and,
3. the elucidation of proposed "anterior stabilizing" signals (Fraser and Stern 2004). These are mediated by continuing BMP antagonism (Hartley et al. 2001), Wnt antagonism by inhibitors such as Frzb1 and Dickkopf1 (Leyns et al. 1997; Wang et al. 1997; Glinka et al. 1998), multifunctional antagonism by members of the Cerberus/DAN family of proteins (Bouwmeester et al. 1996) and the feedback Nodal antagonism by Lefty1 (Thisse et al. 2000; Cheng et al. 2000). These molecules are expressed prior to gastrulation in the extraembryonic endoderm (anterior visceral endoderm (AVE)/hypoblast/foregut endoderm), and during gastrulation in the anterior definitive endoderm and prechordal plate, providing signals that initiate and stabilize anterior pattern, respectively.



## 6.5.1 Cellular and Tissue Basis of Anteroposterior Patterning

### 6.5.1.1 The Hypoblast/Anterior Visceral Endoderm in Anterior Patterning

Comparative studies have suggested that the role of the hypoblast/AVE in anterior patterning is widely conserved, based on analogous gene expression patterns in the AVE and later anterior definitive endoderm, cell migration and anterior signaling activities (rabbit hypoblast, Knoetgen et al. 1999; chicken hypoblast, Foley et al. 2000; teleost YSL, Ho et al. 1999; amphibian foregut/anterior endoderm, Bouwmeester et al. 1996; Jones et al. 1999). Interestingly, long before it was molecularly characterized, the hypoblast received considerable attention as a potential regulator of anteroposterior patterning through the control of cell movements. Classical cell tracking experiments in the chicken showed that the pregastrula epiblast undergoes bilaterally symmetrical “polonaise-like” or “double vortex” cell movements, which head anteriorly along the midline before circling back and converging at the posterior, where the primitive streak will form (Wetzel 1929; Gräper 1929). This pattern of cell movements was shown to be controlled by the hypoblast (Waddington 1930, 1933). In these studies, which were repeated and extended by later authors (e.g., Azar and Eyal-Giladi 1981; Foley et al. 2000), changing the orientation of the underlying hypoblast changed the orientation and placement of the forming primitive streak. Although this effect of the hypoblast on cell movements was clearly recognized by Waddington at the time (Waddington 1933), later experiments focused mainly on presumed signaling/inductive properties of the hypoblast in head and mesoderm induction (Azar and Eyal-Giladi 1981).


In mammals, the AVE has been extensively studied at the cellular and genetic level. The AVE forms prior to gastrulation and is characterized by cells with a distinct columnar morphology (rabbit, Viebahn et al. 1995; mouse, Kimura et al. 2000; Rivera-Pérez et al. 2003) as well as specific gene expression patterns (VE-1 antigen, Rosenquist and Martin 1995; *Hex1*, Thomas and Beddington 1996; *Otx2*, Acampora et al. 1995; *Hhex*, Thomas et al. 1998; *Lhx1*, *Gsc*, *Foxa2*, *Cer1*, Belo et al. 1997). In mouse, these genes are expressed in visceral endoderm cells at the distal tip of the forming egg cylinder (Thomas et al. 1998). Cell labeling with DiI (Thomas et al. 1998) or with transgenic methods (Kimura et al. 2000; Rivera-Pérez et al. 2003; Srinivas et al. 2004), showed that these *Hhex*-expressing AVE cells unilaterally migrate towards the future anterior region of the embryo. Physical ablation of patches of AVE compromised development of the underlying forebrain (Thomas and Beddington 1996), demonstrating a requirement for the AVE in anterior patterning. Also, gene targeting analysis of chimeric embryos showed that a number of transcription factors are required in the AVE for its proper formation and migration and for anterior specification (*Otx2*, Acampora et al. 1995; Rhinn et al. 1998; Perea-Gomez et al. 2001; *Foxa2*, Dufort et al. 1998; *Lhx1*, Shawlot et al. 1999 and *Hhex*, Martinez Barbera et al. 2000 among others).

Despite this requirement for anterior development, several experiments have revealed that the hypoblast and related tissues largely lack head organizing properties. Einsteck implantation of the prospective foregut endoderm in *Xenopus* fails to result in ectopic head induction (Bouwmeester et al. 1996). Similar results were seen in chick and mouse regarding the function of the AVE/hypoblast, which likewise fail to induce anterior structures in transplant assays (Tam and Steiner 1999; Foley et al. 2000). Additionally, removal of the hypoblast/AVE results in ectopic primitive streak and mesoderm formation and anteriorization of the embryo, demonstrating that these tissues are repressors rather than inducers of mesoderm (Bertocchini and Stern 2002) as well as permissive regulators of anterior development (Yamamoto et al. 2004). Thus a more complex picture of anteroposterior patterning is emerging in which the hypoblast/AVE/anterior endoderm self-organizes its own morphogenesis and engages in reciprocal signaling with the epiblast/prospective definitive mesendoderm to control cell fate and morphogenesis in the embryo.

### 6.5.1.2 Formation and Migration of the Hypoblast/AVE

The initial formation of the hypoblast/AVE has been studied with considerable detail in the mouse but is little understood in the chicken. In the latter case, hypoblast cells delaminate from the early epiblast, forming isolated clusters, which merge together from posterior-to-anterior (Stern 2004). While it has never been studied as such, it is likely that this process is analogous to the cell sorting of primitive endoderm from epiblast in the mouse blastocyst. In the mouse, the primitive endoderm in contact with the epiblast (embryonic visceral endoderm) differentiates into the visceral endoderm after implantation, under the control of BMP signaling (Yamamoto et al. 2009). Continuing BMP signals from the epiblast and extraembryonic ectoderm likely inhibit the formation of the AVE at the proximal end of the mouse egg cylinder (Rodriguez 2005; Mesnard et al. 2006). The AVE is ultimately specified through the activation of Nodal/Smad2 signals from the epiblast (Waldrip et al. 1998; Brennan et al. 2001) and the inhibition of BMP/Smad1 signals (Yamamoto et al. 2009).

In the presumptive AVE, Nodal indirectly and directly induces a number of AVE transcription factors as well as the expression of Nodal antagonists *Lefty1* and *Cerberus-like (Cerl)* and the Wnt antagonists *Dickkopf1* and *Sfrp5* (Brennan et al. 2001, see below). The expression of *Lefty1* and *Cerl* in the AVE inhibits the autoregulatory maintenance of Nodal expression in the distal epiblast (Norris et al. 2002). As AVE migration commences and is replaced by posterior visceral endoderm, Nodal and Nodal target gene expression are gradually restricted to the posterior epiblast, mediating mesoderm induction and primitive streak formation (Ding et al. 1998; Brennan et al. 2001; Perea-Gomez et al. 2002; Yamamoto et al. 2004). Thus, anterior migration of the AVE has several consequences for anteroposterior patterning: the maintenance of ectoderm germ layer and anterior neural in the anterior epiblast, and the removal of inhibitory signals for primitive streak/mesendoderm formation (Fig. 6.13).



**Fig. 6.13** Model for origin and role of the AVE in anteroposterior patterning in the mouse. (a) In the peri-implantation blastocyst (E5.0), a subpopulation of primitive endoderm (PE) expressing *Lefty1* and *Cerl* arise stochastically positioned asymmetrically in the distal egg cylinder. This population requires *Nodal* and *TdGF1* in the epiblast and is inhibited by *Bmp4* in the extraembryonic ectoderm. (b) As the conceptus grows after implantation (E5.25), the AVE begins to also express *Wnt* antagonists and is repelled by *Nodal* and *Wnt* signals; the action of *BMP* is limited to the proximal epiblast and PE, allowing migration of AVE in the distal region. (c) As AVE migration proceeds, a second set of *Lefty1*; *Cerl*-expressing cells is induced in the distal VE (*2° AVE*) by *TdGF1*-independent *Nodal* signaling. These cells and the *1° AVE* migrate in a coherent stream toward the presumptive anterior, inhibiting *Nodal* and *Wnt* signaling in the epiblast and specifying the anterior neuroectoderm. Progressive loss of *Nodal* from the anterior limits activity to the posterior, where *Nodal* and *Wnt* maintain and amplify each other's expression through *BMP4*, inducing the primitive streak (PS) in the prospective posterior. *ParE* parietal endoderm, *PE/VE* primitive/visceral endoderm, *ExE* extra-embryonic ectoderm. Images were modified and adapted from: Bedzhov I, Graham SJL, Leung CY, Zernicka-Goetz M (2014) Developmental plasticity, cell fate specification and morphogenesis in the early mouse embryo. *Philosophical Transactions of the Royal Society B: Biological Sciences* 369:20130538. doi:10.1098/rstb.2013.0538 under the terms of the Creative Commons Attribution License CC BY 3.0

Recent experiments have identified considerable complexity in the control of AVE specification and migration. Gain-of-function studies in mouse embryos showed that *Cerl/Lefty1* and *Dkk1* can act as attractants for AVE migration, whereas *Nodal* and *Wnts* can act as repellants (Yamamoto et al. 2004; Kimura-Yoshida et al. 2005). Additionally, it is also now better appreciated that the AVE is heterogeneous, arising from an early sub-population of primitive endoderm in the preimplantation blastocyst, in addition to a later population established from naïve distal visceral endoderm (Takaoka et al. 2006). Importantly, the primary AVE cells (*alias* DVE, in some literature) originate asymmetrically in the visceral endoderm and are necessary to “lead” the migration of the secondary AVE in unilateral anterior migration (Takaoka et al. 2011; Morris et al. 2012a). This complexity of the AVE was first hinted at in mice mutant for the *Nodal* coreceptor *TdGF1*, which fail to express early AVE markers (Ding et al. 1998). Although these do become expressed later, the AVE nevertheless fails to migrate. Reporter gene analyses showed that the *Cerl* and

*Lefty1*-expressing primary AVE arises asymmetrically within the primitive endoderm (E4.0–5.0; Takaoka et al. 2006; Torres-Padilla et al. 2007b), with live imaging of cultured embryos revealing a second population gaining *Cerl*-expression at E5.0 (Torres-Padilla et al. 2007b). Genetic lineage labeling of *Lefty1*-expressing cells confirmed that the secondary AVE arises from progenitors that lack early *Lefty1*-expression (Takaoka et al. 2011).

The leading role of the primary AVE was suggested through microsurgical ablation of the primary AVE, which resulted in a wider defect in remaining VE migration (Miura and Mishina 2007). Importantly, inducible genetic lineage ablation of *Lefty1*-expressing cells clearly showed that the secondary AVE does not migrate anteriorly if early *Lefty1*-expressing primitive endoderm is eliminated (Takaoka et al. 2011). Similar results were seen in live imaging of embryos cultured on a collagen-coated polyacrylamide matrix to mimic implantation (Morris et al. 2012a). In this case, laser ablation of leading “pioneer” cells expressing the highest levels of *Cerl* prevented the migration of the AVE. Interestingly, ablation of an immediate “follower” cell also inhibited migration, suggesting that a coherent stream of interacting cells is necessary for anterior migration (Morris et al. 2012a).

This behavior conceptually resembles “follow-the-leader” collective cell migration strategies evident in epithelial wound healing, lateral line and neural crest cell migration, cancer metastasis and other paradigms (Friedl and Gilmour 2009; Wynn et al. 2012; Cheung et al. 2013; Dalle Nogare et al. 2014; Yamaguchi et al. 2015). However, many of the well-characterized cases involve mesenchymal cells or epithelial-to-mesenchymal transitions, whereas AVE migration occurs by directional intercalation and neighbor exchange within the simple epithelium of the visceral endoderm (Srinivas et al. 2004). Trinkaus performed seminal work on coordinated migration in epithelial sheets and coupled protrusive behavior over 40 years ago (Vaughan and Trinkaus 1966), but the mechanisms controlling follow-the-leader behavior are only beginning to be studied. Recent data in *Xenopus* suggest that cell–cell adhesion between leaders and followers results in mechanical stimulation of cadherins and formation of a cadherin-keratin-plakoglobin complex, which is critical for stimulating protrusive behavior opposite the site of cell–cell contact (Weber et al. 2012). This maintenance of cellular tension as well as chemoattraction by the leading cells is likely to play key roles in this coordinated chain migration.

The initiation of AVE migration is similarly not well understood. Nodal-dependent cell proliferation in the epiblast is required (Stuckey et al. 2011), suggesting that the egg cylinder may need to reach a critical size, possibly to extend the AVE outside the reach of inhibitory signals. Thus, an initial asymmetry in the AVE, established when the embryo is small, would be amplified during growth and would determine the direction of subsequent migration, which is maintained by coordinated cell movements.

## 6.5.2 Molecular Regulation of Anteroposterior Patterning

### 6.5.2.1 Nodal and Wnt Antagonists as Regulators of Anterior Patterning

Clues to the molecular regulation of anteroposterior patterning first came from the characterization of conserved Nodal and Wnt growth factor antagonists expressed in the amphibian organizer endoderm region and in the amniote hypoblast/AVE (see below). *Xenopus cerberus* (*cer*) was identified through differential screening of dorsal lip-specific molecules (Bouwmeester et al. 1996). *Cer* overexpression on the ventral side could induce multiple second axes consisting only of heads (*Kerberos* was the multi-headed guard dog of the underworld in Greek mythology; Hesiod and Evelyn-White 1914). This head-inducing activity was attributed to the ability of *Cer* to antagonize multiple developmental signaling molecules extracellularly, including BMP, Wnt and Nodal proteins (Piccolo et al. 1999), although Nodal antagonism is likely paramount. The activity of *Cer* could largely be recapitulated by co-expression of the Cerberus Nodal antagonist domain (Cerberus-short, *CerS*) with *Chrd*. Additionally, overexpression of *nodal1* (*nodal homolog 1*) after the mid-blastula transition (i.e., from a plasmid vector) resulted in anterior truncations (Piccolo et al. 1999). Additional studies with amniote *Cer1* suggested the anti-Wnt activity is minimal in mouse and chick and that *Cer1* represents mainly a Nodal and BMP antagonist (Marvin et al. 2001). These and studies in zebrafish using *Lefty*, another Nodal antagonist, established that inhibition of Nodal activity is critical for proper patterning of anterior fate in the axial mesendoderm (Thisse et al. 2000).

In addition to anti-Nodal signals, Wnt inhibitors are expressed within the organizer and in the hypoblast/AVE and Wnt signaling is antagonistic to anterior development in a stage-dependent manner. Although Wnt signaling is critical for axis induction, other experiments show that activating Wnt/beta-catenin signaling during gastrulation causes anterior truncations (Christian and Moon 1993; Kelly et al. 1995b; Hoppler et al. 1996) whereas inhibiting late Wnt signaling results in hyperanteriorization (Hoppler et al. 1996; Leyns et al. 1997; Wang et al. 1997; Glinka et al. 1997, 1998). Additionally, inhibition of BMP and Wnt signaling together is needed to induce secondary axes with head structures, whereas induction with BMP antagonists alone generates only trunk structures (i.e., notochord; Glinka et al. 1997). Genetic studies in zebrafish *pcf3* (*headless*, *hdl*) mutants show a reduction in anterior structures and expanded ventrolateral derivatives (Kim et al. 2000), whereas the opposite occurs in *wnt8a* mutants (Kim et al. 2000; Lekven et al. 2001). Wnt antagonists are also required in vivo for anterior patterning. Interference with *Dkk1* function in fish and frogs leads to anterior truncations (Hashimoto et al. 2000; Shinya et al. 2000), mimicking Wnt overexpression. Similarly, genetic deletion of *Dkk1* in the mouse epiblast led to loss of forebrain neural markers (Mukhopadhyay et al. 2001). Additionally, *Dkk1* is a direct target of *Otx2*, a key anterior specifying gene, in both the anterior mesendoderm (Ip et al. 2014) and anterior visceral endoderm (Kimura-Yoshida et al. 2005). And, either overexpression of *Dkk1* or additional deletion of one *Ctnnb1* allele is sufficient to partially rescue loss of *Otx2* (Kimura-Yoshida et al. 2005), suggesting that controlling Wnt levels is a critical role of *Otx2*.

### 6.5.2.2 Self-Regulation of Nodal and Wnt Signaling During Anterior Patterning

As is the case for BMP signaling/antagonism and the organizer, Nodal and Wnt signaling are also precisely regulated through auto-regulation and negative feedback loops controlled by secreted antagonists. Nodal can directly auto-regulate its own expression in the epiblast and visceral endoderm through a Smad2/Foxh1-dependent enhancer in the first intron of the *Nodal* locus (Osada et al. 2000; Norris et al. 2002). Also, *Wnt3* is expressed in the proximal egg cylinder ectoderm and can regulate *Nodal* expression through a Wnt/beta-catenin-responsive proximal ectoderm enhancer (Brennan et al. 2001; Ben-Haim et al. 2006). *Wnt3* is in turn regulated by secreted Bmp4 signals emanating from a population of extra-embryonic ectoderm adjacent to the epiblast. This population is maintained by epiblast Nodal signaling, forming the second feedback loop (Ben-Haim et al. 2006). Nodal signals emanating primarily from the epiblast also induce the AVE at the distal end of the egg cylinder in a Smad2-dependent manner, regulating (both directly and indirectly) the expression of Nodal antagonists *Cer1* and *Lefty1*. These in turn feedback and inhibit Nodal signaling in the underlying epiblast.

*Wnt3* also activates its own expression in the posterior epiblast and visceral endoderm directly through beta-catenin-dependent Wnt signaling (Tortelote et al. 2013) and Wnt signals (likely including Wnt3) directly activate *Dkk1* expression in a number of cell types (Chamorro et al. 2005). Interestingly, the autoregulation of *Wnt3* is evolutionarily ancient, occurring in sponges and in cnidarians (Holstein 2012). Both Wnt/Dkk1 and Nodal/Lefty1 interactions have been suggested to constitute reaction-diffusion systems in vivo, with regard to hair follicle spacing in the case of Wnt (Sick et al. 2006) and to left-right patterning in the case of Nodal (Nakamura et al. 2006). Whether or not these match the strict mathematical assumptions of Turing reaction-diffusion systems (the case for Wnt has been disputed, Meinhardt 2012), it is clear that interacting networks of self-propagating and self-limiting loops of Nodal, Wnt and BMP signaling underlie much of axis formation in vertebrates and indeed likely all animals.

In addition to reciprocal regulation of Nodal and Wnt, there is extensive interaction between Wnt/BMP and Nodal/BMP at the level of signaling integration during antero-posterior patterning. Wnt signaling can potentiate BMP signaling through several possible mechanisms. In the frog, BMP-activated Smad1 can be inhibited by Gsk3b phosphorylation of the linker region, and this in turn can be inhibited by Wnt leading to the perdurance of active Smad1 (Fuentelba et al. 2007). In zebrafish embryos, dorsoventral and anteroposterior patterning occur along a similar time course (Tucker et al. 2008). Correspondingly, manipulation of Wnt signaling can alter the anteroposterior character of Chrd-induced tissues at any time during gastrulation (Hashiguchi and Mullins 2013), suggesting that BMP and Wnt signaling mechanisms are active together, along with FGF and retinoid signaling (Hashiguchi and Mullins 2013). Additionally, in the chick epidermis Wnt signaling can block FGF inhibition of *Bmp4* transcription during neural plate induction (Wilson et al. 2001), although it is unclear whether this interaction is also critical for anteroposterior neural plate patterning.

Nodal and BMP signaling also interact at several different levels. Nodal and certain BMPs can form inactive heterodimeric ligands that fail to stimulate either signaling pathway and exhibit mutual antagonism (Yeo and Whitman 2001). Also, in the frog organizer Nodal and Admp (a BMP ligand) form a self-regulating network to control anteroposterior patterning (Inui et al. 2012). Nodal and Admp can compete for a shared receptor, Acvr2a, to oppositely regulate Nodal and Wnt antagonists in the anterior endoderm. High Nodal levels activate high expression of Nodal antagonists, which subsequently block Nodal-Acvr2a interaction and permit the occupancy by Admp. This alternate pathway limits the expression of Nodal antagonists, allowing Nodal signaling to reengage (Inui et al. 2012). Interestingly, Admp depletion results in the anterior endoderm gaining head-inducing activity in Einsteck transplantation assays, owing to overproduction of Nodal and Wnt antagonists (Inui et al. 2012). The high degree of interrelatedness of these signaling pathways thus can underlie many of the self-organizing properties of early axis formation.

### ***6.5.3 Regulation of Posterior Development by Wnt Signaling***

#### **6.5.3.1 Default Specification of Anterior Neural Fate**

In line with the activation-transformation model, experimental neural induction in *Xenopus* animal caps (ectoderm), either by Noggin treatment (Lamb et al. 1993) or by disaggregation (Grunz and Tacke 1989, 1990), results exclusively in anterior neural induction. Additionally, anteroposterior patterning is maintained in frog embryos lacking mesoderm and endoderm, generated by *vegt* depletion or CerS overexpression (Zhang et al. 1998; Wessely et al. 2001), and in those lacking both organizer and ventral BMP signals (Reversade and De Robertis 2005). Similarly in zebrafish, embryos lacking Nodal signaling and thus mesendoderm also retain anterior identity in the ectoderm (Gritsman et al. 2000). In the mouse, loss of Nodal signaling results in anterior neural specification in the epiblast in the absence of mesendoderm and the AVE (Conlon et al. 1994; Brennan et al. 2001; Camus et al. 2006). The molecules mediating this anterior default state are not known but are hypothesized to be present maternally in fish and amphibians and in the early epiblast of amniotes. Because embryonic stem cells can also acquire anterior neural fates by default (Tropépe et al. 2001; Chambers et al. 2009), it is possible that core pluripotency factors or proteins activated at the onset of epiblast differentiation would be good candidates for anterior specifiers.

#### **6.5.3.2 Graded Wnt Signals and Posterior Specification**

Although many different signaling molecules have been implicated in posterior neural fate specification, current evidence suggests that the graded activity of Wnt signaling is the critical proximate determinant of early posterior identity. Wnt

activity can be seen in *Xenopus* embryos as a gradient of nuclear beta-catenin in the prospective neural plate of the gastrula, with the highest levels posteriorly (Kiecker and Niehrs 2001). Similarly, in the mouse gastrula, posterior gradients of Wnt reporter construct activity have been observed (Maretto et al. 2003; Lewis et al. 2007, 2008; Fossat et al. 2011). Experiments in neuralized *Xenopus* animal caps showed that inducible expression of activated beta-catenin during gastrulation directly induces posterior neural markers (McGrew et al. 1997; Domingos et al. 2001). Additionally, cell–cell contact and FGF signaling are required in this case, suggesting the involvement of a “community-effect” response, as described for muscle formation (Standley and Gurdon 2004). Similar requirements for Wnt and FGF signaling were seen in chicken, where Wnt treatment directly and dose-dependently posteriorizes competent anterior neural tissue explants (Nordstrom et al. 2002). Correspondingly, Wnt inhibition in posterior neural explants causes anteriorization. FGF signaling is also required for posterior specification, but likely in a permissive role, acting in a non-dose-dependent fashion. And loss of FGF signaling does not anteriorize posterior neural fates (Nordstrom et al. 2002). Wnt signaling also has subsequent roles in specifying the posterior portion of major brain subdivisions later in development, in particular distinguishing diencephalon versus telencephalon (Heisenberg et al. 2001; Houart et al. 2002). It is therefore of great interest to determine how this signaling system is compartmentalized and reiterated during multiple aspects of individual tissue and organ development.

At the transcriptional level, Wnt/beta-catenin signaling during posterior specification is thought to proceed through the use of Tcf/Lef proteins as coactivators. This mechanism of gene activation is likely different from early, axis-inducing Wnt signals, which involve mainly derepression of Tcf3 (see Sect. 6.3.1). In mouse development, double homozygous genetic deletion of *Lef1* and *Tcf7(Tcf1)* results in posterior defects in addition to abnormalities in mesoderm specification (Galceran et al. 1999). Similarly in *Xenopus*, *Lef1* and *Tcf1* are required for the late/ventral response to Wnt (Roël et al. 2002; Liu et al. 2005), whereas Tcf3 is dispensable in this regard (Hamilton et al. 2001). Recall that Tcf3 is necessary for the repression of early Wnt target genes in vertebrates. Experiments in mouse and frog have suggested a “Tcf exchange” model for the changes in target gene responses to Wnt signaling prior to and following axis formation. *Lef1* can be transcriptionally activated by Wnt signaling (chick, Skromne and Stern 2001; mouse, Wu et al. 2012a), suggesting that cells experiencing Wnt signals undergo a qualitative change in their potential underlying responses to subsequent Wnt exposure during development. Similarly, early Tcf3 repressor function is inactivated by Wnt-activated kinases and replaced by activating Tcf1 function in both *Xenopus* and in mouse embryonic stem cells (Hikasa et al. 2010; Yi et al. 2011). Thus, the activation of genes during posterior specification is likely to involve both beta-catenin-dependent derepression of Tcf3, followed by replacement and activation by Lef1/Tcf1. In support of this idea, mouse knock-ins in which Tcf3 has been replaced by the deltaNTcf3 construct (non-derepressible) lack *Lef1* expression and show defective late Wnt responses (Wu et al. 2012a).



## 6.6 Axial Morphogenesis

The physical form of the vertebrate axis is brought about during gastrulation. During this process, the presumptive mesoderm and endoderm germ layers are internalized and undergo a set of morphogenetic events that elongate the axis in the anteroposterior direction and establish dorsoventral (mediolateral) organization in the mesoderm. Vertebrate embryos undergo gastrulation in one of two basic ways, depending on the architecture of the embryo. In spherical anamniote embryos, internalization occurs through the blastopore, which circumscribes the marginal zone, and forms the archenteron (primitive gut tube). In the flat blastoderm of amniote embryos, internalization generally occurs through the primitive streak in the posterior of the embryo, forming a trilaminar embryo with the endoderm remaining open to the yolk and forming a tube secondarily.

In recent years, the cellular mechanics of vertebrate gastrulation have been studied in great depth using live imaging approaches. Surprisingly, the underlying cell and tissue mechanics and their molecular regulation are extensively conserved, despite differing patterns and timing of gastrulation movements. Largely based on the seminal work of Trinkaus, Keller, Stern and others, a common set of vertebrate gastrulation movements and cell behaviors have been identified. These have been extensively reviewed in recent years (Keller 1986; Keller and Shook 2004; Stern 2004; Solnica-Krezel 2005; Tam et al. 2006; Shook and Keller 2008; Solnica-Krezel and Sepich 2012) and fall into three main categories, internalization, convergent extension and epiboly. Together, these processes form a “mosaic of regional processes” (Keller 1986) that are exquisitely integrated to accomplish the feat of gastrulation.

Mesendoderm internalization typically initiates by invagination (in-pocketing) in a tissue layer (the epiblast), followed either by involution as an in-rolling sheet of cells over a lip (*Einrollung*; Vogt 1929), or by ingression of individual cells. Invagination and ingression proceed by apical constriction of epiblast cells, and in the case of ingression, epithelial-to-mesenchymal transition and subsequent cell migration. Convergent extension is a thinning and narrowing of a tissue and occurs by mediolateral intercalation of closely packed cells or by migration of cells toward a midline. Cells in the mesendoderm undergoing convergent extension typically elongate along the mediolateral axis and exhibit bipolar protrusive behavior using lateral lamellopodia. The traction generated by this bipolar activity is thought to facilitate cell intercalation. Epiboly is a thinning and spreading/extension of a tissue and involves radial intercalation to form a thinner cell layer, as well as cell shape changes. In spherical embryos, epiboly is essential for the ectoderm to spread and cover the yolk mass during gastrulation. Uniquely in teleost embryos, epiboly encompasses all germ layers and the entire embryonic blastoderm covers the yolk cell.

### 6.6.1 Amphibian Gastrulation

Gastrulation in amphibians initiates at a latitudinal slit at the boundary of equatorial and vegetal regions, forming the dorsal lip of the incipient blastopore. Cell internalization begins with apical constriction of the so-called “bottle cells,” creating the

initial invagination at the blastopore (Keller and Shook 2004). Extension of this cavity during gastrulation creates the archenteron. Internal vegetal rotation cell movements in the presumptive endoderm pull the marginal zone vegetally and initiate internalization by involution at the dorsal lip. Internalization spreads laterally from the dorsal lip eventually forming a circular blastopore, with axial mesoderm entering at the early dorsal lip and more paraxial and lateral mesoderm internalizing from the lateral and ventral blastopore (Keller and Shook 2004).

On the dorsal side, the anterior leading edge of the mesoderm (prechordal/head mesoderm) migrates as a cellular stream along the blastocoel roof (Winklbauer and Nagel 1991; Winklbauer et al. 1992). This migration requires fibronectin fibrils deposited by the blastocoel roof cells (Boucaut et al. 1985; Winklbauer and Nagel 1991; Nagel and Winklbauer 1999). Internalizing mesoderm cells maintain what is termed “tissue separation” with the ectoderm, and cannot intercalate back into the ectoderm (Wacker et al. 2000). This segregation is evident by the presence of Brachet’s Cleft between the mesoderm and ectoderm/neuroectoderm (Nieuwkoop and Florschütz 1950). An analogous cleft separating these tissue layers is found throughout the vertebrates.

In *Xenopus* and other anurans, the majority of the mesoderm arises from deep rather than superficial cells (Nieuwkoop and Florschütz 1950; Keller 1975). Involution of the mesoderm is thus internal in these species and provides the main mechanical forces driving gastrulation. The mesoderm and deep neural ectoderm undergo radial intercalation just prior to involution, occurring first in more anterior cells and later in posterior ones (Keller et al. 1985). Additionally, tissue extension is biased in the anterior–posterior direction, likely because some initial mediolateral intercalation occurs. After involution, cells become elongated perpendicular to the axis of extension and exhibit bipolar protrusive behavior and undergo mediolateral intercalation (Keller and Tibbetts 1989; Shih and Keller 1992b). This behavior drives convergent extension of the marginal zone and is the major biomechanical force behind internalization and elongation of the axis. Mediolateral intercalation behavior begins in more anterior tissue as it involutes and spreads to more posterior axial tissue as well as lateral mesoderm as the blastopore closes.

Closure of the blastopore at the ventral marginal zone is accomplished by convergent thickening, a little understood behavior in which prospective posterior somitic mesoderm converges into a pile of cells that do not extend, although they do so later in neurulation (Keller and Danilchik 1988). In general, the process of convergence can lead either to extension or thickening, or both, with the degree of each outcome depending on the constraints applied by the surrounding tissue. In either case, the progressive expansion of mediolateral intercalation behavior around the blastopore and the linking of forces around the marginal zone is thought to create “hoop stress” that pulls the blastopore lip over the yolk plug and internalizes the endoderm (Keller et al. 2000; Keller and Shook 2004).

Urodele and anuran amphibians undergo significant differences in development that have been long noted but perhaps not well appreciated. The process of gastrulation in salamanders and newts is outwardly similar to the situation described in

*Xenopus*, but has several important distinctions. Mesoderm is derived from the surface layer in urodeles (Vogt 1929). Axial mesoderm undergoes similar involution at the dorsal lip in urodeles, but internalization in the ventral and lateral regions occurs mostly by ingression of individual cells (Shook et al. 2002). This pattern of ingression has been referred to as a “bilateral primitive streak.” In urodeles, lateral and ventral convergent forces are generated by the ingressing cells in these regions, as these cells undergo apical constriction and bring additional cells into the ingression zone (Keller and Shook 2004).

Interestingly, the urodele pattern of dorsal lip involution and lateral ingression is conserved in more primitive organisms including cyclostomes and lungfish (Shook and Keller 2008). It is likely that both the anuran and teleost mechanisms of primary internalization by involution evolved independently within those lineages. These changes in gastrulation may be correlated with the cellular organization of the ectodermal moiety, occurring as a multilayered ectodermal “deep layer” of non-polarized cells, covered by a superficial polarized epithelial layer in the anurans and teleosts, as opposed to the typical “epiblast” of pseudostratified interdigitating epithelial cells (Shook and Keller 2008).

### 6.6.2 Teleost Gastrulation

Similar morphogenetic events occur in the zebrafish/teleost gastrula, albeit in the context of a different embryonic architecture. One of the main features of fish gastrulation is prominent epiboly, in which all three germ layers (as opposed to just ectoderm) undergo extensive radial intercalation as the blastoderm expands to cover the yolk cell. Epiboly in teleost fish is somewhat unusual, and involves interconnections between the yolk cell and the superficial epithelial layer of the blastoderm. Epiboly begins just prior to gastrulation with YSL nuclei migrating vegetally in advance of the blastoderm margin. Cytoskeletal perturbations and in vivo observations suggest that the YSL “tows” the blastoderm margin vegetally (Betchaku and Trinkaus 1978; Solnica-Krezel and Driever 1994).

Internalization initiates by an involution-like “synchronized ingression” around the blastoderm margin, forming a bilayered structure, the germ ring (Adams and Kimmel 2004). Typical ingression is not thought to occur, as teleost gastrulation lacks a classical epithelial-to-mesenchymal transition (EMT) and cell migration characteristic of typical ingression movements (Shook and Keller 2008). Internalizing cells migrate beneath the outer layer toward the animal pole, forming an inner hypoblast (this is mesendoderm, and not homologous to the amniote hypoblast) and outer epiblast (ectoderm). Both layers are covered by a flattened superficial epithelium, the enveloping layer (EVL), which remains extraembryonic and may provide structural support for morphogenetic movements (Kimmel et al. 1990). On the dorsal side of the margin, involuting and single “delaminating” cells form a thicker layer, constituting the embryonic shield (Montero 2005). Also dorsally, a

teleost-specific population of epiblast cells, the dorsal forerunner cells (DFCs) detach and migrate vegetally ahead of the dorsal lip, in association with the margin of the EVL (Cooper and D'Amico 1996). Their function at this stage is unknown, but later these cells will form the Kupffer's vesicle, which functions in left-right asymmetry. Also, similar to the process in anurans, convergent extension elongates the axial mesendoderm as a result of bipolar protrusive behavior and well as dorsally directed migration of more lateral cells (Solnica-Krezel 2005). Teleost thus appear to have independently evolved an amphibian-style method of gastrulation as a solution to concentrated yolk and discoidal cleavage in the egg.

### 6.6.3 *Amniote Gastrulation*

In the mammals and birds, gastrulation initiates at the primitive streak, a groove formed by internalizing cells along the posterior axial midline. Anterior–posterior patterning, and thus the positioning of the primitive streak towards the posterior, is controlled by the anterior migration of the hypoblast/AVE (Sect. 6.5.1). Mesoderm precursors are initially found throughout the posterior epiblast but migrate toward the posterior end prior to primitive streak formation. In the chick, this occurs through the double-vortex polonaise-like cell movements (see Sect. 6.5.1) and similar mechanisms have typically been thought to occur in mammals. However, mouse embryos appear to lack these large-scale cell movements in the epiblast (Williams et al. 2011), possibly owing to a smaller number of cells at the equivalent stage or to constraints of the cup-shaped architecture. Alternatively, the cell movements positioning the streak may occur in the visceral endoderm layer in mouse (Weber et al. 1999). Polonaise-like movements appear to occur in rabbit embryos, which have a flat morphology similar to chick embryos (Halacheva et al. 2011). Interestingly though, in this case, a more localized extreme intercalation cell movement event (“processional cell movements”), predominates (Halacheva et al. 2011). These observations indicate that overall primitive streak positioning in amniotes can result from many different patterns of cell movements depending on the species.

Internalization through the primitive streak involves massive epithelial-to-mesenchymal transition (EMT) and ingression of individual cells to form definitive mesoderm and endoderm (Stern 2004). Early ingressing mesoderm contributes mainly to pharyngeal and cardiac mesoderm anteriorly, as well as extraembryonic mesoderm posteriorly. Definitive endoderm ingresses by mid-streak stages and intercalates into and displaces the hypoblast/visceral endoderm. The primitive streak elongates and progresses anteriorly (distally in the mouse egg cylinder) until mid-gastrulation (late streak stage), approximately to the level of the future hind-brain, and begins to form the anatomically distinct Hensen's node at the anterior tip. Internalization at Hensen's node in many birds and mammals, including humans, resembles that of the amphibian dorsal lip, with invaginating bottle cells and involution of notochord and somitic mesoderm (Shook and Keller 2008). Chick and mouse

embryos however appear to have separately evolved ingression as the main mode of internalization, although dorsal lip involution may be evolutionarily ancestral.

Reptiles (non-avians) represent a useful but little-studied intermediate case. These embryos typically form a dorsal lip perpendicular to the axial midline in the posterior of the blastoderm, through which chordamesoderm involutes. Lateral and ventral cells ingress through a reptilian blastoporal plate, which resembles both the urodele blastopore and amniote primitive streak (Bertocchini et al. 2013).

After its formation, the Hensen's node begins to regress posteriorly and the primitive streak closes in an anterior-to-posterior progression. Notochord and somites form in the mesoderm behind the regressing node. At this point Hensen's node and surrounding cells likely comprise a pluripotent stem cell population that persists throughout regression and into tailbud formation (Wilson et al. 2009). It is often not discussed as such, but amniote primitive streak regression is likely an analogous process to blastopore closure, with mediolateral intercalation behavior driving convergent extension in the axial midline tissues and convergent behavior following ingression at the primitive streak. For the mouse, mediolateral intercalation of mesoderm has been established as a mechanism of axial elongation, and convergent-extension behavior has been demonstrated to initiate in pre-somitic mesoderm cells immediately after they exit the primitive streak (Yen et al. 2009).

In all vertebrates, the embryo lengthens and narrows along the AP axis at the culmination of gastrulation. The neural plate is visible in the dorsal ectoderm and the closed blastopore represents the posterior pole of the embryo. Patterning of the AP and DV axes is essentially accomplished during gastrulation. Although the left-right axis is thus specified topologically, the cellular and molecular mechanisms that regulate left-right asymmetry of the internal organs do not come into play until after gastrulation. These are primarily mediated by a distinct ciliated structure that forms during late gastrulation at the posterior end of the notochord along the gastrocoel roof plate, a part of the archenteron lining (Blum et al. 2014). This structure is derived from surface mesoderm in the late involuting dorsal lip and is homologous in many respects across vertebrates. Monocilia in the posterior notochord region are thought to create a leftward fluid flow, generating asymmetry in Nodal signaling in the left lateral plate mesoderm, leading to morphological left-right asymmetry and the generation of internal bilateral asymmetry in organs such as the heart, gut and lungs (also, see Blum et al. (2014) for review of caveats in chick and pig, which lack nodal cilia (Gros et al. 2009)).

#### 6.6.4 *Molecular Regulation of Axial Morphogenesis*

The control of convergent extension has been extensively studied at the molecular level in *Xenopus* and zebrafish embryos. A host of cellular and genetic experiments have implicated components of the Wnt/PCP pathway as critical regulators of cell shape, cell polarity and activity during convergent extension. In *Xenopus*, overexpression of certain beta-catenin-independent Wnts and dominant-negative Dishevelled 2 (Xdd1; Sokol 1996) were sufficient to block axis elongation without

blocking mesoderm formation. Additionally, mutation or inhibition of *wnt11* function in fish and frogs indicated that Wnt/PCP signaling was indeed required for convergent extension behavior in vivo (Tada and Smith 2000; Heisenberg et al. 2000). Importantly, Wallingford et al. (2000) showed in detailed live imaging of explants that Dvl activity was needed for bipolar cell shape changes and for stabilizing mediolaterally oriented bilateral protrusive activity of lamellopodia in converging and extending mesodermal cells. And similar regions of the protein were required for controlling CE in *Xenopus* as for *Drosophila* PCP, namely the C-terminal DEP domain (Wallingford et al. 2000). Analogous results were found for zebrafish *wnt11* and *wnt5a* mutants, which showed unstable monopolar protrusive activity in dorsally migrating cells (Kilian et al. 2003; Ulrich et al. 2003). Other PCP proteins have also been shown to control cell intercalation by regulating mediolateral lamellopodia in fish and frogs, including Frizzleds, Vangl2, and Prickle (Djiane et al. 2000; Darken et al. 2002; Goto and Keller 2002; Jessen et al. 2002; Wallingford et al. 2002; Veeman et al. 2003b; Takeuchi et al. 2003), as well an extracellular modulator of Wnt, Glypican 4 (*knypek*, Topczewski et al. 2001).

More recent data from amniotes has shown that the regulation of cell shape and the formation of mediolateral lamellopodia are conserved in mouse (Ybot-Gonzalez et al. 2007; Yen et al. 2009; Williams et al. 2014) and chicken (Voiculescu et al. 2007). Additionally, mutations in core Wnt/PCP proteins in mice and humans display axis elongation and neural tube defects, both of which depend on convergent extension (Vangl1/2, Kibar et al. 2001; Murdoch et al. 2001; Montcouquiol et al. 2003; Kibar et al. 2007; Torban et al. 2008; Celsr1, Curtin et al. 2003; and Dvl1/2, Wang et al. 2006). Also, noncore vertebrate Wnt/PCP regulators Ptk7 and Scrib are also required for axis elongation and convergent extension, with Ptk7 likely directly controlling bipolar protrusive activity (Montcouquiol et al. 2003; Lu et al. 2004; Yen et al. 2009). The exact role of Ptk7 is unclear, although genetic experiments in zebrafish have suggested a role in antagonizing Wnt/beta-catenin signaling to allow Wnt/PCP signaling (Hayes et al. 2013). However, beta-catenin-activating roles have also been described suggesting temporal or context-dependent roles.

In addition to controlling cell intercalation during convergent extension, Wnt/PCP signaling is also implicated in cell intercalation prior to primitive streak formation. In the chicken, local cell intercalation in the presumptive primitive streak region prior to gastrulation is thought to result from, and may facilitate the polonaise movements in the epiblast. A number of Wnt/PCP components are expressed in the midline prior to primitive streak formation and inhibition by electroporation of anti-PCP morpholino oligos can inhibit streak morphogenesis (Voiculescu et al. 2007). Additionally, and consistent with the role of the hypoblast in controlling polonaise movements, rotation of the hypoblast can alter Wnt/PCP gene expression, possibly through Fgf8 signaling (Voiculescu et al. 2007). Interestingly, it is likely that ingression itself can function locally to organize polonaise-like movements and pull additional cells into the streak, thus driving ingression in a feed-forward manner (Voiculescu et al. 2014).


The situation is less clear in mammals, as Wnt/PCP components have not been widely implicated in primitive streak formation in the mouse. However, ingression at the primitive streak in the mouse does require the down-regulation of Vangl2 protein,

mediated by Dact1 (Dishevelled-binding antagonist of beta-catenin 1; Suriben et al. 2009). Thus, whereas mice may exhibit a reduced role for cell movement-driven mediolateral intercalation during primitive streak formation (see above), a role for Wnt/PCP components may have been retained at the level of EMT in the forming primitive streak.

Wnt/PCP signaling exerts its effects on intercalating cells by coordinately regulating bipolar cell shape and polarization of actin dynamics to generate tensile forces along the cell's mediolateral axis. The mechanisms that orient this bilateral protrusive activity perpendicular to the elongating anteroposterior axis are not well understood. Experiments using dissociated *Xenopus* animal cap cells treated with graded Activin doses or the recombination of anterior versus posterior notochord explants showed that apposition of tissue from different axial levels was necessary for convergent extension along a perpendicular axis, whereas recombination of equivalent levels produced no elongation (Ninomiya et al. 2004). It is possible that cells could respond to this gradient of Tgfb signaling through changes in cadherin- and protocadherin-mediated cell adhesion during intercalation (Briehner and Gumbiner 1994; Kraft et al. 2012). Similar results have been obtained by opposing gradients of Nodal and BMP activity in zebrafish blastoderm explants (Xu et al. 2014), suggesting an important role for integration of these pathways in controlling elongation. Temporal changes in Nodal signaling are also correlated with the onset of mediolateral intercalation behavior, further suggesting that the local activity of Wnt/PCP may be cued by differences in Tgfb activity.

Tgfb signaling, most likely through Nodal and Nodal-related proteins, is also involved in inducing ingression behavior and bottle cell formation by activating EMT. This can be shown in *Xenopus* by overexpression of Nodal in animal caps, which results in ectopic bottle cell formation (Lustig et al. 1996; Agius et al. 2000; Kurth and Hausen 2000). Also, loss of Nodal signaling dramatically reduces EMT-like cell ingression in zebrafish embryos (Keller et al. 2008). Movement of the hypoblast in amniotes, which expresses Nodal antagonists, is also highly correlated with primitive streak formation in the epiblast formed by those cells left behind. This Nodal-controlled ingression may also be self-regulating, as ingression may facilitate recruitment of additional cells to the streak (see above), which would expose more cells to Nodal signals, further triggering more EMT and ingression (Voiculescu et al. 2014).

Signaling by FGFs also plays a key role in regulating convergent extension and mediolateral intercalation. FGF signaling can regulate *brachyury (t)* expression, which in turn can activate *wnt11* (Tada and Smith 2000). In addition to its role in morphogenesis, FGF signaling also has a well-known role in mesoderm formation (Chap. 7). Experiments with cytoplasmic antagonists of FGF signaling, Sprouty/Spry and Spred, indicate that maternal Sprys antagonize convergent extension behavior by blocking PKC and calcium outputs of FGF signaling (Sivak et al. 2005). Later during gastrulation, zygotic Spred proteins accumulate and antagonize Mapk signaling, allowing the morphogenetic signals to predominate after mesoderm specification has occurred (Sivak et al. 2005). Additionally, FGF regulation of convergent extension is controlled by the Nodal related 3.1 protein through an atypical and little characterized mechanism (Yokota et al. 2003).



**Fig. 6.14** Model for the evolution of axis formation in vertebrates. Highly generalized schematic diagrams of different vertebrate embryos during early (*upper panels*) and middle (*lower panels*) gastrulation. In the basal vertebrates (*left panels*; vegetal views), gastrulation initiates at the organizer with induction by the Nieuwkoop center (high/early Nodal signaling). The initial internalization movements are through involution. By mid-gastrulation, ingression of mesendoderm progresses around the vegetal cells (yolk plug) and forms the nascent blastopore. Involution dorsal mesoderm undergoes convergent extension via mediolateral cell intercalation under the control of Wnt/PCP signaling. During the evolution of amniotes (*middle panels*; top/dorsal views), eggs increased in size and yolk content and began to undergo meroblastic cleavage. Organizer induction by Nieuwkoop center molecules is retained in early gastrulation. Ingression proceeds through the horizontal slit of the blastoporal plate/blastopore and does not circumferentially envelop the non-cleaving yolk (not shown). In the evolution of modern birds and mammals, gastrulation initiates with ingression at the primitive streak, which would be homologous to the later ventrolateral blastopore in ancestral forms. The organizer (Hensen's node) is induced later by Nodal signaling from the middle primitive streak. The heterochrony in the pattern of gastrulation morphogenesis and organizer formation could result from several main events; the hypoblast/anterior endoderm segregating from the epiblast as opposed to forming from cleaving vegetal cells, the loss of early organizer induction, the apparent emergent behavior of polonaise-like movements leading to primitive streak formation and the relatively later induction of Hensen's node by the primitive streak

## 6.7 Concluding Remarks

The overall conservation of axial patterning mechanisms in vertebrates has long been recognized. However, detailed investigations at the cellular, molecular and genetic levels are only now revealing the true extent of this conservation and are also uncovering important differences. The broadest pattern of axis specification involves the generation of animal–vegetal polarity in the oocyte followed by the dorsal localization of determinants in the egg. This *ab ovo* specification mechanism operates throughout the primitive vertebrates, reptiles and possibly the egg-laying mammals. Based on various cell biological, genetic, and morphological observations, the basic pattern of axis generation is the involution of notochordal precursors at the early dorsal lip/organizer and the later ingression of ventrolateral precursors around the lateral blastopore, or “bilateral primitive streaks.”



This pattern exists throughout the anamniotes (ichthyosaurs) and the reptiles and is initially established by cortical rotation in the egg, leading to Wnt/beta-catenin asymmetry, elevated dorsal Nodal signaling and organizer induction and dorsal mesoderm involution (Fig. 6.14). Interestingly, much of this sequence of events is likely specific to the vertebrates. In basal invertebrate chordates, such as *Branchiostoma*, cortical rotation and early dorsal enrichment of beta-catenin do not occur and gastrulation occurs symmetrically via cell ingression, with minimal involution (Zhang et al. 1997; Holland and Onai 2012). Early Nodal asymmetry is present and likely leads to the induction of organizer/BMP antagonist gene expression on the dorsal side of the blastopore (Yu et al. 2007; Holland and Onai 2012). Thus, a key series of events in early vertebrate evolution was likely the appearance of cortical rotation, leading to the restriction of gastrulation initiation dorsally, coupled with a transition to cell involution behavior in the organizer.

In contrast to this early asymmetry mechanism, therian mammals undergo apparent *ab embryo* axis specification, with the early phase of Wnt asymmetry being lost and evolving into new mechanisms of symmetry breaking in the blastocyst, based on cell position and migration in the visceral endoderm. In both *ab ovo* and *ab embryo* cases, the egg and the blastocyst exhibit marked metastability, rendering development with radial symmetry improbable under normal conditions. The evolutionary transitions between these mechanisms are not well understood. Increases in yolk and egg size and the shift to meroblastic cleavage have been suggested to drive the evolution towards primitive streak-based gastrulation (Arendt and Nübler-Jung 1999), although these cannot be the sole factors involved. Reptiles have large eggs and undergo meroblastic cleavage but still form the organizer first and develop a horizontal slit blastopore/blastoporal plate, maintaining the ancestral dorsal involution/lateral ingression pattern. Additionally, teleosts also evolved meroblastic cleavage but this change resulted in additional adaptations to keep a blastoporal gastrulation pattern, rather than evolving primitive streak-like movements. Furthermore, the “primitive streak” evolved independently in birds and in mammals, possibly multiple times, which could explain the differences in cellular mechanisms underlying formation of the primitive streak, as well as observations such as the presence of left-right asymmetries around Hensen’s node and lack of nodal cilia in both chick and pig (Gros et al. 2009; Blum et al. 2014).

This convergent evolution toward the primitive streak could be correlated with changes in the development of the hypoblast/anterior endoderm. In amphibians and reptiles, these cells arise during cleavage from yolky vegetal cells, whereas in birds and mammals the hypoblast develops from cells delaminating or sorting out from the epiblast. Formation of the hypoblast in this manner could be a prerequisite for the emergent behavior of coordinated polonaise-like cell movements in the epiblast, which themselves are likely sufficient to position and initiate the primitive streak (Voiculescu et al. 2007). Additionally, owing to the more prominent role of hypoblast migration in birds and mammals, cell internalization initiates with the primitive streak, not with the organizer. Full organizer formation, in terms of gene expression and inducing activity, does not occur until mid-gastrulation. Thus, there is significant heterochrony in the main pattern of axis formation in birds and mam-

mals, with the appearance of the organizer and cell involution lagging behind that of ventrolateral cell ingression (Fig. 6.14). It is possible that the lessened importance of cortical rotation-like mechanisms led to subtle reorganization of the organizer gene regulatory networks, causing organizer genes such as *Noggin* to be expressed later relative to other dorsal genes.

Mammals exhibit the greatest changes in axis-forming mechanisms within the vertebrates. There is also wide variation in these mechanisms within the mammals themselves. Mammalian eggs have evolved to lose size and yolk content, as well as any notable animal–vegetal asymmetry and cortical rotation-related events in the process. The evolution of implantation also necessitated the early segregation of embryonic and extra-embryonic lineages and the maintenance of embryonic pluripotency. Furthermore, gastrulation in mammals initiates with far fewer cells than in other groups of organisms, ~250 cells in the mouse (compared with >10,000 in the frog and chicken), and is accompanied by rapid cell division. This rapid cell cycle of 2–6 h (Snow 1977) coincides with germ layer and axis patterning in the mammalian embryo, and has sparked evolutionary comparisons to the cleavage stage cell cycle in more primitive animals (O’Farrell et al. 2004). Whether this rapid division is indeed a conserved but temporally deferred event, as has been proposed, is unclear. Alternatively, there may exist a general correlation between pluripotency and shorter G1/G2 phases, with limited cell cycle checkpoints, as is seen in cultured stem cells (Becker et al. 2006).

What are some future prospects for the study of axis formation? After many decades of research, the ideas of symmetry breaking and self-organization in development remain compelling yet incompletely understood. Indeed, the ideas of self-organization and biological “regulation” predate the discovery of the organizer and have been developed into the concepts of the self-regulating embryonic “field” and pattern formation by positional information (Child 1915; Weiss 1926; Nieuwkoop 1967; Wolpert 1969, 1971; Green and Sharpe 2015). The ability of the organizer to undergo self-organization was noted by Spemann and others and clearly implies a complex network of interactions. Recent work showing inverse transcriptional regulation within the BMP regulatory circuitry has shed some light onto these processes. This observation is only the first step toward understanding self-organization at the gene regulatory network level, and additional general principles are likely to be identified. Although the conserved transcriptional control of genes like *Gooseoid* has long been noted, these mechanisms have not necessarily been fully characterized or compared. Remarkably, large and robust organizer gene regulatory networks have not yet been elucidated for any vertebrate organism. It should eventually be possible to generate comparative organizer gene regulatory networks across vertebrates and quantitatively compare their properties. Such analyses might shed light on possible changes in relative onset of organizer activity and also provide insight into the control of cell behaviors during gastrulation, as there have been several evolutionary transitions between involution and ingression in similar tissues.

Additionally, with regard to the complex interactions underlying symmetry breaking events in early vertebrate development, the molecular basis of microtubule orga-

nization during cortical rotation, the prime example of a symmetry-breaking process, remains largely unknown. The cell movements underlying asymmetry in the mammalian blastocyst are similarly mysterious. Both of these areas would likely benefit from advances in live imaging technology that are rendering the molecular activities of cells and tissues more visible and quantifiable. In particular, new methods for culturing and imaging different vertebrate embryos (e.g., Bedzhov and Zernicka-Goetz 2014; Keller et al. 2008) will likely provide novel insights into morphogenesis.

There have recently been rapid advances in whole genome analysis and genome editing technology that will likely transform the study of developmental biology and biology in general. Large-scale gene expression, chromatin analysis and proteomics are becoming commonplace and should yield abundant material for data science analysis. It is probably not a stretch to imagine that real-time whole transcriptome analyses will also become technologically feasible in living cells in the near future. TALEN and Cas9/CRISPR-mediated genome editing are also on the verge of becoming routine, extending mutational analysis to many areas previously intractable. Thus, in the foreseeable future, a variety of “-omics” data should be readily available for any organism, and investigators will have the capability to interrogate the function of any genomic region and immediately read out responses in gene expression and cell behavior.

The lack of detailed genomic and genetic information and methodologies has often been a barrier to work in many nontraditional model organisms. However, newer technologies should soon allow the genetic analysis of axis formation and other processes in organisms rationally chosen based on phylogenetic position or other criteria based on the biological question. Indeed multiple related species could be analyzed in parallel to capture extant variation in developmental mechanisms. Such ideas are not new (Tzika and Milinkovitch 2008) but are much closer to realization. Some interesting candidates for developmental analysis could include tree shrews, owing to their similarity to primates, and echidnas, as an egg-laying mammal model (Tzika and Milinkovitch 2008). The increased availability of human embryonic stem cells is also leading to a better understanding of human development. Gene regulatory networks can be analyzed in embryonic stem cells and induced-pluripotent stem cells and tissue “organoids” are increasingly being used to understand human organ development (e.g., McCracken et al. 2014). Certain ethical considerations would have to be navigated however to study human axis formation in this way. Comparative studies of early development across all vertebrates could be facilitated by reviving the idea of a “standardized vertebrate normal table” (Witschi 1956; Hopwood 2007). Such a standard series could be based on conserved gene expression data, gene regulatory network organization and morphogenetic patterns. Potential in-roads to such a project have already been made with the construction of comparative drawings based on gene expression patterns, the “Molecular Haeckels” (Elinson and Kezmoh 2010).

It is perhaps ironic that just as long-awaited sequenced genomes and targeted genome editing technologies are becoming routinely available in organisms like fish and amphibians, interest in the basic science of development in these organisms is perceived to be waning in favor of human-centered translational research. It remains

to be seen whether this perception is borne out in the long term. An optimist might argue that the processes of asymmetry, axis formation, and self-organization have fascinated scientists and have generated fundamental biological insight for well over 100 years, and should continue to do so long into the future.

**Acknowledgements** The author would like to thank B. Fritsch for assistance with German translations. This work was supported by the University of Iowa.

## References

- Aberle H, Bauer A, Stappert J et al (1997) Beta-catenin is a target for the ubiquitin-proteasome pathway. *EMBO J* 16:3797–3804. doi:[10.1093/emboj/16.13.3797](https://doi.org/10.1093/emboj/16.13.3797)
- Acampora D, Mazan S, Lallemand Y et al (1995) Forebrain and midbrain regions are deleted in *Otx2*<sup>-/-</sup> mutants due to a defective anterior neuroectoderm specification during gastrulation. *Development* 121:3279–3290
- Adams RJ, Kimmel CB (2004) Morphogenetic cellular flows during zebrafish gastrulation. In: Stern CD (ed) *Gastrulation: from cells to embryo*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp 305–316
- Agalliu D, Takada S, Agalliu I et al (2009) Motor neurons with axial muscle projections specified by Wnt4/5 signaling. *Neuron* 61:708–720. doi:[10.1016/j.neuron.2008.12.026](https://doi.org/10.1016/j.neuron.2008.12.026)
- Agius E, Oelgeschlager M, Wessely O et al (2000) Endodermal Nodal-related signals and mesoderm induction in *Xenopus*. *Development* 127:1173–1183
- Ahumada A, Slusarski DC, Liu X et al (2002) Signaling of rat Frizzled-2 through phosphodiesterase and cyclic GMP. *Science* 298:2006–2010. doi:[10.1126/science.1073776](https://doi.org/10.1126/science.1073776)
- Ambrosio AL, Taelman VF, Lee HX et al (2008) Crossveinless-2 is a BMP feedback inhibitor that binds Chordin/BMP to regulate *Xenopus* embryonic patterning. *Dev Cell* 15:248–260. doi:[10.1016/j.devcel.2008.06.013](https://doi.org/10.1016/j.devcel.2008.06.013)
- Amit S, Hatzubai A, Birman Y et al (2002) Axin-mediated CKI phosphorylation of beta-catenin at Ser 45: a molecular switch for the Wnt pathway. *Genes Dev* 16:1066–1076. doi:[10.1101/gad.230302](https://doi.org/10.1101/gad.230302)
- Ancel P, Vintemberger P (1948) Recherches sur le determinisme de la symetrie bilaterale dans l’oeuf des Amphibiens. *Bull Biol Fr Belg (Suppl)* 31:1–182
- Antic, D., Stubbs, J. L., Suyama, K., Kintner, C., Scott, M. P. and Axelrod, J. D. (2010) Planar Cell Polarity Enables Posterior Localization of Nodal Cilia and Left-Right Axis Determination during Mouse and *Xenopus* Embryogenesis. *PLoS ONE* 5, e8999.
- Arendt D, Nubler-Jung K (1999) Rearranging gastrulation in the name of yolk: evolution of gastrulation in yolk-rich amniote eggs. *Mech Dev* 81:3–22
- Ault KT, Dirksen ML, Jamrich M (1996) A novel homeobox gene PV.1 mediates induction of ventral mesoderm in *Xenopus* embryos. *Proc Natl Acad Sci U S A* 93:6415–6420
- Axelrod JD, Miller JR, Shulman JM et al (1998) Differential recruitment of Dishevelled provides signaling specificity in the planar cell polarity and Wingless signaling pathways. *Genes Dev* 12:2610–2622
- Azar Y, Eyal-Giladi H (1981) Interaction of epiblast and hypoblast in the formation of the primitive streak and the embryonic axis in chick, as revealed by hypoblast-rotation experiments. *J Embryol Exp Morphol* 61:133–144
- Bachiller D, Klingensmith J, Kemp C et al (2000) The organizer factors Chordin and Noggin are required for mouse forebrain development. *Nature* 403:658–661. doi:[10.1038/35001072](https://doi.org/10.1038/35001072)
- Bachvarova RF, Skromne I, Stern CD (1998) Induction of primitive streak and Hensen’s node by the posterior marginal zone in the early chick embryo. *Development* 125:3521–3534

- von Baer KE (1828) Über Entwicklungsgeschichte der Thiere. Beobachtung und Reflexion. Gebrüder Bornträger, Königsberg
- Baker J, Beddington R, Harland R (1999) Wnt signaling in *Xenopus* embryos inhibits bmp4 expression and activates neural development. *Genes Dev* 13:3149–3159
- Barker N, Morin P, Clevers H (2000) The Yin-Yang of TCF/beta-catenin signaling. *Adv Cancer Res* 77:1–24
- Barrott JJ, Cash GM, Smith AP et al (2011) Deletion of mouse *Porcn* blocks Wnt ligand secretion and reveals an ectodermal etiology of human focal dermal hypoplasia/Goltz syndrome. *Proc Natl Acad Sci U S A* 108:12752–12757. doi:[10.1073/pnas.1006437108](https://doi.org/10.1073/pnas.1006437108)
- Bartscherer K, Pelte N, Ingelfinger D, Boutros M (2006) Secretion of Wnt ligands requires Evi, a conserved transmembrane protein. *Cell* 125:523–533. doi:[10.1016/j.cell.2006.04.009](https://doi.org/10.1016/j.cell.2006.04.009)
- Bayly R, Axelrod JD (2011) Pointing in the right direction: new developments in the field of planar cell polarity. *Nat Rev Genet* 12:385–391. doi:[10.1038/nrg2956](https://doi.org/10.1038/nrg2956)
- Bánki Ö (1927) Die Langebeziehung der Spermiumeintrittsstelle zur Medianebene und zur ersten Furche nach Versuchen mit örtlicher Vitälfärbung am Axolotl Ei. *Verh. Anat. Ges., Jena*, 36: 198–208
- Becker KA, Ghule PN, Therrien JA et al (2006) Self-renewal of human embryonic stem cells is supported by a shortened G1 cell cycle phase. *J Cell Physiol* 209:883–893. doi:[10.1002/jcp.20776](https://doi.org/10.1002/jcp.20776)
- Beddington RS (1994) Induction of a second neural axis by the mouse node. *Development* 120:613–620
- Bedzhov I, Graham SJL, Leung CY, Zernicka-Goetz M (2014) Developmental plasticity, cell fate specification and morphogenesis in the early mouse embryo. *Philos Trans R Soc B Biol Sci* 369:20130538. doi:[10.1098/rstb.2013.0538](https://doi.org/10.1098/rstb.2013.0538)
- Bedzhov I, Zernicka-Goetz M (2014) Self-organizing properties of mouse pluripotent cells initiate morphogenesis upon implantation. *Cell* 156:1032–1044. doi:[10.1016/j.cell.2014.01.023](https://doi.org/10.1016/j.cell.2014.01.023)
- Belenkaya TY, Han C, Standley HJ et al (2002) *pygopus* Encodes a nuclear protein essential for wingless/Wnt signaling. *Development* 129:4089–4101
- Belo JA, Bouwmeester T, Leyns L et al (1997) Cerberus-like is a secreted factor with neutralizing activity expressed in the anterior primitive endoderm of the mouse gastrula. *Mech Dev* 68:45–57
- Ben-Haim N, Lu C, Guzman-Ayala M et al (2006) The nodal precursor acting via activin receptors induces mesoderm by maintaining a source of its convertases and BMP4. *Dev Cell* 11:313–323. doi:[10.1016/j.devcel.2006.07.005](https://doi.org/10.1016/j.devcel.2006.07.005)
- ten Berge D, Kurek D, Blauwkamp T et al (2011) Embryonic stem cells require Wnt proteins to prevent differentiation to epiblast stem cells. *Nat Cell Biol* 13:1070–1075. doi:[10.1038/ncb2314](https://doi.org/10.1038/ncb2314)
- Bertocchini F, Alev C, Nakaya Y, Sheng G (2013) A little winning streak: the reptilian-eye view of gastrulation in birds. *Dev Growth Differ* 55:52–59. doi:[10.1111/dgd.12014](https://doi.org/10.1111/dgd.12014)
- Bertocchini F, Skromne I, Wolpert L, Stern CD (2004) Determination of embryonic polarity in a regulative system: evidence for endogenous inhibitors acting sequentially during primitive streak formation in the chick embryo. *Development* 131:3381–3390. doi:[10.1242/dev.01178](https://doi.org/10.1242/dev.01178)
- Bertocchini F, Stern CD (2002) The hypoblast of the chick embryo positions the primitive streak by antagonizing nodal signaling. *Dev Cell* 3:735–744
- Bertocchini F, Stern CD (2012) *Gata2* provides an early anterior bias and uncovers a global positioning system for polarity in the amniote embryo. *Development* 139:4232–4238. doi:[10.1242/dev.081901](https://doi.org/10.1242/dev.081901)
- Betchaku T, Trinkaus JP (1978) Contact relations, surface activity, and cortical microfilaments of marginal cells of the enveloping layer and of the yolk syncytial and yolk cytoplasmic layers of fundulus before and during epiboly. *J Exp Zool* 206:381–426. doi:[10.1002/jez.1402060310](https://doi.org/10.1002/jez.1402060310)
- Biechele S, Cockburn K, Lanner F et al (2013) *Porcn*-dependent Wnt signaling is not required prior to mouse gastrulation. *Development* 140:2961–2971. doi:[10.1242/dev.094458](https://doi.org/10.1242/dev.094458)
- Biechele S, Cox BJ, Rossant J (2011) Porcupine homolog is required for canonical Wnt signaling and gastrulation in mouse embryos. *Dev Biol* 355:275–285. doi:[10.1016/j.ydbio.2011.04.029](https://doi.org/10.1016/j.ydbio.2011.04.029)

- Bilic J, Huang Y-L, Davidson G et al (2007) Wnt induces LRP6 signalosomes and promotes dishevelled-dependent LRP6 phosphorylation. *Science* 316:1619–1622. doi:[10.1126/science.1137065](https://doi.org/10.1126/science.1137065)
- Bin-Nun N, Lichtig H, Malyarova A et al (2014) PTK7 modulates Wnt signaling activity via LRP6. *Development* 141:410–421. doi:[10.1242/dev.095984](https://doi.org/10.1242/dev.095984)
- Birsoy B, Kofron M, Schaible K et al (2006) Vg 1 is an essential signaling molecule in *Xenopus* development. *Development* 133:15–20. doi:[10.1242/dev.02144](https://doi.org/10.1242/dev.02144)
- Blitz IL, Cho K W Y (2009) Finding partners: how BMPs select their targets. *Dev Dyn* 238:1321–1331. doi:[10.1002/dvdy.21984](https://doi.org/10.1002/dvdy.21984)
- Blum M, Andre P, Muders K, et al. (2007) Ciliation and gene expression distinguish between node and posterior notochord in the mammalian embryo. *Differentiation*
- Blum M, Feistel K, Thumberger T, Schweickert A (2014) The evolution and conservation of left-right patterning mechanisms. *Development* 141:1603–1613. doi:[10.1242/dev.100560](https://doi.org/10.1242/dev.100560)
- Blum M, Gaunt SJ, Cho KW et al (1992) Gastrulation in the mouse: the role of the homeobox gene goosecoid. *Cell* 69:1097–1106
- Blythe SA, Cha S-W, Tadjuidje E et al (2010) Beta-catenin primes organizer gene expression by recruiting a histone H3 arginine 8 methyltransferase, Prmt2. *Dev Cell* 19:220–231. doi:[10.1016/j.devcel.2010.07.007](https://doi.org/10.1016/j.devcel.2010.07.007)
- Boterenbrood EC, Nieuwkoop PD (1973) The formation of the mesoderm in urodelean amphibians. *Wilhelm Roux' Arch Entwicklunsgmech* 173:319–332. doi:[10.1007/BF00575837](https://doi.org/10.1007/BF00575837)
- Boucaut JC, Darribere T, De Li S, Boulekbache H, Yamada KM, Thiery JP (1985) Evidence for the role of fibronectin in amphibian gastrulation. *Development* 89:211–227
- Bourillot PY, Garrett N, Gurdon JB (2002) A changing morphogen gradient is interpreted by continuous transduction flow. *Development* 129:2167–2180
- Boutros M, Mlodzik M (1999) Dishevelled: at the crossroads of divergent intracellular signaling pathways. *Mech Dev* 83:27–37
- Bouwmeester T, Kim S-H, Sasai Y et al (1996) Cerberus is a head-inducing secreted factor expressed in the anterior endoderm of Spemann's organizer. *Nature* 382:595–601. doi:[10.1038/382595a0](https://doi.org/10.1038/382595a0)
- Brachet A (1904) Recherches expérimentales sur l'oeuf de *Rana fusca*. *Arch Biol* 21:103–160
- Brannon M, Gomperts M, Sumoy L et al (1997) A beta-catenin/XTcf-3 complex binds to the siamois promoter to regulate dorsal axis specification in *Xenopus*. *Genes Dev* 11:2359–2370
- Brannon M, Kimelman D (1996) Activation of Siamois by the Wnt pathway. *Dev Biol* 180:344–347
- Brennan J, Lu CC, Norris DP et al (2001) Nodal signalling in the epiblast patterns the early mouse embryo. *Nature* 411:965–969. doi:[10.1038/35082103](https://doi.org/10.1038/35082103)
- Briehier WM, Gumbiner BM (1994) Regulation of C-cadherin function during activin induced morphogenesis of *Xenopus* animal caps. *J Cell Biol* 126:519–527
- Brown A, Papkoff J, Fung Y et al (1987) Identification of protein products encoded by the proto-oncogene int-1. *Mol Cell Biol* 7:3971–3977
- Brunner E, Peter O, Schweizer L, Basler K (1997) pangolin encodes a Lef-1 homologue that acts downstream of Armadillo to transduce the Wingless signal in *Drosophila*. *Nature* 385:829–833
- Bytynski-Salz H (1937) Trapianti di organizzatore nelle uova di Lampreda. Tip Luigi Niccolai
- Callebaut M (2005) Origin, fate, and function of the components of the avian germ disc region and early blastoderm: role of ooplasmic determinants. *Dev Dyn* 233:1194–1216. doi:[10.1002/dvdy.20493](https://doi.org/10.1002/dvdy.20493)
- Callebaut M, Van Nueten E (1994) Rauber's (Koller's) sickle: the early gastrulation organizer of the avian blastoderm. *Eur J Morphol* 32:35–48
- Callebaut M, Van Nueten E (1995) Gastrulation inducing potencies of endophyll and Rauber's sickle in isolated caudocranially oriented prestreak avian blastoderm quadrants (or fragments) in vitro. *Eur J Morphol* 33:221–235
- Camp E, Sánchez-Sánchez AV, García-España A et al (2009) Nanog regulates proliferation during early fish development. *Stem Cells* 27:2081–2091. doi:[10.1002/stem.133](https://doi.org/10.1002/stem.133)
- Campbell PD, Heim AE, Smith MZ, Marlow FL (2015) Kinesin-1 interacts with Bucky ball to form germ cells and is required to pattern the zebrafish body axis. *Development* 142:2996–3008. doi:[10.1242/dev.124586](https://doi.org/10.1242/dev.124586)

- Camus A, Perea-Gomez A, Moreau A, Collignon J (2006) Absence of Nodal signaling promotes precocious neural differentiation in the mouse embryo. *Dev Biol* 295:743–755. doi:[10.1016/j.ydbio.2006.03.047](https://doi.org/10.1016/j.ydbio.2006.03.047)
- Cha S-W, Tadjuidje E, Tao Q et al (2008) Wnt5a and Wnt11 interact in a maternal Dkk1-regulated fashion to activate both canonical and non-canonical signaling in *Xenopus* axis formation. *Development* 135:3719–3729. doi:[10.1242/dev.029025](https://doi.org/10.1242/dev.029025)
- Cha S-W, Tadjuidje E, White J et al (2009) Wnt11/5a complex formation caused by tyrosine sulfation increases canonical signaling activity. *Curr Biol* 19:1573–1580. doi:[10.1016/j.cub.2009.07.062](https://doi.org/10.1016/j.cub.2009.07.062)
- Chambers SM, Fasano CA, Papapetrou EP et al (2009) Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat Biotechnol* 27:275–280. doi:[10.1038/nbt.1529](https://doi.org/10.1038/nbt.1529)
- Chamorro MN, Schwartz DR, Vonica A et al (2005) FGF-20 and DKK1 are transcriptional targets of beta-catenin and FGF-20 is implicated in cancer and development. *EMBO J* 24:73–84. doi:[10.1038/sj.emboj.7600460](https://doi.org/10.1038/sj.emboj.7600460)
- Chan AP, Kloc M, Larabell CA et al (2007) The maternally localized RNA fatvg is required for cortical rotation and germ cell formation. *Mech Dev* 124:350–363. doi:[10.1016/j.mod.2007.02.001](https://doi.org/10.1016/j.mod.2007.02.001)
- Chang C, Harland RM (2007) Neural induction requires continued suppression of both Smad1 and Smad2 signals during gastrulation. *Development* 134:3861–3872. doi:[10.1242/dev.007179](https://doi.org/10.1242/dev.007179)
- Chang C, Hemmati-Brivanlou A (1998) Cell fate determination in embryonic ectoderm. *J Neurobiol* 36:128–151
- Chazaud C, Rossant J (2006) Disruption of early proximodistal patterning and AVE formation in *Apc* mutants. *Development* 133:3379–3387. doi:[10.1242/dev.02523](https://doi.org/10.1242/dev.02523)
- Cheng AM, Thisse B, Thisse C, Wright CV (2000) The lefty-related factor *Xatv* acts as a feedback inhibitor of nodal signaling in mesoderm induction and L-R axis development in *Xenopus*. *Development* 127:1049–1061
- Cheung KJ, Gabrielson E, Werb Z, Ewald AJ (2013) Collective invasion in breast cancer requires a conserved basal epithelial program. *Cell* 155:1639–1651. doi:[10.1016/j.cell.2013.11.029](https://doi.org/10.1016/j.cell.2013.11.029)
- Child CM (1915) Individuality in organisms (Google Books). The University of Chicago Press, Chicago, IL
- Cho KW, Blumberg B, Steinbeisser H, De Robertis EM (1991) Molecular nature of Spemann's organizer: the role of the *Xenopus* homeobox gene goosecoid. *Cell* 67:1111–1120
- Christian J, Moon R (1993) Interactions between *Xwnt-8* and Spemann organizer signaling pathways generate dorsoventral pattern in the embryonic mesoderm of *Xenopus*. *Genes Dev* 7:13–28
- Christian JL, McMahon JA, McMahon AP, Moon RT (1991) *Xwnt-8*, a *Xenopus* Wnt-1/int-1-related gene responsive to mesoderm-inducing growth factors, may play a role in ventral mesodermal patterning during embryogenesis. *Development* 111:1045–1055
- Chung HM, Malacinski GM (1980) Establishment of the dorsal/ventral polarity of the amphibian embryo: use of ultraviolet irradiation and egg rotation as probes. *Dev Biol* 80:120–133
- Ciemerych MA, Mesnard D, Zernicka-Goetz M (2000) Animal and vegetal poles of the mouse egg predict the polarity of the embryonic axis, yet are nonessential for development. *Development* 127:3467–3474
- Ciruna B, Jenny A, Lee D et al (2006) Planar cell polarity signalling couples cell division and morphogenesis during neurulation. *Nat Cell Biol* 439:220–224. doi:[10.1038/nature04375](https://doi.org/10.1038/nature04375)
- Clavert J (1962) Symmetrization of the egg of vertebrates. *Adv Morphol* 2:27–60
- Clavert J (1961) Développement de la symétrie chez les Vertébrés. *Bull Soc Zool Fr* 86:381–401
- Clevers H, Nusse R (2012) Wnt/ $\beta$ -catenin signaling and disease. *Cell* 149:1192–1205. doi:[10.1016/j.cell.2012.05.012](https://doi.org/10.1016/j.cell.2012.05.012)
- Cliffe A, Hamada F, Bienz M (2003) A role of Dishevelled in relocating Axin to the plasma membrane during wingless signaling. *Curr Biol* 13:960–966
- Cole MF, Johnstone SE, Newman JJ et al (2008) Tcf3 is an integral component of the core regulatory circuitry of embryonic stem cells. *Genes Dev* 22:746–755. doi:[10.1101/gad.1642408](https://doi.org/10.1101/gad.1642408)

- Collavin L, Kirschner MW (2003) The secreted Frizzled-related protein Sizzled functions as a negative feedback regulator of extreme ventral mesoderm. *Development* 130:805–816
- Colozza G, De Robertis EM (2014) Maternal syntabulin is required for dorsal axis formation and is a germ plasm component in *Xenopus*. *Differentiation* 88:17–26. doi:[10.1016/j.diff.2014.03.002](https://doi.org/10.1016/j.diff.2014.03.002)
- Conlon FL, Lyons KM, Takaesu N et al (1994) A primary requirement for nodal in the formation and maintenance of the primitive streak in the mouse. *Development* 120:1919–1928
- Connolly DJ, Patel K, Cooke J (1997) Chick noggin is expressed in the organizer and neural plate during axial development, but offers no evidence of involvement in primary axis formation. *Int J Dev Biol* 41:389–396
- Cook D, Fry MJ, Hughes K et al (1996) Wingless inactivates glycogen synthase kinase-3 via an intracellular signalling pathway which involves a protein kinase C. *EMBO J* 15:4526–4536
- Cooke J, Smith JC (1987) The midblastula cell cycle transition and the character of mesoderm in UV-induced nonaxial *Xenopus* development. *Development* 99:197–210
- Cooper MS, D'Amico LA (1996) A cluster of noninvoluting endocytic cells at the margin of the zebrafish blastoderm marks the site of embryonic shield formation. *Dev Biol* 180:184–198. doi:[10.1006/dbio.1996.0294](https://doi.org/10.1006/dbio.1996.0294)
- Coudreuse DYM (2006) Wnt gradient formation requires retromer function in Wnt-producing cells. *Science* 312:921–924. doi:[10.1126/science.1124856](https://doi.org/10.1126/science.1124856)
- Cruciat C-M, Niehrs C (2013) Secreted and transmembrane wnt inhibitors and activators. *Cold Spring Harb Perspect Biol* 5:a015081. doi:[10.1101/cshperspect.a015081](https://doi.org/10.1101/cshperspect.a015081)
- Csardi G, Nepusz T (2014) igraph: Network analysis and visualization. R package version 0.7
- Cselenyi CS, Jernigan KK, Tahinci E et al (2008) LRP6 transduces a canonical Wnt signal independently of Axin degradation by inhibiting GSK3's phosphorylation of beta-catenin. *Proc Natl Acad Sci U S A* 105:8032–8037. doi:[10.1073/pnas.0803025105](https://doi.org/10.1073/pnas.0803025105)
- Cui Y, Tian Q, Christian JL (1996) Synergistic effects of Vg1 and Wnt signals in the specification of dorsal mesoderm and endoderm. *Dev Biol* 180:22–34. doi:[10.1006/dbio.1996.0281](https://doi.org/10.1006/dbio.1996.0281)
- Curtin JA, Quint E, Tsipouri V et al (2003) Mutation of Celsr1 disrupts planar polarity of inner ear hair cells and causes severe neural tube defects in the mouse. *Curr Biol* 13:1129–1133
- Cuykendall TN, Houston DW (2009) Vegetally localized *Xenopus* trim36 regulates cortical rotation and dorsal axis formation. *Development* 136:3057–3065. doi:[10.1242/dev.036855](https://doi.org/10.1242/dev.036855)
- Dalcq A, Pasteels J (1937) Une conception nouvelle des bases physiologiques de la morphogénèse. *Arch Biol (Liege)* 48:699–710
- Dale L, Howes G, Price B, Smith J (1992) Bone morphogenetic protein 4: a ventralizing factor in early *Xenopus* development. *Development* 115:573–585
- Dale L, Smith JC, Slack JM (1985) Mesoderm induction in *Xenopus laevis*: a quantitative study using a cell lineage label and tissue-specific antibodies. *J Embryol Exp Morphol* 89:289–312
- Dalle Nogare D, Somers K, Rao S et al (2014) Leading and trailing cells cooperate in collective migration of the zebrafish posterior lateral line primordium. *Development* 141:3188–3196. doi:[10.1242/dev.106690](https://doi.org/10.1242/dev.106690)
- Darken R, Scola A, Rakeman A et al (2002) The planar polarity gene strabismus regulates convergent extension movements in *Xenopus*. *EMBO J* 21:976–985
- Darras S, Marikawa Y, Elinson RP, Lemaire P (1997) Animal and vegetal pole cells of early *Xenopus* embryos respond differently to maternal dorsal determinants: implications for the patterning of the organiser. *Development* 124:4275–4286
- Davidson G, Shen J, Huang Y-L et al (2009) Cell cycle control of wnt receptor activation. *Dev Cell* 17:788–799. doi:[10.1016/j.devcel.2009.11.006](https://doi.org/10.1016/j.devcel.2009.11.006)
- Davidson G, Wu W, Shen J et al (2005) Casein kinase I gamma couples Wnt receptor activation to cytoplasmic signal transduction. *Nature* 438:867–872. doi:[10.1038/nature04170](https://doi.org/10.1038/nature04170)
- De Robertis E, Larrain J, Oelgeschlager M, Wessely O (2000) The establishment of Spemann's organizer and patterning of the vertebrate embryo. *Nat Rev Genet* 1:171–181
- De Robertis EM (2009) Spemann's organizer and the self-regulation of embryonic fields. *Mech Dev* 126:925–941. doi:[10.1016/j.mod.2009.08.004](https://doi.org/10.1016/j.mod.2009.08.004)



- De Robertis EM (2006) Spemann's organizer and self-regulation in amphibian embryos. *Nat Rev Mol Cell Biol* 7:296–302. doi:[10.1038/nrm1855](https://doi.org/10.1038/nrm1855)
- De Robertis EM, Sasai Y (1996) A common plan for dorsoventral patterning in Bilateria. *Nature* 380:37–40. doi:[10.1038/380037a0](https://doi.org/10.1038/380037a0)
- De Vries WN, Evsikov AV, Haac BE et al (2004) Maternal beta-catenin and E-cadherin in mouse development. *Development* 131:4435–4445. doi:[10.1242/dev.01316](https://doi.org/10.1242/dev.01316)
- Deans MR, Antic D, Suyama K et al (2007) Asymmetric distribution of Prickle-like 2 reveals an early underlying polarization of vestibular sensory epithelia in the inner ear. *J Neurosci* 27:3139–3147. doi:[10.1523/JNEUROSCI.5151-06.2007](https://doi.org/10.1523/JNEUROSCI.5151-06.2007)
- Delaune E, Lemaire P, Kodjabachian L (2005) Neural induction in *Xenopus* requires early FGF signalling in addition to BMP inhibition. *Development* 132:299–310. doi:[10.1242/dev.01582](https://doi.org/10.1242/dev.01582)
- Dick A, Hild M, Bauer H, Imai Y, Maifeld H, Schier AF, Talbot WS, Bouwmeester T, Hammerschmidt M (2000) Essential role of Bmp7 (snailhouse) and its prodomain in dorsoventral patterning of the zebrafish embryo. *Development* 127:343–354
- Ding J, Yang L, Yan YT et al (1998) Cripto is required for correct orientation of the anterior-posterior axis in the mouse embryo. *Nature* 395:702–707. doi:[10.1038/27215](https://doi.org/10.1038/27215)
- Dirksen ML, Jamrich M (1992) A novel, activin-inducible, blastopore lip-specific gene of *Xenopus laevis* contains a fork head DNA-binding domain. *Genes Dev* 6:599–608
- Djiane A, Riou J, Umbhauer M et al (2000) Role of frizzled 7 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. *Development* 127:3091–3100
- Dobrowolski R, De Robertis EM (2012) Endocytic control of growth factor signalling: multivesicular bodies as signalling organelles. *Nat Rev Mol Cell Biol* 13:53–60. doi:[10.1038/nrm3244](https://doi.org/10.1038/nrm3244)
- Doi JY, Niigaki H, Sone K et al (2000) Distribution of dorsal-forming activity in precleavage embryos of the Japanese newt, *Cynops pyrrhogaster*: effects of deletion of vegetal cytoplasm, UV irradiation, and lithium treatment. *Dev Biol* 223:154–168. doi:[10.1006/dbio.2000.9735](https://doi.org/10.1006/dbio.2000.9735)
- Domingos PM, Itasaki N, Jones CM et al (2001) The Wnt/beta-catenin pathway posteriorizes neural tissue in *Xenopus* by an indirect mechanism requiring FGF signalling. *Dev Biol* 239:148–160. doi:[10.1006/dbio.2001.0431](https://doi.org/10.1006/dbio.2001.0431)
- Dorsky RI, Itoh M, Moon RT, Chitnis A (2003) Two tcf3 genes cooperate to pattern the zebrafish brain. *Development* 130:1937–1947
- Dougan ST, Warga RM, Kane DA et al (2003) The role of the zebrafish nodal-related genes *squint* and *cyclops* in patterning of mesendoderm. *Development* 130:1837–1851
- Du SJ, Purcell SM, Christian JL et al (1995) Identification of distinct classes and functional domains of Wnts through expression of wild-type and chimeric proteins in *Xenopus* embryos. *Mol Cell Biol* 15:2625–2634
- Dufort D, Schwartz L, Harpal K, Rossant J (1998) The transcription factor HNF3beta is required in visceral endoderm for normal primitive streak morphogenesis. *Development* 125:3015–3025
- Dyson S, Gurdon JB (1998) The interpretation of position in a morphogen gradient as revealed by occupancy of activin receptors. *Cell* 93:557–568
- Egger-Adam D, Katanaev VL (2010) The trimeric G protein Go inflicts a double impact on axin in the Wnt/frizzled signaling pathway. *Dev Dyn* 239:168–183. doi:[10.1002/dvdy.22060](https://doi.org/10.1002/dvdy.22060)
- Elinson R, Pasceri P (1989) Two UV-sensitive targets in dorsoanterior specification of frog embryos. *Development* 106:511–518
- Elinson RP (1983) Cytoplasmic phases in the first cell cycle of the activated frog egg. *Dev Biol* 100:440–451
- Elinson RP, del Pino EM (2011) Developmental diversity of amphibians. *WIREs Dev Biol* 1:345–369. doi:[10.1002/wdev.23](https://doi.org/10.1002/wdev.23)
- Elinson RP, Holowacz T (1995) Specifying the dorsoanterior axis in frogs: 70 years since Spemann and Mangold. *Curr Top Dev Biol* 30:253–285
- Elinson RP, Kezmoh L (2010) Molecular Haeckel. *Dev Dyn* 239:1905–1918. doi:[10.1002/dvdy.22337](https://doi.org/10.1002/dvdy.22337)
- Elinson RP, Rowning B (1988) A transient array of parallel microtubules in frog eggs: potential tracks for a cytoplasmic rotation that specifies the dorso-ventral axis. *Dev Biol* 128:185–197

- Erter CE, Solnica-Krezel L, Wright CV (1998) Zebrafish nodal-related 2 encodes an early mesodermal inducer signaling from the extraembryonic yolk syncytial layer. *Dev Biol* 204:361–372. doi:[10.1006/dbio.1998.9097](https://doi.org/10.1006/dbio.1998.9097)
- Eyal-Giladi H (1954) Dynamic aspects of neural induction in amphibia. *Arch Biol (Liege)* 65:179–259
- Eyal-Giladi H, Fabian BC (1980) Axis determination in uterine chick blastodiscs under changing spatial positions during the sensitive period for polarity. *Dev Biol* 77:228–232
- Eyal-Giladi H, Kochav S (1976) From cleavage to primitive streak formation: a complementary normal table and a new look at the first stages of the development of the chick: I. General morphology. *Dev Biol* 49:321–337
- Fainsod A, Steinbeisser H, De Robertis E (1994) On the function of BMP-4 in patterning the marginal zone of the *Xenopus* embryo. *EMBO J* 13:5015–5025
- Faure S, Lee MA, Keller T et al (2000) Endogenous patterns of TGFbeta superfamily signaling during early *Xenopus* development. *Development* 127:2917–2931
- Fekany K, Yamanaka Y, Leung T, Sirotkin HI, Topczewski J, Gates MA, Hibi M, Renucci A, Stemple D, Radbill A et al (1999) The zebrafish bozozok locus encodes Dharma, a homeodomain protein essential for induction of gastrula organizer and dorsoanterior embryonic structures. *Development* 126:1427–1438
- Feldman B, Gates MA, Egan ES et al (1998) Zebrafish organizer development and germ-layer formation require nodal-related signals. *Nature* 395:181–185. doi:[10.1038/26013](https://doi.org/10.1038/26013)
- Fiedler M, Sánchez-Barrena MJ, Nekrasov M et al (2008) Decoding of methylated histone H3 tail by the Pygo-BCL9 Wnt signaling complex. *Mol Cell* 30:507–518. doi:[10.1016/j.molcel.2008.03.011](https://doi.org/10.1016/j.molcel.2008.03.011)
- Flowers GP, Topczewska JM, Topczewski J (2012) A zebrafish Notum homolog specifically blocks the Wnt/ $\beta$ -catenin signaling pathway. *Development* 139:2416–2425. doi:[10.1242/dev.063206](https://doi.org/10.1242/dev.063206)
- Foley AC, Skromne I, Stern CD (2000) Reconciling different models of forebrain induction and patterning: a dual role for the hypoblast. *Development* 127:3839–3854
- Fossat N, Jones V, Khoo P-L et al (2011) Stringent requirement of a proper level of canonical WNT signalling activity for head formation in mouse embryo. *Development* 138:667–676. doi:[10.1242/dev.052803](https://doi.org/10.1242/dev.052803)
- Fraser SE, Stern CD (2004) Early rostrocaudal patterning of the mesoderm and neural plate. In: Stern CD (ed) *Gastrulation*. CSHL Press, New York, pp 389–401
- Friedl P, Gilmour D (2009) Collective cell migration in morphogenesis, regeneration and cancer. *Nat Rev Mol Cell Biol* 10:445–457. doi:[10.1038/nrm2720](https://doi.org/10.1038/nrm2720)
- Fu J, Jiang M, Mirando AJ et al (2009) Reciprocal regulation of Wnt and Gpr177/mouse Wntless is required for embryonic axis formation. *Proc Natl Acad Sci U S A* 106:18598–18603. doi:[10.1073/pnas.0904894106](https://doi.org/10.1073/pnas.0904894106)
- Fuentealba LC, Eivers E, Ikeda A et al (2007) Integrating patterning signals: Wnt/GSK3 regulates the duration of the BMP/Smad1 signal. *Cell* 131:980–993. doi:[10.1016/j.cell.2007.09.027](https://doi.org/10.1016/j.cell.2007.09.027)
- Fujimori T, Kurotaki Y, Miyazaki J-I, Nabeshima Y-I (2003) Analysis of cell lineage in two- and four-cell mouse embryos. *Development* 130:5113–5122. doi:[10.1242/dev.00725](https://doi.org/10.1242/dev.00725)
- Fujisue M, Kobayakawa Y, Yamana K (1993) Occurrence of dorsal axis-inducing activity around the vegetal pole of an uncleaved *Xenopus* egg and displacement to the equatorial region by cortical rotation. *Development* 118:163–170
- Fujisue M, Sakai M, Yamana K (1991) Subcortical rotation and specification of the dorsoventral axis in newt eggs. *Dev Growth Differ* 33:34–51
- Funayama N, Fagotto F, McCrea P, Gumbiner BM (1995) Embryonic axis induction by the armadillo repeat domain of beta-catenin: evidence for intracellular signaling. *J Cell Biol* 128:959–968
- Furushima K, Yamamoto A, Nagano T et al (2007) Mouse homologues of Shisa antagonistic to Wnt and Fgf signalings. *Dev Biol* 306:480–492. doi:[10.1016/j.ydbio.2007.03.028](https://doi.org/10.1016/j.ydbio.2007.03.028)
- Galceran J, Fariñas I, Depew MJ et al (1999) Wnt3a<sup>-/-</sup>-like phenotype and limb deficiency in *Lef1(-/-)Tcf1(-/-)* mice. *Genes Dev* 13:709–717

- Gao B, Song H, Bishop K et al (2011) Wnt signaling gradients establish planar cell polarity by inducing Vangl2 phosphorylation through Ror2. *Dev Cell* 20:163–176. doi:[10.1016/j.devcel.2011.01.001](https://doi.org/10.1016/j.devcel.2011.01.001)
- Gardner RL, Cockcroft DL (1998) Complete dissipation of coherent clonal growth occurs before gastrulation in mouse epiblast. *Development* 125:2397–2402
- Gawantka V, Delius H, Hirschfeld K et al (1995) Antagonizing the Spemann organizer: role of the homeobox gene *Xvent-1*. *EMBO J* 14:6268–6279
- Ge X, Grotjahn D, Welch E et al (2014) Hecate/Grip2a acts to reorganize the cytoskeleton in the symmetry-breaking event of embryonic axis induction. *PLoS Genet* 10:e1004422. doi:[10.1371/journal.pgen.1004422](https://doi.org/10.1371/journal.pgen.1004422)
- Gerhart J (2004) *Symmetry breaking in the egg of Xenopus laevis*. 341–351. Cold Spring Harbor, New York: CSHL Press
- Gerhart J, Danilchik M, Doniach T et al (1989) Cortical rotation of the *Xenopus* egg: consequences for the anteroposterior pattern of embryonic dorsal development. *Development* 107(Suppl):37–51
- Gerhart J, Ubbels G, Black S et al (1981) A reinvestigation of the role of the grey crescent in axis formation in *Xenopus laevis*. *Nature* 292:511–516
- Gerlitz O, Basler K (2002) Wingful, an extracellular feedback inhibitor of Wingless. *Genes Dev* 16:1055–1059. doi:[10.1101/gad.991802](https://doi.org/10.1101/gad.991802)
- Gimlich RL (1986) Acquisition of developmental autonomy in the equatorial region of the *Xenopus* embryo. *Dev Biol* 115:340–352
- Gimlich RL, Gerhart JC (1984) Early cellular interactions promote embryonic axis formation in *Xenopus laevis*. *Dev Biol* 104:117–130
- Giraldez AJ, Copley RR, Cohen SM (2002) HSPG modification by the secreted enzyme Notum shapes the Wingless morphogen gradient. *Dev Cell* 2:667–676
- Glinka A, Delius H, Blumenstock C, Niehrs C (1996) Combinatorial signalling by *Xwnt-11* and *Xnr3* in the organizer epithelium. *Mech Dev* 60:221–231
- Glinka A, Wu W, Delius H et al (1998) Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction. *Nature* 391:357–362. doi:[10.1038/34848](https://doi.org/10.1038/34848)
- Glinka A, Wu W, Onichtchouk D et al (1997) Head induction by simultaneous repression of *Bmp* and *Wnt* signalling in *Xenopus*. *Nature* 389:517–519. doi:[10.1038/39092](https://doi.org/10.1038/39092)
- Godsave SF, Slack JM (1991) Single cell analysis of mesoderm formation in the *Xenopus* embryo. *Development* 111:523–530
- Gore AV, Maegawa S, Cheong A et al (2005) The zebrafish dorsal axis is apparent at the four-cell stage. *Nature* 438:1030–1035. doi:[10.1038/nature04184](https://doi.org/10.1038/nature04184)
- Goto T, Keller R (2002) The planar cell polarity gene *strabismus* regulates convergence and extension and neural fold closure in *Xenopus*. *Dev Biol* 247:165–181
- Grädl D, König A, Wedlich D (2002) Functional diversity of *Xenopus* lymphoid enhancer factor/T-cell factor transcription factors relies on combinations of activating and repressing elements. *J Biol Chem* 277:14159–14171. doi:[10.1074/jbc.M107055200](https://doi.org/10.1074/jbc.M107055200)
- Graff JM, Thies RS, Song JJ et al (1994) Studies with a *Xenopus* BMP receptor suggest that ventral mesoderm-inducing signals override dorsal signals in vivo. *Cell* 79:169–179
- Granier C, Gurchenkov V, Perea-Gomez A et al (2011) Nodal cis-regulatory elements reveal epiblast and primitive endoderm heterogeneity in the peri-implantation mouse embryo. *Dev Biol* 349:350–362. doi:[10.1016/j.ydbio.2010.10.036](https://doi.org/10.1016/j.ydbio.2010.10.036)
- Gray H (1918) *Anatomy of the human body*, 20th edn. Lea and Febiger, Philadelphia, PA
- Gräper L (1929) Die Primitiventwicklung des Hühnchens nach stereokinematographischen Untersuchungen, kontrolliert durch vitale Farbmarkierung und verglichen mit der Entwicklung anderer Wirbeltiere. *Wilhelm Roux' Arch Entwicklungsmech* 116:382–429. doi:[10.1007/BF02145235](https://doi.org/10.1007/BF02145235)
- Green JBA, Sharpe J (2015) Positional information and reaction-diffusion: two big ideas in developmental biology combine. *Development* 142:1203–1211. doi:[10.1242/dev.114991](https://doi.org/10.1242/dev.114991)
- Gritsman K, Talbot WS, Schier AF (2000) Nodal signaling patterns the organizer. *Development* 127:921–932

- Gros J, Feistel K, Viebahn C et al (2009) Cell movements at Hensen's node establish left/right asymmetric gene expression in the chick. *Science* 324:941–944. doi:[10.1126/science.1172478](https://doi.org/10.1126/science.1172478)
- Gross JC, Chaudhary V, Bartscherer K, Boutros M (2012) Active Wnt proteins are secreted on exosomes. *Nat Cell Biol* 14:1036–1045. doi:[10.1038/ncb2574](https://doi.org/10.1038/ncb2574)
- Grunz H (2004) *The vertebrate organizer*. Springer Science & Business Media, Berlin
- Grunz H, Tacke L (1989) Neural differentiation of *Xenopus laevis* ectoderm takes place after disaggregation and delayed reaggregation without inducer. *Cell Differ Dev* 28:211–217
- Grunz H, Tacke L (1990) Extracellular matrix components prevent neural differentiation of disaggregated *Xenopus* ectoderm cells. *Cell Differ Dev* 32:117–123
- Guger KA, Gumbiner BM (1995) beta-Catenin has Wnt-like activity and mimics the Nieuwkoop signaling center in *Xenopus* dorsal-ventral patterning. *Dev Biol* 172:115–125. doi:[10.1006/dbio.1995.0009](https://doi.org/10.1006/dbio.1995.0009)
- Habas R, Dawid IB, He X (2003) Coactivation of Rac and Rho by Wnt/Frizzled signaling is required for vertebrate gastrulation. *Genes Dev* 17:295–309. doi:[10.1101/gad.1022203](https://doi.org/10.1101/gad.1022203)
- Habas R, Kato Y, He X (2001) Wnt/Frizzled activation of Rho regulates vertebrate gastrulation and requires a novel Formin homology protein Daam1. *Cell* 107:843–854
- Haegel H, Larue L, Ohsugi M et al (1995) Lack of beta-catenin affects mouse development at gastrulation. *Development* 121:3529–3537
- Hainski AM, Moody SA (1992) *Xenopus* maternal RNAs from a dorsal animal blastomere induce a secondary axis in host embryos. *Development* 116:347–355
- Halacheva V, Fuchs M, Dönitz J et al (2011) Planar cell movements and oriented cell division during early primitive streak formation in the mammalian embryo. *Dev Dyn* 240:1905–1916. doi:[10.1002/dvdy.22687](https://doi.org/10.1002/dvdy.22687)
- Hamburger V (1988) *The heritage of experimental embryology: Hans Spemann and the organizer*. Oxford University Press, USA
- Hamburger V, Hamilton HL (1951) A series of normal stages in the development of the chick embryo. *J Morphol* 88:231–272. doi:[10.1002/aja.1001950404](https://doi.org/10.1002/aja.1001950404)
- Hamilton FS, Wheeler GN, Hoppler S (2001) Difference in XTcf-3 dependency accounts for change in response to beta-catenin-mediated Wnt signalling in *Xenopus* blastula. *Development* 128:2063–2073
- Hammerschmidt M, Pelegri F, Mullins MC et al (1996a) Mutations affecting morphogenesis during gastrulation and tail formation in the zebrafish, *Danio rerio*. *Development* 123:143–151
- Hammerschmidt M, Serbedzija GN, McMahon AP (1996b) Genetic analysis of dorsoventral pattern formation in the zebrafish: requirement of a BMP-like ventralizing activity and its dorsal repressor. *Genes Dev* 10:2452–2461. doi:[10.1101/gad.10.19.2452](https://doi.org/10.1101/gad.10.19.2452)
- Hansen C, Marion C, Steele K et al (1997) Direct neural induction and selective inhibition of mesoderm and epidermis inducers by Xnr3. *Development* 124:483–492
- Harland R (2008) Induction into the Hall of Fame: tracing the lineage of Spemann's organizer. *Development* 135:3321–3323
- Harland R, Gerhart J (1997) Formation and function of Spemann's organizer. *Annu Rev Cell Dev Biol* 13:611–667
- Harland RM (1994) Neural induction in *Xenopus*. *Curr Opin Genet Dev* 4:543–549
- Harland RM, Grainger RM (2011) *Xenopus* research: metamorphosed by genetics and genomics. *Trends Genet* 27:507–515. doi:[10.1016/j.tig.2011.08.003](https://doi.org/10.1016/j.tig.2011.08.003)
- Hartley KO, Hardcastle Z, Friday RV et al (2001) Transgenic *Xenopus* embryos reveal that anterior neural development requires continued suppression of BMP signaling after gastrulation. *Dev Biol* 238:168–184. doi:[10.1006/dbio.2001.0398](https://doi.org/10.1006/dbio.2001.0398)
- Hashiguchi M, Mullins MC (2013) Anteroposterior and dorsoventral patterning are coordinated by an identical patterning clock. *Development* 140:1970–1980. doi:[10.1242/dev.088104](https://doi.org/10.1242/dev.088104)
- Hashimoto H, Itoh M, Yamanaka Y, Yamashita S, Shimizu T, Solnica-Krezel L, Hibi M, Hirano T (2000) Zebrafish Dkk1 functions in forebrain specification and axial mesendoderm formation. *Dev Biol* 217:138–152
- Hashimoto M, Shinohara K, Wang J et al (2010) Planar polarization of node cells determines the rotational axis of node cilia. *Nat Cell Biol* 12:170–176. doi:[10.1038/ncb2020](https://doi.org/10.1038/ncb2020)

- Hausen P, Riebesell M (1991) The early development of *Xenopus laevis*: an atlas of the histology. Springer, New York
- Hayes M, Naito M, Daulat A et al (2013) Ptk7 promotes non-canonical Wnt/PCP-mediated morphogenesis and inhibits Wnt/ $\beta$ -catenin-dependent cell fate decisions during vertebrate development. *Development* 140(8):1807–1818
- He X, Saint-Jeannet JP, Wang Y et al (1997) A member of the Frizzled protein family mediating axis induction by Wnt-5A. *Science* 275:1652–1654
- He X, Semenov M, Tamai K, Zeng X (2004) LDL receptor-related proteins 5 and 6 in Wnt/ $\beta$ -catenin signaling: arrows point the way. *Development* 131:1663–1677. doi:[10.1242/dev.01117](https://doi.org/10.1242/dev.01117)
- Heasman J (1997) Patterning the *Xenopus* blastula. *Development* 124:4179–4191
- Heasman J, Crawford A, Goldstone K et al (1994) Overexpression of cadherins and underexpression of  $\beta$ -catenin inhibit dorsal mesoderm induction in early *Xenopus* embryos. *Cell* 79:791–803
- Hedge TA, Mason I (2008) Expression of Shisa2, a modulator of both Wnt and Fgf signaling, in the chick embryo. *Int J Dev Biol* 52:81–85. doi:[10.1387/ijdb.072355th](https://doi.org/10.1387/ijdb.072355th)
- Heisenberg CP, Houart C, Take-Uchi M et al (2001) A mutation in the Gsk3-binding domain of zebrafish Masterblind/Axin1 leads to a fate transformation of telencephalon and eyes to dien-cephalon. *Genes Dev* 15:1427–1434. doi:[10.1101/gad.194301](https://doi.org/10.1101/gad.194301)
- Heisenberg CP, Tada M, Rauch GJ et al (2000) Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. *Nature* 405:76–81. doi:[10.1038/35011068](https://doi.org/10.1038/35011068)
- Hemmati-Brivanlou A, Kelly O, Melton DA (1994) Follistatin, an antagonist of activin, is expressed in the Spemann organizer and displays direct neuralizing activity. *Cell* 77:283–295
- Hemmati-Brivanlou A, Melton D (1997) Vertebrate embryonic cells will become nerve cells unless told otherwise. *Cell* 88(1):13–17
- Hemmati-Brivanlou A, Melton DA (1994) Inhibition of activin receptor signaling promotes neuralization in *Xenopus*. *Cell* 77:273–281
- Hernández AR, Klein AM, Kirschner MW (2012) Kinetic responses of  $\beta$ -catenin specify the sites of Wnt control. *Science* 338:1337–1340. doi:[10.1126/science.1228734](https://doi.org/10.1126/science.1228734)
- Hesiod, Evelyn-White HG (1914) Works and days. The Homeric Hymns and Homeric 174–ff
- Hikasa H, Ezan J, Itoh K et al (2010) Regulation of TCF3 by Wnt-dependent phosphorylation during vertebrate axis specification. *Dev Cell* 19:521–532. doi:[10.1016/j.devcel.2010.09.005](https://doi.org/10.1016/j.devcel.2010.09.005)
- Hikasa H, Sokol SY (2013) Wnt signaling in vertebrate axis specification. *Cold Spring Harb Perspect Biol* 5:a007955. doi:[10.1101/cshperspect.a007955](https://doi.org/10.1101/cshperspect.a007955)
- Hild M, Dick A, Rauch GJ et al (1999) The smad5 mutation somitabun blocks Bmp2b signaling during early dorsoventral patterning of the zebrafish embryo. *Development* 126:2149–2159
- Hillman N, Sherman MI, Graham C (1972) The effect of spatial arrangement on cell determination during mouse development. *J Embryol Exp Morphol* 28:263–278
- Hilton E, Rex M, Old R (2003) VegT activation of the early zygotic gene *Xnr5* requires lifting of Tcf-mediated repression in the *Xenopus* blastula. *Mech Dev* 120:1127–1138
- Ho C-Y, Houart C, Wilson SW, Stainier DYC (1999) A role for the extraembryonic yolk syncytial layer in patterning the zebrafish embryo suggested by properties of the hex gene. *Curr Biol* 9:1131–1134. doi:[10.1016/S0960-9822\(99\)80485-0](https://doi.org/10.1016/S0960-9822(99)80485-0)
- Ho RK (1992) Cell movements and cell fate during zebrafish gastrulation. *Dev Suppl* 65–73
- Hoffmann R, Städeli R, Basler K (2005) Pygopus and legless provide essential transcriptional coactivator functions to armadillo/ $\beta$ -catenin. *Curr Biol* 15:1207–1211. doi:[10.1016/j.cub.2005.05.054](https://doi.org/10.1016/j.cub.2005.05.054)
- Hofmann K (2000) A superfamily of membrane-bound O-acyltransferases with implications for wnt signaling. *Trends Biochem Sci* 25:111–112
- Hogan BL (1996) Bone morphogenetic proteins: multifunctional regulators of vertebrate development. *Genes Dev* 10:1580–1594
- Holland LZ, Onai T (2012) Early development of cephalochordates (amphioxus). *WIREs Dev Biol* 1:167–183. doi:[10.1002/wdev.11](https://doi.org/10.1002/wdev.11)
- Holowacz T, Elinson RP (1993) Cortical cytoplasm, which induces dorsal axis formation in *Xenopus*, is inactivated by UV irradiation of the oocyte. *Development* 119:277–285

- Holstein TW (2012) The evolution of the Wnt pathway. *Cold Spring Harb Perspect Biol* 4:a007922. doi:[10.1101/cshperspect.a007922](https://doi.org/10.1101/cshperspect.a007922)
- Holtfreter J (1944) Neural differentiation of ectoderm through exposure to saline solution. *J Exp Zool* 95:307–343. doi:[10.1002/jez.1400950303](https://doi.org/10.1002/jez.1400950303)
- Holwill S, Heasman J, Crawley C, Wylie CC (1987) Axis and germ line deficiencies caused by UV irradiation of *Xenopus* oocytes cultured in vitro. *Development* 100:735–743
- Hoodless PA, Pye M, Chazaud C et al (2001) FoxH1 (Fast) functions to specify the anterior primitive streak in the mouse. *Genes Dev* 15:1257–1271. doi:[10.1101/gad.881501](https://doi.org/10.1101/gad.881501)
- Hoppler S, Brown JD, Moon RT (1996) Expression of a dominant-negative Wnt blocks induction of MyoD in *Xenopus* embryos. *Genes Dev* 10:2805–2817
- Hoppler S, Moon RT (1998) BMP-2/-4 and Wnt-8 cooperatively pattern the *Xenopus* mesoderm. *Mech Dev* 71:119–129
- Hopwood N (2007) A history of normal plates, tables and stages in vertebrate embryology. *Int J Dev Biol* 51:1–26. doi:[10.1387/ijdb.062189nh](https://doi.org/10.1387/ijdb.062189nh)
- Houart C, Caneparo L, Heisenberg C et al (2002) Establishment of the telencephalon during gastrulation by local antagonism of Wnt signaling. *Neuron* 35:255–265
- Houliston E (1994) Microtubule translocation and polymerisation during cortical rotation in *Xenopus* eggs. *Development* 120:1213–1220
- Houliston E, Elinson RP (1991) Evidence for the involvement of microtubules, ER, and kinesin in the cortical rotation of fertilized frog eggs. *J Cell Biol* 114:1017–1028
- Houston DW (2012) Cortical rotation and messenger RNA localization in *Xenopus* axis formation. *WIREs Dev Biol* 1:371–388. doi:[10.1002/wdev.29](https://doi.org/10.1002/wdev.29)
- Houston DW (2013) Regulation of cell polarity and RNA localization in vertebrate oocytes. *Int Rev Cell Mol Biol* 306:127–185. doi:[10.1016/B978-0-12-407694-5.00004-3](https://doi.org/10.1016/B978-0-12-407694-5.00004-3)
- Houston DW, Kofron M, Resnik E et al (2002) Repression of organizer genes in dorsal and ventral *Xenopus* cells mediated by maternal XTcf3. *Development* 129:4015–4025
- Hsieh J-C, Lee L, Zhang L et al (2003) Mesd encodes an LRP5/6 chaperone essential for specification of mouse embryonic polarity. *Cell* 112:355–367
- Huelsken J, Vogel R, Brinkmann V et al (2000) Requirement for beta-catenin in anterior-posterior axis formation in mice. *J Cell Biol* 148:567–578
- Hume CR, Dodd J (1993) Cwnt-8C: a novel Wnt gene with a potential role in primitive streak formation and hindbrain organization. *Development* 119:1147–1160
- Hunt TE (1929) Hensen's node as an organiser in the formation of the chick embryo. *Anat Rec* 42:22
- Imai Y, Gates MA, Melby AE et al (2001) The homeobox genes *vox* and *vent* are redundant repressors of dorsal fates in zebrafish. *Development* 128:2407–2420
- Inomata H, Haraguchi T, Sasai Y (2008) Robust stability of the embryonic axial pattern requires a secreted scaffold for chordin degradation. *Cell* 134:854–865. doi:[10.1016/j.cell.2008.07.008](https://doi.org/10.1016/j.cell.2008.07.008)
- Inui M, Montagner M, Ben-Zvi D et al (2012) Self-regulation of the head-inducing properties of the Spemann organizer. *Proc Natl Acad Sci U S A* 109:15354–15359. doi:[10.1073/pnas.1203000109](https://doi.org/10.1073/pnas.1203000109)
- Ip CK, Fossat N, Jones V et al (2014) Head formation: OTX2 regulates Dkk1 and Lhx1 activity in the anterior mesendoderm. *Development* 141:3859–3867. doi:[10.1242/dev.114900](https://doi.org/10.1242/dev.114900)
- Ishibashi H, Matsumura N, Hanafusa H et al (2008) Expression of Siamois and Twin in the blastula Chordin/Noggin signaling center is required for brain formation in *Xenopus laevis* embryos. *Mech Dev* 125:58–66. doi:[10.1016/j.mod.2007.10.005](https://doi.org/10.1016/j.mod.2007.10.005)
- Ishitani T, Kishida S, Hyodo-Miura J et al (2003) The TAK1-NLK mitogen-activated protein kinase cascade functions in the Wnt-5a/Ca(2+) pathway to antagonize Wnt/beta-catenin signaling. *Mol Cell Biol* 23:131–139
- Ishitani T, Ninomiya-Tsuji J, Nagai S et al (1999) The TAK1-NLK-MAPK-related pathway antagonizes signalling between beta-catenin and transcription factor TCF. *Nature* 399:798–802
- Itasaki N, Jones CM, Mercurio S et al (2003) Wise, a context-dependent activator and inhibitor of Wnt signalling. *Development* 130:4295–4305
- Iwao Y, Yasumitsu K, Narihira M et al (1997) Changes in microtubule structures during the first cell cycle of physiologically polyspermic newt eggs. *Mol Reprod Dev* 47:210–221.

- Izpisua Belmonte JC, De Robertis EM, Storey KG, Stern CD (1993) The homeobox gene gooseoid and the origin of organizer cells in the early chick blastoderm. *Cell* 74:645–659
- Janda CY, Waghray D, Levin AM et al (2012) Structural basis of Wnt recognition by frizzled. *Science* 337:59–64. doi:[10.1126/science.1222879](https://doi.org/10.1126/science.1222879)
- Jernigan KK, Cselenyi CS, Thorne CA et al (2010) Gbetagamma activates GSK3 to promote LRP6-mediated beta-catenin transcriptional activity. *Sci Signal* 3:ra37. doi:[10.1126/scisignal.2000647](https://doi.org/10.1126/scisignal.2000647)
- Jessen JR, Topczewski J, Bingham S et al (2002) Zebrafish trilobite identifies new roles for Strabismus in gastrulation and neuronal movements. *Nat Cell Biol* 4:610–615. doi:[10.1038/ncb828](https://doi.org/10.1038/ncb828)
- Jesuthasan S, Stähle U (1997) Dynamic microtubules and specification of the zebrafish embryonic axis. *Curr Biol* 7:31–42
- Jiang J, Struhl G (1998) Regulation of the Hedgehog and Wingless signalling pathways by the F-box/WD40-repeat protein Slimb. *Nature* 391:493–496
- Jones C, Lyons K, Lapan P et al (1992) DVR-4 (bone morphogenetic protein-4) as a posterior-ventralizing factor in *Xenopus* mesoderm induction. *Development* 115:639–647
- Jones CM, Broadbent J, Thomas PQ et al (1999) An anterior signalling centre in *Xenopus* revealed by the homeobox gene XHex. *Curr Biol* 9:946–954
- Jose Maria Carvajal-Gonzalez MM (2014) Mechanisms of planar cell polarity establishment in *Drosophila*. *F1000Prime Reports*. doi:[10.12703/P6-98](https://doi.org/10.12703/P6-98)
- Joubin K, Stern CD (1999) Molecular interactions continuously define the organizer during the cell movements of gastrulation. *Cell* 98:559–571
- Jullien J, Gurdon JB (2005) Morphogen gradient interpretation by a regulated trafficking step during ligand-receptor transduction. *Genes Dev* 19:2682–2694. doi:[10.1101/gad.341605](https://doi.org/10.1101/gad.341605)
- Kadowaki T, Wilder E, Klingensmith J et al (1996) The segment polarity gene porcupine encodes a putative multitransmembrane protein involved in Wingless processing. *Genes Dev* 10:3116–3128
- Kagermeier-Schenk B, Wehner D, Ozhan-Kizil G et al (2011) Waif1/5 T4 inhibits Wnt/ $\beta$ -catenin signaling and activates noncanonical Wnt pathways by modifying LRP6 subcellular localization. *Dev Cell* 21:1129–1143. doi:[10.1016/j.devcel.2011.10.015](https://doi.org/10.1016/j.devcel.2011.10.015)
- Kageura H (1997) Activation of dorsal development by contact between the cortical dorsal determinant and the equatorial core cytoplasm in eggs of *Xenopus laevis*. *Development* 124:1543–1551
- Kageura H (1990) Spatial distribution of the capacity to initiate a secondary embryo in the 32-cell embryo of *Xenopus laevis*. *Dev Biol* 142:432–438
- Kakugawa S, Langton PF, Zebisch M et al (2015) Notum deacylates Wnt proteins to suppress signalling activity. *Nature* 519:187–192. doi:[10.1038/nature14259](https://doi.org/10.1038/nature14259)
- Kalisz M, Winzi M, Bisgaard HC, Serup P (2012) EVEN-SKIPPED HOMEBOX 1 controls human ES cell differentiation by directly repressing GOOSECOID expression. *Dev Biol* 362:94–103. doi:[10.1016/j.ydbio.2011.11.017](https://doi.org/10.1016/j.ydbio.2011.11.017)
- Katanaev V, Buestorf S (2009) Frizzled proteins are bona fide G protein-coupled receptors. *Nature Precedings* 1–19
- Kato M, Patel MS, Levasseur R et al (2002) Cbfa1-independent decrease in osteoblast proliferation, osteopenia, and persistent embryonic eye vascularization in mice deficient in Lrp5, a Wnt coreceptor. *J Cell Biol* 157:303–314. doi:[10.1083/jcb.200201089](https://doi.org/10.1083/jcb.200201089)
- Kawahara A, Wilm T, Solnica-Krezel L, Dawid IB (2000) Antagonistic role of vega1 and bozozok/dharma homeobox genes in organizer formation. *Proc Natl Acad Sci U S A* 97:12121–12126. doi:[10.1073/pnas.97.22.12121](https://doi.org/10.1073/pnas.97.22.12121)
- Keller PJ, Schmidt AD, Wittbrodt J, Stelzer EHK (2008) Reconstruction of zebrafish early embryonic development by scanned light sheet microscopy. *Science* 322:1065–1069
- Keller R, Danilchik M (1988) Regional expression, pattern and timing of convergence and extension during gastrulation of *Xenopus laevis*. *Development* 103:193–209
- Keller R, Davidson L, Edlund A et al (2000) Mechanisms of convergence and extension by cell intercalation. *Philos Trans R Soc Lond B Biol Sci* 355:897–922. doi:[10.1098/rstb.2000.0626](https://doi.org/10.1098/rstb.2000.0626)
- Keller R, Shook D (2004) Gastrulation in amphibians. In: Stern CD (ed) *Gastrulation: from cells to embryo*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp 171–203

- Keller R, Tibbetts P (1989) Mediolateral cell intercalation in the dorsal, axial mesoderm of *Xenopus laevis*. *Dev Biol* 131:539–549
- Keller RE (1986) The cellular basis of amphibian gastrulation. *Dev Biol* (NY) 2:241–327
- Keller RE (1975) Vital dye mapping of the gastrula and neurula of *Xenopus laevis*: I. Prospective areas and morphogenetic movements of the superficial layer. *Dev Biol* 42(2):222–241
- Keller RE, Danilchik M, Gimlich R, Shih J (1985) The function and mechanism of convergent extension during gastrulation of *Xenopus laevis*. *J Embryol Exp Morphol* 89(Suppl):185–209
- Kelly C, Chin AJ, Leatherman JL et al (2000) Maternally controlled (beta)-catenin-mediated signaling is required for organizer formation in the zebrafish. *Development* 127:3899–3911
- Kelly G, Greenstein P, Erezyilmaz D, Moon R (1995a) Zebrafish wnt8 and wnt8b share a common activity but are involved in distinct developmental pathways. *Development* 121:1787–1799
- Kelly GM, Erezyilmaz DF, Moon RT (1995b) Induction of a secondary embryonic axis in zebrafish occurs following the overexpression of beta-catenin. *Mech Dev* 53:261–273
- Kelly OG, Pinson KI, Skarnes WC (2004) The Wnt co-receptors Lrp5 and Lrp6 are essential for gastrulation in mice. *Development* 131:2803–2815. doi:[10.1242/dev.01137](https://doi.org/10.1242/dev.01137)
- Kelly SJ (1977) Studies of the developmental potential of 4- and 8-cell stage mouse blastomeres. *J Exp Zool* 200:365–376. doi:[10.1002/jez.1402000307](https://doi.org/10.1002/jez.1402000307)
- Kemler R, Hierholzer A, Kanzler B et al (2004) Stabilization of beta-catenin in the mouse zygote leads to premature epithelial-mesenchymal transition in the epiblast. *Development* 131:5817–5824. doi:[10.1242/dev.01458](https://doi.org/10.1242/dev.01458)
- Kemp C, Willems E, Abdo S et al (2005) Expression of all Wnt genes and their secreted antagonists during mouse blastocyst and postimplantation development. *Dev Dyn* 233:1064–1075. doi:[10.1002/dvdy.20408](https://doi.org/10.1002/dvdy.20408)
- Khokha M, Yeh J, Grammer T, Harland R (2005) Depletion of three BMP antagonists from Spemann's organizer leads to a catastrophic loss of dorsal structures. *Dev Cell* 8:401–411
- Kibar Z, Torban E, McDermid JR et al (2007) Mutations in VANGL1 associated with neural-tube defects. *N Engl J Med* 356:1432–1437. doi:[10.1056/NEJMoa060651](https://doi.org/10.1056/NEJMoa060651)
- Kibar Z, Vogan KJ, Groulx N et al (2001) Ltap, a mammalian homolog of *Drosophila* Strabismus/Van Gogh, is altered in the mouse neural tube mutant Loop-tail. *Nat Genet* 28:251–255. doi:[10.1038/90081](https://doi.org/10.1038/90081)
- Kiecker C, Niehrs C (2001) A morphogen gradient of Wnt/beta-catenin signalling regulates anteroposterior neural patterning in *Xenopus*. *Development* 128:4189–4201
- Kikkawa M, Takano K, Shinagawa A (1996) Location and behavior of dorsal determinants during first cell cycle in *Xenopus* eggs. *Development* 122:3687–3696
- Kilian B, Mansukoski H, Barbosa F et al (2003) The role of Ppt/Wnt5 in regulating cell shape and movement during zebrafish gastrulation. *Mech Dev* 120:467–476
- Kim CH, Oda T, Itoh M et al (2000) Repressor activity of Headless/Tcf3 is essential for vertebrate head formation. *Nature* 407:913–916. doi:[10.1038/35038097](https://doi.org/10.1038/35038097)
- Kim GH, Her JH, Han JK (2008) Ryk cooperates with Frizzled 7 to promote Wnt11-mediated endocytosis and is essential for *Xenopus laevis* convergent extension movements. *J Cell Biol* 182:1073–1082. doi:[10.1083/jcb.200710188](https://doi.org/10.1083/jcb.200710188)
- Kim SE, Huang H, Zhao M et al (2013) Wnt stabilization of beta-catenin reveals principles for morphogen receptor-scaffold assemblies. *Science* 340:867–870. doi:[10.1126/science.1232389](https://doi.org/10.1126/science.1232389)
- Kimmel CB, Warga RM, Schilling TF (1990) Origin and organization of the zebrafish fate map. *Development* 108:581–594
- Kimura C, Yoshinaga K, Tian E et al (2000) Visceral endoderm mediates forebrain development by suppressing posteriorizing signals. *Dev Biol* 225:304–321. doi:[10.1006/dbio.2000.9835](https://doi.org/10.1006/dbio.2000.9835)
- Kimura-Yoshida C, Nakano H, Okamura D et al (2005) Canonical Wnt signaling and its antagonist regulate anterior-posterior axis polarization by guiding cell migration in mouse visceral endoderm. *Dev Cell* 9:639–650. doi:[10.1016/j.devcel.2005.09.011](https://doi.org/10.1016/j.devcel.2005.09.011)
- Kinder SJ, Tsang TE, Wakamiya M et al (2001) The organizer of the mouse gastrula is composed of a dynamic population of progenitor cells for the axial mesoderm. *Development* 128:3623–3634



- Kinoshita N, Iioka H, Miyakoshi A, Ueno N (2003) PKC delta is essential for Dishevelled function in a noncanonical Wnt pathway that regulates *Xenopus* convergent extension movements. *Genes Dev* 17:1663–1676
- Kirschner M, Gerhart JC, Hara K, Ubbels GA (1980) Initiation of the cell cycle and establishment of bilateral symmetry in *Xenopus* eggs. *Symp Soc Dev Biol* 38:187–215
- Kishimoto Y, Lee KH, Zon L, Hammerschmidt M, Schulte-Merker S (1997) The molecular nature of zebrafish swirl: BMP2 function is essential during early dorsoventral patterning. *Development* 124:4457–4466
- Klein SL, Moody SA (2015) Early neural ectodermal genes are activated by siamois and twin during blastula stages. *Genesis* 53:308–320. doi:[10.1002/dvg.22854](https://doi.org/10.1002/dvg.22854)
- Kloc M (2009) Teachings from the egg: new and unexpected functions of RNAs. *Mol Reprod Dev* 76:922–932. doi:[10.1002/mrd.21043](https://doi.org/10.1002/mrd.21043)
- Knoetgen H, Viebahn C, Kessel M (1999) Head induction in the chick by primitive endoderm of mammalian, but not avian origin. *Development* 126:815–825
- Kochav S, Eyal-Giladi H (1971) Bilateral symmetry in chick embryo determination by gravity. *Science* 171:1027–1029
- Kofron M, Birsoy B, Houston D et al (2007) Wnt11/beta-catenin signaling in both oocytes and early embryos acts through LRP6-mediated regulation of axin. *Development* 134:503–513. doi:[10.1242/dev.02739](https://doi.org/10.1242/dev.02739)
- Kohn AD, Moon RT (2005) Wnt and calcium signaling:  $\beta$ -catenin-independent pathways. *Cell Calcium* 38:439–446. doi:[10.1016/j.ceca.2005.06.022](https://doi.org/10.1016/j.ceca.2005.06.022)
- Koos DS, Ho RK (1999) The *nieuwkoid/dharma* homeobox gene is essential for *bmp2b* repression in the zebrafish pregastrula. *Dev Biol* 215(2):190–207
- Kozmik Z, Holland LZ, Schubert M et al (2001) Characterization of amphioxus amphivent, an evolutionarily conserved marker for chordate ventral mesoderm. *Genesis* 29:172–179. doi:[10.1002/gene.1021](https://doi.org/10.1002/gene.1021)
- Kraft B, Berger CD, Wallkamm V et al (2012) Wnt-11 and Fz7 reduce cell adhesion in convergent extension by sequestration of PAPC and C-cadherin. *J Cell Biol* 198:695–709. doi:[10.1083/jcb.201110076](https://doi.org/10.1083/jcb.201110076)
- Kramps T, Peter O, Brunner E et al (2002) Wnt/wingless signaling requires BCL9/legless-mediated recruitment of pygopus to the nuclear beta-catenin-TCF complex. *Cell* 109:47–60
- Krasnow R, Adler P (1994) A single frizzled protein has a dual function in tissue polarity. *Development* 120:1883–1893
- Kreuger J, Perez L, Giraldez AJ, Cohen SM (2004) Opposing activities of Dally-like glypican at high and low levels of Wingless morphogen activity. *Dev Cell* 7:503–512. doi:[10.1016/j.devcel.2004.08.005](https://doi.org/10.1016/j.devcel.2004.08.005)
- Ku M, Melton DA (1993) *Xwnt-11*: a maternally expressed *Xenopus* wnt gene. *Development* 119:1161–1173
- Kuhl M, Sheldahl LC, Malbon CC, Moon RT (2000) Ca<sup>2+</sup>/calmodulin-dependent protein kinase II is stimulated by Wnt and Frizzled homologs and promotes ventral cell fates in *Xenopus*. *J Biol Chem* 275:12701–12711
- Kuroda H, Wessely O, De Robertis EM (2004) Neural induction in *Xenopus*: requirement for ectodermal and endomesodermal signals via Chordin, Noggin, beta-Catenin, and Cerberus. *PLoS Biol* 2:E92. doi:[10.1371/journal.pbio.0020092](https://doi.org/10.1371/journal.pbio.0020092)
- Kurth T, Hausen P (2000) Bottle cell formation in relation to mesodermal patterning in the *Xenopus* embryo. *Mech Dev* 97:117–131. doi:[10.1016/S0925-4773\(00\)00428-7](https://doi.org/10.1016/S0925-4773(00)00428-7)
- Kwon H-J, Bhat N, Sweet EM et al (2010) Identification of early requirements for preplacodal ectoderm and sensory organ development. *PLoS Genet* 6:e1001133. doi:[10.1371/journal.pgen.1001133](https://doi.org/10.1371/journal.pgen.1001133)
- Ladher R, Mohun TJ, Smith JC, Snape AM (1996) *Xom*: a *Xenopus* homeobox gene that mediates the early effects of BMP-4. *Development* 122:2385–2394
- Lamb TM, Knecht AK, Smith WC et al (1993) Neural induction by the secreted polypeptide noggin. *Science* 262:713–718

- Larabell C, Rowning B, Wells J et al (1996) Confocal microscopy analysis of living *Xenopus* eggs and the mechanism of cortical rotation. *Development* 122:1281–1289
- Larsen WJ, Schoenwolf GC, Bleyl SB et al (2009) Larsen's human embryology. Churchill Livingstone, Philadelphia, PA
- Lawrence P, Casal J, Struhl G (2002) Towards a model of the organisation of planar polarity and pattern in the *Drosophila* abdomen. *Development* 129:2749–2760
- Lawson KA, Meneses JJ, Pedersen RA (1991) Clonal analysis of epiblast fate during germ layer formation in the mouse embryo. *Development* 113:891–911
- Lawson KA, Pedersen RA (1992) Clonal analysis of cell fate during gastrulation and early neurulation in the mouse. *Ciba Found Symp* 165:3–21, discussion 21–26
- Lee E, Salic A, Kruger R et al (2003) The roles of APC and Axin derived from experimental and theoretical analysis of the Wnt pathway. *PLoS Biol* 1:E10
- Lee HX, Ambrosio AL, Reversade B, De Robertis EM (2006) Embryonic dorsal-ventral signaling: secreted frizzled-related proteins as inhibitors of tolloid proteinases. *Cell* 124:147–159. doi:10.1016/j.cell.2005.12.018
- Lee MA, Heasman J, Whitman M (2001) Timing of endogenous activin-like signals and regional specification of the *Xenopus* embryo. *Development* 128:2939–2952
- Lekven A, Thorpe C, Waxman J, Moon R (2001) Zebrafish *wnt8* encodes two *wnt8* proteins on a bicistronic transcript and is required for mesoderm and neurectoderm patterning. *Dev Cell* 1:103–114
- Lele Z, Nowak M, Hammerschmidt M (2001) Zebrafish *admp* is required to restrict the size of the organizer and to promote posterior and ventral development. *Dev Dyn* 222:681–687. doi:10.1002/dvdy.1222
- Lemaire P, Garrett N, Gurdon JB (1995) Expression cloning of *Siamois*, a *Xenopus* homeobox gene expressed in dorsal-vegetal cells of blastulae and able to induce a complete secondary axis. *Cell* 81:85–94
- Leung T, Bischof J, Söll I et al (2003a) *bozozok* directly represses *bmp2b* transcription and mediates the earliest dorsoventral asymmetry of *bmp2b* expression in zebrafish. *Development* 130:3639–3649
- Leung T, Söll I, Arnold SJ et al (2003b) Direct binding of *Lef1* to sites in the *boz* promoter may mediate pre-midblastula-transition activation of *boz* expression. *Dev Dyn* 228:424–432. doi:10.1002/dvdy.10408
- Lewis SL, Khoo P-L, Andrea De Young R et al (2007) Genetic interaction of *Gsc* and *Dkk1* in head morphogenesis of the mouse. *Mech Dev* 124:157–165. doi:10.1016/j.mod.2006.10.001
- Lewis SL, Khoo PL, De Young RA et al (2008) *Dkk1* and *Wnt3* interact to control head morphogenesis in the mouse. *Development* 135:1791–1801. doi:10.1242/dev.018853
- Leyns L, Bouwmeester T, Kim SH et al (1997) *Frzb-1* is a secreted antagonist of Wnt signaling expressed in the Spemann organizer. *Cell* 88:747–756
- Lim S, Kumari P, Gilligan P et al (2012) Dorsal activity of maternal *squint* is mediated by a non-coding function of the RNA. *Development* 139:2903–2915. doi:10.1242/dev.077081
- Linker C, Stern CD (2004) Neural induction requires BMP inhibition only as a late step, and involves signals other than FGF and Wnt antagonists. *Development* 131:5671–5681. doi:10.1242/dev.01445
- Linnemannstöns K, Ripp C, Honemann-Capito M et al (2014) The PTK7-related transmembrane proteins off-track and off-track 2 are co-receptors for *Drosophila* *Wnt2* required for male fertility. *PLoS Genet* 10:e1004443. doi:10.1371/journal.pgen.1004443
- Little SC, Mullins MC (2006) Extracellular modulation of BMP activity in patterning the dorsoventral axis. *Birth Defects Res C Embryo Today* 78:224–242. doi:10.1002/bdrc.20079
- Liu C, Kato Y, Zhang Z et al (1999a) *beta-Trcp* couples *beta-catenin* phosphorylation-degradation and regulates *Xenopus* axis formation. *Proc Natl Acad Sci U S A* 96:6273–6278
- Liu C, Li Y, Semenov M et al (2002) Control of *beta-catenin* phosphorylation/degradation by a dual-kinase mechanism. *Cell* 108:837–847
- Liu F, van den Broek O, Destrée O, Hoppler S (2005) Distinct roles for *Xenopus* *Tcf/Lef* genes in mediating specific responses to Wnt/*beta-catenin* signalling in mesoderm development. *Development* 132:5375–5385. doi:10.1242/dev.02152

- Liu G, Bafico A, Harris VK, Aaronson SA (2003) A novel mechanism for Wnt activation of canonical signaling through the LRP6 receptor. *Mol Cell Biol* 23:5825–5835
- Liu P, Wakamiya M, Shea MJ et al (1999b) Requirement for Wnt3 in vertebrate axis formation. *Nat Genet* 22:361–365. doi:[10.1038/11932](https://doi.org/10.1038/11932)
- Long WL (1983) The role of the yolk syncytial layer in determination of the plane of bilateral symmetry in the rainbow trout, *Salmo gairdneri* Richardson. *J Exp Zool* 228:91–97
- Lu F-I, Thisse C, Thisse B (2011) Identification and mechanism of regulation of the zebrafish dorsal determinant. *Proc Natl Acad Sci U S A* 108:15876–15880. doi:[10.1073/pnas.1106801108](https://doi.org/10.1073/pnas.1106801108)
- Lu X, Borchers AGM, Jolicœur C et al (2004) PTK7/CCK-4 is a novel regulator of planar cell polarity in vertebrates. *Nature* 430:93–98. doi:[10.1038/nature02677](https://doi.org/10.1038/nature02677)
- Lustig KD, Kroll KL, Sun EE, Kirschner MW (1996) Expression cloning of a *Xenopus* T-related gene (*Xombi*) involved in mesodermal patterning and blastopore lip formation. *Development* 122:4001–4012
- Luther WH (1935) Entwicklungsphysiologische Untersuchungen am Forellenkeim: die Rolle des Organisationszentrums bei der Entstehung der Embryonalanlage. *Biol Zentralbl* 55:114–137
- Lutz H (1949) Sur la production expérimentale de la polyembryone et de la monstruosité double chez les oiseaux. *Arch Anat Microsc Morphol Exp* 39:79–144
- Lyman Gingerich J, Westfall TA, Slusarski DC, Pelegri F (2005) *hecate*, a zebrafish maternal effect gene, affects dorsal organizer induction and intracellular calcium transient frequency. *Dev Biol* 286:427–439. doi:[10.1016/j.ydbio.2005.07.031](https://doi.org/10.1016/j.ydbio.2005.07.031)
- MacDonald BT, Tamai K, He X (2009) Wnt/beta-catenin signaling: components, mechanisms, and diseases. *Dev Cell* 17:9–26. doi:[10.1016/j.devcel.2009.06.016](https://doi.org/10.1016/j.devcel.2009.06.016)
- Malacinski GM, Benford H, Chung HM (1975) Association of an ultraviolet irradiation sensitive cytoplasmic localization with the future dorsal side of the amphibian egg. *J Exp Zool* 191:97–110. doi:[10.1002/jez.1401910110](https://doi.org/10.1002/jez.1401910110)
- Manes ME, Elinson RP (1980) Ultraviolet light inhibits grey crescent formation on the frog egg. Wilhelm Roux's *Arch Dev Biol* 189:73–76
- Manes ME, Elinson RP, Barbieri FD (1978) Formation of the amphibian grey crescent: effects of colchicine and cytochalasin B. Wilhelm Roux's *Arch Dev Biol* 185:99–104
- Mao J, Wang J, Liu B et al (2001) Low-density lipoprotein receptor-related protein-5 binds to Axin and regulates the canonical Wnt signaling pathway. *Mol Cell* 7:801–809
- Marchal L, Luxardi G, Thomé V, Kodjabachian L (2009) BMP inhibition initiates neural induction via FGF signaling and *Zic* genes. *Proc Natl Acad Sci U S A* 106:17437–17442. doi:[10.1073/pnas.0906352106](https://doi.org/10.1073/pnas.0906352106)
- Maretto S, Cordenonsi M, Dupont S et al (2003) Mapping Wnt/beta-catenin signaling during mouse development and in colorectal tumors. *Proc Natl Acad Sci U S A* 100:3299–3304. doi:[10.1073/pnas.0434590100](https://doi.org/10.1073/pnas.0434590100)
- Marikawa Y, Elinson RP (1999) Relationship of vegetal cortical dorsal factors in the *Xenopus* egg with the Wnt/beta-catenin signaling pathway. *Mech Dev* 89:93–102
- Marikawa Y, Li Y, Elinson RP (1997) Dorsal determinants in the *Xenopus* egg are firmly associated with the vegetal cortex and behave like activators of the Wnt pathway. *Dev Biol* 191:69–79. doi:[10.1006/dbio.1997.8710](https://doi.org/10.1006/dbio.1997.8710)
- Marlow F, Topczewski J, Sepich D, Solnica-Krezel L (2002) Zebrafish Rho kinase 2 acts downstream of Wnt11 to mediate cell polarity and effective convergence and extension movements. *Curr Biol* 12:876–884
- Marrari Y, Clarke EJ, Rouvière C, Houliston E (2003) Analysis of microtubule movement on isolated *Xenopus* egg cortices provides evidence that the cortical rotation involves dynein as well as Kinesin Related Proteins and is regulated by local microtubule polymerisation. *Dev Biol* 257:55–70
- Marrari Y, Rouvière C, Houliston E (2004) Complementary roles for dynein and kinesins in the *Xenopus* egg cortical rotation. *Dev Biol* 271:38–48. doi:[10.1016/j.ydbio.2004.03.018](https://doi.org/10.1016/j.ydbio.2004.03.018)
- Marrari Y, Terasaki M, Arrowsmith V, Houliston E (2000) Local inhibition of cortical rotation in *Xenopus* eggs by an anti-KRP antibody. *Dev Biol* 224:250–262. doi:[10.1006/dbio.2000.9773](https://doi.org/10.1006/dbio.2000.9773)
- Martello G, Zacchigna L, Inui M et al (2007) MicroRNA control of Nodal signalling. *Nature* 449:183–188. doi:[10.1038/nature06100](https://doi.org/10.1038/nature06100)

- Martindale MQ (2005) The evolution of metazoan axial properties. *Nat Rev Genet* 6:917–927. doi:[10.1038/nrg1725](https://doi.org/10.1038/nrg1725)
- Martinez Barbera JP, Clements M, Thomas P et al (2000) The homeobox gene *Hex* is required in definitive endodermal tissues for normal forebrain, liver and thyroid formation. *Development* 127:2433–2445
- Martyn U, Schulte-Merker S (2003) The ventralized *ogon* mutant phenotype is caused by a mutation in the zebrafish homologue of *Sizzled*, a secreted Frizzled-related protein. *Dev Biol* 260:58–67
- Marvin MJ, Di Rocco G, Gardiner A et al (2001) Inhibition of Wnt activity induces heart formation from posterior mesoderm. *Genes Dev* 15:316–327. doi:[10.1101/gad.855501](https://doi.org/10.1101/gad.855501)
- Maung S, Jenny A (2011) Planar cell polarity in *Drosophila*. *Organogenesis* 7(3):165–179
- McCracken KW, Catá EM, Crawford CM et al (2014) Modelling human development and disease in pluripotent stem-cell-derived gastric organoids. *Nature* 516:400–404. doi:[10.1038/nature13863](https://doi.org/10.1038/nature13863)
- McGrew LL, Hoppeler S, Moon RT (1997) Wnt and FGF pathways cooperatively pattern antero-posterior neural ectoderm in *Xenopus*. *Mech Dev* 69:105–114
- McMahon AP, Moon RT (1989) Ectopic expression of the proto-oncogene *int-1* in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* 58:1075–1084
- Mei W, Jin Z, Lai F et al (2013) Maternal *Dead-End1* is required for vegetal cortical microtubule assembly during *Xenopus* axis specification. *Development* 140:2334–2344. doi:[10.1242/dev.094748](https://doi.org/10.1242/dev.094748)
- Meinhardt H (2012) Turing's theory of morphogenesis of 1952 and the subsequent discovery of the crucial role of local self-enhancement and long-range inhibition. *Interface Focus* 2:407–416. doi:[10.1098/rsfs.2011.0097](https://doi.org/10.1098/rsfs.2011.0097)
- Melby AE, Beach C, Mullins M, Kimelman D (2000) Patterning the early zebrafish by the opposing actions of *bozozok* and *vox/vent*. *Dev Biol* 224(2):275–285
- Melton DA (1987) Translocation of a localized maternal mRNA to the vegetal pole of *Xenopus* oocytes. *Nature* 328:80–82. doi:[10.1038/328080a0](https://doi.org/10.1038/328080a0)
- Meneghini M, Ishitani T, Carter J et al (1999) MAP kinase and Wnt pathways converge to down-regulate an HMG-domain repressor in *Caenorhabditis elegans*. *Nature* 399:793–797
- Merrill BJ, Pasolli HA, Polak L et al (2004) *Tcf3*: a transcriptional regulator of axis induction in the early embryo. *Development* 131:263–274. doi:[10.1242/dev.00935](https://doi.org/10.1242/dev.00935)
- Mesnard D, Guzman-Ayala M, Constam DB (2006) *Nodal* specifies embryonic visceral endoderm and sustains pluripotent cells in the epiblast before overt axial patterning. *Development* 133:2497–2505. doi:[10.1242/dev.02413](https://doi.org/10.1242/dev.02413)
- Messenger NJ, Kabitschke C, Andrews R et al (2005) Functional specificity of the *Xenopus* T-domain protein *Brachyury* is conferred by its ability to interact with *Smad1*. *Dev Cell* 8:599–610. doi:[10.1016/j.devcel.2005.03.001](https://doi.org/10.1016/j.devcel.2005.03.001)
- Metcalfe C, Bienz M (2011) Inhibition of GSK3 by Wnt signalling—two contrasting models. *J Cell Sci* 124:3537–3544. doi:[10.1242/jcs.091991](https://doi.org/10.1242/jcs.091991)
- Mieszczanek J, de la Roche M, Bienz M (2008) A role of *Pygopus* as an anti-repressor in facilitating Wnt-dependent transcription. *Proc Natl Acad Sci U S A* 105:19324–19329. doi:[10.1073/pnas.0806098105](https://doi.org/10.1073/pnas.0806098105)
- Mikels AJ, Nusse R (2006) Purified *Wnt5a* protein activates or inhibits beta-catenin-TCF signaling depending on receptor context. *PLoS Biol* 4:e115. doi:[10.1371/journal.pbio.0040115](https://doi.org/10.1371/journal.pbio.0040115)
- Miller J, Rowling B, Larabell C et al (1999) Establishment of the dorsal-ventral axis in *Xenopus* embryos coincides with the dorsal enrichment of *dishevelled* that is dependent on cortical rotation. *J Cell Biol* 146:427–437
- Mintz B (1964) Gene expression in the morula stage of mouse embryos, as observed during development of *t12/t12* lethal mutants in vitro. *J Exp Zool* 157:267–272
- Mintzer KA, Lee MA, Runke G, Trout J, Whitman M, Mullins MC (2001) *Lost-a-fin* encodes a type I BMP receptor, *Alk8*, acting maternally and zygotically in dorsoventral pattern formation. *Development* 128:859–869

- Miura S, Mishina Y (2007) The DVE changes distal epiblast fate from definitive endoderm to neuroectoderm by antagonizing nodal signaling. *Dev Dyn* 236:1602–1610. doi:[10.1002/dvdy.21166](https://doi.org/10.1002/dvdy.21166)
- Mizuno T, Yamaha E, Kuroiwa A, Takeda H (1999) Removal of vegetal yolk causes dorsal deficiencies and impairs dorsal-inducing ability of the yolk cell in zebrafish. *Mech Dev* 81:51–63
- Mohamed OA, Clarke HJ, Dufort D (2004) Beta-catenin signaling marks the prospective site of primitive streak formation in the mouse embryo. *Dev Dyn* 231:416–424. doi:[10.1002/dvdy.20135](https://doi.org/10.1002/dvdy.20135)
- Molenaar M, van de Wetering M, Oosterwegel M et al (1996) XTcf-3 transcription factor mediates beta-catenin-induced axis formation in *Xenopus* embryos. *Cell* 86:391–399
- Montcouquiol M, Rachel RA, Lanford PJ et al (2003) Identification of *Vangl2* and *Scrb1* as planar polarity genes in mammals. *Nature* 423:173–177. doi:[10.1038/nature01618](https://doi.org/10.1038/nature01618)
- Montero JA (2005) Shield formation at the onset of zebrafish gastrulation. *Development* 132:1187–1198. doi:[10.1242/dev.01667](https://doi.org/10.1242/dev.01667)
- Moon RT, Campbell RM, Christian JL et al (1993) Xwnt-5A: a maternal Wnt that affects morphogenetic movements after overexpression in embryos of *Xenopus laevis*. *Development* 119:97–111
- Moos M, Wang S, Krinks M (1995) Anti-dorsalizing morphogenetic protein is a novel TGF-beta homolog expressed in the Spemann organizer. *Development* 121:4293–4301
- Morgan TH, Boring AM (1903) The relation of the first plane of cleavage and the grey crescent to the median plane of the embryo of the frog. *Wilhelm Roux' Arch Entwicklungsmech* 16:680–690. doi:[10.1007/BF02301271](https://doi.org/10.1007/BF02301271)
- Morgan TH, Tsuda U (1894) The orientation of the frog's egg. *Q J Microsc Sci* 35:373–405
- Morkel M, Huelsken J, Wakamiya M et al (2003) Beta-catenin regulates Cripto- and Wnt3-dependent gene expression programs in mouse axis and mesoderm formation. *Development* 130:6283–6294. doi:[10.1242/dev.00859](https://doi.org/10.1242/dev.00859)
- Morris SA, Grewal S, Barrios F et al (2012a) Dynamics of anterior-posterior axis formation in the developing mouse embryo. *Nat Commun* 3:673. doi:[10.1038/ncomms1671](https://doi.org/10.1038/ncomms1671)
- Morris SA, Guo Y, Zernicka-Goetz M (2012b) Developmental plasticity is bound by pluripotency and the Fgf and Wnt signaling pathways. *Cell Rep* 2:756–765. doi:[10.1016/j.celrep.2012.08.029](https://doi.org/10.1016/j.celrep.2012.08.029)
- Morris SA, Teo RTY, Li H et al (2010) Origin and formation of the first two distinct cell types of the inner cell mass in the mouse embryo. *Proc Natl Acad Sci U S A* 107:6364–6369. doi:[10.1073/pnas.0915063107](https://doi.org/10.1073/pnas.0915063107)
- Mukhopadhyay M, Shtrom S, Rodriguez-Esteban C et al (2001) Dickkopf1 is required for embryonic head induction and limb morphogenesis in the mouse. *Dev Cell* 1:423–434
- Murdoch JN, Doudney K, Paternotte C et al (2001) Severe neural tube defects in the loop-tail mouse result from mutation of *Lpp1*, a novel gene involved in floor plate specification. *Hum Mol Genet* 10:2593–2601
- Myers DC, Sepich DS, Solnica-Krezel L (2002) Bmp activity gradient regulates convergent extension during zebrafish gastrulation. *Dev Biol* 243:81–98. doi:[10.1006/dbio.2001.0523](https://doi.org/10.1006/dbio.2001.0523)
- Na J, Lykke-Andersen K, Torres-Padilla M-E, Zernicka-Goetz M (2007) Dishevelled proteins regulate cell adhesion in mouse blastocyst and serve to monitor changes in Wnt signaling. *Dev Biol* 302:40–49. doi:[10.1016/j.ydbio.2006.08.036](https://doi.org/10.1016/j.ydbio.2006.08.036)
- Nagai, H., Sezaki, M., Kakiguchi, K., Nakaya, Y., Lee, H. C., Ladher, R., Sasanami, T., Han, J. Y., Yonemura, S. and Sheng, G. (2015) Cellular analysis of cleavage-stage chick embryos reveals hidden conservation in vertebrate early development. *Development* 142, 1279–1286.
- Nagel M, Winklbauer R (1999) Establishment of substratum polarity in the blastocoel roof of the *Xenopus* embryo. *Development* 126:1975–1984
- Nakamura T, Mine N, Nakaguchi E et al (2006) Generation of robust left-right asymmetry in the mouse embryo requires a self-enhancement and lateral-inhibition system. *Dev Cell* 11:495–504. doi:[10.1016/j.devcel.2006.08.002](https://doi.org/10.1016/j.devcel.2006.08.002)
- Nakayama T, Snyder MA, Grewal SS et al (1998) *Xenopus* Smad8 acts downstream of BMP-4 to modulate its activity during vertebrate embryonic patterning. *Development* 125:857–867
- Newport G (1851) On the impregnation of the Ovum in the Amphibia. (First Series). *Philos Trans R Soc Lond B Biol Sci* 141:169–242

- Newport G (1854) Researches on the impregnation of the Ovum in the Amphibia; and on the early stages of development of the Embryo. (Third Series). *Philos Trans R Soc Lond B Biol Sci* 144:229–244
- Nguyen VH, Schmid B, Trout J et al (1998) Ventral and lateral regions of the zebrafish gastrula, including the neural crest progenitors, are established by a *bmp2b/swirl* pathway of genes. *Dev Biol* 199:93–110. doi:[10.1006/dbio.1998.8927](https://doi.org/10.1006/dbio.1998.8927)
- Niehrs C, Keller R, Cho KW, De Robertis EM (1993) The homeobox gene *goosecoid* controls cell migration in *Xenopus* embryos. *Cell* 72:491–503
- Nieuwkoop P, Nigtevecht G (1954) Neural activation and transformation in explants of competent ectoderm under the influence of fragments of anterior notochord in Urodeles. *J Embryol Exp Morphol* 2:175–193
- Nieuwkoop PD (1999) The neural induction process; its morphogenetic aspects. *Int J Dev Biol* 43:615–623
- Nieuwkoop PD (1952) Activation and organization of the central nervous system in amphibians. Part III. Synthesis of a new working hypothesis. *J Exp Zool* 120:83–108
- Nieuwkoop PD (1967) The “organization centre”: II. Field phenomena, their origin and significance. *Acta Biotheor* 17:151–177
- Nieuwkoop PD, Florschütz PA (1950) Quelques caractères spéciaux de la gastrulation et de la neurulation de l’oeuf de *Xenopus laevis*, Daud, et de quelques autres Anoures. *Arch Biol* 61:113–150
- Nieuwkoop PD, Sutasurya LA (1979) Primordial germ cells in the chordates: embryogenesis and phylogenesis
- Nikaido M, Tada M, Saji T, Ueno N (1997) Conservation of BMP signaling in zebrafish mesoderm patterning. *Mech Dev* 61:75–88
- Ninomiya H, Elinson RP, Winklbauser R (2004) Antero-posterior tissue polarity links mesoderm convergent extension to axial patterning. *Nature* 430:364–367. doi:[10.1038/nature02620](https://doi.org/10.1038/nature02620)
- Nishimura T, Honda H, Takeichi M (2012) Planar cell polarity links axes of spatial dynamics in neural-tube closure. *Cell* 149:1084–1097. doi:[10.1016/j.cell.2012.04.021](https://doi.org/10.1016/j.cell.2012.04.021)
- Nojima H, Rothhamel S, Shimizu T et al (2010) Syntabulin, a motor protein linker, controls dorsal determination. *Development* 137:923–933. doi:[10.1242/dev.046425](https://doi.org/10.1242/dev.046425)
- Nojima H, Shimizu T, Kim C-H et al (2004) Genetic evidence for involvement of maternally derived Wnt canonical signaling in dorsal determination in zebrafish. *Mech Dev* 121:371–386. doi:[10.1016/j.mod.2004.02.003](https://doi.org/10.1016/j.mod.2004.02.003)
- Nordstrom U, Jessell T, Edlund T (2002) Progressive induction of caudal neural character by graded Wnt signaling. *Nat Neurosci* 5:525–532
- Norris DP, Brennan J, Bikoff EK, Robertson EJ (2002) The *Foxh1*-dependent autoregulatory enhancer controls the level of Nodal signals in the mouse embryo. *Development* 129:3455–3468
- Nusse R, Varmus H (1982) Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. *Cell* 31:99–109
- O’Farrell PH, Stumpff J, Su TT (2004) Embryonic cleavage cycles: how is a mouse like a fly? *Curr Biol* 14:R35–R45
- Oelgeschläger M, Kuroda H, Reversade B, De Robertis EM (2003) Chordin is required for the Spemann organizer transplantation phenomenon in *Xenopus* embryos. *Dev Cell* 4:219–230
- Ohkawara B, Yamamoto T, Tada M, Ueno N (2003) Role of glypican 4 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. *Development* 130:2129–2138
- Olson DJ, Oh D, Houston DW (2015) The dynamics of plus end polarization and microtubule assembly during *Xenopus* cortical rotation. *Dev Biol* 401:249–263. doi:[10.1016/j.ydbio.2015.01.028](https://doi.org/10.1016/j.ydbio.2015.01.028)
- Onichtchouk D, Chen YG, Dosch R et al (1999) Silencing of TGF-beta signalling by the pseudo-receptor BAMBI. *Nature* 401:480–485. doi:[10.1038/46794](https://doi.org/10.1038/46794)
- Onichtchouk D, Gawantka V, Dosch R et al (1996) The *Xvent-2* homeobox gene is part of the BMP-4 signalling pathway controlling [correction of controlling] dorsoventral patterning of *Xenopus* mesoderm. *Development* 122:3045–3053

- Oppenheimer JM (1934a) Experimental studies on the developing perch (*Perca flavescens* Mitchell). *Exp Biol Med* 31:1123–1124. doi:[10.3181/00379727-31-7465P](https://doi.org/10.3181/00379727-31-7465P)
- Oppenheimer JM (1934b) Experiments on early developing stages of fundulus. *Proc Natl Acad Sci U S A* 20:536–538
- Osada SI, Saijoh Y, Frisch A et al (2000) Activin/nodal responsiveness and asymmetric expression of a *Xenopus* nodal-related gene converge on a FAST-regulated module in intron 1. *Development* 127:2503–2514
- Ozair MZ, Kintner C, Brivanlou AH (2012) Neural induction and early patterning in vertebrates. *WIREs Dev Biol* 2:479–498. doi:[10.1002/wdev.90](https://doi.org/10.1002/wdev.90)
- Panáková D, Sprong H, Marois E et al (2005) Lipoprotein particles are required for Hedgehog and Wingless signalling. *Nature* 435:58–65. doi:[10.1038/nature03504](https://doi.org/10.1038/nature03504)
- Papanayotou C, Benhaddou A, Camus A et al (2014) A novel nodal enhancer dependent on pluripotency factors and smad2/3 signaling conditions a regulatory switch during epiblast maturation. *PLoS Biol* 12:e1001890. doi:[10.1371/journal.pbio.1001890](https://doi.org/10.1371/journal.pbio.1001890)
- Papkoff J, Brown A, Varmus H (1987) The int-1 proto-oncogene products are glycoproteins that appear to enter the secretory pathway. *Mol Cell Biol* 7:3978–3984
- Parameswaran M, Tam P (1995) Regionalisation of cell fate and morphogenetic movement of the mesoderm during mouse gastrulation. *Dev Genet* 17:16–28
- Parfitt D-E, Zernicka-Goetz M (2010) Epigenetic modification affecting expression of cell polarity and cell fate genes to regulate lineage specification in the early mouse embryo. *Mol Biol Cell* 21:2649–2660. doi:[10.1091/mbc.E10-01-0053](https://doi.org/10.1091/mbc.E10-01-0053)
- Parker D, Jemison J, Cadigan K (2002) Pygopus, a nuclear PHD-finger protein required for Wingless signaling in *Drosophila*. *Development* 129:2565–2576
- Pedersen RA, Wu K, Balakier H (1986) Origin of the inner cell mass in mouse embryos: cell lineage analysis by microinjection. *Dev Biol* 117:581–595
- Pelegri F, Maischein HM (1998) Function of zebrafish beta-catenin and TCF-3 in dorsoventral patterning. *Mech Dev* 77:63–74
- Pera EM, Ikeda A, Eivers E, De Robertis EM (2003) Integration of IGF, FGF, and anti-BMP signals via Smad1 phosphorylation in neural induction. *Genes Dev* 17:3023–3028. doi:[10.1101/gad.1153603](https://doi.org/10.1101/gad.1153603)
- Peradziryi H, Kaplan NA, Podleschny M et al (2011) PTK7/Otk interacts with Wnts and inhibits canonical Wnt signalling. *EMBO J* 30:3729–3740. doi:[10.1038/emboj.2011.236](https://doi.org/10.1038/emboj.2011.236)
- Perea-Gomez A, Lawson KA, Rhinn M et al (2001) Otx2 is required for visceral endoderm movement and for the restriction of posterior signals in the epiblast of the mouse embryo. *Development* 128:753–765
- Perea-Gomez A, Vella FDJ, Shawlot W et al (2002) Nodal antagonists in the anterior visceral endoderm prevent the formation of multiple primitive streaks. *Dev Cell* 3:745–756
- Petersen CP, Reddien PW (2011) Polarized notum activation at wounds inhibits Wnt function to promote planarian head regeneration. *Science* 332:852–855. doi:[10.1126/science.1202143](https://doi.org/10.1126/science.1202143)
- Piao S, Lee S-H, Kim H et al (2008) Direct inhibition of GSK3beta by the phosphorylated cytoplasmic domain of LRP6 in Wnt/beta-catenin signaling. *PLoS One* 3:e4046. doi:[10.1371/journal.pone.0004046](https://doi.org/10.1371/journal.pone.0004046)
- Piccolo S, Agius E, Leyns L et al (1999) The head inducer Cerberus is a multifunctional antagonist of Nodal, BMP and Wnt signals. *Nature* 397:707–710. doi:[10.1038/17820](https://doi.org/10.1038/17820)
- Piccolo S, Sasai Y, Lu B, De Robertis EM (1996) Dorsoventral patterning in *Xenopus*: inhibition of ventral signals by direct binding of chordin to BMP-4. *Cell* 86:589–598
- Piepenburg O, Grimmer D, Williams PH, Smith JC (2004) Activin redux: specification of mesodermal pattern in *Xenopus* by graded concentrations of endogenous activin B. *Development* 131:4977–4986. doi:[10.1242/dev.01323](https://doi.org/10.1242/dev.01323)
- Pinho S, Simonsson PR, Trevers KE et al (2011) Distinct steps of neural induction revealed by Asterix, Obelix and TrkC, genes induced by different signals from the organizer. *PLoS One* 6:e19157. doi:[10.1371/journal.pone.0019157](https://doi.org/10.1371/journal.pone.0019157)

- Piotrowska-Nitsche K, Perea-Gomez A, Haraguchi S, Zernicka-Goetz M (2005) Four-cell stage mouse blastomeres have different developmental properties. *Development* 132:479–490. doi:[10.1242/dev.01602](https://doi.org/10.1242/dev.01602)
- Plachta N, Bollenbach T, Pease S, Fraser SE (2011) Oct4 kinetics predict cell lineage patterning in the early mammalian embryo. *Nat Cell Biol* 13:117–123
- Plouhinec J-L, Zakin L, Moriyama Y, De Robertis EM (2013) Chordin forms a self-organizing morphogen gradient in the extracellular space between ectoderm and mesoderm in the *Xenopus* embryo. *Proc Natl Acad Sci U S A* 110:20372–20379. doi:[10.1073/pnas.1319745110](https://doi.org/10.1073/pnas.1319745110)
- Pukrop T, Gradl D, Henningfeld KA et al (2001) Identification of two regulatory elements within the high mobility group box transcription factor XTcf-4. *J Biol Chem* 276:8968–8978. doi:[10.1074/jbc.M007533200](https://doi.org/10.1074/jbc.M007533200)
- Rankin SA, Kormish J, Kofron M et al (2011) A gene regulatory network controlling *hhex* transcription in the anterior endoderm of the organizer. *Dev Biol* 351:297–310. doi:[10.1016/j.ydbio.2010.11.037](https://doi.org/10.1016/j.ydbio.2010.11.037)
- Rauch G, Hammerschmidt M, Blader P et al (1997) *Wnt5* is required for tail formation in the zebrafish embryo. *Cold Spring Harb Symp Quant Biol* 62:227–234
- Rawat VPS, Arseni N, Ahmed F et al (2010) The vent-like homeobox gene *VENTX* promotes human myeloid differentiation and is highly expressed in acute myeloid leukemia. *Proc Natl Acad Sci U S A* 107:16946–16951. doi:[10.1073/pnas.1001878107](https://doi.org/10.1073/pnas.1001878107)
- Rebagliati MR, Toyama R, Haffter P, Dawid IB (1998) *cyclops* encodes a nodal-related factor involved in midline signaling. *Proc Natl Acad Sci U S A* 95:9932–9937
- Rebagliati MR, Weeks DL, Harvey RP, Melton DA (1985) Identification and cloning of localized maternal RNAs from *Xenopus* eggs. *Cell* 42:769–777
- Reversade B, De Robertis EM (2005) Regulation of *ADMP* and *BMP2/4/7* at opposite embryonic poles generates a self-regulating morphogenetic field. *Cell* 123:1147–1160. doi:[10.1016/j.cell.2005.08.047](https://doi.org/10.1016/j.cell.2005.08.047)
- Reversade B, Kuroda H, Lee H et al (2005) Depletion of *Bmp2*, *Bmp4*, *Bmp7* and Spemann organizer signals induces massive brain formation in *Xenopus* embryos. *Development* 132:3381–3392. doi:[10.1242/dev.01901](https://doi.org/10.1242/dev.01901)
- Rex M, Hilton E, Old R (2002) Multiple interactions between maternally-activated signalling pathways control *Xenopus* nodal-related genes. *Int J Dev Biol* 46:217–226
- Rhinn M, Dierich A, Shawlot W et al (1998) Sequential roles for *Otx2* in visceral endoderm and neuroectoderm for forebrain and midbrain induction and specification. *Development* 125:845–856
- Rivera-Pérez JA, Mager J, Magnuson T (2003) Dynamic morphogenetic events characterize the mouse visceral endoderm. *Dev Biol* 261:470–487
- Rodriguez TA (2005) Induction and migration of the anterior visceral endoderm is regulated by the extra-embryonic ectoderm. *Development* 132:2513–2520. doi:[10.1242/dev.01847](https://doi.org/10.1242/dev.01847)
- Roël G, Hamilton FS, Gent Y et al (2002) *Lef-1* and *Tcf-3* transcription factors mediate tissue-specific *Wnt* signaling during *Xenopus* development. *Curr Biol* 12:1941–1945
- Roose J, Molenaar M, Peterson J et al (1998) The *Xenopus* *Wnt* effector *XTcf-3* interacts with Groucho-related transcriptional repressors. *Nature* 395:608–612
- Rosenquist TA, Martin GR (1995) Visceral endoderm-1 (VE-1): an antigen marker that distinguishes anterior from posterior embryonic visceral endoderm in the early post-implantation mouse embryo. *Mech Dev* 49:117–121
- Rothbacher U, Laurent MN, Deardorff MA et al (2000) Dishevelled phosphorylation, subcellular localization and multimerization regulate its role in early embryogenesis. *EMBO J* 19:1010–1022. doi:[10.1093/emboj/19.5.1010](https://doi.org/10.1093/emboj/19.5.1010)
- Roux W (1888) Beiträge zur Entwicklungsmechanik des Embryo. V. *Virchows Arch Pathol Anat* 114:113–153
- Roux W (1885) Ueber die bestimmung der hauptrichtungen des froschembryo im ei und über die erste theilung des froscheies. *Zeitschrift* 20:1–54
- Roux W (1887) Beiträge zur Entwicklungsmechanik des Embryo. *Arch Mikrosk Anat* 29:157–211



- Roux W (1903) Über die Ursachen der Bestimmung der Hauptrichtungen des Embryo im Froschei. *Anat Anz* 23:65–183
- Rowning BA, Wells J, Wu M et al (1997) Microtubule-mediated transport of organelles and localization of beta-catenin to the future dorsal side of *Xenopus* eggs. *Proc Natl Acad Sci U S A* 94:1224–1229
- Rugh R (1951) The frog, its reproduction and development. The Blakiston Co., Philadelphia, PA
- Ruiz i Altaba A, Choi T, Melton DA (1991) Expression of the Xhox3 homeobox protein in *Xenopus* embryos: blocking its early function suggests the requirement of Xhox3 for normal posterior development (axial pattern/central nervous system/embryonic mesoderm/homeobox gene/*Xenopus laevis*). *Dev Growth Differ* 33:651–669. doi:10.1111/j.1440-169X.1991.00651.x
- Saint-Jeannet JP, Huang S, Duprat AM (1990) Modulation of neural commitment by changes in target cell contacts in *Pleurodeles waltl*. *Dev Biol* 141:93–103
- Sala M (1955) Distribution of activating and transforming influences in the archenteron roof during the induction of the nervous system in amphibians. *PNAS* 58:635–647
- Salic AN, Kroll KL, Evans LM, Kirschner MW (1997) Sizzled: a secreted Xwnt8 antagonist expressed in the ventral marginal zone of *Xenopus* embryos. *Development* 124:4739–4748
- Sampath K, Rubinstein AL, Cheng AM et al (1998) Induction of the zebrafish ventral brain and floorplate requires cyclops/nodal signalling. *Nature* 395:185–189. doi:10.1038/26020
- Sander V, Reversade B, De Robertis EM (2007) The opposing homeobox genes Goosecoid and Vent1/2 self-regulate *Xenopus* patterning. *EMBO J* 26:2955–2965. doi:10.1038/sj.emboj.7601705
- Saneyoshi T, Kume S, Amasaki Y, Mikoshiba K (2002) The Wnt/calcium pathway activates NF-AT and promotes ventral cell fate in *Xenopus* embryos. *Nature* 417:295–299. doi:10.1038/417295a
- Sasai Y, Lu B, Steinbeisser H, De Robertis EM (1995) Regulation of neural induction by the Chd and Bmp-4 antagonistic patterning signals in *Xenopus*. *Nature* 376:333–336. doi:10.1038/376333a0
- Sasai Y, Lu B, Steinbeisser H et al (1994) *Xenopus* chordin: a novel dorsalizing factor activated by organizer-specific homeobox genes. *Cell* 79:779–790
- Scerbo P, Girardot F, Vivien C et al (2012) Ventx factors function as nanog-like guardians of developmental potential in *Xenopus*. *PLoS One* 7:e36855. doi:10.1371/journal.pone.0036855.s008
- Scerbo P, Markov GV, Vivien C et al (2014) On the origin and evolutionary history of NANOG. *PLoS One* 9:e85104. doi:10.1371/journal.pone.0085104
- Schambony A, Wedlich D (2007) Wnt-5A/Ror2 regulate expression of XPAPC through an alternative noncanonical signaling pathway. *Dev Cell* 12:779–792. doi:10.1016/j.devcel.2007.02.016
- Scharf SR, Gerhart JC (1983) Axis determination in eggs of *Xenopus laevis*: a critical period before first cleavage, identified by the common effects of cold, pressure and ultraviolet irradiation. *Dev Biol* 99:75–87
- Scharf SR, Gerhart JC (1980) Determination of the dorsal-ventral axis in eggs of *Xenopus laevis*: complete rescue of uv-impaired eggs by oblique orientation before first cleavage. *Dev Biol* 79:181–198
- Scharf SR, Rowning B, Wu M, Gerhart JC (1989) Hyperdorsoanterior embryos from *Xenopus* eggs treated with D<sub>2</sub>O. *Dev Biol* 134:175–188
- Schmid B, Fürthauer M, Connors SA et al (2000) Equivalent genetic roles for bmp7/snailhouse and bmp2b/swirl in dorsoventral pattern formation. *Development* 127:957–967
- Schmidt JE, von Dassow G, Kimelman D (1996) Regulation of dorsal-ventral patterning: the ventralizing effects of the novel *Xenopus* homeobox gene Vox. *Development* 122:1711–1721
- Schneider S, Steinbeisser H, Warga RM, Hausen P (1996) Beta-catenin translocation into nuclei demarcates the dorsalizing centers in frog and fish embryos. *Mech Dev* 57:191–198
- Schohl A, Fagotto F (2002) Beta-catenin, MAPK and Smad signaling during early *Xenopus* development. *Development* 129:37–52
- Schroeder KE, Condic ML, Eisenberg LM, Yost HJ (1999) Spatially regulated translation in embryos: asymmetric expression of maternal Wnt-11 along the dorsal-ventral axis in *Xenopus*. *Dev Biol* 214:288–297. doi:10.1006/dbio.1999.9426
- Schroeder MM, Gard DL (1992) Organization and regulation of cortical microtubules during the first cell cycle of *Xenopus* eggs. *Development* 114:699–709

- Schultze O (1899) Ueber das erste auftreten der bilateralen symmetrie im verlauf der entwicklung. Arch Mikrosk Anat 55:171–201
- Seidel F (1956) Nachweis eines Zentrums zur Bildung der Keimscheibe im Säugetierei. Naturwissenschaften 43:306–307
- Seleiro EA, Connolly DJ, Cooke J (1996) Early developmental expression and experimental axis determination by the chicken Vg1 gene. Curr Biol 6:1476–1486
- Seoane J, Lagna G, Montalvo E, Hemmati-Brivanlou A (2000) OAZ uses distinct DNA- and protein-binding zinc fingers in separate BMP-Smad and Olf signaling pathways. Cell 100(2):229–240
- Shah SB, Skromne I, Hume CR, Kessler DS, Lee KJ, Stern CD, Dodd J (1997) Misexpression of chick Vg1 in the marginal zone induces primitive streak formation. Development 124:5127–5138
- Shawlot W, Wakamiya M, Kwan KM et al (1999) Lim1 is required in both primitive streak-derived tissues and visceral endoderm for head formation in the mouse. Development 126:4925–4932
- Sheldahl L, Park M, Malbon C, Moon R (1999) Protein kinase C is differentially stimulated by Wnt and Frizzled homologs in a G-protein-dependent manner. Curr Biol 9:695–698
- Sheldahl L, Slusarski D, Pandur P et al (2003) Dishevelled activates Ca<sup>2+</sup> flux, PKC, and CamKII in vertebrate embryos. J Cell Biol 161:769–777
- Sheng G, dos Reis M, Stern CD (2003) Churchill, a zinc finger transcriptional activator, regulates the transition between gastrulation and neurulation. Cell 115:603–613
- Shih J, Keller R (1992a) The epithelium of the dorsal marginal zone of *Xenopus* has organizer properties. Development 116:887–899
- Shih J, Keller R (1992b) Patterns of cell motility in the organizer and dorsal mesoderm of *Xenopus laevis*. Development 116:915–930
- Shimizu K, Gurdon JB (1999) A quantitative analysis of signal transduction from activin receptor to nucleus and its relevance to morphogen gradient interpretation. Proc Natl Acad Sci U S A 96:6791–6796
- Shinya M, Eschbach C, Clark M et al (2000) Zebrafish Dkk1, induced by the pre-MBT Wnt signaling, is secreted from the prechordal plate and patterns the anterior neural plate. Mech Dev 98:3–17
- Shook DR, Keller R (2008) Epithelial type, ingression, blastopore architecture and the evolution of chordate mesoderm morphogenesis. J Exp Zool B Mol Dev Evol 310:85–110. doi:10.1002/jez.b.21198
- Shook DR, Majer C, Keller R (2002) Urodeles remove mesoderm from the superficial layer by subduction through a bilateral primitive streak. Dev Biol 248:220–239. doi:10.1006/dbio.2002.0718
- Shy BR, Wu C-I, Khramtsova GF et al (2013) Regulation of Tcf7l1 DNA binding and protein stability as principal mechanisms of Wnt/ $\beta$ -catenin signaling. Cell Rep 4:1–9. doi:10.1016/j.celrep.2013.06.001
- Sick S, Reinker S, Timmer J, Schlake T (2006) WNT and DKK determine hair follicle spacing through a reaction-diffusion mechanism. Science 314:1447–1450. doi:10.1126/science.1130088
- Siegfried E, Chou T, Perrimon N (1992) wingless signaling acts through zeste-white 3, the Drosophila homolog of glycogen synthase kinase-3, to regulate engrailed and establish cell fate. Cell 71:1167–1179
- Siegfried E, Wilder E, Perrimon N (1994) Components of wingless signalling in Drosophila. Nature 367:76–80
- Simeoni I, Gurdon JB (2007) Interpretation of BMP signaling in early *Xenopus* development. Dev Biol 308:82–92. doi:10.1016/j.ydbio.2007.05.010
- Sivak J, Petersen L, Amaya E (2005) FGF signal interpretation is directed by sprouty and spread proteins during mesoderm formation. Dev Cell 8:689–701. doi:10.1016/j.devcel.2005.02.011
- Skirkanich J, Luxardi G, Yang J et al (2011) An essential role for transcription before the MBT in *Xenopus laevis*. Dev Biol 357:478–491. doi:10.1016/j.ydbio.2011.06.010
- Skromne I, Stern CD (2001) Interactions between Wnt and Vg1 signalling pathways initiate primitive streak formation in the chick embryo. Development 128:2915–2927
- Skromne I, Stern CD (2002) A hierarchy of gene expression accompanying induction of the primitive streak by Vg1 in the chick embryo. Mech Dev 114:115–118

- Slack JMW (1991) From egg to embryo: regional specification in early development. Cambridge University Press, Cambridge
- Slusarski D, Yang-Snyder JA, Busa W, Moon R (1997a) Modulation of embryonic intracellular Ca<sup>2+</sup> signaling by Wnt-5A. *Dev Biol* 182:114–120
- Slusarski DC, Corces VG, Moon RT (1997b) Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature* 390:410–413. doi:[10.1038/37138](https://doi.org/10.1038/37138)
- Smith JC (1989) Mesoderm induction and mesoderm-inducing factors in early amphibian development. *Development* 105:665–677
- Smith JC, Dale L, Slack JM (1985) Cell lineage labels and region-specific markers in the analysis of inductive interactions. *J Embryol Exp Morphol* 89(Suppl):317–331
- Smith W, Harland R (1992) Expression cloning of noggin, a new dorsalizing factor localized to the Spemann organizer in *Xenopus* embryos. *Cell* 70:829–840
- Smith WC, Harland RM (1991) Injected Xwnt-8 RNA acts early in *Xenopus* embryos to promote formation of a vegetal dorsalizing center. *Cell* 67:753–765
- Smith WC, McKendry R, Ribisi S, Harland RM (1995) A nodal-related gene defines a physical and functional domain within the Spemann organizer. *Cell* 82:37–46
- Snow M (1977) Gastrulation in the mouse: growth and regionalization of the epiblast. *Development* 42:293–303
- Sokol S (1996) Analysis of Dishevelled signalling pathways during *Xenopus* development. *Curr Biol* 6:1456–1467
- Sokol S, Christian JL, Moon RT, Melton DA (1991) Injected Wnt RNA induces a complete body axis in *Xenopus* embryos. *Cell* 67:741–752
- Solnica-Krezel L (2005) Conserved patterns of cell movements during vertebrate gastrulation. *Curr Biol* 15:R213–R228. doi:[10.1016/j.cub.2005.03.016](https://doi.org/10.1016/j.cub.2005.03.016)
- Solnica-Krezel L, Driever W (1994) Microtubule arrays of the zebrafish yolk cell: organization and function during epiboly. *Development* 120:2443–2455
- Solnica-Krezel L, Sepich DS (2012) Gastrulation: making and shaping germ layers. *Annu Rev Cell Dev Biol* 28:687–717. doi:[10.1146/annurev-cellbio-092910-154043](https://doi.org/10.1146/annurev-cellbio-092910-154043)
- Spemann H (1938) Embryo development and induction. Yale University Press, New Haven
- Spemann H (1931) Über den Anteil von Implantat und Wirtskeim an der Orientierung und Beschaffenheit der induzierten Embryonalanlage. *Wilhelm Roux' Arch Entwicklungsmech* 123:389–517. doi:[10.1007/BF01380646](https://doi.org/10.1007/BF01380646)
- Spemann H (1918) Über die Determination der ersten Organanlagen des Amphibienembryo I–VI. *Wilhelm Roux' Arch Entwicklungsmech* 43:448–555. doi:[10.1007/BF02267308](https://doi.org/10.1007/BF02267308)
- Spemann H (1921) Die Erzeugung tierischer Chimären durch heteroplastische embryonale Transplantation zwischen Triton cristatus und taeniatus. *Wilhelm Roux' Arch Entwicklungsmech* 48:533–570. doi:[10.1007/BF02554578](https://doi.org/10.1007/BF02554578)
- Spemann H (1903) Entwicklungsphysiologische Studien am Triton-Ei. *Wilhelm Roux' Arch Entwicklungsmech* 16:551–631. doi:[10.1007/BF02301267](https://doi.org/10.1007/BF02301267)
- Spemann H, Mangold (1924) Über Induktion von Embryonenanlagen durch Implantation artfremder Organisatoren. *Wilhelm Roux' Arch EntwMech Org* 100:599–638
- Spratt NT, Haas H (1960) Integrative mechanisms in development of the early chick blastoderm. I. Regulative potentiality of separated parts. *J Exp Zool* 145:97–137
- Srinivas S, Rodriguez T, Clements M et al (2004) Active cell migration drives the unilateral movements of the anterior visceral endoderm. *Development* 131:1157–1164. doi:[10.1242/dev.01005](https://doi.org/10.1242/dev.01005)
- Standley HJ, Gurdon JB (2004) The community effect in *Xenopus* development. In: H. Grunz (Ed.) *The vertebrate organizer*. Springer, Berlin, Heidelberg, pp 73–91
- Stern CD (1990) The marginal zone and its contribution to the hypoblast and primitive streak of the chick embryo. *Development* 109:667–682
- Stern CD (2004) Gastrulation in the chick. From Cells to Embryo, Gastrulation
- Stern CD (2006) Neural induction: 10 years on since the “default model”. *Curr Opin Cell Biol* 18:692–697. doi:[10.1016/j.ceb.2006.09.002](https://doi.org/10.1016/j.ceb.2006.09.002)
- Stern CD (2005) Neural induction: old problem, new findings, yet more questions. *Development* 132:2007–2021. doi:[10.1242/dev.01794](https://doi.org/10.1242/dev.01794)

- Stern CD, Downs KM (2012) The hypoblast (visceral endoderm): an evo-devo perspective. *Development* 139:1059–1069. doi:[10.1242/dev.070730](https://doi.org/10.1242/dev.070730)
- Strähle U, Jesuthasan S (1993) Ultraviolet irradiation impairs epiboly in zebrafish embryos: evidence for a microtubule-dependent mechanism of epiboly. *Development* 119:909–919
- Streit A, Lee KJ, Woo I et al (1998) Chordin regulates primitive streak development and the stability of induced neural cells, but is not sufficient for neural induction in the chick embryo. *Development* 125:507–519
- Streit A, Stern CD (1999) Establishment and maintenance of the border of the neural plate in the chick: involvement of FGF and BMP activity. *Mech Dev* 82:51–66
- Strutt D (2001) Asymmetric localization of frizzled and the establishment of cell polarity in the *Drosophila* wing. *Mol Cell* 7:367–375
- Stuckey DW, Clements M, Di-Gregorio A et al (2011) Coordination of cell proliferation and anterior-posterior axis establishment in the mouse embryo. *Development* 138:1521–1530. doi:[10.1242/dev.063537](https://doi.org/10.1242/dev.063537)
- Sudou N, Yamamoto S, Ogino H, Taira M (2012) Dynamic in vivo binding of transcription factors to cis-regulatory modules of *cer* and *gsc* in the stepwise formation of the Spemann-Mangold organizer. *Development* 139:1651–1661. doi:[10.1242/dev.068395](https://doi.org/10.1242/dev.068395)
- Suriben R, Kivimäe S, Fisher DAC et al (2009) Posterior malformations in *Dact1* mutant mice arise through misregulated *Vangl2* at the primitive streak. *Nat Genet* 41:977–985. doi:[10.1038/ng.435](https://doi.org/10.1038/ng.435)
- Suzuki A, Thies R, Yamaji N et al (1994) A truncated bone morphogenetic protein receptor affects dorsal-ventral patterning in the early *Xenopus* embryo. *Proc Natl Acad Sci U S A* 91:10255–10259
- Suzuki A, Ueno N, Hemmati-Brivanlou A (1997) *Xenopus* *msx1* mediates epidermal induction and neural inhibition by BMP4. *Development* 124:3037–3044
- Tabansky I, Lenarcic A, Draft RW et al (2013) Developmental bias in cleavage-stage mouse blastomeres. *Curr Biol* 23:21–31. doi:[10.1016/j.cub.2012.10.054](https://doi.org/10.1016/j.cub.2012.10.054)
- Tada M, Smith J (2000) *Xwnt11* is a target of *Xenopus* Brachyury: regulation of gastrulation movements via *Dishevelled*, but not through the canonical Wnt pathway. *Development* 127:2227–2238
- Taelman VF, Dobrowolski R, Plouhinec J-L et al (2010) Wnt signaling requires sequestration of glycogen synthase kinase 3 inside multivesicular endosomes. *Cell* 143:1136–1148. doi:[10.1016/j.cell.2010.11.034](https://doi.org/10.1016/j.cell.2010.11.034)
- Tahinci E, Symes K (2003) Distinct functions of Rho and Rac are required for convergent extension during *Xenopus* gastrulation. *Dev Biol* 259:318–335
- Taira M, Jamrich M, Good PJ, Dawid IB (1992) The LIM domain-containing homeo box gene *Xlim-1* is expressed specifically in the organizer region of *Xenopus* gastrula embryos. *Genes Dev* 6:356–366
- Takahashi S, Yokota C, Takano K et al (2000) Two novel nodal-related genes initiate early inductive events in *Xenopus* Nieuwkoop center. *Development* 127:5319–5329
- Takaoka K, Hamada H (2012) Cell fate decisions and axis determination in the early mouse embryo. *Development* 139:3–14. doi:[10.1242/dev.060095](https://doi.org/10.1242/dev.060095)
- Takaoka K, Yamamoto M, Hamada H (2011) Origin and role of distal visceral endoderm, a group of cells that determines anterior-posterior polarity of the mouse embryo. *Nat Cell Biol* 13:743–752. doi:[10.1038/ncb2251](https://doi.org/10.1038/ncb2251)
- Takaoka K, Yamamoto M, Shiratori H et al (2006) The mouse embryo autonomously acquires anterior-posterior polarity at implantation. *Dev Cell* 10:451–459. doi:[10.1016/j.devcel.2006.02.017](https://doi.org/10.1016/j.devcel.2006.02.017)
- Takeuchi M, Nakabayashi J, Sakaguchi T et al (2003) The Prickle-related gene in vertebrates is essential for gastrulation cell movements. *Curr Biol* 13:674–679
- Tam PP, Loebel DA, Tanaka SS (2006) Building the mouse gastrula: signals, asymmetry and lineages. *Curr Opin Genet Dev* 16:419–425. doi:[10.1016/j.gde.2006.06.008](https://doi.org/10.1016/j.gde.2006.06.008)
- Tam PP, Steiner KA (1999) Anterior patterning by synergistic activity of the early gastrula organizer and the anterior germ layer tissues of the mouse embryo. *Development* 126:5171–5179
- Tamai K, Semenov M, Kato Y et al (2000) LDL-receptor-related proteins in Wnt signal transduction. *Nature* 407:530–535
- Tamai K, Zeng X, Liu C et al (2004) A mechanism for Wnt coreceptor activation. *Mol Cell* 13:149–156

- Tanaka K, Okabayashi K, Asashima M et al (2000) The evolutionarily conserved porcupine gene family is involved in the processing of the Wnt family. *Eur J Biochem* 267:4300–4311
- Tao Q, Yokota C, Puck H et al (2005) Maternal wnt11 activates the canonical wnt signaling pathway required for axis formation in *Xenopus* embryos. *Cell* 120:857–871. doi:[10.1016/j.cell.2005.01.013](https://doi.org/10.1016/j.cell.2005.01.013)
- Tarbashevich K, Koebernick K, Pieler T (2007) XGRIP2.1 is encoded by a vegetally localizing, maternal mRNA and functions in germ cell development and anteroposterior PGC positioning in *Xenopus laevis*. *Dev Biol* 311:554–565. doi:[10.1016/j.ydbio.2007.09.012](https://doi.org/10.1016/j.ydbio.2007.09.012)
- Tarkowski AK (1959) Experiments on the development of isolated blastomeres of mouse eggs. *Nature* 184:1286–1287. doi:[10.1038/1841286a0](https://doi.org/10.1038/1841286a0)
- Tarkowski AK (1961) Mouse chimaeras developed from fused eggs. *Nature* 190:857–860
- Tarkowski AK, Wróblewska J (1967) Development of blastomeres of mouse eggs isolated at the 4- and 8-cell stage. *J Embryol Exp Morphol* 18:155–180
- Terasaki M, Chen LB, Fujiwara K (1986) Microtubules and the endoplasmic reticulum are highly independent structures. *J Cell Biol* 103:1557–1568
- Thisse B, Wright CV, Thisse C (2000) Activin- and Nodal-related factors control antero-posterior patterning of the zebrafish embryo. *Nature* 403:425–428. doi:[10.1038/35000200](https://doi.org/10.1038/35000200)
- Thomas P, Beddington R (1996) Anterior primitive endoderm may be responsible for patterning the anterior neural plate in the mouse embryo. *Curr Biol* 6:1487–1496
- Thomas PQ, Brown A, Beddington RS (1998) Hex: a homeobox gene revealing peri-implantation asymmetry in the mouse embryo and an early transient marker of endothelial cell precursors. *Development* 125:85–94
- Tolwinski NS, Wehrl M, Rives A et al (2003) Wg/Wnt signal can be transmitted through arrow/ LRP5,6 and Axin independently of Zw3/Gsk3beta activity. *Dev Cell* 4:407–418
- Topczewski J, Sepich DS, Myers DC et al (2001) The zebrafish glypican knypek controls cell polarity during gastrulation movements of convergent extension. *Dev Cell* 1:251–264
- Topol L, Jiang X, Choi H et al (2003) Wnt-5a inhibits the canonical Wnt pathway by promoting GSK-3-independent beta-catenin degradation. *J Cell Biol* 162:899–908. doi:[10.1083/jcb.200303158](https://doi.org/10.1083/jcb.200303158)
- Torban E, Patenaude A-M, Leclerc S et al (2008) Genetic interaction between members of the Vangl family causes neural tube defects in mice. *Proc Natl Acad Sci U S A* 105:3449–3454. doi:[10.1073/pnas.0712126105](https://doi.org/10.1073/pnas.0712126105)
- Torlopp A, Khan MAF, Oliveira NMM et al (2014) The transcription factor Pitx2 positions the embryonic axis and regulates twinning. *eLife Sci* 3:e03743. doi:[10.7554/eLife.03743](https://doi.org/10.7554/eLife.03743)
- Torres MA, Yang-Snyder JA, Purcell SM et al (1996) Activities of the Wnt-1 class of secreted signaling factors are antagonized by the Wnt-5A class and by a dominant negative cadherin in early *Xenopus* development. *J Cell Biol* 133:1123–1137
- Torres-Padilla M-E, Parfitt D-E, Kouzarides T, Zernicka-Goetz M (2007a) Histone arginine methylation regulates pluripotency in the early mouse embryo. *Nature* 445:214–218. doi:[10.1038/nature05458](https://doi.org/10.1038/nature05458)
- Torres-Padilla M-E, Richardson L, Kolasinska P et al (2007b) The anterior visceral endoderm of the mouse embryo is established from both preimplantation precursor cells and by de novo gene expression after implantation. *Dev Biol* 309:97–112. doi:[10.1016/j.ydbio.2007.06.020](https://doi.org/10.1016/j.ydbio.2007.06.020)
- Tortelote GG, Hernández-Hernández JM, Quaresma AJC et al (2013) Wnt3 function in the epiblast is required for the maintenance but not the initiation of gastrulation in mice. *Dev Biol* 374:164–173. doi:[10.1016/j.ydbio.2012.10.013](https://doi.org/10.1016/j.ydbio.2012.10.013)
- Tran LD, Hino H, Quach H et al (2012) Dynamic microtubules at the vegetal cortex predict the embryonic axis in zebrafish. *Development* 139:3644–3652. doi:[10.1242/dev.082362](https://doi.org/10.1242/dev.082362)
- Trindade M, Tada M, Smith JC (1999) DNA-binding specificity and embryological function of Xom (Xvent-2). *Dev Biol* 216:442–456. doi:[10.1006/dbio.1999.9507](https://doi.org/10.1006/dbio.1999.9507)
- Tropepe V, Hitoshi S, Sirard C et al (2001) Direct neural fate specification from embryonic stem cells: a primitive mammalian neural stem cell stage acquired through a default mechanism. *Neuron* 30:65–78

- Tsunekawa N, Naito M, Sakai Y et al (2000) Isolation of chicken vasa homolog gene and tracing the origin of primordial germ cells. *Development* 127:2741–2750
- Tucker JA, Mintzer KA, Mullins MC (2008) The BMP signaling gradient patterns dorsoventral tissues in a temporally progressive manner along the anteroposterior axis. *Dev Cell* 14:108. doi:[10.1016/j.devcel.2007.11.004](https://doi.org/10.1016/j.devcel.2007.11.004)
- Tung TC, Wu SC, Tung Y (1962) Experimental studies on neural induction in *Amphioxus*. *Sci Sin* 11:805
- Tzika A, Milinkovitch MC (2008) A pragmatic approach for selecting evo-devomodel species in amniotes. In: Minelli A, Fusco G (eds) *Evolving pathways; key themes in evolutionary developmental biology*. Cambridge University Press, Cambridge, pp 119–140
- Ulrich F, Concha M, Heid P et al (2003) *Slb/Wnt11* controls hypoblast cell migration and morphogenesis at the onset of zebrafish gastrulation. *Development* 130:5375–5384
- Vallier L, Mendjan S, Brown S et al (2009) *Activin/Nodal* signalling maintains pluripotency by controlling *Nanog* expression. *Development* 136:1339–1349. doi:[10.1242/dev.033951](https://doi.org/10.1242/dev.033951)
- van Amerongen R, Fuerer C, Mizutani M, Nusse R (2012) *Wnt5a* can both activate and repress *Wnt/β-catenin* signaling during mouse embryonic development. *Dev Biol* 369:101–114. doi:[10.1016/j.ydbio.2012.06.020](https://doi.org/10.1016/j.ydbio.2012.06.020)
- van den Heuvel M, Harryman-Samos C, Klingensmith J et al (1993) Mutations in the segment polarity genes *wingless* and *porcupine* impair secretion of the *wingless* protein. *EMBO J* 12:5293–5302
- Varelas X, Miller BW, Sopko R et al (2010) The Hippo pathway regulates *Wnt/beta-catenin* signaling. *Dev Cell* 18:579–591. doi:[10.1016/j.devcel.2010.03.007](https://doi.org/10.1016/j.devcel.2010.03.007)
- Vaughan RB, Trinkaus JP (1966) Movements of epithelial cell sheets in vitro. *J Cell Sci* 1:407–413
- Veeman M, Axelrod JD, Moon R (2003a) A second canon. Functions and mechanisms of *beta-catenin-independent Wnt* signaling. *Dev Cell* 5:367–377
- Veeman MT, Slusarski DC, Kaykas A et al (2003b) Zebrafish *Prickle*, a modulator of noncanonical *Wnt/Fz* signaling, regulates gastrulation movements. *Curr Biol* 13:680–685. doi:[10.1016/S0960-9822\(03\)00240-9](https://doi.org/10.1016/S0960-9822(03)00240-9)
- VerMilyea MD, Maneck M, Yoshida N et al (2011) Transcriptome asymmetry within mouse zygotes but not between early embryonic sister blastomeres. *EMBO J* 30:1841–1851. doi:[10.1038/emboj.2011.92](https://doi.org/10.1038/emboj.2011.92)
- Viebahn C (2001) Hensen's node. *Genesis* 29:96–103
- Viebahn C, Mayer B, Hrabé de Angelis M (1995) Signs of the principle body axes prior to primitive streak formation in the rabbit embryo. *Anat Embryol* 192:159–169
- Vincent J, Oster G, Gerhart J (1986) Kinematics of gray crescent formation in *Xenopus* eggs: the displacement of subcortical cytoplasm relative to the egg surface. *Dev Biol* 113:484–500
- Vincent JP, Gerhart JC (1987) Subcortical rotation in *Xenopus* eggs: an early step in embryonic axis specification. *Dev Biol* 123:526–539
- Vincent SD, Dunn NR, Hayashi S et al (2003) Cell fate decisions within the mouse organizer are governed by graded *Nodal* signals. *Genes Dev* 17:1646–1662. doi:[10.1101/gad.1100503](https://doi.org/10.1101/gad.1100503)
- Vinson CR, Adler PN (1987) Directional non-cell autonomy and the transmission of polarity information by the *frizzled* gene of *Drosophila*. *Nature* 329:549–551. doi:[10.1038/329549a0](https://doi.org/10.1038/329549a0)
- Vintemberger P, Clavert J (1959) On the determinism of bilateral symmetry in birds: XI. The moment of embryonic axis determination, according to the results of our experiences in rotation of the chicken egg in the uterus. *C R Seances Soc Biol Fil* 153:661–665
- Vogt W (1929) Gestaltungsanalyse am Amphibienkeim mit Örtlicher Vitalfärbung. *Wilhelm Roux' Arch Entwicklungsmech* 120:384–706. doi:[10.1007/BF02109667](https://doi.org/10.1007/BF02109667)
- Voiculescu O, Bertocchini F, Wolpert L et al (2007) The amniote primitive streak is defined by epithelial cell intercalation before gastrulation. *Nature* 449:1049–1052. doi:[10.1038/nature06211](https://doi.org/10.1038/nature06211)
- Voiculescu O, Bodenstein L, Lau I-J, Stern CD (2014) Local cell interactions and self-amplifying individual cell ingression drive amniote gastrulation. *eLife Sci* 3:e01817
- Wacker S, Grimm K, Joos T, Winklbauer R (2000) Development and control of tissue separation at gastrulation in *Xenopus*. *Dev Biol* 224:428–439. doi:[10.1006/dbio.2000.9794](https://doi.org/10.1006/dbio.2000.9794)
- Waddington CH (1940) *Organisers and genes*. Cambridge University Press, Cambridge UK
- Waddington CH (1956) *Principles of embryology*. The Macmillan Company, New York

- Waddington CH (1930) Developmental mechanics of chicken and duck embryos. *Nature* 125:924–925. doi:[10.1038/125924b0](https://doi.org/10.1038/125924b0)
- Waddington CH (1934) Experiments on Embryonic induction. III. A note on inductions by chick primitive streak transplanted to the rabbit Embryo. *J Exp Biol* 11:224–227
- Waddington CH (1936) Organizers in mammalian development: Abstract. *Nature*
- Waddington CH (1937) Experiments on determination in the rabbit embryo. *Arch Biol* 48:273–290
- Waddington CH (1933) Induction by the endoderm in birds. Wilhelm Roux' Arch Entwicklungsmech 128:502–521. doi:[10.1007/BF00649862](https://doi.org/10.1007/BF00649862)
- Waldrip WR, Bikoff EK, Hoodless PA et al (1998) Smad2 signaling in extraembryonic tissues determines anterior-posterior polarity of the early mouse embryo. *Cell* 92:797–808
- Wallingford JB, Goto T, Keller R, Harland RM (2002) Cloning and expression of *Xenopus* Prickle, an orthologue of a Drosophila planar cell polarity gene. *Mech Dev* 116:183–186
- Wallingford JB, Rowling BA, Vogeli KM et al (2000) Dishevelled controls cell polarity during *Xenopus* gastrulation. *Nature* 405:81–85. doi:[10.1038/35011077](https://doi.org/10.1038/35011077)
- Wang J, Hamblet NS, Mark S et al (2006) Dishevelled genes mediate a conserved mammalian PCP pathway to regulate convergent extension during neurulation. *Development* 133:1767–1778. doi:[10.1242/dev.02347](https://doi.org/10.1242/dev.02347)
- Wang S, Krinks M, Lin K et al (1997) Frzb, a secreted protein expressed in the Spemann organizer, binds and inhibits Wnt-8. *Cell* 88:757–766
- Wang Y, Nathans J (2007) Tissue/planar cell polarity in vertebrates: new insights and new questions. *Development* 134:647–658. doi:[10.1242/dev.02772](https://doi.org/10.1242/dev.02772)
- Wawersik S, Evola C, Whitman M (2005) Conditional BMP inhibition in *Xenopus* reveals stage-specific roles for BMPs in neural and neural crest induction. *Dev Biol* 277:425–442. doi:[10.1016/j.ydbio.2004.10.002](https://doi.org/10.1016/j.ydbio.2004.10.002)
- Weaver C, Farr GH, Pan W et al (2003) GBP binds kinesin light chain and translocates during cortical rotation in *Xenopus* eggs. *Development* 130:5425–5436. doi:[10.1242/dev.00737](https://doi.org/10.1242/dev.00737)
- Weaver C, Kimelman D (2004) Move it or lose it: axis specification in *Xenopus*. *Development* 131:3491–3499. doi:[10.1242/dev.01284](https://doi.org/10.1242/dev.01284)
- Weber RJ, Pedersen RA, Wianny F et al (1999) Polarity of the mouse embryo is anticipated before implantation. *Development* 126:5591–5598
- Weber GF, Bjerke MA, DeSimone DW (2012) A mechanoresponsive cadherin-keratin complex directs polarized protrusive behavior and collective cell migration. *Dev Cell* 22:104–115. doi:[10.1016/j.devcel.2011.10.013](https://doi.org/10.1016/j.devcel.2011.10.013)
- Weeks DL, Melton DA (1987) A maternal mRNA localized to the vegetal hemisphere in *Xenopus* eggs codes for a growth factor related to TGF-beta. *Cell* 51:861–867
- Weidinger G, Stebler J, Slanchev K et al (2003) dead end, a novel vertebrate germ plasm component, is required for zebrafish primordial germ cell migration and survival. *Curr Biol* 13:1429–1434
- Weise A, Bruser K, Elfert S et al (2010) Alternative splicing of Tcf7l2 transcripts generates protein variants with differential promoter-binding and transcriptional activation properties at Wnt/beta-catenin targets. *Nucleic Acids Res* 38:1964–1981. doi:[10.1093/nar/gkp1197](https://doi.org/10.1093/nar/gkp1197)
- Weiss PA (1926) Morphodynamik: Ein Einblick in die Gesetze der organischen Gestaltung an Hand von experimentellen Ergebnissen. In: Schaxel J (ed) Abhandlungen zur theoretischen Biologie, vol 23. Gebrüder Borntraeger, Berlin
- Wessely O, Agius E, Oelgeschlager M et al (2001) Neural induction in the absence of mesoderm: beta-catenin-dependent expression of secreted BMP antagonists at the blastula stage in *Xenopus*. *Dev Biol* 234:161–173. doi:[10.1006/dbio.2001.0258](https://doi.org/10.1006/dbio.2001.0258)
- Westfall T, Brimeyer R, Twedt J et al (2003) Wnt-5/pipetail functions in vertebrate axis formation as a negative regulator of Wnt/beta-catenin activity. *J Cell Biol* 162:889–898
- Wetzel R (1929) Untersuchungen am Hühnchen. Die Entwicklung des Keims Während der Ersten Beiden Bruttage. In: Untersuchungen am Hühnchen. Springer, Berlin, Heidelberg, pp 188–321
- Willert K, Brown J, Danenberg E et al (2003) Wnt proteins are lipid-modified and can act as stem cell growth factors. *Nature* 423:448–452

- Williams M, Burdsal C, Periasamy A et al (2011) Mouse primitive streak forms in situ by initiation of epithelial to mesenchymal transition without migration of a cell population. *Dev Dyn* 241:270–283. doi:[10.1002/dvdy.23711](https://doi.org/10.1002/dvdy.23711)
- Williams M, Yen W, Lu X, Sutherland A (2014) Distinct apical and basolateral mechanisms drive planar cell polarity-dependent convergent extension of the mouse neural plate. *Dev Cell* 29:34–46. doi:[10.1016/j.devcel.2014.02.007](https://doi.org/10.1016/j.devcel.2014.02.007)
- Wills AE, Choi VM, Bennett MJ et al (2010) BMP antagonists and FGF signaling contribute to different domains of the neural plate in *Xenopus*. *Dev Biol* 337:335–350. doi:[10.1016/j.ydbio.2009.11.008](https://doi.org/10.1016/j.ydbio.2009.11.008)
- Wilson E (1928) *The cell in development and heredity*, 3rd edn. The Macmillian Company, New York
- Wilson PA, Hemmati-Brivanlou A (1995) Induction of epidermis and inhibition of neural fate by Bmp-4. *Nature* 376:331–333. doi:[10.1038/376331a0](https://doi.org/10.1038/376331a0)
- Wilson SI, Rydstrom A, Trimborn T et al (2001) The status of Wnt signalling regulates neural and epidermal fates in the chick embryo. *Nature* 411:325–330. doi:[10.1038/35077115](https://doi.org/10.1038/35077115)
- Wilson V, Olivera-Martinez I, Storey KG (2009) Stem cells, signals and vertebrate body axis extension. *Development* 136:1591–1604. doi:[10.1242/dev.021246](https://doi.org/10.1242/dev.021246)
- Winklbauer R, Nagel M (1991) Directional mesoderm cell migration in the *Xenopus* gastrula. *Dev Biol* 148:573–589
- Winklbauer R, Selchow A, Nagel M, Angres B (1992) Cell interaction and its role in mesoderm cell migration during *Xenopus* gastrulation. *Dev Dyn* 195:290–302
- Winnier G, Blessing M, Labosky PA, Hogan BL (1995) Bone morphogenetic protein-4 is required for mesoderm formation and patterning in the mouse. *Genes Dev* 9:2105–2116
- Witschi E (1956) Proposals for an international agreement on normal stages in vertebrate embryology. In: XIV international congress of zoology, Copenhagen, pp 260–262
- Wolda SL, Moody CJ, Moon RT (1993) Overlapping expression of Xwnt-3A and Xwnt-1 in neural tissue of *Xenopus laevis* embryos. *Dev Biol* 155:46–57. doi:[10.1006/dbio.1993.1005](https://doi.org/10.1006/dbio.1993.1005)
- Wolpert L (1969) Positional information and the spatial pattern of cellular differentiation. *J Theor Biol* 25:1–47
- Wolpert L (1971) Positional information and pattern formation. *Curr Top Dev Biol* 6:183–224
- Wöhrle S, Wallmen B, Hecht A (2007) Differential control of Wnt target genes involves epigenetic mechanisms and selective promoter occupancy by T-cell factors. *Mol Cell Biol* 27:8164–8177. doi:[10.1128/MCB.00555-07](https://doi.org/10.1128/MCB.00555-07)
- Wu C-I, Hoffman JA, Shy BR et al (2012a) Function of Wnt/ $\beta$ -catenin in counteracting Tcf3 repression through the Tcf3- $\beta$ -catenin interaction. *Development* 139:2118–2129. doi:[10.1242/dev.076067](https://doi.org/10.1242/dev.076067)
- Wu G, Huang H, Garcia Abreu J, He X (2009) Inhibition of GSK3 phosphorylation of beta-catenin via phosphorylated PPPSPXS motifs of Wnt coreceptor LRP6. *PLoS One* 4:e4926. doi:[10.1371/journal.pone.0004926](https://doi.org/10.1371/journal.pone.0004926)
- Wu S-Y, Shin J, Sepich DS, Solnica-Krezel L (2012b) Chemokine GPCR signaling inhibits  $\beta$ -catenin during zebrafish axis formation. *PLoS Biol* 10:e1001403. doi:[10.1371/journal.pbio.1001403](https://doi.org/10.1371/journal.pbio.1001403)
- Wylie CC, Kofron M, Payne C et al (1996) Maternal beta-catenin establishes a “dorsal signal” in early *Xenopus* embryos. *Development* 122:2987–2996
- Wynn ML, Kulesa PM, Schnell S (2012) Computational modelling of cell chain migration reveals mechanisms that sustain follow-the-leader behaviour. *J R Soc Interface* 9:1576–1588. doi:[10.1002/dvdy.22612](https://doi.org/10.1002/dvdy.22612)
- Xanthos JB, Kofron M, Tao Q et al (2002) The roles of three signaling pathways in the formation and function of the Spemann Organizer. *Development* 129:4027–4043
- Xu PF, Houssin N, Ferri-Lagneau KF et al (2014) Construction of a vertebrate embryo from two opposing morphogen gradients. *Science* 344:87–89. doi:[10.1126/science.1248252](https://doi.org/10.1126/science.1248252)
- Xue Y, Zheng X, Huang L, et al. (2014) Organizer-derived Bmp2 is required for the formation of a correct Bmp activity gradient during embryonic development. *Nat Commun*. doi:[10.1038/ncomms4766](https://doi.org/10.1038/ncomms4766)



- Yabe T, Shimizu T, Muraoka O, Bae YK, Hirata T, Nojima H, Kawakami A, Hirano T, Hibi M (2003) Ogon/secreted frizzled functions as a negative feedback regulator of Bmp signaling. *Development* 130:2705–2716
- Yamada T (1938) Induktion der sekundären Embryonalanlage im Neunaugenkeim. *Okajimas Folia Anatomica Japonica*
- Yamaguchi N, Mizutani T, Kawabata K, Haga H (2015) Leader cells regulate collective cell migration via Rac activation in the downstream signaling of integrin  $\beta$ 1 and PI3K. *Sci Rep* 5:7656. doi:[10.1038/srep07656](https://doi.org/10.1038/srep07656)
- Yamamoto A, Nagano T, Takehara S et al (2005) Shisa promotes head formation through the inhibition of receptor protein maturation for the caudalizing factors, Wnt and FGF. *Cell* 120:223–235. doi:[10.1016/j.cell.2004.11.051](https://doi.org/10.1016/j.cell.2004.11.051)
- Yamamoto H (1999) Phosphorylation of axin, a Wnt signal negative regulator, by glycogen synthase kinase-3 $\beta$  regulates its stability. *J Biol Chem* 274:10681–10684. doi:[10.1074/jbc.274.16.10681](https://doi.org/10.1074/jbc.274.16.10681)
- Yamamoto M, Beppu H, Takaoka K et al (2009) Antagonism between Smad1 and Smad2 signaling determines the site of distal visceral endoderm formation in the mouse embryo. *J Cell Biol* 184:323–334. doi:[10.1038/361543a0](https://doi.org/10.1038/361543a0)
- Yamamoto M, Meno C, Sakai Y et al (2001) The transcription factor FoxH1 (FAST) mediates Nodal signaling during anterior-posterior patterning and node formation in the mouse. *Genes Dev* 15:1242–1256. doi:[10.1101/gad.883901](https://doi.org/10.1101/gad.883901)
- Yamamoto M, Saijoh Y, Perea-Gomez A et al (2004) Nodal antagonists regulate formation of the anteroposterior axis of the mouse embryo. *Nature* 428:387–392. doi:[10.1038/nature02418](https://doi.org/10.1038/nature02418)
- Yang J, Tan C, Darken RS et al (2002) Beta-catenin/Tcf-regulated transcription prior to the mid-blastula transition. *Development* 129:5743–5752
- Yao L-C, Blitz IL, Peiffer DA et al (2006) Schnurri transcription factors from *Drosophila* and vertebrates can mediate Bmp signaling through a phylogenetically conserved mechanism. *Development* 133:4025–4034. doi:[10.1242/dev.02561](https://doi.org/10.1242/dev.02561)
- Ybot-Gonzalez P, Savery D, Gerrelli D et al (2007) Convergent extension, planar-cell-polarity signalling and initiation of mouse neural tube closure. *Development* 134:789–799. doi:[10.1242/dev.000380](https://doi.org/10.1242/dev.000380)
- Yen WW, Williams M, Periasamy A et al (2009) PTK7 is essential for polarized cell motility and convergent extension during mouse gastrulation. *Development* 136:2039–2048. doi:[10.1242/dev.030601](https://doi.org/10.1242/dev.030601)
- Yeo C, Whitman M (2001) Nodal signals to Smads through Cripto-dependent and Cripto-independent mechanisms. *Mol Cell* 7:949–957
- Yi F, Pereira L, Hoffman JA et al (2011) Opposing effects of Tcf3 and Tcf1 control Wnt stimulation of embryonic stem cell self-renewal. *Nat Cell Biol* 13:762–770. doi:[10.1038/ncb2283](https://doi.org/10.1038/ncb2283)
- Yi F, Pereira L, Merrill BJ (2008) Tcf3 functions as a steady-state limiter of transcriptional programs of mouse embryonic stem cell self-renewal. *Stem Cells* 26:1951–1960. doi:[10.1634/stemcells.2008-0229](https://doi.org/10.1634/stemcells.2008-0229)
- Yin C, Kiskowski M, Pouille PA et al (2008) Cooperation of polarized cell intercalations drives convergence and extension of presomitic mesoderm during zebrafish gastrulation. *J Cell Biol* 180:221–232. doi:[10.1016/S1534-5807\(04\)00060-7](https://doi.org/10.1016/S1534-5807(04)00060-7)
- Yokota C, Kofron M, Zuck M et al (2003) A novel role for a nodal-related protein; Xnr3 regulates convergent extension movements via the FGF receptor. *Development* 130:2199–2212
- Yoshikawa S, McKinnon R, Kokel M, Thomas J (2003) Wnt-mediated axon guidance via the *Drosophila* Derailed receptor. *Nature* 422:583–588
- Yu J-K, Satou Y, Holland ND et al (2007) Axial patterning in cephalochordates and the evolution of the organizer. *Nature* 445:613–617. doi:[10.1038/nature05472](https://doi.org/10.1038/nature05472)
- Yuge M, Kobayakawa Y, Fujisue M, Yamana K (1990) A cytoplasmic determinant for dorsal axis formation in an early embryo of *Xenopus laevis*. *Development* 110:1051–1056
- Zecca M, Basler K, Struhl G (1996) Direct and long-range action of a wingless morphogen gradient. *Cell* 87:833–844

- Zeng X, Huang H, Tamai K et al (2008) Initiation of Wnt signaling: control of Wnt coreceptor Lrp6 phosphorylation/activation via frizzled, dishevelled and axin functions. *Development* 135:367–375. doi:[10.1242/dev.013540](https://doi.org/10.1242/dev.013540)
- Zeng X, Tamai K, Doble B et al (2005) A dual-kinase mechanism for Wnt co-receptor phosphorylation and activation. *Nature* 438:873–877. doi:[10.1038/nature04185](https://doi.org/10.1038/nature04185)
- Zernicka-Goetz M (1998) Fertile offspring derived from mammalian eggs lacking either animal or vegetal poles. *Development* 125:4803–4808
- Zernicka-Goetz M (2013) Development: do mouse embryos play dice? *Curr Biol* 23:R15–R17. doi:[10.1016/j.cub.2012.10.032](https://doi.org/10.1016/j.cub.2012.10.032)
- Zernicka-Goetz M, Morris SA, Bruce AW (2009) Making a firm decision: multifaceted regulation of cell fate in the early mouse embryo. *Nat Rev Genet* 10:467–477. doi:[10.1038/nrg2564](https://doi.org/10.1038/nrg2564)
- Zhang J, Houston DW, King ML et al (1998) The role of maternal VegT in establishing the primary germ layers in *Xenopus* embryos. *Cell* 94:515–524
- Zhang S, Holland ND, Holland LZ (1997) Topographic changes in nascent and early mesoderm in amphioxus embryos studied by DiI labeling and by in situ hybridization for a Brachyury gene. *Dev Genes Evol* 206:532. doi:[10.1007/s004270050083](https://doi.org/10.1007/s004270050083)
- Zhang X, Abreu JG, Yokota C et al (2012) Tiki1 is required for head formation via Wnt cleavage-oxidation and inactivation. *Cell* 149:1565–1577. doi:[10.1016/j.cell.2012.04.039](https://doi.org/10.1016/j.cell.2012.04.039)
- Zhang X, Cheong S-M, Amado NG et al (2015) Notum is required for neural and head induction via wnt deacylation, oxidation, and inactivation. *Dev Cell* 32:719–730. doi:[10.1016/j.devcel.2015.02.014](https://doi.org/10.1016/j.devcel.2015.02.014)
- Zimmerman LB, De Jesus-Escobar J, Harland R (1996) The Spemann organizer signal noggin binds and inactivates bone morphogenetic protein 4. *Cell* 86:599–606
- Ziomek CA, Johnson MH (1980) Cell surface interaction induces polarization of mouse 8-cell blastomeres at compaction. *Cell* 21:935–942
- Zisckind N, Elinson R (1990) Gravity and microtubules in dorsoventral polarization of the *Xenopus* egg. *Dev Growth Differ* 32:575–581