

Approximation Algorithm for the Minimum Directed Tree Cover

Viet Hung Nguyen

LIP6, Université Pierre et Marie Curie Paris 6, 4 place Jussieu, Paris, France

Abstract. Given a directed graph G with non negative cost on the arcs, a directed tree cover of G is a rooted directed tree such that either head or tail (or both of them) of every arc in G is touched by T . The minimum directed tree cover problem (DTCP) is to find a directed tree cover of minimum cost. The problem is known to be NP -hard. In this paper, we show that the weighted Set Cover Problem (SCP) is a special case of DTCP. Hence, one can expect at best to approximate DTCP with the same ratio as for SCP. We show that this expectation can be satisfied in some way by designing a purely combinatorial approximation algorithm for the DTCP and proving that the approximation ratio of the algorithm is $\max\{2, \ln(D^+)\}$ with D^+ is the maximum outgoing degree of the nodes in G .

1 Introduction

Let $G = (V, A)$ be a directed graph with a (non negative) cost function $c : A \Rightarrow \mathbb{Q}_+$ defined on the arcs. Let $c(u, v)$ denote the cost of the arc $(u, v) \in A$. A *directed tree cover* is a weakly connected subgraph $T = (U, F)$ such that

1. for every $e \in A$, F contains an arc f intersecting e , i.e. f and e have an end-node in common.
2. T is a rooted branching.

The *minimum directed tree cover problem* (DTCP) is to find a directed tree cover of minimum cost. Several related problems to DTCP have been investigated, in particular:

- its undirected counterpart, the minimum tree cover problem (TCP) and
- the tour cover problem in which T is a tour (not necessarily simple) instead of a tree. This problem has also two versions: undirected (ToCP) and directed (DToCP).

We discuss first about TCP which has been intensively studied in recent years. The TCP is introduced in a paper by Arkin et al. [1] where they were motivated by a problem of locating tree-shaped facilities on a graph such that all the nodes are dominated by chosen facilities. They proved the NP -hardness of TCP by observing that the unweighted case of TCP is equivalent to the *connected vertex cover* problem, which in fact is known to be as hard (to approximate)

as the vertex cover problem [10]. Consequently, DTCP is also NP -hard since the TCP can be easily transformed to an instance of DTCP by replacing every edge by the two arcs of opposite direction between the two end-nodes of the edge. In their paper, Arkin et al. presented a 2-approximation algorithm for the unweighted case of TCP, as well as 3.5-approximation algorithm for general costs. Later, Konemann et al. [11] and Fujito [8] independently designed a 3-approximation algorithm for TCP using a bidirected formulation. They solved a linear program (of exponential size) to find a vertex cover U and then they found a Steiner tree with U as the set of terminals. Recently, Fujito [9] and Nguyen [13] propose separately two different approximation algorithms achieving 2 the currently best approximation ratio. Actually, the algorithm in [13] is expressed for the TCP when costs satisfy the triangle inequality but one can suppose this for the general case without loss generality. The algorithm in [9] is very interesting in term of complexity since it is a primal-dual based algorithm and thus purely combinatorial. In the prospective section of [11] and [9], the authors presented DTCP as a wide open problem for further research on the topic. In particular, Fujito [9] pointed out that his approach for TCP can be extended to give a 2-approximation algorithm for the unweighted case of DTCP but falls short once arbitrary costs are allowed.

For ToCP, a 3-approximation algorithm has been developed in [11]. The principle of this algorithm is similar as for TCP, i.e. it solved a linear program (of exponential size) to find a vertex cover U and then found a traveling salesman tour over the subgraph induced by U . Recently, Nguyen [14] considered DToCP and extended the approach in [11] to obtain a $2 \log_2(n)$ -approximation algorithm for DToCP. We can similarly adapt the method in [11] for TCP to DTCP but we will have to find a directed Steiner tree with U a vertex cover as the terminal set. Using the best known approximation algorithm by Charikar et al. [4] for the minimum Steiner directed tree problem, we obtain a ratio of $(1 + \sqrt{|U|})^{2/3} \log^{1/3}(|U|)$ for DTCP which is worse than a logarithmic ratio.

In this paper, we improve this ratio by giving a logarithmic ratio approximation algorithm for DTCP. In particular, we show that the weighted Set Cover Problem (SCP) is a special case of DTCP and the transformation is approximation preserving. Based on the known complexity results for SCP, we can only expect a logarithmic ratio for the approximation of DTCP. Let D^+ be the maximum outgoing degree of the nodes in G , we design a primal-dual $\max\{2, \ln(D^+)\}$ -approximation algorithm for DTCP which is thus somewhat best possible.

The paper is organized as follows. In the remaining of this section, we will define the notations that will be used in the papers. In Section 2, we present an integer formulation and state a primal-dual algorithm for DTCP. Finally, we prove the validity of the algorithm and its approximation ratio.

Let us introduce the notations that will be used in the paper. Let $G = (V, A)$ be a digraph with vertex set V and arc set A . Let $n = |V|$ and $m = |A|$. If $x \in \mathbb{Q}^{|A|}$ is a vector indexed by the arc set A and $F \subseteq E$ is a subset of arcs, we use $x(F)$ to denote the sum of values of x on the arcs in F , $x(F) = \sum_{e \in F} x_e$. Similarly, for a vector $y \in \mathbb{Q}^{|V|}$ indexed by the nodes and $S \subseteq V$ is a subset of

nodes, let $y(S)$ denote the sum of values of y on the nodes in the set S . For a subset of nodes $S \subseteq V$, let $A(S)$ denote the set of the arcs having both end-nodes in S . Let $\delta^+(S)$ (respectively $\delta^-(S)$) denote the set of the arcs having only the tail (respectively head) in S . We will call $\delta^+(S)$ the *outgoing cut* associated to S , $\delta^-(S)$ the *ingoing cut* associated to S . For two subset $U, W \subset V$ such that $U \cap W = \emptyset$, let $(U : W)$ be the set of the arcs having the tail in U and the head in W . For $u \in V$, we say v an *outneighbor* (respectively *inneighbor*) of u if $(u, v) \in A$ (respectively $(v, u) \in A$). For the sake of simplicity, in clear contexts, the singleton $\{u\}$ will be denoted simply by u .

For an arc subset F of A , let $V(F)$ denote the set of end-nodes of all the arcs in F . We say F covers a vertex subset S if $F \cap \delta^-(S) \neq \emptyset$. We say F is a cover for the graph G if for all arc $(u, v) \in A$, we have $F \cap \delta^-(\{u, v\}) \neq \emptyset$.

When we work on more than one graph, we specify the graph in the index of the notation, e.g. $\delta_G^+(S)$ will denote $\delta^+(S)$ in the graph G . By default, the notations without indication of the the graph in the index are applied on G .

2 Minimum r -Branching Cover Problem

Suppose that T is a directed tree cover of G rooted in $r \in V$, i.e. T is a branching, $V(T)$ is a vertex cover in G and there is a directed path in T from r to any other node in $V(T)$. In this case, we call T , a *r -branching cover*. Thus, DTCP can be divided into n subproblems in which we find a minimum r -branching cover for all $r \in V$. By this observation, in this paper, we will focus on approximating the minimum r -branching cover for a specific vertex $r \in V$. An approximation algorithm for DTCP is then simply resulted from applying n times the algorithm for the minimum r -branching cover for each $r \in V$.

2.1 Weighted Set Cover Problem as a Special Case

Let us consider any instance \mathcal{A} of the weighted Set Cover Problem (SCP) with a set $E = \{e_1, e_2, \dots, e_p\}$ of ground elements, and a collection of subsets $S_1, S_2, \dots, S_q \subseteq E$ with corresponding non-negative weights w_1, w_2, \dots, w_q . The objective is to find a set $I \subseteq \{1, 2, \dots, q\}$ that minimizes $\sum_{i \in I} w_i$, such that $\bigcup_{i \in I} S_i = E$.

We transform this instance to an instance of the minimum r -branching cover problem in some graph G_1 as follows. We create a node r , q nodes S_1, S_2, \dots, S_q and q arcs (r, S_i) with weight w_i . We then add $2p$ new nodes e_1, \dots, e_p and e'_1, \dots, e'_p . If $e_k \in S_i$ for some $1 \leq k \leq p$ and $1 \leq i \leq q$, we create an arc (S_i, e_k) with weight 0 (or a very insignificant positive weight). At last, we add an arc (e_k, e'_k) of weight 0 (or a very insignificant positive weight) for each $1 \leq k \leq p$.

Lemma 1. *Any r -branching cover in G_1 correspond to a set cover in \mathcal{A} of the same weight and vice versa.*

Proof. Let us consider any r -branching cover T in G_1 . Since T should cover all the arcs (e_k, e'_k) for $1 \leq k \leq n$, T contains the nodes e_k . By the construction of

G_1 , these nodes are connected to r uniquely through the nodes S_1, \dots, S_q with the corresponding cost w_1, \dots, w_q . Clearly, the nodes S_i in T constitute a set cover in \mathcal{A} of the same weight as T . It is then easy to see that any set cover in \mathcal{A} correspond to r -branching cover in G_1 of the same weight.

Let D_r^+ be the maximum outgoing degree of the nodes (except r) in G_1 . We can see that $D_r^+ = p$, the number of ground elements in \mathcal{A} . Hence, we have

Corollary 1. *Any $f(D_r^+)$ -approximation algorithm for the minimum r -branching cover problem is also an $f(p)$ -approximation algorithm for SCP where f is a function from \mathbb{N} to \mathbb{R} .*

Note that the converse is not true. As a corollary of this corollary, we have the same complexity results for the minimum r -branching cover problem as known results for SCP [12,7,15,2]. Precisely,

Corollary 2

- *If there exists a $c \ln(D_r^+)$ -approximation algorithm for the minimum r -branching cover problem where $c < 1$ then $NP \subseteq DTIME(n^{O(\ln^k(D_r^+))})$.*
- *There exists some $0 < c < 1$ such that if there exists a $c \log(D_r^+)$ -approximation algorithm for the minimum r -branching cover problem, then $P = NP$.*

Note that this result does not contradict the Fujito’s result about an approximation ratio 2 for the unweighted DTCP because in our transformation we use arcs of weight 0 (or a very insignificant fractional positive weight) which are not involved in an instance of unweighted DTCP.

Hence in some sense, the $\max\{2, \ln(D_r^+)\}$ approximation algorithm that we are going to describe in the next sections seems to be best possible for the general weighted DTCP.

3 Integer Programming Formulation for Minimum r -Branching Cover

We use a formulation inspired from the one in [11] designed originally for the TCP. The formulation is as follows: for a fixed root r , define \mathcal{F} to be the set of all subsets S of $V \setminus \{r\}$ such that S induces at least one arc of A ,

$$\mathcal{F} = \{S \subseteq V \setminus \{r\} \mid A(S) \neq \emptyset\}.$$

Let T be the arc set of a directed tree cover of G containing r , T is thus a branching rooted at r . Now for every $S \in \mathcal{F}$, at least one node, saying v , in S should belong to $V(T)$. By definition of directed tree cover there is a path from r to v in T and as $r \notin S$, this path should contain at least one arc in $\delta^-(S)$. This allows us to derive the following *cut* constraint which is valid for the DTCP:

$$\sum_{e \in \delta^-(S)} x_e \geq 1 \text{ for all } S \in \mathcal{F}$$

This leads to the following IP formulation for the minimum r -branching cover.

$$\begin{aligned} & \min \sum_{e \in A} c(e)x_e \\ & \sum_{e \in \delta^-(S)} x_e \geq 1 \text{ for all } S \in \mathcal{F} \\ & x \in \{0, 1\}^A. \end{aligned}$$

A trivial case for which this formulation has no constraint is when G is a r -rooted star but in this case the optimal solution is trivially the central node r with cost 0.

Replacing the integrity constraints by

$$x \geq 0,$$

we obtain the linear programming relaxation. We use the $DTC(G)$ to denote the convex hull of all vectors x satisfying the constraints above (with integrity constraints replaced by $x \geq 0$). We express below the dual of $DTC(G)$:

$$\begin{aligned} & \max \sum_{S \in \mathcal{F}} y_S \\ & \sum_{S \in \mathcal{F} \text{ s.t. } e \in \delta^-(S)} y_S \leq c(e) \text{ for all } e \in A \\ & y_S \geq 0 \text{ for all } S \in \mathcal{F} \end{aligned}$$

4 Approximating the Minimum r -Branching Cover

4.1 Preliminary Observations and Algorithm Overview

Preliminary observations. As we can see, the minimum r -branching cover is closely related to the well-known minimum r -arborescence problem which finds a minimum r -branching spanning all the nodes in G . Edmonds [6] gave a linear programming formulation for this problem which consists of the cut constraints for all the subsets $S \subseteq V \setminus \{r\}$ (not limited to $S \in \mathcal{F}$). He designed then a primal-dual algorithm (also described in [5]) which repeatedly keeps and updates a set A_0 of zero reduced cost and the subgraph G_0 induced by A_0 and at each iteration, tries to cover a chosen strongly connected component in G_0 by augmenting (as much as possible with respect to the current reduced cost) the corresponding dual variable. The algorithm ends when all the nodes are reachable from r in G_0 . The crucial point in the Edmonds' algorithm is that when there still exist nodes not reachable from r in G_0 , there always exists in G_0 a strongly connected component to be covered because we can choose trivial strongly connected components which are singletons. We can not do such a thing for minimum r -branching cover because a node can be or not belonging to a r -branching cover. But we shall see that if G_0 satisfies a certain conditions, we can

use an Edmonds-style primal-dual algorithm to find a r -branching cover and to obtain a G_0 satisfying such conditions, we should pay a ratio of $\max\{2, \ln(n)\}$. Let us see what could be these conditions. A node j is said *connected to* another node i (resp. a connected subgraph B) if there is a path from i (resp. a node in B) to j . Suppose that we have found a vertex cover U and a graph G_0 , we define an *Edmonds connected subgraph* as a non-trivial connected (not necessarily strongly) subgraph B not containing r of G_0 such that given any node $i \in B$ and for all $v \in B \cap U$, v is connected to i in G_0 . Note that any strongly connected subgraph not containing r in G_0 which contains at least a node in U is an Edmonds connected subgraph. As in the definition, for an Edmonds connected subgraph B , we will also use abusively B to denote its vertex set.

Theorem 1. *If for any node $v \in U$ not reachable from r in G_0 , we have*

- *either v belongs to an Edmonds connected subgraph of G_0 ,*
- *or v is connected to an Edmonds connected subgraph of G_0 .*

then we can apply an Edmonds-style primal-dual algorithm completing G_0 to get a r -branching cover spanning U without paying any additional ratio.

Proof. We will prove that if there still exist nodes in U not reachable from r in G_0 , then there always exists an Edmonds connected subgraph, say B , uncovered, i.e. $\delta_{G_0}^-(B) = \emptyset$. Choosing any node $v_1 \in U$ not reachable from r in G_0 , we can see that in both cases, the Edmonds connected subgraph, say B_1 , of G_0 containing v_1 or to which v_1 is connected, is not reachable from r . In this sense we suppose that B_1 is maximal. If B_1 is uncovered, we have done. If B_1 is covered then it should be covered by an arc from a node $v_2 \in U$ not reachable from r because if $v_2 \notin U$ then $B_1 \cup \{v_2\}$ induces an Edmonds connected subgraph which contradicts the fact that B_1 is maximal. Similarly, we should have $v_2 \neq v_1$ because otherwise $B_1 \cup \{v_1\}$ induces an Edmonds connected subgraph. We continue this reasoning with v_2 , if this process does not stop, we will meet another node $v_3 \in U \setminus \{v_1, v_2\}$ not reachable from r and so on As $|U| \leq n - 1$, this process should end with an Edmonds connected subgraph B_k uncovered.

We can then apply a primal-dual Edmonds-style algorithm (with respect to the reduced cost modified by the determination of U and G_0 before) which repeatedly cover in each iteration an uncovered Edmonds connected subgraph in G_0 until every node in U is reachable from r . By definition of Edmonds connected subgraphs, in the output r -branching cover, we can choose only one arc entering the chosen Edmonds connected subgraph and it is enough to cover the nodes belonging to U in this subgraph.

Algorithm overview. Based on the above observations on $DTC(G)$ and its dual, we design an algorithm which is a composition of 3 phases. Phases I and II determine G_0 and a vertex cover U satisfying the conditions stated in Theorem 1. The details of each phase is as follows:

- Phase I is of a primal-dual style which tries to cover the sets $S \in \mathcal{F}$ such that $|S| = 2$. We keep a set A_0 of zero reduced cost and the subgraph G_0 induced by A_0 . A_0 is a cover but does not necessarily contain a r -branching

cover. We determine after this phase a vertex cover (i.e. a node cover) set U of G . Phase I outputs a partial solution T_0^1 which is a directed tree rooted in r spanning the nodes in U reachable from r in G_0 . It outputs also a dual feasible solution y .

- Phase II is executed only if A_0 does not contain a r -branching cover, i.e. there are nodes in U determined in Phase I which are not reachable from r in G_0 . Phase II works with the reduced costs issued from Phase I and tries to make the nodes in U not reachable from r in G_0 , either reachable from r in G_0 , or belong or be connected to an Edmonds connected subgraph in G_0 . Phase II transforms this problem to a kind of Set Cover Problem and solve it by a greedy algorithm. Phase II outputs a set of arcs T_0^2 and grows the dual solution y issued from Phase I (by growing only the zero value components of y).
- Phase III is executed only if $T_0^1 \cup T_0^2$ is not a r -branching cover. Phase III applies a primal-dual Edmonds-style algorithm (with respect to the reduced cost issued from Phases I and II) which repeatedly cover in each iteration an uncovered Edmonds connected subgraph in G_0 until every node in U is reachable from r .

4.2 Initialization

Set \mathcal{B} to be the collection of the vertex set of all the arcs in A which do not have r as an end vertex. In other words, \mathcal{B} contains all the sets of cardinality 2 in \mathcal{F} , i.e. $\mathcal{B} = \{S \mid S \in \mathcal{F} \text{ and } |S| = 2\}$. Set the dual variable to zero, i.e. $y \leftarrow 0$ and set the reduced cost \bar{c} to c , i.e. $\bar{c} \leftarrow c$. Set $A_0 \leftarrow \{e \in A \mid \bar{c}(e) = 0\}$. Let $G_0 = (V_0, A_0)$ be the subgraph of G induced by A_0 .

During the algorithm, we will keep and update constantly a subset of $T_0 \subseteq A_0$. At this stage of initialization, we set $T_0 \leftarrow \emptyset$.

During Phase I, we also keep updating a dual feasible solution y that is initialized at 0 (i.e. all the components of y are equal to 0). The dual solution y is not necessary in the construction of a r -branching cover but we will need it in the proof for the performance guarantee of the algorithm.

4.3 Phase I

In this phase, we will progressively expand A_0 so that it covers all the sets in \mathcal{B} . In the mean time, during the expansion of A_0 , we add the vertex set of newly created strongly connected components of G_0 to \mathcal{B} .

Phase I repeatedly do the followings until \mathcal{B} becomes empty.

1. select a set $S \in \mathcal{B}$ which is not covered by A_0 .
2. select the cheapest (reduced cost) arc(s) in $\delta^-(S)$ and add it (them) to A_0 . A_0 covers then S . Let α denote the reduced cost of the cheapest arc(s) chosen above, then we modify the reduced cost of the arcs in $\delta^-(S)$ by subtracting α from them. Set $y_S \leftarrow \alpha$.
3. Remove S from \mathcal{B} and if we detect a strongly connected component K in G_0 due to the addition of new arcs in A_0 , in the original graph G , we add the set $V(K)$ to \mathcal{B} .

Proposition 1. *After Phase I, A_0 is a cover.*

Proof. As we can see, Phase I terminates when \mathcal{B} becomes empty. That means the node sets of the arcs, which do not have r as an end-node, are all covered by A_0 . Also all the strongly connected components in G_0 are covered. \square

At this stage, if for any node v there is a path from r to v in G_0 , we say that v is *reachable* from r . Set T_0 to be a directed tree (rooted in r) in G_0 spanning the nodes reachable from r . T_0 is chosen such that for each strongly connected component K added to \mathcal{B} in Phase I, there is exactly one arc in T_0 entering K , i.e. $|\delta^-(K) \cap T_0| = 1$. If the nodes reachable from r in G_0 form a vertex cover, then T_0 is a r -branching cover and the algorithm stops. Otherwise, it goes to Phase II.

4.4 Phase II

Let us consider the nodes which are not reachable from r in G_0 . We divide them into three following categories:

- The nodes i such that $|\delta_{G_0}^-(i)| = 0$, i.e. there is no arc in A_0 entering i . Let us call these nodes *source nodes*.
- The nodes i such that $|\delta_{G_0}^-(i)| = 1$, i.e. there is exactly one arc in A_0 entering i . Let us call these nodes *sink nodes*.
- The nodes i such that $|\delta_{G_0}^-(i)| \geq 2$, i.e. there is at least two arcs in A_0 entering i . Let us call these nodes *critical nodes*.

Proposition 2. *The set of the source nodes is a stable set.*

Proof. Suppose that the converse is true, then there is an arc (i, j) with i, j are both source nodes. As $\delta_{G_0}^-(i) = \delta_{G_0}^-(j) = \emptyset$, we have $\delta_{G_0}^-(\{i, j\}) = \emptyset$. Hence, (i, j) is not covered by A_0 . Contradiction.

Corollary 3. *The set U containing the nodes reachable from r in G_0 after Phase I, the sink nodes and the critical nodes is a vertex cover (i.e. a node cover) of G .*

Proposition 3. *For any sink node j , there is at least one critical node i such that j is connected to i in G_0 .*

Proof. Let the unique arc in $\delta_{G_0}^-(j)$ be (i_1, j) . Since this arc should be covered by A_0 , $\delta_{G_0}^-(i_1) \neq \emptyset$. If $|\delta_{G_0}^-(i_1)| \geq 2$ then i_1 is a critical node and we have done. Otherwise, i.e. $|\delta_{G_0}^-(i_1)| = 1$ and i_1 is a sink node. Let (i_2, i_1) be the unique arc in $\delta_{G_0}^-(i_1)$, we repeat then the same reasoning for (i_2, i_1) and for i_2 . If this process does not end with a critical node, it should meet each time a new sink node not visited before (It is not possible that a directed cycle is created since then this directed cycle (strongly connected component) should be covered in Phase I and hence at least one of the nodes on the cycle has two arcs entering it, and is therefore critical). As the number of sink nodes is at most $n - 1$, the process can not continue infinitely and should end at a stage k ($k < n$) with i_k is a critical node. By construction, the path $i_k, i_{k-1}, \dots, i_1, j$ is a path in G_0 from i_k to j .

A critical node v is said to be *covered* if there is at least one arc $(w, v) \in A_0$ such that w is not a source node, i.e. w can be a sink node or a critical node or a node reachable from r . Otherwise, we say v is *uncovered*.

Proposition 4. *If all critical nodes are covered then for any critical node v , one of the followings is verified:*

- *either v belongs to an Edmonds connected subgraph of G_0 or v is connected to an Edmonds connected subgraph of G_0 ,*
- *there is a path from r to v in G_0 , i.e. v is reachable from r in G_0 .*

Proof. If v is covered by a node reachable from r , we have done. Otherwise, v is covered by sink node or by another critical node. From Proposition 3 we derive that in the both cases, v will be connected to a critical node w , i.e. there is a path from w to v in G_0 . Continue this reasoning with w and so on, we should end with a node reachable from r or a critical node visited before. In the first case v is reachable from r . In the second case, v belongs to a directed cycle in G_0 if we have revisited v , otherwise v is connected to a directed cycle in G_0 . The directed cycle in the both cases is an Edmonds connected subgraph (because it is strongly connected) and it can be included in a greater Edmonds connected subgraph.

Lemma 2. *If all critical nodes are covered then for any node $v \in U$ not reachable from r in G_0 ,*

- *either v belongs to an Edmonds connected subgraph of G_0 ,*
- *or v is connected to an Edmonds connected subgraph of G_0 .*

Proof. The lemma is a direct consequence of Propositions 3 and 4.

The aim of Phase II is to cover all the uncovered critical nodes. Let us see how to convert this problem into a weighted SCP and to solve the latter by adapting the well-known greedy algorithm for weighted SCP.

A source node s is *zero connecting* a critical node v (reciprocally v is *zero connected from s*) if $(s, v) \in A_0$. If $(s, v) \notin A_0$ but $(s, v) \in A$ then s is *positively connecting* v (reciprocally v is *positively connected from s*).

Suppose that at the end of Phase I, there are k uncovered critical nodes v_1, v_2, \dots, v_k and p source nodes s_1, s_2, \dots, s_p . Let $S = \{s_1, s_2, \dots, s_p\}$ denote the set of the source nodes.

Remark 1. An uncovered critical node v can be only covered:

- by directly an arc from a sink node or another critical node to v ,
- or via a source node s connecting (zero or positively) v , i.e. by two arcs: an arc in $\delta^-(s)$ and the arc (s, v) .

Remark 1 suggests us that we can consider every critical node v as a ground element to be covered in a Set Cover instance and the subsets containing v could be the singleton $\{v\}$ and any subset containing v of the set of the critical

nodes connecting (positive or zero) from s . The cost of the the singleton $\{v\}$ is the minimum reduced cost of the arcs from a sink node or another critical node to v . The cost of a subset T containing v of the set of the critical nodes connecting from s is the minimum reduced cost of the arcs in $\delta^-(s)$ plus the sum of the reduced cost of the arcs (s, w) for all $w \in T$.

Precisely, in Phase II, we proceed to cover all the uncovered critical nodes by solving by the greedy algorithm the following instance of the Set Cover Problem:

- The ground set contains k elements which are the critical nodes v_1, v_2, \dots, v_k .
- The subsets are

Type I For each source node s_i for $i = 1, \dots, p$, let $\mathcal{C}(s_i)$ be the set of all the critical nodes connected (positively or zero) from s_i . The subsets of Type I associated to s_i are the subsets of $\mathcal{C}(s_i)$ ($\mathcal{C}(s_i)$ included). To define their cost, we define

$$\bar{c}(s_i) = \begin{cases} \min\{\bar{c}(e) \mid e \in \delta^-(s_i)\} & \text{if } \delta^-(s_i) \neq \emptyset, \\ +\infty & \text{otherwise} \end{cases}$$

Let us choose an arc $e_{s_i} = \operatorname{argmin}\{\bar{c}(e) \mid e \in \delta^-(s_i)\}$ which denotes an arc entering s_i of minimum reduced cost. Let T be any subset of type I associated to s_i , we define $\bar{c}(T)$ the cost of T as $\bar{c}(T) = \bar{c}(s_i) + \sum_{v \in T} \bar{c}(s_i, v)$.

Let us call the arc subset containing the arc e_{s_i} and the arcs (s_i, v) for all $v \in T$ uncovered, *the covering arc subset of T* .

Type II the singletons $\{v_1\}, \{v_2\}, \dots, \{v_k\}$. We define the cost of the singleton $\{v_i\}$,

$$\bar{c}(v_i) = \begin{cases} \min\{\bar{c}(w, v_i) \mid \text{where } w \text{ is not a source node, i.e. } w \in V \setminus S\} & \text{if } (V \setminus S : \{v_i\}) \neq \emptyset, \\ +\infty & \text{otherwise} \end{cases}$$

Let us choose an arc $e_{v_i} = \operatorname{argmin}\{\bar{c}(w, v_i) \mid \text{where } w \text{ is not a source node, i.e. } w \in V \setminus S\}$, denotes an arc entering v_i from a non source node of minimum reduced cost. Let the singleton $\{e_{v_i}\}$ be *the covering arc subset of $\{v_i\}$* .

We will show that we can adapt the greedy algorithm solving this set cover problem to our primal-dual scheme. In particular, we will specify how to update dual variables et the sets A_0 and T_0 in each iteration of the greedy algorithm. The sketch of the algorithm is explained in Algorithm 1.

Note that in Phase II, contrary to Phase I, the reduced costs \bar{c} are not to be modified and all the computations are based on the reduced costs \bar{c} issued from Phase I. In the sequel, we will specify how to compute the most efficient subset Δ and update the dual variables.

For $1 \leq i \leq p$ let us call \mathcal{S}_i the collection of all the subsets of type I associated to s_i . Let \mathcal{S} be the collection of all the subsets of type I and II.

Algorithm 1. Greedy algorithm for Phase II

```

1 while there exist uncovered critical nodes do
2   Compute the most efficient subset  $\Delta$  ;
3   Update the dual variables and the sets  $A_0$  and  $T_0$ ;
4   Change the status of the uncovered critical nodes in  $\Delta$  to covered ;
5 end

```

Computing the most efficient subset. Given a source node s_i , while the number of subsets in \mathcal{S}_i can be exponential, we will show in the following that computing the most efficient subset in \mathcal{S}_i is can be done in polynomial time. Let us suppose that there are i_q critical nodes denoted by $v_{s_i}^{i_1}, v_{s_i}^{i_2}, \dots, v_{s_i}^{i_q}$ which are connected (positively or zero) from s_i . In addition, we suppose without loss of generality that $\bar{c}(s_i, v_{s_i}^{i_1}) \leq \bar{c}(s_i, v_{s_i}^{i_2}) \leq \dots \leq \bar{c}(s_i, v_{s_i}^{i_q})$. We compute f_i and S_i which denote respectively the best efficiency and the most effecient set in \mathcal{S}_i by the following algorithm.

Step 1. Suppose that $v_{s_i}^{i_h}$ is the first uncovered critical node met when we scan the critical nodes $v_{s_i}^{i_1}, v_{s_i}^{i_2}, \dots, v_{s_i}^{i_q}$ in this order.
 Set $S_i \leftarrow \{v_{s_i}^{i_h}\}$. Set $\bar{c}(S_i) \leftarrow \bar{c}(s_i) + \bar{c}(s_i, v_{s_i}^{i_h})$.
 Set $d_i \leftarrow 1$. Set $f_i \leftarrow \frac{\bar{c}(S_i)}{d_i}$ and $\Delta_i \leftarrow S_i$.

Step 2. We add progressively uncovered critical nodes $v_{s_i}^{i_j}$ for $j = h + 1, \dots, i_q$ to S_i while this allows to increase the efficiency of S_i :
 For $j = h + 1$ to i_q , if $v_{s_i}^{i_j}$ is uncovered and $f_i > \frac{\bar{c}(S_i) + \bar{c}(s_i, v_{s_i}^{i_j})}{d_i + 1}$ then $f_i \leftarrow \frac{\bar{c}(S_i) + \bar{c}(s_i, v_{s_i}^{i_j})}{d_i + 1}$, $d_i \leftarrow d_i + 1$ and $S_i \leftarrow S_i \cup \{v_{s_i}^{i_j}\}$.

Set $i_{min} \leftarrow \operatorname{argmin}\{f_i \mid s_i \text{ is a source node}\}$.

Choose the most efficient subset among $S_{i_{min}}$ and the singletons of type II for which the computation of efficiency is straightforward. Set Δ to be most efficient subset and set $d \leftarrow |\Delta|$ the number of the uncovered critical nodes in Δ .

Updating the dual variables and the sets A_0 and T_0

Let $g = \max\{|T| \mid T \in \mathcal{S}\}$ and let $H_g = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{g}$.

Remark 2. $g \leq D_r^+$.

Given a critical node v , let p_v denote the number of source nodes connecting v . Let $s_1^v, s_2^v, \dots, s_{p_v}^v$ be these source nodes such that $\bar{c}(s_1^v, v) \leq \bar{c}(s_2^v, v) \leq \dots \leq \bar{c}(s_{p_v}^v, v)$. We define $S_v^j = \{v, s_1^v, \dots, s_j^v\}$ for $j = 1, \dots, p_v$. We can see that for $j = 1, \dots, p_v$, $S_v^j \in \mathcal{F}$. Let $y_{S_v^j}$ be the dual variable associated to the cut constraints $x(\delta^-(S_v^j)) \geq 1$. The dual variables will be updated as follows. For each critical node v uncovered in Δ , we update the value of $y_{S_v^j}$ for $j = 1, \dots, p_v$ for that $\sum_{j=1}^{p_v} y_{S_v^j} = \frac{\bar{c}(\Delta)}{H_g \times d}$. This updating process saturates progressively the arcs (s_j^v, v) for $j = 1, \dots, p_v$. Details are given in Algorithm 2. We add to A_0 and to T_0 the arcs in the covering arc subset of Δ .

Algorithm 2. Updating the dual variables

```

1  $j \leftarrow 1$  ;
2 while ( $j < p_v$ ) and ( $\bar{c}(s_v^{j+1}, v) < \frac{\bar{c}(\Delta)}{H_g \times d}$ ) do
3      $y_{S_v^j} \leftarrow \bar{c}(s_v^{j+1}, v) - \bar{c}(s_v^j, v)$ ;
4      $j \leftarrow j + 1$  ;
5 end
6 if  $\bar{c}(s_v^{p_v}, v) < \frac{\bar{c}(\Delta)}{H_g \times d}$  then
7      $y_{S_v^{p_v}} \leftarrow \frac{\bar{c}(\Delta)}{H_g \times d} - \bar{c}(s_v^{p_v}, v)$ ;
8 end

```

Let us define \mathcal{T} as the set of the subsets T such that y_T is made positive in Phase II.

Lemma 3. *The dual variables which were made positive in Phase II respect the reduced cost issued from Phase I.*

Proof. For every $T \in \mathcal{T}$, the arcs in $\delta^-(T)$ can only be either an arc in $\delta^-(s_i)$ with s_i is a source node or an arc in $\delta^-(v)$ with v is a critical node. Hence, we should show that for every arc (u', u) with u is either a critical node or a source node, we have

$$\sum_{T \in \mathcal{T} \text{ s.t. } u \in T} y_T \leq \bar{c}(u', u)$$

– u is a critical node v and u' is the source node s_j^v . The possible subsets $T \in \mathcal{T}$ such that $(s_j^v, v) \in \delta^-(T)$ are the sets S_v^1, \dots, S_v^{j-1} . By Algorithm 2, we can see that

$$\sum_{k=1}^{j-1} y_{S_v^k} \leq \bar{c}(s_j^v, v).$$

– u is a critical node v and $u' \in V \setminus S$. By definition of $\bar{c}(v)$, we have $\bar{c}(u', u) \geq \bar{c}(v)$. By analogy with the Set Cover problem, the dual variables made positive in Phase II respect the cost of the singleton $\{v\}$. Hence

$$\sum_{T \in \mathcal{T} \text{ s.t. } v \in T} y_T \leq \bar{c}(v) \leq \bar{c}(u', u)$$

– u is source node and $u' \in V \setminus S$. For each critical node w such that $(u, w) \in A$, we suppose that $u = s_w^{i(u,w)}$ where $1 \leq i(u, w) \leq p_w$. Let

$$T_u = \{w \mid w \text{ is a critical node, } (u, w) \in A \text{ and } y_{S_w^{i(u,w)}} > 0\}$$

We can see that $T_u \in \mathcal{S}$ and $\bar{c}(T_u) = \bar{c}(u) + \sum_{w \in T_u} \bar{c}(u, w)$. Suppose that l is the total number of iterations in Phase II. We should show that

$$\sum_{k=1}^l \sum_{w \in T_u \cap \Delta_k} \left(\frac{\bar{c}(\Delta_k)}{H_g \times d_k} - \bar{c}(u, w) \right) \leq \bar{c}(u) \tag{1}$$

where Δ_k is the subset which has been chosen in k^{th} iteration. Let a_k be the number of uncovered critical nodes in T_u at the beginning of the k^{th} iteration. We have then $a_1 = |T_u|$ and $a_{l+1} = 0$. Let A_k be the set of previously uncovered critical nodes of T_u covered in the k^{th} iteration. We immediately find that $|A_k| = a_k - a_{k+1}$. By Algorithm 1, we can see that at the k^{th} iteration $\frac{\bar{c}(\Delta_k)}{H_g \times d_k} \leq \frac{\bar{c}(T_u)}{H_g \times a_k}$. Since $|A_k| = a_k - a_{k+1}$ then

$$\sum_{w \in T_u \cap \Delta_k} \left(\frac{\bar{c}(\Delta_k)}{H_g \times d_k} \right) - \sum_{w \in T_u \cap \Delta_k} \bar{c}(u, w) \leq \frac{\bar{c}(T_u)}{H_g} \times \frac{a_k - a_{k+1}}{a_k} - \sum_{w \in T_u \cap \Delta_k} \bar{c}(u, w)$$

Hence,

$$\begin{aligned} \sum_{k=1}^l \sum_{w \in T_u \cap \Delta_k} \left(\frac{\bar{c}(\Delta_k)}{H_g \times d_k} - \bar{c}(u, w) \right) &\leq \frac{\bar{c}(T_u)}{H_g} \sum_{k=1}^l \frac{a_k - a_{k+1}}{a_k} - \sum_{k=1}^l \sum_{w \in T_u \cap \Delta_k} \bar{c}(u, w) \\ &\leq \frac{\bar{c}(T_u)}{H_g} \sum_{k=1}^l \left(\frac{1}{a_k} + \frac{1}{a_k - 1} + \dots + \frac{1}{a_{k+1} - 1} \right) \\ &\quad - \sum_{k=1}^l \sum_{w \in T_u \cap \Delta_k} \bar{c}(u, w) \\ &\leq \frac{\bar{c}(T_u)}{H_g} \sum_{i=1}^{a_1} \frac{1}{i} - \sum_{k=1}^l \sum_{w \in T_u \cap \Delta_k} \bar{c}(u, w) \\ &\leq \bar{c}(T_u) - \sum_{k=1}^l \sum_{w \in T_u \cap \Delta_k} \bar{c}(u, w) = \bar{c}(u). \end{aligned}$$

Let $T_0^2 \subset T_0$ the set of the arcs added to T_0 in Phase II. For each $e \in T_0^2$, let $c_2(e)$ be the part of the cost $c(e)$ used in Phase II.

Theorem 2

$$c_2(T_0) = \sum_{e \in T_0^2} c_2(e) \leq H_g \sum_{T \in \mathcal{T}} y_T \leq \ln(D_r^+) \sum_{T \in \mathcal{T}} y_T$$

Proof. By Algorithm 2, at the k^{th} iteration, a subset Δ_k is chosen and we add the arcs in the covering arc subset of Δ_k to T_0^2 for all $v \in \Delta_k$. Let T_0^{2k} be covering arc subset of Δ_k . We can see that $c_2(T_0^{2k}) = \sum_{e \in T_0^{2k}} \bar{c}_e = \bar{c}(\Delta_k)$. In this iteration, we update the dual variables in such a way that for each critical node $v \in \Delta_k$, $\sum_{j=1}^{p_v} y_{S_v^j} = \frac{\bar{c}(\Delta_k)}{H_g \times d_k}$ with $d_k = |\Delta_k|$. Together with the fact that $\bar{c}(\Delta_k) = \bar{c}(w_k) + \sum_{v \in \Delta_k} \bar{c}(w_k, v)$ we have $\sum_{v \in \Delta_k} \sum_{j=1}^{p_v} y_{S_v^j} = \frac{\bar{c}(\Delta_k)}{H_g} = \frac{c_2(T_0^{2k})}{H_g}$. By summing over l be the number of iterations in Phase II, we obtain

$$\sum_{T \in \mathcal{T}} y_T = \sum_{k=1}^l \sum_{v \in \Delta_k} \sum_{j=1}^{p_v} y_{S_v^j} = \sum_{k=1}^l \frac{\bar{c}(\Delta_k)}{H_g} = \sum_{k=1}^l \frac{c_2(T_0^{2k})}{H_g} = \frac{c_2(T_0)}{H_g}$$

which proves that $c_2(T_0) = H_g \sum_{T \in \mathcal{T}} y_T$. By Remark 2, we have $g \leq D_r^+$ and $H_g \approx \ln g$, hence $c_2(T_0) \leq \ln(D_r^+) \sum_{T \in \mathcal{T}} y_T$. \square

4.5 Phase III

We perform Phase III if after Phase II, there exist nodes in U not reachable from r in G_0 . By Lemma 2, they belong or are connected to some Edmonds connected subgraphs of G_0 . By Theorem 1, we can apply an Edmonds-style primal-dual algorithm which tries to cover uncovered Edmonds connected subgraphs of G_0 until all nodes in U reachable from r . The algorithm repeatedly choosing uncovered Edmonds connected subgraph and adding to A_0 the cheapest (reduced cost) arc(s) entering it. As the reduced costs have not been modified during Phase II, we update first the reduced cost \bar{c} with respect to the dual variables made positive in Phase II.

Algorithm 3. Algorithm for Phase III

- 1 Update the reduced cost \bar{c} with respect to the dual variables made positive in Phase II;
 - 2 **repeat**
 - 3 Choose B an uncovered Edmonds connected subgraph ;
 - 4 Let y_B be the associated dual variable to B ;
 - 5 Set $\bar{c}(B) \leftarrow \min\{\bar{c}_e \mid e \in \delta^-(B)\}$; Set $y_B \leftarrow \bar{c}(B)$;
 - 6 **foreach** $e \in \delta^-(B)$ **do**
 - 7 $\bar{c}_e \leftarrow \bar{c}_e - \bar{c}(B)$;
 - 8 **end**
 - 9 Update A_0, G_0 and T_0 (see below);
 - 10 **until** every nodes in U reachable from r ;
-

For updating A_0 , at each iteration, we add all the saturated arcs belonging to $\delta^-(B)$ to A_0 . Among these arcs, we choose only one arc (u, v) with $v \in B$ to add to T_0 with a preference for a u connected from r in G_0 . In the other hand, we delete the arc (x, v) with $x \in B$ from T_0 . We then add to T_0 an directed tree rooted in v in G_0 spanning B . If there are sink nodes directly connected to B , i.e. the path from a critical node $w \in B$ to these nodes contains only sink nodes except w . We also add all such paths to T_0 .

Lemma 4. *After Phase III, T_0 is a r -branching cover.*

Proof. We can see that after Phase III, for any critical node or a sink node v , there is a path containing only the arcs in T_0 from r to v and there is exactly one arc in $\delta^-(v) \cap T_0$.

4.6 Performance Guarantee

We state now a theorem about the performance guarantee of the algorithm.

Theorem 3. *The cost of T_0 is at most $\max\{2, \ln(D_r^+)\}$ times the cost of an optimal r -branching cover.*

Proof. Suppose that T^* is an optimal r -branching cover of G with respect to the cost c . First, we can see that the solution y built in the algorithm is feasible dual solution. Hence $c^T y \leq c(T^*)$. Let \mathcal{B} be the set of all the subsets B in Phase I and Phase III (B is either a subset of cardinality 2 in \mathcal{F} or a subset such that the induced subgraph is a strongly connected component or an Edmonds connected subgraph in G_0 at some stage of the algorithm). Recall that we have defined \mathcal{T} as the set of the subsets T such that y_T is made positive in Phase II. We have then $c^T y = \sum_{B \in \mathcal{B}} y_B + \sum_{T \in \mathcal{T}} y_T$. For any arc e in T_0 , let us divide the

cost $c(e)$ into two parts: $c_1(e)$ the part saturated by the dual variables y_B with $B \in \mathcal{B}$ and $c_2(e)$ the part saturated by the dual variables y_T with $T \in \mathcal{T}$. Hence $c(T_0) = c_1(T_0) + c_2(T_0)$. By Theorem 2, we have $c_2(T_0) \leq \ln(D_r^+) \sum_{T \in \mathcal{T}} y_T$ (note that the replacing in Phase III of an arc (x, v) by another arc (u, v) with $v \in B_i$ do not change the cost $c_2(T_0)$). Let us consider any set $B \in \mathcal{B}$ by the algorithm, B is the one of the followings:

- $|B| = 2$. As T_0 is a branching so that for all vertex $v \in V$, we have $|\delta^-(v) \cap T_0| \leq 1$. Hence, $|\delta^-(B) \cap T_0| \leq 2$.
- B is a vertex set of a strongly connected component or an Edmonds connected subgraph in G_0 . We can see obviously that by the algorithm $|\delta^-(B) \cap T_0| = 1$.

These observations lead to the conclusion that $c_1(T_0) \leq 2 \sum_{B \in \mathcal{B}} y_B$. Hence

$$\begin{aligned} c(T_0) &= c_1(T_0) + c_2(T_0) \leq 2 \sum_{B \in \mathcal{B}} y_B + \ln(D_r^+) \sum_{T \in \mathcal{T}} y_T \\ &\leq \max\{2, \ln(D_r^+)\} c^T y \leq \max\{2, \ln(D_r^+)\} c(T^*). \end{aligned}$$

Corollary 4. *We can approximate the DTCP within a $\max\{2, \ln(D^+)\}$ ratio.*

5 Final Remarks

The paper has shown that the weighted Set Cover Problem is a special case of the Directed Tree Cover Problem and the latter can be approximated with a ratio of $\max\{2, \ln(D^+)\}$ (where D^+ is the maximum outgoing degree of the nodes in G) by a primal-dual algorithm. Based on known complexity results for weighted Set Cover, in one direction, this approximation seems to be best possible.

In our opinion, an interesting question is whether the same techniques can be applied to design a combinatorial approximation algorithm for Directed Tour Cover. As we have seen in Introduction section, a $2 \log_2(n)$ -approximation algorithm for Directed Tour Cover has been given in [14], but this algorithm is not combinatorial.

References

1. Arkin, E.M., Halldórsson, M.M., Hassin, R.: Approximating the tree and tour covers of a graph. *Information Processing Letters* 47, 275–282 (1993)
2. Arora, S., Sudan, M.: Improved Low-Degree Testing and Its Applications. In: *Proceedings of STOC 1997*, pp. 485–495 (1997)
3. Bock, F.: An algorithm to construct a minimum spanning tree in a directed network. In: *Developments in Operations Research*, pp. 29–44. Gordon and Breach, NY (1971)
4. Charikar, M., Chekuri, C., Cheung, T., Dai, Z., Goel, A., Guha, S., Li, M.: Approximation Algorithms for Directed Steiner Problems. *Journal of Algorithms* 33, 73–91 (1999)
5. Chu, Y.J., Liu, T.H.: On the shortest arborescence of a directed graph. *Science Sinica* 14, 1396–1400 (1965)
6. Edmonds, J.: Optimum branchings. *J. Research of the National Bureau of Standards* 71B, 233–240 (1967)
7. Feige, U.: A threshold of $\ln n$ for approximating set cover. *Journal of the ACM* 45, 634–652 (1998)
8. Fujito, T.: On approximability of the independent/connected edge dominating set problems. *Information Processing Letters* 79, 261–266 (2001)
9. Fujito, T.: How to Trim an MST: A 2-Approximation Algorithm for Minimum Cost Tree Cover. In: Bugliesi, M., Preneel, B., Sassone, V., Wegener, I. (eds.) *ICALP 2006*. LNCS, vol. 4051, pp. 431–442. Springer, Heidelberg (2006)
10. Garey, M.R., Johnson, D.S.: The rectilinear Steiner-tree problem is NP complete. *SIAM J. Appl. Math.* 32, 826–834 (1977)
11. Könemann, J., Konjevod, G., Parekh, O., Sinha, A.: Improved Approximations for Tour and Tree Covers. *Algorithmica* 38, 441–449 (2003)
12. Lund, C., Yannakakis, M.: On the hardness of approximating minimization problems. *Journal of the ACM* 41, 960–981 (1994)
13. Nguyen, V.H.: Approximation algorithms for metric tree cover and generalized tour and tree covers. *RAIRO Operations Research* 41(3), 305–315 (2007)
14. Nguyen, V.H.: A $2 \log_2(n)$ -Approximation Algorithm for Directed Tour Cover. In: *Proceedings of COCOA 2009*. LNCS, vol. 5573, pp. 208–218. Springer, Heidelberg (2009)
15. Raz, R., Safra, R.: A sub-constant error-probability low-degree test, and a sub-constant error-probability PCP characterization of NP. In: *Proceedings of STOC 1997*, pp. 475–484 (1997)