

Jae Young Jang

Contents

10.1	Principles of Endoscopic Diagnosis of Gastric Polyps	170
10.1.1	Definitions	170
10.1.2	Clinical Manifestations	171
10.1.3	Endoscopic Features	171
10.2	Nonneoplastic Polyps	173
10.2.1	Hyperplastic Polyps	173
10.2.2	Fundic Gland Polyps	175
10.2.3	Inflammatory Fibroid Polyp	176
10.2.4	Xanthoma	178
10.2.5	Hamartomatous Polyps and Polyps Associated with Polyposis Syndrome	179
10.3	Neoplastic Polyps	182
10.3.1	Adenoma/Dysplasia	182
10.3.2	Gastric Polyps and Familial Adenomatous Polyposis	183
	References	195

Gastric polyps can be defined as luminal lesions projecting above the mucosal surface. Gastric polyps are detected in 2–3 % of upper gastrointestinal endoscopies. Gastric polyps are generally divided into nonneoplastic and neoplastic types. About 80–90 % of gastric polyps are nonneoplastic. Hyperplastic polyps, fundic gland polyps, inflammatory fibroid polyps, and hamartomatous polyps are nonneoplastic. Hyperplastic polyps may be sessile or pedunculated. Larger hyperplastic polyps may have small erosions or ulcerations on the surface. Fundic gland polyps appear usually as multiple, glassy, transparent, sessile polyps. Inflammatory fibroid polyps are well-circumscribed solitary, small sessile, or pedunculated lesions with regular pit pattern. Hamartomatous polyps are usually pedunculated, round polyps with hyperemic and exudative surfaces. Dysplastic epithelium is defined as noninvasive, neoplastic epithelium. Gastric adenomas are defined as circumscribed, polypoid dysplastic lesions. Flat tubular adenomas are also called dysplasias. Endoscopically, adenomas/dysplasias are reddish or whitish discolored, often with a multilobulated surface, contrasting with smooth and atrophic adjacent mucosa. The surface may be smooth or superficially eroded.

J.Y. Jang
 Division of Gastroenterology,
 Department of Internal Medicine,
 College of Medicine, Kyung Hee University, Seoul, Korea
 e-mail: jjjang@khu.ac.kr

10.1 Principles of Endoscopic Diagnosis of Gastric Polyps

10.1.1 Definitions

Gastric polyps can be defined broadly as luminal lesions projecting above the plane of the mucosal surface. It is

recommended that discrete protrusions identified on radiology or endoscopy be referred to as polypoid lesions until histological diagnoses are made. Various subtypes of gastric polyps are recognized; they are generally divided into non-neoplastic and neoplastic. About 80–90 % of gastric polyps are nonneoplastic. They can be divided into epithelial and nonepithelial types (Table 10.1).

Table 10.1 Classification of gastric polyps

	Epithelial polyp	Nonepithelial polyp
Nonneoplastic polyps	Hyperplastic polyp Fundic gland polyp Inflammatory fibroid polyp Hamartomatous polyp Cronkhite-Canada syndrome	
Neoplastic polyps	Adenoma/dysplasia Low grade High grade Carcinoma (primary or secondary) Carcinoid tumor	
Miscellaneous lesions		Xanthelasma Lymphoid hyperplasia/lymphoma Mesenchymal stromal tumors Gastrointestinal tumors Smooth muscle tumors Glomus tumor Neural tumors Schwannoma/neuroma Ganglioneuromas Granular cell tumor Other rare tumors Lipoma/liposarcoma Rhabdomyosarcoma Fibrous histiocytoma Vascular Hemangioma/lymphangioma Hemangiosarcoma

10.1.2 Clinical Manifestations

Gastric polyps are detected in 2–3 % of upper gastrointestinal endoscopic examinations, often incidentally. They only rarely produce symptoms, such as gastrointestinal bleeding and delayed gastric emptying. Rarely, a prolapsing antral polyp may obstruct the gastric outlet.

10.1.3 Endoscopic Features

They are usually small, with a diameter of less than 1–2 cm. The endoscopic appearance of gastric polyps is variable,

ranging from slightly raised plaques to soft multilobulated nodules to, more rarely, broad-based or sessile lesions. In noninvasive neoplastic lesions and carcinomas with invasion of the lamina propria, called superficial lesions, flat or even depressed morphologies below the contour of the mucosa may be observed. In pedunculated polyps, the base is narrow; in sessile polyps, the base and the top of the lesion have the same diameter. Intermediate and broad-based forms are called semi-pedunculated. Non-protruding or non-polypoid neoplastic lesions include ulcers and the so-called flat lesions. In the latter situation, the lesion, compared with the adjacent mucosa, is either slightly elevated or completely flat or depressed (Table 10.2, Fig. 10.1) [1].

Table 10.2 Neoplastic lesions with superficial morphology (Paris endoscopic classification)

	Morphology	
Polypoid	Ip	Pedunculated
	Is	Sessile
Non-polypoid	IIa	Slightly elevate
	IIb	Flat
	IIc	Slightly depressed
	III	Excavated (ulcer)




Fig. 10.1 Morphological classification of gastric polyps. (a) Ip polyp, (b) Is polyp, (c) Ila polyp, (d) I Ib polyp, (e) I Ic polyp

10.2 Nonneoplastic Polyps

10.2.1 Hyperplastic Polyps

These benign lesions are the second most common type of gastric polyp after fundic gland polyps. They are generally thought to result from excessive regeneration following mucosal damage and, consequently, commonly occur in chronic *Helicobacter pylori*-associated gastritis, in

pernicious anemia, adjacent to ulcers and erosions, or at gastroenterostomy sites. Hyperplastic polyps can occur throughout the stomach. They are usually less than 1.5 cm in diameter, may be single or multiple, and may be sessile or pedunculated. Usually, the pits of polyp surface are enlarged and hyperemic in color [2]. With larger polyps, the overlying mucosa is often red and friable, and there may be a small erosion or ulceration on the tip of the polyp (Fig. 10.2).




Fig. 10.2 Hyperplastic polyps. (a) A 6-mm, Is-type polyp. (b) A 7-mm, Ip-type polyp. (c) A 2-cm, Ip-type polyp with hemorrhagic change. (d) A 2-cm, Ip-typed, hyperemic polyp. (e–g) Ip-type polyps with erosions covered with exudate and/or hemorrhagic changes




Fig.10.2 (continued)

10.2.2 Fundic Gland Polyps

Fundic gland polyps are seen as single, multiple, sessile lesions that occur uniquely in the fundus and upper body of the stomach. The morphology is characteristic, with cystic

transformation of the gland lined by parietal cells and chief cells. Endoscopically, they appear as glassy, transparent, sessile polyps, less than 1 cm in diameter (Fig. 10.3). These are usually multiple and may be found in males and females of any age [3].




Fig. 10.3 Fundic gland polyp. (a, b) Several 2–3-mm diminutive, glassy, transparent, sessile polyps. (c) A 5-mm subpedunculated, transparent polyp at the gastric body

10.2.3 Inflammatory Fibroid Polyp

Inflammatory fibroid polyps may be found throughout the gastrointestinal tract, but they are most common in the antrum and pyloric region. These polyps are characterized by the proliferation of spindle cells, small blood vessels, and

inflammatory cells, often dominated by eosinophils. Endoscopically, they are well-circumscribed, solitary, small sessile or pedunculated lesions with regular pit patterns. With larger polyps, there may be erosions on the surface of the lesion, and they are often observed as subepithelial tumors with bridging folds (Fig. 10.4).




Fig. 10.4 Inflammatory fibroid polyp. (a, b) Is-type, round polyp of similar surface color as the surroundings, with no discoloration. (c) Iia-type, round polyp with small erosion. (d, e) Is-type polyp with surface erosion observed as a subepithelial tumor with bridging folds




Fig.10.4 (continued)

10.2.4 Xanthoma

These clinically insignificant lesions are found with increasing age and are often associated with chronic gastritis and

intestinal metaplasia. Grossly, they are single or multiple, 1–2 mm in diameter, round or oval, well-circumscribed, yellow, macular, or nodular lesions (Fig. 10.5). They are found most frequently along the lesser curvature.




Fig. 10.5 Xanthoma. (a, b) A 3-mm flat, round, yellowish macular lesion at the body. (c) A 5-mm yellowish polypoid lesion

10.2.5 Hamartomatous Polyps and Polyps Associated with Polyposis Syndrome

Gastric hamartomatous polyps are composed of hyperplastic glands lined by foveolar-type epithelium and separated by

branching cores of smooth muscle, with atrophy of the deep glandular components. Endoscopic features of hamartomatous polyps are pedunculated round polyps with hyperemic and exudative surfaces (Fig. 10.6).




Fig. 10.6 Hamartomatous polyp. (a, b) A 6–8-mm sessile polyps with exudative changes at the body. (c) A 1.5-cm hyperemic, exudative, pedunculated polyp at the antrum

10.2.5.1 Peutz-Jeghers Polyp

Gastric polyps occur in approximately 49 % of Peutz-Jeghers polyposis patients. However, they may also occur sporadically. Rare reports of gastric carcinoma in Peutz-Jeghers polyposis have been described.

10.2.5.2 Juvenile Polyp

Gastric juvenile polyps are rarely observed and often occur within the onset of juvenile polyposis, either of the stomach alone or of the entire gastrointestinal tract. The gastric polyps are usually multiple (Fig. 10.7). They may present at any age. Juvenile polyposis is associated with an increased risk of cancer, particularly in the colon, but the stomach also may be at risk.




Fig. 10.7 Juvenile polyposis in the stomach. Multiple, variable sized, and sessile polyps are noted at the whole stomach

10.2.5.3 Cowden's Disease Polyps

Cowden's disease is also known as multiple hamartoma syndrome. The patients have an increased risk of breast and thyroid carcinoma. The incidence of gastrointestinal polyps is estimated to be close to 75 % of patients.

10.2.5.4 Cronkhite-Canada Polyp

These polyps occur usually in conjunction with lesions in other parts of the gastrointestinal tract. They are indistinguishable from juvenile and hyperplastic polyps (Fig. 10.8) and can be diagnosed only in the presence of clinical evidence of alopecia, nail atrophy, or hyperpigmentation.




Fig. 10.8 Cronkhite-Canada polyps in the stomach. Multiple, slightly elevated, and hyperemic polyps are noted at the entire stomach

10.3 Neoplastic Polyps

10.3.1 Adenoma/Dysplasia

10.3.1.1 Definition

Dysplastic epithelium is defined as noninvasive, neoplastic epithelium. Gastric adenomas are defined by the World Health Organization as circumscribed, polypoid lesions composed of tubular and/or villous structures, lined by dysplastic epithelium [4]. In Western literature, flat tubular adenomas are frequently called dysplasias. In the Vienna consensus classification [5] for intramucosal neoplasia, the

terms adenoma and dysplasia are both replaced by “intraepithelial neoplasia” (Table 10.3).

10.3.1.2 Clinical Significance

Gastric adenomas/dysplasias are precancerous lesions, comparable to colonic adenomas. Gastric adenoma/dysplasia can be subdivided according to the degree of dysplasia (low or high grade). The risk of malignancy in gastric adenomas/dysplasias is related to size, degree of dysplasia, and villosity of growth pattern. High-grade dysplasia has been identified in close proximity to 40–100 % of early gastric cancers and 5–80 % of advanced adenocarcinomas [6].

Table 10.3 The revised Vienna classification of gastrointestinal epithelial neoplasia

Category	Diagnosis
1	Negative for neoplasia
2	Indefinite for neoplasia
3	Mucosal low-grade neoplasia Low-grade adenoma Low-grade dysplasia
4	Mucosal high-grade neoplasia 4.1 High-grade adenoma/dysplasia 4.2 Noninvasive carcinoma (carcinoma in situ) 4.3 Suspicious for invasive carcinoma 4.4 Intramucosal carcinoma
5	Submucosal invasion by carcinoma

10.3.1.3 Endoscopic Features

Adenomas/dysplasias are reddish or whitish discolored, often with a multilobulated surface, contrasting with smooth and atrophic adjacent mucosa. The surface may be smooth or superficially eroded (Figs. 10.9, 10.10, 10.11, and 10.12). The presence of erosion or ulceration is thought to correlate with an increased risk of cancer. Similarly, superficial erosion and atypia may increase the likelihood of a gastric cancer being present. According to a recent study, a characteristic white opaque substance [7] was found along the mucosal surface when an elevated gastric adenoma was observed with magnified endoscopy with narrow band imaging (NBI, Fig. 10.13). High-grade adenoma/dysplasia tends to be associated with

a more eroded surface, hyperemic changes, and depressed type than low-grade adenoma/dysplasia (Fig. 10.14).

10.3.2 Gastric Polyps and Familial Adenomatous Polyposis

Familial adenomatous polyposis is inherited as an autosomal dominant trait, characterized by the development of multiple adenomatous polyps in the colon. Fundic gland polyps are found in 13–53 % of patients; adenomas, which are more common in the antrum, are found in 6–12 %.




Fig. 10.9 Low-grade adenoma. (a, b) Ila-type, whitish discolored, flat, elevated lesions at the angle. (c–e) Antrum. (f–i) Body




Fig. 10.9 (continued)




Fig. 10.10 Low-grade adenoma. (a–c) IIa-type, whitish discolored, lobulated, elevated lesions at the antrum. (d–e) Angle. (f–h) Body




Fig. 10.10 (continued)




Fig. 10.11 Low-grade dysplasia. (a–c) IIB-type, whitish discolored, flat, well-circumscribed lesions at the antrum. (d–e) Angle




Fig. 10.12 Low-grade dysplasia. Indistinct margined dysplasias due to marked atrophy and intestinal metaplasia around the lesions are often observed. Indigo carmine chromoendoscopy delineated the lesion more

clearly. (a, c, e, g, i, k, m) White light endoscopy. (b, d, f, h, j, l, n) Indigo carmine chromoendoscopy




Fig. 10.12 (continued)




Fig.10.12 (continued)




Fig. 10.13 White opaque substances in NBI. (a, c) A whitish discolored low-grade dysplasia at the antrum. (b, d) White opaque substances along the pit at the magnified endoscopy with NBI




Fig. 10.14 High-grade adenoma/dysplasia. (a) A 1.0-cm Is-type polypoid lesion with surface exudate and hyperemic change. (b, c) A whitish discolored, elevated lesions at the antrum. (d) A 4-mm flat erosion. (e) A 2-cm elevated lesion with central erosive and hyperemic change

at the angle. (f) A 1.5-cm flat, well-circumscribed, hyperemic lesion at the body. (g–k) Whitish discolored lesions with erosion. (l, m) Depressed lesions. (n, o) Lesions with depression and hyperemic change




Fig. 10.14 (continued)




Fig.10.14 (continued)

References

1. The Paris endoscopic classification of superficial neoplastic lesions: esophagus, stomach, and colon: November 30 to December 1, 2002. *Gastrointest Endosc.* 2003;58:S3–43.
2. Oberhuber G, Stolte M. Gastric polyps: an update of their pathology and biological significance. *Virchows Arch.* 2000;437:581–90.
3. Park do Y, Lauwers GY. Gastric polyps: classification and management. *Arch Pathol Lab Med.* 2008;132:633–40.
4. Goldstein NS, Lewin KJ. Gastric epithelial dysplasia and adenoma: historical review and histological criteria for grading. *Hum Pathol.* 1997;28:127–33.
5. Dixon MF. Gastrointestinal epithelial neoplasia: Vienna revisited. *Gut.* 2002;51:130–1.
6. Lauwers GY, Srivastava A. Gastric preneoplastic lesions and epithelial dysplasia. *Gastroenterol Clin North Am.* 2007;36:813–29.
7. Yao K, Iwashita A, Tanabe H, et al. White opaque substance within superficial elevated gastric neoplasia as visualized by magnification endoscopy with narrow-band imaging: a new optical sign for differentiating between adenoma and carcinoma. *Gastrointest Endosc.* 2008;68:574–80.