
A Review of the Geology, Structural Controls, and Tectonic Setting of Copahue Volcano, Southern Volcanic Zone, Andes, Argentina

1

A. Folguera, E. Rojas Vera, L. Vélez, J. Tobal, D. Orts, M. Agosto, A. Caselli and V.A. Ramos

Abstract

Copahue Volcano lies in the Southern Volcanic Zone of the Andes Mountains, although its geology and local structural controls differ from nearby active volcanic centers. Most of its geology is substantially older than active volcanoes at these latitudes, as the postglacial component is relatively minor. The basement of Copahue Volcano, represented by the Agrio Caldera products and its basal sections, accumulated in extensional depocenters when the arc narrowed from a broad geometry on both sides of the Andes to its present configuration. Initial stages comprise early Pliocene basaltic-andesitic eruptions associated with extensional (trans-tensional?) processes that ended with the formation of a series of rhombohedral calderas that emitted important amounts of ignimbrites in latest Pliocene-early Pleistocene time. Copahue Volcano concentrates the Pleistocene activity of one of these calderas, the Agrio Caldera, before the emplacement and development of the Present arc front to the west. Volcano morphology reflects this particular evolution, looking more degraded than Antuco, Callaqui and Lonquimay volcanoes located immediately to the west in the arc front. Most of Copahue's volume is early Pleistocene in age, showing a thin resurfacing cover in synglacial (>27 ka) and postglacial times. A synglacial stage occurred mainly to the east of Copahue Volcano toward the caldera interior in a series of independent, mostly monogenetic centers. Postglacial eruptions occurred

A. Folguera (✉) · E. Rojas Vera · L. Vélez · J. Tobal
D. Orts · M. Agosto · V.A. Ramos
Facultad de Ciencias Exactas y Naturales, Instituto
de Estudios Andinos (IDEAN), Universidad de
Buenos Aires-CONICET, Buenos Aires, Argentina
e-mail: folguera@gl.fcen.uba.ar

A. Caselli
Facultad de Ciencias Exactas y Naturales,
Universidad de Río Negro e Instituto de Estudios
Andinos (IDEAN), Universidad de Buenos Aires,
Buenos Aires, Argentina

as both central and fissural emissions reactivating the old Pleistocene conduits. Its particular geological record and eastern longitudinal position indicate that Copahue was probably part of the late Pliocene-Pleistocene arc mostly developed in the axial and eastern Andes. Narrowing and westward retraction of the arc front, proposed in previous works for the last 5 Ma at 38°S, could have been the result of the eastward migration of the asthenospheric wedge during slab steepening. Reasons for this long-lived eruptive history at Copahue volcano could be related to the particular geometry of the active Liquiñe-Ofqui dextral strike-slip fault system that runs through the arc front from south to north when penetrates the retroarc area at the latitude of Copahue volcano. This behavior could be due to the collision of the oceanic Mocha plateau at these latitudes, as recently proposed. This jump and related deflection would have produced local transtensional deformation associated with abundant emissions of syn- and post-glacial products that could have partially resurfaced this long-lived center.

Keywords

Southern volcanic zone · Transtension · Northern Liquiñe-Ofqui fault zone

1.1 Introduction

Copahue volcano has been studied due to its frequent eruptions since the 1990s (see Delpino and Bermúdez 1993, 1995; González Ferrán 1994; Naranjo and Polanco 2004). Structural and tectonic models associated with this center proposed since then are summarized in this review, although it still lacks a robust stratigraphic framework over which these hypotheses are supported. The few available isotopic ages are K-Ar (Linares et al. 1999) and have significant errors (see Melnick et al. 2006a). Available geological maps are based on these ages and proposed lateral correlations that could be inappropriate for volcanic environments. Dellapé and Pando (1975) and Pesce (1989) were the first attempts to represent the different units of the Agrio Caldera, where the Copahue Volcano is hosted. Latter representations were more focused on studying the associated structural controls

than the volcanic stratigraphy itself (Folguera and Ramos 2000; Melnick et al. 2006a; Rojas Vera et al. 2009a).

The most outstanding feature associated with Copahue volcano is the Agrio Caldera (Pesce 1989), also denominated in other works as the Caviahue Caldera. This 15 × 20 km depression was initially interpreted as the product of strong glacial erosion (Groeber 1921) based on the finding of presumably related diamictites. However, these were recently reinterpreted as large mass wasting deposits emplaced from the perimeters of the caldera towards its interior (Hermanns et al. 2011). The identification of this depression during the 1920s constituted an important discovery since no remote images existed at that time. Its linkage with volcanic processes waited until the late 1980s when Pesce (1989) identified the perimeter of the caldera and the location of Copahue volcano in one of the main collapse scars. Other resurgent centers,

partially coeval to Copahue volcano, have been identified along the caldera perimeters that have been erupted in the last 1 Ma (Linares et al. 1999).

A description of the state of knowledge of the tectonic setting associated with Copahue volcano and the Agrio Caldera is presented, including a discussion about the peculiarities of this center, which is morphologically distinctive and not aligned with the arc front at these latitudes.

1.2 Tectonic Framework at 37–39°S and Sublithospheric Structure

Recently released seismic tomography developed after the Maule earthquake has shed light on the sublithospheric structure of the Southern Volcanic Zone where Copahue volcano is located (Fig. 1.1) (Pesicek et al. 2012). The Nazca plate is illuminated as a zone of fast P-wave velocities, with some complexity previously not noted. Between




Fig. 1.1 **a b** Pliocene-Early Pleistocene and Pleistocene-Holocene arc activity in the Main Andes between 37–39°S (Lara et al. 2001; Lara and Folguera 2006), and seismic tomography of the retroarc zone modified from Pesicek et al. (2012). **c d** Pale blue and blue colors indicate P-wave velocity contours that are interpreted as subducted oceanic lithosphere while white dots indicate the Benioff zone (Pesicek et al. 2012). Note that along 39°S a steeper configuration of the subducted Nazca plate is not coincident with the present

seismogenic zone. The thick dashed line in **a** and **b** indicates the tearing in the subducted Nazca plate interpreted by Pesicek et al. (2012). The Copahue Volcano is indicated with a triangle in yellow. Note that, as depicted in **B**, arc volcanoes with Holocene activity between 37° and 41°S are mainly emplaced on the western Andean slope, in comparison with late Pliocene to early Pleistocene centers developed on both sides of the Andes (see text for further details)

37° and 39°S, a tear zone appears to have developed in the subducting Nazca plate, leaving a steeper subduction angle to the south (Fig. 1.1). This tearing likely developed prior to 5–3 Ma, since the interplate seismogenic zone at 39°S, indicating the present fate of the subducted slab, is not coincident with the steeper zone of fast P-wave velocities (Fig. 1.1). Then, the present geometry of the subducting Nazca plate is considerably shallower than that steeper configuration, showing a physical continuation to the north (Pesicek et al. 2012). Thus a change in subduction geometry would have been achieved in the last 5–3 Ma between 37° and 39°S from a steeper configuration to a shallower one.

This important change has a correlation with the pattern of arc activity between 36° and 40°S. Lara et al. (2001) demonstrated that, over the last 2 Ma the arc passed from a broad and diffuse morphology, covering a wide area from the current active arc to the retroarc zone to the east, to become contracted to a narrow band in the west at the axial Andean zone (Fig. 1.1).

This westward retraction was accompanied by within-plate volcanism in the Loncopué extensional trough in the last 2 Ma (Rojas Vera et al. 2010, 2013). Eruptions at the eastern retroarc zone during Quaternary time cover an area that is coincident at surface with the Nazca slab tearing described by Pesicek et al. (2012).

Shallow structure through this segment is characterized by two distinctive systems: (i) Along the arc front, the Liquiñe-Ofqui fault system accommodates lateral displacements imposed by the oblique convergence between the Nazca and South American plates (Lavenue and Cembrano 1999) from the triple junction among the South American, Nazca and Antarctica plates (~46°S) to the Mandolegüe volcanic lineament (~38°S) (Radic 2010). From this point to the north, strike-slip displacements are absorbed at the western retroarc zone over the Argentinian territory by the Antifñir-Copahue fault system (Fig. 1.2; Folguera et al. 2004). Copahue volcano is located at the transition zone between these two neotectonic systems. (ii) To the east, over the eastern retroarc area, Pliocene to Quaternary within-plate

volcanic rocks have been related to extensional deformation (Fig. 1.3) (Kay et al. 2006; Folguera et al. 2006). Normal faults affect postglacial volcanic products, indicating young deformation of the upper crust more than 300 km east of the arc front (Fig. 1.3; Folguera et al. 2006; Rojas Vera et al. 2010). Crustal attenuation is identified beneath the arc and retroarc in coincidence with the area of Pliocene to Quaternary volcanic eruptions at 39°S and extensional deformation using receiver function techniques (Yuan et al. 2006). Density models derived from gravity data predict this area is elongate through the whole retroarc area parallel to the arc (Fig. 1.3; Folguera et al. 2012).

This crustal attenuation and associated retroarc magmatism have been explained by steepening of the Nazca plate during westward arc retraction about 5–2 Ma and consequent injection of hot asthenospheric material, after a period (13–5 Ma) of shallow subduction in the area (Kay et al. 2006; Folguera et al. 2007). This injection of hot asthenosphere material has been recently illuminated through magnetotelluric analysis (Burd et al. 2013).

1.2.1 The Mandolegüe Volcanic Lineament

This 60–70-km-long volcanic lineament is formed by the partial amalgamation of seven main volcanic centers, which from west to east, are the Callaqui Volcano, Copahue Volcano, Las Mellizas volcanic center, Trolón Volcano, Bayo Dome, and Huecú and Mandolegüe monogenetic basaltic fields (Figs. 1.4 and 1.5). This volcanic lineament is interposed between the northern termination of the Liquiñe-Ofqui and the Antifñir-Copahue fault systems along a NE direction (Fig. 1.4).

Mechanisms and ages of the different volcanic vents are highly variable along the strike of the Mandolegüe volcanic lineament (Fig. 1.5). The youngest products are dominant in the eastern sector where the Huecú and Mandolegüe volcanic fields were emplaced. There, monogenetic fields with fissural vents and, more locally,




Fig. 1.2 Tectonic framework of the Southern Volcanic Zone (modified from Hervé 1976; Cembrano et al. 1996, 2002; Hervé 1994; López Escobar et al. 1995; Lavenu and Cembrano 1999; Folguera et al. 2004). Note the location of the Mandolegüe volcanic lineament in the site where the Copahue Volcano is hosted at the northern termination

of the strike-slip Liquiñe-Ofqui fault zone and the transition to the Antiñir-Copahue fault system. Note also that the Mandolegüe volcanic lineament coincides with the site of inception of the oceanic Mocha transform zone and associated plateau into the Chilean trench

basaltic calderas are associated with broad basaltic fields that are circumscribed by the present drainage network (Fig. 1.5). This fluvial morphology has been flooded in postglacial times and probably also in historical times by reduced lacustrine basins as a response of the youngest eruptions hosted in the southern flank of the Mandolegüe volcanic lineament (Fig. 1.5) (Groeber 1928; Rojas Vera et al. 2009b).

West of the 70°40'W meridian, the average age of the different units becomes older, with pre-glacial volcanic products associated with mostly central polygenetic volcanic vents. Polygenetic stratovolcanoes are represented by the Trolón, Callaqui and Copahue volcanoes. The three centers have postglacial lava flows associated with fissural and central mechanisms, although most of its structure is pre-glacial in age (Pesce 1989; Linares et al. 1999).

In particular, the active Callaqui volcano (Fig. 1.4) is a stratovolcano elongated in a N60°E direction (Moreno and Lahsen 1986). Here, post-glacial eruptions concentrated along a 700-m-long fissure formed by 22 connected individual vents, producing an elliptical 15-km-long and 8 km-wide stratocone, as can be seen in the digital elevation model (Fig. 1.4).

Copahue volcano to the east occupies the southwestern corner of the 15 × 20 km Agrio Caldera (Pesce 1989). This has been formed over a volcanic plateau of 5–4 Ma andesitic flows associated with the Cola de Zorro-Hualcupén Formations that constitute gently sloped to flat sections up to 1,500 m thick (Fig. 1.5) (Vergara and Muñoz 1982; Niemeyer and Muñoz 1982; Muñoz Bravo et al. 1989; Suárez and Emparán 1997). The thickest sections of these packages have been described as controlled by normal




Fig. 1.3 Attenuated crust at the retroarc zone determined from a receiver function profile at 39°S (Yuan et al. 2006) where a plume-like feature is impacting against the base of the lithosphere (Burd et al. 2013) and gravimetric expression. Note that most of the volcanic plateaux at the

arc-retroarc zone are coincident with the area of high gravimetric anomalies interpreted from 3D density models as caused by crustal attenuation (Folguera et al. 2012). Gravimetric values are in mGals

faults in a broad region from 36 to 39°S, defining half-graben geometries (Folguera et al. 2006; Melnick et al. 2006b).

Locally, these sections around Copahue volcano determine a broad volcanic morphology that is dipping outward from a central point situated at the 71°W–37°50'S intersection in the center of the Agrio Caldera (Fig. 1.5). A thin cover of pyroclastic deposits of the Riscos Bayos Formation dated at 1.98 Ma (Pesce 1989; Linares et al. 1999; Mazzoni and Licitra 2000) is irregularly distributed through the external slopes of this feature, being interpreted as the outflow

ignimbrites associated with the caldera collapse (Fig. 1.6). Ignimbritic sheets of the Las Mellizas Formation occupy the inner sector of the caldera and were dated at 2.5 Ma (Pesce 1989; Melnick et al. 2006a). The northern rim of the Agrio Caldera is occupied by lava fields associated with monogenetic vents aligned along fissures and have been gathered in an informal category denominated “*Basaltos de Fondo de Valle*” (Pesce 1989) (Fig. 1.6). These lava flows, dated at 1.6–0.8 Ma (Linares et al. 1999), are controlled by the glacial morphology and project through glacial tributaries beyond the rims of the




Fig. 1.4 Digital Elevation Model that shows the surficial expression of the Callaqui-Copahue-Mandolegüe

volcanic lineament interposed between the Liquiñe-Ofqui and the Callaqui-Mandolegüe fault systems

Agrio Caldera, feeding lava fields corresponding to the El Huecú volcanic field (Fig. 1.5).

Activity less than 1 Ma at Copahue volcano occurred within the Agrio Caldera (Fig. 1.5). The volcano is formed by the amalgamation of several centers along a NE fissure, having 8–9 well preserved craters, similarly the neighboring Callaqui Volcano (Moreno and Lahsen 1986) (Fig. 1.4). The easternmost crater along the Copahue Volcano fissure has concentrated part of the postglacial eruptions as well as the 1990s and the 2000s eruptions. This center has an older section intruded by the 0.9 Ma dacitic Pucón Mahuida dome, which is in turn covered in an erosional unconformity by a younger section that can be divided in two parts, a lower pre-glacial section (Copahue Stage 1; Fig. 1.6) and an upper synglacial and postglacial section (Copahue Stage 2; Fig. 1.6). The upper synglacial section is eastwardly displaced with respect to the central Copahue vents, constituting small centers

over the southern flank of the volcano and smaller accumulations of hundred of meters of pillow lavas as independent centers. Even smaller volumes can be tracked up to the eastern rim of the caldera following a W–NW track (Fig. 1.5).

Postglacial vents are related to the volcano edifice itself along fissural systems hosted in its northern slope and central vents associated with the easternmost apical crater of the stratocone (Copahue Stage 3; Melnick et al. 2006a) (Figs. 1.5 and 1.6). Additionally, postglacial explosive products have been identified in the caldera perimeter east of the active crater (Polanco et al. 2000). The age of these products is less than 30–27 ka based on the determination of the final retreat of the ice sheet in the area. The last glacial retreat at these latitudes has been addressed from five different lines of evidence: (i) in 25.6 ± 1.2 and 23.3 ± 0.6 ka by whole-rock $^{40}\text{Ar}/^{39}\text{Ar}$, dating postglacial lava flows in the




Fig. 1.5 Geology of the Callaqui-Copahue-Mandolegüe volcanic lineament, from the Agrio Caldera to the El Huecú volcanic field. Geology of the Copahue Volcano is based on Pesce (1989), JICA (1992), Folguera and Ramos (2000), Melnick et al. (2006a) and Rojas Vera et al. (2009b). Note that most of the postglacial volcanic eruptions have been emplaced over the retroarc area in the El Huecú volcanic field, while the Copahue Volcano and Agrio Caldera have minimum resurfacing during this stage




Fig. 1.6 Stratigraphic chart of the Agrio Caldera and Copahue Volcano (after Melnick et al. 2006a; based on Pesce 1989 and Linares et al. 1999)

Laguna del Maule area immediately to the north (36°S, 70°30'W) (Singer et al. 2000); in 30 ka by dating by cosmogenic ^{36}Cl the postglacial Varvarco rock avalanche deposit (36°26'S, 70°36'W) (Costa and González Díaz 2007); in 25–30 ka by whole-rock $^{40}\text{Ar}/^{39}\text{Ar}$, dating postglacial lava flows from the Chillán Volcano (37°S)

(Dixon et al. 1999); in 30–35 ka by ^{10}Be , dating moraine deposits in the Rucachoroi valley (39°S) (Zech et al. 2008); and finally and more locally in 30 ka by ^{14}C , dating glaciofluvial deposits from Copahue volcano circumscribed to a glacial morphology (37°50'S, 71°03'W) (Bermúdez and Delpino 1999).

1.3 Structure of Copahue Volcano and Agrio Caldera

Copahue volcano is situated at the intersection point between two nearly perpendicular structural systems, the two WNW parallel Caviahue grabens and the NE contractional Chancho-Co hill (Fig. 1.7). The Caviahue grabens run through the southern section of the Agrio Caldera from the eastern inner wall to the northern Liquiñe-Ofqui fault system. These constitute symmetrical structures that broaden to the east where they are clearly separated by a small horst, whose deepest part is occupied by the Agrio lake. This extensional depression affects 2.5 Ma ignimbrites and lava flows of the Las Mellizas Formation and controls the emplacement of most of the synglacial volcanic rocks in the region (Fig. 1.7) (Melnick et al. 2006a).

The Caviahue grabens parallel the southern wall of the Agrio Caldera and are formed by a

broken structural lineament composed of WNW and NW north-facing scarps affecting early Pliocene sections of the Cola de Zorro-Hualcupén Formations (Fig. 1.7). The northern flank of the caldera is formed by similarly oriented fault systems that, together with a degraded north-facing scarp north of Copahue town, delimits the Trolope graben where 1.6–0.8 Ma lava flows of the “*Basaltos de Fondo de Valle*” were emplaced (Fig. 1.7). North and south inner walls of the Agrio Caldera are tied by NNE and NW normal fault systems constituting the eastern west-facing wall of the Agrio Caldera (Fig. 1.7). The western caldera wall is not well defined south of its intersection with the Caviahue grabens. Pesce (1989) had proposed that this fault section would have been erased by later caldera-forming processes. This caldera perimeter, composed of kilometer-long fault segments, has a rhomboidal shape with a maximum axis of 20 km in the NW direction and 15 km in the




Fig. 1.7 Structural map of the Mandolegüe volcanic lineament between Copahue Volcano and northern Loncopué trough (modified from Folguera et al. 2004). Two main structural trends interact in the Agrio Caldera, a NE

trend across the Copahue Volcano composed of reverse and strike slip structures of the Lomín and Chancho-Co structural systems and a WNW pattern of normal faults that define the two western Caviahue grabens

NE direction that suggests a tectonic origin (Folguera and Ramos 2000).

The Chancho-Co hill is north of Copahue Volcano and corresponds to a SW vergent structure that folded and uplifted Las Mellizas Formation in a broad structure in the NW extreme of the Agrio Caldera (Fig. 1.7). This anticline is affected at its southern part by the Caviahue grabens, which displace the Pliocene to Pleistocene sections beneath the basal lavas of Copahue Volcano. This relation suggests a chronology of events that starts with the uplift of the Chancho-Co hill followed by extension that led to the Caviahue grabens formation. However, SW-vergent reverse faults that uplift the Chancho-Co hill affect and displace a Pleistocene glacial pavement defining very fresh scarps, which is indicative of young deformation (Fig. 1.8). Additionally, part of these structures propagates toward Copahue town, where late Pliocene rocks are thrust over Pleistocene till deposits (Rojas Vera et al. 2009a).

Extinct and active hot springs of the Agrio Caldera are associated with the Chancho-Co structural system (see Varekamp et al. 2001, for a recent synthesis). Las Máquinas, Maquinitas, Copahue, Anfiteatro and Chancho-Co hot springs are aligned through the Chancho-Co hill frontal structures. In particular, the last lies in the upper Trapa Trapa river on the Chilean side at the intersection point between the Chancho-Co and the Caviahue structural systems.

South of the Chancho-Co hill, associated reverse faults propagate as southeast-facing scarps into the <1 Ma volcanic rocks and post-glacial products of Copahue volcano (Figs. 1.9 and 1.10). These structures act as conduits of fissural postglacial lavas flowing through the northern slope of the volcano towards the Trapa Trapa river in Chile (Figs. 1.7 and 1.10).

These two structural systems are spatially associated with indicators of ongoing deep activity in the caldera interior revealed using geophysical tools. In particular, Ibáñez et al. (2008) have measured shallow seismic activity since 2004 along the broadest and axial section

of the Caviahue grabens (Fig. 1.11). This area does not show surficial evidence of young deformation, being fault scarps strongly affected by glacial abrasion, indicating that the structure formed at least in Pleistocene time since it is affecting latest Pliocene products. Additionally, radar interferometry data have shown deflation processes previously to 2011, at the intersection point between the Caviahue grabens and the Chancho-Co hill, where fissural postglacial products are emplaced in the Trapa Trapa upper valley, (Fig. 1.11) (Vélez et al. 2011). This anomaly is shifted with respect to the axial part of Copahue volcano and coincident with an area of normal faults affecting the youngest products at its northern slope. This deflation could indicate that collapse of the Chancho-Co structural system through WNW normal structures remains active. Mass-wasting phenomena associated with the northern slope of the volcano (Figs. 1.7 and 1.10) occurred prior to the postglacial fissural eruptions and coincide with the area of noted deformation, indicating slope instability.

After 2011 a new inflationary process started with an anomaly fairly coincident with the previous slope deflation (Fig. 1.12; Vélez et al., this book). However, in a closer view, it is noted that the area that accumulated nearly 5 cm of vertical displacements in half a year shows an elliptical shape (Fig. 1.12) aligned with the scarp that is affecting and controlling the emission of post-glacial lava products (Fig. 1.10). The interferogram also shows displacements through the structures that are associated with the Chancho-Co hill suggesting its reactivation (Fig. 1.12).

1.4 Tectonic Evolution of Copahue Volcano and Agrio Caldera

Copahue volcano, as part of the Southern Volcanic Zone, has evolved through two distinctive southern Andean deformational stages. The description of these processes can explain




Fig. 1.8 Neotectonics in the basement rocks of the Copahue Volcano flanking the Chancho-Co hill. **a** Aerial picture where spatial relation between fault scarps associated with the Chancho Co hill and Copahue town

are displayed (arrows indicate the scarp traces). **b** Axial zone of the Chancho-Co hill where a glacial pavement is folded and broken by a reverse fault. **c** Detail of **b**

variable mechanics through time and distribution of the structures that were developed previously to the emplacement of Copahue volcano, particularly during the Agrio Caldera formation, and synchronously. This center is located in the drainage divide area to the east of the present arc front and its basal section is older than these neighbor active volcanoes. Additionally, the

basal section of Copahue volcano, particularly when considering the Agrio Caldera as part of its evolution, is more similar in age to the extinct Pliocene to Pleistocene centers emplaced on both sides of the Andes at these latitudes. Then, Copahue volcano could be considered as part of this old configuration with the exception that it has undergone a thin and restricted resurfacing in




Fig. 1.9 Block diagram constructed with Aster image draped on DEM (modified from Rojas Vera et al. 2009a), where neotectonic contractional structure of the Chancho-Co hill is represented in the basement of

Copahue Volcano. Compare these with ground deformation contours determined from InSar studies shown in Véllez et al. (2011)

syn- and post-glacial times. During this stage, early Pleistocene to postglacial sections have been uplifted as a contractional structure in the caldera interior. This uplift constitutes the northern continuation of the Liquiñe-Ofqui fault system, denoting a profound change in deformational mechanisms in the last 1 Ma.

1.4.1 The Pliocene to Pleistocene Evolution

The basement of Copahue Volcano and the Agrio Caldera is formed by the Cola de Zorro and Las Mellizas Formations volcanic sections. These are hosted in quadrangular-rhomboidal depocenters delimited by normal faults that contain sections up to 1,500 m (Folguera et al. 2006; Melnick et al. 2006b; Rojas Vera et al. 2009c). Younger products in the region are also associated with extensional/transensional mechanisms in the last 2–1 Ma time period, both in the Andean and extra-Andean domains. These

processes were contemporaneous to the retraction and narrowing of the volcanic arc in the last 1 Ma from the eastern to the western Andean slope (Fig. 1.1). Even though this process was initially defined as a simple westward migration of the arc front (Stern 1989; Muñoz and Stern 1988, 1989), in the last decade it was redefined on the light of new isotopic ages as an arc narrowing, from a broad geometry through both Andean slopes in Pliocene to Pleistocene times to a narrower configuration in Pleistocene to Holocene times (Fig. 1.1) (Lara et al. 2001).

This arc narrowing coincides with the area of Pliocene to early Quaternary extensional deformation between 36 and 39°S that affected the arc and retroarc areas. In particular, the Loncopué trough, considered the major extensional depocenter in the area with more than 1,500 meters accumulated in the last 5 Ma (Rojas Vera et al. 2010), is partially superimposed on the crustal attenuation processes identified in the retroarc zone from receiver function analyses by Yuan




Fig. 1.10 Above Picture of the SE facing scarp affecting the northern slope of Copahue Volcano as part of the structures that are flanking the Chanco-Co hill (below).

Below Oblique view (ikonos image taken from Google Earth) and morphological expression of the fault associated with postglacial lava flows and mass wasting deposits

et al. (2006) (Fig. 1.13). This crustal attenuation zone has produced a broad positive gravity anomaly associated with the emplacement of mantle rocks in shallower levels beneath the retroarc zone that has a latitudinal extent similar to the arc narrowing registered in the last 1 Ma and to the extensional deformed zone and related within-plate volcanic series. Asthenospheric upwelling beneath the steepening subduction zone proposed from gravity models coincides

with the anomalies in ^{13}C and ^3He detected by Augusto et al. (2013) in the hot springs located north of the Copahue Volcano that are indicative of a relatively near mantle source.

The steeper configuration of the subducted Nazca plate at these latitudes (Fig. 1.1) revealed by seismic tomographies (Pesicek et al. 2012) is considered to have developed prior to 2 Ma, and could indicate the broadening of the asthenospheric wedge coeval to crustal attenuation and




Fig. 1.11 Structural map of the Agrio Caldera and Copahue Volcano (modified from Rojas Vera et al. 2009a), with 2004s superimposed seismic events (small yellow dots; Ibáñez et al. 2008), ground deformation contours determined from satellite interferometry previous to 2011 (2002–2007) when a new inflationary process started (see next figure) (Vélez et al. 2011) and ^3He and

^{13}C anomalies determined in hot springs by Agosto et al. (2013). White stars indicate main eruptive synglacial vents. Note that synglacial activity is not coincident with the volcanic axial zone but slightly displaced to the northeast where neotectonic features have been described. Note also that seismic events are aligned with the axial zone of the Caviahue grabens

extensional processes registered in the area (Fig. 1.13). Agrio Caldera and lower sections of Copahue volcano were coeval to these processes and were circumscribed to extensional depocenters.

1.4.2 Processes Younger Than 1 Ma

WNW contractional structures involve glacial deposits and postglacial products of Copahue volcano (Fig. 1.8). Additionally, these control the emplacement of postglacial products. These structures are aligned with the Liquiñe-Ofqui fault system to the south, which abandons the arc front and penetrates at these latitudes to the Argentinian Andean slope (Figs. 1.7 and 1.14).

Reverse structures that affected postglacial products have subordinate right-lateral components displacing Quaternary deposits and landforms and producing small pull-apart depocenters (Folguera et al. 2004; Rojas Vera et al. 2009c). These structures determine the ENE Mandolegüe fault system, where the Chancho-Co hill occupies the western edge. To the north, these structural trends bend to the NW Antiñir-Copahue fault system trending through the eastern Andean front (Fig. 1.14). This system has accommodated maximum shortening in the 1.4–0.8 Ma time period, showing minor reactivations in postglacial times (Folguera et al. 2004). Its development is spatially coincident with the area of the inception of the Mocha transfer zone in the




Fig. 1.12 Inflationary processes detected by InSAR over the NE slope of Copahue Volcano since 2011 (Vélez et al. this issue). This inflation accelerated since August–October 2011 and continues after 2012–13 eruptive cycle, accumulating nearly 5 cm in 6 months. Arrows indicate fault traces of the neotectonic structure affecting

the northern slope of the volcano (see Fig. 1.10). Note that the area of higher displacements describes an ellipse with a major axis in the W–E direction surrounding the normal fault affecting postglacial products. Note also that curves of iso-displacement seem to be deflected in the north when crossing active faults in the Chancho-Co hill

Chilean trench in the last 3 Ma that could have produced an eastward wave of out-of-sequence contractional deformation through the forearc and western retroarc zones. The area of collision of the Mocha transfer zone coincides with a segment where the subducted slab penetrates with an angle 8° shallower in comparison to neighboring areas (Krawczyk et al. 2006; Tašárová 2004). Seismic tomography illuminates this shallow segment that would have been produced in the last 2–3 Ma and replaced a steeper previous configuration (Pesicek et al. 2012), showing a strong change in the subduction regime.

In this context, the Mandolegüe fault system could have officiated as a transfer zone between the Liquiñe–Ofqui and the Antiñir–Copahue fault

systems that would have operated accommodating shortening and lateral displacements on opposite sides of the Andean drainage divide (Fig. 1.14) (Folguera and Ramos 2009). This transfer zone forms a 70–80-km-long volcanic lineament that can be explained by the collision of the southern part of an oceanic plateau whose size would be similar to the Mocha plateau presently subducted south of 38°S . InSAR data show that this structure near Copahue volcano concentrates present deformation. Seismic tomography may show a steeper subduction configuration concomitant with older products and a shallower one contemporaneous to late contraction and transpressional deformation at the arc and retroarc areas.




Fig. 1.13 Retraction of the arc front during the Pliocene and Quaternary time as a function of the steepening of the Nazca plate following Stern (1989)s hypotheses (modified from Folguera et al. 2007). Attenuated crust is determined from the receiver function analyses of Yuan

et al. (2006). Note that highly conductive crustal masses from magnetotelluric studies of Brasse and Soyer (2001) are associated with the retroarc area immediately south of the Copahue Volcano area




Fig. 1.14 Hypothesis related to the latest Pliocene to Quaternary collision of the Mocha plateau and inception

of contractional and transtensional deformation in the arc zone (modified from Folguera et al. 2009)

References

- Agusto M, Tassi F, Caselli A, Vaselli O, Rouwet D, Capaccioni B, Caliro S, Chiodini G, Darrah T (2013) Gas geochemistry of the magmatic-hydrothermal fluid reservoir in the Copahue-Caviahue volcanic complex (Argentina). *J Volcanol Geoth Res* 257:44–56
- Bermúdez A, Delpino D (1999) Erupciones subglaciales y en contacto con hielo en la región volcánica de Copahue, Neuquén: Salta, Argentina. XIV Congreso Geológico Argentino, Córdoba, Argentina 2:250–253
- Brasse H, Soyer W (2001) A magnetotelluric study in the Southern Chilean Andes. *Geophys Res Lett* 28 (19):3757–3760
- Burd A, Booker J, Mackie R, Favetto A, Pomposiello M (2013) Three-dimensional electrical conductivity in the mantle beneath the Payún Matrú Volcanic Field in the Andean back-arc of Argentina near 36.5°S: decapitation of a mantle plume by resurgent upper mantle shear during slab steepening? *Geophys J Internat* (in press)
- Cembrano J, Hervé F, Lavenu A (1996) The Liqueñe Ofqui fault zone: a long-lived intra-arc fault system in southern Chile. *Tectonophysics* 259:55–66
- Cembrano J, Lavenu A, Reynolds P, Arancibia G, López G, Sanhueza A (2002) Late Cenozoic transpressional ductile deformation north of the Nazca-South America-Antarctica triple junction. *Tectonophysics* 354:289–314
- Costa CH, González Díaz EF (2007) Age constraints and paleoseismic implication of rock-avalanches in the Northern Patagonian Andes Argentina. *J S Am Earth Sci* 24(1):48–57. doi:10.1016/j.jsames.2007.03.001
- Dellapé D, Pando G (1975) Relevamiento geológico de la cuenca geotérmica de Copahue. Yacimientos Petrolíferos Fiscales Unpublished Report 524. Buenos Aires
- Delpino D, Bermúdez A (1993) La actividad volcánica del volcán Copahue durante 1992. Erupción con emisión de azufre piroclástico. Provincia de Neuquén. XII Congreso Geológico Argentino, Mendoza, Argentina 4:292–301
- Delpino D, Bermúdez A (1995) Eruptions of pyroclastic sulphur at crater lake of Copahue volcano, Argentina. XXI International union of geodesy and geophysics. Boulder, USA, B410
- Dixon HJ, Murphy MD, Sparks SJ, Chávez R, Naranjo JA, Dunkley PN, Young SR, Gilbert JS, Pringue MR (1999) The geology of Nevados de Chillán volcano Chile. *Rev Geol Chile* 26(2):227–253
- Folguera A, Ramos VA (2000) Control estructural del volcán Copahue (38°S–71°O): implicancias tectónicas para el arco volcánico cuaternario (36–39°S). *Rev Asoc Geol Arg* 55:229–244
- Folguera A, Ramos VA (2009) Collision of the Mocha fracture zone and a less than 4 Ma old wave of orogenic uplift in the Andes (36°–38°S). *Lithosphere* 1(6):364–369
- Folguera A, Ramos V, Hermanns R, Naranjo J (2004) Neotectonics in the foothills of the Southernmost Central Andes (37°–38°S). Evidence of the strike-slip displacement along the Antifir-Copahue fault zone. *Tectonics* 23 TC 5008
- Folguera A, Zapata T, Ramos VA (2006) Late Cenozoic Extension and the evolution of the Neuquén Andes.

- In: Kay SM, Ramos VA (eds) Late Cretaceous to Recent magmatism and tectonism of the Southern Andean margin at the latitude of the Neuquen basin (36–39°S). *Geol S Am S* 407:267–285
- Folguera A, Introcaso A, Giménez M, Ruiz F, Martínez P, Tunstall C, García Morabito E, Ramos VA (2007) Crustal attenuation in the Southern Andean retroarc determined from gravimetric studies (38°–39°30'S): the Lonco-Luán asthenospheric anomaly. *Tectonophysics* 439:129–147. doi:10.1016/j.tecto.2007.04.001
- Folguera A, Alasonati Tašárová Z, Götze H-J, Rojas Vera E, Giménez M, Ramos VA (2012) Retroarc extension in the last 6 Ma in the South-Central Andes (36°S–40°S) evaluated through a 3-D gravity modeling. *J S Am Earth Sci* 40:23–37
- González-Ferrán O (1994) *Volcanes de Chile*. Instituto Geográfico Militar, Santiago, pp 640
- Groeber P (1921) La región de Copahue y su glaciación diluvial. *Rev Soc Arg Estud Geogr* 1:92–110
- Groeber P (1928) Traslado del vulcanismo de la falda oriental de la cordillera hacia la ladera occidental. *An Soc Arg Estud Geogr* 3(1):210–218
- Hermanns RL, Folguera A, Penna I, González Díaz E, Fauque L, Niedermann S (2011) Landslide dams in the Central Andes of Argentina (northern Patagonia and the Argentine northwest). In: Evans SG, Hermanns RL, Strom A, Scarascia Mugnozza G (eds) *Lecture notes in the earth sciences V 133: Natural and artificial rockslide dams*. Springer, Berlin, pp 147–177
- Hervé M (1976) Estudio geológico de la falla Liqueñe-Reloncaví en el área de Liqueñe: antecedentes de un movimiento transcurrente (Provincia de Vadivía). I Congreso Geológico Chileno, *Actas B*:39–56
- Hervé F (1994) The southern Andes between 39° and 44° S latitude: the geological signature of a transpressive tectonic regime related to a magmatic arc. In: Reutter KJ, Scheuber E, Wigger PJ (eds) *Tectonics of the Southern Central Andes*. Springer, Berlin, pp 243–248
- Ibáñez J, Del Pezzo E, Bengoa C, Caselli A, Badi G, Almendros J (2008) Volcanic tremor and local earthquakes at Copahue volcanic complex, southern Andes, Argentina. *J Volcanol Geoth Res* 174:284–294
- JICA (Japan International Cooperation Agency) (1992) The feasibility study on the Northern Neuquén geothermal development project. Unpublished Ente Provincial de Energía de la Provincial del Neuquén, pp 89
- Kay SM, Burns WM, Copeland PC, Mancilla O (2006) Upper Cretaceous to Holocene magmatism and evidence for transient Miocene shallowing of the Andean subduction zone under the northern Neuquén Basin. In: Kay S, Ramos VA (eds) *Late Cretaceous to Recent magmatism and tectonism of the Southern Andean margin at the latitude of the Neuquen basin (36–39°S)*. *Geol S Am S* 407:19–60
- Krawczyk C, Mechie J, Tašárová Z, Lüth S, Stiller M, Brasse H, Echter HP, Bataille K, Wigger P, Arane-da M (2006) Geophysical signatures and active tectonics at the southern central Chilean margin. In: Oncken G et al (eds) *The andes—active subduction orogeny: frontiers in earth sciences 1*. Springer, Berlin Heidelberg New York, pp 171–192
- Lara L, Rodríguez C, Moreno H, Pérez de Arce C (2001) Geocronología K-Ar y geoquímica del vulcanismo plioceno superior-pleistoceno en los Andes del sur (39°–42°S). *Rev Geol Chile* 28:67–91
- Lara L, Folguera A (2006) Plio-Quaternary narrowing of the volcanic arc at Southern Andes (37°–41°S), southwestern margin of Neuquén Basin: geochronologic and field tectonic data evaluated. In: Kay S, Ramos VA (eds) *Late Cretaceous to Recent magmatism and tectonism of the Southern Andean margin at the latitude of the Neuquen basin (36–39°S)*. *Geol S Am S* 407:299–315
- Lavenu A, Cembrano J (1999) Compressional and transpressional stress pattern for Pliocene and Quaternary brittle deformation in fore-arc and intra-arc zones (Andes of Central and Southern Chile). *J Struct Geol* 21:1669–1691
- Linares E, Osters HA, Mas L (1999) Cronología Potasio-Argón del complejo efusivo Copahue-Caviahue, Provincia de Neuquén. *Rev Asoc Geol Arg* 54(3):240–247
- López-Escobar L, Cembrano J, Moreno H (1995) Geochemistry and tectonics of the Chilean Southern Andes Quaternary volcanism (37°–46°S). *Rev Geol Chile* 22(2):219–234
- Mazzoni MM, Licita D (2000) Significado estratigráfico y volcanológico de ignimbritas neógenas con composición intermedia en la zona del lago Caviahue, Neuquén. *Rev Asoc Geol Arg* 55(3):188–200
- Melnick D, Folguera A, Ramos V (2006a) Structural control on arc volcanism: the Caviahue-Copahue complex, Central to Patagonian Andes transition (38°S). *J S Am Earth Sci* 22:66–88
- Melnick D, Rosenau M, Folguera A, Echter H (2006b) Neogene Tectonics of the Western flank of the Neuquén Andes, 37°–39°30'S. In: Kay S, Ramos VA (eds) *Late Cretaceous to Recent magmatism and tectonism of the Southern Andean margin at the latitude of the Neuquen basin (36–39°S)*. *Geol S Am S* 407:73–95
- Moreno H, Lahsen A (1986) El volcán Callaqui: ejemplo de vulcanismo fisural en los Andes del Sur. *Rev Asoc Geol Arg* 42:1–8
- Muñoz J, Stern C (1988) The Quaternary volcanic belt of the southern continental margin of South America: transverse structural and petrochemical variations across the segment between 38° and 39°S. *J S Am Earth Sci* 1(2):147–161
- Muñoz J, Stern C (1989) Alkaline magmatism within the segment 38°–39°S of the Plio-Quaternary volcanic belt of the Southern South American continental margin. *J Geophys Res* 94:4545–4560
- Muñoz Bravo J, Stern C, Bermúdez A, Delpino D, Dobbs MF, Frey FA (1989) El vulcanismo plio-cuaternalario a través de los 38° y 39°S de los Andes. *Rev Asoc Geol Arg* 44(1–4):270–286

- Naranjo JA, Polanco E (2004) The 2000 AD eruption of Copahue Volcano, Southern Andes. *Rev Geol Chile* 31:279–292
- Niemeyer H, Muñoz J (1982) Hoja Laguna de La Laja, Región del Bío-Bío, scale 1:250.000, Servicio Nacional de Geología y Minería
- Pesce A (1989) Evolución volcánico-tectónica del complejo efusivo Copahue-Caviahue y su modelo geotérmico preliminar. *Rev Asoc Geol Arg* 44:307–327
- Pesicek J, Engdahl E, Thurber C, DeShon H, Lange D (2012) Mantle subducting slab structure in the region of the 2010 M8.8 Maule earthquake (30–40°S), Chile. *Geophys J Int* 191:317–324
- Polanco E, Naranjo JA, Young S, Moreno H (2000) Volcanismo Explosivo Holoceno en la cuenca del alto Bío-Bío, Andes del Sur (37°45'–38°30'S). IX Congreso Geológico Chileno, Puerto Varas, Chile 2:59–61
- Radic J (2010) Las cuencas cenozoicas y su control en el volcanismo de los complejos Nevados de Chillán y Copahue-Callaqui (36°–39°S). *Andean Geol* 37 (1):220–246
- Rojas Vera E, Folguera A, Spagnuolo M, Ramos VA (2009a) La neotectónica del arco volcánico a la latitud del volcán Copahue (38°S). In: *Geología de los Andes del Sur*. *Rev Asoc Geol Arg* 65(1):204–214
- Rojas Vera E, Folguera A, Ramos VA (2009b) Estratigrafía del sector central de la cuenca de Loncopué: El depocentro cuaternario del Huecú (sector occidental de la cuenca neuquina). En: *Geología de la Cuenca Neuquina*. *Rev Asoc Geol Arg* 65(2):400–412
- Rojas Vera E, Folguera A, Giménez M, Martínez P, Ruiz F, Ramos VA (2009c) Evolución tectónica de la fosa de Loncopué: Estructura del depocentro cuaternario del Huecú y su relación con la sedimentación y el volcanismo. *Rev Asoc Geol Arg* 64(2):213–229
- Rojas Vera E, Folguera A, Zamora Valcarce G, Giménez M, Ruiz F, Martínez P, Bottesi G, Ramos VA (2010) Neogene to Quaternary extensional reactivation of a fold and thrust belt: the Agrio belt in the Southern Central Andes and its relation to the Loncopué trough (38°–39°S). *Tectonophysics* 92(1–4):279–294
- Rojas Vera E, Folguera A, Zamora Valcarce G, Bottesi G, Ramos VA (2013) Structure and development of the Andean system between 36–39°S. *J Geodyn* 73:34–52
- Singer B, Hildreth W, Vincze Y (2000) 40Ar/39Ar evidence for early deglaciation of central Chilean Andes. *Geophys Res Lett* 27:1663–1666. doi:10.1029/1999GL011065
- Stern C (1989) Pliocene to present migration of the volcanic front, Andean Southern Volcanic Front. *Rev Geol Chile* 16(2):145–162
- Suárez M, Emparán G (1997) Hoja Curacautín, Regiones de la Araucanía y del Bío-Bío, scale 1:250,000, Servicio Nacional de Geología y Minería
- Tašárová Z (2004) Gravity data analysis and interdisciplinary 3D modelling of a convergent plate margin (Chile, 36–42°S) (PhD thesis): Berlin, Germany, Freie Universität Berlin, pp 187
- Varekamp J, Ouimette A, Herman S, Bermúdez A, Delpino D (2001) Hydrothermal element fluxes from Copahue, Argentina: a beehive volcano in turmoil. *Geology* 29(11):1059–1062
- Vélez M, Euillades P, Caselli A, Blanco M, Martínez Díaz J (2011) Deformation of Copahue volcano: inversion of InSar data using a genetic algorithm. *J Volcanol Geoth Res* 202:117–126
- Vergara M, Muñoz J (1982) La Formación Cola de Zorro en la alta cordillera Andina Chilena (36°–39° Lat. S), sus características petrográficas y petrológicas: una revisión. *Rev Geol Chile* 17:31–46
- Yuan X, Asch G, Bataille K, Bock G, Bohm M, Ehtler H, Kind R, Oncken O, Wölbern I (2006) Deep seismic images of the Southern Andes. In: Kay S, Ramos VA (eds) *Late Cretaceous to Recent magmatism and tectonism of the Southern Andean margin at the latitude of the Neuquen basin (36–39°S)*. *Geol S Am S* 407:61–72
- Zech R, May J-H, Kull C, Ilgner J, Kubik PW, Veit H (2008) Timing of the late Quaternary glaciations in the Andes from ~15 to 40°S. *J Quaternary Sci* 23 (6):635–647. doi:10.1002/jqs.1200