

Chapter 1

Introduction of Ultra-Weak Chemiluminescence



Xiangnan Dou, Syed Niaz Ali Shah, and Jin-Ming Lin

Abstract This chapter gives an overview of the basic principle of chemiluminescence and ultra-weak chemiluminescence. The development of various ultra-weak CL systems will be introduced. The ultra-weak chemiluminescence based on reactive oxygen species will be focused in this chapter. Ultra-weak chemiluminescence hinders its application in analytical chemistry due to its low luminous efficiency. The roles of nanoparticles in enhancing these ultra-weak CL systems will also be reviewed.

Keywords Chemiluminescence · Ultra-weak chemiluminescence · Principle · Application

1.1 Introduction

Chemiluminescence (CL) is light emitted from a chemical reaction. The requirements for chemiluminescence are (1) The chemical reaction must provide enough excitation energy to enable the electron to transition from the ground state to the excited state, and (2) The energy from the chemical reaction can be accepted by at least one substance and generate an excited state; (3) the electronically excited products have chemiluminescence quantum yield to release photon or transfer its energy to other luminophores [1]. CL is typically observed during redox reaction processes. The chemiluminescence quantum yield, Φ_{CL} , is determined by the efficiency of generating the excited state product and the efficiency of an emitted photon of the excited state product. Φ_{CL} does not usually surpass 0.1–0.001% in most circumstances. Only a few CL systems, such as the unique substituted oxiamide or dioxetane intermediate, may achieve a yield of 30% [2]. High quantum yields ranging from 1 to 100% of luminescence emerges during oxidation of the luciferin, mediated by the enzyme luciferase in living cells. Luciferin and luciferase are substrates and enzymes that interact to produce visible light. It is the enzyme with a unique microenvironment and

X. Dou · S. N. A. Shah · J.-M. Lin (✉)
Department of Chemistry, Tsinghua University, Beijing 100084, China
e-mail: jmlin@mail.tsinghua.edu.cn

reactive properties that are frequently used as a reaction medium in bioluminescence. The quantum yields of the excited state species and emission in bioluminescent emission based on an enzymatic reaction are quite high. This means that some ultra-weak CL systems could be produced as CL reaction media, the quantum yield of CL may be improved.

The CL technique has the advantage of having mild response conditions and being simple to manage. CL provides several distinct advantages as an analytical method:

- (1) Unlike fluorescence and UV–visible analysis, CL does not require an external light source, making it simple to integrate with other equipment like chromatography and capillary electrophoresis.
- (2) The sensitivity of CL analysis is very high. CL excitation energy derives from chemical reactions (mostly redox reactions). As a result, CL analysis lowers the signal's effect on the source stability by eliminating Rayleigh and Raman scattering. A low level of noise and a high signal-to-noise ratio is achieved by the CL method, which has a detection limit as low as attomole (10^{-18}) or even zeptomole (10^{-21}).
- (3) CL analysis may be used to detect substances involved in CL processes, including CL substrates, catalysts, inhibitors, and sensitizers.

During the last 40 years, many analysts have become interested in CL analysis because of its great sensitivity and ease of instrumentation. In contrast to the fluorescence approach, the exciting light source for CL is not required. As a result, the CL has found widespread application in clinical chemistry, biochemistry, and environmental chemistry, etc. The most evident manifestation of the rapid development is the large number of articles devoted solely to CL [3–10]. Because of its extraordinarily high sensitivity and other advantages, analytical methods comprising CL have recently attracted a lot of interest in a variety of sectors. Luminol, lucigenin, and oxalate esters are common CL systems [10, 11]. The CL method's exceptional sensitivity, however, presented a selectivity problem. Although various new CL systems are being developed for the identification of analytes [12–15], most CL systems could not be employed to directly determine the analytes. These limited CL systems only meet the needs of researchers for practical application. It's worth noting that the scarcity of functional CL systems is a key roadblock to CL's mainstream adoption. The quest for a new CL system is critical for expanding the scope of CL methods. Taking note of the fact that many redox reactions generate relatively faint light, the efforts on finding new CL systems is the main focus. The introduction of sensitizers or catalysts has already been shown to dramatically improve the CL properties of such weak CL systems [16, 17].

Many analysts worked hard to increase the CL intensity as well as the selectivity. Many new CL systems, both homogeneous and heterogeneous, have been created because of their work.

1.2 Principle of Chemiluminescence

CL can be produced by a chemical reaction that releases between 41 and 72 kcal per mole energy on its way to the product [18, 19]. There are three primary phases in the CL response. The first is the production of an intermediate that can react to provide enough energy to emit a visible light photon. The development of an electrically excited state is the next step. The CL reactions are distinguished from the rest of the chemical reactions by this step. The excited state's emission of light is the final stage [19]. The molecule's vibrational and rotational shifts are frequently accompanied by the electronic transition. Transitions from π bonding to π^* anti-bonding orbital ($\pi - \pi^*$) or from a non-bonding to an anti-bonding orbital ($n - \pi^*$) are the most common in organic compounds. CL denotes the return of an electron to its ground state accompanied by the emission of a photon. Excited molecules can also lose energy through a variety of mechanisms including chemical processes, collisional deactivation, internal conversion, and inter-system crossing. As a result, when these radiationless processes compete with CL, they are considered undesirable from an analytical perspective. The chemical interaction between the molecules A and B, as well as the energy transfer during the reaction, is depicted in Fig. 1.1.

In the first step, the reaction of A and B forms an excited state compound (C^*) which has a quantum yield Φ_{EX} .



We can see from Fig. 1.1 that the excited molecule (C^*) can emit CL in two different ways.

- (1) In the first case, C^* is a luminescence emitter that may be viewed through the luminescence emission.






Fig. 1.1 The basic principle of chemiluminescence

- (2) When the excited molecule C^* transferred its energy to the fluorescent molecule (F), the emission was increased by a more sensitive fluorescence. As a result, F is referred to as a CL enhancer, and the process is referred to as an enhancement CL reaction.



As soon as the CL reaction was initiated in the first instance. The quantum yield (CL) is the fraction of molecules that produce a photon when they return to their ground state. It is calculated as the product of two ratios:

$$\Phi_{CL} = \Phi_{EX} \cdot \Phi_{EM} \quad (1.5)$$

Traditional CL systems often have very poor quantum yields, either because of heat release or because of quenching from the surrounding environment. To avoid this difficulty, it is recommended that a fluorescent chemical with a reasonably high quantum yield be added to the reaction mixture to facilitate the transfer of energy to the fluorescent species [18]. As its name implies, this type of CL system emits its ultimate emission from an excited state created by direct oxidation of the excitation step. It is referred to as the direct CL system. The indirect CL system, on the other hand, is a CL system in which the final emission occurs indirectly because of energy transfer from the first created excited state, as opposed to the direct CL system [19]. Recently, nanomaterials (NMs) have been widely used to increase the CL intensity of classical systems, which is a significant advancement.

Inorganic reactions in CL systems produce weak CL emissions because of the low quantum yield of the inorganic reactions. For analytical applications, then, it is important to increase the CL intensity level. Compounds such as NMs, transition elements, and/or reaction techniques such as catalytic and energy transfer processes have been proposed to improve the CL signals of these compounds [20]. Currently, there is a considerable deal of interest in CL signal enhancement [21]. The research orientation in CL is partially devoted to the development of novel CL systems and partly to the application of these systems to real samples, with the former being the primary focus. This type of research can be primarily focused on discovering new CL reagents, such as substrates, sensitizers, and catalysts [22].

In general, the CL reaction is a redox reaction [23], which involves the participation of a variety of reactive oxygen species (ROS) [24, 25]. Hydroxyl radical ($\cdot\text{OH}$), superoxide radical ($\cdot\text{O}_2^-$), and substrate radicals, as well as singlet oxygen ($^1\text{O}_2$), are all examples of ROS [24–26]. Increased ROS generation is a frequently used technique for increasing the CL intensity of a system [27]. The peroxide-induced CL systems have garnered considerable attention and are widely used in a variety of fields. Hydrogen peroxide (H_2O_2), peroxydicarbonate, peroxydisulfate, and peroxyxynitrite are all examples of peroxides [28]. H_2O_2 has a critical role

in the CL responses [23]. The breakdown of H_2O_2 in a CL reaction generates reactive oxygen species (ROS) [13, 29]. Many CL emissions from a system are possible because of the system's multiple CL paths. Multiple CL pathways can occur concurrently in the same system. The quenching of CL signals occurs when a CL pathway is blocked. CL emission may have increased because of the closure of a less efficient CL pathway [24]. To detect and track these radicals in a CL system, electron paramagnetic resonance (EPR) and free radical scavengers are utilized.


1.3 Development of Ultra-Weak CL Systems

CL provides several advantages for analytical chemistry applications, including high sensitivity, a broad linear range, simple and inexpensive apparatus, and minimal background noise. However, many CL systems use certain conventional reagents, including luminol, lucigenin, acridinium, AMPPD, and $Ru(bpy)_3^{2+}$, the majority of which are organic molecules. We have attempted to build various new CL systems based on inorganic reactions throughout the last two decades, notwithstanding the difficulty of directly detecting light emissions using CL approaches due to the poor quantum yield. The book focuses on the construction of ultra-weak CL systems using peroxydicarbonate, peroxydicarbonate/peroxydicarbonate, peroxydicarbonate, periodate, or hydrogen peroxide as starting materials. Additionally, it will address ultra-weak CL that has been boosted or amplified by nanoparticles, quantum dots, or layered double hydroxide (LDH), as well as aggregation-induced emission (AIE).




1.3.1 Hydrogen Peroxide-Based Chemiluminescence

1.3.1.1 Fenton/Fenton like System

The Fenton reaction, which produces hydroxyl radicals, is commonly utilized in the practice of scientific research as a source of radicals. It is the Fenton reaction that produces the light radiation that may be measured by a contemporary luminometer. The metal (M)-catalyzed breakdown of H_2O_2 results in the formation of the $\bullet OH$ radical [30].





Initially, the Fenton reaction resulted in the creation of reactive oxygen species (ROS), which then resulted in the emission of CL. When the Fenton reagent is introduced, a flash of light can be seen. The singlet oxygen (1O_2) is the emissive species that emits luminescence. The following is an example of the ROS production response resulting from the Fenton reaction [31]. Lin et al. used CDs for the enhancement of the Fenton system [32, 33].



1.3.1.2 Peroxynitrite (ONOO^-) CL System

Peroxynitrite (ONOO^-) is a substantial nitric oxide derivative that has a wide range of uses in several fields. Even though peroxynitrite has a half-life of 10–20 ms, it is sufficiently stable to permeate biological membranes, diffuse one to two cell diameters, and interact with many essential biomolecules [34].

The coupling of $\text{NO}\bullet$ and O_2^- radicals in biological systems is most likely the source of ONOO^- (Eq. 1.10) [35]. Furthermore, ONOO^- can be produced by mixing acidified H_2O_2 with nitrite in real-time. Nitrite reacts with hydrogen peroxide in an acidic medium to form ONOOH (Eq. 1.11), which is unstable and can be decomposed into ONOO^- . Weak chemiluminescence was observed during the decomposition of ONOO^- .



Several studies described how fluorescent compounds and NMs such as CdTe QDs, Mg–Al carbonate LDHs, CDs, and Au NPs improve the CL from ONOO^- [36–38]. Carbonate has been shown to react with ONOO^- to form the adduct ONOOCO_2^- , which decomposes fast into $\text{NO}_2\bullet$ and the carbonate radical ($\text{CO}_3^{\bullet-}$), according to the literature. It is the conversion of the $\text{CO}_3^{\bullet-}$ radical into CO_2 in its excited form that causes the CL emission. CDs were shown to significantly increase the CL from the peroxynitrous acid (ONOOH) and CO_3^{2-} CL system described by Lin et al. [39] (Fig. 1.2).

1.3.1.3 Peroxysulfite (HSO_4^-) CL System

Early in the twentieth century, the H_2O_2 – HSO_3^- CL phenomena were identified by Stauff [40]. A peroxysulfite compound (HSO_4^- or HOOSO_2^-) is generated during the interaction of HSO_3^- with hydrogen peroxide. This compound is unstable and can be converted into the radicals $\text{SO}_3^{\bullet-}$ and $\text{OH}\bullet$ in the presence of H_2O_2 . There








Fig. 1.2 Schematic of the CL process in the carbon nanodots–NaNO₂–H₂O₂–Na₂CO₃ system using a schematic diagram [39]

are many CL emitting molecules, such as SO₂* and ¹O₂ that have been produced through radical reactions [41, 42].



Xue et al. investigated the influence of CDs on the $\text{H}_2\text{O}_2\text{-HSO}_3^-$ CL system and discovered that they had an enhanced effect [43]. In the $\text{H}_2\text{O}_2\text{-NaHSO}_3$ system, the CDs aided in the production of the emitter species SO_2^* and $^1\text{O}_2$ by facilitating the creation of these species (Fig. 1.3a). The CL of the $\text{H}_2\text{O}_2\text{-NaHSO}_3$ system was also improved using plasmonic luminous core-shell nanocomposites. According to [44], the combination of chemically produced excited states with surface plasmons results in an amplifying effect (Fig. 1.3b).

As a result of its high oxidation potential and proclivity to react through oxygen transfer, peroxydisulfate (HSO_4^-) is one of the most promising oxidants for organic molecules in use today. Comparative to other common reagents such as peroxydisulfate and peroxymonophosphoric acid, it has also been shown to be a powerful oxidizing agent. Chaps. 5 and 6 will discuss the CL of HSO_4^- in further detail.







Fig. 1.3 a The mechanism of the CL reaction in the $\text{H}_2\text{O}_2\text{-NaHSO}_3$ CDs system [43]; b the enhancement of CL intensity from the $\text{H}_2\text{O}_2\text{-NaHSO}_3$ system by plasmonic luminous nanocomposites [44]

1.3.1.4 Peroxymonocarbonate CL System

Peroxymonocarbonate (HCO_4^-) has been reported as a product of the interaction of hydrogen peroxide with bicarbonate since the 1980s. The breakdown of HCO_4^- resulted in the formation of CL. Carbonate radical was created as a result of it. The CL emitters includes singlet oxygen ($^1\text{O}_2$), singlet oxygen molecular pair [$(\text{O}_2)_2^*$], and excited double ($(\text{CO}_2)_2^*$).



Many articles are focused on improving the peroxymonocarbonate system's ultra-weak CL. Cu/Ni metal NPs were used to increase the ultra-weak CL utilizing surface plasmons [45]. Protein-stabilized zinc/copper nanoclusters have been shown to greatly increase the CL of peroxymonocarbonate [46]. The catalysis of Zn/Cu@BSA nanoclusters and the metal surface plasmon-linked emission of excited $(\text{CO}_2)_2^*$ resulted in the enhanced CL.

1.3.1.5 Periodate CL System

The periodate produces soft oxidation products and has been used for the micro-analysis of different compounds [18, 47]. The periodate reacts with H_2O_2 in acidic, neutral, and basic mediums to produce $^1\text{O}_2$ as an emissive species [48]. Various researchers used different enhancers to amplify the CL intensity of this system [49]. Lin et al. [50] used sodium dodecylbenzene sulfonate (SDBS) modified layered double hydroxides to enhance the CL intensity of the $\text{IO}_4^- - \text{H}_2\text{O}_2$ system (Fig. 1.4a). Our group used nitrogen-doped CDs for the enhancement of this system [51, 52]. Shah et al. [53] reviewed the details on the improvement and application of the periodate-peroxide CL system recently.






Fig. 1.4 **a** The mechanism of CL enhancement by sodium dodecylbenzene sulfonate (SDBS)-modified LDHs [50]. **b** Detailed about the improvement and application of the periodate-peroxide CL system [53]

1.3.2 The Role of NPs in the CL System

Because of their possible application in CL, nanomaterials (NMs) with good optical and electrical characteristics have gotten a lot of attention. Many studies have attempted to employ nanoparticles (NPs) to boost CL in various ultra-weak CL systems, including $\text{H}_2\text{O}_2\text{-NaHCO}_3$, $\text{H}_2\text{O}_2\text{-NaIO}_4$, $\text{H}_2\text{O}_2\text{-NaNO}_2$, $\text{H}_2\text{O}_2\text{-NaClO}$, $\text{H}_2\text{O}_2\text{-NaHSO}_3$, and others [52, 54, 55]. NPs can directly participate in redox reactions or operate as catalysts or energy acceptors in CL processes, to boost CL intensity. The role of NPs in the CL system is summarized below.





1.3.2.1 Direct CL

Excitons being a bound state of the introduced electrons and holes in the conduction and valence bands of the nanomaterials, respectively, is usually generated during a CL reaction involving nanomaterials. The relaxation of excitons resulted in the NPs-based CL. The CL system of CdTe QDs–calcein– $\text{K}_3\text{Fe}(\text{CN})_6$ was developed by Zhang et al. [56], in which CdTe QDs directly participating in the reaction by oxidized to excited CdTe QDs by $\text{K}_3\text{Fe}(\text{CN})_6$. The excited CdTe QDs are capable of transferring their energy to the calcein acceptor through a process known as energy transfer.



Lin et al. discovered carbon dots could react with acidic KMnO_4 and cerium (IV) to emit CL [57]. Such oxidants could cause holes to form in the CDs. The injection of holes into carbon dots by an oxidant increases the number of holes in the CDs and speeds up electron–hole annihilation, resulting in energy release in the form of CL emission.

Li et al. [58] discovered that CdTe/CdS/ZnS QDs exhibit novel CL performance with KIO_4 . The direct oxidation of CdTe/CdS/ZnS QDs by dissolved O_2 produced a superoxide radical ($\bullet\text{O}_2^-$), $\text{QD}(e_{1se}^-)$ and $\text{QD}(h_{1sh}^+)$ excitons were generated by injecting electrons and holes from $\bullet\text{O}_2^-$ and $\bullet\text{OH}$ radicals, respectively, resulting in QDs^* in the excited state.



1.3.2.2 CL Catalyst

Many research groups have investigated the catalytic effect of NMs. The NMs-based catalytic CL system was developed as an analytical chemical sensor. The catalytic effect or enhancement of CL processes by NMs is attributed to the formation of reactive oxygen radicals such as $\bullet\text{OH}$ radical, and $\bullet\text{O}_2^-$ radical, which are initiated by catalysts or enhancers. The NMs, particularly those containing transition metals, exhibit a significant catalytic outcome on H_2O_2 -based CL systems, such as luminol– H_2O_2 and lucigenin– H_2O_2 .

Furthermore, CL responses varied depending on the morphology and composition of catalytic NMs. Take, for example, the work of Li et al. [59], who investigated the CL response of triangular Au NPs in the luminol– H_2O_2 system and contrasted it with the reaction of spherical Au NPs. According to the researchers, when compared to spherical Au NPs, triangular Au NPs exhibited much higher catalytic activity due to a higher surface-to-volume ratio and a greater number of active surface sites. These

triangular Au NPs exhibited improved activity in the formation of reactive oxygen radicals and electron transferability on their surfaces. It has also been demonstrated that the CL system of luminol–H₂O₂ catalyzed by triangular Au NPs has excellent analytical application for the detection of aminothiols.

Because of the redox feature of the distinct electron and hole states of QDs, they can act as a catalyst in CL systems. With QDs, the injection of holes from a strong oxidizer or electrons from a reducer can alter the valence and conduction bands of the QDs. During the injection procedure, ROS such as $\cdot\text{OH}$, $\cdot\text{O}_2^-$ radicals will be created. By radical reactions, these radical species can oxidize CL reagents, resulting in increased CL, or generating CL emitters. L-cysteine capped Mn-doped ZnS QDs have previously been shown to improve CL of the H₂O₂-NaClO system [60]. The increased CL of NaClO–H₂O₂ is due to the sensitizer's catalysis in creating reactive intermediates as well as the electron-transfer mechanism [53].

The CL's efficiencies were affected by the types of reactants and catalysts used. Chen et al. [61] demonstrated improved CL of the H₂O₂–NaHCO₃ system with the addition of CdTe QDs. The excited CdTe QDs were produced by combining the hole and electron injected QDs, which cause the enhancement of CL. They investigated the size effect of various catalyst types on the CL of the H₂O₂–NaHCO₃ system. CdTe QDs were compared to CdSe and CdS QDs with emission wavelengths of 549 and 607 nm, respectively. They discovered that CdTe QDs with a larger size than CdSe and CdS quantum dots at the same concentration could significantly increase CL intensity.

1.4 Conclusion

The purpose of this chapter is to introduce the basic principles of chemiluminescence and the development of ultra-weak CL systems, focusing on the CL from the decomposition of peroxynitrite (ONOO⁻), peroxymonocarbonate (HCO₄⁻), peroxysulfite (HSO₄⁻), and periodate (IO₄⁻), as well as the application of NPs to develop new reactive oxygen (ROS) CL systems and enhance the weak CL systems. More information on the CL of these inorganic compounds can be found in Chaps. 2, 3, 4, 5, 6, 7, 8, 9, 10 and 11 of this book.

References

1. Musfeldt JL (1998) Luminescence of solids edited by D. R. Vij. Plenum Press: New York. J Am Chem Soc 121:4310–4310. <https://doi.org/10.1021/ja985728i>
2. Cadenas E (1982) Chemical and biological generation of excited states. Herausgegeben von W. Adam und G. Cilento. Academic Press, New York. Angew Chem 96:452. <https://doi.org/10.1002/ange.19840960631>
3. Cormier M (2013) Chemiluminescence and bioluminescence. Springer Science and Business Media

4. Andreotti PE (1988) Chemiluminescence: principles and applications in biology and medicine. In: Campbell AK (ed) Ellis Horwood, Chichester, UK. <https://doi.org/10.1002/jps.2600780920>
5. Blum LJ (1997) Bio- and chemi-luminescent sensors
6. Chen H, Gao Q, Li J, Lin J-M (2016) Graphene materials-based chemiluminescence for sensing. *J Photochem Photobiol C: Photochem Rev* 27:54–71. <https://doi.org/10.1016/j.jphotochemrev.2016.04.003>
7. Shah SNA, Lin J-M (2017) Recent advances in chemiluminescence based on carbonaceous dots. *Adv Colloid Interface Sci* 241:24–36. <https://doi.org/10.1016/j.cis.2017.01.003>
8. Yang M, Huang J, Fan J, Du J, Pu K, Peng X (2020) Chemiluminescence for bioimaging and therapeutics: recent advances and challenges. *Chem Soc Rev* 49:6800–6815. <https://doi.org/10.1039/DOCS00348D>
9. Al Mughairy B, Al-Lawati HAJ (2020) Recent analytical advancements in microfluidics using chemiluminescence detection systems for food analysis. *TrAC, Trends Anal Chem* 124:115802. <https://doi.org/10.1016/j.trac.2019.115802>
10. Birks JW (1989) Chemiluminescence and photochemical reaction detection in chromatography. VCH Publishers, New York
11. Rongen HAH, Hoetelmans RMW, Bult A, Van Bennekom WP (1994) Chemiluminescence and immunoassays. *J Pharm Biomed Anal* 12:433–462. [https://doi.org/10.1016/0731-7085\(94\)80027-8](https://doi.org/10.1016/0731-7085(94)80027-8)
12. Yu W, Zhao L (2021) Chemiluminescence detection of reactive oxygen species generation and potential environmental applications. *TrAC, Trends Anal Chem* 136:116197. <https://doi.org/10.1016/j.trac.2021.116197>
13. Dou X, Zhang Q, Shah SNA, Khan M, Uchiyama K, Lin J-M (2019) MoS₂-quantum dot triggered reactive oxygen species generation and depletion: responsible for enhanced chemiluminescence. *Chem Sci* 10:497–500. <https://doi.org/10.1039/C8SC03511C>
14. Aslan K, Geddes CD (2009) Metal-enhanced chemiluminescence: advanced chemiluminescence concepts for the 21st century. *Chem Soc Rev* 38:2556–2564. <https://doi.org/10.1039/b807498b>
15. Chen J, Qiu H, Zhao S (2020) Fabrication of chemiluminescence resonance energy transfer platform based on nanomaterial and its application in optical sensing, biological imaging and photodynamic therapy. *TrAC, Trends Anal Chem* 122:115747. <https://doi.org/10.1016/j.trac.2019.115747>
16. Ishimaru N, Lin J-M, Yamada M (1998) Luminol-free chlorine chemiluminescence in an oil-in-water microemulsion medium. *Anal Commun* 35:67–69. <https://doi.org/10.1039/A708423D>
17. Tsukada S, Miki H, Lin J-M, Suzuki T, Yamada M (1998) Chemiluminescence from fluorescent organic compounds induced by cobalt(II) catalyzed decomposition of peroxomonosulfate. *Anal Chim Acta* 371:163–170. [https://doi.org/10.1016/S0003-2670\(98\)00356-0](https://doi.org/10.1016/S0003-2670(98)00356-0)
18. Evmiridis NP, Vlessidis AG, Thanasoulas NC (2007) Chemical analysis through CL-Detection Assisted By Periodate Oxidation. *Bioinorg Chem Appl* 2007:092595. <https://doi.org/10.1155/2007/92595>
19. Schuster GB (1979) Chemiluminescence of organic peroxides. Conversion of ground-state reactants to excited-state products by the chemically initiated electron-exchange luminescence mechanism. *Acc Chem Res* 12:366–373. <https://doi.org/10.1021/ar50142a003>
20. Chen H, Lin L, Li H, Lin J-M (2014) Quantum dots-enhanced chemiluminescence: mechanism and application. *Coord Chem Rev* 263:86–100. <https://doi.org/10.1016/j.ccr.2013.07.013>
21. Gnaïm S, Scomparin A, Eldar-Boock A, Bauer CR, Satchi-Fainaro R, Shabat D (2019) Light emission enhancement by supramolecular complexation of chemiluminescence probes designed for bioimaging. *Chem Sci* 10:2945–2955. <https://doi.org/10.1039/c8sc05174g>
22. Song H, Su Y, Zhang L, Lv Y (2019) Quantum dots-based chemiluminescence probes: an overview. *Luminescence* 34:530–543. <https://doi.org/10.1002/bio.3633>
23. Dou X, Zheng Y, Uchiyama K, Lin J-M (2016) Fluorescent carbon nanoparticles: Mimicking hydrogen peroxide properties in a chemiluminescence system. *Chem Commun* 52:14137–14140. <https://doi.org/10.1039/c6cc07285b>

24. Ma AJ, Chang YT, Lin WY (2013) A kinetic treatment of stopped-flow time courses for multiple chemiluminescence of the KIO_4 -luminol- Mn^{2+} system. *Luminescence* 28:355–362. <https://doi.org/10.1002/bio.2389>
25. Huang TY, Lin WY (2011) A stopped-flow study of the dual chemiluminescence for the luminol- KIO_4 - Mn^{2+} system in strong alkaline solutions. *Luminescence* 26:118–124. <https://doi.org/10.1002/bio.1193>
26. Chen X, Tian X, Shin I, Yoon J (2011) Fluorescent and luminescent probes for detection of reactive oxygen and nitrogen species. *Chem Soc Rev* 40:4783–4804. <https://doi.org/10.1039/c1cs15037e>
27. Chen YC, Lin WY (2010) Enhancement of chemiluminescence of the KIO_4 -luminol system by gallic acid, acetaldehyde and Mn^{2+} : application for the determination of catecholamines. *Luminescence* 25:43–49. <https://doi.org/10.1002/bio.1141>
28. Lin Z, Chen H, Lin J-M (2013) Peroxide induced ultra-weak chemiluminescence and its application in analytical chemistry. *Analyst* 138:5182–5193. <https://doi.org/10.1039/c3an00910f>
29. Zhang M, Yao Q, Guan W, Lu C, Lin J-M (2014) Layered double hydroxide-supported carbon dots as an efficient heterogeneous Fenton-like catalyst for generation of hydroxyl radicals. *J Phys Chem C* 118:10441–10447. <https://doi.org/10.1021/jp5012268>
30. Lu C, Song G, Lin J-M (2006) Reactive oxygen species and their chemiluminescence-detection methods. *TrAC, Trends Anal Chem* 25:985–995. <https://doi.org/10.1016/j.trac.2006.07.007>
31. Aristova N, Ivanova I, Trofimova S, Knyazev D, Piskarev I (2011) Influence of luminol on the chemiluminescence intensity in Fenton's reaction. *High Energy Chem* 45:505–509. <https://doi.org/10.1134/S0018143911060038>
32. Shah SNA, Lin L, Zheng Y, Zhang D, Lin J-M (2017) Redox cycling of iron by carbon dot enhanced chemiluminescence: mechanism of electron-hole induction in carbon dot. *Phys Chem Chem Phys* 19:21604–21611. <https://doi.org/10.1039/C7CP03724D>
33. Shah SNA, Dou X, Khan M, Uchiyama K, Lin J-M (2019) N-doped carbon dots/ H_2O_2 chemiluminescence system for selective detection of Fe^{2+} ion in environmental samples. *Talanta* 196:370–375. <https://doi.org/10.1016/j.talanta.2018.12.091>
34. Pacher P, Beckman JS, Liaudet L (2007) Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 87:315–424. <https://doi.org/10.1152/physrev.00029.2006>
35. Radi R, Cosgrove TP, Beckman J, Freeman B (1993) Peroxynitrite-induced luminol chemiluminescence. *Biochem J* 290:51–57. <https://doi.org/10.1042/bj2900051>
36. Dong S, Guan W, Lu C (2013) Quantum dots in organo-modified layered double hydroxide framework-improved peroxynitrous acid chemiluminescence for nitrite sensing. *Sens Actuators, B Chem* 188:597–602. <https://doi.org/10.1016/j.snb.2013.07.060>
37. Adegoke O, Nyokong T (2013) Probing the sensitive and selective luminescent detection of peroxynitrite using thiol-capped CdTe and CdTe@ ZnS quantum dots. *J Lumin* 134:448–455. <https://doi.org/10.1016/j.jlumin.2012.08.002>
38. Zhang H, Zhang L, Lu C, Zhao L, Zheng Z (2012) CdTe nanocrystals-enhanced chemiluminescence from peroxynitrous acid-carbonate and its application to the direct determination of nitrite. *Spectrochim Acta Part A Mol Biomol Spectrosc* 85:217–222. <https://doi.org/10.1016/j.saa.2011.09.063>
39. Lin Z, Dou X, Li H, Ma Y, Lin J-M (2015) Nitrite sensing based on the carbon dots-enhanced chemiluminescence from peroxynitrous acid and carbonate. *Talanta* 132:457–462. <https://doi.org/10.1016/j.talanta.2014.09.046>
40. Stauff J (1967) Jaeschke W (1975) A chemiluminescence technique for measuring atmospheric trace concentrations of sulfur dioxide. *Atmos Environ* 9:1038–1039. [https://doi.org/10.1016/0004-6981\(75\)90028-1](https://doi.org/10.1016/0004-6981(75)90028-1)
41. Zheng Y, Dou X, Li H, Lin J-M (2016) Bisulfite induced chemiluminescence of gC3N4 nanosheets and enhanced by metal ions. *Nanoscale* 8:4933–4937. <https://doi.org/10.1039/C5NR08943C>
42. Shah SNA, Zheng Y, Li H, Lin J-M (2016) Chemiluminescence character of ZnS quantum dots with bisulphite-hydrogen peroxide system in acidic medium. *J Phys Chem C* 120:9308–9316. <https://doi.org/10.1021/acs.jpcc.6b01925>

43. Xue W, Lin Z, Chen H, Lu C, Lin J-M (2011) Enhancement of ultraweak chemiluminescence from reaction of hydrogen peroxide and bisulfite by water-soluble carbon nanodots. *J Phys Chem C* 115:21707–21714. <https://doi.org/10.1021/jp207554t>
44. Chen H, Xue W, Lu C, Li H-F, Zheng Y, Lin J-M (2013) Plasmonic luminescent core-shell nanocomposites-enhanced chemiluminescence arising from the decomposition of peroxomonosulfite. *Spectrochim Acta Part A Mol Biomol Spectrosc* 116:355–360. <https://doi.org/10.1016/j.saa.2013.07.057>
45. Chen H, Li R, Li H, Lin J-M (2012) Plasmon-assisted enhancement of the ultraweak chemiluminescence using Cu/Ni metal nanoparticles. *J Phys Chem C* 116:14796–14803. <https://doi.org/10.1021/jp303092d>
46. Chen H, Lin L, Li H, Li J, Lin J-M (2015) Aggregation-induced structure transition of protein-stabilized zinc/copper nanoclusters for amplified chemiluminescence. *ACS Nano* 9:2173–2183. <https://doi.org/10.1021/acs.nano.5b00141>
47. Vlessidis AG, Evmiridis NP (2009) Periodate oxidation and its contribution to instrumental methods of micro-analysis—a review. *Anal Chim Acta* 652:85–127. <https://doi.org/10.1016/j.aca.2009.06.065>
48. Li M, Lee SH, Bae ZU, Lee KP, Park YC, Lee MS (2004) Optical sensor for the determination of glucose based on KIO₄ chemiluminescence detection. *J Fluoresc* 14:597–601. <https://doi.org/10.1023/B:JOFL.0000039346.35328.d3>
49. Shah SNA, Shah AH, Dou X, Khan M, Lin L, Lin J-M (2019) Radical-triggered chemiluminescence of phenanthroline derivatives: an insight into radical-aromatic interaction. *ACS Omega* 4:15004–15011. <https://doi.org/10.1021/acs.omega.9b01785>
50. Zhang M, Han D, Lu C, Lin J-M (2012) Organo-modified layered double hydroxides switch-on chemiluminescence. *J Phys Chem C* 116:6371–6375. <https://doi.org/10.1021/jp300879c>
51. Shah SNA, Li H, Lin J-M (2016) Enhancement of periodate-hydrogen peroxide chemiluminescence by nitrogen doped carbon dots and its application for the determination of pyrogallol and gallic acid. *Talanta* 153:23–30. <https://doi.org/10.1016/j.talanta.2016.02.056>
52. Zheng Y, Zhang D, Shah SNA, Li H, Lin J-M (2017) Ultra-weak chemiluminescence enhanced by facilely synthesized nitrogen-rich quantum dots through chemiluminescence resonance energy transfer and electron hole injection. *Chem Commun* 53:5657–5660. <https://doi.org/10.1039/C7CC02041D>
53. Shah SNA, Khan M, Rehman ZU (2020) A prolegomena of periodate and peroxide chemiluminescence. *TrAC, Trends Anal Chem* 122:115722. <https://doi.org/10.1016/j.trac.2019.115722>
54. Li R, Chen H, Li Y, Lu C, Lin J-M (2012) Enhancing effect of alcoholic solvent on hydrosulfite-hydrogen peroxide chemiluminescence system. *J Phys Chem A* 116:2192–2197. <https://doi.org/10.1021/jp300012t>
55. Wu J, Wang X, Lin Y, Zheng Y, Lin J-M (2016) Peroxynitrous-acid-induced chemiluminescence detection of nitrite based on microfluidic chip. *Talanta* 154:73–79. <https://doi.org/10.1016/j.talanta.2016.03.062>
56. Kang J, Li X, Geng J, Han L, Tang J, Jin Y, Zhang Y (2012) Determination of hyperin in seed of *Cuscuta chinensis* Lam. by enhanced chemiluminescence of CdTe quantum dots on calcein/K₃Fe (CN)₆ system. *Food Chem* 134:2383–2388. <https://doi.org/10.1016/j.foodchem.2012.04.055>
57. Lin Z, Xue W, Chen H, Lin J-M (2012) Classical oxidant induced chemiluminescence of fluorescent carbon dots. *Chem Commun* 48:1051–1053. <https://doi.org/10.1039/c1cc15290d>
58. Li Y, Zheng Y-Z, Zhang D-K, Li H-F, Ma Y, Lin J-M (2017) Enhanced chemiluminescence from reactions between CdTe/CdS/ZnS quantum dots and periodate. *Chin Chem Lett* 28:184–188. <https://doi.org/10.1016/j.ccl.2016.07.020>
59. Li Q, Liu F, Lu C, Lin J-M (2011) Aminothiols sensing based on fluorosurfactant-mediated triangular gold nanoparticle-catalyzed luminol chemiluminescence. *J Phys Chem C* 115:10964–10970. <https://doi.org/10.1021/jp200711a>
60. Zhou Y, Chen H, Ogawa N, Lin J-M (2011) Chemiluminescence from NaClO–H₂O₂ and enhanced by L-cysteine capped Mn-doped ZnS quantum-dots. *J Lumin* 131:1991–1997. <https://doi.org/10.1016/j.jlumin.2011.04.019>

61. Chen H, Lin L, Lin Z, Guo G, Lin J-M (2010) Chemiluminescence arising from the decomposition of peroxydicarbonate and enhanced by CdTe quantum dots. *J Phys Chem A* 114:10049–10058. <https://doi.org/10.1021/jp104060x>