

---

# Castanea sativa

---

## Scientific Name

*Castanea sativa* Mill.

---

## Synonyms

*Castanea castanea* (L.) H. Karst. nom inval., *Castanea prolifera* (K. Koch) Hickel, *Castanea sativa* f. *discolor* Vuk., *Castanea sativa* var. *hamulata* A. Camus, *Castanea sativa* var. *microcarpa* Lavalille, *Castanea sativa* var. *prolifera* K. Koch, *Castanea sativa* var. *spicata* Husn., *Castanea sativa* var. *typica* Seemen, nom. inval., *Castanea vesca* Gaertn., *Castanea vulgaris* Lam., *Fagus castanea* L. (basonym), *Fagus castanea* var. *variegata* Weston, *Fagus procera* Salisb.

---

## Family

Fagaceae

---

## Common/English Names

Chestnut, Chestnut Tree, Edible Chestnut, European Chestnut, Italian Chestnut, Marron, Portuguese Chestnut, Spanish Chestnut, Sweet Chestnut.

---

## Vernacular Names

**Afrikaans:** Kastaiing;

**Brazil:** Castanha Européia, Castanha Portuguesa (Portuguese);

**Bulgarian:** Kecteh;

**Chinese:** Ou Zhou Li, Wu Shu Li;

**Croatian:** Kestenjaste Boje;

**Czech:** Kaštanovník Jedlý, Kaštanovník Setý;

**Danish:** Ægte Kastanie, Ægte Kastanje;

**Dutch:** EuropeseKastanje, Kastanje, Kastanjeboom, Tamme Kastanje, Tamme Kastanjeboom, Tame Kastanjeboom Soort;

**Eastonian:** Harilik Kastanipuu;

**Finnish:** Aito Kastanja, Jalokastanja;

**French:** Châtaigne, Châtaigner Commun, Châtaignier, Châtaignier Commun Marron, Marron Comestible;

**German:** Cheschtene, Ceste, Echte Kastanie, Edelkastanie, Edel-Kastanie, Edelkastanienbaum, Eßkastanie, Eßkastanienbaum, Ess-Kastanie, Essbare Kastanie, Keschte, Marone, Maroni;

**Greek:** Kastania;

**Hungarian:** Édes Gesztenye, Szelídsztenye;

**Icelandic:** Kastaníuhmeta;

**Indonesian:** Berangan;

**Italian:** Castagno, Castagno Comune, Castagno Domestico, Marone;

**Japanese:** Yooroppa Guri;

**Latvian:** Kastanis;

**Lithuanian:** Kaštonas;

**Malaysia:** Buah Berangan;

**Norwegian:** Edelkastanje, Ekte Kastanje, Kastanje;

**Polish:** Kasztan Jadalny;

**Portuguese:** Castanheiro-Comum, Castanheiro, Reboleiro;

**Romanian:** Castan, Castană;

**Russian:** Kashtan Nastoishchii (Kaštan Nastojaščij), Kashtan Posevnoi (Kaštan Posevnoj);

**Serbian:** Kesten;

**Slovačina:** Evropski, Pravi, Pravi Kostanj, Žlahtni Kostanj;

**Slovincina:** Gaštan Jedlý;

**Spanish:** Castaña, Castaño, Castaño Común, Castaño Regoldo, Regoldo;

**Swedish:** Äkta Kastanj, Kastanje;

**Turkish:** Kestane Ağacı;

**Vietnamese:** Cây Hạt Dẻ.

tolerant but adequate moisture is required for good growth and a good nut harvest.

---

## Edible Plant Parts and Uses

Chestnuts can be eaten raw or dried but are usually eaten roasted, fried or cooked (boiled or steamed). Roasting, frying or cooking brings out the delicious, sweet chestnut flavour and floury texture. The dried or cooked nuts are used in confectionery, pastries, chocolates, puddings, desserts and cakes. They are used for flour, bread making, as a cereal substitute, coffee substitute, a thickener in soups and other cookery uses, as well as for fattening pig stock. A sugar can be extracted from it. The Italian and Corsican polenta (type of porridge) is made with sweet chestnut flour. A local variety of Corsican beer also uses chestnuts.

---

## Origin/Distribution

A species of chestnut originally native to the Mediterranean in south-eastern Europe to Caucasus in Asia Minor. Wild or naturalized populations occur throughout southern Europe, northern Africa and southwestern Asia. It is cultivated in mild temperate regions in Europe and in the southern hemisphere and some subtropical regions.

---

## Agroecology

Chestnut requires a mild cool climate Mediterranean or sub-temperate climate with good annual rainfall of 750–1,200 mm. The tree is frost sensitive. Sub-zero temperatures are injurious to the tree. The tree thrives in full sun on deep well-drained, fertile, sandy or loamy soils as it has a deep root system. It is highly tolerant of acidic soils, gravelly or stony soils but intolerant of calcareous soils, heavy clays and impermeable soils. The soil pH range is from 4 to 6.5 with an optimum from 5.5 to 6.5. It will tolerate partial shade under forest conditions. Adult trees are drought

---

## Botany

A medium to large, deciduous tree growing to 15–35 m high with a spreading crown and trunk diameter reaching 2 m and deeply fissured bark (Plate 1). Branchlets are tomentose. Leaves are elliptic to ovate-lanceolate, 14–28 cm × 5–9 cm, with 11–14 pairs of nerves more prominent on the under surface, serrated margin, acuminate tip, oblique base, coriaceous, glabrous, pale green and puberulous on the under surfaces (Plate 2). Inflorescences or catkins are unisexual (male) or androgynous with female flowers (cupules) at the base of an otherwise male inflorescence. Staminate catkins are pendulous, pubescent, creamy yellow, consists of male flowers in dense cymules. Male flowers are apetalous, with 4–6 (–9), scale-like, connate or distinct sepals, filiform filaments with dorsifixed or versatile anthers opening by longitudinal slits; and with or without a rudimentary pistil. Female inflorescences of 1–7 or more flowers subtended individually or collectively by a spiny protective cupule formed from numerous fused bracts, arranged individually or in small groups along an axis or at base of an androgynous inflorescence. Female flower



**Plate 1** Tree habit with spreading crown



**Plate 3** Chestnut catkins



**Plate 4** Spiny chestnut cupules



**Plate 2** Elliptic leaves with serrated margins



**Plate 5** Harvested chestnuts

with 1–7 perianth, 1 pistil, inferior ovary with 3–6(–9) locules, style and carpels as many as locules; placentation axile; ovules anatropous, 2 per locule. Fruit is 1.3–2.5 cm in diameter, with each spiny cupule or burr (Plate 3) consisting of 1–4

nuts. Seed (nut) is brown to reddish-brown, usually solitary by abortion and is non-endospermic with a large embryo (Plate 5).

## Nutritive/Medicinal Properties

Food value of raw peeled, European chestnuts per 100 g edible portion was reported as follows (USDA 2011): water 52.00 g, energy 196 kcal (820 kJ), protein 1.63 g, total lipid (fat) 1.25 g, ash 0.96 g, carbohydrate 44.17 g; minerals – calcium 19 mg, iron, 0.94 mg, magnesium 30 mg, phosphorus 38 mg, potassium 484 mg, sodium 2 mg, zinc 0.49 mg, copper 0.418 mg, manganese 0.336 mg, vitamins – vitamin C (total ascorbic acid) 40.2 mg, thiamine 0.144 mg, riboflavin 0.016 mg, niacin 1.102 mg, pantothenic acid 0.476 mg, vitamin b-6 0.352 mg, folate (total) 58 mcg, vitamin A 26 IU; lipids – fatty acids (total saturated) 0.235 g, 14:0 (myristic acid) 0.005 g, 16:0 (palmitic acid) 0.212 g, 18:0 (stearic acid) 0.012 g; fatty acids (total monounsaturated) 0.430 g, 16:1 undifferentiated (palmitoleic acid) 0.012 g, 18:1 undifferentiated (oleic acid) 0.413 g, 20:1 (gadoleic acid) 0.005 g; fatty acids (total polyunsaturated) 0.493 g, 18:2 undifferentiated (linoleic acid) 0.440 g, 18:3 undifferentiated (linolenic acid) 0.053 g; amino acids – tryptophan 0.018 g, threonine 0.058 g, isoleucine 0.064 g, leucine 0.096 g, lysine 0.096 g, methionine 0.038 g, cystine 0.052 g, phenylalanine 0.069 g, tyrosine 0.045 g, valine 0.091 g, arginine 0.116 g, histidine 0.045 g, alanine 0.109 g, aspartic acid 0.281 g, glutamic acid 0.210 g, glycine 0.084 g, proline 0.086 g and serine 0.081 g.

In an analysis conducted in Turkey (Yıldız et al. 2009), the following nutrient data was reported for wild chestnut (*C. sativa*): moisture 54.84%, crude lipid 2.24%, crude protein, 8.93%, crude fibre, 3.92%, crude energy 4,046 kcal/100 g, ash 2.078%, dry matter 45.16%, total carbohydrate 11.21%, and ether soluble extract 4.44%. Zinc 5099.4 mg/kg, manganese 3031.9 mg/kg, sodium 1058.6 mg/kg and calcium 308.5 mg/kg were established as major minerals of chestnut fruit. Aluminium 21.52 mg/kg, boron 15.87 mg/kg, arsenic 7.82 mg/kg, bismuth 1.54 mg/kg, copper 0.12 mg/kg, chromium 5.72 mg/kg, strontium 15.22 mg/kg and titanium 768.72 mg/kg, iron 38.15 mg/kg were found in minor amounts.

Chestnut with a low fat content, completely free of cholesterol, a low sodium and high potassium content, moderate but high quality protein content, and rich in energy, vitamin C and amino acids especially lysine, tryptophan and sulphur containing amino acids provide a balanced and quality food (Bounous et al. 2000). Chestnuts also contained folate and vitamin Bs and A and the macro minerals. Chestnut was found to be characterized by the presence of seven organic acids: oxalic, cis-aconitic, citric, ascorbic, malic, quinic and fumaric acids (Ribeiro et al. 2007). Roasting, boiling and frying procedures led to significant reduction of total organic acids contents. Moisture was the major component in four Portuguese chestnut cultivars followed by carbohydrates, protein and fat, resulting in an energetic value lower than 195 kcal/100 g of fresh fruit (Barreira et al. 2009). Fatty acids (FA) profiling revealed a clear prevalence of C18:1 and C18:2, two FA very well-known due to their beneficial effects on human health, e.g., in the prevention of cardiovascular diseases. Triacylglycerols (TAG) profiling revealed that OLL, PLL, OOL and POL were the major compounds (O=oleic acid, P=palmitic acid, L = linoleic acid).

Chestnuts are mostly consumed as processed forms, and the different types of processing clearly affect the nutrient and non-nutrient composition of the fruits. Chestnut fruit industrial processing could be divided into four major stages: harvest, post-harvest storage (during 3 months at  $\pm 0^{\circ}\text{C}$ ), industrial peeling (by flame or fire – “brûlage” – at temperatures of 800–1,000 $^{\circ}\text{C}$ ) and freezing (tunnel with a CO<sub>2</sub> flow at  $-65^{\circ}\text{C}$  during 15–20 minutes) (De Vasconcelos et al. 2010b). Starch and hemicelluloses were the predominant polysaccharides but free sugars, mainly sucrose (saccharose), were also present. Vitamin C and vitamin E (predominantly  $\delta$ -tocopherol) were present in the fruits with lower levels of carotenoids (lutein and lutein esters). The fibre and free sugars contents were found to increase during industrial processing, while the interaction of cultivar  $\times$  processing stage had the greatest influence on the starch content. Various phenolics, mainly gallic and ellagic

acids and lower levels of ellagitannins, were also detected. De Vasconcelos et al. (2010c) also found that potassium and phosphorous were predominant in the fruits of all cultivars for both harvest years. Calcium, magnesium, iron, zinc and manganese were also present. Fruits from both harvest years had a significant content of free sugars, with sucrose predominating, and these sugars were more affected by the processing stage. Significant levels of lutein, lutein esters,  $\gamma$ -tocopherol and vitamin C were also found in the chestnut fruits. The contents of vitamin C and carotenoids were adversely affected during industrial processing.

The cooking processes was found to significantly affect primary and secondary metabolite composition of chestnuts (Gonçalves et al. 2010). Roasted chestnuts had higher protein contents, insoluble and total dietary fibre and lower fat contents whilst boiled chestnuts had lower protein, but higher fat contents. Cooking increased citric acid contents, especially in roasted chestnuts. In contrast, raw chestnuts had higher malic acid contents than cooked chestnuts. Moreover, roasted chestnuts had significantly higher gallic acid and total phenolic contents, and boiled chestnuts had higher gallic and ellagic acids contents, when compared to raw chestnuts. The data confirmed cooked chestnuts to be a good source of organic acids and phenolics and with low fat contents, properties that associated with positive health benefits.

### **Other Phytochemicals**

All three Portuguese cultivars were found to have high moisture contents but were low in ash, crude fat, and crude protein contents, with high starch and low fiber contents (De Vasconcelos et al. 2007). The free amino acid contents, including various essential amino acids, varied depending on the cultivar. All three cultivars also had a significant content of polyphenolics with gallic acid; ellagic acid was predominant among hydrolyzable and condensed tannins. Chestnuts have become increasingly important with respect to

human health, for example, as an alternative gluten-free flour source.

The chestnut fruit processing had been reported to generate large amounts of residues as pericarp (outer shell; 8.9–13.5%) and integument (inner shell; 6.3–10.1%) and studies showed that these materials clearly had the potential as sources of valuable co-products (de Vasconcelos et al. 2010a). The analyses of the pericarp and integument of four Portuguese chestnut cultivars revealed significant contents of total phenolics, low molecular weight phenolics (gallic and ellagic acid), condensed tannins and ellagitannins including castalagin, vescalagin, acutissimin A and acutissimin B. The integument tissues had the highest levels of total phenolics and condensed tannins. The most efficient extraction solvent for the total phenolics, total condensed tannins and low molecular weight phenolics was 70:30 acetone:water at 20°C.

Some low molecular weight phenolic compounds and hydrolyzable tannins were found in hardwood extracts of *Castanea sativa* before and after toasting (Sanz et al. 2010). The low molecular weight phenolic compounds were lignin constituents as the acids gallic, protocatechuic, vanillic, syringic, ferulic, and ellagic, the aldehydes protocatechuic, vanillic, syringic, coniferylic, and sinapic, and the coumarin scopoletin. Vescalagin and castalagin were the main ellagitannins, and acutissimin was also identified in the wood. Some gallotannins were tentatively identified, including different isomers of di, tri, tetra, and pentagalloyl glucopyranose, and di and trigalloyl-hexahydroxydiphenoyl glucopyranose, comprising 20 different compounds, as well as some ellagic derivatives such as ellagic acid deoxyhexose, ellagic acid dimer dehydrated, and valoneic acid dilactone. These ellagic derivatives as well as some galloyl and hexahydroxydiphenoyl derivatives were tentatively identified for the first time in chestnut wood. Seasoned and toasted chestnut wood showed a very different balance between lignin derivatives and tannins because toasting resulted in the degradation of tannins and the formation of low molecular

weight phenolic compounds from lignin degradation. A chlorine-free environmentally-friendly process was developed for extraction of 4-O-methylglucuronoxylans (MGX) from *Castanea sativa* hardwood (Barbat et al. 2010). Chestnut sawdust was first delignified using metal-phthalocyanine or porphyrin in presence of hydrogen peroxide. Then, MGXs were easily extracted by hot water.

The parts of chestnut such as: seed, peeled seed, brown seed shell, red internal seed shell, leaves, catkin, spiny bur, as well as the new and old chestnut bark were found to have varying amounts of total phenolics, total flavonoids and tannins (Živković et al. 2009a, b). The highest content of total phenolic compounds (3.28)% GAE (Gallic acid equivalent) was found in dry extract of catkin, while the lowest content (0.42)% GAE was obtained for the dry extract of the seeds. The phenolic contents (% GAE) in other tissues were as follows: peeled chestnut 0.59%, brown seed shell 1.19%, red internal seed shell 2.82%, leaf 1.40%, new chestnut bark 3.00%, old chestnut bark 1.70%, spiny burs 0.49%. The total flavonoid content % CE (catechin equivalent) in various plant parts were reported as: seeds 0.17%, brown seed shell 0.65%, peeled chestnut 0.09%, red internal seed shell 1.44%, catkin 0.60%, leaf 1.40%, new chestnut bark 0%, old chestnut bark 0.69%, spiny burs 0.13%. The total condensed tannins (CT) content was highest in red internal seed shell 15.29% CE (vanillin assay) and 3.12% CT (acid butanol assay). The total condensed tannins content in other plant parts were estimated as follows: seeds 0.39% CE (vanillin assay) 0.88% CT (butanol assay); peeled chestnut 0% CE (vanillin assay), 0% CT (acid butanol assay); brown seed shell 2.78% CE (vanillin assay), 1.67% CT (acid butanol assay); catkin 0.49% CE (vanillin assay), 0.95% CT (acid butanol assay); leaf 0% CE (vanillin assay) 0.08% CT (acid butanol assay); new chestnut bark 3.91% CE (vanillin assay), 1.89% CT (acid butanol assay); old chestnut bark 0.76% CE (vanillin assay), 0.58% CT (acid butanol assay); and spiny burs 0% CE (vanillin assay), 0.08% (acid butanol assay).

## Antioxidant Activity

Studies had shown that the leaves had antioxidant properties. The leaf extract and an ethyl acetate fraction, which contained a high level of total phenolic compounds (29.1 g/100 g) demonstrated high antioxidant potentials, equivalent to at least those of reference compounds quercetin and vitamin E and standard extracts (pycnogenol, from French *Pinus maritima* bark, and grape marc extract) (Calliste et al. 2005). In another study, five phenolic compounds were identified in the leaf ethanol:water (7:3) extract, namely chlorogenic acid, ellagic acid, rutin, isoquercitrin and hyperoside (Almeida et al. 2008a). The content of total phenolics for *C. sativa* was 284 of gallic acid equivalents (GAE)/g of lyophilized extract. The extract presented a high potency to scavenge the tested reactive species viz. reactive oxygen species (ROS) such as superoxide radical, peroxy radical, hydrogen peroxide, singlet oxygen and reactive nitrogen species (RNS) namely nitric oxide and peroxynitrite; all the  $IC_{50}$ s being found at the  $\mu\text{g/ml}$  level. The  $IC_{50}$  found for the iron chelation and DPPH (1,1-diphenyl-2-picrylhydrazyl) scavenging assays were 132.94 and 12.58  $\mu\text{g/ml}$ , respectively.

The water soluble extracts obtained from leaves, catkins, and outer brown peel of *Castanea sativa* showed high antioxidant activity in scavenging  $\cdot\text{OH}$  and DPPH radical (Živković et al. 2009b). The highest content of total phenolic compounds ((4.24)% of GAE) and flavonoids ((2.41)% of GAE) were determined in dry extract of outer brown peel of Lovran's Marrone cultivar. The TF/TP (total flavonoid/total phenolic) ratio was from 11.86 for the extract of peeled chestnut to 56.83% determined in the outer brown peel of Lovran's Marrone cultivar. All extracts, except for sweet chestnut catkins, showed the ability to protect liposomes from peroxidation. Phenolic compounds, as active antioxidants, have the ability to enter and protect cell membranes from lipid peroxidation, thus overcoming the body's refractory response to the antioxidant supplements in the diet. It was shown that phenolics were easily accessible natural antioxidants that could be used as food supplements or for the

treatment of pathophysiological conditions related to oxidative stress. In a recent study, all the assayed by-products (almond green husks, chestnut skins and chestnut leaves) revealed good antioxidant properties, with very low  $EC_{50}$  values (lower than 380  $\mu\text{g/ml}$ ), particularly for lipid peroxidation inhibition (lower than 140  $\mu\text{g/ml}$ ) (Barreira et al. 2010). The correlation between the bioactive compounds (total phenols and flavonoids) and DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity, reducing power, inhibition of  $\beta$ -carotene bleaching and inhibition of lipid peroxidation in pig brain tissue through formation of thiobarbituric acid reactive substances, was also obtained. Although, all the assayed by-products proved to have a high potential of application in new antioxidants formulations, chestnut skins and leaves demonstrated better results. Studies proposed that glycerine with a pH around 5 was best vehicle for topical formulations incorporating *C. sativa* leaf extract with antioxidant activity for the prevention of photoaging and oxidative-stress-mediated diseases (Almeida et al. 2010).

Gamma irradiation is a possible feasible alternative to substitute the traditional quarantine chemical fumigation with methyl bromide to extend post-harvest shelf-life of food. Results of studies appeared to indicate that storage favoured chestnuts antioxidant potential as assessed by DPPH (2,2-diphenyl-1-picrylhydrazyl) radical-scavenging activity, reducing power and inhibition of  $\beta$ -carotene bleaching capacity (Antonio et al. 2011). In addition, the application of gamma irradiation also appeared to be advantageous for antioxidant activity, independently of the dose used (0.27 or 0.54 kGy).

### **Photoprotective Activity**

The in-vivo patch test carried out on 20 volunteers showed that, with respect to irritant effects, chestnut leaf extract could be regarded as safe for topical application (Almeida et al. 2008b). Topical application of natural antioxidants had proven to be effective in protecting the skin against ultraviolet-mediated oxidative damage

and to provide a straightforward way to strengthen the endogenous protection system. Further studies proposed that glycerine with a pH around 5 was the best vehicle for topical formulations incorporating *C. sativa* leaf extract with antioxidant activity for the prevention of photoaging of the skin and oxidative-stress-mediated diseases (Almeida et al. 2010).

### **Anticancer Activity**

The concentration of ellagic acid, a naturally occurring inhibitor of carcinogenesis, in chestnut fruits and bark was generally increased after hydrolysis due to the presence of ellagitannins in the crude extract (Vekiari et al. 2008). The concentration varied between 0.71 and 21.6 mg/g (d.w.) in non-hydrolyzed samples, and between 2.83 and 18.4 mg/g (d.w.) in hydrolyzed samples. In chestnut fruits, traces of ellagic acid were present in the seed, with higher concentrations in the pellicle and pericarp. However, all fruit tissues had lower concentrations of ellagic acid than the bark.

Xylans, 4-O-methylglucuronoxylan (MGX) and a homoxylan (HX), were purified from delignified holocellulose alkaline extracts of *Castanea sativa* (Barbat et al. 2008). The xylan, MGX inhibited the proliferation of A431 human epidermoid carcinoma cells with an  $IC_{50}$  value of 50  $\mu\text{M}$  (Moine et al. 2007; Barbat et al. 2008). In addition, this xylan inhibited A431 cell migration and invasion. Preliminary experiments showing that secretion of metalloproteinases MMP2 and MMP9 by A431 tumour cells was inhibited by the purified MGX strongly suggesting that this mechanism of action may play a role in its anti-migration and anti-invasive properties.

### **Antigastroenteritic Activity**

Seeds of *C. sativa* were reported to be used in paediatrics for treatment of gastroenteritis and as a gluten-free diet in cases of coeliac disease (Živković et al. 2009a, b). A new pyrrole alkaloid, methyl-(5-formyl-1*H*-pyrrole-2-yl)-4-hydroxybutyrate, was

isolated from sweet chestnut seeds (Hiermann et al. 2002).

### **Antibacterial Activity**

An ethyl acetate extract of *C. sativa* was shown to have pronounced antibacterial effects against seven of the eight strains of Gram-positive and Gram-negative bacteria used (MIC in the range of 64–256 µg/ml and MBC in the range of 256–512 µg/ml) (Basile et al. 2000). The active fraction contained rutin, hesperidin, quercetin, apigenin, morin, naringin, galangin and kaempferol. The highest bactericidal activity was shown by quercetin, rutin and apigenin. was tested.

Staphylococcal quorum sensing is encoded by the AGR locus and is responsible for the production of δ-hemolysin (Quave et al. 2011). Studies found that the extracts from three medicinal plants (*Ballota nigra*, *Castanea sativa*, and *Sambucus ebulus*) exhibited a dose-dependent response in the production of δ-hemolysin, indicating anti-quorum sensing activity in a pathogenic methicillin-resistant *Staphylococcus aureus* (MRSA) isolate.

### **Antithrombin Activity**

Of selected Slovak medicinal plants, the methanol extract of *C. sativa* exhibited the most distinct inhibition activity to thrombin with  $IC_{50} = 73.2$  µg/ml (Jedinak et al. 2010). The methanol extract also had strong inhibition activity to protease trypsin with  $IC_{50}$  values below 10 µg/ml.

### **Antidiarrhoeal Activity**

Stop Fitán®, a dietary supplement based on the bioactive purified natural extract of chestnut (*Castanea sativa*) wood and *Saccharomyces boulardii*, a nonpathogenic yeast strain has been proposed as a co-adjuvant in the therapy of diarrhea (Budriesi et al. 2010). Studies showed that natural extract of chestnut wood rich in hydrolyzable tannins, exerted spasmolytic effects in guinea pig ileum and proximal colon. The findings, coupled

with the antibacterial, antiviral, and antispasmodic properties of tannins, suggest that the combination of chestnut tannins and *S. boulardii* may be relevant to treat diarrhea by Stop Fitán®.

### **Hypersensitivity Problem**

Only occasionally does chestnut cause hypersensitivity. There were only a few reported cases, depending on cross-reactivity in previously latex-hypersensitive patients. The case of oral allergy syndrome to chestnut appeared to be a manifestation of immediate IgE-dependent hypersensitivity resulting from direct contact between food and the oral mucosa (Antico 1996).

### **Traditional Medicinal Uses**

In the Middle Ages the raw seeds were found useful in the treatment of heart disorders. *C. sativa* leaves were used in folk medicine as a tea in France to treat whooping cough and diarrhoea. Beside the leaves, the bark is also a good source of tannins. They are antiinflammatory, expectorant, tonic and astringent. The astringent activity is useful in the treatment of bleeding and diarrhoea. Leaf infusions are employed in respiratory diseases and are a common therapy for whooping cough, fevers and ague. The leaves are also employed in the treatment of rheumatism, to ease lower back pains and to relieve stiff muscles and joints. A decoction is used as gargle for treating sore throats. A hair shampoo can be made from infusing leaves and fruit husks. Leaves are used in homeopathy for therapy of depression and fatigue.

### **Other Uses**

The Spanish chestnut is a magnificent huge shade tree for parks, estates and avenues and sometimes planted for erosion control in Mediterranean countries. The flowers provide good forage for bees. A blackish-brown dye is extracted from the leaves and the bark and also an oil which is medicinal. Tannin is obtained from the bark and



used in tanning. The wood, leaves and seed husks also contain tannin. The seed meal can be used as a source of starch, for fattening stock and also for whitening linen cloth. A hair shampoo is made from the leaves and the husks of the fruits. The wood is good for carpentry, turnery, furniture, barrels, roof beams, props, basketry and fence posts. It is also a very good fuel.

---

## Comments

Chestnut tree responds well to coppicing, a traditional method of woodland management.

---

## Selected References

- Almeida IF, Fernandes E, Lima JL, Costa PC, Bahia MF (2008a) Protective effect of *Castanea sativa* and *Quercus robur* leaf extracts against oxygen and nitrogen reactive species. *J Photochem Photobiol B* 91(2–3):87–95
- Almeida IF, Valentão P, Andrade PB, Seabra RM, Pereira TM, Amaral MH, Costa PC, Bahia MF (2008b) In vivo skin irritation potential of a *Castanea sativa* (Chestnut) leaf extract, a putative natural antioxidant for topical application. *Basic Clin Pharmacol Toxicol* 103(5):461–467
- Almeida IF, Costa PC, Bahia MF (2010) Evaluation of functional stability and batch-to-batch reproducibility of a *Castanea sativa* leaf extract with antioxidant activity. *AAPS PharmSciTech* 11(1):120–125
- Antico A (1996) Oral allergy syndrome induced by chestnut (*Castanea sativa*). *Ann Allergy Asthma Immunol* 76(1):37–40
- Antonio AL, Fernandes A, Barreira JC, Bento A, Botelho ML, Ferreira IC (2011) Influence of gamma irradiation in the antioxidant potential of chestnuts (*Castanea sativa* Mill.) fruits and skins. *Food Chem Toxicol* 49(9):1918–1923
- Bailey LH (1976) *Hortus third. A concise dictionary of plants cultivated in the United States and Canada.* Liberty Hyde Bailey Hortorium, Cornell University, Wiley, London, 1312 pp
- Barbat A, Gloaguen V, Moine C, Sainte-Catherine O, Kraemer M, Rogniaux H, Ropartz D, Krausz P (2008) Structural characterization and cytotoxic properties of a 4-O-methylglucuronoxylan from *Castanea sativa*. 2. Evidence of a structure-activity relationship. *J Nat Prod* 71(8):1404–1409
- Barbat A, Gloaguen V, Sol V, Krausz P (2010) Aqueous extraction of glucuronoxylans from chestnut wood: new strategy for lignin oxidation using phthalocyanine or porphyrin/H<sub>2</sub>O<sub>2</sub> system. *Bioresour Technol* 101(16):6538–6544
- Barreira JC, Casal S, Ferreira IC, Oliveira MB, Pereira JA (2009) Nutritional, fatty acid and triacylglycerol profiles of *Castanea sativa* Mill. cultivars: a compositional and chemometric approach. *J Agric Food Chem* 57(7):2836–2842
- Barreira JCM, Ferreira ICFR, Oliveira MPP, Pereira JA (2010) Antioxidant potential of chestnut *Castanea sativa* L., and almond, *Prunus dulcis* L., by-products. *Food Sci Technol Int* 16:209–216
- Basile A, Sorbo S, Giordano S, Ricciardi L, Ferrara S, Montesano D, Castaldo Cobianchi R, Vuotto ML, Ferrara L (2000) Antibacterial and allelopathic activity of extract from *Castanea sativa* leaves. *Fitoterapia* 71(Suppl 1):S110–S116
- Bounous G, Botta R, Beccaro G (2000) The chestnut: the ultimate energy source. Nutritional value and alimentary benefits. *Nucis – Information Bulletin of the Research Network on Nuts (FAO-CIHEAM)* 9:44–50
- Budriesi R, Ioan P, Micucci M, Micucci E, Limongelli V, Chiarini A (2010) Stop Fitan: antispasmodic effect of natural extract of chestnut wood in guinea pig ileum and proximal colon smooth muscle. *J Med Food* 13(5):1104–1110
- Calliste CA, Trouillas P, Allais DP, Duroux JL (2005) *Castanea sativa* Mill. leaves as new sources of natural antioxidant: an electronic spin resonance. *J Agric Food Chem* 53(2):282–288
- Chevallier A (1996) *The encyclopedia of medicinal plants.* Dorling Kindersley, London, 336 pp
- De Vasconcelos MCBM, Bennett RN, Rosa EAS, Ferreira-Cardoso JV (2007) Primary and secondary metabolite composition of kernels from three cultivars of Portuguese chestnut (*Castanea sativa* Mill.) at different stages of industrial transformation. *J Agric Food Chem* 55(9):3508–3516
- De Vasconcelos MCBM, Bennett RN, Quideau S, Jacquet R, Rosa EAS, Ferreira-Cardoso JV (2010a) Evaluating the potential of chestnut (*Castanea sativa* Mill.) fruit pericarp and integument as a source of tocopherols, pigments and polyphenols. *Ind Crops Res* 31(2):301–311
- De Vasconcelos MCBM, Bennett R, Rosa EAS, Ferreira-Cardoso JV, Nunes F (2010b) Industrial processing of chestnut fruits (*Castanea sativa* Mill.) – effects on nutrients and phytochemicals. *Acta Horti (ISHS)* 866:611–617
- De Vasconcelos MCBM, Nunes F, Viguera CG, Bennett RN, Rosa EAS, Ferreira-Cardoso JV (2010c) Industrial processing effects on chestnut fruits (*Castanea sativa* Mill.) 3. Minerals, free sugars, carotenoids and antioxidant vitamins. *Int J Food Sci Technol* 45(3):496–505
- Gonçalves B, Borges P, Costa HS, Bennett R, Santos M, Silva AP (2010) Metabolite composition of chestnut (*Castanea sativa* Mill.) upon cooking: Proximate analysis, fibre, organic acids and phenolics. *Food Chem* 122(1):154–160
- Hiermann A, Kedwani S, Schramm HW, Seger C (2002) A new pyrrole alkaloid from seeds of *Castanea sativa*. *Fitoterapia* 73:22–27

- Jedinak A, Valachova M, Maliar T, Sturdik E (2010) Antiprotease activity of selected Slovak medicinal plants. *Pharmazie* 65(2):137–140
- Miller G, Miller DD, Jaynes RA (1996) Chestnuts. In: Janick J, Moore JN (eds) *Fruit breeding*, vol 3, Nuts. Wiley, New York, 278 pp
- Mills SY (1985) *The dictionary of modern herbalism*. Thorsons, Wellingborough
- Moine C, Krausz P, Chaleix V, Sainte-Catherine O, Kraemer M, Gloaguen V (2007) Structural characterization and cytotoxic properties of a 4-O-methylglucuronoxylan from *Castanea sativa*. *J Nat Prod* 70(1):60–66
- Quave CL, Plano LR, Bennett BC (2011) Quorum sensing inhibitors of *Staphylococcus aureus* from Italian medicinal plants. *Planta Med* 77(2):188–195
- Ribeiro B, Rangel J, Valentão P, Andrade PB, Pereira JA, Böлке H, Seabra RM (2007) Organic acids in two Portuguese chestnut (*Castanea sativa* Miller) varieties. *Food Chem* 100(2):504–508
- Rutter PA, Miller G, Payne JA (1990) Chestnuts (*Castanea*). *Acta Hort* 290(2):761–788
- Sanz M, Cadahía E, Esteruelas E, Muñoz AM, Fernández de Simón B, Hernández T, Estrella I (2010) Phenolic compounds in chestnut (*Castanea sativa* Mill.) heartwood. Effect of toasting at cooperage. *J Agric Food Chem* 58(17):9631–9640
- U.S. Department of Agriculture, Agricultural Research Service (2011) USDA National Nutrient Database for standard reference, release 24. Nutrient Data Laboratory Home Page. <http://www.ars.usda.gov/ba/bhnrc/ndl>
- Vekiari SA, Gordon MH, García-Macías P, Labrinea H (2008) Extraction and determination of ellagic acid content in chestnut bark and fruit. *Food Chem* 110(4):1007–1011
- Yıldiz MU, Özcan MM, Çalır S, Demir F, Er F (2009) Physico-chemical properties of wild chestnut (*Castanea sativa* Mill.) fruit grown in Turkey. *World Appl Sci J* 6(3):365–372
- Živković J, Mujić I, Zeković Z, Nikolić G, Vidović S, Mujić A (2009a) Extraction and analysis of condensed tannins in *Castanea sativa* Mill. *J Cent Eur Agric* 10(3):283–288
- Živković J, Zeković Z, Mujić I, Vesna Tumbas V, Cvetković D, Spasojević I (2009b) Antioxidant properties of chestnut phenolics. *Food Technol Biotechnol* 47(4):421–427