

P.L. Smedley and David G. Kinniburgh

**Contents**

12.1 Introduction .....	279
12.2 Sources of Arsenic in the Natural Environment .....	280
12.3 Arsenic in Groundwater .....	283
12.4 Mineral-Water Interactions .....	298
12.5 Common Features of High-Arsenic Groundwater Provinces .....	300
12.6 Concluding Remarks .....	304
References .....	304

**12.1 Introduction**

Awareness of the problems associated with arsenic in drinking water and the environment has grown significantly over the last two decades or so and today an enormous literature exists documenting its occurrence, behaviour and impacts in many places across the globe. The mobilisation of arsenic in the environment occurs through a complex combination of natural biogeochemical reactions and human interactions. Most recognised problems are generated by mobilisation and transport under natural conditions, but mobilisation has also been caused, or exacerbated, by mining, fossil-fuel combustion and use of synthetic arsenical compounds (pesticides, herbicides, crop desiccants and arsenic-based additives in livestock feed). Arsenical pesticides and herbicides have been used much less over the last few decades, and more recent restrictions have been imposed on the use of arsenic in wood preservation (e.g. European Communities' Directive 2003/2/EC), but the legacy of such sources may still pose a localised threat to the environment.

Human exposure to arsenic occurs through a number of pathways, including air, food, water and soil. The relative impacts of these vary depending on local circumstances but of the potential pathways, drinking water poses one of the greatest threats to human health as borne out by the large number of documented case histories from around the world. The concentrations of arsenic in drinking water are very variable, depending on nature of source (surface water, groundwater, rainwater) and local conditions, and observed ranges vary over several orders of magnitude. Excepting localised sources of anthropogenic contamination, the highest aqueous arsenic concentrations are usually found in groundwaters because of the high solid/solution ratios found in aquifers. Groundwaters therefore pose the greatest overall threat to health. Groundwaters with arsenic concentrations sufficiently high to be detrimental to humans, or with already detectable health impacts have been reported in Argentina, Bangladesh, Burkina Faso, Cambodia, Chile, China, India,

---

P.L. Smedley (✉) • D.G. Kinniburgh  
British Geological Survey, Oxfordshire, UK  
e-mail: [pls@bgs.ac.uk](mailto:pls@bgs.ac.uk)

Hungary, Laos, Mexico, Nepal, Romania, Spain, Taiwan, Thailand and Vietnam and occasional problems are found in many other countries.

Some of the groundwater arsenic problems have been recognised for a considerable time. Probably the earliest cases of health effects from arsenic contamination of drinking water were recognised in a mining area of Poland in the 1890s. Here, contamination of water supplies by oxidation of arsenic-bearing sulphide minerals produced localised health problems (see Tseng et al. 1968). In central Argentina, arsenic-related health problems were first documented in 1917 and problems in Taiwan were also first identified in the 1910s. In each of these cases, solutions to the problems were sought primarily through engineering which for the most part provided alternative supplies of surface water or of treated water. As a result, geochemical investigations into the processes controlling the arsenic mobilisation in these areas were not carried out until much more recently and awareness of the problems observed in impacted regions such as the Bengal Basin, Cambodia and Vietnam have developed many years later.

European and US EPA regulations implement the current recommended WHO provisional guideline value for arsenic in drinking water of  $10 \mu\text{g L}^{-1}$ . However, many countries continue to use the WHO pre-1993 guideline value of  $50 \mu\text{g L}^{-1}$  as their national standard, in part because of the difficulties of testing low concentrations and in part due to difficulties with compliance. Concentrations of  $10 \mu\text{g L}^{-1}$  and  $50 \mu\text{g L}^{-1}$  are therefore both still used as reference points for the testing and reporting of arsenic.

This chapter reports the current state of knowledge on the occurrence of arsenic in natural waters and their host rocks across the world and attempts to describe what are currently understood to be the main biogeochemical processes that control its mobilisation in the environment (see also Chaps. 2, 24, and 25, this volume).

## 12.2 Sources of Arsenic in the Natural Environment

### 12.2.1 Minerals

Arsenic occurs as a major constituent in more than 200 minerals, including elemental arsenic, arsenides, sulphides, oxides, arsenates and arsenites. Most are ore minerals or their alteration products. These minerals are relatively rare in the natural environment. Among the most common occurrences in ore zones are arsenian pyrite ( $\text{Fe}(\text{S,As})_2$ ), arsenopyrite ( $\text{FeAsS}$ ), löllingite ( $\text{FeAs}_2$ ), realgar ( $\text{AsS}$ ), orpiment ( $\text{As}_2\text{S}_3$ ), cobaltite ( $\text{CoAsS}$ ), niccolite ( $\text{NiAs}$ ) and scorodite ( $\text{FeAsO}_4 \cdot 2\text{H}_2\text{O}$ ). Arsenian pyrite ( $\text{Fe}(\text{As,S})_2$ ) is probably the most important source of arsenic in ore zones (Deditius et al. 2008; Savage et al. 2000) and concentrations

**Table 12.1** Typical ranges of arsenic concentrations in rock-forming and accessory minerals

Mineral	Arsenic concentration range ( $\text{mg kg}^{-1}$ )
<b>Sulphide minerals</b>	
Pyrite	100–130,000
Pyrrhotite	5–100
Marcasite	20–126,000
Galena	5–10,000
Sphalerite	5–17,000
Chalcopyrite	10–5,000
<b>Oxide minerals</b>	
Haematite	up to 160
Fe(III) oxyhydroxide	up to 76,000
Magnetite	2.7–41
Ilmenite	<1
<b>Silicate minerals</b>	
Quartz	0.4–1.3
Feldspar	<0.1–2.1
Biotite	1.4
Amphibole	1.1–2.3
Olivine	0.08–0.17
Pyroxene	0.05–0.8
<b>Carbonate minerals</b>	
Calcite	1–260
Dolomite	<3
Siderite	<3
<b>Sulphate minerals</b>	
Gypsum/anhydrite	<1–6
Barite	<1–12
Jarosite	34–2,000
<b>Other minerals</b>	
Apatite	<1–1,000
Halite	<3–30
Fluorite	<2

Barker et al. (2009), Di Benedetto et al. (2006), Savage et al. (2005), Smedley and Kinniburgh (2002) and references therein

in excess of 10 wt.% (up to 19 wt.%) As have been reported (e.g. Barker et al. 2009; Blanchard et al. 2007; Reich and Becker 2006). Reich and Becker (2006) concluded from first-principles calculations that pyrite and marcasite polymorphs can contain up to 6 wt.% As in solid solution, beyond which unmixing to separate pyrite/marcasite and arsenopyrite occurs. The oxidation state of As in arsenian pyrite is most typically  $\text{As}^{1-}$  in substitution for S, although recent occurrences of  $\text{As}^{3+}$ -bearing pyrite ( $(\text{Fe,As})\text{S}_2$ ) have been documented in some hydrothermal gold deposits, signalling precipitation in more oxidising conditions than in the more common  $\text{As}^{1-}$  form (Chouinard et al. 2005; Deditius et al. 2008).

Other rock-forming sulphide minerals, including chalcopyrite and galena, also contain often high concentrations of arsenic (Table 12.1) as well as a range of other trace elements (transition metals, Cd, Pb, Ag, Au, Sb, P, W and Mo). Zonation of trace elements within sulphide grains is common.

Besides being an important component of ore bodies, pyrite is also formed in low-temperature sedimentary environments under sulphide-reducing conditions. Such authigenic pyrite plays an important role in present-day geochemical cycles, being present in the sediments of rivers, lakes, oceans and aquifers. Authigenic realgar has also been observed in shallow aquifer sediments (O'Day et al. 2004). In sulphidic solutions, soluble arsenic will likely be incorporated into pyrite through surface arsenite adsorption (Bostick and Fendorf 2003). Pyrite is unstable in aerobic systems and oxidises to form iron oxides with release of associated arsenic. A lower stability of arsenian pyrite compared to low-As pyrite also potentially favours As release (Blanchard et al. 2007; Savage et al. 2000). The presence of pyrite in mineralised veins is responsible for the production of acid mine drainage, and for the common presence of arsenic problems around coal mines and areas of intensive coal burning.

Arsenic is also found in large concentrations in many oxide minerals and hydrous metal oxides, either as part of the mineral structure or as adsorbed species. Concentrations in Fe oxides can reach several weight percent (Table 12.1), particularly where they form as the oxidation products of primary As-bearing iron sulphide minerals. Adsorption of arsenic to hydrous iron oxides is particularly strong, depending on redox speciation, pH, solute compositions and iron-oxide form; adsorbed loadings can be great even where concentrations in solution are low. Adsorption to hydrous Al and Mn oxides may also be important if present in quantity (Goldberg 1986; Stollenwerk 2003). Sorption to the edges of clays may occur, although the loadings are much smaller on a weight basis than for the iron oxides (Smedley and Kinniburgh 2002).

Concentrations in phosphate minerals can be relatively high, for example up to 1,000 mg kg<sup>-1</sup> in apatite (Table 12.1). However, phosphate minerals are much less abundant than oxide minerals and so make a correspondingly small contribution to the arsenic concentration in most sediments. Arsenic tends to be present at much smaller concentrations in the common rock-forming minerals, although low concentrations are invariably present. Most common silicate minerals (including quartz, feldspar, micas, amphiboles) contain around 1 mg kg<sup>-1</sup> or less (Table 12.1). Concentrations of arsenic in carbonate minerals are also usually low, of the order of a few mg kg<sup>-1</sup>, although Di Benedetto et al. (2006) found up to 257 mg kg<sup>-1</sup> in Italian travertines.

### 12.2.2 Rocks, Sediments and Soils

Arsenic occurs ubiquitously but at variable concentrations in rocks, unconsolidated sediments and soils. Average crustal abundance is around 1.5 mg kg<sup>-1</sup>. Igneous rocks usually

have concentrations of around 5 mg/kg or less. Volcanic glasses have on average around 6 mg kg<sup>-1</sup> (Smedley and Kinniburgh 2002). Metamorphic rocks tend to have concentrations that reflect those of their igneous and sedimentary precursors. Most also contain around 5 mg kg<sup>-1</sup> or less. Pelitic rocks (slates, phyllites) typically have the highest concentrations (Table 12.2), with on average of around 18 mg kg<sup>-1</sup> (see also Chap. 2, this volume).

Concentrations in sedimentary rocks are typically in the range 5–10 mg kg<sup>-1</sup> (Webster 1999). Average sediments are enriched in arsenic relative to igneous and metamorphic rocks because they contain greater quantities of minerals with high adsorbed arsenic loads. Of the sediments, sands and sandstones have lower concentrations which reflect the dominance of low-arsenic minerals (i.e. quartz and feldspars). Average sandstone arsenic concentrations are around 4 mg kg<sup>-1</sup>.

Argillaceous deposits have a broader range of concentrations than sandstones (Table 12.2), being on average around 13 mg kg<sup>-1</sup>. Higher values reflect the larger proportion of sulphide minerals, oxides, organic matter and clays present. Black shales have concentrations at the upper end of the range, principally because of their enhanced pyrite content (Table 12.2). Marine clays appear to have higher concentrations than non-marine equivalents. This may also be due to a higher proportion of pyrite in offshore pelagic sediments.

German organic-rich shales have arsenic concentrations of 100–900 mg kg<sup>-1</sup> (Riedel and Eikmann 1986). Some coal samples have very high concentrations, up to 35,000 mg kg<sup>-1</sup> (Belkin et al. 2000) (Table 12.2), although lower concentrations of 2.5–17 mg kg<sup>-1</sup> are more typical. Some of the highest observed arsenic concentrations are found in ironstones and Fe-rich rocks. Concentrations of arsenic up to 2,900 mg kg<sup>-1</sup> have been reported for such rocks (Table 12.2, references in Smedley and Kinniburgh 2002). Phosphorites are also relatively enriched in arsenic, with concentrations up to ca. 400 mg kg<sup>-1</sup>. Data for carbonate rock types usually indicate relatively low arsenic concentrations (Tables 12.1 and 12.2).

Unconsolidated sediments have compositions which do not differ significantly from their indurated equivalents. Much evidence indicates that muds and clays have higher concentrations than most sands and carbonates. Values are typically 3–10 mg kg<sup>-1</sup>, depending on texture and mineralogy (Table 12.2). High concentrations reflect abundance of pyrite or iron oxides. There is often a significant positive correlation between the iron and arsenic concentrations in sediments. High concentrations are also common in mineralised areas. Placer deposits in streams can have concentrations reflecting the abundance of sulphide minerals. Average arsenic concentrations for stream sediments in England and Wales have been reported in the range 5–8 mg kg<sup>-1</sup> (Table 12.2, references in Smedley and Kinniburgh 2002), although a higher median value (13.4 mg kg<sup>-1</sup>) has been computed

**Table 12.2** Typical ranges of arsenic concentrations in rocks, sediments, soils and other superficial deposits

Rock/sediment type	As concentration range (mg kg <sup>-1</sup> )
<b>Igneous rocks</b>	
Ultrabasic rocks	0.03–15.8
Basic rocks	0.06–113
Intermediate rocks	0.09–13.4
Acidic rocks	0.2–15
<b>Metamorphic rocks</b>	
Quartzite	2.2–7.6
Hornfels	0.7–11
Phyllite/slate	0.5–143
Schist/gneiss	<0.1–18.5
Amphibolite and greenstone	0.4–45
<b>Sedimentary rocks</b>	
Marine shale/mudstone	3–15 (up to 490)
Shale (Mid-Atlantic Ridge)	48–361
Non-marine shale/mudstone	3.0–12
Sandstone	0.6–120
Limestone/dolomite	0.1–20.1
Phosphorite	0.4–188
Iron formations and Fe-rich sediment	1–2,900
Evaporites (gypsum/anhydrite)	0.1–10
Coals	0.3–35,000
<b>Unconsolidated sediments</b>	
Alluvial sand (Bangladesh)	1.0–6.2
Alluvial mud/clay (Bangladesh)	2.7–14.7
River bed sediments (Bangladesh)	1.2–5.9
Lake sediments	0.5–44
Glacial till	1.9–170
World average river sediments	5
Stream and lake silt	<1–72
Stream sediment, England and Wales	5–8
Loess silts, Argentina	5.4–18
Continental margin sediments	2.3–8.2
<b>Soils</b>	
Mixed soils	0.1–55
Peaty and bog soils	2–36
Peat	up to 9
Acid sulphate soils	1.5–45
Soils near sulphide deposits	2–8,000
<b>Contaminated superficial deposits</b>	
Mining-contaminated lake sediment	80–1,104
Mining-contaminated reservoir sediment	100–800
Mine tailings	396–2,000
Soils and tailings-contaminated soil	120–52,600
Industrially polluted inter-tidal sediments	0.38–1,260
Orchard soils	366–732
Soils below chemicals factory	1.3–4,770

From sources summarised in Smedley and Kinniburgh (2002)

from a more recent survey in eastern England (BGS data, unpublished). In the Bengal Basin, river sediments with concentrations averaging 2.0, 2.8 and 3.5 mg kg<sup>-1</sup> were found for samples from the Ganges, Brahmaputra and Meghna rivers respectively (Datta and Subramanian 1997).

Cook et al. (1995) found concentrations in Canadian lake sediments ranging between 0.9 and 44 mg kg<sup>-1</sup> but noted that the highest concentrations were present up to a few kilometres down-slope of mineralised areas. They also found concentrations in glacial till of 1.9–170 mg kg<sup>-1</sup>

(Table 12.2) and noted the highest concentrations down-ice of mineralised areas.

In soils, concentrations of arsenic are of the order of 5–15 mg kg<sup>-1</sup>, with an average of 7.2 mg kg<sup>-1</sup> quoted for world soils and 7.4 mg kg<sup>-1</sup> for American soils (Table 12.2, references in Smedley and Kinniburgh 2002). Peats and bog soils can have higher concentrations (average 13 mg kg<sup>-1</sup>; Table 12.2), but this is principally because of an increased prevalence of sulphide mineral phases under the reduced conditions. Shotyk (1996) found up to 9 mg As kg<sup>-1</sup> in peat profiles and in a profile with lower mineral content, i. e. purer peat, the As content was 1 mg kg<sup>-1</sup> or lower.

Acid sulphate soils can be generated by the oxidation of pyrite in sulphide-rich terrains including mineralised areas and dewatered mangrove swamps and as such can be relatively enriched in arsenic. Acid sulphate soils from Canada have arsenic concentrations up to 45 mg kg<sup>-1</sup> (Dudas 1984). Gustafsson and Tin (1994) found similarly high concentrations (up to 41 mg kg<sup>-1</sup>) in acid sulphate soils from the Mekong delta of Vietnam.

Additional arsenic inputs to soils may be derived locally from industrial sources such as smelting and fossil-fuel combustion products and agricultural sources such as pesticides and phosphate fertilisers. Concentrations in the range 366–732 mg kg<sup>-1</sup> have been found in orchard soils following historical application of arsenical pesticides to fruit crops (Table 12.2). Long-term use of phosphate fertilisers may also add arsenic to soil.

Concentrations of arsenic in sediments and soils contaminated by the products of mining activity, including tailings and effluent, can have arsenic concentrations orders of magnitude higher than under natural conditions. Tailings piles and tailings-contaminated soils can contain up to several thousands of mg kg<sup>-1</sup> (Table 12.2). The values reflect not only increased abundance of primary arsenic-rich sulphide minerals, but also secondary iron arsenates and iron oxides formed as reaction products of the original ore minerals. The primary sulphide minerals are susceptible to oxidation in the tailings pile and the secondary minerals have varying solubility under oxidising conditions in groundwaters and surface waters. Scorodite (FeAsO<sub>4</sub>·2H<sub>2</sub>O) is a common sulphide oxidation product and its solubility is likely to control arsenic concentrations in such environments. Secondary arsenolite (As<sub>2</sub>O<sub>3</sub>) is also often represented in such environments.

## 12.3 Arsenic in Groundwater

### 12.3.1 Aqueous Speciation

Analytical capability for aqueous arsenic species has developed rapidly and modern automated hyphenated chromatographic techniques (e.g. Bednar et al. 2004) have improved on traditional methods using hydride generation,


repeat measurements and determination of species by difference. Sensitive detection can be achieved with methods such as ICP-MS and this coupled with field-based (solid-phase extraction) or laboratory-based (e.g. HPLC) techniques allows for the direct measurement of aqueous organic forms alongside the more abundant inorganic As(III) and As(V). Field-based measurements counteract potential problems with instability in sample storage although significant problems with matrix effects using solid-phase extraction (Bednar et al. 2004) mean that this technique can be unsuitable for analysis of e.g. SO<sub>4</sub>-rich acid minewaters, saline geothermal waters or alkaline, SO<sub>4</sub>- and oxyanion-rich groundwaters, at least without large sample dilutions.

Arsenic can occur in the environment in a number of oxidation states (−3, −1, 0, +3 and +5) but in natural waters is found principally in inorganic form as arsenite (As(III)) or arsenate (As(V)), depending on redox conditions. Arsenic can form aqueous complexes with carbonate ligands, and As(III) with reduced sulphur (thioarsenites), which may be significant in some groundwaters.

Organic arsenic species produced by microbiological methylation reactions may be present in surface waters and may even dominate in organic-rich soil solutions, but are rarely present in quantity in groundwaters (Bednar et al. 2004; Chen et al. 1995; Das et al. 1995; Del Razo et al. 1990; Kondo et al. 1999; Lin et al. 1998). Exceptions potentially occur locally in cases of industrial or agricultural pollution. In practice concentrations of organic compounds in soil solutions rarely exceed a few micrograms per litre (Huang and Matzner 2006).

The ratio of As(III) to As(V) in groundwaters depends upon the abundance of the redox-active solids, including organic carbon and iron/manganese oxide, the flux of potential oxidants (oxygen, nitrate and sulphate) and on microbial activity. In geothermal systems, As(III) usually forms the main solute component but this can oxidise rapidly in surface conditions when mixed with surface water and especially when facilitated by bacteria (Wilkie and Hering 1998). Arsenic(III) also dominates in strongly reducing aquifers, typified by Fe(III) and sulphate reduction, as expected from the redox sequence (Berg et al. 2007; BGS and DPHE 2001; Chen et al. 1994; Das et al. 1995; Postma et al. 2007; Smedley et al. 2003; Stollenwerk et al. 2007; Tandukar et al. 2005; van Geen et al. 2006; Zheng et al. 2005). By contrast, in oxidising systems, solute arsenic is dominated by As(V) (Altamirano Espinoza and Bundschuh 2009; Bhattacharya et al. 2005; Del Razo et al. 1990; Robertson 1989; Smedley et al. 2002, 2007). The strong oxidising capacity of Mn oxides has been well-documented and implicated in arsenic redox speciation in several systems (Amirbahman et al. 2006; Oscarson et al. 1981a, b; Ying et al. 2011).

In natural systems, thermodynamic equilibrium in As species is often not achieved although abundant evidence indicates that reaction kinetics can be accelerated significantly by microbial activity, which have been shown to both oxidise



**Fig. 12.1** pe-pH predominance diagrams for the As-Fe system. (a) Aqueous As species only; (b) aqueous Fe species only; (c) As species where hydrous ferric oxide (*HFO*) has been allowed to precipitate when stable and sorb As, and (d) the same as (c) except that goethite has been assumed to be the stable iron-oxide mineral. Adsorbed species

concentrations have been calculated using the CD-MUSIC model. Where appropriate, total concentrations of  $Fe = 3e - 3 \text{ M}$  and  $As = 1e - 4 \text{ M}$ . A background concentration of 0.1 M NaCl was assumed in all cases. No reduced Fe minerals such as magnetite have been allowed to precipitate. All calculations were made by Phreeqc and plotted using PhreePlot

arsenite (Hering and Kneebone 2002; Wilkie and Hering 1998) and reduce arsenate (Oremland and Stolz 2003).

Sorption plays a key role in arsenic speciation and iron oxides in particular have a strong control on aqueous concentrations. Figure 12.1 shows pe-pH predominance diagrams for the As-Fe system. Figure 12.1a, b both show the redox and pH sensitivity of both aqueous As and Fe, with a range of protonated and hydrolysed species present over the natural pH range. Iron tends to be insoluble in oxic, near-neutral pH systems and forms a variety of iron oxides depending on the precise conditions and their evolution.

Hydrous ferric oxide (HFO), a high-surface-area form of iron oxide, often forms when Fe is precipitated rapidly. This oxide is able to adsorb As on its surface and the adsorbed As (HFO-As) can become the dominant form of As (Fig. 12.1c). HFO is subject to both acid dissolution at low pH and reductive dissolution at low pe (Fig. 12.1c) which result in the release of As to solution. At high pH, As is desorbed from HFO due to increasing electrostatic repulsion of the negatively charged oxide surface and the  $AsO_4^{3-}$  species, again leading to higher concentrations in solution. HFO tends to transform slowly to more stable

forms of iron oxide with lower specific surface area, such as goethite, with time. Goethite still has a significant adsorption affinity for As, albeit somewhat different from HFO due to the different crystal structure. Goethite is less prone than HFO to both acid dissolution and reductive dissolution and so the Goethite-As field occupies a correspondingly large area on the predominance diagram at low pH and low  $p_e$  (Fig. 12.1d). However, the lower specific surface area of goethite compared with HFO means that adsorption is not so great at high pH and so in goethite-dominated systems the aqueous As species tend to predominate at high pH.

Although these predominance diagrams illustrate some important geochemical interactions of Fe and As, they by no means tell the whole story. There are many other important interactions that have been omitted from these diagrams. For example, reduced iron oxides such as magnetite can form at low  $p_e$  and these too will tend to adsorb As (Dixit and Hering 2003). Also, significant concentrations of sulphide may be present in reduced systems leading to the potential formation of pyrite and other Fe-S minerals, again with scope for the adsorption of As on the mineral surface or the incorporation of As into the mineral structure as a solid solution. Nonetheless, the diagrams go some way to explaining observed occurrences of high-arsenic groundwater in some strongly reducing aquifers and at extremes of pH.

### 12.3.2 Arsenic Abundance and Distribution in Groundwater

Aquifers have high solid:solution ratios, typically 3–20 kg L<sup>-1</sup> and as a result, groundwaters within them are especially vulnerable to water-rock interaction and arsenic release. In addition, aquifers more often have the physical and chemical conditions favourable for arsenic mobilisation and transport than is the case in surface waters. Despite this, the occurrence of high arsenic concentrations in groundwaters is the exception rather than the rule. Background concentrations in groundwater are in most countries less than 10 µg L<sup>-1</sup> and often substantially lower. Nonetheless, values quoted in the literature show a very large range from <0.5 to 15,000 µg L<sup>-1</sup>, i.e. more than four orders of magnitude.

Mobilisation of arsenic in solution is favoured especially under oxidising conditions at high pH and under strongly reducing conditions. Evaporative concentration can also increase arsenic (and other element) concentrations substantially and may be important in some arid areas. Additional arsenic problems are encountered in some geothermal areas and in many areas of sulphide mineralisation and mining. Documented cases of industrial arsenic pollution (including those from agriculture) also exist. While these may be severe locally, occurrences are relatively rare and can usually be anticipated.

Investigations worldwide have revealed a number of major aquifers with significant groundwater arsenic

problems. The hydrogeological and geochemical conditions in these affected aquifers vary, although some common unifying features are apparent. Particular areas of risk are large alluvial and delta plains and sedimentary inland basins, the latter especially in arid and semi-arid areas. Geologically young (Quaternary) aquifers are particularly prone to developing and preserving high-arsenic groundwater. Alluvial and delta plains with recognised groundwater arsenic problems include the Bengal Basin (Bangladesh, India), Mekong Valley (Cambodia, Laos, Vietnam), Yellow River Plain (China), Terai (Nepal), Red River delta (Vietnam), Chianan and Lanyang Plains (Taiwan) and Great Hungarian Plain (Hungary and Romania). Those from inland and closed basins include the Chaco-Pampean Plain (Argentina), and other plains of Chile, Mexico and south-western USA (Fig. 12.2). In many of these areas, significant numbers of wells have arsenic concentrations of several hundreds of micrograms per litre, with occasional sources in the milligram-per-litre range. Arsenic-related health problems have been recognised in the resident populations in numerous areas. Reconnaissance surveys of groundwater quality in other areas such as parts of Burma and Pakistan have also revealed small numbers of wells with concentrations of arsenic above drinking-water limits. Documentation of the affected aquifers of Burma is so far limited. The regions of the world having major aquifers with recognised arsenic problems are outlined below and categorised in terms of their environmental conditions. Distributions of arsenic in the environment related to geothermal activity as well as mining and mineralisation are also described.

### 12.3.3 Alluvial Plains and Deltas

#### 12.3.3.1 Bengal Basin (Bangladesh, India)

The change from use of surface water or dug-well water to groundwater accessed by hand-pumped tubewells began in India and Bangladesh during the 1960s and accelerated significantly from the 1980s onwards. This led to a vast increase in the access of rural populations to what was considered a superior and safe source of drinking water from the readily available groundwater resources contained in the shallow aquifers. The discovery in 1983 in West Bengal (Garai et al. 1984) and in 1993 in Bangladesh (BGS and DPHE 2001) of problems with arsenic in the groundwater changed the situation significantly. Access of rural populations to improved supplies of drinking-water was down from the 90% coverage in the early 1990s to 68% in 2004 (Rosenboom 2004). The discoveries led to the instigation of major programmes of well testing and resource assessment, health surveillance, water-supply mitigation, and research.

Concentrations of arsenic in the groundwaters of the Bengal Basin span some four orders of magnitude, with occasional extremes above 1 mg L<sup>-1</sup>. Recognised health



**Fig. 12.2** Occurrence of documented arsenic problems in groundwater (arsenic  $>50 \mu\text{g L}^{-1}$ ) in major aquifers and environmental problems related to mining and geothermal sources

problems include skin disorders and internal cancers. The high arsenic concentrations occur in groundwaters within unconsolidated grey micaceous alluvial and deltaic Holocene sands and silts associated with the Ganges, Brahmaputra and Meghna river systems. The presence of poorly-permeable surficial overbank deposits (BGS and DPHE 2001; Chakraborti et al. 2001; McArthur et al. 2004; van Geen et al. 2008), particularly in the lower sections of the basin, together with both solid organic matter and DOC in the groundwater, has resulted in development of strongly reducing aquifer conditions, aided by microbial reactions. Groundwater compositions are typified by high concentrations of dissolved Fe, Mn,  $\text{HCO}_3^-$ ,  $\text{NH}_4^+$ , P and DOC as well as high As, and low concentrations of  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$ , though the correlations between these parameters are often poor. Methane has also been detected in some groundwaters (Ahmed et al. 1998). Many studies have observed that dissolved arsenic concentrations peak in the typical depth range 15–50 m (BGS and DPHE 2001; Harvey et al. 2002; Klump et al. 2006; McArthur et al. 2004; Zheng et al. 2005), although high concentrations ( $>50 \mu\text{g L}^{-1}$ ) are observed at depths up to about 150 m below surface.


A national survey of arsenic in groundwater (BGS and DPHE 2001), using some 3,500 groundwater samples, found that 27% of samples from the Holocene shallow aquifer ( $<150$  m depth) contained arsenic at concentrations exceeding  $50 \mu\text{g L}^{-1}$ , and 46% exceeded  $10 \mu\text{g L}^{-1}$  (Fig. 12.3). Subsequent arsenic analyses from the Bangladesh Arsenic

Mitigation and Water Supply Program (BAMWSP 2005) showed that of almost five million tube wells tested nationally by the mid 2000s, some 30% had arsenic concentrations above  $50 \mu\text{g L}^{-1}$ . The results from these two datasets compare reasonably well. Each demonstrates a very variable distribution of arsenic regionally across Bangladesh, with the greatest proportion of exceedances occurring in groundwaters from the southern and south-eastern parts of the country.

A UNICEF/DPHE survey of groundwaters from southern Bangladesh (Rosenboom 2004), found of some 317,000 tubewells tested, 66% contained arsenic with concentrations above  $50 \mu\text{g L}^{-1}$ ; 10% had  $<10 \mu\text{g L}^{-1}$ . Van Geen et al. (2003b) also found, from a survey of 6,000 tube wells in Araihasar, south-central Bangladesh, that some 75% of tube wells (depth range 15–30 m) had arsenic concentrations above  $50 \mu\text{g L}^{-1}$ . The higher percentage exceedances of these studies reflect their location in some of the worst-affected areas.

The BGS and DPHE (2001) study estimated, on the basis of the population at the time (125 million estimate), that up to 35 million people were drinking groundwater with arsenic concentrations above  $50 \mu\text{g L}^{-1}$ , and 57 million were drinking water with concentrations above  $10 \mu\text{g L}^{-1}$ . This represents by far the most severely affected population recognised globally. Figures will have been modified somewhat by subsequent mitigation efforts although this is to be offset by population growth (161 million 2012 estimate).





**Fig. 12.3** Smoothed contour map of arsenic concentrations in shallow groundwater (<math><150\text{ m}</math>) from Bangladesh and histograms of the distributions of arsenic in three special study areas (BGS and DPHE

2001). Low concentrations indicated in blue in northern Bangladesh denote the locations of the Tista Fan, and Madhupur and Barind Tracts

In India, the first evidence of arsenic-related health problems emerged in West Bengal (Bhattacharya et al. 1997; Chowdhury et al. 1997; Das et al. 1994), where it is estimated that about 6.5 million people are drinking water with arsenic concentrations greater than  $50\ \mu\text{g L}^{-1}$  (Mukherjee et al. 2006). High-arsenic groundwaters have subsequently also been identified in alluvial aquifers in the states of Bihar, Tripura, Uttar Pradesh, Jharkhand and Assam (Chakraborti et al. 2003; Mukherjee et al. 2006). Sedimentary aquifers potentially at

risk from high arsenic concentrations in groundwater also occur in the neighbouring states of Meghalaya and Mizoram. In 2003, the Indian Bureau of Standards revised the national limit for arsenic in drinking water from  $50$  to  $10\ \mu\text{g L}^{-1}$ , in recognition of the severity of the problem and response to changing international regulations.

High-arsenic groundwaters have been found in up to nine districts of West Bengal, the five worst affected being Malda, Murshidabad, Nadia, 24 North Parganas and 24

South Parganas (Chakraborti et al. 2004; McArthur et al. 2004; Rahman et al. 2001). These cover up to 38 000 km<sup>2</sup>. High arsenic concentrations occur typically in the depth range 10–80 m (McArthur et al. 2004). Chakraborti et al. (2001) found from 90,000 water analyses in the Holocene aquifers of West Bengal that some 34% exceeded 50 µg L<sup>-1</sup> and 55% exceeded 10 µg L<sup>-1</sup>.

In both Bangladesh and West Bengal, groundwaters from older Pleistocene aquifers usually have low arsenic concentrations (<10 µg L<sup>-1</sup>). These sediments, dominantly orange-brown and more oxic than the overlying Holocene deposits, occur at variable depths. BGS and DPHE (2001) distinguished the aquifers using a 150-m depth cut-off, although van Geen et al. (2003b) documented Pleistocene deposits in the Araihasar area of Bangladesh at depths as shallow as 30 m (range 30–120 m). Pleistocene deposits also occur at surface within the distinct structural units of the Madhupur and Barind Tracts of Bangladesh (BGS and DPHE 2001) and give rise to the low mapped arsenic concentrations in shallow groundwaters there (Fig. 12.3).

Dug wells also usually have low arsenic concentrations, and offer a potential alternative supply of drinking water. However, Rosenboom (2004) found that 11% of shallow dug wells tested in the UNICEF/DPHE survey had concentrations above 50 µg L<sup>-1</sup>. Dug wells are also more prone to bacterial contamination and sustainability problems during the dry season.

Arsenic in the Holocene aquifer sediments typically has concentrations in the range <1–20 mg/kg (BGS and DPHE 2001; Chakraborti et al. 2001; Swartz et al. 2004) which are not unusual by world standards. The arsenic has been found or postulated to occur in association with Fe(III)- or mixed Fe(II)–Fe(III) oxides (BGS and DPHE 2001; Harvey et al. 2002; Horneman et al. 2004; McArthur et al. 2001; Swartz et al. 2004), phyllosilicate minerals (Breit et al. 2001) and sulfide minerals (Nickson et al. 2000; Polizzotto et al. 2006). Most studies concur that iron oxides are an important control on arsenic mobilisation in the Bengal Basin groundwaters, although opinions are divided on the precise mechanisms involved. Competition for adsorption sites between dissolved arsenic species and other constituents such as phosphate is also likely to exert a control. Paucity of sulphate in the groundwaters over large parts of the region appears contributory to the limited sequestration of arsenic by iron sulphide under the strongly reducing conditions.

Redox changes in the aquifers involve reduction of As(V) to As(III) with resultant changes in sorption behaviour. Reductive dissolution of iron oxide and reductive desorption of arsenic are processes likely to result in increased solute arsenic concentrations, exacerbated by competition for binding sites between arsenic and especially phosphate. Such reactions appear to be only part of a complex sequence of diagenetic transformations in the sediments arising from the

onset of reducing conditions. Reactions involve changes in iron oxide structure, redox status and surface properties over time, with transformation to phases such as magnetite as well as precipitation of siderite and vivianite having been invoked (e.g. BGS and DPHE 2001; Horneman et al. 2004; Swartz et al. 2004). Microbial mediation of the redox reactions and arsenic cycling has been demonstrated (Islam et al. 2004).

Several studies have found much higher ratios of Fe(II)/FeT in the Holocene Bengal sediments compared to the underlying Pleistocene deposits (Horneman et al. 2004; Polizzotto et al. 2006; van Geen et al. 2006). Iron oxides in the latter appear dominated by goethite and haematite and explain the prominent colour change from grey to orange-brown in passing from the Holocene to the Pleistocene sediments (e.g. van Geen et al. 2003b). Holocene sediments have also been found to contain much higher concentrations of phosphate-extractable (van Geen et al. 2006; Zheng et al. 2005) and oxalate-extractable (BGS and DPHE 2001) arsenic and iron than Pleistocene aquifer sediments, which point to an increased proportion of labile arsenic in Holocene deposits. Nonetheless, some studies have concluded that conditions in the Holocene aquifers are too strongly reducing to be consistent with in-situ Fe(III) reduction to Fe(II) and propose that arsenic is released instead from iron oxides in surficial soils and sediments and transported to depth as a plume by groundwater flow (Polizzotto et al. 2005, 2006).

The nature of organic matter involved in the redox cycling is also a matter of debate. Peat deposits (McArthur et al. 2001) and peaty strata marginal to peat basins (McArthur et al. 2004) have been invoked as important sources of organic matter to drive the redox reactions. Others cite the importance of disseminated organic matter in the sediments and dissolved in water (BGS and DPHE 2001). In these cases, rapid rates of sedimentation accommodate the organic carbon and can prevent its rapid oxidation. Others have concluded that introduction of anthropogenic carbon through drawdown induced by groundwater pumping has been responsible for catalysing the reduction reactions within the aquifers (Harvey et al. 2002, 2006), although this conclusion has not been universally accepted (e.g. Klump et al. 2006; van Geen et al. 2003a). In West Bengal, the lack of detectable thermotolerant coliforms in high-arsenic groundwaters has also been used as an argument against anthropogenic organic matter being the driver for iron reduction and arsenic release at depth (McArthur et al. 2004).

Groundwater flow and flushing has doubtless had an impact on arsenic mobilisation and distributions, both under natural conditions and by modern groundwater pumping (Michael and Voss 2008). Natural hydraulic gradients in the aquifers are low and groundwater flow strongly controlled by local permeability variations. Groundwater arsenic concentrations show a relationship with surface soil and sediment texture, which suggests a larger degree of aquifer recharge, and flushing, in areas with higher surface permeability (van Geen et al. 2006).

It is also likely that longer-term groundwater flow throughout the Quaternary has resulted in flushing of Pleistocene relative to Holocene deposits, especially since palaeo-head gradients would have been greater during pre-Holocene glacial lowstand periods (BGS and DPHE 2001). However, major abstractions of groundwater over the last few decades, particularly that for irrigation use, have changed flow patterns significantly. Many recent studies have shown that shallow groundwaters (<30 m or so) in the Bengal Basin often have short residence times, on the scale of years to decades (e.g. Klump et al. 2006), likely induced by pumping. Pumping-induced flow and mixing have the potential to either increase or decrease solute arsenic concentrations with time.

In recent years, many studies have focussed on the sustainability of the Pleistocene low-arsenic aquifer in respect of potential drawdown of contaminated water following increased abstraction. Modelling has concluded that the groundwater arsenic concentrations in the deep aquifer are likely to remain low provided abstraction is limited to domestic use by hand-pumps, partly because of the comparatively small volumes involved and partly due to the instigation of a hydraulic barrier in the shallow aquifer by irrigation pumping from there (Burgess et al. 2010; Michael and Voss 2008).

### 12.3.3.2 Nepal

Groundwater is abundant in the Quaternary alluvial sediments of the lowland Terai region of southern Nepal and is an important resource for domestic and agricultural use. Groundwater from tubewells supplies some 11 million people (Chitrakar and Neku 2001). Both shallow and deep aquifers occur throughout most of the Terai region, the deep aquifer being artesian. Quaternary alluvium also infills several intermontaine basins in Nepal, most notably that of the Kathmandu Valley of central Nepal.

A number of surveys of groundwater quality in the Terai region have revealed the presence of arsenic in samples from shallow tubewells (<50 m depth) (Gurung et al. 2005) and some 500,000 people are believed to be exposed to drinking water with arsenic concentrations greater than  $50 \mu\text{g L}^{-1}$  (Pokhrel et al. 2009). Arsenicosis problems have been identified in some areas (Maharjan et al. 2007). By 2004, some 245,000 tubewells had been tested for arsenic, of which 3% were found to exceed  $50 \mu\text{g L}^{-1}$ , the highest observed concentration being  $400 \mu\text{g L}^{-1}$  (Tandukar et al. 2005). From testing in 17 of the 20 Terai districts, the Nepal Red Cross Society (NRCS), also found 3% of groundwater sources sampled having concentrations above  $50 \mu\text{g L}^{-1}$ , the highest observed concentration being  $205 \mu\text{g L}^{-1}$ . The worst-affected districts are Rautahat, Nawalparasi, Parsa and Bara in the central area of the Terai. Other surveys have revealed the occurrence of high-arsenic groundwaters

in the River Bagmati area. Concentrations in the Kathmandu Valley appear mostly low (Gurung et al. 2007; Pant 2011).

The high arsenic concentrations in the Terai occur in anaerobic groundwaters with characteristics similar to those found in the Holocene aquifers of the Bengal Basin (high Fe, Mn,  $\text{NH}_4$ , DOC and low  $\text{SO}_4$ ,  $\text{NO}_3$ ). Dissolved arsenic is dominated by As(III) (Bhattacharya et al. 2003; Tandukar et al. 2005). Deeper tubewells in the Terai appear to have lower arsenic concentrations (generally  $<10 \mu\text{g L}^{-1}$ ).

### 12.3.3.3 Taiwan

Health problems due to arsenic in groundwater were first reported in Taiwan between 1910 and 1920 (Chen and Liu 2007) although awareness of the link with drinking water was delayed until the 1960s (Tseng et al. 1968). Blackfoot disease (BFD, a peripheral vascular disorder with similarities to gangrene) is a well-publicised health problem of the region and is most likely linked to past occurrence of arsenic in the drinking water, although the high humic-acid concentration in groundwaters of the region has been cited as an additional causal factor (Lu et al. 1990). A number of other diseases, including internal cancers, have also been described. High groundwater arsenic concentrations have been recognised in both the south-west Chianan Plain (Chen and Liu 2007; Tseng et al. 1968) and north-east Lanyang Plain (Hsu et al. 1997; Lewis et al. 2007), although BFD occurrence appears restricted to the south-west (Lewis et al. 2007).

Groundwater samples from Chianan Plain have arsenic concentrations in the range  $10\text{--}1,800 \mu\text{g L}^{-1}$  with a significant number being more than  $400 \mu\text{g L}^{-1}$  (Chen and Liu 2007). These occur in groundwaters from deep artesian wells (mostly 80–250 m) abstracted from late Pleistocene alluvial and deltaic sediments. The groundwaters are strongly reducing with evidence of dissolved sulphide and  $\text{CH}_4$  in some samples. Groundwaters from the top 50–80 m are also reducing but are saline with  $\text{SO}_4$  concentrations of  $2,700\text{--}6,000 \text{mg L}^{-1}$ , associated with marine deposition and subsequent evaporation (Chen and Liu 2007). Groundwaters abstracted from Lanyang are also artesian, but of shallower depth (typical range 16–40 m) (Hsu et al. 1997). Here, arsenic concentrations reach up to  $600 \mu\text{g L}^{-1}$  (Hsu et al. 1997). In each area, the groundwaters are strongly reducing, dissolved As being dominated by As(III) (Chen et al. 1994). Groundwater samples taken from shallow open dug wells have low arsenic concentrations (Guo et al. 1994).

### 12.3.3.4 Cambodia, Laos and Vietnam

The discovery of high concentrations of arsenic in groundwaters from Cambodia, Laos and Vietnam is, perhaps surprisingly, relatively recent. Development of groundwater from tubewells in these countries has increased significantly

since the 1990s (Berg et al. 2006; Chanpiwat et al. 2011; Polyá et al. 2005) and this recent change is perhaps responsible for the delayed recognition. High arsenic concentrations are found in groundwater from Holocene sediments. These are deposited by the Mekong River in all three countries, the Red River in Vietnam and the Bassac River (Buschmann et al. 2007) and the Tonle Sap (Berg et al. 2007) in Cambodia.

In Cambodia, 10 affected provinces have been identified but the most severely impacted is Kandal Province where extremes up to  $3,500 \mu\text{g L}^{-1}$  have been reported (Berg et al. 2007; Buschmann et al. 2007; Sampson et al. 2008). Berg et al. (2006) found 20% of samples analysed exceeding  $50 \mu\text{g L}^{-1}$ . The groundwaters occur mainly in groundwater from wells  $>16$  m deep (Polyá et al. 2005). In Laos, more limited data are available but concentrations along the Mekong Valley have been reported in the range  $<0.5$ – $278 \mu\text{g L}^{-1}$  (Chanpiwat et al. 2011).

In Vietnam, aquifers in both the Mekong delta and Red River delta are exploited for water supply and the total number of tubewells in use could be of the order of one million (Berg et al. 2007). Aquifers exploited are of both Holocene and Pleistocene age. In the Vietnamese Mekong delta, the shallow aquifer is affected by salinity and is not used for drinking water. Groundwater for irrigation and drinking is therefore abstracted from depths of 150–250 m from Neogene deposits. Holocene aquifers in the northern Vietnamese part of the Mekong delta, upstream of the zone affected by saline intrusion, have arsenic concentrations in the range  $<1$ – $845 \mu\text{g L}^{-1}$  (Berg et al. 2007).

In the Red River delta, concentrations up to  $3,000 \mu\text{g L}^{-1}$  occur (Berg et al. 2007; Nguyen et al. 2009) in an area which includes the city of Hanoi, particularly the southern part. Holocene sediments up to about 30 m thick form the shallow aquifer, below which, Pleistocene sands and gravels occur. These are separated in places by a clay layer several metres thick, although ‘windows’ exist which allow hydraulic continuity between the two. Holocene sand deposits are largely capped by surficial confining overbank clays (typically some 2–10 m thick) (Postma et al. 2007). Both Holocene and Pleistocene aquifers are used for water supply. It is estimated that around one million people are drinking water in the delta area with an As concentration greater than  $50 \mu\text{g L}^{-1}$  (Winkel et al. 2011).

Arsenic-affected groundwaters in these countries are reducing (with high concentrations of Fe, Mn and DOC; dissolved As overwhelmingly dominated by As(III)) (Chanpiwat et al. 2011; Rowland et al. 2007). In Vietnam, high concentrations of  $\text{NH}_4$  and  $\text{CH}_4$  are also found in some areas (Berg et al. 2007; Postma et al. 2007). As elsewhere in Asia, arsenic mobilisation in the Holocene aquifers has been linked to reductive dissolution of iron oxides in the sediments. In the Red River delta, these have been taken to

be relatively crystalline oxide phases such as haematite or goethite (Postma et al. 2007). As in Bangladesh however, there are proponents of the origin of the arsenic lying in surface soils and sediments with subsequent transport to the Holocene aquifer (Polizzotto et al. 2008). In an extensive survey of groundwater in the Red River delta, Winkel et al. (2011) have documented the drawdown of As from the Holocene to the Pleistocene aquifer as a result of long-term heavy abstraction of groundwater.

To date, limited numbers of arsenicosis patients have been recognized in Cambodia, probably due to relatively recent introduction of tubewell water as a source of drinking water (Berg et al. 2007). However, the first cases were diagnosed (Kandal Province) in 2006 (Sampson et al. 2008). Few cases have been reported in Vietnam and to the best of our knowledge no arsenic-health problems have been documented in Laos.

### 12.3.3.5 Northern China

Arsenic (concentrations in excess of the Chinese national standard of  $50 \mu\text{g L}^{-1}$ ) has been found in groundwaters from a number of areas in northern China, including Xinjiang, Shanxi, Jilin and Liaoning Provinces as well as Inner Mongolia (Fig. 12.2) (Bian et al. 2012; Deng et al. 2009; Ning et al. 2007; Smedley et al. 2003; Xie et al. 2009). The earliest problems were recognised in Xinjiang Province where concentrations of  $40$ – $750 \mu\text{g L}^{-1}$  were found in deep artesian groundwater (wells to 660 m deep) from the Quaternary Dzungaria (Junggar) Basin (Wang and Huang 1994). Under artesian conditions, As concentrations increased with depth; shallow (non-artesian) groundwaters had concentrations between  $<10$  and  $68 \mu\text{g L}^{-1}$ .

In the Datong and Jinzhong Basins of Shanxi Province, arsenic concentrations up to  $4,400 \mu\text{g L}^{-1}$  have been found (Guo et al. 2003; Xie et al. 2008, 2009). Datong Basin groundwaters have highest observed concentrations in the depth range 15–60 m (Xie et al. 2008), present under usually strongly reducing conditions (Xie et al. 2009). Groundwaters from the Hetao Plain of Inner Mongolia also have high concentrations, up to  $1,350 \mu\text{g L}^{-1}$  (Guo et al. 2001), being highest in boreholes with a depth range of 15–30 m (open dug wells of 3–5 m depth had low concentrations).

In the Huhhot Basin, part of the Tumet Plain of Inner Mongolia, arsenic also occurs in strongly reducing groundwaters from mainly Holocene alluvial and lacustrine aquifers. Problems are worst in the lowest-lying parts of the basin (Smedley et al. 2003). These groundwaters also have characteristically high concentrations of dissolved Fe, Mn, DOC,  $\text{HCO}_3$  and  $\text{NH}_4$  and low concentrations of nitrate and sulphate (Luo et al. 1997; Smedley et al. 2003). The arsenic, at concentrations up to  $1,500 \mu\text{g L}^{-1}$ , is again dominated by As(III). Unlike the Bengal Basin, deep pre-Holocene aquifers also have relatively high, though less extreme,

arsenic concentrations (up to  $300 \mu\text{g L}^{-1}$ ). Many of these waters have associated high concentrations of humic acid which may have an impact on arsenic mobilisation, as well as health effects. Unusually, groundwater from several hand-dug wells in the low-lying part of the Huhhot Basin also has relatively high arsenic concentrations, up to  $550 \mu\text{g L}^{-1}$ . This may owe to the maintenance of relatively reducing conditions at shallow depths as a result of high organic content (solid and dissolved) (Smedley et al. 2003).

Documented health problems derive especially from Xinjiang Province and Inner Mongolia. The problems are manifested by skin lesions, including melanosis and hyperkeratosis, vascular disorders and lung, skin and bladder cancer.

### 12.3.3.6 Hungary and Romania

Concentrations of arsenic above  $50 \mu\text{g L}^{-1}$  have been identified in groundwaters from alluvial sediments in the southern part of the Great Hungarian Plain (part of the Pannonian Basin) of Hungary and neighbouring parts of Romania (Fig. 12.2). Concentrations up to  $150 \mu\text{g L}^{-1}$  (average  $32 \mu\text{g L}^{-1}$ ) have been found by Varsányi et al. (1991). The Plain, some 110,000  $\text{km}^2$  in area, consists of a thick sequence of subsiding Quaternary sediments. Groundwaters vary from Ca-Mg- $\text{HCO}_3$ -type in the recharge areas of the basin margins to Na- $\text{HCO}_3$ -type in the low-lying discharge regions. Groundwaters in deep parts of the basin (80–560 m depth) with high arsenic concentrations are reducing and many have reported high concentrations of dissolved organic matter (humic acid quoted as up to  $20 \text{mg L}^{-1}$ ) (Varsányi et al. 1991). The groundwaters have largest arsenic concentrations in the lowest parts of the basin, where the sediment is fine-grained. Gurzau and Gurzau (2001) reported concentrations up to  $176 \mu\text{g L}^{-1}$  in the associated aquifers of neighbouring Romania.

## 12.3.4 Inland Basins in Arid and Semi-arid Areas

### 12.3.4.1 Mexico

In terms of arsenic occurrence, the Principal aquifer of Comarca Lagunera in the states of Coahuila and Durango is the best-documented region in Mexico. The population exposed to arsenic in drinking water with  $>50 \mu\text{g L}^{-1}$  has been estimated at around 400,000 (Del Razo et al. 1990). Health problems related to arsenic were first recognized there in the early 1960s and symptoms include melanosis, keratosis and skin cancer. Groundwater from a large number of wells has been found with arsenic concentrations well in excess of the national standard for drinking water (currently  $25 \mu\text{g L}^{-1}$ ). Concentrations have been found in the range  $3$ – $740 \mu\text{g L}^{-1}$ , around half being above  $50 \mu\text{g L}^{-1}$  (Del Razo et al. 1990, 1994; IMTA 1992; Rosas et al. 1999). Highest

concentrations are found in the San Pedro, Tlahualilo and Francisco I. Madero areas (Fig. 12.4). The region has been subject to heavy groundwater abstraction and water levels have fallen significantly over the last century. The water table lies more than 100 m below surface in the centre of the depression.

The groundwaters of Comarca Lagunera are predominantly oxidising with neutral to high pH. Del Razo et al. (1990) quoted pH values for groundwaters in the range 6.3–8.9. They also noted that most ( $>90\%$ ) of the groundwater samples investigated had arsenic present predominantly as As(V). The groundwater also has high concentrations of fluoride (up to  $3.7 \text{mg L}^{-1}$ ) (Cebrián et al. 1994). The high arsenic concentrations occur in the low-lying parts of the basin which act as zones of groundwater discharge. Being an internal drainage basin the discharge area contains groundwater with significant residence times; radiocarbon dating and supporting stable-isotopic measurements reveal groundwater model ages up to 30,000 years in the low-lying areas (Brouste et al. 1997) (Fig. 12.4). Convergent flow within the discharge zones likely facilitates accumulation of solutes such as arsenic.

Arsenic problems have also been identified in groundwaters from the states of Sonora and Chihuahua in northern Mexico. Wyatt et al. (1998) found concentrations in Sonora in the range  $2$ – $305 \mu\text{g L}^{-1}$  and observed similar correlations with fluoride, which reached up to  $7.4 \text{mg L}^{-1}$ . In Chihuahua, arsenic concentrations reached  $650 \mu\text{g L}^{-1}$  in saline and F-rich groundwater from wells up to 250 m deep (Junta Municipal 2000).


### 12.3.4.2 Chile

Health problems related to arsenic in drinking water were first recognised in northern Chile in 1962. Symptoms included skin-pigmentation changes, keratosis, skin cancer, cardiovascular problems and respiratory disease. More recently, chronic arsenic ingestion has been linked to lung and bladder cancer. Arsenic exposure dated back from the period 1955–1970. Both surface waters and groundwaters from Administrative Region II (incorporating the cities of Antofagasta, Calama and Tocopilla) of northern Chile (Cáceres et al. 1992) have high concentrations. The region is arid, being part of the Atacama Desert, and water resources are limited. High As concentrations are accompanied by high salinity and concentrations of boron and lithium. This in part relates to evaporation but is also significantly affected by geothermal inputs from the El Tatio geothermal field. Arsenic concentrations below  $100 \mu\text{g L}^{-1}$  in surface waters and groundwaters are apparently quite rare, and concentrations up to  $21,000 \mu\text{g L}^{-1}$  have been found. Karcher et al. (1999) quoted ranges of  $100$ – $1,000 \mu\text{g L}^{-1}$  in untreated surface waters and groundwaters (average  $440 \mu\text{g L}^{-1}$ ).


The affected groundwaters of Chile are predominantly oxidising with As(V) the dominant species. The aquifers are composed of volcanic rocks and volcanogenic sediments, but the arsenic sources are not well-characterised. In Antofagasta, concentrations of arsenic in sediments are ca.  $3.2 \text{ mg kg}^{-1}$  (Cáceres et al. 1992). Sediments from the Rio Loa and its tributaries have much higher concentrations ( $26\text{--}2,000 \text{ mg kg}^{-1}$ ) as a result of geothermal inputs from the river system (Romero et al. 2003). Additional exposure to arsenic from the smelting of copper ore has also been noted in northern Chile (Cáceres et al. 1992).

### 12.3.4.3 Argentina

The Chaco-Pampean Plain of central Argentina constitutes perhaps one of the largest regions of high-arsenic groundwaters known, covering around one million  $\text{km}^2$ . High concentrations of arsenic have been found in parts of the provinces of Córdoba, La Pampa, Santa Fe, Buenos Aires, Tucumán, San Luis, Chaco and Santiago Del Estero. An estimated two million people live in these As-affected areas (Castro de Esparza 2009). Symptoms typical of chronic arsenic poisoning, including skin lesions and some internal cancers, have been recorded (Bates et al. 2004;



**Fig. 12.4** (a) Distributions of arsenic in the groundwaters of Comarca Lagunera, Mexico (Data from IMTA 1992). (b) Groundwater bodies distinguished by residence time (Data from Brouste et al. 1997)




**Fig. 12.4** (continued)

Hopenhayn-Rich et al. 1996). The high-arsenic groundwaters are mostly from Quaternary deposits of silt-grade loess, with intermixed rhyolitic or dacitic volcanic ash (Nicolli et al. 1989, 2010; Smedley et al. 2002). The sediments display abundant evidence of post-depositional diagenetic changes under semi-arid climatic conditions, with common occurrences of palaeosols and calcrete.

A large range of As concentrations is observed in the groundwaters, with many samples exceeding  $1,000 \mu\text{g L}^{-1}$ . Nicolli et al. (1989) reported concentrations in groundwaters from Córdoba in the range  $6\text{--}11,500 \mu\text{g L}^{-1}$  (median  $255 \mu\text{g L}^{-1}$ ). Smedley et al. (2002) found concentrations in La Pampa Province in the range  $<4\text{--}5,280 \mu\text{g L}^{-1}$  (median  $145 \mu\text{g L}^{-1}$ ). Nicolli et al. (2010) found concentrations in

Tucumán Province of  $11\text{--}1,660 \mu\text{g L}^{-1}$  (median  $46 \mu\text{g L}^{-1}$ ). Blanes et al. (2011) reported concentrations up to  $1,070 \mu\text{g L}^{-1}$  in groundwater from Chaco Province, Bhattacharya et al. (2006) found concentrations of  $7\text{--}15,000 \mu\text{g L}^{-1}$  in Santiago Del Estero and Farías et al. (2003) found concentrations of  $10\text{--}593 \mu\text{g L}^{-1}$  in groundwater from four Pampean provinces.

The groundwaters from the Chaco-Pampean loess plain have a large salinity range, with often high pH and alkalinity and are predominantly oxidising. Smedley et al. (2002) found pH values of 7.0–8.7 in groundwaters from La Pampa; Nicolli et al. (2010) found values of 6.3–9.2 in Tucumán. The high-As Pampean groundwaters are usually of Na-HCO<sub>3</sub> type. Under the overwhelmingly oxic conditions of the groundwaters, As(V) usually dominates



**Fig. 12.5** Variations in total arsenic concentration with pH, alkalinity, fluoride, vanadium, boron, molybdenum and uranium in groundwater from La Pampa and Tucumán provinces, Argentina (Data from Nicolli et al. 2010; Smedley et al. 2002; Warren 2001)

(Bhattacharya et al. 2006; Smedley et al. 2002). Generation of the Na-HCO<sub>3</sub> high-pH waters is achieved through silicate hydrolysis (e.g. albite weathering), carbonate reaction and associated ion exchange.

Arsenic mobility in the Chaco-Pampean groundwaters is likely controlled to a large extent by pH-dependent sorption to metal oxides in the sediments. Positive correlations between dissolved As and pH in the groundwaters (Fig. 12.5) are consistent with weaker sorption of As(V) to iron oxide surfaces at high pH (Bhattacharya et al. 2005; Smedley et al. 2002). Presence of Mn(IV) oxides, along with low concentrations of solid organic carbon (Sracek et al. 2009) are likely contributory in maintaining oxic conditions through the aquifer.

A unifying feature of the Chaco-Pampean loess sediments is the presence of rhyolitic volcanic ash either intermixed with the sediments or in thin discrete layers. The components of the ash may constitute the ultimate sources of the dissolved As, though linkages are lost through subsequent weathering and redistribution. Dissolved As is also generally well-correlated with other anion and oxyanion elements (F, V, HCO<sub>3</sub>, Mo and B; Fig. 12.5), some of these

elements having very high concentrations. Volcanic ash may also be the ultimate source of these species (Nicolli et al. 1989).

#### 12.3.4.4 Nicaragua

Alluvial aquifers of the Sébaco Valley of western Nicaragua contain groundwater with occasional high As concentrations. Altmirano Espinoza and Bundschuh (2009) reported concentrations in the range 10–122 μg L<sup>-1</sup> and cited an earlier (2005) UNICEF survey in Jinotega municipality in which concentrations up to 1,200 μg L<sup>-1</sup> were found. In the village of El Zapote, a well drilled in 1994 and tested for As in 1996 had a recorded concentration of 1,320 μg L<sup>-1</sup>. Residents drinking the water displayed skin and respiratory disorders (Gomez et al. 2009) and the well was closed in 1996. The groundwater in the El Zapote well derived from Tertiary volcanic rocks which also form the precursors of the Quaternary alluvial aquifer in the valley. The groundwaters of the Sébaco Valley have pHs in the range 6–8, are oxic, and arsenic is dominantly As(V) (Altmirano Espinoza and Bundschuh 2009). The higher-As groundwaters appear to be of Na-HCO<sub>3</sub> composition.



Morales et al. (2009) also found high As concentrations in groundwater from neighbouring Esteli Department, with observed concentrations up to  $115 \mu\text{g L}^{-1}$ , an apparent positive correlation between As and pH and similar dominance of As(V).

#### 12.3.4.5 Spain

Cenozoic aquifers of the Duero Basin of north-central Spain have recently been found to contain arsenic with concentrations up to  $613 \mu\text{g L}^{-1}$ . Sediments are mixed, carbonate-rich alluvial and lacustrine deposits and water sources within them include springs, shallow wells and boreholes up to 500 m deep (Gomez et al. 2006). Groundwaters are mainly oxidic but with variable redox potentials and often high salinity and  $\text{SO}_4$  concentrations. Groundwater pH ranges from 5.9 to 10.6 and the highest arsenic concentrations generally occur in waters with pH >8. Correlations have been found between arsenic and both vanadium and tungsten and, in shallow wells, also with uranium which reaches concentrations up to  $216 \mu\text{g L}^{-1}$ . Concentrations of boron are also high in some (Gomez et al. 2006).

#### 12.3.4.6 South-West USA

Concentrations of arsenic in most groundwaters of the USA are low. From some 20,000 analyses of groundwater samples taken from public-supply wells across the USA, Focazio et al. (2000) found that 11% exceeded  $10 \mu\text{g L}^{-1}$  and 2% exceeded  $50 \mu\text{g L}^{-1}$  (55% contained  $1 \mu\text{g L}^{-1}$  or less). However, high concentrations have been found in several states, as shown for example by the USGS map of groundwater arsenic occurrence (Ryker 2001). Some of the most widespread and best-documented cases occur in the south-western states of Nevada, California and Arizona. Other aquifers in Maine, Michigan, Minnesota, South Dakota, Oregon, Texas, Oklahoma and Wisconsin have groundwater with concentrations of arsenic exceeding  $10 \mu\text{g L}^{-1}$  (e.g. Ayotte et al. 2006; Focazio et al. 2000; Scanlon et al. 2009; Schreiber et al. 2000; Welch et al. 2000). From the high-arsenic areas, various sources and processes have been cited, including natural dissolution/desorption reactions, geothermal water and mining activity. High concentrations occur under both reducing and oxidising conditions in different areas. Evaporative concentration is thought to be an important process in the more arid areas. Human influences including effects of groundwater pumping have also been cited as causal in arsenic mobilisation (Ayotte et al. 2011).

In Nevada, arsenic concentrations over  $100 \mu\text{g/L}$  have been found in a large number of private groundwater supplies (Fontaine 1994). Concentrations exceeding  $100 \mu\text{g L}^{-1}$  with extremes up to  $2,600 \mu\text{g L}^{-1}$  have also been found in shallow groundwaters from the southern Carson Desert (Welch et al. 1988). These are largely present under reducing conditions.

The groundwaters also have associated high pH (>8) and high concentrations of phosphorus (some  $>4 \text{mg L}^{-1}$ ) and uranium ( $>100 \mu\text{g L}^{-1}$ ), concluded to be largely concentrated by evaporation, combined with the influence of redox and desorption processes involving metal oxides.

In California, a large range of arsenic concentrations ( $<1\text{--}2,600 \mu\text{g L}^{-1}$ ) has been found in groundwater from the Tulare Basin of the San Joaquin Valley (Fujii and Swain 1995). Redox conditions are variable and high arsenic concentrations are found in both oxidising and reducing conditions. The proportion of groundwater arsenic present as As(III) increases with increasing well depth. The groundwaters from the Basin are often strongly affected by evaporation with resulting high concentrations of total dissolved solids. Many also have high concentrations of Se (up to  $1,000 \mu\text{g L}^{-1}$ ), U (up to  $5,400 \mu\text{g L}^{-1}$ ), B (up to  $73,000 \mu\text{g L}^{-1}$ ) and Mo (up to  $15,000 \mu\text{g L}^{-1}$ ) (Fujii and Swain 1995).

In Arizona, Robertson (1989) also noted the occurrence of high arsenic concentrations in some groundwaters under oxidising conditions (dissolved oxygen  $3\text{--}7 \text{mg L}^{-1}$ ) in alluvial aquifers of the Basin and Range Province. Only limited analysis of arsenic species was carried out, but results obtained suggested that the arsenic was present predominantly as As(V). The dissolved arsenic correlated positively with pH, Mo and V as well as Se and F. Groundwater pH values were in the range 6.9–9.3. Arsenic concentrations in the sediments of the Basin and Range Province were in the range  $2\text{--}88 \text{mg kg}^{-1}$ . Oxidising conditions persist in the aquifers down to significant depths (600 m) despite the groundwaters having very long residence times (up to 10,000 years). The high arsenic (and other oxyanion) concentrations are a feature of the closed basins of the province.

### 12.3.5 Sulphide Mineralisation and Mining-Related Arsenic Problems

Arsenic problems have long been recognised in association with zones of ore mineralisation as a result of the high concentrations of arsenic in sulphide minerals. Oxidation of the sulphide minerals, exacerbated by mining activity, potentially leads to the release of arsenic and transition and heavy metals. Oxidation reactions are catalysed by bacteria, notably *Thiobacillus ferrooxidans*, *Thiobacillus thiooxidans* and *Leptospirillum ferrooxidans* (Schrenk et al. 1998). Oxidation releases  $\text{SO}_4$  and associated trace elements and generates acidity which maintains high aqueous concentrations of many released trace metals. One of the highest arsenic concentrations ever reported,  $850,000 \mu\text{g L}^{-1}$ , is from a mine seep at Iron Mountain, California (Nordstrom and Alpers 1999). Plumlee et al. (1999) also found arsenic concentrations in USA mine drainage ranging up to  $340,000 \mu\text{g L}^{-1}$ .

Under oxidising conditions, iron released during pyrite oxidation typically precipitates as an iron(III) oxide. Trace elements likewise coprecipitate with, or adsorb onto, these iron oxides. Water is also neutralised in the presence of carbonate minerals. As a result, many mine waters have near-neutral pH values (e.g. Welch et al. 2000) and many have relatively low iron and arsenic concentrations. The considerable capacity of iron oxides to adsorb arsenic means that even in areas where high dissolved arsenic concentrations occur, they are usually restricted to the local area around the zone of oxidation, typically a few kilometres or less. Nonetheless, contamination of soils, sediments and vegetation in mining and mineralised areas can be substantial.

Most mining-impacted water sources are not used for potable supply, although a few notable exceptions exist and have given rise to localised health problems. Arsenic health problems related to mining activity have been well-documented in Ron Phibun District in Nakhon Si Thammarat Province of southern Thailand. The area lies within the South-East Asian Tin Belt and has been mined for several generations. Arsenic-related health problems were first recognised there in 1987. By the late 1990s, around 1,000 people had been diagnosed with skin disorders, particularly in and close to Ron Phibun town (Williams 1997). Arsenic concentrations up to  $5,000 \mu\text{g L}^{-1}$  have been found in shallow groundwaters from Quaternary alluvial sediments that have been extensively dredged during mining operations. Deeper groundwaters from an older limestone aquifer were apparently less contaminated (Williams et al. 1996), although a few high arsenic concentrations occurred, presumably also as a result of contamination from the mine workings. The mobilisation of arsenic is related to the oxidation of arsenopyrite, exacerbated by the former tin-mining activities. The recent appearance in groundwater has occurred during post-mining groundwater rebound (Williams 1997).

A relatively recent discovery has also been made of arsenic-related health problems in northern Burkina Faso. In rural boreholes close to the town of Ouahigouya, concentrations of As in the range  $<0.5\text{--}1,630 \mu\text{g L}^{-1}$  have been documented following recognition of occurrences of melanosis, keratosis and skin cancer in some village populations. The highest concentrations occurred in boreholes in close proximity to mineralised veins containing arsenopyrite and other sulphide minerals which are hosted within Precambrian (Birimian) metamorphic rocks (Smedley et al. 2007) (Fig. 12.6). The arsenic was overwhelmingly present as As(V), signifying oxidation of the original As-hosting sulphide minerals. Although mining for gold takes place in Burkina Faso, this is chiefly on a small (artisanal) scale and there is no evidence that the mining operations per se have been responsible for the arsenic mobilisation. During


the 1970s, health problems of suspected drinking water origin were also identified in a village close to the town of Mogtedo in central Burkina Faso. High concentrations of As ( $200\text{--}1,600 \mu\text{g L}^{-1}$ ) were found in three borehole samples. The residents were relocated and further investigations of As in Birkinabé groundwater ceased until the renewed interest during the mid 2000s.

Environmental arsenic damage has also been documented in geologically analogous areas of neighbouring Ghana, where gold mining occurs on a much larger scale in the Ashanti Region. To date, there is little evidence of detrimental effects on human health through drinking-water exposure. Gold reserves have long been mined in the region where the gold is also associated with sulphide mineralisation, particularly arsenopyrite. Around the town of Obuasi, high arsenic concentrations have been found in soils close to the mines and treatment works. Some high concentrations were also found in river waters close to the mining activity. However, many of the groundwaters had low arsenic concentrations and the highest observed were not proximal to mining activity but appeared to be natural occurrences under mildly reducing aquifer conditions.

Arsenic contamination from mining activities has been identified in numerous areas of the USA (Welch et al. 2000). Groundwater from some areas has very high arsenic concentrations locally (up to  $48,000 \mu\text{g L}^{-1}$ ). Well-documented cases include the Fairbanks gold-mining district of Alaska, the Coeur d'Alene Pb-Zn-Ag mining area of Idaho, Bunker Hill mine in Idaho, Leviathan Mine in California, Mother Lode in California, Kelly Creek Valley in Nevada, Clark Fork river area in Montana and Lake Oahe in South Dakota.

Elsewhere, documented cases of mining-related arsenic contamination include the Zimapán Valley (Mexico), Baja California (Mexico), Lavrion (Greece), Minas Gerais (Brazil), Styria (Austria), Zloty Stok (Poland), Oruro Province (Bolivia), west Devon (England), eastern Zimbabwe, Korea, Sarawak (Malaysia) (Smedley and Kinniburgh 2002) and Aznalcollar (Spain) (Fig. 12.2). Doubtless, arsenic problems also exist in many other undocumented mining areas.

Increased concentrations of dissolved arsenic have also been found in parts of the world with local mineralisation which has not been mined. In Wisconsin, USA, arsenic and other trace-element problems in groundwater have arisen as a result of the oxidation of sulphide minerals (pyrite and marcasite) present as a discrete secondary cement horizon in the regional Ordovician sandstone aquifer. Concentrations of arsenic up to  $12,000 \mu\text{g L}^{-1}$  have been found in the well waters (Schreiber et al. 2000). The oxidation appears to have been promoted by groundwater abstraction which has led to the lowering of the piezometric surface at a rate of around  $0.6 \text{ m year}^{-1}$  since the 1950s, resulting in a partial



**Fig. 12.6** Map of the distribution of arsenic in groundwater from the Ouahigouya area of Burkina Faso (From Smedley et al. 2007). High concentrations are associated closely with zones of sulphide mineralisation, which are also subject to artisanal gold mining

dewatering of the aquifer. The high arsenic concentrations were observed where the piezometric surface intersects, or lies close to, the sulphide cement horizon. Boyle et al. (1998) also recorded naturally high arsenic concentrations (up to  $580 \mu\text{g L}^{-1}$ ) in groundwaters from an area of sulphide mineralisation in Bowen Island, British Columbia.

### 12.3.6 Geothermal Sources

The common occurrence of high concentrations of arsenic in geothermal fluids has been recognised for a long time. Geothermal areas with documented high concentrations include the USA, Japan, New Zealand, Chile, Argentina, Ecuador,

Kamchatka, France, Italy and Dominica (Fig. 12.2). One of the largest and best-documented geothermal systems is that of Yellowstone National Park, USA where arsenic concentrations up to  $7,800 \mu\text{g L}^{-1}$  have been found in hot springs and geysers (Thompson and Demonge 1996). Geothermal inputs have also given rise to high concentrations of arsenic (up to  $370 \mu\text{g L}^{-1}$ ) in waters of the Madison River, USA (Nimick et al. 1998). Others include Honey Lake Basin, California (arsenic up to  $2,600 \mu\text{g L}^{-1}$ ), Coso Hot Springs, California (up to  $7,500 \mu\text{g L}^{-1}$ ), Imperial Valley, California (up to  $15,000 \mu\text{g L}^{-1}$ ), Long Valley, California (up to  $2,500 \mu\text{g L}^{-1}$ ), Lassen Volcanic National Park, California (up to  $27,000 \mu\text{g L}^{-1}$ ), Steamboat Springs, Nevada (up to  $2,700 \mu\text{g L}^{-1}$ ) and Geyser Bight, Umnak Island, Alaska (up to  $3,800 \mu\text{g L}^{-1}$ ) (Welch et al. 2000). Geothermal inputs from Long Valley, California are believed responsible for the relatively high concentrations ( $20 \mu\text{g L}^{-1}$ ) of arsenic in the Los Angeles Aqueduct which supplies water for the city of Los Angeles (Wilkie and Hering 1998). Geothermal inputs also contribute significantly to the high dissolved arsenic concentrations (up to  $20 \text{mg L}^{-1}$ ) in Mono Lake, California (Maest et al. 1992).

In New Zealand, geothermal waters occur with arsenic concentrations up to  $9,000 \mu\text{g L}^{-1}$  (Webster 2003). River and lake waters receiving inputs from the Wairakei, Ohaaki, Orakei Korako and Atiamuri geothermal fields have concentrations up to  $121 \mu\text{g L}^{-1}$  (Robinson et al. 1995). Concentrations diminish significantly downstream from the geothermal inputs.

Arsenic concentrations in the range  $45,000$ – $50,000 \mu\text{g L}^{-1}$  have been found in geothermal waters from the El Tatio system in the Antofagasta region of Chile (Ellis and Mahon 1977). The geothermal area lies between the volcanoes of the Andes and the Serrania de Tucle. Romero et al. (2003) found concentrations in waters from the Rio Loa and its tributaries in the range  $120$ – $10,000 \mu\text{g L}^{-1}$  as a result of inputs from the El Tatio geothermal system.

In Papallacta Lake, Ecuador, arsenic of geothermal origin has a range between  $1,090$  and  $7,850 \mu\text{g L}^{-1}$ . The lake is fed by the Tambo River which in turn receives inflows from hot springs along its route. Speciation of As in water from two hot springs which were Fe-rich and highly reducing showed a dominance of inorganic As(III) (Cumbal et al. 2009). In contrast, two cool springs with higher redox potentials showed a dominance of As(V). The variations indicate oxidation of discharging geothermal fluids over time and mixing with oxygenated surface water.

Arsenic concentrations in the range  $100$ – $5,900 \mu\text{g L}^{-1}$  have been also been found in geothermal waters from Kamchatka (White et al. 1963) and in the range  $500$ – $4,600 \mu\text{g L}^{-1}$  in geothermal fields from Kyushu, Japan (Yokoyama et al. 1993).

White et al. (1963) also reported arsenic concentrations in geothermal waters from Iceland. While concentrations were high compared to most groundwaters (range  $50$ – $120 \mu\text{g L}^{-1}$ ),

these are much lower than the values found in many other geothermal systems described above. This may be related to the fact that the geothermal fluids in the Icelandic volcanic field are associated with oceanic basaltic magmas, hence having low arsenic concentrations (Webster and Nordstrom 2003). Typical high-arsenic geothermal fluids are associated with acidic volcanic systems in continental settings. In this case, higher arsenic concentrations may be derived from the interaction of geothermal fluids with the continental crust, particularly argillaceous sediments (Nordstrom 2002) in which the element is known to be preferentially partitioned. High arsenic concentrations have not been documented in geothermal systems associated with other volcanic provinces of dominantly basaltic composition, whether oceanic or continental (e.g. Hawaii, East African Rift, Reimann et al. 2003).

Hot springs commonly show a positive correlation between arsenic and chloride (e.g. Welch et al. 1988; Wilkie and Hering 1998). This association was noted by Webster and Nordstrom (2003) to relate to their similar behaviour during sub-surface boiling and phase separation as both elements partition preferentially into the liquid phase.

## 12.4 Mineral-Water Interactions

### 12.4.1 Adsorbed Arsenic in Sediments

Of the arsenic problem aquifers identified, it appears to be those hosted in young sediments that are most vulnerable to the development of high-arsenic groundwater on a regional scale. In these, surface reactions are important controls on the trace-element chemistry of the groundwaters. The major minerals which adsorb arsenic (as both arsenate and arsenite) in sediments are the metal oxides, particularly those of iron, aluminium and manganese. Iron oxides are relatively abundant in most sediments and are commonly produced by the weathering of primary minerals. In freshwater sediments and silicate clays, they often constitute about 50% of the iron present (Manning and Goldberg 1997). In sandy aquifers, the iron oxides are probably the most important adsorbents of arsenic because of their relative abundance, strong binding affinity and high specific surface area, especially the freshly-formed amorphous oxides. These are also particularly sensitive to a changing geochemical environment, acid dissolution and reductive dissolution, as well as to changes in mineral structure and crystallinity. Although the affinity for arsenate adsorption is strong, arsenite has been shown to adsorb more strongly to HFO and goethite than arsenate under alkaline conditions ( $\text{pH} > 7$ ) (Dixit and Hering 2003). Aluminium oxides can be expected to play a significant role in arsenic adsorption when present in quantity (Manning and Goldberg 1997). Iron sulphide minerals ( $\text{FeS}$ ,  $\text{FeS}_2$ ) may also adsorb significant quantities of arsenic and it is likely the limited

supply of  $\text{SO}_4$  in reducing groundwater that restricts the removal of arsenic from solution in many of the high-arsenic aquifers of south-east Asia (O'Day et al. 2004).

The role of carbonates as arsenic sorbents has also been investigated recently. Experiments by So et al. (2008) and Alexandratos et al. (2007) found that arsenate sorbs readily to calcite surfaces, the affinity depending on alkalinity, pH and ionic strength. Experiments by Goldberg and Glaubig (1988) also observed adsorption of As(V) on calcite, and Romero et al. (2004) speculated that calcite retarded arsenate in the carbonate aquifer of Zimapán, Mexico. By contrast, So et al. (2008) found that arsenite did not sorb significantly to calcite. These observations suggest that calcite would have little effect on arsenic mobility in reducing conditions but could be important in carbonate aquifers under oxic conditions.

#### 12.4.2 Reduced Sediments and the Role of Iron Oxides

A recognised sequence of reduction reactions occurs when sediments are buried and the environment becomes anaerobic (Berner 1981). Such reaction sequences are common in sediments from a wide variety of environments (e.g. Postma et al. 2007). The processes causing changes in iron redox chemistry are particularly important since they can directly affect the mobility of arsenic. The reductive dissolution of hydrous iron oxides and/or the release of adsorbed and coprecipitated arsenic are key to this process. The redox sequence begins with the consumption of oxygen and an increase in dissolved  $\text{CO}_2$  from the decomposition of organic matter.  $\text{NO}_3^-$  decreases by reduction to  $\text{NO}_2^-$  and ultimately to the gases  $\text{N}_2\text{O}$  and  $\text{N}_2$ . Insoluble manganese (IV) oxides dissolve by reduction to soluble  $\text{Mn}^{2+}$  and hydrous ferric oxides are reduced to  $\text{Fe}^{2+}$ . Subsequent reactions involve  $\text{SO}_4^{2-}$  reduction to  $\text{S}^{2-}$ , then  $\text{CH}_4$  production from fermentation and methanogenesis, and finally reduction of  $\text{N}_2$  to  $\text{NH}_4^+$ . The position of As(V) reduction in the sequence relative to Fe reduction has been disputed but is expected before  $\text{SO}_4^{2-}$  reduction. During sulphate reduction, the sulphide produced reacts with any available iron to produce FeS and ultimately pyrite,  $\text{FeS}_2$ . Iron is often more abundant than sulphur so that excess iron results beyond that which can be converted to pyrite. In the presence of carbonate or phosphate, dissolved Fe(II) may precipitate as siderite (e.g. Postma et al. 2007) or vivianite (Swartz et al. 2004), each of which has potential to remove As from solution (Islam et al. 2005b).

There is also evidence for solid-state transformations of the iron oxides under reducing conditions. This is reflected by a change from red/orange/brown to grey/green/blue

coloration. Analysis of the Fe(II) and Fe(III) contents of iron oxides from reduced lake and aquifer sediments often indicate the presence of mixed Fe(II)-Fe(III) oxides (Davison 1993; Fendorf et al. 2010). Green rusts have been identified or suspected in anoxic soils and sediments (Cummings et al. 1999; O'Day et al. 2004; Taylor 1980). These consist of a range of green-coloured mixed Fe(II)-Fe(III) hydroxide minerals. Authigenic magnetite ( $\text{Fe}_3\text{O}_4$ ) has also been identified in anaerobic sediments, formed by microbial transformations (Cummings et al. 2000; Fredrickson et al. 1998). Magnetite formation has also been established under reducing conditions in the laboratory (Benner et al. 2002; Dixit and Hering 2003; Guerin and Blakemore 1992; Islam et al. 2005b). Under strongly reducing conditions magnetite is unstable and eventually dissolves. In the presence of high concentrations of  $\text{H}_2\text{S}$ , magnetite converts slowly to pyrite on a scale of centuries or more.

These studies of iron oxides in reducing environments indicate the complexities involved in the sequence of events taking place when Fe(III) oxides are subjected to strongly reducing conditions. The changes are evidently substantial and can result in the partial dissolution of the oxides and their transformation to completely new mineral phases, as well as to changes in their arsenic sorptive capacities. Even quite small changes in arsenic binding could have a large impact on porewater arsenic concentrations because of the large solid/solution ratio found in sediments. Therefore, understanding the changes to the nature of iron oxide minerals in sedimentary environments is an important part of understanding the processes leading to arsenic mobilisation in sedimentary environments.

#### 12.4.3 Role of Microbes

It has become increasingly clear that microbes play an important role in arsenic speciation and mobilisation. They can be significant catalysts in the oxidation of arsenite, respiration of arsenate and methylation and volatilisation reactions (Lloyd and Oremland 2006; Oremland and Stolz 2005; Oremland et al. 2004). The microbiological transformations either provide sources of energy or act as detoxifying mechanisms. A number of chemoautotrophs oxidise As(III) by using oxygen, nitrate or ferric iron as a terminal electron acceptor and  $\text{CO}_2$  as their sole carbon source. Some heterotrophs are also capable of As(III) oxidation using organic carbon. Conversely, many prokaryotes (e.g. *Bacillus selentireducens* and *Bacillus arsenicoselantis*) are capable of As(V) respiration or As(III) oxidation (Oremland et al. 2002; Oremland and Stolz 2003; Pederick et al. 2007). Arsenic can also be released indirectly as a

result of other microbially-induced redox reactions. For example, the dissimilatory Fe-reducing bacterium *Shewanella alga* reduces Fe(III) to Fe(II) in scorodite ( $\text{FeAsO}_4 \cdot 2\text{H}_2\text{O}$ ), releasing As(V) in the process (Cummings et al. 1999). The Fe(III)-reducing bacteria *Geobacter* and *Geothrix* have also been shown to affect arsenic cycling in reduced aquifer sediments (Islam et al. 2005a, b).

Thermophilic bacteria and cyanobacteria have been recognised in geothermal settings and affect arsenic speciation and mineral precipitation in such systems (Webster and Nordstrom 2003). Streams affected by geothermal inputs may also be influenced by microbial activity. In geothermally-fed waters of Hot Creek, California, Wilkie and Hering (1998) concluded that the oxidation of As(III) was bacterially-controlled and that in abiotic systems, this oxidation reaction was much slower. Extremophiles have also adapted to the alkaline and saline conditions in the geothermally-influenced Mono Lake of California (Oremland et al. 2002, 2004).

Organic matter plays a key role in many microbial reactions in sediments and the type of organic matter present affects its reactivity and hence bioavailability. Organic matter has been shown to act both as an energy source for Fe- and As-reducing bacteria and as an electron shuttle enhancing microbial reactions (Islam et al. 2005a; Rowland et al. 2007).

## 12.5 Common Features of High-Arsenic Groundwater Provinces

### 12.5.1 The Sources of Arsenic

In the cases where affected groundwaters are found close to obvious geological or industrial sources rich in arsenic (geothermal springs, drainage from mineralised and mining areas, specific contaminant sources), it is clear that the anomalously high arsenic concentrations in the source region are responsible. The extent of this contamination is usually highly localised because the geochemical conditions within most aquifers do not favour arsenic mobilisation on a regional scale. Areas affected by geothermal activity are potentially more widespread since in this case mobilisation of arsenic is not required: arsenic is already present in solution and the size of geothermal reservoirs can be large. This probably accounts for why high-arsenic surface waters are normally located in geothermal areas. Perhaps more remarkable is the way in which very high concentrations of arsenic, up to several  $\text{mg L}^{-1}$ , are found in groundwaters from areas with unremarkable concentrations of arsenic in the source rocks. In aquifers with extensive areas of high-arsenic groundwater, this appears to be the rule rather than the exception. Most of these cases arise in aquifers derived from relatively young sediments, often consisting

of alluvium or loess where the total sediment arsenic concentrations are typically in the range  $1\text{--}20 \text{ mg kg}^{-1}$  (e.g. BGS and DPHE 2001; Postma et al. 2007; Swartz et al. 2004).

Of critical importance is the fact that drinking-water limits for arsenic are very low in relation to the overall abundance of arsenic in the environment. Fortunately, most arsenic is normally retained in the solid phase and does not constitute a problem for potable water supplies. However, it only takes a very small percentage of this 'solid' arsenic to dissolve or desorb to give rise to a serious groundwater problem.

### 12.5.2 Mobilisation of Arsenic

Two key factors appear to be involved in the formation of high-As groundwaters on a regional scale: firstly, some form of geochemical trigger releases arsenic from the aquifer solid phase into the groundwater. Secondly, the released arsenic remains in the groundwater and must not be flushed away. There are a number of possible geochemical triggers. In mining and mineralised areas, oxidation of sulphide ores is triggered by influxes of oxygen or other oxidising agents. This may follow a lowering of the water table or change in hydrogeological regime or excavation by mining. In most arsenic-affected aquifers, the most important trigger appears to be the desorption or dissolution of arsenic from oxide minerals, particularly iron oxides. An important feature of this process is that the initial adjustment to environmental changes is probably quite rapid since adsorption reactions are surface reactions. The rate-limiting factors are probably those that control the major changes in pH, redox condition and associated water-quality parameters of the aquifer. These are in part related to physical factors such as the rate of diffusion of gases through the sediment, the rate of sedimentation, the extent of microbiological activity and the rates of chemical reactions. Many of these are likely to be rapid on a geological time scale. Dissolution reactions are relatively slow but even oxide dissolution is rapid on a geological time scale and can be observed over the course of weeks or even days in flooded soils (Masscheleyn et al. 1991). A qualification is that if diagenetic changes to the oxide mineral structure take place or if burial of sediment occurs, there could be a slow release of arsenic over a much longer time scale. The rates of release are likely to be most rapid initially and to diminish over time. Once the diagenetic readjustment has taken place and the sediments have equilibrated with their new environment, there should be little further release of arsenic.

The geochemical triggers involved in arsenic mobilisation could arise for a number of reasons. These are discussed further below.

**Table 12.3** Summary of documented high-arsenic groundwater provinces and their typical chemical and hydrogeological characteristics

Groundwater environment/aquifer type	Examples	Typical aquifer conditions	Typical chemical features of high-arsenic groundwaters	Likely mechanisms of arsenic mobilisation
<b>Strongly reducing groundwater</b>	Alluvial/deltaic aquifers of the Bengal Basin, Mekong Valley, northern China, Taiwan, Nepal, Great Hungarian Plain	<b>Young (Quaternary) sediments, often slow groundwater flow, low-lying parts of aquifers; rapidly accumulated sediments</b>	High Fe ( $>1 \text{ mg L}^{-1}$ ), Mn ( $>0.5 \text{ mg L}^{-1}$ ), $\text{NH}_4\text{-N}$ ( $>1 \text{ mg L}^{-1}$ ), high $\text{HCO}_3$ ( $>500 \text{ mg L}^{-1}$ ); low $\text{NO}_3\text{-N}$ ( $<1 \text{ mg L}^{-1}$ ), low $\text{SO}_4$ ( $<5 \text{ mg L}^{-1}$ ). Often high P ( $>0.5 \text{ mg L}^{-1}$ ). Sometimes high concentrations of dissolved organic matter, including humic acid. Arsenic dominated by As(III)	Reductive desorption of As from metal oxides and reductive dissolution of Fe oxides. Competition between arsenic and other anionic species (especially P)
Oxic groundwater, high pH	Inland basins or closed basins (arid and semi-arid areas): Argentina, Mexico, parts of Nicaragua, western USA, Chile, Spain	Young (Quaternary) sediments, slow groundwater flow, low-lying parts of aquifers; often (not always) volcanic components present	pH typically $>8$ , high $\text{HCO}_3$ ( $>500 \text{ mg L}^{-1}$ ), low Fe, Mn, often correspondingly high F, U, B, V, Mo, Se. Some groundwaters have high salinity due to evaporation. Arsenic dominated by As(V)	Desorption of As and other oxyanion-forming elements from metal oxides, especially of Fe and Mn; volcanic ash may be ultimate source for some locations
Geothermally-influenced groundwater	<b>Parts of Kamchatka, Chile, Argentina, Ecuador, western USA, Japan, New Zealand</b>	Any aquifers affected by geothermal inputs, especially in rift zones	High Si, B, Li, often high salinity (Na, Cl); high pH ( $>7$ ); increased groundwater temperature; As speciation variable	Mixing of fresh groundwater with geothermal solutions
Groundwater from sulphide mineralised/mining areas	Parts of Burkina Faso, Canada, USA, Thailand, Korea, Poland, Greece, Ghana, Zimbabwe, England	Groundwater in fractures in crystalline rocks or alluvial placer deposits	Oxidising or mildly reducing conditions possible; high $\text{SO}_4$ concentrations (typically hundreds of $\text{mg L}^{-1}$ or higher); acidic (unless buffered by carbonate minerals); often increased concentrations of other trace metals (Ni, Pb, Zn, Cu, Cd). Arsenic speciation variable	Oxidation of sulphide minerals

### 12.5.3 Desorption of Arsenic at High pH Under Oxidising Conditions

Under the aerobic and acidic to near-neutral conditions typical of many natural environments, arsenic is strongly adsorbed by oxide minerals as the arsenate ion and the concentrations in solution are low. The non-linear nature of the adsorption isotherm for arsenate (Dzombak and Morel 1990) ensures that the amount of arsenic adsorbed is often relatively large, even when dissolved concentrations of arsenic are low. With an increase in pH, especially above pH 8.5, arsenate tends to desorb from oxide surfaces, thereby increasing the concentration in solution (Dzombak and Morel 1990). The impact of this is magnified by the high solid/solution ratios typical of unconsolidated aquifers.

The presence of other ions in the system can further impact on dissolved arsenic concentrations because other ions such as phosphate compete for sorption sites on the oxide (e.g. HFO, goethite) and will reduce the arsenic

loading on the solid surface. At high pH, phosphate is also released (Dzombak and Morel 1990). Other potential competing ions include bicarbonate, silicate and vanadium (Appelo et al. 2002; Smedley et al. 2005; Swedlund and Webster 1999). Bicarbonate is often the dominant anion in high-arsenic groundwaters and concentrations can be high, frequently exceeding  $500 \text{ mg L}^{-1}$ , and occasionally above  $1,000 \text{ mg L}^{-1}$  (Table 12.3). Concentrations of silica can also be high in some oxic arid groundwaters; high concentrations of vanadium are more unusual but have been found for example in Argentina (Smedley et al. 2002, 2005).

There are several reasons why the pH might increase but the most important in the present context is the uptake of protons by mineral weathering and ion-exchange reactions, together with the effect of evaporation in arid and semi-arid regions. A pH increase is commonly associated with the development of salinity and the salinisation of soils. Inputs of high-pH geothermal waters may be important in maintaining high arsenic concentrations in some alkaline

lakes. Desorption at high pH is the most likely mechanism for the development of groundwater-arsenic problems under oxidising conditions such as those observed in Argentina, Mexico and parts of the USA for example, and would account for the observed positive correlation of arsenic concentrations with increasing pH (e.g. Fig. 12.5). Increases in pH also induce the desorption of a wide variety of other oxyanions including those of V, U, Mo and B which are observed in many high-pH groundwaters (Fig. 12.5) (Bhattacharya et al. 2005; Smedley et al. 2002, 2005). These specifically-adsorbed anions all interact with adsorption sites on the oxides in a competitive way and so influence the extent of binding of each other.

By contrast, some cations, because of their positive charge, may promote the adsorption of negatively charged arsenate (Wilkie and Hering 1996). Calcium and magnesium are likely to be the most important cations in this respect because of their abundance in most natural waters and their +2 charge. Changes in surface charge and sorption capacity due to Ca/Na ion exchange have been implicated recently in the mobilisation of arsenate in the Southern High Plains aquifer, Texas (Scanlon et al. 2009). Other cations could also have an influence:  $\text{Fe}^{2+}$  may be important in reduced waters and  $\text{Al}^{3+}$  in acidic waters.

The aridity described above enables high pH values to be maintained. Aridity also minimises the flushing of released arsenic. It also allows the build-up of high chloride and fluoride concentrations. Other high-pH environments (up to pH 8.3), particularly open-system calcareous environments, are likely to be too well flushed to allow released arsenic to accumulate.

#### 12.5.4 Arsenic Desorption and Dissolution Due to a Change to Reducing Conditions

The onset of strongly reducing conditions, sufficient to enable iron(III) and probably sulphate reduction to take place is another well-documented trigger for the release of arsenic. Arsenic may be released by reductive dissolution, or reductive desorption, or a combination of the two. No clear relationship between dissolved concentrations of Fe(II) or As are expected as a result (Pedersen et al. 2006). Such processes occur in large alluvial systems, especially broad lowland meander belts and braided channels, and in prograding deltas. Microbial reactions are key and the organic carbon content and reactivity in the buried sediment can determine the rate at which reducing conditions are created (Postma et al. 2007; Rowland et al. 2007). Freshly-produced soil organic matter is reactive and readily decomposes; the presence of even small quantities can consume all of the available electron acceptors (e.g. dissolved oxygen, nitrate and sulphate) in the system. Solid-phase Fe

(III) in minerals may moderate the rate of reduction. Reducing conditions are maintained as long as the diffusion and convection of dissolved oxygen and other oxidants from the surface is less rapid than their consumption. This is facilitated if there is a confining layer of fine-grained material close to the surface. This commonly occurs in large deltas where fine-grained overbank deposits overlie coarser-grained alluvial deposits.

A change in the redox state of adsorbed arsenic from As(V) to As(III) also occurs during the diagenetic process. Arsenite is found as a dominant redox state in many reducing sediments (e.g. Rowland et al. 2005). Some studies have found As reduction coinciding with Fe(III) reduction (Herbel and Fendorf 2005). Others have concluded that reductive dissolution of iron oxides precedes the reduction of As(V) to As(III) (Islam et al. 2004).

The changing As redox state will also affect a large number of competing reactions. Phosphate-arsenite competition, for example, is likely to be less important than phosphate-arsenate competition. Competition with  $\text{HCO}_3^-$  and organic matter is possible (Appelo et al. 2002; Stachowicz et al. 2007). There is also the potential for arsenite-arsenate competition.

#### 12.5.5 Changes in the Structure of Oxide Minerals

Disordered and fine-grained iron oxides, including HFO, lepidocrocite, schwertmannite and magnetite, are common products of the early stages of weathering. Freshly-precipitated HFO is extremely fine-grained with a specific surface area of  $300 \text{ m}^2 \text{ g}^{-1}$  or more. HFO gradually transforms to more ordered structures such as goethite or haematite with larger crystal sizes and reduced surface areas. Goethite typically has specific surface areas of  $150 \text{ m}^2 \text{ g}^{-1}$  or less and those for haematite are even less (Cornell and Schwertmann 1996). This ageing reaction can take place rapidly in the laboratory but the rate in nature may be inhibited by the presence of other ions, particularly strongly adsorbed ions such as aluminium, phosphate, sulphate, arsenate, bicarbonate and silicate (Cornell and Schwertmann 1996). A reduction in surface area ultimately decreases the amount of As(V) adsorption possible on a weight-for-weight basis. If the site density (site  $\text{nm}^{-2}$ ) and binding affinities of the adsorbed ions remain constant, then as the specific surface area of the oxide mineral is reduced, some of the adsorbed ions may be desorbed (Dixit and Hering 2003).

Under strongly reducing conditions, additional processes could operate which lead to a change in the binding affinity for arsenic. In iron oxides, some of the surface iron could be reduced from Fe(III) to Fe(II) to produce a mixed-valence oxide such as magnetite. This has been shown to have variable effects on arsenic release, especially depending on



microbiological interactions. Several studies have shown that biotransformation of fresh iron oxides can lead to a net reduction in the release of As, at least in the short term until specific surface area is reduced to a point where As binding sites are diminished or reductive dissolution of the oxide occurs (Herbel and Fendorf 2006; Kocar et al. 2006; Pedersen et al. 2006; Tufano and Fendorf 2008). Herbel and Fendorf (2006) observed that in experiments with HFO reduction, *Sulfospirillum barnesii*, a bacterium capable of both Fe(III) and As(V) reduction, high concentrations of Fe(II) were induced which were capable of transforming ferrihydrite to goethite and magnetite, and although As was released initially, release diminished upon mineral transformation. Nonetheless, laboratory and field evidence suggests that at micromolar concentrations of arsenic, freshly-formed HFO binds more arsenic than the more crystalline goethite on a mole of Fe basis (de Vitre et al. 1991) and so a reduction in affinity is more likely overall (Dixit and Hering 2003; Herbel and Fendorf 2005).

In most soils and sediments, total As concentrations tend to correlate with total Fe. In Bangladesh, areas with high-As groundwaters tend to correspond with areas in which the sediments contain relatively large amounts of phosphate- or oxalate-extractable iron (BGS and DPHE 2001; van Geen et al. 2006; Zheng et al. 2005). These observations provide indirect support for the importance of labile iron oxides in releasing arsenic to solution. It is likely that the soils and sediments most sensitive to arsenic release on reduction and ageing are those in which iron oxides are abundant, in which HFO is initially a major fraction of the iron oxides present, and in which other arsenic-sorbing minerals are absent or scarce.

### 12.5.6 Mineral Dissolution

Mineral dissolution reactions tend to be most rapid under extremes of pH and Eh. For example, iron oxides dissolve under strongly acidic conditions and under strongly reducing conditions. Minor elements, including arsenic, present either as adsorbed (labile) arsenic or as irreversibly-bound (non-labile) arsenic will also tend to be released during the dissolution. This can explain, at least in part, the presence of high arsenic concentrations in acid mine drainage and in strongly reducing groundwaters. Reductive dissolution of Fe(III) oxides accounts for the high Fe(II) content of anaerobic waters.

As described above, one of the most pertinent mineral dissolution processes in respect of arsenic mobilisation is the oxidation of sulphide minerals. Pyrite, the most abundant of these minerals, can be an important source of arsenic, especially where it is freshly exposed by mining or by lowering of the water table. In extreme cases, this can lead to highly acidic groundwaters rich in sulphate, iron and trace metals. As the dissolved iron is neutralised and oxidised, it

precipitates as HFO with resultant adsorption and coprecipitation of dissolved As(V). In this sense, pyrite oxidation is not a very efficient mechanism for releasing arsenic into water.

### 12.5.7 Arsenic Transport Through Aquifers

The geochemical triggers described above are capable of releasing arsenic into groundwater but are not in themselves sufficient to account for the distribution of high-arsenic groundwaters observed in various parts of the world. The released arsenic must necessarily also have not been flushed away or diluted by groundwater flow. The rate of arsenic release must be set against the accumulated flushing of the aquifer during the period of release. The rocks of most aquifers used for drinking water are millions of years old and yet contain groundwater that may be at most a few thousand years old. Hence, a large number of pore volumes of fresh water will have passed through the aquifer over its history. This is also the case in most young aquifers with actively flowing groundwater. By contrast, many alluvial and deltaic aquifers are composed of relatively young sediments. Where groundwater flow is slow, these can contain relatively old groundwater. High groundwater arsenic concentrations can occur on a regional scale when geochemical conditions capable of mobilising arsenic are combined with hydrogeological conditions which prevent its removal.

The Quaternary period has seen considerable changes in climate and global sea level. Variation in groundwater piezometric levels over this period would have induced large variations in base levels of erosion and in groundwater flow regimes and rates. During the last glaciation, some 21,000–13,500 years ago, sea levels would have been up to 130 m below present mean sea level. This was a worldwide phenomenon and would have affected all then existing coastal aquifers. Continental and closed basin aquifers on the other hand would not have been affected. The increased hydraulic gradient in coastal aquifers during the glacial period would have resulted in correspondingly large groundwater flows and extensive flushing. The arsenic in groundwater in these older aquifers would therefore tend to have been flushed away. The deep unsaturated zone would also have led to more extensive oxidation of the shallower horizons with possible increased sorption of arsenic to Fe(III) oxides. Aquifers younger than around 7,000 years old, i.e. of Holocene age, will not have been subjected to this increased flushing. Such conditions are applicable to the southern Asian aquifers where Holocene high-As aquifers overlie older Pleistocene aquifers with low As concentrations. Differing flushing histories are likely contributory, in combination with different aquifer redox characteristics (grey vs. red-brown aquifer sands), to the differing aqueous As distributions observed.

The evidence suggests that in the Bengal Basin, Pleistocene aquifers have thus far been largely protected from contamination by As-rich groundwater from the overlying Holocene aquifers by clay aquitards that separate the two. However, that could change if drawdown of groundwater is induced by heavy pumping of the deep aquifer (Michael and Voss 2008). Such drawdown appears to have already affected the Pleistocene aquifer of the Red River delta of Vietnam, where intervening clay aquitards are thinner or in places absent (Winkel et al. 2011) and the aquifer is contaminated in places as a result of a century of heavy groundwater abstraction. Perhaps even more vulnerable are the currently low-As zones identified in Holocene deposits as lateral flow induced by pumping is more favourable than vertical flow in the stratified sediments (Fendorf et al. 2010).

At a local scale, small variations in relief or in drainage patterns may dictate local flow patterns and hence the distribution of arsenic-rich groundwater. Low-lying areas favour deposition of fine-grained deposits and are prone to flooding, under which reducing conditions can prevail and groundwater flow is restricted. Small-scale topographic depressions can also be the locations of seasonal discharge which restrict arsenic flushing. Such accumulations of dissolved As in topographic depressions have been observed for example in Argentina, China, Bangladesh, Hungary and Mexico (Nath et al. 2010; Smedley et al. 2002, 2003; Varsányi et al. 1991). It is clear that flat low-lying areas, particularly large deltas and inland basins, are prone to potentially high-arsenic groundwaters since they combine many of the risk factors identified above.

## 12.6 Concluding Remarks

This account has attempted to characterise the distribution of arsenic in the environment, and to describe the main geochemical controls on its speciation and mobilisation. We have also summarised the documented arsenic-rich areas and highlighted the main types of environment under which arsenic concentrations in waters are expected to be problematic. Doubtless there are other areas of the world where arsenic problems are yet to be recognised. However, increased awareness of the need for testing, increased health surveillance and increased recognition of the various risk factors associated with high-arsenic water mean that these are likely to be diminished in number. There are however, other areas where problems have been recognised but the scale is yet to be determined.

Environmental research has progressed significantly since the first aquifers were mapped or cases of arsenicosis recognised, although uncertainties remain in some of the key biogeochemical processes involved. On a simple level, mobilisation of arsenic relates to changes in speciation and

mineral solubility as a result of pH and redox changes in aquifers and the environment. But the processes involve a complex interplay of microbial interactions, ageing and diagenesis of metal oxides, ion competitive effects, mineral solubilities, presence of multiple minerals, nature and reactivity of organic matter, and flow.

One of the greatest hindrances to development is the enormous degree of spatial heterogeneity in aqueous As concentrations recognised in many if not all affected aquifers, often seemingly in aquifers with unremarkable concentrations of solid-phase arsenic. Such heterogeneity, on various scales, requires major testing and monitoring operations and hinders prediction of water quality for future drilling programmes. In the efforts to provide alternative sources of low-arsenic water to affected populations, significant challenges also exist to avoid risk substitution. Such risks include less-sustainable supplies, bacterial and other contamination of surface sources or future arsenic contamination of thus-far low-arsenic aquifers.

Groundwater provides an important source of drinking water to many millions of people globally. While this chapter has focused on areas where groundwater arsenic concentrations are often high, it is to be remembered that these are the exceptions. Groundwater more commonly provides a safe and reliable form of drinking water. In reducing exposure to the potentially fatal effects of faecal contamination, the use of groundwater for improving community health in developing countries has been highly successful. With regards to arsenic, most wells in most aquifers are likely to be uncontaminated, even when the groundwaters contain high concentrations of dissolved iron. It is also important to understand why these groundwaters are not affected. It appears that it is only when a number of critical geochemical and hydrogeological factors are combined that high-arsenic groundwaters occur.

*See Also the Following Chapters.* Chapter 24 (Environmental Medicine), Chapter 25 (Environmental Pathology), Chapter 33 (Modeling Groundwater Flow and Quality)

## References

- Ahmed KM, Hoque M, Hasan MK, Ravenscroft P, Chowdhury LR (1998) Occurrence and origin of water well methane gas in Bangladesh. *J Geol Soc India* 51:697–708
- Alexandratos VG, Elzinga EJ, Reeder RJ (2007) Arsenate uptake by calcite: macroscopic and spectroscopic characterization of adsorption and incorporation mechanisms. *Geochim Cosmochim Acta* 71:4172–4187
- Altamirano Espinoza M, Bundschuh J (2009) Natural arsenic groundwater contamination of the sedimentary aquifers of southwestern Sebaco valley, Nicaragua. In: Bundschuh J, Armienta MA, Birkle P, Bhattacharya P, Matschullat J, Mukherjee AB (eds) *Natural arsenic in groundwaters of Latin America*. CRC Press/Balkema, Leiden, pp 109–122

- Amirbahman A, Kent DB, Curtis GP, Davis JA (2006) Kinetics of sorption and abiotic oxidation of arsenic(III) by aquifer materials. *Geochim Cosmochim Acta* 70:533–547
- Appelo CAJ, Van der Weiden MJJ, Tournassat C, Charlet L (2002) Surface complexation of ferrous iron and carbonate on ferrihydrite and the mobilization of arsenic. *Environ Sci Technol* 36:3096–3103
- Ayotte JD, Nolan BT, Nucklos JR, Cantor KP, Robinson GR, Baris D, Hayes L, Kargas M, Bress W, Silverman DT, Lubin JH (2006) Modeling the probability of arsenic in groundwater in New England as a tool for exposure assessment. *Environ Sci Technol* 40:3578–3585
- Ayotte JD, Szabo Z, Focazio MJ, Eberts SM (2011) Effects of human-induced alteration of groundwater flow on concentrations of naturally-occurring trace elements at water-supply wells. *Appl Geochem* 26:747–762
- BAMWSP (2005) Bangladesh Arsenic Mitigation Water Supply Project (BAMWSP). Upazila wise summary results. <http://www.bwssp.org/BAMWSPContents/Survey%20Result/Upazila%20Summary.pdf>
- Barker SLL, Hickey KA, Cline JS, Dipple GM, Kilburn MR, Vaughan JR, Longo AA (2009) Unclenching invisible gold: use of nanoSIMS to evaluate gold, trace elements and sulfur isotopes in pyrite from Carlin-type gold deposits. *Econ Geol* 104:897–904
- Bates MN, Rey OA, Biggs ML, Hopenhayn C, Moore LE, Kalman D, Steinmaus C, Smith AH (2004) Case-control study of bladder cancer and exposure to arsenic in Argentina. *Am J Epidemiol* 159:381–389
- Bednar AJ, Garbarino JR, Burkhardt MR, Ranville JF, Wildeman TR (2004) Field and laboratory arsenic speciation methods and their application to natural-water analysis. *Water Res* 38:355–364
- Belkin HE, Zheng B, Finkelman RB (2000) Human health effects of domestic combustion of coal in rural China: a causal factor for arsenic and fluorine poisoning. In: The Second World Chinese conference on geological sciences, extended abstracts with programs, Stanford, pp 522–524
- Benner SG, Hansel CM, Wielinga BW, Barber TM, Fendorf S (2002) Reductive dissolution and biomineralization of iron hydroxide under dynamic flow conditions. *Environ Sci Technol* 36:1705–1711
- Berg M, Giger W, Tran HC, Pham HV, Trang PTK, Schertenleib R (2006) Extent and severity of arsenic pollution in Vietnam and Cambodia. In: Naidu R, Smith E, Owens G, Bhattacharya P, Nadebaum P (eds) *Managing arsenic in the environment – from soil to human health*. CSIRO Publishing, Collingwood
- Berg M, Stengel C, Trang PTK, Hung Viet P, Sampson ML, Leng M, Samreth S, Fredericks D (2007) Magnitude of arsenic pollution in the Mekong and Red River Deltas – Cambodia and Vietnam. *Sci Total Environ* 372:413–425
- Berner RA (1981) A new geochemical classification of sedimentary environments. *J Sedim Petrol* 51:0359–0365
- BGS, DPHE (2001) Arsenic contamination of groundwater in Bangladesh. In: Kinniburgh DG, Smedley PL (eds) *BGS technical report WC/00/19*, 4 vols. British Geological Survey, Keyworth (see [www.bgs.ac.uk/Arsenic/Bangladesh](http://www.bgs.ac.uk/Arsenic/Bangladesh))
- Bhattacharya P, Chatterjee D, Jacks G (1997) Occurrence of arsenic-contaminated groundwater in alluvial aquifers from delta plains, Eastern India: options for safe drinking water supply. *Water Resour Dev* 13:79–92
- Bhattacharya P, Tandukar N, Neku A, Valero AA, Mukherjee AB, Jacks G (2003) Geogenic arsenic in groundwaters from Terai alluvial plain of Nepal. *J Phys IV* 107:173–176
- Bhattacharya P, Claesson M, Fagerberg J, Bundschuh J, Storniolo ADR, Martin RA, Thir JM, Sracek O (2005) Natural arsenic in the groundwater of the alluvial aquifers of Santiago del Estero Province, Argentina. In: Bundschuh J, Bhattacharya P, Chandrasekharam D (eds) *Natural arsenic in groundwater: occurrence, remediation and management*. Balkema, Leiden, pp 57–65
- Bhattacharya P, Claesson M, Bundschuh J, Sracek O, Fagerberg J, Jacks G, Martin RA, Storniolo AD, Thir JM (2006) Distribution and mobility of arsenic in the Rio Dulce alluvial aquifers in Santiago del Estero Province, Argentina. *Sci Total Environ* 358:97–120
- Bian J, Tang J, Zhang L, Ma H, Zhao J (2012) Arsenic distribution and geological factors in the western Jilin province, China. *J Geochem Explor* 112:347–356
- Blanchard M, Alfredsson M, Brodholt J, Wright K, Catlow CRA (2007) Arsenic incorporation into FeS<sub>2</sub> pyrite and its influence on dissolution: a DFT study. *Geochim Cosmochim Acta* 71:624–630
- Blanes PS, Buchhamer EE, Gimenez MC (2011) Natural contamination with arsenic and other trace elements in groundwater of the Central-West region of Chaco, Argentina. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 46:1197–1206
- Bostick BC, Fendorf S (2003) Arsenite sorption on troilite (FeS) and pyrite (FeS<sub>2</sub>). *Geochim Cosmochim Acta* 67:909–921
- Boyle DR, Turner RJW, Hall GEM (1998) Anomalous arsenic concentrations in groundwaters of an island community, Bowen Island, British Columbia. *Environ Geochem Health* 20:199–212
- Breit GN, Foster AL, Sanzalone RF, Yount JC, Whitney JW, Welch AH, Islam MK, Islam MN (2001) Arsenic cycling in eastern Bangladesh: the role of phyllosilicates. *Geol Soc Am Abstr* 32: A192
- Brouste L, Marlin C, Dever L (1997) Geochemistry and residence time estimation of groundwater from the upper aquifer of the Chihuahua desert (Comarca Lagunera, Northern Mexico). *Appl Geochem* 12:775–786
- Burgess WG, Hoque MA, Michael HA, Voss CI, Breit GN, Ahmed KM (2010) Vulnerability of deep groundwater in the Bengal Aquifer System to contamination by arsenic. *Nat Geosci* 3:83–87
- Buschmann J, Berg M, Stengel C, Sampson ML (2007) Arsenic and manganese contamination of drinking water resources in Cambodia: coincidence of risk areas with low relief topography. *Environ Sci Technol* 41:2146–2152
- Cáceres L, Gruttner E, Contreras R (1992) Water recycling in arid regions: Chilean case. *Ambio* 21:138–144
- Castro de Esparza ML (2009) The presence of arsenic in drinking water in Latin America and its effect on public health. In: Bundschuh J, Armienta MA, Birkle P, Bhattacharya P, Matschullat J, Mukherjee AB (eds) *Natural arsenic in groundwaters of Latin America*. CRC Press/Balkema, Leiden, pp 17–29
- Cebrián ME, Albores MA, Garci-Vargas G, Del Razo LM, Ostrosky-Wegman P (1994) Chronic arsenic poisoning in humans. In: Nriagu JO (ed) *Arsenic in the environment, Part II: Human health and ecosystem effects*. Wiley, New York, pp 93–107
- Chakraborti D, Basu GK, Biswas BK, Chowdhury UK, Rahman MM, Paul K, Chowdhury TR, Chanda CR, Lodh D, Ray SL (2001) Characterization of arsenic-bearing sediments in the Gangetic delta of West Bengal, India. In: Chappell WR, Abernathy CO, Calderon RL (eds) *Arsenic exposure and health effects IV*. Elsevier, Amsterdam, pp 27–52
- Chakraborti D, Mukherjee SC, Pati S, Sengupta MK, Rahman MM, Chowdhury UK, Lodh D, Chanda CR, Chakraborty AK, Basul GK (2003) Arsenic groundwater contamination in Middle Ganga Plain, Bihar, India: a future danger? *Environ Health Perspect* 111:1194–1201
- Chakraborti D, Sengupta MK, Rahman MM, Ahamed S, Chowdhury UK, Hossain MA, Mukherjee SC, Pati S, Saha KC, Dutta RN, Quamruzzaman Q (2004) Groundwater arsenic contamination and its health effects in the Ganga-Meghna-Brahmaputra plain. *J Environ Monit* 6:74N–83N
- Chanpiwat P, Sthiannopkao S, Cho KH, Kim K-W, San V, Suvanthong B, Vongthavady C (2011) Contamination by arsenic and other trace elements of tube-well water along the Mekong River in Lao PDR. *Environ Pollut* 159:567–576

- Chen KY, Liu TK (2007) Major factors controlling arsenic occurrence in the groundwater and sediments of the Chianan coastal plain, SW Taiwan. *Terr Atmos Ocean Sci* 18:975–994
- Chen SL, Dzung SR, Yang MH, Chlu KH, Shieh GM, Wal CM (1994) Arsenic species in groundwaters of the Blackfoot disease areas, Taiwan. *Environ Sci Technol* 28:877–881
- Chen SL, Yeh SJ, Yang MH, Lin TH (1995) Trace element concentration and arsenic speciation in the well water of a Taiwan area with endemic Blackfoot disease. *Biol Trace Elem Res* 48:263–274
- Chitrakar RL, Neku A (2001) The scenario of arsenic in drinking water in Nepal. Department of Water Supply & Sewerage, Kathmandu, p 11
- Chouinard A, Paquette J, Williams-Jones AE (2005) Crystallographic controls on trace-element incorporation in auriferous pyrite from the Pascua epithermal high-sulfidation deposit, Chile-Argentina. *Can Mineral* 43:951–963
- Chowdhury TR, Manal BK, Samanta G, Basu GK, Chowdhury PP, Chanda CR, Karan NK, Lodh D, Dhar RK, Das D, Saha KC, Chakraborti D (1997) Arsenic in groundwater in six districts of West Bengal, India: the biggest arsenic calamity in the world: the status report up to August 1995. In: Abernathy CO, Calderon RL, Chappell WR (eds) *Arsenic: exposure and health effects*. Chapman and Hall, London, pp 93–111
- Cook SJ, Levson VM, Giles TR, Jackaman W (1995) A comparison of regional lake sediment and till geochemistry surveys – a case-study from the Fawnie Creek area, Central British Columbia. *Explor Min Geol* 4:93–110
- Cornell RM, Schwertmann U (1996) The iron oxides: structure, properties, reactions, occurrence and uses. VCH, Weinheim
- Cumbal L, Bundschuh J, Aguirre ME, Murgueitio E, Tipan I, Chavez C (2009) The origin of arsenic in waters and sediments from Papallacta lake area in Ecuador. In: Bundschuh J, Armentia MA, Birkle P, Bhattacharya P, Matschullat J, Mukherjee AB (eds) *Natural arsenic in groundwaters of Latin America*. CRC Press/Balkema, Boca Raton
- Cummings DE, Caccavo F, Fendorf S, Rosenzweig RF (1999) Arsenic mobilization by the dissimilatory Fe(III)-reducing bacterium *Shewanella alga* BrY. *Environ Sci Technol* 33:723–729
- Cummings DE, March AW, Bostick B, Spring S, Caccavo F Jr, Fendorf S, Rosenzweig RF (2000) Evidence for microbial Fe(III) reduction in anoxic, mining-impacted lake sediments (Lake Coeur d'Alene, Idaho). *Appl Environ Microbiol* 66:154–162
- Das D, Chatterjee A, Samanta G, Mandal B, Chowdhury TR, Chowdhury PP, Chanda C, Basu G, Lodh D, Nandi S, Chakraborty T, Mandal S, Bhattacharya SM, Chakraborti D (1994) Arsenic contamination in groundwater in 6 districts of West Bengal, India – the biggest arsenic calamity in the world. *Analyst* 119:N168–N170
- Das D, Chatterjee A, Mandal BK, Samanta G, Chakraborti D, Chanda B (1995) Arsenic in ground-water in 6 districts of West Bengal, India – the biggest arsenic calamity in the world. 2. Arsenic concentration in drinking-water, hair, nails, urine, skin-scale and liver-tissue (biopsy) of the affected people. *Analyst* 120:917–924
- Datta DK, Subramanian V (1997) Texture and mineralogy of sediments from the Ganges-Brahmaputra-Meghna river system in the Bengal basin, Bangladesh and their environmental implications. *Environ Geol* 30:181–188
- Davison W (1993) Iron and manganese in lakes. *Earth Sci Rev* 34:119–163
- de Vitre R, Belzile N, Tessier A (1991) Speciation and adsorption of arsenic on diagenetic iron oxyhydroxides. *Limnol Oceanogr* 36:1480–1485
- Deditius AP, Utsunomiya S, Renock D, Ewing RC, Ramana CV, Becker U, Kesler SE (2008) A proposed new type of arsenian pyrite: composition, nanostructure and geological significance. *Geochim Cosmochim Acta* 72:2919–2933
- Del Razo LM, Arellano MA, Cebrian ME (1990) The oxidation states of arsenic in well-water from a chronic arsenicosis area of Northern Mexico. *Environ Pollut* 64:143–153
- Del Razo LM, Hernandez JL, Garcia-Vargas GG, Ostrosky-Wegman P, Cortinas de Nava C, Cebrian ME (1994) Urinary excretion of arsenic species in a human population chronically exposed to arsenic via drinking water. A pilot study. In: Chappell WR, Abernathy CO, Cothorn CR (eds) *Arsenic exposure and health*. Science and Technology Letters, Northwood, pp 91–100
- Deng Y, Wang Y, Ma T (2009) Isotope and minor element geochemistry of high arsenic groundwater from Hangjinhouqi, the Hetuo Plain, Inner Mongolia. *Appl Geochem* 24:587–599
- Di Benedetto F, Costagliola P, Benvenuti M, Lattanzi P, Romanelli M, Tanelli G (2006) Arsenic incorporation in natural calcite lattice: evidence from electron spin echo spectroscopy. *Earth Planet Sci Lett* 246:458–465
- Dixit S, Hering JG (2003) Comparison of arsenic(V) and arsenic(III) sorption onto iron oxide minerals: implications for arsenic mobility. *Environ Sci Technol* 37:4182–4189
- Dudas MJ (1984) Enriched levels of arsenic in post-active acid sulfate soils in Alberta. *Soil Sci Soc Am J* 48:1451–1452
- Dzombak DA, Morel FMM (1990) Surface complexation modelling – hydrous ferric oxide. Wiley, New York
- Ellis AJ, Mahon WAJ (1977) *Chemistry and geothermal systems*. Academic, New York
- Farias SS, Casa VA, Vazquez C, Ferpozzi L, Pucci GN, Cohen IM (2003) Natural contamination with arsenic and other trace elements in ground waters of Argentine Pampean Plain. *Sci Total Environ* 309:187–199
- Fendorf S, Michael HA, van Geen A (2010) Spatial and temporal variations of groundwater arsenic in South and Southeast Asia. *Science* 328:1123–1127
- Focazio MJ, Welch AH, Watkins SA, Helsel DR, Horng MA (2000) A retrospective analysis of the occurrence of arsenic in groundwater resources of the United States and limitations in drinking water supply characterizations. U.S. Geological Survey, Denver, p 11
- Fontaine JA (1994) Regulating arsenic in Nevada drinking water supplies: past problems, future challenges. In: Chappell WR, Abernathy CO, Cothorn CR (eds) *Arsenic exposure and health*. Science and Technology Letters, Northwood, pp 285–288
- Fredrickson JK, Zachara JM, Kennedy DW, Dong H, Onstott TC, Hinman NW, Li S-M (1998) Biogenic iron mineralization accompanying the dissimilatory reduction of hydrous ferric oxide by a groundwater bacterium. *Geochim Cosmochim Acta* 62:3239–3257
- Fujii R, Swain WC (1995) Areal distribution of selected trace elements, salinity, and major ions in shallow ground water, Tulare Basin, Southern San Joaquin Valley, California. US geological survey water-resources investigations report 95-4048. USGS
- Garai R, Chakraborty A, Dey S, Saha K (1984) Chronic arsenic poisoning from tube-well water. *J Indian Med Assoc* 82:34–35
- Goldberg S (1986) Chemical modeling of arsenate adsorption on aluminum and iron oxide minerals. *Soil Sci Soc Am J* 50:1154–1157
- Goldberg S, Glaubig RA (1988) Anion sorption on a calcareous, montmorillonitic soil – arsenic. *Soil Sci Soc Am J* 52:1297–1300
- Gomez JJ, Lillo J, Sahun B (2006) Naturally occurring arsenic in groundwater and identification of the geochemical sources in the Duero Cenozoic Basin, Spain. *Environ Geol* 50:1151–1170
- Gomez ML, Blarasin MT, Martinez DE (2009) Arsenic and fluoride in a loess aquifer in the central area of Argentina. *Environ Geol* 57:143–155
- Guerin WF, Blakemore RP (1992) Redox cycling of iron supports growth and magnetite synthesis by *Aquaspirillum magnetotacticum*. *Appl Environ Microbiol* 58:1102–1109

- Guo HR, Chen CJ, Greene HL (1994) Arsenic in drinking water and cancers: a brief descriptive view of Taiwan studies. In: Chappell WR, Abernathy CO, Cothorn CR (eds) Arsenic exposure and health. Science and Technology Letters, Northwood, pp 129–138
- Guo XJ, Fujino Y, Kaneko S, Wu KG, Xia YJ, Yoshimura T (2001) Arsenic contamination of groundwater and prevalence of arsenical dermatosis in the Hetao plain area, Inner Mongolia, China. *Mol Cell Biochem* 222:137–140
- Guo HM, Wang YX, Shepeizer GM, Yan SL (2003) Natural occurrence of arsenic in shallow groundwater, Shanyin, Datong Basin, China. *J Environ Sci Health A Toxic/Hazard Subst Environ Eng* 38:2565–2580
- Gurung J, Ishiga H, Khadka MS (2005) Geological and geochemical examination of arsenic contamination in groundwater in the Holocene Terai Basin, Nepal. *Environ Geol* 49:98–113
- Gurung JK, Ishiga H, Khadka MS, Shrestha NR (2007) The geochemical study of fluvio-lacustrine aquifers in the Kathmandu Basin (Nepal) and the implications for the mobilization of arsenic. *Environ Geol* 52:503–517
- Gurzau ES, Gurzau AE (2001) Arsenic exposure from drinking groundwater in Transylvania, Romania: an overview. In: Chappell WR, Abernathy CO, Calderon RL (eds) Arsenic exposure and health effects IV. Elsevier, Amsterdam, pp 181–184
- Gustafsson JP, Tin NT (1994) Arsenic and selenium in some Vietnamese acid sulfate soils. *Sci Total Environ* 151:153–158
- Harvey CF, Swartz CH, Badruzzaman ABM, Keon-Blute N, Yu W, Ali MA, Jay J, Beckie R, Niedan V, Brabander D, Oates PM, Ashfaq KN, Islam S, Hemond HF, Ahmed MF (2002) Arsenic mobility and groundwater extraction in Bangladesh. *Science* 298:1602–1606
- Harvey CF, Ashfaq KN, Yu W, Badruzzaman ABM, Ali MA, Oates PM, Michael HA, Neumann RB, Beckie R, Islam S, Ahmed MF (2006) Groundwater dynamics and arsenic contamination in Bangladesh. *Chem Geol* 228:112–136
- Herbel M, Fendorf S (2005) Transformation and transport of arsenic within ferric hydroxide coated sands upon dissimilatory reducing bacterial activity. In: O'Day PA, Vlassopoulos D, Meng X, Benning LG (eds) Advances in arsenic research: integration of experimental and observational studies and implications for mitigation. American Chemical Society, Washington, DC, pp 77–90
- Herbel M, Fendorf S (2006) Biogeochemical processes controlling the speciation and transport of arsenic within iron coated sands. *Chem Geol* 228:16–32
- Hering J, Kneebone PE (2002) Biogeochemical controls on arsenic occurrence and mobility in water supplies. In: Frankenberger W (ed) Environmental chemistry of arsenic. Marcel Dekker, New York, pp 155–181
- Hopenhayn-Rich C, Biggs ML, Fuchs A, Bergoglio R, Tello EE, Nicolli H, Smith AH (1996) Bladder-cancer mortality associated with arsenic in drinking water in Argentina. *Epidemiology* 7:117–124
- Horneman A, van Geen A, Kent D, Mathe PE, Zheng Y, Dhar RK, O'Connell S, Hoque M, Aziz Z, Shamsudduha M, Seddique AA, Ahmad KM (2004) Decoupling of As and Fe release to Bangladesh groundwater under reducing conditions. Part 1: Evidence from sediment profiles. *Geochim Cosmochim Acta* 68:3459–3473
- Hsu KH, Froines JR, Chen CJ (1997) Studies of arsenic ingestion from drinking water in northeastern Taiwan: chemical speciation and urinary metabolites. In: Abernathy CO, Calderon RL, Chappell WR (eds) Arsenic exposure and health effects. Chapman and Hall, London, pp 190–209
- Huang JH, Matzner E (2006) Dynamics of organic and inorganic arsenic in the solution phase of an acidic fen in Germany. *Geochim Cosmochim Acta* 70:2023–2033
- IMTA (1992) Estudio hidroquímico e isotópico del acuífero granular de la Comarca Lagunera. Instituto Mexicano de Tecnología del Agua, Cuernavaca, México
- Islam FS, Gault AG, Boothman C, Polya DA, Charnock JM, Chatterjee D, Lloyd JR (2004) Role of metal-reducing bacteria in arsenic release from Bengal delta sediments. *Nature* 430:68–71
- Islam FS, Boothman C, Gault AG, Polya DA, Lloyd JR (2005a) Potential role of the Fe(III)-reducing bacteria *Geobacter* and *Geothrix* in controlling arsenic solubility in Bengal delta sediments. *Mineral Mag* 69:865–875
- Islam FS, Pederick RL, Gault AG, Adams LK, Polya DA, Charnock JM, Lloyd JR (2005b) Interactions between the Fe(III)-reducing bacterium *Geobacter sulfurreducens* and arsenate, and capture of the metalloid by biogenic Fe(II). *Appl Environ Microbiol* 71:8642–8648
- Junta Municipal (2000) Prospección hidrogeológica e hidrogeoquímica en el bolsón de la Mesilla, Chihuahua (Zona de Conejos-Medanos), Report Técnico No DS GEO-011/00. Junta Municipal de Agua y Saneamiento de Juárez
- Karcher S, Caceres L, Jekel M, Contreras R (1999) Arsenic removal from water supplies in Northern Chile using ferric chloride coagulation. *J Chart Inst Water Environ Manage* 13:164–169
- Klump S, Kipfer R, Cirpka OA, Harvey CF, Brennwald MS, Ashfaq KN, Badruzzaman ABM, Hug SJ, Imboden DM (2006) Groundwater dynamics and arsenic mobilization in Bangladesh assessed using noble gases and tritium. *Environ Sci Technol* 40:243–250
- Kocar BD, Herbel MJ, Tufano KJ, Fendorf S (2006) Contrasting effects of dissimilatory iron(III) and arsenic(V) reduction on arsenic retention and transport. *Environ Sci Technol* 40:6715–6721
- Kondo H, Ishiguro Y, Ohno K, Nagase M, Toba M, Takagi M (1999) Naturally occurring arsenic in the groundwaters in the southern region of Fukuoka Prefecture, Japan. *Water Res* 33:1967–1972
- Lewis C, Ray DT, Chiu KK (2007) Primary geologic sources of arsenic in the Chianan Plain (Blackfoot disease area) and the Lanyang Plain of Taiwan. *Int Geol Rev* 49:947–961
- Lin TH, Huang YL, Wang MY (1998) Arsenic species in drinking water, hair, fingernails, and urine of patients with blackfoot disease. *J Toxicol Environ Health A* 53:85–93
- Lloyd JR, Oremland RS (2006) Microbial transformations of arsenic in the environment: from soda lakes to aquifers. *Elements* 2:85–90
- Lu FJ, Shih SR, Liu TM, Shown SH (1990) The effect of fluorescent humic substances existing in the well water of Blackfoot disease endemic areas in Taiwan on prothrombin time and activated partial thromboplastin time in vitro. *Thromb Res* 57:747–753
- Luo ZD, Zhang YM, Ma L, Zhang GY, He X, Wilson R, Byrd DM, Griffiths JG, Lai S, He L, Grumski K, Lamm SH (1997) Chronic arsenicosis and cancer in Inner Mongolia – consequences of well-water arsenic levels greater than 50  $\mu\text{g l}^{-1}$ . In: Abernathy CO, Calderon RL, Chappell WR (eds) Arsenic exposure and health effects. Chapman and Hall, London, pp 55–68
- Maest AS, Pasilis SP, Miller LG, Nordstrom DK (1992) Redox geochemistry of arsenic and iron in Mono Lake, California, USA. In: Kharaka Y, Maest A (eds) Proceedings of the seventh international symposium on water-rock interaction, Park City, Utah, USA, 13–18 July 1992. A.A. Balkema, Rotterdam, pp 507–511
- Maharjan M, Watanabe C, Ahmad SA, Umezaki M, Ohtsuka R (2007) Mutual interaction between nutritional status and chronic arsenic toxicity due to groundwater contamination in an area of Terai, lowland Nepal. *J Epidemiol Community Health* 61:389–394
- Manning BA, Goldberg S (1997) Adsorption and stability of arsenic (III) at the clay mineral-water interface. *Environ Sci Technol* 31:2005–2011
- Masscheleyn PH, DeLaune RD, Patrick WH (1991) Effect of redox potential and pH on arsenic speciation and solubility in a contaminated soil. *Environ Sci Technol* 25:1414–1419
- McArthur JM, Ravenscroft P, Safiulla S, Thirwall MF (2001) Arsenic in groundwater: testing pollution mechanisms for sedimentary aquifers in Bangladesh. *Water Resour Res* 37:109–117

- McArthur JM, Banerjee DM, Hudson-Edwards KA, Mishra R, Purohit R, Ravenscroft P, Cronin A, Howarth RJ, Chatterjee A, Talukder T, Lowry D, Houghton S, Chadha DK (2004) Natural organic matter in sedimentary basins and its relation to arsenic in anoxic ground water: the example of West Bengal and its worldwide implications. *Appl Geochem* 19:1255–1293
- Michael HA, Voss CI (2008) Evaluation of the sustainability of deep groundwater as an arsenic-safe resource in the Bengal Basin. *Proc Natl Acad Sci U S A* 105:8531–8536
- Morales L, Puigdomenech C, Punti A, Torres E, Canyellas C, Cortina JL, Sancha AM (2009) Arsenic and water quality of rural community wells in San Juan de Limay, Nicaragua. In: Bundschuh J, Armienta MA, Birkle P, Bhattacharya P, Matschullat J, Mukherjee AB (eds) *Natural arsenic in groundwaters of Latin America*. CRC Press/Balkema, Leiden, pp 123–128
- Mukherjee AB, Bhattacharya P, Jacks G, Banerjee DM, Ramanathan CM, Chandrashekharam D, Chatterjee D, Naidu R (2006) Groundwater arsenic contamination in India. In: Naidu R, Smith E, Owens G, Bhattacharya P, Nadebaum P (eds) *Managing arsenic in the environment – from soil to human health*. CSIRO Publishing, Collingwood
- Nath B, Mallik SB, Stueben D, Chatterjee D, Charlet L (2010) Electrical resistivity investigation of the arsenic affected alluvial aquifers in West Bengal, India: usefulness in identifying the areas of low and high groundwater arsenic. *Environ Earth Sci* 60:873–884
- Nguyen VA, Bang S, Pham Hung V, Kim K-W (2009) Contamination of groundwater and risk assessment for arsenic exposure in Ha Nam province, Vietnam. *Environ Int* 35:466–472
- Nickson RT, McArthur JM, Ravenscroft P, Burgess WG, Ahmed KM (2000) Mechanism of arsenic release to groundwater, Bangladesh and West Bengal. *Appl Geochem* 15:403–413
- Nicolli HB, Suriano JM, Peral MAG, Ferpozzi LH, Baleani OA (1989) Groundwater contamination with arsenic and other trace-elements in an area of the Pampa, province of Cordoba, Argentina. *Environ Geol Water Sci* 14:3–16
- Nicolli HB, Bundschuh J, Garcia JW, Falcon CM, Jean J-S (2010) Sources and controls for the mobility of arsenic in oxidizing groundwaters from loess-type sediments in arid/semi-arid dry climates Evidence from the Chaco-Pampean plain (Argentina). *Water Res* 44:5589–5604
- Nimick DA, Moore JN, Dalby CE, Savka MW (1998) The fate of geothermal arsenic in the Madison and Missouri Rivers, Montana and Wyoming. *Water Resour Res* 34:3051–3067
- Ning Z, Lobdell DT, Kwok RK, Liu Z, Zhang S, Ma C, Riediker M, Mumford JL (2007) Residential exposure to drinking water arsenic in Inner Mongolia, China. *Toxicol Appl Pharmacol* 222: 351–356
- Nordstrom DK (2002) Public health – worldwide occurrences of arsenic in ground water. *Science* 296:2143–2144
- Nordstrom DK, Alpers CN (1999) Negative pH, efflorescent mineralogy, and consequences for environmental restoration at the Iron Mountain superfund site, California. *Proc Natl Acad Sci U S A* 96:3455–3462
- O'Day PA, Vlassopoulos D, Root R, Rivera N (2004) The influence of sulfur and iron on dissolved arsenic concentrations in the shallow subsurface under changing redox conditions. *Proc Natl Acad Sci U S A* 101:13703–13708
- Oremland RS, Stolz JF (2003) The ecology of arsenic. *Science* 300:93–94
- Oremland RS, Stolz JF (2005) Arsenic, microbes and contaminated aquifers. *Trends Microbiol* 13:45–49
- Oremland R, Newman D, Kail B, Stolz J (2002) Bacterial respiration of arsenate and its significance in the environment. In: Frankenberger W (ed) *Environmental chemistry of arsenic*. Marcel Dekker, New York, pp 273–295
- Oremland RS, Stolz JF, Hollibaugh JT (2004) The microbial arsenic cycle in Mono Lake, California. *FEMS Microbiol Ecol* 48:15–27
- Oscarson DW, Huang PM, Defosse C, Herbillon A (1981a) Oxidative power of Mn(IV) and Fe(III) oxides with respect to As(III) in terrestrial and aquatic environments. *Nature* 291:50–51
- Oscarson DW, Huang PM, Liaw WK (1981b) Role of manganese in the oxidation of arsenite by freshwater lake sediments. *Clays Clay Miner* 29:219–225
- Pant BR (2011) Ground water quality in the Kathmandu valley of Nepal. *Environ Monit Assess* 178:477–485
- Pederick RL, Gault AG, Charnock JM, Polya DA, Lloyd JR (2007) Probing the biogeochemistry of arsenic: response of two contrasting aquifer sediments from Cambodia to stimulation by arsenate and ferric iron. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 42:1763–1774
- Pedersen HD, Postma D, Jakobsen R (2006) Release of arsenic associated with the reduction and transformation of iron oxides. *Geochim Cosmochim Acta* 70:4116–4129
- Plumlee GS, Smith KS, Montour MR, Ficklin WH, Mosier EL (1999) Geologic controls on the composition of natural waters and mine waters draining diverse mineral-deposit types. In: Filipek LH, Plumlee GS (eds) *Reviews in economic geology*, vol 6B. Society of Economic Geologists, Littleton, pp 373–432
- Pokhrel D, Bhandari BS, Viraraghavan T (2009) Arsenic contamination of groundwater in the Terai region of Nepal: an overview of health concerns and treatment options. *Environ Int* 35:157–161
- Polizzotto ML, Harvey CF, Sutton SR, Fendorf S (2005) Processes conducive to the release and transport of arsenic into aquifers of Bangladesh. *Proc Natl Acad Sci U S A* 102:18819–18823
- Polizzotto ML, Harvey CF, Li GC, Badruzzaman B, Ali A, Newville M, Sutton S, Fendorf S (2006) Solid-phases and desorption processes of arsenic within Bangladesh sediments. *Chem Geol* 228:97–111
- Polizzotto ML, Kocar BD, Benner SG, Sampson M, Fendorf S (2008) Near-surface wetland sediments as a source of arsenic release to ground water in Asia. *Nature* 454:505–508
- Polya DA, Gault AG, Diebe N, Feldman P, Rosenboom JW, Gilligan E, Fredericks D, Milton AH, Sampson M, Rowland HAL, Lythgoe PR, Jones JC, Middleton C, Cooke DA (2005) Arsenic hazard in shallow Cambodian groundwaters. *Mineral Mag* 69:807–823
- Postma D, Larsen F, Hue NTM, Duc MT, Viet PH, Nhan PQ, Jessen S (2007) Arsenic in groundwater of the Red River floodplain, Vietnam: controlling geochemical processes and reactive transport modeling. *Geochim Cosmochim Acta* 71:5054–5071
- Rahman MM, Chowdhury UK, Mukherjee SC, Mondal BK, Paul K, Lodh D, Biswas BK, Chanda CR, Basu GK, Saha KC, Roy S, Das R, Palit SK, Quamruzzaman Q, Chakraborti D (2001) Chronic arsenic toxicity in Bangladesh and West Bengal, India – a review and commentary. *J Toxicol Clin Toxicol* 39:683–700
- Reich M, Becker U (2006) First-principles calculations of the thermodynamic mixing properties of arsenic incorporation into pyrite and marcasite. *Chem Geol* 225:278–290
- Reimann C, Bjorvatn K, Frengstad B, Melaku Z, Tekle-Haimanot R, Siewers U (2003) Drinking water quality in the Ethiopian section of the East African Rift Valley I – data and health aspects. *Sci Total Environ* 311:65–80
- Riedel FN, Eikmann T (1986) Natural occurrence of arsenic and its compounds in soils and rocks. *Wiss Umwelt* 3–4:108–117
- Robertson FN (1989) Arsenic in groundwater under oxidizing conditions, south-west United States. *Environ Geochem Health* 11:171–185
- Robinson B, Outred H, Brooks R, Kirkman J (1995) The distribution and fate of arsenic in the Waikato River System, North Island, New Zealand. *Chem Spec Bioavailab* 7:89–96
- Romero L, Alonso H, Campano P, Fanfani L, Cidu R, Dadea C, Keegan T, Thornton I, Farago M (2003) Arsenic enrichment in waters and sediments of the Rio Loa (Second Region, Chile). *Appl Geochem* 18:1399–1416

- Romero FM, Armienta MA, Carrillo-Chavez A (2004) Arsenic sorption by carbonate-rich aquifer material, a control on arsenic mobility at Zimapan, Mexico. *Arch Environ Contam Toxicol* 47:1–13
- Rosas I, Belmont R, Armienta A, Baez A (1999) Arsenic concentrations in water, soil, milk and forage in Comarca Lagunera, Mexico. *Water Air Soil Pollut* 112:133–149
- Rosenboom JW (2004) Not just red or green. An analysis of arsenic data from 15 upazilas in Bangladesh. APSU, Arsenic Policy Support Unit, Bangladesh
- Rowland HAL, Gault AG, Charnock JM, Polya DA (2005) Preservation and XANES determination of the oxidation state of solid-phase arsenic in shallow sedimentary aquifers in Bengal and Cambodia. *Mineral Mag* 69:825–839
- Rowland HAL, Pederick RL, Polya DA, Pancost RD, Van Dongen BE, Gault AG, Vaughan DJ, Bryant C, Anderson B, Lloyd JR (2007) The control of organic matter on microbially mediated iron reduction and arsenic release in shallow alluvial aquifers, Cambodia. *Geobiology* 5:281–292
- Ryker SJ (2001) Mapping arsenic in groundwater: a real need, but a hard problem – why was the map created? *Geotimes* 46:34–36
- Sampson ML, Bostick B, Chiew H, Hagan JM, Shantz A (2008) Arsenicosis in Cambodia: case studies and policy response. *Appl Geochem* 23:2977–2986
- Savage KS, Tingle TN, O'Day PA, Waychunas GA, Bird DK (2000) Arsenic speciation in pyrite and secondary weathering phases, Mother Lode Gold District, Tuolumne County, California. *Appl Geochem* 15:1219–1244
- Savage KS, Bird DK, O'Day PA (2005) Arsenic speciation in synthetic jarosite. *Chem Geol* 215:473–498
- Scanlon BR, Nicot JP, Reedy RC, Kurtzman D, Mukherjee A, Nordstrom DK (2009) Elevated naturally occurring arsenic in a semi-arid oxidizing system, Southern High Plains aquifer, Texas, USA. *Appl Geochem* 24:2061–2071
- Schreiber ME, Simo JA, Freiberg PG (2000) Stratigraphic and geochemical controls on naturally occurring arsenic in groundwater, eastern Wisconsin, USA. *Hydrogeol J* 8:161–176
- Schrenk MO, Edwards KJ, Goodman RM, Hamers RJ, Banfield JF (1998) Distribution of *Thiobacillus ferrooxidans* and *Leptospirillum ferrooxidans*: implications for generation of acid mine drainage. *Science* 279:1519–1522
- Shotyk W (1996) Natural and anthropogenic enrichments of As, Cu, Pb, Sb, and Zn in ombrotrophic versus minerotrophic peat bog profiles, Jura Mountains, Switzerland. *Water Air Soil Pollut* 90:375–405
- Smedley PL, Kinniburgh DG (2002) A review of the source, behaviour and distribution of arsenic in natural waters. *Appl Geochem* 17:517–568
- Smedley PL, Nicolli HB, Macdonald DMJ, Barros AJ, Tullio JO (2002) Hydrogeochemistry of arsenic and other inorganic constituents in groundwaters from La Pampa, Argentina. *Appl Geochem* 17:259–284
- Smedley PL, Zhang M-Y, Zhang G-Y, Luo Z-D (2003) Mobilisation of arsenic and other trace elements in fluvio-lacustrine aquifers of the Huhhot Basin, Inner Mongolia. *Appl Geochem* 18:1453–1477
- Smedley PL, Kinniburgh DG, Macdonald DMJ, Nicolli HB, Barros AJ, Tullio JO, Pearce JM, Alonso MS (2005) Arsenic associations in sediments from the loess aquifer of La Pampa, Argentina. *Appl Geochem* 20:989–1016
- Smedley PL, Knudsen J, Maiga D (2007) Arsenic in groundwater from mineralised Proterozoic basement rocks of Burkina Faso. *Appl Geochem* 22:1074–1092
- So HU, Postma D, Jakobsen R, Larsen F (2008) Sorption and desorption of arsenate and arsenite on calcite. *Geochim Cosmochim Acta* 72:5871–5884
- Sracek A, Novak M, Sulovsky P, Martin R, Bundschuh J, Bhattacharya P (2009) Mineralogical study of arsenic-enriched aquifer sediments at Santiago del Estero, Northwest Argentina. In: Bundschuh J, Armienta MA, Birkle P, Bhattacharya P, Matschullat J, Mukherjee AB (eds) *Natural arsenic in groundwaters of Latin America*. CRC Press/Balkema, Leiden
- Stachowicz M, Hiemstra T, Van Riemsdijk WH (2007) Arsenic-bicarbonate interaction on goethite particles. *Environ Sci Technol* 41:5620–5625
- Stollenwerk KG (2003) Geochemical processes controlling transport of arsenic in groundwater: a review of adsorption. In: Stollenwerk KG (ed) *Arsenic in ground water: geochemistry and occurrence*. Kluwer, Boston, pp 67–100
- Stollenwerk KG, Breit GN, Welch AH, Yount JC, Whitney JW, Foster AL, Uddin MN, Majumder RK, Ahmed N (2007) Arsenic attenuation by oxidized aquifer sediments in Bangladesh. *Sci Total Environ* 379:133–150
- Swartz CH, Blute NK, Badruzzaman B, Ali A, Brabander D, Jay J, Besancon J, Islam S, Hemond HF, Harvey CF (2004) Mobility of arsenic in a Bangladesh aquifer: inferences from geochemical profiles, leaching data, and mineralogical characterization. *Geochim Cosmochim Acta* 68:4539–4557
- Swedlund PJ, Webster JG (1999) Adsorption and polymerisation of silicic acid on ferrihydrite, and its effect on arsenic adsorption. *Water Res* 33:3413–3422
- Tandukar N, Bhattacharya P, Jacks G, Valero AA (2005) Naturally occurring arsenic in groundwater of Terai region in Nepal and mitigation options. In: Bundschuh J, Bhattacharya P, Chandrasekharam D (eds) *Natural arsenic in groundwater: occurrence, remediation and management*. A.A. Balkema, Leiden, pp 41–48
- Taylor RM (1980) Formation and properties of the Fe(II)Fe(III) hydroxy-carbonate and its possible significance in soil formation. *Clay Miner* 15:369–382
- Thompson JM, Demonge JM (1996) Chemical analyses of hot springs, pools, and geysers from Yellowstone National Park, Wyoming, and vicinity, 1980–1993. USGS Open-File Report
- Tseng WP, Chu HM, How SW, Fong JM, Lin CS, Yeh S (1968) Prevalence of skin cancer in an endemic area of chronic arsenicosis in Taiwan. *J Natl Cancer Inst* 40:453–463
- Tufano KJ, Fendorf S (2008) Confounding impacts of iron reduction on arsenic retention. *Environ Sci Technol* 42:4777–4783
- van Geen A, Zheng Y, Stute M, Ahmed KM (2003a) Comment on “Arsenic mobility and groundwater extraction in Bangladesh” (II). *Science* 300:584C
- van Geen A, Zheng Y, Versteeg R, Stute M, Horneman A, Dhar R, Steckler M, Gelman A, Small C, Ahsan H, Graziano JH, Hussain I, Ahmed KM (2003b) Spatial variability of arsenic in 6000 tube wells in a 25 km<sup>2</sup> area of Bangladesh. *Water Resour Res* 39, art. no.-1140
- van Geen A, Zheng Y, Cheng Z, Aziz Z, Horneman A, Dhar RK, Mailloux B, Stute M, Weinman B, Goodbred S, Seddique AA, Hope MA, Ahmed KM (2006) A transect of groundwater and sediment properties in Arai-hazar, Bangladesh: further evidence of decoupling between As and Fe mobilization. *Chem Geol* 228:85–96
- van Geen A, Zheng Y, Goodbred S, Horneman A, Aziz Z, Cheng Z, Stute M, Mailloux B, Weinman B, Hoque MA, Seddique AA, Hossain MS, Chowdhury SH, Ahmed KM (2008) Flushing history as a hydrogeological control on the regional distribution of arsenic in shallow groundwater of the Bengal Basin. *Environ Sci Technol* 42:2283–2288
- Varsányi I, Fodré Z, Bartha A (1991) Arsenic in drinking water and mortality in the southern Great Plain, Hungary. *Environ Geochem Health* 13:14–22
- Wang L, Huang J (1994) Chronic arsenism from drinking water in some areas of Xinjiang, China. In: Nriagu JO (ed) *Arsenic in the environment, Part II: Human health and ecosystem effects*. Wiley, New York, pp 159–172

- Warren C (2001) Hydrogeology and groundwater quality of Los Pereyras, Tucuman, Argentina, University College. University College London, London
- Webster JG (1999) Arsenic. In: Marshall CP, Fairbridge RW (eds) *Encyclopaedia of geochemistry*. Chapman and Hall, London, pp 21–22
- Webster JG (2003) Arsenic adsorption on natural riverine suspended particulate material (SPM). *Abstr Pap Am Chem Soc* 226:U583
- Webster JG, Nordstrom DK (2003) Geothermal arsenic. In: Stollenwerk KG (ed) *Arsenic in ground water: geochemistry and occurrence*. Kluwer, Boston, pp 101–125
- Welch AH, Lico MS, Hughes JL (1988) Arsenic in ground-water of the Western United States. *Ground Water* 26:333–347
- Welch AH, Westjohn DB, Helsel DR, Wanty RB (2000) Arsenic in ground water of the United States: occurrence and geochemistry. *Groundwater* 38:589–604
- White DE, Hem JD, Waring GA (1963) Chemical composition of sub-surface waters. In: Fleischer M (ed) *Data of geochemistry*, 6th edn. Geological Survey, Washington, DC, pp 1–67
- Wilkie JA, Hering JG (1996) Adsorption of arsenic onto hydrous ferric oxide: effects of adsorbate/adsorbent ratios and co-occurring solutes. *Colloids Surf A Physicochem Eng Asp* 107:97–110
- Wilkie JA, Hering JG (1998) Rapid oxidation of geothermal arsenic (III) in streamwaters of the eastern Sierra Nevada. *Environ Sci Technol* 32:657–662
- Williams M (1997) Mining-related arsenic hazards: Thailand case-study. British Geological Survey technical report WC/97/49, 36 pp
- Williams M, Fordyce F, Pajitprapapon A, Charoenchaisri P (1996) Arsenic contamination in surface drainage and groundwater in part of the southeast Asian tin belt, Nakhon Si Thammarat Province, southern Thailand. *Environ Geol* 27:16–33
- Winkel LHE, Pham Thi Kim T, Vi Mai L, Stengel C, Amini M, Nguyen Thi H, Pham Hung V, Berg M (2011) Arsenic pollution of groundwater in Vietnam exacerbated by deep aquifer exploitation for more than a century. *Proc Natl Acad Sci U S A* 108:1246–1251
- Wyatt CJ, Fimbres C, Romo L, Mendez RO, Grijalva M (1998) Incidence of heavy metal contamination in water supplies in Northern Mexico. *Environ Res* 76:114–119
- Xie X, Wang Y, Su C, Liu H, Duan M, Xie Z (2008) Arsenic mobilization in shallow aquifers of Datong Basin: hydrochemical and mineralogical evidences. *J Geochem Explor* 98:107–115
- Xie X, Ellis A, Wang Y, Xie Z, Duan M, So C (2009) Geochemistry of redox-sensitive elements and sulfur isotopes in the high arsenic groundwater system of Datong Basin, China. *Sci Total Environ* 407:3823–3835
- Ying SC, Kocar BD, Griffis SD, Fendorf S (2011) Competitive microbially and Mn oxide mediated redox processes controlling arsenic speciation and partitioning. *Environ Sci Technol* 45: 5572–5579
- Yokoyama T, Takahashi Y, Tarutani T (1993) Simultaneous determination of arsenic and arsenious acids in geothermal water. *Chem Geol* 103:103–111
- Zheng Y, van Geen A, Stute M, Dhar R, Mo Z, Cheng Z, Horneman A, Gavrieli I, Simpson HJ, Versteeg R, Steckler M, Grazioli-Venier A, Goodbred S, Shahnewaz M, Shamsudduha M, Hoque MA, Ahmed KM (2005) Geochemical and hydrogeological contrasts between shallow and deeper aquifers in two villages of Araihasar, Bangladesh: implications for deeper aquifers as drinking water sources. *Geochim Cosmochim Acta* 69:5203–5218