

Combinatorial Homotopy Categories



Carles Casacuberta and Jiří Rosický

Abstract A model category is called combinatorial if it is cofibrantly generated and its underlying category is locally presentable. As shown in recent years, homotopy categories of combinatorial model categories share useful properties, such as being well generated and satisfying a very general form of Ohkawa’s theorem.

Keywords Combinatorial model category · Cofibrantly generated · Locally presentable · Well generated · Brown representability

1 Introduction

The term “combinatorial” in topology classically refers to discrete methods or, more specifically, to the use of polyhedra, simplicial complexes or cell complexes in order to deal with topological problems [17, 32].

In the context of Quillen model categories in homotopy theory [25], those called *combinatorial* are, by definition, the cofibrantly generated ones whose underlying category is locally presentable. For example, simplicial sets are combinatorial, but topological spaces are not. As a consequence of this fact, certain constructions involving homotopy colimits, such as Bousfield localizations, may seem intricate if one works with topological spaces while they have become standard technology in the presence of combinatorial models [2, 6, 12].

One key feature of combinatorial model categories is that they admit presentations in terms of generators and relations; in fact, as shown by Dugger in [11], they

C. Casacuberta
Facultat de Matemàtiques i Informàtica, Universitat de Barcelona (UB),
Gran Via de les Corts Catalanes 585, 08007 Barcelona, Spain
e-mail: carles.casacuberta@ub.edu

J. Rosický (✉)
Department of Mathematics and Statistics, Faculty of Sciences, Masaryk
University, Kotlářská 2, 611 37 Brno, Czech Republic
e-mail: rosicky@math.muni.cz

© Springer Nature Singapore Pte Ltd. 2020
T. Ohsawa and N. Minami (eds.), *Bousfield Classes and Ohkawa’s Theorem*,
Springer Proceedings in Mathematics & Statistics 309,
https://doi.org/10.1007/978-981-15-1588-0_4

are Quillen equivalent to localizations of categories of simplicial presheaves with respect to sets of maps. Moreover, for each combinatorial model category \mathcal{K} there exist cardinals λ for which \mathcal{K} admits fibrant and cofibrant replacement functors that preserve λ -filtered colimits and λ -presentable objects, and the class of weak equivalences is closed under λ -filtered colimits [4, 11, 28].

Cofibrantly generated model categories admit weak generators [13, 26]. Combinatorial model categories are, in addition, well generated in the sense of [18, 21]. This fact links the study of combinatorial model categories with the theory of triangulated categories in useful ways. For instance, it was shown in [8] that localizing subcategories of triangulated categories with combinatorial models are coreflective assuming a large-cardinal axiom (Vopěnka’s principle), and similarly colocalizing subcategories are reflective.

In this article we show that a suitably restricted Yoneda embedding [1, 28] gives a way to implement Ohkawa’s argument [24] in the homotopy category of any combinatorial model category, not necessarily stable. Ohkawa’s original theorem becomes then a special case, since the homotopy category of spectra admits combinatorial models [15]. Thus we prove that, if \mathcal{K} is a pointed strongly λ -combinatorial model category (see Sect. 3 below for details) then there is only a set of distinct kernels of endofunctors $H: \mathcal{K} \rightarrow \mathcal{K}$ preserving λ -filtered colimits and the zero object.

This statement (and our method of proof) is a variant of the main result in [9], where Ohkawa’s theorem was broadly generalized. In independent work, Stevenson used abelian presheaves over compact objects to prove that Ohkawa’s theorem holds in compactly generated tensor triangulated categories [31], and Iyengar and Krause extended this result to well generated tensor triangulated categories [16].

Our approach shows that Ohkawa’s theorem is valid in the categories of motivic spaces and motivic spectra over any Noetherian base scheme of finite dimension [19], and also in categories of modules over (ordinary or motivic) ring spectra, since such categories have combinatorial models. Therefore, for example, Ohkawa’s theorem holds in the derived category of motives over any field k of characteristic zero, since these are modules over a motivic Eilenberg–Mac Lane spectrum [27].

2 Combinatorial Model Categories

The notion of a combinatorial model category was introduced by Jeff Smith in unpublished work made in the decade of 1990. The name refers to the fact that the underlying category and its model structure are both controlled by sets of sufficiently small objects and maps between them, in the precise sense that we next define. Further details and additional motivation can be found in [1, 4, 11–13].

For a regular cardinal λ , a small category \mathcal{A} is *λ -filtered* if every diagram in \mathcal{A} of cardinality smaller than λ has a cocone. An object A of a category \mathcal{C} is called *λ -presentable* if the hom-functor $\mathcal{C}(A, -): \mathcal{C} \rightarrow \mathbf{Set}$ preserves λ -filtered colimits. For example, a group (or a module over a ring) is λ -presentable if and only if it admits a presentation with less than λ generators and less than λ relations.

A category \mathcal{C} is *locally λ -presentable* if it is cocomplete and has a set \mathcal{A} of λ -presentable objects such that every object of \mathcal{C} is a λ -filtered colimit of objects of \mathcal{A} . A category is *locally presentable* if it is locally λ -presentable for some regular cardinal λ . The category of sets is locally \aleph_0 -presentable, since every set is the colimit of the inclusions of its finite subsets. As shown in [1, Corollary 3.7], every variety of finitary algebras is locally \aleph_0 -presentable. Many more examples arise from the fact that every functor category from a small category to a locally presentable category is locally presentable [1, Corollary 1.54].

A model category \mathcal{K} is *cofibrantly generated* if it has a set \mathcal{I} of cofibrations such that the trivial fibrations of \mathcal{K} are those morphisms having the right lifting property with respect to \mathcal{I} , and a set \mathcal{J} of trivial cofibrations such that the fibrations of \mathcal{K} are those morphisms having the right lifting property with respect to \mathcal{J} , and moreover \mathcal{I} and \mathcal{J} permit the small object argument, that is, their domains are small relative to transfinite compositions of pushouts of elements of \mathcal{I} and \mathcal{J} respectively. The category of simplicial sets is cofibrantly generated with \mathcal{I} the set of inclusions $\partial\Delta[n] \hookrightarrow \Delta[n]$ for $n \geq 0$ and \mathcal{J} the set of inclusions $\Delta^k[n] \hookrightarrow \Delta[n]$ for $n \geq 0$ and $0 \leq k \leq n$; see [12, 13] for notation and a proof.

A model category is called *combinatorial* if it is locally presentable and cofibrantly generated. By a *combinatorial homotopy category* we mean a homotopy category of a combinatorial model category.

Every locally presentable category \mathcal{C} can be viewed as a combinatorial homotopy category because the trivial model structure on \mathcal{C} (that is, the one in which every morphism is both a cofibration and a fibration, and the weak equivalences are the isomorphisms) is cofibrantly generated by the argument given in [30, Example 4.6]. In general, combinatorial homotopy categories are far from being locally presentable themselves, but they behave in some sense like a homotopy-theoretical version of those.

A model category \mathcal{K} is called *λ -combinatorial* for a regular cardinal λ if it is locally λ -presentable and cofibrantly generated by morphisms between λ -presentable objects. Then the functors giving factorizations of morphisms in \mathcal{K} into cofibrations followed by trivial fibrations or into trivial cofibrations followed by fibrations can be chosen to be λ -accessible, that is, preserving λ -filtered colimits. Details are given in [28, Proposition 3.1].

3 Restricted Yoneda Embedding

Let \mathcal{C} be a category and \mathcal{A} a small full subcategory of \mathcal{C} . The *restricted Yoneda embedding*

$$E_{\mathcal{A}} : \mathcal{C} \longrightarrow \mathbf{Set}^{\mathcal{A}^{\text{op}}}$$

sends every object X of \mathcal{C} to the hom-set $\mathcal{C}(-, X)$ restricted to \mathcal{A} . Thus $E_{\mathcal{A}}$ is full and faithful on morphisms whose domain is an object of \mathcal{A} .

The subcategory \mathcal{A} is called a *generator* of \mathcal{C} if $E_{\mathcal{A}}$ is faithful, and a *strong generator* if $E_{\mathcal{A}}$ is faithful and conservative, that is, reflecting isomorphisms. We say that \mathcal{A} is a *weak generator* if $E_{\mathcal{A}}$ reflects isomorphisms whose codomain is the terminal object of \mathcal{C} . This means that an object of \mathcal{C} is terminal whenever its image under $E_{\mathcal{A}}$ is terminal; hence the objects in a weak generator of \mathcal{C} form a *left weakly adequate* set in the sense of [26].

Recall from [13, 25] that if \mathcal{K} is a model category then its homotopy category $\text{Ho } \mathcal{K}$ is obtained by quotienting the full subcategory \mathcal{K}_{cf} of objects that are fibrant and cofibrant by the homotopy relation on morphisms. Each choice of a fibrant replacement functor R_f and a cofibrant replacement functor R_c on \mathcal{K} yields an essentially surjective functor

$$P: \mathcal{K} \longrightarrow \text{Ho } \mathcal{K}, \tag{1}$$

namely the composite $R_c R_f: \mathcal{K} \rightarrow \mathcal{K}_{cf}$ followed by the projection $\mathcal{K}_{cf} \rightarrow \text{Ho } \mathcal{K}$.

It was shown in [13, Theorem 7.3.1] that, if \mathcal{I} is a set of generating cofibrations in a pointed cofibrantly generated model category \mathcal{K} , then the cofibres of morphisms in \mathcal{I} form a weak generator of $\text{Ho } \mathcal{K}$. The assumption that \mathcal{K} be pointed can be removed if \mathcal{K} has a set \mathcal{I} of generating cofibrations between cofibrant objects, in which case the domains and codomains of morphisms in \mathcal{I} form a weak generator of $\text{Ho } \mathcal{K}$, as shown in [26, Theorem 1.2].

We also recall that a small full subcategory \mathcal{A} of a category \mathcal{C} is called *dense* if every object X in \mathcal{C} is a colimit of its canonical diagram with respect to \mathcal{A} . This is equivalent to $E_{\mathcal{A}}$ being full and faithful; see [1, Proposition 1.26]. Correspondingly, $E_{\mathcal{A}}$ is full if and only if \mathcal{A} is *weakly dense* in the sense that every object X is a weak colimit of its canonical diagram with respect to \mathcal{A} . Finally, $E_{\mathcal{A}}$ is full and conservative if and only if every X is a minimal weak colimit of its canonical diagram with respect to \mathcal{A} . Recall that a weak colimit $(\delta_d: Dd \rightarrow X)$ of a diagram $D: \mathcal{D} \rightarrow \mathcal{C}$ is called *minimal* if every morphism $f: X \rightarrow X$ such that $f \circ \delta_d = \delta_d$ for each $d \in \mathcal{D}$ is an isomorphism [10].

Theorem 3.1 *If \mathcal{K} is a combinatorial model category, then there exist arbitrarily large regular cardinals λ such that \mathcal{K} has the following properties:*

1. \mathcal{K} is locally λ -presentable.
2. There is a small weak generator of $\text{Ho } \mathcal{K}$ consisting of λ -presentable objects.
3. There are fibrant and cofibrant replacement functors R_f and R_c on \mathcal{K} that preserve λ -filtered colimits and λ -presentable objects.

Proof If \mathcal{K} is combinatorial, then, according to [11, Corollary 1.2], there is a zig-zag of Quillen equivalences into another combinatorial model category \mathcal{M} where all objects are cofibrant. Consequently, the domains and codomains of morphisms in a set of generating cofibrations for \mathcal{M} form a weak generator of the homotopy category $\text{Ho } \mathcal{M}$ by [26, Theorem 1.2]. Since the latter is equivalent to $\text{Ho } \mathcal{K}$, it follows that $\text{Ho } \mathcal{K}$ also has a small weak generator \mathcal{A} .

As \mathcal{K} is locally presentable, there are arbitrarily large regular cardinals μ such that \mathcal{K} is locally μ -presentable, by [1, Theorem 1.20]. Thus we can choose μ big enough so that \mathcal{K} is locally μ -presentable and cofibrantly generated by morphisms between μ -presentable objects, and, furthermore, the objects in the chosen weak generator \mathcal{A} are μ -presentable. Then, as shown in the proof of [28, Proposition 3.1], there are μ -accessible functors giving factorizations of morphisms in \mathcal{K} into cofibrations followed by trivial fibrations and into trivial cofibrations followed by fibrations. In particular we can pick a fibrant replacement functor R_f and a cofibrant replacement functor R_c that are μ -accessible. Moreover, using [1, Theorem 2.19] or [11, Proposition 7.2], we can pick a regular cardinal $\lambda \geq \mu$ such that R_f and R_c preserve both λ -filtered colimits and λ -presentable objects. \square

Definition 3.2 We call a model category \mathcal{K} *strongly λ -combinatorial* if it is combinatorial and λ satisfies the conditions stated in Theorem 3.1.

For a regular cardinal λ , let \mathcal{K} be a strongly λ -combinatorial model category and denote by \mathcal{K}_λ a small full subcategory of representatives of all isomorphism classes of λ -presentable objects. Here and in what follows we assume that fibrant and cofibrant replacement functors R_f and R_c have been chosen on \mathcal{K} so that they preserve λ -filtered colimits and λ -presentable objects.

Let $\text{Ho } \mathcal{K}_\lambda$ denote the full image of the composition

$$\mathcal{K}_\lambda \longleftarrow \mathcal{K} \xrightarrow{P} \text{Ho } \mathcal{K},$$

where P is the composite $R_c R_f$ followed by the canonical projection as in (1), and denote by $P_\lambda: \mathcal{K}_\lambda \rightarrow \text{Ho } \mathcal{K}_\lambda$ the domain and codomain restriction of P .

Consider the restricted Yoneda embedding

$$E_\lambda: \text{Ho } \mathcal{K} \longrightarrow \mathbf{Set}^{(\text{Ho } \mathcal{K}_\lambda)^{\text{op}}},$$

for which the composite $E_\lambda P$ preserves λ -presentable objects.

The next two results follow from [28, Proposition 5.1 and Corollary 5.2].

Theorem 3.3 *Let \mathcal{K} be a strongly λ -combinatorial model category for a regular cardinal λ . Then the composite*

$$\mathcal{K} \xrightarrow{P} \text{Ho } \mathcal{K} \xrightarrow{E_\lambda} \mathbf{Set}^{(\text{Ho } \mathcal{K}_\lambda)^{\text{op}}}$$

preserves λ -filtered colimits.

Corollary 3.4 *If \mathcal{K} is strongly λ -combinatorial, then $E_\lambda P \cong \text{Ind}_\lambda P_\lambda$.*

Here Ind_λ denotes free cocompletion with respect to λ -filtered colimits [1, Definition 2.25], so $\text{Ind}_\lambda P_\lambda$ is a functor from \mathcal{K} to $\text{Ind}_\lambda \text{Ho } \mathcal{K}_\lambda$. The statement of Corollary 3.4 means that E_λ factorizes through the inclusion

$$\mathrm{Ind}_\lambda \mathrm{Ho} \mathcal{K}_\lambda \subseteq \mathbf{Set}^{(\mathrm{Ho} \mathcal{K}_\lambda)^{\mathrm{op}}},$$

and its codomain restriction, which we keep denoting by E_λ , makes the composite $E_\lambda P$ isomorphic to $\mathrm{Ind}_\lambda P_\lambda$.

If the model category \mathcal{K} is pointed, then $\mathrm{Ind}_\lambda \mathrm{Ho} \mathcal{K}_\lambda$ is also pointed and E_λ preserves the zero object 0 , since $E_\lambda 0$ is terminal and it is also initial because 0 is λ -presentable and E_λ is full and faithful on morphisms with domain in $\mathrm{Ho} \mathcal{K}_\lambda$.

Corollary 3.5 *If \mathcal{K} is a strongly λ -combinatorial model category, the codomain restriction $E_\lambda: \mathrm{Ho} \mathcal{K} \rightarrow \mathrm{Ind}_\lambda \mathrm{Ho} \mathcal{K}_\lambda$ preserves coproducts.*

Proof Pick a cofibrant replacement functor R_c preserving λ -filtered colimits and λ -presentable objects. Note that P preserves coproducts between cofibrant objects and every object in $\mathrm{Ho} \mathcal{K}$ is isomorphic to PX for some cofibrant object X in \mathcal{K} . Hence, using Corollary 3.4 it suffices to show that $\mathrm{Ind}_\lambda P_\lambda$ preserves coproducts between cofibrant objects. Since each coproduct is a λ -filtered colimit of λ -small coproducts and $\mathrm{Ind}_\lambda P_\lambda$ preserves λ -filtered colimits, we have to prove that $\mathrm{Ind}_\lambda P_\lambda$ preserves λ -small coproducts between cofibrant objects. Let $\coprod_{i \in I} K_i$ be such a coproduct, so that the cardinality of I is smaller than λ . Since the functor R_c preserves λ -filtered colimits and λ -presentable objects, each K_i is a λ -filtered colimit of cofibrant λ -presentable objects. Let $D_i: \mathcal{D}_i \rightarrow \mathcal{K}_\lambda$ denote the corresponding diagrams, so that $K_i \cong \mathrm{colim} D_i$. Then $\coprod_{i \in I} K_i$ is a colimit of a λ -filtered diagram whose values are coproducts $\coprod_{i \in I} D_i d_i$ with $d_i \in \mathcal{D}_i$, and each such coproduct $\coprod_{i \in I} D_i d_i$ is λ -presentable as the cardinality of I is smaller than λ . Since the functor $\mathrm{Ind}_\lambda P_\lambda$ preserves λ -filtered colimits and P_λ preserves λ -small coproducts of cofibrant objects, the result is proved. \square

Definition 3.6 Let \mathcal{C} be a category with coproducts and λ a cardinal. An object S of \mathcal{C} is λ -small if for every morphism $f: S \rightarrow \coprod_{i \in I} X_i$ there is a subset J of I of cardinality less than λ such that f factorizes as

$$S \longrightarrow \coprod_{j \in J} X_j \longrightarrow \coprod_{i \in I} X_i,$$

where the second morphism is the subcoproduct injection.

We also say that \aleph_0 -small objects are *compact*. This terminology is due to Neeman [21], who found how compactness should be defined for uncountable cardinals in triangulated categories. His definition was simplified by Krause in [18]. They considered compactness in additive categories but the definition makes sense in general. Consider classes \mathcal{S} of λ -small objects in a category \mathcal{C} with coproducts such that for every morphism $f: S \rightarrow \coprod_{i \in I} X_i$ with $S \in \mathcal{S}$ there exist morphisms $g_i: S_i \rightarrow X_i$ for which $S_i \in \mathcal{S}$ for all $i \in I$ and f factorizes through

$$\coprod_{i \in I} g_i: \coprod_{i \in I} S_i \longrightarrow \coprod_{i \in I} X_i.$$

Since the collection of such classes is closed under unions, there is a greatest class with this property. Its objects are called λ -compact.

Proposition 3.7 *If \mathcal{K} is a strongly λ -combinatorial model category, then all objects in $\text{Ho } \mathcal{K}_\lambda$ are λ -compact in $\text{Ho } \mathcal{K}$.*

Proof Choose fibrant and cofibrant replacement functors R_f and R_c preserving λ -filtered colimits and λ -presentable objects, and let $P : \mathcal{K} \rightarrow \text{Ho } \mathcal{K}$ be as in (1). Suppose given a morphism $g : PA \rightarrow \coprod_{i \in I} PK_i$ in $\text{Ho } \mathcal{K}$ where A is in \mathcal{K}_λ . According to Corollary 3.5, we have

$$E_\lambda g : E_\lambda PA \longrightarrow \coprod_{i \in I} E_\lambda PK_i.$$

Due to the fact that $E_\lambda P$ preserves λ -presentable objects, $E_\lambda PA$ is λ -presentable in $\text{Ind}_\lambda \text{Ho } \mathcal{K}_\lambda$. Since each coproduct is a λ -filtered colimit of λ -small subcoproducts, $E_\lambda g$ factorizes through some $\coprod_{j \in J} E_\lambda PK_j$ where J has cardinality smaller than λ . Since E_λ is full and faithful on morphisms with domain in $\text{Ho } \mathcal{K}_\lambda$, we obtain a factorization of g through $\coprod_{j \in J} PK_j$ and therefore we conclude that PA is λ -small.

Moreover, the argument used in the proof of Corollary 3.5 shows in a similar way that $E_\lambda g$ factors through some coproduct $\coprod_{j \in J} E_\lambda PD_j d_j$ where J has cardinality smaller than λ and $D_j d_j$ is in \mathcal{K}_λ for all j . Using again the fact that E_λ is full and faithful on morphisms with domain in $\text{Ho } \mathcal{K}_\lambda$, we find a factorization of g through $\coprod_{j \in J} PD_j d_j$. Hence PA is indeed λ -compact. \square

Definition 3.8 A category with coproducts is called *well λ -generated* if it has a small weak generator consisting of λ -compact objects. It is called *well generated* if it is well λ -generated for some λ .

For example, every locally λ -presentable category is well λ -generated.

The following result was proved in [28, Proposition 6.10] with the additional assumption that \mathcal{K} was stable, which is not necessary.

Theorem 3.9 *If \mathcal{K} is a strongly λ -combinatorial model category, then $\text{Ho } \mathcal{K}$ is well λ -generated.*

Proof Since, by assumption, there is a small weak generator of $\text{Ho } \mathcal{K}$ whose objects are λ -presentable, $\text{Ho } \mathcal{K}_\lambda$ weakly generates $\text{Ho } \mathcal{K}$. The rest follows from Proposition 3.7. \square

As a corollary one infers Neeman's result in [22] that, for any Grothendieck abelian category \mathcal{A} , the derived category $D(\mathcal{A})$ is well generated.

4 Ohkawa's Theorem

For an endofunctor $H : \mathcal{K} \rightarrow \mathcal{K}$ (not necessarily preserving weak equivalences) on a model category \mathcal{K} , we consider the composition

$$\mathcal{K} \xrightarrow{H} \mathcal{K} \xrightarrow{P} \text{Ho } \mathcal{K},$$

where P is defined as in (1). The class of objects X in \mathcal{K} such that PHX is the terminal object in $\text{Ho } \mathcal{K}$ will be called the *kernel* of H and will be denoted by $\ker H$. Hence, if \mathcal{K} is pointed and 0 denotes the zero object in \mathcal{K} and also its image in $\text{Ho } \mathcal{K}$, then $\ker H$ consists of objects X in \mathcal{K} such that $PHX = 0$.

In this section we prove the following result.

Theorem 4.1 *Suppose that \mathcal{K} is a pointed strongly λ -combinatorial model category. Then there is only a set of distinct kernels of endofunctors $H: \mathcal{K} \rightarrow \mathcal{K}$ preserving λ -filtered colimits and the zero object.*

Proof Consider the restricted Yoneda embedding as given by Corollary 3.4,

$$E_\lambda: \text{Ho } \mathcal{K} \longrightarrow \text{Ind}_\lambda \text{Ho } \mathcal{K}_\lambda.$$

For a morphism $f: E_\lambda S \rightarrow E_\lambda PHA$ with $A \in \mathcal{K}_\lambda$ and $S \in \text{Ho } \mathcal{K}_\lambda$, let us denote by $T_H(f)$ the set of all morphisms $t: A \rightarrow B$ in \mathcal{K}_λ such that the composite

$$E_\lambda S \xrightarrow{f} E_\lambda PHA \xrightarrow{E_\lambda PHt} E_\lambda PHB$$

is the zero morphism, that is, $E_\lambda PHt \circ f$ factors through the zero object.

Next, we denote

$$J(H) = \{T_H(f) \mid f: E_\lambda S \rightarrow E_\lambda PHA \text{ with } A \in \mathcal{K}_\lambda \text{ and } S \in \text{Ho } \mathcal{K}_\lambda\}.$$

We are going to prove that if $J(H_1) = J(H_2)$ then $\ker H_1 = \ker H_2$, assuming that H_1 and H_2 preserve λ -filtered colimits and the zero object. Thus suppose that $J(H_2) \subseteq J(H_1)$ and let $X \in \ker H_1$. In order to prove that $PH_2X = 0$, it is enough to show that every morphism $E_\lambda G \rightarrow E_\lambda PH_2X$ factors through the zero object if G is in $\text{Ho } \mathcal{K}_\lambda$, since $\text{Ho } \mathcal{K}_\lambda$ is a weak generator of $\text{Ho } \mathcal{K}$ and E_λ is full and faithful on morphisms whose domain is in $\text{Ho } \mathcal{K}_\lambda$.

Assume given such a morphism $f: E_\lambda G \rightarrow E_\lambda PH_2X$. Since the category \mathcal{K} is locally λ -presentable, $X \cong \text{colim}(D: \mathcal{D} \rightarrow \mathcal{K}_\lambda)$ for a certain λ -filtered diagram D . Since $E_\lambda PH_2$ preserves λ -filtered colimits by Theorem 3.3, we then have

$$E_\lambda PH_2X \cong \text{colim} \left(\mathcal{D} \xrightarrow{D} \mathcal{K}_\lambda \xrightarrow{PH_2} \text{Ho } \mathcal{K} \xrightarrow{E_\lambda} \text{Ind}_\lambda \text{Ho } \mathcal{K}_\lambda \right).$$

Since $E_\lambda G$ is λ -presentable, f factors through $\hat{f}: E_\lambda G \rightarrow E_\lambda PH_2Dd$ for some $d \in \mathcal{D}$. Note that the set $T_{H_2}(\hat{f})$ is nonempty, since the morphism $Dd \rightarrow 0$ is in it as H_2 preserves the zero object. Consequently, the assumption that $J(H_2) \subseteq J(H_1)$ implies that $T_{H_2}(\hat{f}) \in J(H_1)$. This means that there exist an object $V \in \text{Ho } \mathcal{K}_\lambda$ and a morphism $g: E_\lambda V \rightarrow E_\lambda PH_1Dd$ such that $T_{H_2}(\hat{f}) = T_{H_1}(g)$.

Now, since $X \in \ker H_1$, we have $E_\lambda PH_1X = 0$. However,

$$E_\lambda PH_1 X \cong \operatorname{colim} \left(\mathcal{D} \xrightarrow{D} \mathcal{K}_\lambda \xrightarrow{PH_1} \operatorname{Ho} \mathcal{K} \xrightarrow{E_\lambda} \operatorname{Ind}_\lambda \operatorname{Ho} \mathcal{K}_\lambda \right),$$

and, since $E_\lambda V$ is λ -presentable, there is a morphism $\delta: d \rightarrow d'$ in \mathcal{D} such that

$$E_\lambda V \xrightarrow{g} E_\lambda PH_1 Dd \xrightarrow{E_\lambda PH_1 D\delta} E_\lambda PH_1 Dd'$$

factors through the zero object. Hence $D\delta \in T_{H_1}(g)$. Therefore $D\delta \in T_{H_2}(\hat{f})$ and this implies that $f: E_\lambda G \rightarrow E_\lambda PH_2 X$ factors through the zero object, as we wanted to show.

Finally, since there is only a set of distinct sets $J(H)$, the theorem is proved. \square

Ohkawa's theorem [24, Theorem 2] is a special case of Theorem 4.1. Recall that two (reduced) homology theories E_* and F_* on spectra are said to be *Bousfield equivalent* if the class of E_* -acyclic spectra coincides with the class of F_* -acyclic spectra. A spectrum X is called *E_* -acyclic* if $E_*(X) = 0$.

Corollary 4.2 *There is only a set of Bousfield equivalence classes of representable homology theories on spectra.*

Proof The homotopy category of spectra admits a combinatorial model category \mathcal{K} ; for instance, symmetric spectra over simplicial sets [15]. For each cofibrant spectrum E we consider the endofunctor on \mathcal{K} defined as $H_E X = E \wedge R_c X$ where R_c is a cofibrant replacement functor preserving filtered colimits. Since smashing with E has a right adjoint, H_E preserves filtered colimits. Moreover, a spectrum X is in $\ker H_E$ if and only if X is E_* -acyclic. Hence Theorem 4.1 implies that there is only a set of distinct kernels of endofunctors of the form H_E . \square

5 Generalized Brown Representability

In this section we prove other properties of combinatorial homotopy categories related to results in [28]. Note that if \mathcal{C} is a locally λ -presentable category with the trivial model structure then the functor $E_\lambda: \mathcal{C} \rightarrow \operatorname{Ind}_\lambda \mathcal{C}_\lambda$ is an isomorphism.

Definition 5.1 A strongly λ -combinatorial model category \mathcal{K} is called *λ -Brown on morphisms* if $E_\lambda: \operatorname{Ho} \mathcal{K} \rightarrow \operatorname{Ind}_\lambda \operatorname{Ho} \mathcal{K}_\lambda$ is full. It is called *λ -Brown on objects* if E_λ is essentially surjective. Finally, \mathcal{K} is called *λ -Brown* if it is λ -Brown both on objects and on morphisms.

Let us remark the following facts:

- (i) A locally finitely presentable stable combinatorial model category is ω -Brown if it is Brown in the sense of [14], where ω denotes the first infinite ordinal.

- (ii) Whenever \mathcal{K} is strongly ω -combinatorial and E_ω is full then E_ω is essentially surjective as well. In fact, by Corollary 3.4, $\text{Ind}_\omega P_\omega$ is full; since each object of $\text{Ind}_\omega \mathcal{K}_\omega$ can be obtained by taking successive colimits of smooth chains [1] and P_ω is essentially surjective on objects, $\text{Ind}_\omega P_\omega$ is essentially surjective on objects too. Hence \mathcal{K} is ω -Brown on objects. This argument does not work for $\lambda > \omega$ because, in the proof, we need colimits of chains of cofinality ω .
- (iii) E_λ is full if and only if $\text{Ho } \mathcal{K}_\lambda$ is weakly dense in $\text{Ho } \mathcal{K}$.

The homotopy category $\text{Ho } \mathcal{K}$ of any model category \mathcal{K} has weak colimits and weak limits. Weak colimits are constructed from coproducts and homotopy pushouts in the same way as colimits are constructed from coproducts and pushouts. A homotopy pushout of

$$PC \xleftarrow{Pg} PA \xrightarrow{Pf} PB$$

is a commutative diagram

$$\begin{array}{ccc} PA & \xrightarrow{Pf_1} & PB_1 \\ \downarrow Pg_1 & & \downarrow P\bar{g} \\ PC_1 & \xrightarrow{P\bar{f}} & PE \end{array}$$

where $f = f_2 \circ f_1$ and $g = g_2 \circ g_1$ are factorizations of f and g , respectively, into a cofibration followed by a trivial fibration, and

$$\begin{array}{ccc} A & \xrightarrow{f_1} & B_1 \\ \downarrow g_1 & & \downarrow \bar{g} \\ C_1 & \xrightarrow{\bar{f}} & E \end{array}$$

is a pushout in \mathcal{K} . The following definition is taken from [5].

Definition 5.2 A functor $H: \mathcal{C} \rightarrow \mathcal{D}$ will be called *nearly full* if for each commutative triangle

$$\begin{array}{ccc} HA & \xrightarrow{Hh} & HC \\ & \searrow f & \nearrow Hg \\ & & HB \end{array}$$

there is a morphism $\bar{f}: A \rightarrow B$ in \mathcal{C} such that $H\bar{f} = f$.

Proposition 5.3 *A strongly λ -combinatorial model category \mathcal{K} is λ -Brown on morphisms if and only if the functor $E_\lambda: \text{Ho } \mathcal{K} \rightarrow \text{Ind}_\lambda \text{Ho } \mathcal{K}_\lambda$ is nearly full.*

Proof Sufficiency is evident because any full functor is nearly full. Let \mathcal{K} be a strongly λ -combinatorial model category and assume that E_λ is nearly full. Consider an object K in \mathcal{K} and express it as a λ -filtered colimit $(\delta_d: Dd \rightarrow K)$ of its canonical diagram $D: \mathcal{D} \rightarrow \mathcal{K}_\lambda$. This means that we have

$$\begin{array}{ccccc}
 & Dd & & & \\
 & \downarrow u_e & \searrow v_d & & \\
 \coprod_{e: d \rightarrow d'} Dd & \xrightarrow[p]{q} & \coprod_d Dd & \xrightarrow{g} & K \\
 \uparrow u_e & & \uparrow v_{d'} & & \\
 Dd & \xrightarrow{D_e} & Dd' & &
 \end{array}$$

where g is given by a pushout

$$\begin{array}{ccc}
 \coprod_d Dd & \xrightarrow{g} & K \\
 \uparrow (p, \text{id}) & & \uparrow g \\
 \left(\coprod_e Dd \right) \amalg \left(\coprod_d Dd \right) & \xrightarrow{(q, \text{id})} & \coprod_d Dd.
 \end{array}$$

If we replace the pushout above by a homotopy pushout, we get $(\bar{\delta}_d: Dd \rightarrow \bar{K})$. It is not a cocone in \mathcal{K} but $(P\bar{\delta}_d: PDd \rightarrow P\bar{K})$ is a standard weak colimit [10] in $\text{Ho } \mathcal{K}$, and there is a comparison morphism $t: \bar{K} \rightarrow K$ such that $t \circ \bar{\delta}_d = \delta_d$ for each d . Since $H_\lambda = \text{Ind}_\lambda P_\lambda$ preserves λ -filtered colimits, there is a morphism $u: H_\lambda K \rightarrow H_\lambda \bar{K}$ such that $u \circ H_\lambda \delta_d = H_\lambda \bar{\delta}_d$ for each d . Then $H_\lambda t \circ u = \text{id}$ because

$$H_\lambda t \circ u \circ H_\lambda \delta_d = H_\lambda (t \circ \bar{\delta}_d) = H_\lambda \delta_d.$$

Now, since E_λ is nearly full, there is $\bar{u}: PK \rightarrow P\bar{K}$ such that $u = E_\lambda \bar{u}$.

Consider a morphism $h: H_\lambda K_1 \rightarrow H_\lambda K_2$. Let u_1, t_1, u_2, t_2 be as u, t above for K_1 and K_2 . There is a cocone $(\gamma_d: PD_1 d \rightarrow P\bar{K}_2)$ from PD_1 such that

$$E_\lambda \gamma_d = u_2 \circ h \circ H_\lambda \delta_{1d}: H_\lambda D_1 d \rightarrow H_\lambda \bar{K}_2$$

for each d in \mathcal{D}_1 . Thus there is a morphism $\bar{h}: \bar{K}_1 \rightarrow \bar{K}_2$ such that $\bar{h} \circ P\bar{\delta}_{1d} = \gamma_d$ for each d in \mathcal{D}_1 . Hence

$$E_\lambda \bar{h} \circ u_1 \circ H_\lambda \delta_{1d} = E_\lambda \bar{h} \circ H_\lambda \bar{\delta}_{1d} = E_\lambda \gamma_d = u_2 \circ h \circ H_\lambda \delta_{1d}$$

for each d in \mathcal{D}_1 . Thus $E_\lambda \bar{h} \circ u_1 = u_2 \circ h$. Putting $h' = Pt_2 \circ \bar{h} \circ \bar{u}_1$, we obtain

$$E_\lambda h' = E_\lambda (Pt_2 \circ \bar{h}) \circ u_1 = H_\lambda t_2 \circ u_2 \circ h = h,$$

which proves that E_λ is full. \square

Remark 5.4 In Proposition 5.3 it suffices to assume that E_λ is full on split monomorphisms. This means that $h = \text{id}$ in Definition 5.2.

The proof of the following result is given in [28, Proposition 6.4].

Proposition 5.5 *If \mathcal{K} is a combinatorial stable model category, then E_λ reflects isomorphisms for arbitrarily large regular cardinals λ .*

Remark 5.6 If E_λ is full and reflects isomorphisms then each object of $\text{Ho } \mathcal{K}$ is a minimal weak colimit of its canonical diagram with respect to $\text{Ho } \mathcal{K}_\lambda$.

One could ask if every combinatorial stable model category is λ -Brown for arbitrarily large regular cardinals λ , as discussed in [28, 29]. This fact would have important consequences [23], but it is unfortunately not true. The first counterexample was given in [7], and in [3] a large class was found of combinatorial stable model categories which are not λ -Brown for any λ . An obstruction theory for generalized Brown representability in triangulated categories was developed in [20], with special focus on derived categories of rings.

Acknowledgements This article has been written as a contribution to the proceedings of the memorial conference for Professor Tetsusuke Ohkawa held at the University of Nagoya in 2015. The content of Sect. 4 is based on previous joint work of the authors with Javier Gutiérrez published in [9]. We also appreciate useful discussions with George Raptis. The authors were supported by the Grant Agency of the Czech Republic under grant P201/12/G028, the Agency for Management of University and Research Grants of Catalonia with references 2014SGR 114 and 2017 SGR 585, and the Spanish Ministry of Economy and Competitiveness under AEI/FEDER research grants MTM2013-42178-P and MTM2016-76453-C2-2-P, as well as grant MDM-2014-0445 awarded to the Barcelona Graduate School of Mathematics.

References

1. Adámek, J., Rosický, J.: Locally Presentable and Accessible Categories. London Mathematical Society Lecture Note Series, vol. 189. Cambridge University Press, Cambridge (1994)
2. Barwick, C.: On left and right model categories and left and right Bousfield localizations. Homol. Homotopy Appl. **12**, 245–320 (2010)

3. Bazzoni, S., Šťovíček, J.: On the abelianization of derived categories and a negative solution to Rosický's problem. *Compos. Math.* **149**, 125–147 (2013)
4. Beke, T.: Sheafifiable homotopy model categories. *Math. Proc. Cambridge Philos. Soc.* **129**, 447–475 (2000)
5. Beke, T., Rosický, J.: Abstract elementary classes and accessible categories. *Ann. Pure Appl. Logic* **163**, 2008–2017 (2012)
6. Bousfield, A.K.: Homotopical localizations of spaces. *Am. J. Math.* **119**, 1321–1354 (1997)
7. Braun, G., Göbel, R.: Splitting kernels into small summands. *Israel J. Math.* **188**, 221–230 (2012)
8. Casacuberta, C., Gutiérrez, J.J., Rosický, J.: Are all localizing subcategories of stable homotopy categories coreflective? *Adv. Math.* **252**, 158–184 (2014)
9. Casacuberta, C., Gutiérrez, J.J., Rosický, J.: A generalization of Ohkawa's theorem. *Compos. Math.* **150**, 893–902 (2014)
10. Christensen, J.D.: Ideals in triangulated categories: phantoms, ghosts and skeleta. *Adv. Math.* **136**, 284–339 (1998)
11. Dugger, D.: Combinatorial model categories have presentations. *Adv. Math.* **164**, 177–201 (2001)
12. Hirschhorn, P.S.: *Model Categories and Their Localizations*. Mathematical Surveys and Monographs, vol. 99. American Mathematical Society, Providence (2003)
13. Hovey, M.: *Model Categories*. Mathematical Surveys and Monographs, vol. 63. American Mathematical Society, Providence (1999)
14. Hovey, M., Palmieri, J.H., Strickland, N.P.: *Axiomatic Stable Homotopy Theory*. Memoirs of the American Mathematical Society, vol. 128, no. 610. American Mathematical Society, Providence (1997)
15. Hovey, M., Shipley, B., Smith, J.H.: Symmetric spectra. *J. Am. Math. Soc.* **13**, 149–208 (2000)
16. Iyengar, S.B., Krause, H.: The Bousfield lattice of a triangulated category and stratification. *Math. Z.* **273**, 1215–1241 (2013)
17. Kan, D.M.: On c.s.s. complexes. *Am. J. Math.* **79**, 449–476 (1957)
18. Krause, H.: On Neeman's well generated triangulated categories. *Doc. Math.* **6**, 121–126 (2001)
19. Morel, F., Voevodsky, V.: \mathbb{A}^1 -homotopy theory of schemes. *Publ. Math. IHÉS* **90**, 45–143 (1999)
20. Muro, F., Raventós, O.: Transfinite Adams representability. *Adv. Math.* **292**, 111–180 (2016)
21. Neeman, A.: *Triangulated Categories*. Annals of Mathematics Studies, vol. 148. Princeton University Press, Princeton (2001)
22. Neeman, A.: On the derived category of sheaves on a manifold. *Doc. Math.* **6**, 483–488 (2001)
23. Neeman, A.: Brown representability follows from Rosický's theorem. *J. Topol.* **2**, 262–276 (2009)
24. Ohkawa, T.: The injective hull of homotopy types with respect to generalized homology functors. *Hiroshima Math. J.* **19**, 631–639 (1989)
25. Quillen, D.: *Homotopical Algebra*. Lecture Notes in Mathematics, vol. 43. Springer, Berlin (1967)
26. Raptis, G.: On the cofibrant generation of model categories. *J. Homotopy Relat. Struct.* **4**, 245–253 (2009)
27. Röndigs, O., Østvær, P.A.: Modules over motivic cohomology. *Adv. Math.* **219**, 689–727 (2008)
28. Rosický, J.: Generalized Brown representability in homotopy categories. *Theory Appl. Categ.* **14**, 451–479 (2005)
29. Rosický, J.: Generalized Brown representability in homotopy categories: Erratum. *Theory Appl. Categ.* **20**, 18–24 (2008)
30. Rosický, J.: On combinatorial model categories. *Appl. Categ. Struct.* **17**, 303–316 (2009)
31. Stevenson, G.: An extension of Dwyer's and Palmieri's proof of Ohkawa's theorem on Bousfield classes (2011) (Unpublished manuscript)
32. Whitehead, J.H.C.: Combinatorial homotopy; Part I. *Bull. Am. Math. Soc.* **55**, 213–245 (1949), Part II. *Bull. Am. Math. Soc.* **55**, 453–496 (1949)