

## RANK ESTIMATES IN A CLASS OF SEMIPARAMETRIC TWO-SAMPLE MODELS\*

DOROTA M. DABROWSKA<sup>1</sup>, KJELL A. DOKSUM<sup>1\*\*</sup> AND RYOZO MIURA<sup>2</sup>

<sup>1</sup>*Department of Statistics, University of California, Berkeley, CA 94720, U.S.A.*

<sup>2</sup>*Osaka City University, Sumiyoshi-ku, Osaka 558, Japan*

(Received December 16, 1987; revised July 14, 1988)

**Abstract.** We consider a two-sample semiparametric model involving a real parameter  $\theta$  and a nuisance parameter  $F$  which is a distribution function. This model includes the proportional hazard, proportional odds, linear transformation and Harrington-Fleming models (1982, *Biometrika*, **69**, 533-546). We propose two types of estimates based on ranks. The first is a rank approximation to Huber's  $M$ -estimates (1981, *Robust Statistics*, Wiley) and the second is a Hodges-Lehmann type rank inversion estimate (1963, *Ann. Math. Statist.*, **34**, 598-611). We obtain asymptotic normality and efficiency results. The estimates are consistent and asymptotically normal generally but fully efficient only for special cases.

*Key words and phrases:* Semiparametric transformation models,  $M$ -estimates based on ranks, Hodges-Lehmann estimates.

### 1. Introduction

We consider the two sample problem where  $X_1, \dots, X_m$  and  $Y_1, \dots, Y_n$  are independent random samples from populations with continuous distribution functions  $F$  and  $G$ , respectively. Many of the models in which rank (partial, marginal) likelihood methods are useful can be put in the form

$$(1.1) \quad F(t) = D(H(t), \theta_1), \quad G(t) = D(H(t), \theta_2),$$

where  $H(t)$  is an unknown continuous distribution function,  $D(u, \theta)$  is a known continuous distribution function on  $(0, 1)$ , and  $\theta_1$  and  $\theta_2$  are in some parameter set  $\Theta$ .

---

\*Research partially supported by National Science Foundation Grant DMS-86-02083 and National Institute of General Medical Sciences Grant SSS-Y1RO1GM35416-01.

\*\*Now visiting professor at Department of Statistics, Harvard University, Cambridge, MA 02138, U.S.A.

For inference based on rank likelihood, the above model is equivalent to the model obtained by using the distributions of  $U_i = F(X_i) = D(H(X_i), \theta_1)$  and  $V_j = F(Y_j) = D(H(Y_j), \theta_1)$ . These distributions are

$$\tilde{F}(u) = u, \quad u \in (0, 1), \quad \tilde{G}(v) = D(D^{-1}(v, \theta_1), \theta_2), \quad v \in (0, 1).$$

In the case where  $\{D(u, \theta): \theta \in \Theta\}$  is a group under composition satisfying  $D(u, 1) = u$ ,  $D(D^{-1}(u, \theta_1), \theta_2) = D(u, \theta)$ ,  $\theta = \theta_2/\theta_1$ , we can write

$$(1.2) \quad \tilde{F}(u) = u, \quad u \in (0, 1), \quad \tilde{G}(v) = D(v, \theta), \quad v \in (0, 1).$$

From this point on we assume that (1.2) is satisfied. The distribution function  $F$  is treated as a nuisance parameter and we consider the problem of estimating  $\theta$ . This model goes back to Lehmann (1953), and includes the following models that have important applications in survival analysis, reliability, and other areas.

*Example 1.1. (Proportional hazard model)* If  $F$  and  $G$  have proportional hazards, then  $D(v, \theta) = 1 - [1 - v]^{1/\theta}$ ,  $\theta > 0$ . Lehmann (1953) and Savage (1956) considered testing in this model. Cox (1972, 1975) developed estimation procedures in a much more general regression problem with censored data.

*Example 1.2. (Proportional odds model)* For any continuous distribution  $H$  the odds rate is defined by  $r_H = H/(1 - H)$ . If  $F$  and  $G$  have proportional odds rates, in the sense that  $r_G(t) = \theta^{-1}r_F(t)$ , then  $D(v, \theta) = v[(1 - v)\theta + v]^{-1}$ . This model has been considered by Ferguson (1967) and Bickel (1986) in the two-sample case and in more general regression models by Bennett (1983) and Pettitt (1984), among others.

*Example 1.3. (Proportional  $\gamma$ -odds model)* For any continuous distribution function  $H$ , the  $\gamma$ -odds rate is defined by

$$r_{H,\gamma}(t) = \begin{cases} \{[1 - H(t)]^{-\gamma} - 1\}/\gamma, & \gamma > 0, \\ -\log [1 - H(t)], & \gamma = 0. \end{cases}$$

If  $F$  and  $G$  have proportional  $\gamma$ -odds, then

$$D(v, \theta) = \begin{cases} 1 - \left\{ \frac{\theta(1 - v)^\gamma}{1 - (1 - v)^\gamma + \theta(1 - v)^\gamma} \right\}^{1/\gamma}, & \gamma > 0, \\ 1 - (1 - v)^{1/\theta}, & \gamma = 0. \end{cases}$$

This model, which has been considered by Harrington and Fleming (1982),

Clayton and Cuzick (1986), Bickel (1986) and Dabrowska and Doksum (1988), reduces to Example 1.1 when  $\gamma = 0$ , and to Example 1.2 when  $\gamma = 1$ .

*Example 1.4. (Transformation shift model)* Let  $Q$  be any known continuous distribution function which is strictly increasing on the whole real line. If  $F$  and  $G$  satisfy the transformation shift model where for some increasing transformation  $h$ ,  $X_i$  and  $Y_j$  can be written

$$(1.3) \quad \begin{aligned} h(X_i) &= \mu_1 + \varepsilon_i, & i &= 1, \dots, m, \\ h(Y_j) &= \mu_2 + \varepsilon_j, & j &= 1, \dots, n, \end{aligned}$$

with  $\mu_2 - \mu_1 = \log \theta$ ,  $N = m + n$  and  $\varepsilon_1, \dots, \varepsilon_N$  independent with distribution function  $Q$ , then  $D(v, \theta) = Q(Q^{-1}(v) - \log \theta)$ . To see this, set  $h(t) = Q^{-1}(F(t)) + \mu_1$ . Then (1.3) is equivalent to (1.2) with  $D(v, \theta) = Q(Q^{-1}(v) - \log \theta)$ . This is an extension of the power transformation model where  $h(t)$  is of the form  $\text{sign}(t)|t|^\lambda$  or  $[\text{sign}(t)|t|^\lambda - 1]/\lambda$ , and where  $Q$  is the standard normal distribution function (see Anscombe and Tukey (1954), Tukey (1957), Box and Cox (1964), Bickel and Doksum (1981) and Doksum (1987)). This transformation shift model reduces to the proportional odds model if we take  $Q$  to be the logistic distribution function  $L(x) = 1/[1 + e^{-x}]$ .

Theory and methods for dealing with semiparametric models and partial likelihood have been developed by Begun (1981), Begun *et al.* (1983), Begun and Wellner (1983), Wellner (1986) and Wong (1986), among others. However, these methods do not lead to tractable efficient scores or tractable efficient estimates for any of the above models except the proportional hazard model. For arbitrary  $\theta_0$  and under certain regularity conditions, Bickel (1986) obtained the asymptotically optimal rank test for testing  $H_0: \theta = \theta_0$ , vs  $H_1: \theta > \theta_0$ . His regularity conditions are satisfied by the  $\gamma$ -odds model with  $\gamma \geq 1$ . However, the optimal test statistic is a non-linear rank statistic whose value can be obtained only after solving certain functional equations numerically on the computer. Attempts to extend these methods to obtain estimates that are asymptotically efficient in the semiparametric sense have not yet succeeded. Doksum (1987), and Dabrowska and Doksum (1988) propose a resampling scheme for computing maximum partial (rank) likelihood estimators in general semiparametric transformation regression models with censored data. These estimates perform well in Monte Carlo simulation studies, but the theoretical properties are difficult to establish. Clayton and Cuzick (1986) proposed estimates for the proportional  $\gamma$ -odds model that also apply to regression and censored data, but the theoretical properties of these estimates are also not well understood.

Methods based on local (near  $\theta = 1$ ) approximations to the likelihood have been developed by Pettitt (1984), Doksum (1987), and Dabrowska and Doksum (1988). These estimates are asymptotically normal for  $\theta$  in neighbourhoods of  $\theta = 1$ , however, they are not consistent for fixed  $\theta \neq 1$ .

In this paper we consider the uncensored case and introduce two classes of estimates. The advantage of these estimates is that they are relatively simple to implement in practice and their properties can readily be derived and understood. Moreover, they are based on intuitive estimation equations that are immediate extensions of the familiar  $M$  and  $R$  estimate equations. Although the new estimates are not fully efficient for all values of  $\theta$ , such fully efficient estimates are not available for  $\gamma \neq 0$ .

In Section 2 we introduce estimates of  $\theta$  that can be regarded as rank approximations to Huber's (1981)  $M$ -estimates based on score function  $\psi$ . We show asymptotic normality of these RAM (Rank Approximate  $M$ ) estimates. We compare these estimates with the asymptotically optimal estimates for  $F$  known, and find that for a certain range of parameter values, not much efficiency is lost. In fact, the score function  $\psi$  can be chosen so that the estimate is fully efficient at  $\theta = 1$ .

In Section 3 we introduce estimates of  $\theta$  based on the Hodges-Lehmann (1963) rank inversion idea and obtain asymptotic normality of these estimates. For appropriate choices of the score functions, these estimates have the same asymptotic distribution as the RAM estimates.

## 2. $M$ -estimates based on ranks

In this section, we introduce estimates that, in an approximate sense, are  $M$ -estimates based on ranks. We start by assuming that the distribution  $F$  of the  $X$ 's is known and introduce  $M$ -estimates of  $\theta$  that depend on  $F$ . Let

$$U_i = F(X_i), \quad V_j = F(Y_j), \quad i = 1, \dots, m, \quad j = 1, \dots, n.$$

The joint distribution of  $U_1, \dots, U_m, V_1, \dots, V_n$  is  $\prod_{i=1}^m u_i \prod_{j=1}^n D(v_j, \theta)$ , so that  $(V_1, \dots, V_n)$  is sufficient for  $\theta$ . Let  $\psi(v, \theta)$  be a function which is monotone decreasing in  $\theta$ , and satisfies  $E_{\theta_0} \psi(V, \theta_0) = 0$  where  $V$  is distributed according to  $D(v, \theta_0)$  and  $\theta_0$  is the true parameter value. An  $M$ -estimate (see Huber (1981)) of  $\theta$  is defined as a solution to the equation  $\Sigma \psi(V_j, \theta) = 0$ .

Let  $N = m + n$ . When  $F$  is unknown, we define  $\hat{F}(u) = mF_m(u)/(m+1)$  for  $u \in [X_{(1)}, X_{(m)}]$ ,  $\hat{F}(u) = 1/(N+1)$  for  $u < X_{(1)}$  and  $\hat{F}(u) = N/(N+1)$  for  $u > X_{(m)}$ . Here  $X_{(1)}$  and  $X_{(m)}$  denote the first and the last order statistics of the  $X_i$ 's and  $F_m(u) = m^{-1} \# \{i: X_i \leq u\}$ .

We set  $\hat{V}_j = \hat{F}(Y_j)$ , and we let  $\hat{\theta}$  be any "solution" to

$$\sum_{j=1}^n \psi(\hat{V}_j, \theta) = 0.$$

More precisely,  $\hat{\theta}$  is any value in the interval  $[\theta^*, \theta^{**}]$  where  $\theta^* = \sup \{\theta: \Sigma \psi(\hat{V}_j, \theta) > 0\}$  and  $\theta^{**} = \inf \{\theta: \Sigma \psi(\hat{V}_j, \theta) < 0\}$ .

In terms of the rank likelihood,  $\hat{V}_1, \dots, \hat{V}_n$  are sufficient for  $\theta$ . To see this, let  $Y_{(1)} < \dots < Y_{(n)}$  be the ordered statistics and let  $K_{(j)} = m\hat{F}(Y_{(j)})$ . Then it is easy to check that  $K_{(1)}, \dots, K_{(n)}$  are equivalent to the ranks. Thus we call  $\hat{\theta}$  a *RAM (Rank Approximate M)* estimate.

Here is an example where we get an explicit formula for  $\hat{\theta}$ .

*Example 2.1.* Assuming  $F$  is known, the  $M$ -estimate based on  $\psi(v, \theta) = -\theta - \log(1-v)$  is the MLE (Maximum Likelihood Estimate) for the proportional hazard model of Example 1.1. This estimate is  $n^{-1} \Sigma [-\log(1-F(Y_j))]$ . The corresponding RAM estimate of  $\theta$  which applies when  $F$  is unknown is  $\hat{\theta} = n^{-1} \sum_{j=1}^n [-\log(1-\hat{F}(Y_j))]$ .

We return to the general case and show asymptotic normality of the RAM estimates. We assume throughout that the limits  $\pi_0 = \lim_{N \rightarrow \infty} (m/N)$  and  $\pi_1 = \lim_{N \rightarrow \infty} (n/N)$  exist and are strictly between 0 and 1. Further, we assume that  $\psi$  is continuously differentiable in  $u$  and we set  $\psi'(u, \theta) = (\partial/\partial u)[\psi(u, \theta)]$  and  $d(u, \theta) = (\partial/\partial u)[D(u, \theta)]$ . Define

$$\begin{aligned} \lambda(\theta) &= \int_0^1 \psi(u, \theta) d(u, \theta_0) du, & \lambda_1(\theta) &= \int_0^1 \psi_1(u, \theta) du, \\ \psi_1(u, \theta) &= \int_0^u \psi'(v, \theta) d(v, \theta_0) dv, \\ \sigma^2(\theta) &= \pi_0^{-1} \left\{ \int_0^1 \psi_1^2(u, \theta) du - \lambda_1^2(\theta) \right\} + \pi_1^{-1} \left\{ \int_0^1 \psi^2(u, \theta) d(u, \theta_0) du - \lambda^2(\theta) \right\}. \end{aligned}$$

Assume that for  $\theta$  in a neighbourhood of the true parameter value  $\theta_0$ , the following assumptions hold:

(A.1)  $|\psi(\cdot, \theta)| = O(r^a)$  and  $|\psi'(\cdot, \theta)| = O(r^{a+1})$  where  $r(u) = [u(1-u)]^{-1}$  and  $a = 1/2 - \tau$  for some  $0 < \tau < 1/2$ .

(A.2)  $\int r(u)^{1-\eta} d(u, \theta) du < \infty$  uniformly in  $\theta$  for some  $0 < \eta < \tau$ .

(A.3)  $m \int u^m d(u, \theta) du = O(1)$  and  $m \int (1-u)^m d(u, \theta) du = O(1)$  uniformly in  $\theta$  and  $m$ .

Assumption (A.3) is satisfied whenever the density  $d(u, \theta)$  is bounded uniformly in  $\theta$  in a neighbourhood of  $\theta_0$ . In particular, it holds for the proportional hazard model with  $\theta_0 \leq 1$  and  $\gamma$ -rate models of Examples 1.2 and 1.3. Assumption (A.2) is satisfied in all three examples.

**THEOREM 2.1.** *Suppose that  $\psi'(u, \theta)$  and  $\lambda(\theta)$  are continuous in  $\theta$  for  $\theta$  in a neighbourhood of the true parameter value  $\theta_0$  and suppose that  $\theta_0$  is the unique point with  $\lambda(\theta_0) = 0$ . Assume that  $\sigma^2(\theta)$  is finite, nonzero and continuous in a neighbourhood of  $\theta_0$  and that (A.1), (A.2) and (A.3) hold. Then  $\sqrt{N}\lambda(\hat{\theta})$  is asymptotically normally distributed with mean zero and variance  $\sigma^2(\theta_0)$ . In addition, if  $\lambda'(\theta_0)$  exists and  $\lambda'(\theta_0) < 0$ , then  $\sqrt{N}(\hat{\theta} - \theta_0)$  is asymptotically normal with mean zero and variance  $\sigma^2(\theta_0)/[\lambda'(\theta_0)]^2$ .*

**PROOF.** First we note that asymptotically all  $\tilde{\theta} \in [\theta^*, \theta^{**}]$  are equivalent, so it is enough to consider  $\theta^*$ .

By the assumed monotonicity of  $\psi(u, \theta)$  in  $\theta$ , the function  $\lambda(\theta)$  is monotone decreasing. Let  $y$  be fixed. Since  $\lambda(\theta_0) = 0$  and  $\lambda$  is continuous, for  $N$  sufficiently large, there is  $\theta_N$  such that  $y = -\sqrt{N}\lambda(\theta_N)$ . In fact,  $\theta_N = -\lambda^{-1}(-y/\sqrt{N})$ . Let  $G_n$  denote the empirical distribution function of the  $Y$ 's. Then

$$\begin{aligned} P(-\sqrt{N}\lambda(\theta^*) < y) &= P(\theta^* < \theta_N) \\ &= P\left[\frac{\sqrt{N}\left[\int\psi(\hat{F}, \theta_N)dG_n - \lambda(\theta_N)\right]}{\sigma(\theta_N)} \leq \frac{y}{\sigma(\theta_N)}\right] \\ &\rightarrow \Phi\left[\frac{y}{\sigma(\theta_0)}\right]. \end{aligned}$$

To see this we note that

$$\begin{aligned} (2.1) \quad \sqrt{N}\left[\int\psi(\hat{F}, \theta_N)dG_n - \lambda(\theta_N)\right] \\ = \sqrt{N}\left[\int(F_m - F)\psi'(F, \theta_N)dG + \int\psi(F, \theta_N)d(G_n - G)\right] + r_N, \end{aligned}$$

where  $r_N$  is a remainder term. We have

$$\begin{aligned} &\int(F_m - F)\psi'(F, \theta_N)dG \\ &= m^{-1}\sum_{i=1}^m\int[I(X_i < x) - F(x)]\psi'(F(x), \theta_N)d(F(x), \theta_0)dF(x) \\ &= \sum_{i=1}^m A_{im}, \end{aligned}$$

which is a sum of independent identically distributed (iid) random variables.

By Assumption (A.1),  $|A_{im}| = O(1)r^{1/2-\tau+\eta}(F(X_i))\int r^{1-\eta}(u)d(u, \theta_0)du$ . By

Assumption (A.2), in a neighbourhood of  $\theta_0$  the deterministic part of this upper bound is uniformly bounded from above. Further, for some  $\tau_1 > 0$  the random part of this bound has an absolute moment of order  $2 + \tau_1$ , which is uniformly bounded above.

Further,  $\int \psi(F, \theta_N) d(G_n - G)$  is a sum of iid random variables. By Assumptions (A.1) and (A.2), they have a finite absolute moment of order  $2 + \tau_2$  for some  $\tau_2 > 0$ . Berry-Esseen's theorem completes the proof of the asymptotic normality of the first two terms. The remainder term  $r_N$  is considered in Section 4.

Note that when  $\theta_0 = 1$ , the asymptotic variance of  $\hat{\theta}$  reduces to

$$(\pi_0^{-1} + \pi_1^{-1}) E \psi^2(U, 1) \left/ \left[ E \frac{\partial}{\partial \theta} \psi(U, \theta) \Big|_{\theta=1} \right]^2 \right.,$$

where  $U$  is uniform on  $(0, 1)$ . This is exactly the same as the asymptotic variance of the  $M$ -estimate for the model (1.1) with  $H$  known. Thus if we choose

$$\psi(v, \theta) = \frac{\partial}{\partial \theta} \log d(v, \theta),$$

then  $\hat{\theta}$ , in addition to being consistent and asymptotically normal for general  $\theta_0$ , is asymptotically efficient when  $\theta_0 = 1$ .

If we consider the RAM estimate  $\hat{\theta} = n^{-1} \sum_{j=1}^n [-\log(1 - \hat{F}(Y_j))]$ , we find that in the proportional hazard model the asymptotic variance is  $\pi_0^{-1} \theta(2 - \theta)^{-1} + \pi_1^{-1} \theta^2$  for  $\theta \leq 1$ . We can compare this estimate with the Cox partial likelihood estimate. The usual parametrization in the Cox proportional hazard model is in terms of  $\beta = \ln \theta$ . If  $\beta^*$  is the Cox partial likelihood estimate, then from Efron (1977) we find that its asymptotic variance is given by

$$\left[ \int_0^1 \{ \pi_1^{-1} + \pi_0^{-1} \theta u^{(\theta-1)/\theta} \}^{-1} du \right]^{-1}.$$

The asymptotic variance of  $\hat{\beta}_0 = \log \hat{\theta}$  where  $\hat{\theta}$  is the RAM estimate is given by

$$\pi_0^{-1} \theta^{-1} (2 - \theta)^{-1} + \pi_1^{-1},$$

for  $\theta \leq 1$ . For  $\pi_0 = \pi_1 = 1/2$  the asymptotic relative efficiency of  $\hat{\beta}_0$  with respect to  $\beta^*$  is equal to 1, 0.951, 0.863, 0.757 and 0.647 for  $\theta = 1, 1/2, 1/4,$

1/8 and 1/16.

*Example 2.2. (Proportional  $\gamma$ -odds model)* Our main application is to the proportional  $\gamma$ -rate model,  $\gamma > 0$ . The  $\psi$  function corresponding to the MLE for the  $F$  known case is

$$(2.2) \quad \psi(u, \theta) = \gamma^{-1} - (1 + \gamma^{-1}) \frac{(1-u)^\gamma \theta}{1 - (1-u)^\gamma + \theta(1-u)^\gamma}.$$

Using Theorem 2.1, we find that the asymptotic variance of the RAM estimate simplifies when  $\gamma$  is of the form  $\gamma = 1/k$ , where  $k \geq 1$  is an integer. In this case the asymptotic variance is

$$\frac{\theta^2}{\pi_1} \frac{2+k}{k} + \frac{\theta^2}{(1-\theta)^2} \frac{1}{\pi_0} \left\{ \frac{(1+k)^2}{k} \theta^{k+2} \sum_{i=0}^{3+k} \binom{3+k}{i} \cdot \frac{1}{i+k} \left( \frac{1-\theta}{\theta} \right)^i - \left( \frac{k+\theta}{k} \right)^2 \right\}.$$

For  $\gamma = k = 1$ , the proportional odds model, the asymptotic variance is  $3\theta^2\pi_1^{-1} + (0.2)\theta\pi_0^{-1}(4 + 7\theta + 4\theta^2)$  while for  $k = 2$ , it equals  $2\pi_1^{-1}\theta^2 + \theta\pi_0^{-1}\{18 + 23\theta + 12\theta^2 + 3\theta^3\}/28$ .

To judge the performance of the RAM estimate based on (2.1), we compute the efficiency of this estimate with respect to the MLE for the  $\gamma$ -odds model with  $F$  known and the  $X$ 's and  $Y$ 's distributed as  $D(H(x), \theta_1)$  and  $D(H(y), \theta_2)$ , respectively, where  $D$  is given in Example 1.3, and  $\theta = \theta_2/\theta_1$ . For  $m = n$ , this efficiency is given by  $e(\hat{\theta}_{\text{RAM}}, \hat{\theta}_{\text{MLE}}) = 60\theta(8 + 44\theta + 8\theta^2)^{-1}$  for  $k = 1$  and  $112\theta(18 + 79\theta + 12\theta^2 + 3\theta^3)^{-1}$  for  $k = 2$ .

Here is a brief table of these efficiencies.

We see from Table 1 that when  $\gamma = 1$  and  $1/2$ ,  $\hat{\theta}$  is quite efficient for  $\theta$  in the range (0.5, 2). The efficiency increases as  $\gamma$  increases. In fact, it is easy to show that as  $\gamma \rightarrow \infty$  the efficiency tends to one for all  $\theta$ . The efficiency given is a lower bound on the efficiency of  $\hat{\theta}$  with respect to the asymptotically optimal estimate based on the ranks.

Table 1. The asymptotic relative efficiency of the RAM estimate with respect to the MLE for the parametric  $\gamma$ -odds model.

| $\gamma \backslash \theta$ | .25  | .5   | .75  | .9   | 1 | 1.1  | 1.25 | 1.5  | 1.75 | 2    | 4    |
|----------------------------|------|------|------|------|---|------|------|------|------|------|------|
| 1                          | .769 | .938 | .989 | .999 | 1 | .999 | .993 | .978 | .959 | .938 | .769 |
| 0.5                        | .726 | .920 | .985 | .998 | 1 | .998 | .990 | .968 | .937 | .903 | .624 |



### 3. Rank-inversion estimates

In this section, we introduce rank-inversion estimates based on the ideas of Hodges-Lehmann (1963). Again, we start by assuming that  $F$  is known and let  $U_i$ ,  $V_j$  and  $D(u, \theta)$  be as in Section 2. In particular, we assume that  $D(u, \theta)$  is monotone decreasing in  $\theta$ . Note that  $U_1, \dots, U_m$ ,  $D(V_1, \theta), \dots, D(V_n, \theta)$  all have the same distribution when  $\theta = \theta_0$ , where  $\theta_0$  is the true value of the parameter. Let  $R_i(\theta)$  denote the rank of  $U_i$  among  $U_1, \dots, U_m, D(V_1, \theta), \dots, D(V_n, \theta)$ , and let

$$T_N(\theta) = m^{-1} \sum_{i=1}^m J_N \left( \frac{R_i(\theta)}{N+1} \right),$$

denote a linear rank function with monotone increasing score function  $J_N$ . For  $F$  known, the Hodges-Lehmann estimate of  $\theta$  is obtained by solving  $T_N(\theta) = \int_0^1 J(u) du$  for  $\theta$ , where  $J(u)$  is the limit of  $J_N(u)$ . Without loss of generality, we assume  $\int_0^1 J(u) du = 0$ .

Suppose now that  $F$  is unknown. Let  $X_{(1)} < \dots < X_{(m)}$  be the vector of order statistics of  $X_i$ 's. Let  $\tilde{F}$  be defined by

$$\tilde{F}(u) = \frac{u + iX_{(i+1)} - (i+1)X_{(i)}}{(m+1)(X_{(i+1)} - X_{(i)})},$$

for  $X_{(i)} \leq u \leq X_{(i+1)}$ ,  $i = 1, \dots, m-1$ . Thus on the interval  $[X_{(1)}, X_{(m)}]$ ,  $\tilde{F}$  is a linearized version of the right-continuous distribution function  $mF_m/(m+1)$ , where  $F_m(u) = m^{-1} \# \{i: X_i \leq u\}$ . Further, let  $Y_{(1)}$  and  $Y_{(n)}$  be the first and the last order statistics of the  $Y_j$ 's. If  $Y_{(1)} < X_{(1)}$  or  $Y_{(n)} > X_{(m)}$ , then we extend  $\tilde{F}$  to the interval  $[\min(X_{(1)}, Y_{(1)}), \max(X_{(m)}, Y_{(n)})]$  linearly with  $\tilde{F}(Y_{(1)}) = 1/(N+1)$  if  $Y_{(1)} < X_{(1)}$  and  $\tilde{F}(Y_{(n)}) = N/(N+1)$  if  $Y_{(n)} > X_{(m)}$ .

Further, let  $\tilde{R}_i(\theta)$  be the rank of  $\tilde{F}(X_i)$  among  $\tilde{F}(X_1), \dots, \tilde{F}(X_m)$ ,  $D(\tilde{F}(Y_1), \theta), \dots, D(\tilde{F}(Y_n), \theta)$ . Let  $\tilde{\theta}_R$  be any "solution" to

$$\tilde{T}_N(\theta) = m^{-1} \sum_{i=1}^m J_N \left( \frac{\tilde{R}_i(\theta)}{N+1} \right) = 0.$$

More precisely, let  $\tilde{\theta}_R$  be any point in  $[\theta_R^*, \theta_R^{**}]$  where  $\theta_R^* = \sup \{\theta: \tilde{T}_N(\theta) < 0\}$  and  $\theta_R^{**} = \inf \{\theta: \tilde{T}_N(\theta) > 0\}$ . Similar estimates have also been considered by Doksum and Nabeya (1984) and Miura (1985).

*Example 3.1.* Assuming  $F$  is known, the Hodges-Lehmann type rank estimate based on  $J(u) = 2u - 1$  is asymptotically optimal for the proportional odds model. Let  $L(x) = 1/[1 + e^{-x}]$  be the logistic distribution func-

tion and note that if we set  $W_i = L^{-1}(F(X_i))$  and  $Z_j = L^{-1}(F(Y_j))$ , then  $W_i$  and  $Z_j$  follow a logistic shift model ( $W_i \sim L(w)$ ,  $Z_j \sim L(z - \log \theta)$ ) with parameter  $\log \theta$ . Since the ranks are invariant under the increasing transformation  $L^{-1}$ , it follows that the Hodges-Lehmann estimate of  $\theta$  is

$$\tilde{\theta}_{HL} = \exp \left\{ \text{median}_{i,j} (Z_j - W_i) \right\} .$$

The corresponding  $\tilde{\theta}_R$ , which is appropriate when  $F$  is unknown, is

$$\tilde{\theta}_R = \exp \left\{ \text{median}_{i,j} (L^{-1}(\tilde{F}(Y_j)) - L^{-1}(\tilde{F}(X_i))) \right\} .$$

We return to the general case and show the asymptotic normality of  $\sqrt{N}(\tilde{\theta}_R - \theta_0)$ . From (1.2), we have

$$(3.1) \quad D(D^{-1}(u, \theta), \delta) = D(u, \delta/\theta), \quad D(u, 1) = u .$$

Let

$$\begin{aligned} \dot{D}(u, \theta) &= \frac{\partial}{\partial \theta} D(u, \theta) \quad \eta = \int_0^1 J'(u) \dot{D}(u, 1) du , \\ \tau^2(\theta) &= \pi_1^{-1} \int_0^1 J^2(u) du + \pi_0^{-1} \left[ \int_0^1 \alpha^2(u) du - \left( \int_0^1 \alpha(u) du \right)^2 \right] , \end{aligned}$$

where  $\alpha$  is defined by

$$\frac{d\alpha(u)}{du} = J'(D(u, \theta)) [d(u, \theta)]^2 .$$

Further, let

$$\begin{aligned} G_n^\theta(u) &= n^{-1} \# [j: D(\tilde{F}(Y_j), \theta) \leq u] = G_n \tilde{F}^{-1} D(u, \theta^{-1}) , \\ H_N^\theta(u) &= \{m F_m \tilde{F}^{-1}(u) + n G_n^\theta(u)\} / (N + 1) , \\ G^\theta(u) &= D(D^{-1}(u, \theta), \theta_0) = D(u, \theta_0/\theta) , \end{aligned}$$

(by (3.1)) and  $H^\theta(u) = \pi_0 u + \pi_1 G^\theta(u)$ . In terms of these functions, we have

$$\tilde{T}_N(\theta) = m^{-1} \sum_{i=1}^m J_N \left( \frac{\tilde{R}_i(\theta)}{N+1} \right) = \int J_N(H_N^\theta(\tilde{F})) dF_m = \int J_N(H_N^\theta) dF_m \tilde{F}^{-1} .$$

Assume

$$(B.1) \quad r_{1N} = \sqrt{N} \int \{J_N(H_N^\theta) - J(H_N^\theta)\} dF_m \tilde{F}^{-1} \xrightarrow{P} 0 \text{ as } N \rightarrow \infty \text{ uniformly}$$

for  $\theta$  in a neighbourhood of  $\theta_0$ .

Moreover,

(B.2)  $J$  is a differentiable function with bounded continuous derivative  $J'$ , and  $0 < \int J^2(u) du < \infty$ .

Finally, we assume that the limits  $\pi_0 = \lim_{N \rightarrow \infty} (m/N)$  and  $\pi_1 = \lim_{N \rightarrow \infty} (n/N)$  exist and are strictly between 0 and 1.

**THEOREM 3.1.** *If  $D(u, \theta)$  is decreasing in  $\theta$ , and if the preceding conditions hold, then  $\sqrt{N}(\tilde{\theta}_R - \theta_0)$  has asymptotically a normal distribution with mean zero and variance  $\theta_0^2 \tau^2(\theta_0)/\eta^2$ .*

**PROOF.** As in the case of Hodges-Lehmann (1963),  $\theta_R^*$ ,  $\theta_R^{**}$  and any point between them, such as  $\tilde{\theta}_R$ , will have the same asymptotic distribution. Further,

$$P(\sqrt{N}(\theta_R^*/\theta_0 - 1) \leq t) = P(\sqrt{N} \tilde{T}_N(\delta) > 0),$$

where  $\delta = \theta_0(1 + t/\sqrt{N})$ . We have

$$\begin{aligned} \sqrt{N} \tilde{T}_N(\delta) &= \sqrt{N} \int J_N(H_N^\delta(u)) dF_m \tilde{F}^{-1}(u) \\ &= \sqrt{N} \int J(H_N^\delta(u)) d[F_m \tilde{F}^{-1}(u) - u] + \sqrt{N} \int [J(H_N^\delta(u)) - J(H^\delta(u))] du \\ &\quad + \sqrt{N} \int [J(H^\delta(u)) - J(u)] du + r_{1N} = I_1 + I_2 + I_3 + r_{1N}. \end{aligned}$$

Note that  $I_1 = -\sqrt{N} \int [F_m \tilde{F}^{-1}(u) - u] dJ(H_N^\delta(u))$ . This term is bounded in absolute value by  $\sup \sqrt{N} |F_m \tilde{F}^{-1}(u) - u| \int |J'(u)| du$ , which tends to zero since  $\sup |F_m \tilde{F}^{-1}(u) - u| \leq 2/(m+1)$  and  $\int |J'(u)| du < \infty$  by (B.2).

The second term can be written as

$$I_2 = \sqrt{N} \int J'(H^\delta(u)) [H_N^\delta(u) - H^\delta(u)] du + r_{2N},$$

where

$$r_{2N} = \sqrt{N} \int (H_N^\delta - H^\delta) \{J'(\tilde{H}^\delta) - J'(H^\delta)\} du,$$

and  $\tilde{H}^\delta$  assumes values between  $H_N^\delta$  and  $H^\delta$ . We have

$$\begin{aligned} (3.2) \quad &\sqrt{N} \int J'(H^\delta(u)) [H_N^\delta(u) - H^\delta(u)] du \\ &= \sqrt{N} \int J'(H^\delta(u)) [m/(N+1) F_m \tilde{F}^{-1}(u) - \pi_0 u] du \end{aligned}$$

$$\begin{aligned}
& + \sqrt{N} \int J'(H^\delta(u))(n/(N+1)) \\
& \quad \cdot [G_n \tilde{F}^{-1} D^{-1}(u, \delta) - G \tilde{F}^{-1} D^{-1}(u, \delta)] du \\
& + \sqrt{N} \int J'(H^\delta(u)) [n/(N+1) G \tilde{F}^{-1} D^{-1}(u, \delta) \\
& \quad - \pi_1 G F^{-1} D^{-1}(u, \delta)] du .
\end{aligned}$$

The first term converges in probability to 0 by an argument similar to  $I_1$ . The next two terms of this expansion converge weakly to

$$\pi_1 \left\{ \pi_1^{-1/2} \int J'(u) B_2(u) du - \pi_0^{-1/2} \int J'(D(u, \theta_0)) d^2(u, \theta_0) B_1(u) du \right\},$$

where  $B_1$  and  $B_2$  are independent Brownian bridges. This follows since  $\sqrt{n}(G_n - G)$  and  $\sqrt{m}(\tilde{F}^{-1}; -F^{-1})$  converge weakly to  $B_2(G)$  and  $-f(F^{-1})B_1(u)$ , respectively and  $\sup |\tilde{F}^{-1} - F^{-1}| = o_P(1)$ . The standard Skorokhod construction yields the desired result. Further,  $r_{2N} \xrightarrow{P} 0$  by (B.2),  $\sup |H_N^\delta - H^\delta| \xrightarrow{P} 0$  and  $\sup \sqrt{N}(H_N^\delta - H^\delta) = O_P(1)$ .

Finally,

$$I_3 = \sqrt{N} \int J'(u) \pi_1 \{D(u, \theta_0/\delta) - u\} du + r_{4N} \rightarrow -\pi_1 t \int_0^1 J'(u) \dot{D}(u, 1) du,$$

and

$$r_{4N} = \sqrt{N} \int [J'(\tilde{H}^\delta) - J'(u)] \pi_1 [D(u, \theta_0/\delta) - u] du,$$

where  $\tilde{H}^\delta$  assumes values between  $H^\delta$  and  $H \equiv u$ . This term converges to 0 by assumption (B.2) and Taylor expansion of  $D(u, \theta_0/\delta)$ .

If we let  $J(u) = [1 + \gamma^{-1}](1 - u)^\gamma - \gamma^{-1}$ ,  $\gamma \geq 1$ , then in the proportional  $\gamma$ -odds model,  $\hat{\theta}_R$  given in Example 3.1 will have the same asymptotic variance as the RAM estimate based on (2.2) (see Example 2.2).

#### 4. Proof of Theorem 2.1: remainder terms

The remainder term  $r_N$  in (2.1) is given by

$$r_N = \sqrt{N} \int \{\psi(\hat{F}, \theta_N) - \psi(F, \theta_N)\} dG_n - \sqrt{N} \int (F_m - F) \psi'(F, \theta_N) dG.$$

For small  $\gamma \in (0, 1)$  define  $S_\gamma = [F^{-1}(\gamma), F^{-1}(1 - \gamma)]$ . Further, let  $\Delta_N = [Y_{(1)}, Y_{(n)}]$  and  $E_N = [X_{(1)}, X_{(m)}]$  where  $X_{(1)}$ ,  $X_{(m)}$ ,  $Y_{(1)}$  and  $Y_{(n)}$  are the first and the last order statistics among  $X_i$ 's and  $Y_j$ 's. Then, after some algebra,

$$r_N = \sum_{i=1}^7 r_{iN} \text{ where}$$

$$\begin{aligned}
r_{1N} &= \sqrt{N} \int_{\Delta_N \cap S_\gamma} \{\psi'(F^*, \theta_N) - \psi'(F, \theta_N)\}(\hat{F} - F) dG_n, \\
r_{2N} &= \sqrt{N} \int_{\Delta_N \cap S_\gamma} \psi'(F, \theta_N)(\hat{F} - F_m) dG_n, \\
r_{3N} &= \sqrt{N} \int_{\Delta_N \cap S_\gamma} \psi'(F, \theta_N)(F_m - F) d(G_n - G), \\
r_{4N} &= -\sqrt{N} \int_{\Delta_N^c \cup S_\gamma^c} \psi'(F, \theta_N)(F_m - F) dG, \\
r_{5N} &= \sqrt{N} \int_{\Delta_N \cap S_\gamma^c \cap E_N} \psi'(F^*, \theta_N)(\hat{F} - F) dG_n, \\
r_{6N} &= \sqrt{N} \int_{\Delta_N \cap S_\gamma^c \cap E_N^c} \psi(\hat{F}, \theta_N) dG_n, \\
r_{7N} &= -\sqrt{N} \int_{\Delta_N \cap S_\gamma^c \cap E_N^c} \psi(F, \theta_N) dG_n.
\end{aligned}$$

Here  $F^*$  is a random function assuming values between  $F$  and  $\hat{F}$ . We shall show that for any fixed  $\gamma$ ,  $r_{1N}$ ,  $r_{2N}$ ,  $r_{3N}$  and  $r_{6N}$  converge in probability to 0 and  $r_{4N}$ ,  $r_{5N}$  and  $r_{7N}$  converge in probability to 0 as  $\gamma \rightarrow 0$  and  $N \rightarrow \infty$ .

LEMMA 4.1. *For fixed  $\gamma$ ,  $r_{1N} \xrightarrow{P} 0$  as  $N \rightarrow \infty$ .*

PROOF. Given  $\gamma \in (0, 1)$  let  $\Omega_{\gamma N} = \{\omega: \sup_{S_\gamma} |\hat{F} - F| < \gamma/2\}$ . Then  $r_{1N} = I(\Omega_{\gamma N})r_{1N} + I(\Omega_{\gamma N}^c)r_{1N}$ . The second term converges in probability to 0, by the Glivenko-Cantelli theorem. Further, we have  $\sup_{S_\gamma} \sqrt{m}|\hat{F} - F| = O_p(1)$ . The function  $\psi'(u, \theta_N)$  is uniformly continuous on  $[\gamma/2, 1 - \gamma/2]$  and  $|F^* - F| \leq |\hat{F} - F|$ . Therefore, by the Glivenko-Cantelli theorem,

$$I(\Omega_{\gamma N})|r_{1N}| \leq O_p(1) \sup_{S_\gamma} |\psi'(F^*, \theta_N) - \psi'(F, \theta_N)| \xrightarrow{P} 0.$$

LEMMA 4.2. *For fixed  $\gamma$ ,  $r_{2N} \xrightarrow{P} 0$  as  $N \rightarrow \infty$ .*

PROOF. This follows since  $|\hat{F} - F_m| \leq 1/(m+1)$  and  $\int_{S_\gamma} \psi'(F) dG_n \xrightarrow{P} \int_{S_\gamma} \psi'(F) dG$ .

LEMMA 4.3. *For fixed  $\gamma$ ,  $r_{3N} \xrightarrow{P} 0$  as  $N \rightarrow \infty$ .*

PROOF. For each positive integer  $k$  define a function  $\chi_k$  on  $[0, 1]$  by

$$\chi_k(0) = 0 ,$$

$$\chi_k(s) = (i-1)/k \quad \text{for} \quad (i-1)/k < s \leq i/k \quad i = 1, \dots, k .$$

Then, by Lemma 4.3 of Ruymgaart *et al.* (1972)

$$\sup \sqrt{m} | F_m F^{-1} \chi_k(F) - \chi_k(F) - F_m + F | \xrightarrow{P} 0 ,$$

as  $k, m \rightarrow \infty$ . Furthermore,  $|r_{3N}| \leq \sum_{i=1}^3 r_{3iN}$  where

$$\begin{aligned} r_{31N} &= \sqrt{N} \int_{\Delta_N \cap S_\gamma} |(F_m - F) \psi'(F, \theta_N) \\ &\quad - (F_m F^{-1} \chi_k(F) - \chi_k(F)) \psi'(\chi_k(F), \theta_N)| dG_n , \\ r_{32N} &= \sqrt{N} \int_{\Delta_N \cap S_\gamma} |(F_m - F) \psi'(F, \theta_N) \\ &\quad - (F_m F^{-1} \chi_k(F) - \chi_k(F)) \psi'(\chi_k(F), \theta_N)| dG , \\ r_{33N} &= \left| \sqrt{N} \int_{\Delta_N \cap S_\gamma} (F_m F^{-1} \chi_k(F) - \chi_k(F)) \psi'(\chi_k(F), \theta_N) d(G_n - G) \right| . \end{aligned}$$

The proof is similar to Corollary 5.5 in Ruymgaart *et al.* (1972). Given  $\varepsilon > 0$  there exist constants  $M$  and  $\eta_{kN} \rightarrow 0$  as  $k, N \rightarrow \infty$  such that the sets

$$\begin{aligned} \Omega_N &= \{ \sup \sqrt{m} | F_m - F | \leq M \} , \\ \Omega_{kN} &= \{ \sup \{ \sqrt{m} | F_m - F - F_m F^{-1} \chi_k(F) + \chi_k(F) | \leq \eta_{kN} \} , \end{aligned}$$

have probability at least  $1 - \varepsilon$ . Further, the function  $\psi'(u, \theta_N)$  is bounded by  $M_\gamma$  and uniformly continuous on  $[0, 1 - \gamma]$ . Finally

$$\xi_{k\gamma N} = \sup_{S_\gamma} | \psi'(F, \theta_N) - \psi'(\chi_k(F), \theta_N) | \rightarrow 0 .$$

Therefore, for  $i = 1, 2$

$$I(\Omega_{kN} \cap \Omega_N) r_{3iN} \leq (\eta_{kN} M_\gamma + M \xi_{k\gamma N}) \sqrt{N/m} \rightarrow 0 .$$

Finally, for each  $\omega \in \Omega_N$ , the integrand of  $r_{33N}$  is a step function assuming value  $a_{ikN}$ ,  $|a_{ikN}| \leq M(M_\gamma + \xi_{k\gamma N}) \sqrt{N/m}$  on the interval  $[F^{-1}((i-1)/k), F^{-1}(i/k)] = R_{ikN}$  and

$$I(\Omega_N) r_{33N} = \left| \sum_{i=1}^k a_{ikN} \int_{R_{ikN}} d(G_n - G) \right|$$

$$\leq 2kM(M_\gamma + \xi_{k\gamma N})\sqrt{N/m} \sup |G_n - G| \xrightarrow{P} 0,$$

as  $N \rightarrow \infty$ .

LEMMA 4.4. *As  $\gamma \downarrow 0$  and  $N \rightarrow \infty$ ,  $r_{4N} \xrightarrow{P} 0$ .*

PROOF. Let  $\varepsilon > 0$  be fixed. By Theorem A1 of Shorack (1972), there exists a constant  $M$  such that the set

$$\Omega_N = \{\sqrt{m}(F_m - F) \leq Mr^{-1/2+\tau-\eta}(F)\},$$

has probability at least  $1 - \varepsilon$ . By Assumption (A.1)

$$(4.1) \quad EI(\Omega_N)|r_{4N}| \leq O(1)M\sqrt{N/m} \int_{\Delta_N^c \cup S_\gamma^c} r(F)^{1-\eta} dG.$$

Let us consider the integral

$$(4.2) \quad \int_{S_\gamma^c} r(u)^{1-\eta} d(u, \theta) du.$$

By Assumption (A.2), we can find a value  $\tilde{\gamma}$  of  $\gamma$  such that (4.2) is less than  $\varepsilon$  provided  $\gamma \leq \tilde{\gamma}$ . For this  $\tilde{\gamma}$  there exists  $\tilde{N}$  such that  $P(\Delta_N \supset S_{\tilde{\gamma}}) > 1 - \varepsilon$  provided  $N > \tilde{N}$ . It follows that the integral on the right-hand side of (4.1) is less than  $\varepsilon$  with probability larger than  $1 - \varepsilon$  for  $\gamma \leq \tilde{\gamma}$  and  $N \geq \tilde{N}$ . Thus,  $r_{4N} \xrightarrow{P} 0$  as  $\gamma \downarrow 0$  and  $N \rightarrow \infty$ .

LEMMA 4.5. *As  $\gamma \downarrow 0$  and  $N \rightarrow \infty$ ,  $r_{5N} \xrightarrow{P} 0$ .*

PROOF. Let  $\varepsilon > 0$  be fixed. By Lemmas 6.1 and 6.2 in Ruymgaart *et al.* (1972), there exist constants  $M_1$  and  $M_2$  such that the sets  $\Omega_{1N} = \{\sqrt{m}|\hat{F} - F| \leq M_1 r^{-1/2+\tau-\eta}(F)$  on  $E_N\}$  and  $\Omega_{2N} = \{r^{a+1}(F^*) \leq M_2 r^{a+1}(F)$  on  $E_N\}$  have probability at least  $1 - \varepsilon$ . Then

$$EI(\Omega_{1N} \cap \Omega_{2N})|r_{5N}| \leq O(1)M_1 M_2 \sqrt{N/m} \int_{S_\gamma^c} r^{1-\eta}(F) dG.$$

The right-hand side converges to 0 as  $\gamma \downarrow 0$ .

LEMMA 4.6. *For fixed  $\gamma$ ,  $r_{6N} \xrightarrow{P} 0$  as  $N \rightarrow \infty$ .*

PROOF. Let  $\varepsilon > 0$  be fixed. Given  $\gamma \in (0, 1)$ , we can find  $\tilde{N}$  such that the set  $\Omega_N = \{S_\gamma \subset E_N\}$  has probability at least  $1 - \varepsilon$  for  $N > \tilde{N}$ . By Assumption (A.1),  $EI(\Omega_N)|r_{6N}| \leq O((N/m)^{1-\tau})m^{1-\tau}P(Y \in E_N^c)$ . But

$$P(Y \in E_N^c) = \int u^m d(u, \theta_N) du + \int (1-u)^m d(u, \theta_N) du .$$

By Assumption (A.3),  $EI(\Omega_N) |r_{6N}| \rightarrow 0$ .

LEMMA 4.7. As  $\gamma \downarrow 0$  and  $N \rightarrow \infty$ ,  $r_{7N} \xrightarrow{P} 0$ .

PROOF. By Assumption (A.1),

$$|r_{7N}| \leq \sqrt{N} \int_{A_N \cap S_\gamma^c \cap E_N^c} r(F)^a dG_n .$$

Hölders inequality yields

$$E|r_{7N}| \leq \sqrt{N/m} \{mP(Y_i \in E_N^c)\}^{1/2} \left\{ \int_{S_\gamma^c} r(F)^{2a} dG \right\}^{1/2} .$$

By Assumptions (A.2) and (A.3),  $mP(Y_i \in E_N^c) = O(1)$  and the integral on the right-hand side converges to 0 as  $\gamma \downarrow 0$ .

## REFERENCES

- Anscombe, F. J. and Tukey, J. W. (1954). The criticism of transformation, unpublished manuscript.
- Begun, J. M. (1981). *A Class of Rank Estimators of Relative Risk*, Institute of Statistics Mimeo Series No. 1381, University of North Carolina at Chapel Hill.
- Begun, J. M. and Wellner, J. A. (1983). Asymptotic efficiency of relative risk estimates, *Contributions to Statistics: Essays in Honor of Norman L. Johnson*, (ed. P. K. Sen), North-Holland, Amsterdam.
- Begun, J. M., Hall W. J., Huang, W. M. and Wellner, J. A. (1983). Information and asymptotic efficiency in parametric—nonparametric models, *Ann. Statist.*, **11**, 432–452.
- Bennett, S. (1983). Log-logistic regression models for survival data, *Appl. Statist.*, **32**, 165–171.
- Bickel, P. J. (1986). Efficient testing in a class of transformation models, Tech. Report, University of California, Berkeley.
- Bickel, P. J. and Doksum, K. A. (1981). An analysis of transformations revisited, *J. Amer. Statist. Assoc.*, **76**, 293–311.
- Box, G. E. P. and Cox, D. R. (1964). An analysis of transformation, *J. Roy. Statist. Soc. Ser. B*, **26**, 211–252.
- Clayton, D. and Cuzick, J. (1986). The semiparametric Pareto model for regression analysis of survival times, *Proc. ISI*, Amsterdam.
- Cox, D. R. (1972). Regression models and life tables, *J. Roy. Statist. Soc. Ser. B*, **34**, 187–202.
- Cox, D. R. (1975). Partial likelihood, *Biometrika*, **62**, 269–276.
- Dabrowska, D. M. and Doksum, K. A. (1988). Partial likelihood in transformation models with censored data, *Scand. J. Statist.*, **15**, 1–23.
- Doksum, K. A. (1987). An extension of partial likelihood methods for proportional hazard model to general transformation models, *Ann. Statist.*, **15**, 325–345.



- Doksum, K. A. and Nabeya, S. (1984). Estimation in proportional hazard and log-linear models, *J. Statist. Plann. Inference*, **9**, 297–303.
- Efron, B. (1977). The efficiency of Cox's likelihood function for censored data, *J. Amer. Statist. Assoc.*, **72**, 557–565.
- Ferguson, T. S. (1967). *Mathematical Statistics*, Academic Press, New York.
- Harrington, D. P. and Fleming, T. R. (1982). A class of rank test procedures for censored survival data, *Biometrika*, **69**, 533–546.
- Hodges, J. L. Jr. and Lehmann, E. L. (1963). Estimates of location based on rank tests, *Ann. Math. Statist.*, **34**, 598–611.
- Huber, P. J. (1981). *Robust Statistics*, Wiley, New York.
- Lehmann, E. L. (1953). The power of rank tests, *Ann. Math. Statist.*, **24**, 23–43.
- Miura, R. (1985). Hodges-Lehmann type estimates and Lehmann's alternative models: A special lecture presented at the annual meeting of Japanese Mathematical Society, the Division of Statistical Mathematics, April, 1985 (in Japanese).
- Pettitt, A. N. (1984). Proportional odds model for survival data and estimates using ranks, *J. Roy. Statist. Soc. Ser. C*, **33**, 169–175.
- Ruymgaart, F. H., Shorack, G. R. and Van Zwet, W. R. (1972). Asymptotic normality of nonparametric tests for independence, *Ann. Math. Statist.*, **43**, 1122–1135.
- Savage, I. R. (1956). Contributions to the theory of rank order statistics: The two sample case, *Ann. Math. Statist.*, **31**, 415–426.
- Shorack, G. R. (1972). Functions of order statistics, *Ann. Math. Statist.*, **43**, 412–427.
- Tukey, J. W. (1957). On the comparative anatomy of transformations, *Ann. Math. Statist.*, **28**, 602–632.
- Wellner, J. A. (1986). Semiparametric models; progress and problems, *Proc. ISI*, Amsterdam.
- Wong, W. H. (1986). Theory of partial likelihood, *Ann. Statist.*, **14**, 88–123.