

## CASE REPORT

Consolato Sergi · Jürgen Weitz · Walter J. Hofmann  
 Peter Sinn · Andrea Eckart · Gerd Otto  
 Philipp A. Schnabel · Herwart F. Otto

## ***Aspergillus* endocarditis, myocarditis and pericarditis complicating necrotizing fasciitis**

### **Case report and subject review**

Received: 30 April 1996 / Accepted: 3 July 1996

**Abstract** A patient with *Aspergillus* endocarditis, myocarditis and pericarditis is described. A 55-year-old man developed necrotizing fasciitis of the lower abdominal wall, pelvis and right thigh. Despite aggressive surgical débridement and antibiotic coverage, the patient died of multisystem organ failure. Autopsy revealed *Aspergillus* thromboulcerative endocarditis, myocarditis and pericarditis, acute necrotizing fungal bronchopneumonia and mycotic dissemination to brain, kidney and thyroid gland. A review of the literature showed that in the absence of open-heart surgery *Aspergillus* endocarditis and myocarditis are very uncommon.

**Key words** Necrotizing fasciitis · Endocarditis · Myocarditis · Epicarditis · *Aspergillus*

### **Introduction**

Necrotizing fasciitis is a relatively rare life-threatening soft tissue infection, frequently caused by a combination of multiple aerobic and anaerobic bacteria and characterized by rapidly spreading neutrophilic and lymphomonocytic cell inflammation, necrosis of the superficial fascia and subcutaneous tissue, and fibrinous necrosis of arterial and venous walls [32]. The release of exotoxins and endogenous cytokines by microorganisms trigger

Dedicated to Prof. Dr. G. Seifert, emeritus Director of the Institute of Pathology, University of Hamburg, on the occasion of his 75th birthday

C. Sergi · W.J. Hofmann · P. Sinn · P.A. Schnabel  
 H. F. Otto (✉)  
 Institute of Pathology, University of Heidelberg,  
 D-69120 Heidelberg, Germany  
 Tel.: (49) 6221-56 26 00, Fax: (49) 6221-56 52 51

J. Weitz · G. Otto  
 Department of Surgery, University of Heidelberg, Heidelberg,  
 Germany

A. Eckart  
 Institute of Neuropathology, University of Heidelberg,  
 Heidelberg, Germany

systemic effects, including renal, heart, and respiratory failure [29, 36]. In this milieu, patients with necrotizing fasciitis may be at elevated risk for overwhelming fungal infections.


Here we describe the postmortem findings of a patient affected with necrotizing fasciitis who presented with *Aspergillus* endo-, myo- and pericarditis, acute necrotizing fungal bronchopneumonia and mycotic dissemination to brain, kidney and thyroid gland.

### **Clinical history**

A 55-year-old man, who was non-diabetic and negative for human immunodeficiency virus (HIV), was admitted to a local hospital for treatment of a perianal abscess. Surgical drainage was performed on the same day and a first-generation cephalosporin therapy was commenced. On the 2nd postoperative day the patient developed fever, and cultures of débrided tissue revealed *Staphylococcus aureus* and *Escherichia coli*, but without pulmonary manifestation by chest X-ray. On the 5th postoperative day the patient became hypotensive and was transferred to the Department of Surgery at Heidelberg University Hospital. Physical examination and computed tomography scanning showed necrotizing fasciitis of the lower abdominal wall, pelvis, perineum, retroperitoneum and proximal internal regions of the right thigh. Débridement, drainage and a sigmoidostomy were performed. Tissue cultures disclosed *Escherichia coli* and *Enterococci*, and necrotizing fasciitis was confirmed by histological investigation. Third-generation cephalosporins and aminoglycosid were started, and three consecutive surgical revisions with extensive débridement were performed. Despite maximum supportive therapy the patient developed bradyarrhythmia on the 19th postoperative day and died of multisystem organ failure. No serum precipitating antibodies to antigens of *Aspergillus* species were found on the 7th and 12th postoperative days.

### **Pathological findings**

At autopsy, there was an extensive, phlegmonous necrotizing soft tissue inflammation with its main focus in the pelvic wall, spreading to the lower abdominal wall, retroperitoneum and perineum and to proximal regions of the right thigh. Microscopically, no bacterial or fungal



**Fig. 1** **a** Section at the level of the transition zone between the left ventricular parietal endocardium and the aortic valve endocardium of the left coronary leaflet, showing an ulcerating and polypoid vegetation with bouquets of 45° angle dichotomous branching septate hyphae, characteristic for *Aspergillus* typus. This invades the underlying myocardium and is disseminating in the bloodstream. GMS,  $\times 50$ . **b** Myocardium of the posterior wall of the left ventricle, demonstrating infected thromboemboli in the muscular tissue. GMS,  $\times 80$ . **c** *Aspergillus* spores on the surface of the epicardium. H & E,  $\times 30$ . **d** Fungal colonies in layers in a bronchial wall. GMS,  $\times 100$

colonies were identified. The heart showed fibrinous epicarditis and marked dilatation of all chambers. A area of small ulcerating and polypoid vegetation was seen at the level of the transition zone from the left ventricular parietal endocardium to the aortic valve endocardium of the left coronary leaflet. The mitral, tricuspid and pulmonary valves were unremarkable. The anterior and posterior walls of both ventricles were mottled by multiple haemorrhagic lesions and small foci of fibrosis. Histologically, at all cardiac levels bouquets of 45° angle dichotomous branching septate hyphae were detected. These were characteristic for *Aspergillus* spp. and gave a positive reaction with PAS and GMS stains but a negative one with the Gram stain (Fig. 1a–c). Numerous infected thromboemboli were seen in the intramural coronary arteries, with coagulation necrosis of the adjacent myocardium in all chambers, in the interventricular septum and

in the anterior papillary muscle. Both lungs had evidence of acute necrotizing fungal bronchopneumonia (Fig. 1d), but no *Aspergillus* spores were localized in the mediastinum. Mucosa-associated lymphatic tissue in the lung was markedly reduced. Numerous accumulations of branching fungi of the *Aspergillus* were also seen in the thyroid gland, in the left kidney, where a fungal thrombus obliterated an intralobular artery with subsequent infarction of the parenchyma supplied, and in the brain, which displayed a multifocal subacute fungal vasculitis. Micronodular cirrhosis and a splenic tumour were also found.

## Discussion

The incidence of *Aspergillus* endocarditis has increased notably over the past few decades in step with the greater frequency of cardiovascular surgery [33]. In contrast, *Aspergillus* endocarditis in patients who have not had cardiac surgery is extremely rare. Woods et al. collected only 28 cases described in the literature and added 1 of their own [42]. Our computer-assisted review of the literature revealed 10 additional cases of *Aspergillus* endocarditis without open-heart surgery [1, 3, 7, 8, 12, 14, 18, 19, 28, 35]. To the best of our knowledge only 1 other case of mycotic dissemination by *Aspergillus* endocarditis has been reported in a patient affected with necrotizing fasci-

**Table 1** Diseases that may be complicated by *Aspergillus* endocarditis in 40 patients with no history of cardiac surgery (IVDA intravenous drug abuse)

Primary disease	No. of cases	References
Haematological malignancy	13	[4, 14, 16, 18, 23, 25, 27, 28, 30, 35, 40, 41]
Other malignancies	2	[31, 40]
HIV infection and/or IVDA	5	[8, 12, 13, 19, 24]
Organ transplantation <sup>a</sup>	3	[15, 40, 42]
Chronic alcoholism	2	[5, 11]
Tuberculosis <sup>b</sup>	2	[22, 38]
Necrotizing fasciitis	1	Present case
Other <sup>c</sup>	6	[6, 17, 20, 40, 43]
None	6	[1, 3, 7, 10, 37, 39]

<sup>a</sup> Includes 2 kidney transplant recipients [15, 40] and 1 liver transplant recipient [42]

<sup>b</sup> Includes 1 patient with miliary tuberculosis [22] and 1 with sili-cotuberculosis [38]

<sup>c</sup> Includes 1 patient each with prematurity and "nonsterile" exchange transfusion [20], uveitis [17], mediastinal fibrosis [6], fulminant hepatitis [40], rheumatic heart disease [43], and polytrauma [43]

itis [9]. In that report the authors described the clinical findings without necroscopic evidence of a 60-year-old woman afflicted with diabetes mellitus and chronic renal failure requiring haemodialysis, who developed a fatal *Aspergillus* infection following débridement for necrotizing fasciitis.

Table 1 shows the primary diseases that may be complicated by *Aspergillus* endocarditis in patients without a history of cardiac surgery. The micronodular cirrhosis present in our patient suggests chronic alcohol intake, and alcoholic subjects are considered to have a greater predisposition to infectious diseases than other people. It is probable that in our patient alcohol abuse was the most important predisposing condition, a necessary but not sufficient factor for the development of *Aspergillus* endocarditis. Zimmerman attributed the dissemination of fungus in the body to a combination of three factors, all of which were present in this case. These are a general lowering of resistance by a debilitating disease, a local point of entry for the fungus, and the use of antibiotics, causing suppression of the normal bacterial flora and inflammatory response [43, 44]. Subsequent studies have supported this and emphasized the role of steroids in the suppression of inflammatory response [21, 33].

*Aspergillus* spp. are airborne organisms that are ubiquitous in the environment. It therefore seems probable that in our patient the route of entry could have been the respiratory tract, with subsequent spread. An attempt at cinical detection of the fungus was unsuccessful, but this is not surprising since blood cultures fail to reveal *Aspergillus* or are positive in less than 11% of patients with cardiac aspergillosis, broncho-alveolar lavage with culture discloses only approximately 50% of those with pulmonary aspergillosis, and *Aspergillus* antibody tests have limited value in immunocompromised patients [2, 21, 33, 42].

With regard to the course of the necrotizing fasciitis, it is difficult to say whether the outcome would have been any different if the diagnosis had been made rapidly. Upon arrival at the hospital, the patient was already extremely ill from the systemic manifestations of the disease. The best outcome in patients with necrotizing fasciitis has been associated with extensive operative procedures in the first 24–48 h after onset of the illness [26]. This patient was already at an advanced state of the illness at the time of presentation. Although the soft tissue infection and the mediators of systemic illness were removed, a complete reversal of the multiorgan failure was not possible, because disseminated aspergillosis dominated the postoperative course.

**Acknowledgements** We wish to thank Mrs. Wegener for the computer-assisted review of the literature and for assisting with data analysis, Mr. J Moyers for preparing the photographs, Dr. M. Meier for the critical review of the paper and Mrs. U. Sturm and Mrs. J. Scherer for their expert technical assistance. In addition, the authors would like to extend their thanks to Dr. A. Nabokov for his help in translating the Russian literature.

## References

- Aisenfarb JC, Dupre-Minet M, Asseman P, Chammas E, Lesenne M, Thery C (1993) Endocardite aspergillaire primitive. Arch Mal Coeur Vaiss 86: 259–261
- Blanc P, Hoffman P, Michaels JF, Bernard E, Vinti H, Morand P, Loubière R (1990) L'atteinte cardiaque des patients porteurs du virus de l'immunosuppression humaine (VIH). A propos de 38 cas. Ann Cardiol Angeiol (Paris) 39: 519–525
- Bogner JR, Lufll S, Middeke M, Spengel F (1990) Erfolgreiche medikamentöse Therapie einer Aspergillen-Endokarditis. Dtsch Med Wochenschr 115: 1833–1837
- Buchbinder NA, Roberts WC (1972) Active infective endocarditis confined to mural endocarditis. A study of six necropsy patients. Arch Pathol 93: 435–440
- Caplan HI, Frisch E, Houghton JD, Climo MS, Natsios GA (1968) *Aspergillus fumigatus* endocarditis. Report of a case diagnosed during life. Ann Intern Med 68: 378–385
- Cohen DM, Goggans EA (1971) Sclerosing mediastinitis and terminal valvular endocarditis caused by fungus suggestive of *Aspergillus* species. Am J Clin Pathol 56: 91–96
- Cortet B, Richard R, Deprez X, Lucet L, Flipo RM, Le Loët X, Duquesnoy B, Delcambre B (1994) *Aspergillus* spondylodiscitis: successful conservative treatment in 9 cases. J Rheumatol 21: 1287–1291
- Cox JN, di Dio F, Pizzolato GP, Lerch R, Pochon N (1990) *Aspergillus* endocarditis and myocarditis in a patient with the acquired immunodeficiency syndrome (AIDS). A review of the literature. Virchows Arch [A] 417: 255–259
- Falsey AR, Goldsticker RD, Ahern MJ (1990) Fatal subcutaneous aspergillosis following necrotizing fasciitis: a case report. Yale J Biol Med 63: 9–13
- Garcia Gomez R, Valdespino Estrada A, Lopez Ortiz R (1981) Endocarditis por *Aspergillus*. Informe de un caso tratado quirurgicamente con éxito. Arch Inst Cardiol Mex 51: 549–553
- Grigor'ev BA (1984) Generalized aspergillosis (in Russian). Arkh Patol 46: 63–66
- Henochowicz S, Mustafa M, Lawrinson WE, Pistole M, Lindsay J (1985) Cardiac aspergillosis in acquired immune deficiency syndrome. Am J Cardiol 55: 1239–1240
- Hofman P, Gari-Toussaint M, Bernard E, Michiels JF, Gibelin P, Le Fichoux Y, Morand P, Loubière R (1992) Myocardites fungiques au cours du syndrome d'immunodéficience acquise. Arch Mal Coeur Vaiss 85: 203–208

14. Hurley J, McGovern E (1994) Rupture of a sinus of Valsalva aneurysm due to *Aspergillus* endocarditis. *J Cardiovasc Surg* 35: 75–77
15. Kaplan R, Duncalf D, Cizmar S (1981) *Aspergillus* pancarditis and cardiac arrest during anesthesia. *Anesth Analg* 60: 440–444
16. Kirschstein RL, Sidransky H (1956) Mycotic endocarditis of the tricuspid valve due to *Aspergillus flavus*. *Arch Pathol* 62: 103–106
17. Kotwal MR, Rinchhen CZ (1981) Primary aspergillosis with multisystem dissemination (letter). *Lancet* I: 562
18. Kuijper PMM, Kuijper EJ, Tweel JG van den, Lelie J van der (1992) *Aspergillus fumigatus*, a rare cause of fatal coronary artery occlusion. *Infection* 20: 45–47
19. Light JT Jr, Hendrickson M, Sholes WM, Portnoy DA, Bell WH III, Kerstein MD (1991) Acute aortic occlusion secondary to *Aspergillus* endocarditis in an intravenous drug abuser. *Ann Vasc Surg* 5: 271–275
20. Luke JL, Bolande RP, Gross S (1963) Generalized aspergillosis and *Aspergillus* endocarditis in infancy. Report of a case. *Pediatrics* 31: 115–122
21. Medical Staff Conference (1971) The spectrum of fungal endocarditis. *Calif Med* 115: 34–40
22. Meyer RD, Fox ML (1973) *Aspergillus* endocarditis. Therapeutic failure of amphotericin B. *Arch Intern Med* 132: 102–106
23. Mikulski SM, Love LJ, Bergquist EJ, Hargadon MT, Applefeld MM, Mergner W (1979) *Aspergillus* vegetative endocarditis and complete heart block in a patient with acute leukemia. *Chest* 76: 473–476
24. Mullen P, Jude C, Borkon M, Porterfield J, Walsh TJ (1986) *Aspergillus* mural endocarditis. Clinical and echocardiographic diagnosis. *Chest* 90: 451–452
25. Nishiura T, Miyazaki Y, Oritani K, Tominaga N, Tomiyama Y, Katagiri S, Kanayama Y, Yonezawa T, Tarui S, Yamada T, Sakurai M, Kume H, Okudaira M (1986) *Aspergillus* vegetative endocarditis complicated with schizocytic hemolytic anemia in a patient with acute lymphocytic leukemia. *Acta Haematol* 76: 60–62
26. Patiño JF, Castro D (1991) Necrotizing lesions of soft tissues: a review. *World J Surg* 15: 235–239
27. Peterson SP, Schiller N, Stricker RB (1984) Failure of two-dimensional echocardiography to detect *Aspergillus* endocarditis. *Chest* 85: 291–294
28. Ramos Fernandez V, Prieto Rodriguez M, Paradis Alos A, Lopez Chulia F, Salom Fuster JV, Vera Sempere FJ (1993) *Aspergilosis* diseminada angioinvasiva: diagnostico necropsico en pacientes leucosicos. *An Med Interna* 10: 337–340
29. Rea WJ, Wyrick WJ Jr (1970) Necrotizing fasciitis. *Ann Surg* 172: 957–964
30. Ribera JM, Ferrer O, Blade J, Valls V, Carreras E, Granena A, Montserrat E, Rozman C (1983) Endocarditis por *Aspergillus* en un paciente con leucemia aguda linfoblastica en remision. *Med Clin (Barc)* 81: 436–439
31. Riou B, Rimailho A, Sinico M, Richard C, Auzepy P (1985) Endocardite droite au cours d'une aspergillose disseminée. *Ann Med Interne (Paris)* 136: 398–400
32. Rouse TM, Malangoni MA, Schulte WJ (1982) Necrotizing fasciitis: a preventable disaster. *Surgery* 92: 765–770
33. Rubinstein E, Lang R (1995) Fungal endocarditis. *Eur Heart J* 16: 84–89
34. Rubinstein E, Noriega ER, Simberkoff MS, Holzman R, Rahal JJ (1975) Fungal endocarditis: analysis of 24 cases and review of the literature. *Medicine (Baltimore)* 54: 331–344
35. Schwartz DA (1989) *Aspergillus* pancarditis following bone marrow transplantation for chronic myelogenous leukemia. *Chest* 95: 1338–1339
36. Stevens DL (1992) Invasive group A streptococcus infections. *Clin Infect Dis* 14: 2–11
37. Swensson EE, Willman VL, Peterson GJ (1986) Acute aortic occlusion from aspergillosis in a healthy patient with survival. *J Vasc Surg* 4: 187–191
38. Vasin VA, Papkov VG, Libiainen TP, Vasin IV (1987) Aspergillotic septic endocarditis of the mitral and trigeminal valves as a complication of silicotuberculosis (in Russian). *Arkh Patol* 49: 50–53
39. Vishniavsky N, Sagar KB, Markowitz SM (1983) *Aspergillus fumigatus* endocarditis on a normal heart valve. *South Med J* 76: 506–508
40. Walsh TJ, Hutchins GM (1979) *Aspergillus* mural endocarditis. *Am J Clin Pathol* 71: 640–644
41. Welsh RA, Buchness JM (1955) *Aspergillus* endocarditis, myocarditis and lung abscesses. Report of a case. *Am J Clin Pathol* 25: 782–786
42. Woods GL, Wood RP, Shaw BW Jr (1989) *Aspergillus* endocarditis in patients without prior cardiovascular surgery: report of a case in a liver transplant recipient and review. *Rev Infect Dis* 11: 263–272
43. Zimmerman LE (1950) *Candida* and *Aspergillus* endocarditis with comments on the role of antibiotics in dissemination of fungus disease. *Arch Pathol* 50: 591–605
44. Zimmerman LE (1955) Fatal fungus infections complicating other diseases. *Am J Clin Pathol* 25: 46–65