

*Short Communication***Construction and Properties of a New Cloning Vehicle, Allowing Direct Screening for Recombinant Plasmids**

Ulrich Rüter

Institute for Genetics, University of Cologne, Weyertal 121, D-5000 Cologne 41, Federal Republic of Germany

Summary. pUR2, a certified B2(EK2) vector, allows easy isolation of variants containing cloned *EcoRI*-fragments. Bacteria harboring plasmids without inserts make blue colonies on indicator-plates, whilst those harboring recombinant plasmids make white colonies.

Introduction

In most gene cloning experiments replica-plating (Lederberg and Lederberg, 1952) has to be used to detect recombinant plasmids. In this communication I describe the construction of a cloning vehicle allowing screening for recombinant plasmids in one step. This was achieved by combining a reduced pBR322-plasmid (Bolivar et al., 1977) with a fragment containing the *lac*-regulatory region and part of the *Z*-gene.

Bacteria, carrying the regulatory region and the *Z*-gene of the *lac*-operon, can be induced by isopropyl-thiogalactoside (IPTG) to express β -galactosidase. This enzyme hydrolyses the colourless 5-bromo-4-chloro-indolyl- β -D-galactoside (x-gal) to a blue insoluble indigo derivative (Davies and Jacob, 1968). A small deletion in the *Z*-gene (M15) lacking the codons for amino acid residues 11–41 (Langley et al., 1975a) destroys the ability of β -galactosidase to hydrolyze x-gal. Ullmann, Jacob and Monod (1967) reported that the α -peptide (amino acid residues 3–92 of β -galactosidase, Langley et al., 1975b) has donor activity in intracistronic complementation and leads to reactivation of defective M15- β -galactosidase. α -complementation can be demonstrated by blue colonies on indicator-plates containing IPTG and x-gal.

Gronenborn and Messing (1978) have introduced a unique *EcoRI*-site in M13mp1 at a position corresponding to the fifth amino acid of the α -peptide.

M13mp1 is a M13-derivative carrying the *lac*-promotor, operator and part of the *Z*-gene coding for α -peptide (Messing et al., 1977). The name of the modified phage is M13mp2; cloning in the *EcoRI*-site of this phage leads often to inactivation of the α -peptide i.e. always when nonsense codons or frameshifts are introduced or created. Bacteria-strains carrying the deletion Δ M15 transformed with such hybrid phages show white colonies (plaques) on x-gal containing indicator-plates. In that way recombinants can be detected easily and the expression of the cloned DNA can be controlled by the *lac*-regulatory region. Because the phage M13mp2 has not yet been certified as a B2 safety vector for cloning experiments I constructed a plasmid with similar advantages for recombinant-screening.

Methods and Results

Cleaving of pBR322 (Bolivar et al., 1977) with *EcoRI* and *PvuII* results in two fragments, a 2067bp fragment containing the gene coding for tetracycline resistance and a 2294bp fragment containing the ampicillin resistance gene and the origin of replication (Sutcliffe, 1978). The *EcoRI*-site was destroyed during incubation of the cleaved DNA for about one hour with DNA-polymerase I in presence of all four nucleotides. The DNA was then treated with T4-DNA-ligase to close the 2294bp fragment to a circle. *E. coli* strain CSH51 ($\Delta(lac, pro) ara, thi, strA, \phi 80dlac^+$; Miller, 1972) was transformed using a slightly modified method of Mandel and Higa (1970). Plasmids were prepared from ampicillin resistant and tetracycline sensitive bacteria. Plasmid-DNA isolated in that way shows a mobility in agarose gels equivalent to approximately 2300bp. The plasmid, called pUR1, is *EcoRI* resistant and can be cleaved once by *PstI* and *HindIII* and twice by *HaeII*.




Fig. 1. Restriction analysis of pUR2 by agarose gel electrophoresis. The agarose concentration in the gel was 2%. Lanes: 1, pUR2 digested with *EcoRI*; 2, digested with *TaqI*; 3, digested with *TaqI* and *EcoRI*; 4, digested with *HaeII*; 5, digested with *HaeII* and *EcoRI*; 6, digested with *HindII* and *EcoRI*; 7, digested with *PstI* and *EcoRI*; 8, an *AvaI*, *PstI* and *PvuII* digest of pBR322 was used as fragment-size-marker

In the *lac*-region of M13mp2 (Gronenborn and Messing, 1978) are two *HaeII*-sites, one at the end of the *I*-gene and the other at the position corresponding to amino acid residue 59 in the α -peptide-part of the *Z*-gene. Digestion of M13mp2 with *HaeII* produces this *lac*-fragment (391bp, W. Gilbert, pers. communication) and 5 other fragments of different sizes. To clone the 391 bp fragment in pUR1, M13mp2-DNA was cleaved with *HaeII*. pUR1 was partially digested with *HaeII* and then mixed with the digested M13mp2-DNA. After treatment with T4-DNA-ligase strain CSH51 was transformed with the ligated DNA. Bacteria were plated on ampicillin-x-gal-plates and blue colonies were picked for further analysis. Blue colonies were expected to arise when more than 20 copies of *lac*-operator-containing plasmid titrated out *lac*-repressor. Initially it was not known whether the first 59 amino acids of β -galactosidase were sufficient for donor activity in the α -complementation-test, therefore a *Lac*⁺-strain was used for the selection. When strain BMH 71-18 ($\Delta(lac, pro) thi, F'lac^{\Delta}Z\Delta M15, pro^+$) was transformed by plasmids isolated from blue colonies, all colonies were blue on ampicillin-x-gal-IPTG-plates. DNA-analysis (Fig. 1) indicated that only the 391bp fragment is cloned in pUR1. This shows that the first 59 amino acids of β -galactosidase are sufficient for α -peptide activity in this plasmid. The plasmid is approximately 2700bp large in agarose gels. This new plasmid is called pUR2.




Fig. 2. A circular restriction map of pUR2. The position of restriction sites are drawn on a circular map divided into 0.1 kilobases

I have cloned different *EcoRI*-fragments in pUR2; recombinant plasmids can be detected in white colonies of BMH 71-18 on ampicillin-x-gal-IPTG-plates. Cloning of some small DNA-fragments (smaller than 100bp) led to light blue colonies. These small inserts may not change the reading frame of the α -peptide and may not code for stop-codons in the same frame.

The mobilisation-frequency of pUR2 was compared to pBR322. Both plasmids have the same mobilisation-frequency (data not shown). As a result pUR2 was certified as a B2 (EK2) vector in West-Germany. For use as a B2 (EK2)-system a suitable host-strain has been constructed (a modified χ 1776 (Curtiss et al., 1977) carrying the *lac Z* deletion Δ M15) and this strain has been submitted for certification.

Acknowledgements. I wish to express my thanks to P. Betteridge, B. Gronenborn and B. Müller-Hill for critically reading the manuscript. This work was supported by a grant from Deutsche Forschungsgemeinschaft through SFB74 to B. Müller-Hill.

References

- Bolivar, F., Rodriguez, R.L., Greene, P.J., Betlach, H.C., Heynacker, H.K., Boyer, H.W.: Construction and characterisation of a new cloning vehicle II. A multipurpose cloning system. *Gene* 2, 95-113 (1977)
- Curtiss III, R., Pereira, D.A., Hsu, J.C., Hull, S.C., Clark, J.E., Maturin, L.J., Sr., Goldschmidt, R., Moddy, R., Inoue, M., Alexander, L.: In: Recombinant molecules, Miles International Symposium Series (eds. Beers, R.F., Bassett, E.G.), pp. 45-56. New York: Raven Press, 1977

- Davies, J., Jacob, F.: Genetic mapping of the regulator and operator genes of the *lac* operon. *J. Mol. Biol.* **36**, 413–417 (1968)
- Gronenborn, B., Messing, J.: Methylation of single-stranded DNA *in vitro* introduces new restriction endonuclease cleavage sites. *Nature* **272**, 375–377 (1978)
- Langley, K.E., Fowler, A.V., Zabin, I.: Amino acid sequence of β -galactosidase IV. Sequence of an α -complementing cyanogen bromide peptide, residues 3 to 92. *J. Biol. Chem.* **250**, 2587–2592 (1975b)
- Langley, K.E., Villarejo, M.R., Fowler, A.V., Zamenhof, P.J., Zabin, I.: Molecular basis of β -galactosidase α -complementation. *Proc. Natl. Acad. Sci. U.S.A.* **72**, 1254–1257 (1975a)
- Lederberg, J., Lederberg, E.M.: Replica plating and indirect selection of bacterial mutants. *J. Bacteriol.* **63**, 399–406 (1952)
- Mandel, M., Higa, A.: Calcium-dependent bacteriophage DNA infection. *J. Mol. Biol.* **53**, 159–162 (1970)
- Messing, J., Gronenborn, B., Müller-Hill, B., Hofschneider, P.H.: Filamentous coliphage M13 as a cloning vehicle: insertion of a *Hind*II fragment of the *lac* regulatory region in M13 replicative form *in vitro*. *Proc. Natl. Acad. Sci. U.S.A.* **74**, 3642–3646 (1977)
- Miller, J.H.: *Experiments in molecular genetics*. New York: Cold Spring Harbor Laboratory 1972
- Sutcliffe, J.G.: pBR322 restriction map derived from the DNA sequence: accurate DNA size markers up to 4361 nucleotide long. *Nucleic Acids Res.* **5**, 2721–2728 (1978)
- Ullmann, A., Jacob, F., Monod, J.: Characterisation by *in vitro* complementation of a peptide corresponding to an operator-proximal segment of the β -galactosidase structural gene of *Escherichia coli*. *J. Mol. Biol.* **24**, 339–343 (1967)

Communicated by E. Bautz

Received February 8, 1980

Note Added in Proof

Variants of pUR2 have been constructed, containing either a *Hind*III- or a *Bam*HI-site in addition to the *Eco*RI-site, allowing the same screening for recombinant plasmids.