

Erratum

Plasma lactate concentration increases as a parabola with delay during ramp exercise

François Péronnet¹, R. Hugh Morton²

¹ Département d'éducation physique, Université de Montréal, CP 6128-A, Montreal, Canada, H3C 3J7

² Department of Statistics, School of Mathematical and Information Sciences, Massey University, PB 11222, Palmerston North, New Zealand

Eur J Appl Physiol (1994) 68:228–233

We unfortunately overlooked two mistakes in Table 1. The values given for a/S were actually those of a , with S (the extramuscular lactate space) assumed to be 18 l. The correct values of a in $\text{mmol} \cdot \text{l}^{-1} \cdot \text{min}^{-2}$ are indicated below. In addition, the fourth column of the table in fact showed the value for the delay δ , in minutes. We apologize for these mistakes, which escaped our attention.

Table 1. Results of the analysis of the lactate curve

Subjects	$[\text{La}^-]_0$ ($\text{mmol} \cdot \text{l}^{-1}$)	a/S ($\text{mmol} \cdot \text{l}^{-1} \cdot \text{min}^{-2}$)	δ (min)	r	$\text{MRS}^{0.5}$ ($\text{mmol} \cdot \text{l}^{-1}$)
1	0.87	0.191	7.17	0.996	0.123
2	0.99	0.115	4.67	0.999	0.082
3	1.27	0.130	5.87	0.994	0.062
4	1.04	0.220	7.70	0.998	0.109
5	0.73	0.147	5.74	0.995	0.103
Mean	0.98	0.161	6.23	—	0.096
SD	0.20	0.044	1.21	—	0.024

$[\text{La}^-]_0$, Lactate concentration at rest; a/S , rate of increase in lactate concentration per litre of extramuscular space; δ , delay between the onset of exercise and the linear increase in R_{e} ; r , correlation coefficient; MRS, mean residual square around the fitting curve