

Methanol metabolism in thermotolerant methylotrophic *Bacillus* strains involving a novel catabolic NAD-dependent methanol dehydrogenase as a key enzyme

N. Arfman¹, E. M. Watling², W. Clement^{1,*}, R. J. van Oosterwijk¹, G. E. de Vries¹, W. Harder¹, M. M. Attwood², and L. Dijkhuizen¹

¹ Department of Microbiology, University of Groningen, NL-9751 NN Haren, The Netherlands

² Department of Microbiology, University of Sheffield, Sheffield S10 2TN, UK

Abstract. The enzymology of methanol utilization in thermotolerant methylotrophic *Bacillus* strains was investigated. In all strains an immunologically related NAD-dependent methanol dehydrogenase was involved in the initial oxidation of methanol. In cells of *Bacillus* sp. C1 grown under methanol-limiting conditions this enzyme constituted a high percentage of total soluble protein. The methanol dehydrogenase from this organism was purified to homogeneity and characterized. In cell-free extracts the enzyme displayed biphasic kinetics towards methanol, with apparent K_m values of 3.8 and 166 mM. Carbon assimilation was by way of the fructose-1,6-bisphosphate aldolase cleavage and transketolase/transaldolase rearrangement variant of the RuMP cycle of formaldehyde fixation. The key enzymes of the RuMP cycle, hexulose-6-phosphate synthase (HPS) and hexulose-6-phosphate isomerase (HPI), were present at very high levels of activity. Failure of whole cells to oxidize formate, and the absence of formaldehyde- and formate dehydrogenases indicated the operation of a non-linear oxidation sequence for formaldehyde via HPS. A comparison of the levels of methanol dehydrogenase and HPS in cells of *Bacillus* sp. C1 grown on methanol and glucose suggested that the synthesis of these enzymes is not under coordinate control.

Key words: *Bacillus* – Methanol – Methylotrophic bacilli – Thermotolerant bacilli – Methylotrophy – Alcohol dehydrogenase – Methanol dehydrogenase – RuMP cycle of formaldehyde fixation – Hexulose-6-phosphate synthase

Over the years it has become clear that Gram-negative methylotrophic bacteria without exception employ a

* Present address: DMV Campina bv, P.O. Box 13, NL-5460 BA Veghel, The Netherlands

Offprint requests to: L. Dijkhuizen

Abbreviations: RuMP, ribulose monophosphate; HPS, hexulose-6-phosphate synthase; HPI, hexulose-6-phosphate isomerase; MDH, methanol dehydrogenase; ADH, alcohol dehydrogenase; PQQ, pyrroloquinoline quinone; DTT, dithiothreitol; NBT, nitroblue-tetrazolium; PMS, phenazine methosulphate; DCPIP, dichlorophenol indophenol

pyrroloquinoline quinone (PQQ) dependent methanol dehydrogenase (MDH; EC 1.1.99.8) for the initial oxidation of methanol (Anthony 1982, 1986; Duine et al. 1978, 1986; de Vries 1986). The situation in Gram-positive methylotrophic bacteria is less clear as relatively few of these organisms have been isolated and studied in detail thus far (Dijkhuizen and Levering 1987). In the methanol-utilizing actinomycete *Nocardia* sp. 239 (Hazeu et al. 1983), the presence of a novel NAD-dependent, PQQ-containing methanol dehydrogenase (nMDH) has been reported (Duine et al. 1984). In this organism nMDH activity resides in a multienzyme complex together with NAD-dependent formaldehyde dehydrogenase and NADH dehydrogenase.

Recently we described the isolation of thermotolerant methylotrophic *Bacillus* strains, employing an NAD-dependent methanol dehydrogenase for methanol oxidation and the RuMP cycle of formaldehyde fixation (Dijkhuizen et al. 1988). NAD-dependent alcohol dehydrogenases (ADH; EC 1.1.1.1) are widely distributed in nature (Sund and Theorell 1963; Brändén et al. 1975) and draw considerable attention, amongst others for phylogenetic reasons (Duester et al. 1986). In general, these enzymes possess a low affinity, if any, for methanol (e.g., Steinbüchel and Schlegel 1984; Rella et al. 1987; Sheehan et al. 1988; Verduyn et al. 1988). One example is the NAD-dependent alcohol dehydrogenase from *Bacillus stearothermophilus* DSM 2334 (Sheehan et al. 1988; Dowds et al. 1988) with apparent K_m values for methanol and ethanol of 20 mM and 82 μ M, respectively. This organism, however, is not able to grow on methanol as a carbon source.

In this paper a detailed study of the enzymology of methanol utilization in thermotolerant methylotrophic *Bacillus* strains, and the purification and characterization of the methanol dehydrogenase from *Bacillus* sp. C1 is presented.

Materials and Methods

Organisms

The organisms used, *Bacillus* sp. C1, PB1, AR2, TS1, TS2 and TS4, and their maintenance have been described before (Dijkhuizen et al. 1988). *Bacillus* sp. S1 and TF are related strains (Al-Awadhi et al. 1988).

Media and cultivation

For batch cultivation conical flasks filled to 25% of the volume with a 50 mM methanol- or 10 mM glucose-mineral salts medium (Dijkhuizen et al. 1988) of pH 7.5 were incubated in a shaking waterbath. If required the medium was supplied with 1 ml/l of a vitamin solution, containing (mg/l): biotin, 100; thiamin · HCl, 100; riboflavin, 100; pyridoxal phosphate, 100; panthotenate, 100; nicotinic acid amide, 100; p-aminobenzoic acid, 20; folic acid, 10; vitamin B12, 10; lipoic acid, 10. The methylotrophic *Bacillus* strains were grown at 50°C and the Gram-negative methylotroph *Xanthobacter* sp. 25a (Croes et al. 1986) at 30°C. For continuous cultivation of *Bacillus* sp. C1 the organism was grown in a methanol-limited (50 mM methanol) chemostat on the mineral medium described by Levering et al. (1981). The temperature was controlled at 50°C and the pH was maintained at 7.3 by automatic adjustment with 1 M NaOH.

Preparation of extracts

Cells from early- to mid-exponential phase of growth were harvested by centrifugation at $3,800 \times g$ for 10 min at 4°C, washed twice and resuspended in 50 mM potassium phosphate buffer pH 7.5, containing 5 mM MgSO₄ (buffer A). Cells were disrupted in the presence of 5 mM dithiothreitol (DTT) by passage through a French pressure cell operating at 1.4×10^5 kN/m². Low-speed cell-free extracts were obtained by centrifugation at $3,800 \times g$ for 10 min at 4°C and used for the determination of NAD(P)H oxidase and NADH dehydrogenase activities. Cell walls and membrane particles were subsequently removed by centrifugation at $25,000 \times g$ for 20 min at 4°C. The supernatants, which contained 5–10 mg of protein/ml, were used as crude extracts for enzyme assays and for the purification of methanol dehydrogenase. For enzyme localization studies extracts were centrifuged at $150,000 \times g$ for 60 min at 4°C, yielding a high speed supernatant and a high speed pellet. The pellet was washed twice with buffer A. Extracts and samples obtained in various purifications steps (see below), were stored at –80°C in the presence of 5 mM DTT.

Enzyme assays

All assays with *Bacillus* sp. C1 and *Bacillus* sp. TS4 were performed at 50°C, using prewarmed buffer solutions.

Methanol oxidation rates of whole cells (early exponential growth phase) were determined polarographically, using a Clark-type oxygen electrode (Biological Oxygen Monitor, Yellow Springs Instruments Co., Yellow Springs, OH, USA). Whole cells rapidly lost methanol oxidizing activity, both on ice and at room temperature. The following procedure was therefore adopted which resulted in reproducible activities. Cells were washed once at room temperature in buffer A and assayed immediately. The reaction mixture (3 ml) contained: potassium phosphate buffer pH 7.5, 150 µmol; MgSO₄, 15 µmol; and cell suspension (0.2–0.5 mg cell dry weight). After air saturation and temperature equilibration at 50°C, the reaction was started with 150 µmol of the alcohol to be tested. At 50°C, 100% air saturation corresponds to 0.187 mMol O₂/l H₂O.

Spectrophotometric assays were performed with a Perkin-Elmer 124 spectrophotometer. The PQQ-containing methanol dehydrogenase (MDH; EC 1.1.99.8), present in

all Gram-negative methylotrophs studied, was assayed according to Ghosh and Quayle (1981), by following DCPIP reduction at 600 nm (molar absorption coefficient of 22×10^3 M⁻¹ · cm⁻¹). The reaction mixture (1 ml) contained: Tris · HCl pH 9.0, 125 µmol; ammonium chloride, 150 µmol; PMS, 1.25 µmol; DCPIP, 80 µmol; potassium cyanide, 5 µmol; extract. The reaction was started by the addition of 2 µmol methanol.

The NAD-dependent, PQQ-containing methanol dehydrogenase (nMDH), as observed in *Nocardia* sp. 239, was assayed by following DCPIP reduction at 600 nm, according to Duine et al. (1984). The reaction mixture (1 ml) contained tetrasodium pyrophosphate buffer pH 9.0, 100 µmol; ammonium chloride, 120 µmol; NAD, 2.5 µmol; DCPIP, 40 nmol; extract. The reaction was started by the addition of 2 µmol methanol.

Various methods were tested to measure activity of the NAD-dependent methanol dehydrogenase in the methylotrophic *Bacillus* strains (see Results section). The following procedure was adopted: Alcohol-dependent NADH production was measured at 50°C, by following the increase in absorbance at 340 nm, in a reaction mixture (1 ml) containing: glycine/KOH buffer pH 9.5, 100 µmol; MgSO₄, 5 µmol; DTT, 1 µmol; NAD, 1 µmol; extract. After temperature equilibration the reaction was started with 500 µmol of the alcohol to be tested. Dilution of extracts resulted in loss of methanol dehydrogenase activity present in *Bacillus* strains, but could be largely circumvented by using buffer A containing 20% sucrose.

The formaldehyde reductase activity (the activity of NAD-dependent methanol dehydrogenase in the reverse direction) was measured by following the formaldehyde-dependent oxidation of NADH at 340 nm. The reaction mixture (1 ml) contained: potassium phosphate buffer pH 6.7, 50 µmol; DTT, 1 µmol; NADH, 150 nmol; extract. After temperature equilibration the reaction was started with 10 µmol formaldehyde.

NADH dehydrogenase activity was measured by following DCPIP reduction at 600 nm in a reaction mixture (1 ml) containing: potassium phosphate buffer pH 8.0, 50 µmol; MgCl₂, 5 µmol; DCPIP, 80 nmol; low-speed cell-free extract. After temperature equilibration at 50°C the reaction was started with 200 nmol NADH.

NAD(P)H oxidase activities were measured at 340 nm in a reaction mixture (1 ml) containing: potassium phosphate buffer pH 7.0, 50 µmol; MgCl₂, 5 µmol; low-speed cell-free extract. The reaction was started with 200 nmol NAD(P)H.

Glucose-6-phosphate isomerase (EC 5.3.1.9) was assayed in a reaction mixture (1 ml) containing: potassium phosphate buffer pH 8.5, 50 µmol; fructose-6-phosphate, 5 µmol; NADP, 0.4 µmol; glucose-6-phosphate dehydrogenase, 1 unit. The reaction was started by the addition of extract.

Hexulose-6-phosphate isomerase activity was determined using a modification of the assay for hexulose-6-phosphate synthase (Levering et al. 1981) in that hexulose-6-phosphate isomerase from *Methylophilus methylotrophus* was omitted.

The following enzymes were assayed according to published methods. NAD-dependent formaldehyde dehydrogenase, EC 1.2.1.1 (van Dijken et al. 1976); NAD-dependent formate dehydrogenase, EC 1.2.1.2, and NAD-independent formate dehydrogenase, EC 1.2.–.– (Dijkhuizen et al. 1978); hexulose-6-phosphate synthase (HPS; spec-

trophotometric assay of Levering et al. 1981); NADP-dependent glucose-6-phosphate dehydrogenase, EC 1.1.1.49, NADP-dependent 6-phosphogluconate dehydrogenase, EC 1.1.1.44, 6-phosphofructokinase, EC 2.7.1.11, combined activity of 6-phosphogluconate dehydratase, EC 4.2.1.12, and KDPG aldolase, EC 4.1.2.14 (van Dijken and Quayle 1977); fructose-1,6-bisphosphate aldolase, EC 4.1.2.13 (van Dijken et al. 1978); phosphoriboisomerase, EC 5.3.1.6, ribulose phosphate 3-epimerase, EC 5.1.3.1, transketolase, EC 2.2.1.1, transaldolase, EC 2.2.1.2, sedoheptulose-1,7-bisphosphatase, EC 3.1.3.66 (Levering et al. 1982).

Purification of methanol dehydrogenase from Bacillus sp. C1

Methanol dehydrogenase was purified from cells grown at a low dilution rate ($D = 0.10 \text{ h}^{-1}$) in a methanol-limited chemostat ($S_R = 50 \text{ mM}$ methanol). The viscosity of the crude extract was lowered by adding protamine sulphate (10% solution) to a final concentration of 1 mg per 10 mg protein. The resulting precipitate (nucleic acids) was removed by centrifugation at $25,000 \times g$ for 20 min at 4°C . The supernatant was desalted in buffer B (20 mM Tris · HCl pH 7.5/5 mM MgSO_4 /5 mM β -mercaptoethanol) by Sephadex G25 gel filtration (Bio-Rad PD10 column) and subjected to anion-exchange chromatography. The sample was applied onto a Fast Flow Q-Sepharose column (volume 25 ml; washed and equilibrated with buffer B), connected to a Pharmacia FPLC system. Proteins were eluted at a flow rate of 1.0 ml/min by applying a linear 0–1.0 M KCl gradient in buffer B. Formaldehyde reductase containing fractions (4 ml) were pooled and prepared for hydrophobic interaction chromatography by adding ammonium sulphate to a final concentration of 1 M. Samples (13–14 mg of protein) were subsequently applied onto a FPLC Phenyl-Superose HR5/5 column (Pharmacia), equilibrated with 50 mM Tris · HCl pH 7.0/5 mM MgSO_4 /5 mM β -mercaptoethanol containing 1.0 M $(\text{NH}_4)_2\text{SO}_4$ (buffer C). Bound protein was eluted at a flow rate of 0.4 ml/min with a linear 1.0–0 M $(\text{NH}_4)_2\text{SO}_4$ gradient in buffer C. Fractions of 0.5 ml were collected. The active fractions were pooled and stored in small aliquots at -80°C .

Determination of relative molecular mass

The molecular weight of native methanol dehydrogenase was determined on a Superose 6 gel filtration column (equilibrated with 100 mM Tris · HCl pH 7.5/5 mM MgSO_4 /5 mM β -mercaptoethanol), using gel filtration standards (1,350–670,000 range) of Bio-Rad, Richmond, USA. Samples (200 μl , containing approximately 200 μg of protein) of both crude extract and purified methanol dehydrogenase were used.

Electrophoretic methods

Denaturing gel electrophoresis (SDS-PAGE) was performed as described by Laemmli and Favre (1973) with the following calibration proteins as molecular weight references: phosphorylase A, 94,000; humane transferrin, 80,000; bovine serum albumin, 68,000; catalase, 58,000; fumarase, 50,000; citrate synthase, 46,000; carboanhydrase, 31,000. Gels were stained with Coomassie brilliant blue G-250. Western blotting was performed as described by Towbin et al. (1979) using the alkaline phosphatase-linked-antibody

procedure. Following gel electrophoresis under non-denaturing conditions alcohol dehydrogenase from baker's yeast (Boehringer, Mannheim, FRG), the NAD-dependent methanol dehydrogenase from *Bacillus* sp. C1 and the PQQ-dependent methanol dehydrogenase from *Xanthobacter* sp. 25a (Janssen et al. 1987) were visualized by an alcohol-dependent tetrazolium activity staining. The enzyme preparations were loaded on a 6% polyacrylamide gel in 100 mM glycine/Tris pH 9.0. After electrophoresis (100 V, constant current), the gels were stained by incubation with methanol (1%) or ethanol (1%) in staining buffer at 45°C . Staining buffer contained (per 100 ml): glycine · KOH pH 9.5, 10 mmol; MgSO_4 , 0.5 mmol; NAD, 0.1 mmol; nitroblue-tetrazolium (NBT), 20 mg; PMS, 8 mg.

Protein determination

Protein was determined by the methods of Lowry et al. (1951; low-speed cell-free extracts only) and Bradford (1976) using bovine serum albumin as a standard.

Formaldehyde and PQQ determination

Formaldehyde was assayed with the method of Avigad (1983). PQQ was determined by measuring the reconstitution of quinoprotein alcohol dehydrogenase from *Pseudomonas testosteroni* (Groen et al. 1986).

Preparation of antiserum

Antibodies against purified methanol dehydrogenase (Phenyl-Superose pool, Table 1; Fig. 2, lane m) were raised in rabbits by subcutaneous injections (500 μg) in complete Freund's adjuvant. The specificity of the antibodies were tested by Ouchterlony double diffusion tests and Western blotting.

Electron microscopy and immunocytochemistry

Electron microscopy and immunogold labeling of NAD-dependent methanol dehydrogenase were performed as described by Douma et al. (1985).

Results

Methanol oxidation

Cells of *Bacillus* sp. C1 grown on methanol in batch culture oxidized methanol at a high rate (1,100–1,500 $\text{nmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ of protein). Only short chain (C1–C4) primary alcohols were oxidized, but compared to methanol (100%) at lower relative rates (ethanol, 90%; n-propanol, 57%, n-butanol, 53%). Secondary alcohols and formate were not oxidized. While the organism oxidized various alcohols, it only grew on methanol. Most probably, *Bacillus* sp. C1 is not able to metabolize aldehydes other than formaldehyde.

Reduced (dithionite) minus oxidized (ferricyanide) cytochrome spectra of low-speed cell-free extracts of methanol- or glucose-grown *Bacillus* sp. C1 revealed the presence of cytochromes of the c- (552 nm), b- (558 nm) and a-type (603 nm). Potassium cyanide was a potent inhibitor of methanol oxidation (> 90% inhibition by 2 μM in low-speed extracts). Interestingly, also antimycin A acted as an inhibi-

tor of methanol oxidation, both in whole cells (50% inhibition by 60 μM antimycin A) and in low-speed cell-free extracts (> 90% inhibition of both NADH and methanol oxidation by 65 μM antimycin A). These observations indicate that electrons derived from methanol oxidation in *Bacillus* sp. C1 are donated to the electron transport chain at, or before, the level of cytochrome b. In contrast, the PQQ-dependent methanol dehydrogenase in Gram-negative methylotrophs interacts with the electron transport chain at the level of cytochrome c (Anthony 1982).

All attempts to measure an ammonium-requiring, dye-linked (PMS/DCPIP) methanol dehydrogenase in extracts of the methylotrophic bacilli failed. Also activity of an NAD-dependent, dye-linked (DCPIP), PQQ-containing methanol dehydrogenase, using the assay as developed for the actinomycete *Nocardia* sp. 239 (Duine et al. 1984), could not be detected. Incubation of extracts of methanol-grown cells of *Bacillus* sp. C1 with methanol nevertheless resulted in production of formaldehyde (700–800 $\text{nmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ of protein), provided methanol was present at relatively high concentrations (approx. 500 mM). When using these much higher methanol concentrations in the assay as developed for *Nocardia* sp. 239, we also were able to detect methanol-dependent DCPIP reduction (360 $\text{nmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ of protein) by extracts of *Bacillus* sp. C1. Omission of NAD from the two assays described above had no effect on the level of activity. Both activities, however, were completely lost when extracts were desalted on a Sephadex G25 column, but could be effectively restored by adding NAD to the reaction mixture. It thus became apparent that NAD in fact was required for methanol oxidation by *Bacillus* sp. C1 and that this had been masked because of the presence of sufficiently high levels of NAD in the extracts. Extracts of *Bacillus* sp. C1 subsequently were found to catalyze a methanol-dependent reduction of NAD that could be measured directly at 340 nm. Maximal activities were observed at methanol concentrations of 500 mM. This activity was ammonium-independent, but was strongly stimulated by addition of Mg^{2+} -ions (5 mM) to the assay buffer. Under optimized conditions (i.e., at 50°C and pH 9.5), extract of cells of *Bacillus* sp. C1 grown in batch culture were found to possess methanol dehydrogenase activities of 1,000–1,200 $\text{nmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ of protein. These high methanol dehydrogenase activities, however, were only observed with undiluted extracts, after preincubation (5 min) of samples and buffers at the assay temperature of 50°C. Starting the reaction with NAD instead of methanol resulted in prolonged acceleration phases and lower final activities. Dilution of samples generally resulted in loss of methanol dehydrogenase activities (only about 10% of the specific activity remained for instance when extract was diluted tenfold in assay buffer) but could be largely overcome by including sucrose in the buffer solutions used. The methanol dehydrogenase activity was not proportional to the amount of extract added to the assay mixtures. Rather, the specific activity increased with increasing protein concentrations. This effect was also observed when sucrose or glycerol were included in the assay buffer. The extracts also catalyzed the reverse reaction, i.e. NADH-dependent formaldehyde reduction (1,200 $\text{nmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ protein). The formaldehyde reductase reaction was not sensitive to dilution inactivation, neither was it affected by Mg^{2+} -ions. Moreover, the formaldehyde reductase activity was clearly proportional to the amount of protein

added. We therefore decided to use this assay during enzyme purification and for localization studies.

Following gel electrophoresis of extracts of methanol-grown cells of *Bacillus* sp. C1 and *Xanthobacter* sp. 25a, and *Saccharomyces cerevisiae* ADH under non-denaturing conditions, the alcohol dehydrogenases present were visualized by staining for activity with methanol and ethanol as substrates. With both substrates the *Bacillus* sp. C1 activity was found to reside in a single band, the electrophoretic mobility of which was clearly different from that of either yeast ADH or methanol dehydrogenase from *Xanthobacter* sp. 25a (Fig. 1). The active band from *Bacillus* sp. C1 extract was excised from the gel and subjected to SDS-PAGE. A single band of approximately 43,000 was observed (results not shown).

Purification of methanol dehydrogenase from Bacillus sp. C1

The methanol dehydrogenase was purified to homogeneity from cells of *Bacillus* sp. C1 grown in chemostat culture under methanol-limitation at $D = 0.10 \text{ h}^{-1}$. Extracts of these cells, subjected to SDS-PAGE, possessed a very dominant 43,000 protein band, which suggested that the methanol dehydrogenase constituted already a high percentage of total cell protein. The purification (shown in Table 1 and Fig. 2) was followed with both the formaldehyde reductase and methanol dehydrogenase assays. During ion-exchange chromatography the formaldehyde reductase eluted at 300 mM KCl, resulting in a 2.3-fold increase in specific activity. In this step the NADH dehydrogenase activity eluted at 500 mM KCl. Hydrophobic interaction chromatography turned out to be an effective second purification step, with formaldehyde reductase eluting at relatively low ionic strength (300 mM $(\text{NH}_4)_2\text{SO}_4$), indicating that the protein is rather hydrophobic in nature. The final preparation, after purification to homogeneity (Fig. 2, lane m), possessed a specific formaldehyde reductase activity of 19.6 $\mu\text{mol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ of protein. The 4.5-fold purification obtained indicates that the enzyme may constitute up to about 22% of total soluble protein at $D = 0.10 \text{ h}^{-1}$ in methanol-limited chemostats, assuming that no inactive enzyme is present in the final preparation. The molecular weight of purified native enzyme was estimated to be 300,000 \pm 30,000 by gel filtration chromatography. Most likely it is a hexamer or octamer with subunits of 43,000.

The methanol dehydrogenase and formaldehyde reductase activities were stable in cell-free extracts. Storage during 4 days at 4°C, –20°C or –80°C in the absence of protease inhibitors did not result in a significant loss of enzyme activity, provided DTT was present (< 5% loss). However, during purification of the enzyme a remarkable difference was observed between the recoveries of methanol dehydrogenase and formaldehyde reductase activities. After the Q-Sepharose and Phenyl-Superose column steps, 34% of the original formaldehyde reductase activity was recovered whereas the preparation had almost completely lost methanol dehydrogenase activity (> 95% loss). Addition of sucrose, glycerol, iron ions, zinc ions or DTT did not prevent this specific loss of methanol dehydrogenase activity.

Pyrroloquinoline quinone analysis

Nocardia sp. 239 was found to excrete PQQ into the culture medium during growth on methanol (Hazeu et al. 1983).




Fig. 1. Zymogram of extracts of *Xanthobacter* sp. 25a (PQQ-MDH; lane a; *Bacillus* sp. C1, lanes b, d) and purified baker's yeast ADH (lanes c, e). Samples were loaded on a native polyacrylamide gel (6%) and stained with methanol (lanes a–c) or ethanol (d–e) as the substrate

Fig. 2A, B. SDS-polyacrylamide (10%) gel electrophoresis of extracts of eight different methanol-grown *Bacillus* strains (A), and samples obtained during purification of methanol dehydrogenase from cells of *Bacillus* sp. C1 grown under methanol-limiting conditions ($D = 0.10 \text{ h}^{-1}$) in continuous culture (B). A Lanes a–j: *Bacillus* sp. C1, AR2, PB1, TS1, TS2, TS4, marker proteins (see Methods, minus carboanhydrase), *Bacillus* sp. S1, TF, respectively. B Lanes k–o: Crude extract, Q-Sepharose pool, Phenyl-Superose pool, marker proteins (see Methods), baker's yeast ADH

Table 1

Purification of methanol dehydrogenase from *Bacillus* sp. C1

Sample	Protein (mg)	Total activity (units) ^a	Specific activity (units/mg)	Recovery (%)	Fold purification
Crude Extract	320	1,408	4.4	100	1.0
Q-Sepharose pool	54.6	541	9.9	38	2.3
Phenyl-Superose pool	24.6	482	19.6	34	4.5

^a Formaldehyde reductase assay

One unit of activity is defined as the amount of enzyme catalyzing the oxidation of 1 μmol of NADH per min

This organism employs a novel MDH whose activity is both PQQ- and NAD-dependent (Duine et al. 1984). The possible involvement of PQQ in methanol oxidation in *Bacillus* sp. C1 was studied by analyzing extracts and purified formaldehyde reductase for the presence of this cofactor. Neither the purified enzyme nor the extract contained PQQ. Only if PQQ is covalently bound to the enzyme can its presence have been overlooked. However, the assay is very sensitive and PQQ is a very stable compound. It thus appears likely that methanol oxidation in *Bacillus* sp. C1 is performed by a PQQ-independent, NAD-dependent methanol dehydrogenase.

Kinetics of methanol oxidation

Oxidation rates of washed cell suspension of *Bacillus* sp. C1 at various methanol concentrations followed normal Michaelis-Menten kinetics. A K_s value of 2.6 mM methanol was calculated from Eadie-Hofstee plots (Fig. 3A).

Compared to whole cells, the methanol dehydrogenase activity in extracts (and partially purified preparations) displayed a similar substrate specificity (C1–C4 primary alcohols only), although its relative activity towards methanol was considerably lower (methanol, 33%; ethanol, 100%; n-propanol, 71%; n-butanol 87%). The enzyme re-

quired NAD(H) as coenzyme, no activity was observed with NADP(H).

Because purified preparations lost methanol dehydrogenase activities, the kinetics of methanol oxidation was further studied in extracts. In extracts the enzyme displayed biphasic kinetics towards methanol (apparent K_m values of 3.8–166 mM; Fig. 3B) and ethanol (apparent K_m values 7.5 and 70 mM; not shown). Biphasic kinetics was not observed with whole cells of *Bacillus* sp. C1 (see above). Thus, methanol oxidation in vivo becomes saturated at lower methanol concentrations, which results in a relatively low K_s value. Biphasic kinetics was also observed towards NAD (apparent K_m values 15 and 190 μM), but not for formaldehyde (apparent K_m value 2.0 mM) in the reverse reaction.

Subcellular localization of methanol dehydrogenase

The subcellular localization of methanol dehydrogenase and NADH dehydrogenase was studied by comparing the levels of these enzymes in cell-free extracts, in high speed supernatants (soluble fractions) and in high speed pellets (membrane fractions). Approximately 70% of the NADH dehydrogenase activity appeared to be membrane associated, while the remaining activity was recovered in the supernatant. Most probably, NADH dehydrogenase has a loose




Fig. 3. Eadie-Hofstee plots of methanol oxidation by whole cells (A), and NAD-dependent methanol dehydrogenase activities in extracts (B; 40 μ g of protein per assay) of *Bacillus* sp. C1




Fig. 4. Immunogold labeling of methanol dehydrogenase in whole cells of methanol-grown *Bacillus* sp. C1. Magnification: $\times 38,500$

association with the cytoplasmic membrane and part of the enzyme is solubilized during cell rupture. NADH oxidase activity, as a control, was exclusively found in the membrane fraction (90% recovery). The supernatant contained more than 90% of the total formaldehyde reductase activity. These results suggest that the methanol dehydrogenase in *Bacillus* sp. C1 is a soluble cytoplasmic enzyme. The possibility remained, however, that the enzyme was loosely bound to the inner surface of the cytoplasmic membrane. This was further tested by immunogold labeling experiments. Gold labeling of the methanol dehydrogenase in *Bacillus* sp. C1 resulted in a scattering of gold particles over the cytoplasm (Fig. 4), thus providing unambiguous evidence that it functions as a soluble enzyme *in vivo*.

Enzyme profiles in *Bacillus* sp. C1 and TS4

SDS-PAGE of extracts of 8 methylotrophic bacilli grown on methanol in batch culture revealed (Fig. 2) that a protein species of approx. 43,000 dominates in all organisms (slightly smaller in size in *Bacillus* sp. TF). Western blots with antibodies directed against the purified methanol dehydrogenase from *Bacillus* sp. C1 showed that all methylotrophic *Bacillus* strains studied harboured an immunologically related methanol dehydrogenase with subunits of approximately 43,000 (not shown). High activities of NAD-dependent methanol dehydrogenase and the key enzymes of the RuMP cycle, hexulose-6-phosphate synthase (HPS) and hexulose-6-phos-

phate isomerase (HPI), were present in all methylotrophic *Bacillus* strains under investigation. The enzyme data obtained with *Bacillus* sp. C1 and TS4 are shown in Table 2. A comparison of the levels of methanol dehydrogenase and HPS in glucose and methanol-grown cells of *Bacillus* sp. C1 suggests that the synthesis of these enzymes is not regulated coordinately. No activities of NAD-(in)dependent formaldehyde or formate dehydrogenases could be detected in *Bacillus* strains C1 and TS4. This, and the failure of methanol-grown whole cells to oxidize formate, indicated that the organisms do not possess a linear pathway for the oxidation of methanol to CO_2 . Energy generation presumably proceeds via the so-called dissimilatory RuMP cycle, involving the enzymes glucose-6-phosphate isomerase, glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase, all of which were present at high activities. The dehydrogenases involved were strictly NAD-dependent, although the high NADPH oxidase and negligible NADPH oxidase activities indicate that NADH is the preferred electron donor for the electron transport chain in these organisms. The enzymes of the Entner-Doudoroff pathway were absent, whereas high activities were detected of 6-phosphofructokinase and FBP aldolase, indicating that the organisms employ the Embden-Meyerhof route for sugar phosphate cleavage. Regeneration of the formaldehyde acceptor molecule ribulose-5-phosphate, from fructose-6-phosphate and glyceraldehyde-3-phosphate, involved the enzymes transketolase, transaldolase, phosphoriboisomerase and ribulose phosphate 3-epimerase (TK/TA rearrangement variant). Sedoheptulose-1,7-bisphosphatase activity could not be detected. The pathway of methanol metabolism in the *Bacillus* strains investigated is shown schematically in Fig. 5.

Discussion

Although the presence of NAD-dependent alcohol dehydrogenases (ADH; EC 1.1.1.1) has been reported in some methylotrophic bacteria, e. g. in *Hyphomicrobium* X (Brooke and Attwood 1983) and in organism PAR (Bellion and Wu 1978), this enzyme only appeared to function in ethanol catabolism. All Gram-negative methylotrophic bacteria studied employ a PQQ-containing methanol dehydrogenase (MDH, EC 1.1.99.8; Duine et al. 1978) for methanol oxidation. In this group of organisms MDH is located in the periplasmic space. Evidence is emerging that in Gram-positive bacteria, which do not possess a periplasmic space, methanol oxidation is catalyzed by enzymes clearly different from MDH. Previously, Duine et al. (1984), reported the involvement of an NAD-dependent, PQQ-containing methanol dehydrogenase in the actinomycete *Nocardia* sp. 239. Although this enzyme requires NAD for activity, methanol oxidation did not result in the production of free NADH. In the present study a cytoplasmic, NAD-dependent methanol dehydrogenase was identified as the methanol-oxidizing enzyme in *Bacillus* strains. No evidence was obtained for the presence of PQQ in *Bacillus* sp. C1 and methanol oxidation clearly resulted in production of free NADH. The characteristic properties of MDH, nMDH, ADH and the NAD-dependent methanol dehydrogenase as observed in *Bacillus* sp. C1 are summarized in Table 3. The data indicate that the methylotrophic bacterium *Bacillus* sp. C1 employs a novel NAD-dependent methanol dehydrogenase for the conver-

Table 2
Specific activities
(nmol · min⁻¹ · mg⁻¹ of
protein) of various enzymes
in extracts of methanol-
grown *Bacillus* sp. C1 and
Bacillus sp. TS4

Enzyme	Activity in	
	<i>Bacillus</i> sp. C1	<i>Bacillus</i> sp. TS4
Methanol oxidation rate (whole cells)	1,100–1,500 (75) ^a	1,100–1,500
1. Formaldehyde reductase	1,220 (880) ^a	1,265
Formaldehyde dehydrogenase	nd	nd
Formate dehydrogenase	nd	nd
2. Hexulose-6-phosphate synthase	17,300 (515) ^a	19,400
3. Hexulose-6-phosphate isomerase	5,950	6,700
4. Glucose-6-phosphate isomerase	1,415	1,480
5. Glucose-6-phosphate dehydrogenase (NADP)	475 (135) ^a	930
6. 6-Phosphogluconate dehydrogenase (NADP)	385	670
7. 6-Phosphofructokinase	600	670
8. Fructose-1,6-bisphosphate aldolase	370	1,190
9. Transketolase	710 (460) ^a	1,110 (140) ^a
10. Transaldolase	60 (9) ^a	140 (60) ^a
11. Phosphoriboisomerase	80	105
12. Ribulose phosphate 3-epimerase	775	650
KDPG aldolase/6-PG dehydratase	nd	nd
NADH oxidase	630	590
NADPH oxidase	25	30
NADH dehydrogenase	740	510

^a Glucose grown cells
nd not detectable




Fig. 5. Schematic representation of methanol metabolism in thermotolerant methylotrophic *Bacillus* strains. The various reactions, indicated by numbers, are catalyzed by the enzymes listed in Table 2. Abbreviations: *DHAP*, dihydroxyacetone-3-phosphate; *GAP*, glyceraldehyde-3-phosphate; *E4P*, erythrose-4-phosphate; *Ru5P*, ribulose-5-phosphate; *Ri5P*, ribose-5-phosphate; *X5P*, xylulose-5-phosphate; *H6P*, hexulose-6-phosphate; *FBP*, fructose-1,6-bisphosphate; *S7P*, sedoheptulose-7-phosphate

sion of methanol into formaldehyde. Inhibition studies revealed that electrons from methanol oxidation are donated to the electron transport chain at, or above, the level of cytochrome b. This also explains the relatively high growth yield of the *Bacillus* strains on methanol (Dijkhuizen et al. 1988).

In general the rate of an enzymatic reaction is favoured when reaction products are immediately further metabolized. This becomes especially important when thermodynamically unfavourable reactions have to be carried out. Multienzymes, as well as enzyme complexes are then thought to play a role (Srere 1987). The redox potential of the couple methanol/formaldehyde ($E^{\circ} = -182$ mV) strongly favours the formaldehyde reductase reaction. It is therefore not sur-

prising that methanol oxidation in methylotrophic bacteria apparently requires the presence of specially adapted enzymes. In Gram-negative strains, a modifier protein is involved in the regulation of MDH activity by decreasing its affinity for formaldehyde, which is also a substrate for the enzyme in vitro (Page and Anthony 1986). In *Nocardia* sp. 239, an enzyme complex is formed between the nMDH and NAD-dependent formaldehyde dehydrogenase and NADH dehydrogenase, which further metabolize the reaction products (Duine et al. 1984). The results obtained in the present study show that *Bacillus* sp. C1 employs a novel methanol dehydrogenase with a relatively low affinity for methanol. The question therefore has to be asked how the organism manages to maintain the high turnover rate of methanol into formaldehyde required for fast growth on the C₁ compound methanol (Attwood and Quayle 1984). First of all, the problem of low affinity of the methanol dehydrogenase for methanol is at least partly overcome by synthesizing high levels of the enzyme. In methanol-limited cells, the enzyme may constitute up to 22% of total soluble protein ($D = 0.10$ h⁻¹). Secondly, the high HPS as well as NADH dehydrogenase activities observed in these bacteria may well ensure that in vivo the methanol dehydrogenase reaction is favoured over the formaldehyde reductase reaction. Maintenance of a relatively high internal pH would also pull the reaction towards formaldehyde and NADH.

Various observations indicate that the methanol dehydrogenase present in *Bacillus* sp. C1 in fact is (part of) a rather complex system. In extracts, but not with whole cells, biphasic kinetics was observed for methanol oxidation. During enzyme purification it was noticed that the enzyme rapidly lost methanol dehydrogenase activity. Moreover, the dehydrogenase activity of the enzyme, but not the reductase activity, appeared to be very sensitive to dilution. Dilution inactivation could be largely overcome via osmotic stabilization, which might be an indication that the methanol dehydrogenase in *Bacillus* sp. C1 resides in a protein complex. None of the MDH and ADH enzymes described so far require Mg²⁺-ions for activity. Since the reverse reaction does not depend on the presence of Mg²⁺-ions, it appears likely

Table 3. Comparison of characteristic properties of alcohol dehydrogenases from various sources

	Methanol dehydrogenase in methylotrophic bacteria			ADH from <i>Saccharomyces cerevisiae</i> (EC 1.1.1.1)
	Gram-negative (MDH; EC 1.1.99.8)	Gram-positive		
		<i>Nocardia</i> sp. 239	<i>Bacillus</i> sp. Cl	
<i>Substrates</i>				
Alcohols	Lower primary alcohols	Methanol	Lower primary alcohols	Lower primary alcohols, but not methanol
Formaldehyde	+	—	—	—
<i>Assay</i>	PMS/DCPIP (A_{600})	NAD/DCPIP (A_{600})	NAD (A_{340})	NAD (A_{340})
NH ₄ ⁺ -requirement	+	+	—	—
Mg ²⁺ -stimulation	—	—	+	—
NAD-dependence	—	+	+	+
PQQ-cofactor	+	+	—	—
K_m [methanol]	5–30 μ M	nd	3.8–166 mM	nu
<i>Localization</i>	Periplasm	Cytoplasm?	Cytoplasm	Cytoplasm
<i>Structure</i>	1 or 2 subunits of 56,000–76,000	nd	6–8 subunits of 43,000	4 subunits of 40,000

nu, not utilized

nd, not determined

that Mg²⁺ is not active at the catalytic site but instead involved in activation of the methanol dehydrogenase reaction. When using the optimized methanol dehydrogenase assay described in the Methods section, it was noted that the reaction velocity did not increase linearly with the amount of extract or partially purified enzyme. This phenomenon was observed with methanol as well as ethanol as a substrate. In contrast, the reductase reaction always displayed a linear relationship between velocity and enzyme concentration.

Further studies will be required in order to clarify at least some of these points.

Acknowledgements. The research was carried out in the framework of contract No. BAP-0267-NL of the Biotechnology Action Programme of the Commission of the European Communities and the Programme Committee for Industrial Biotechnology (The Netherlands). E. M. Watling was supported by the SERC Biotechnology Directorate. Thanks are due to P. W. van Ophem (Delft University of Technology) for performing the PQQ assays, K. Sjollem and M. Veenhuis (University of Groningen) for performing the electron microscopy and immunochemistry studies, H. R. Moezelaar for performing some of the enzyme assays, and D. R. Kremer for valuable discussions.

References

- Al-Awadhi N, Egli T, Hamer G (1988) Growth characteristics of a thermotolerant methylotrophic *Bacillus* sp. (NCIB 12522) in batch culture. *Appl Microbiol Biotechnol* 29:485–493
- Anthony C (1982) The biochemistry of methylotrophs. Academic Press, London
- Anthony C (1986) Bacterial oxidation of methane and methanol. *Adv Microb Physiol* 27:113–210
- Attwood MM, Quayle JR (1984) Formaldehyde as a central metabolite of methylotrophic metabolism. In: Crawford RL, Hanson RS (eds) *Microbial growth on C₁ compounds*. American Society for Microbiology, Washington DC, pp 315–323
- Avigad G (1983) A simple spectrophotometric determination of formaldehyde and other aldehydes: Application to periodate-oxidized glycol systems. *Anal Biochem* 134:499–504
- Bellion E, Wu GTS (1978) Alcohol dehydrogenases from a facultative methylotrophic bacterium. *J Bacteriol* 135:252–258

- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
- Brändén CI, Jörnvall H, Eklund H, Furugren B (1975) Alcohol dehydrogenases. In: Boyer PD (ed) *The enzymes*, 3rd edn, vol 11A. Academic Press, New York, pp 103–190
- Brooke AG, Attwood MM (1983) Regulation of enzyme synthesis during growth of *Hyphomicrobium* X on mixtures of methylamine and ethanol. *J Gen Microbiol* 129:2399–2404
- Croes GM, Levering PR, Dijkhuizen L (1986) Regulation of methanol metabolism and carbon dioxide fixation in *Xanthobacter autotrophicus* strain 25a. *Abstr 5th Int Symp on Microbial Growth on C₁ Compounds*. Haren. Free University Press, Amsterdam, p 13
- Dijken JP van, Quayle JR (1977) Fructose metabolism in four *Pseudomonas* species. *Arch Microbiol* 114:281–286
- Dijken JP van, Oostra-Demkes GJ, Otto R, Harder W (1976) S-Formylglutathione: the substrate for formate dehydrogenase in methanol-utilizing yeasts. *Arch Microbiol* 111:77–84
- Dijken JP van, Harder W, Beardsmore AJ, Quayle JR (1978) Dihydroxyacetone: an intermediate in the assimilation of methanol by yeasts? *FEMS Microbiol Lett* 4:97–102
- Dijkhuizen L, Knight M, Harder W (1978) Metabolic regulation in *Pseudomonas oxalaticus* OX1. Autotrophic and heterotrophic growth on mixed substrates. *Arch Microbiol* 116:77–83
- Dijkhuizen L, Levering PR (1987) Metabolic regulation in facultative methylotrophs. In: Verseveld HW van, Duine JA (eds) *Microbial growth on C₁ compounds*. Nijhoff, Dordrecht, pp 95–104
- Dijkhuizen L, Arfman N, Attwood MM, Brooke AG, Harder W, Watling EM (1988) Isolation and initial characterization of thermotolerant methylotrophic *Bacillus* strains. *FEMS Microbiol Lett* 52:209–214
- Douma AC, Veenhuis M, de Koning W, Evers M, Harder W (1985) Dihydroxyacetone synthase is localized in the peroxisomal matrix of methanol-grown *Hansenula polymorpha*. *Arch Microbiol* 143:237–243
- Dowds BCA, Sheehan MC, Bailey CJ, McConnell DJ (1988) Cloning and characterization of the gene for a methanol-utilising alcohol dehydrogenase from *Bacillus stearothermophilus*. *Gene* 68:11–22
- Duester G, Jörnvall H, Hatfield GW (1986) Intron-dependent evolution of the nucleotide-binding domains within alcohol dehydrogenase and related enzymes. *Nucleic Acids Res* 14:1931–1941

- Duine JA, Frank J, Westerling J (1978) Purification and properties of methanol dehydrogenase from *Hyphomicrobium* X. *Biochim Biophys Acta* 524:277–287
- Duine JA, Frank J, Berkhout MPJ (1984) NAD-dependent, PQQ-containing methanol dehydrogenase: a bacterial dehydrogenase in a multienzyme complex. *FEBS Lett* 168:217–221
- Duine JA, Frank J, Jongejan JA (1986) PQQ and quinoprotein enzymes in microbial oxidations. *FEMS Microbiol Rev* 32:165–178
- Ghosh R, Quayle JR (1981) Purification and properties of the methanol dehydrogenase from *Methylophilus methylotrophus*. *Biochem J* 199:245–250
- Groen B, Kleef MAG van, Duine JA (1986) Quinohaemoprotein alcohol dehydrogenase apoenzyme from *Pseudomonas testosteroni*. *Biochem J* 234:611–615
- Hazeu W, Bruyn JC de, Dijken JP van (1983) *Nocardia* sp. 239, a facultative methanol utilizer with the ribulose monophosphate pathway of formaldehyde fixation. *Arch Microbiol* 135:205–210
- Janssen DB, Keuning S, Witholt B (1987) Involvement of a quinoprotein alcohol dehydrogenase and an NAD-dependent aldehyde dehydrogenase in 2-chloroethanol metabolism in *Xanthobacter autotrophicus* GJ10. *J Gen Microbiol* 133:85–92
- Laemmli UK, Favre K (1973) Maturation of the head of bacteriophage T4. I. DNA packaging events. *J Mol Biol* 80:575–599
- Levering PR, Dijken JP van, Veenhuis M, Harder W (1981) *Arthrobacter* P1, a fast growing versatile methylotroph with amine oxidase as a key enzyme in the metabolism of methylated amines. *Arch Microbiol* 129:72–80
- Levering PR, Dijkhuizen L, Harder W (1982) Enzymatic evidence for the operation of the FBP aldolase cleavage and TK/TA rearrangement variant of the RuMP cycle in *Arthrobacter* P1. *FEMS Microbiol Lett* 14:257–261
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265–275
- Page MD, Anthony C (1986) Regulation of formaldehyde oxidation by the methanol dehydrogenase modifier proteins of *Methylophilus methylotrophus* and *Pseudomonas* AM1. *J Gen Microbiol* 132:1553–1563
- Rella R, Raia CA, Pensa M, Pisani FM, Gambacorta A, De Rosa M, Rossi M (1987) A novel archaeobacterial NAD⁺-dependent alcohol dehydrogenase. Purification and properties. *Eur J Biochem* 167:475–479
- Sheehan MC, Bailey CJ, Dowds BCA, McConnell DJ (1988) A new alcohol dehydrogenase, reactive towards methanol, from *Bacillus stearothermophilus*. *Biochem J* 252:661–666
- Srere PA (1987) Complexes of sequential metabolic enzymes. *Annu Rev Biochem* 56:89–124
- Steinbüchel A, Schlegel HG (1984) A multifunctional fermentative alcohol dehydrogenase from the strict aerobe *Alcaligenes eutrophus*: purification and properties. *Eur J Biochem* 141:555–564
- Sund H, Theorell H (1963) Alcohol dehydrogenases. In: Boyer PD, Lardy H, Myrbäck K (eds) *The enzymes*, 2nd edn, vol 7A. Academic Press, New York, pp 25–83
- Towbin H, Staehelin T, Gordon J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc Natl Acad Sci USA* 76:4350–4354
- Verduyn C, Breedveld GJ, Scheffers WA, Dijken JP van (1988) Substrate specificity of alcohol dehydrogenase from the yeasts *Hansenula polymorpha* CBS 4732 and *Candida utilis* CBS 621. *Yeast* 4:143–148
- Vries GE de (1986) Molecular biology of bacterial methanol oxidation. *FEMS Microbiol Rev* 39:235–258

Received January 30, 1989/Accepted April 28, 1989