

# Hausdorff Measure and the Navier-Stokes Equations

Vladimir Scheffer

Department of Mathematics, Stanford University, Stanford, California 94305, USA

**Abstract.** Solutions to the Navier-Stokes equations are continuous except for a closed set whose Hausdorff dimension does not exceed two.

## 1. Informal Statement of Results

Let  $v: R^3 \rightarrow R^3$  be a divergence free, square integrable vector field on 3-space. We will show that there exists a function  $u: R^3 \times R^+ \rightarrow R^3$  ( $R^+ = \{t: t > 0\}$  is time) which is a weak solution to the Navier-Stokes equations of incompressible fluid flow with viscosity = 1 and initial conditions  $v$ , and which satisfies the following: There exists a set  $S \subset R^3 \times R^+$  such that the two dimensional Hausdorff measure of  $S$  is finite,  $(R^3 \times R^+) - S$  is an open set, and the restriction of  $u$  to  $(R^3 \times R^+) - S$  is a continuous function.

The above will be derived as a consequence of a more general theorem in which  $u$  satisfies a weak form of the Navier-Stokes equations with an external force  $f: R^3 \times R^+ \rightarrow R^3$  which is divergence free with the property  $f(x, t) \cdot u(x, t) \leq 0$ .

## 2. Notation and Complete Statement of Results

Hausdorff measure is defined in [2, p. 171]. We set  $R^+ = \{t \in R: t > 0\}$  and  $B(a, r) = \{x \in R^3: |x - a| \leq r\}$  for all  $a \in R^3$  and  $r > 0$ . The norm  $|\cdot|$  is always euclidean norm and  $\|\cdot\|_p$  is the  $L^p$  norm. Open and closed intervals are denoted  $(a, b)$  and  $[a, b]$ , respectively. If  $f: X \rightarrow R$  and  $A \subset X$  then  $\sup(f, A)$  is the supremum of  $f$  over  $A$  and  $\text{spt}(f)$  is the closure of  $\{x: f(x) \neq 0\}$ . If  $f$  and  $g$  are functions defined on a subset of  $R^3 \times R$ ,  $h$  is a function on  $R^3$ , and  $k$  is a function on  $R$ , then we set

$$(f * g)(x, t) = \iint f(y, s)g(x - y, t - s)dyds,$$

$$(f * h)(x, t) = \int f(y, t)h(x - y)dy,$$

$$(f * k)(x, t) = \int f(x, s)k(t - s)ds$$

whenever the integrals make sense. If  $X = R^3$ ,  $X = R$ , or  $X = R^3 \times R^+$ , we let  $C^\infty(X, R)$  be the set of infinitely differentiable functions  $f: X \rightarrow R$ . In addition,  $C_0^\infty(X, R)$  is the

set of all functions in  $C^\infty(X, R)$  which are zero outside of some compact set. We also set  $D^\infty(R^3 \times R^+, R) = \{f \in C^\infty(R^3 \times R^+, R) : \text{spt}(f) \subset R^3 \times [a, b] \text{ for some } 0 < a < b < \infty\}$ . If  $f$  is a distribution defined on an open subset of  $R^3 \times R$  then  $D_{i_1}f, D_{i_2}f, \dots$  etc. are the distribution partial derivatives  $(\partial/\partial x_{i_1})f, (\partial^2/\partial x_{i_1}\partial x_{i_2})f$  with respect to the variables  $x_1, x_2, x_3$  of  $R^3$ . The partial derivative of  $f$  with respect to the  $R$  variable of  $R^3 \times R$  is denoted  $D_t f$ . We also set  $Df = (D_1f, D_2f, D_3f), \Delta f = D_{ii}f$  (repeated indices are summed), and  $\text{div}(f) = D_i f_i$  in case the range of  $f$  is  $R^3$ . Similar definitions are made for distributions defined on  $R^3$  and  $R$ .

An absolute constant is a positive constant that is independent of all the parameters in this paper. The letter  $C$  always denotes an absolute constant. The value of  $C$  changes from line to line (e.g.  $2C \leq C$ ). When an absolute constant is denoted by a letter other than  $C$ , its value remains fixed.

The statements below (Parts 1 and 2) are called *Hypothesis I*:

*Part. 1.* We have a Lebesgue measurable function  $u: R^3 \times R^+ \rightarrow R^3$  (a time dependent velocity vector field), a Lebesgue measurable and locally integrable function  $p: R^3 \times R^+ \rightarrow R$  (pressure), and a constant  $0 < L < \infty$  such that

$$\text{div}(u) = 0, \tag{2.1}$$

$$\int_{R^3} |u(x, t)|^2 dx \leq L \text{ for almost every } t \in R^+, \tag{2.2}$$

the distribution  $Du$  is a square integrable function satisfying

$$\int_0^\infty \int_{R^3} |Du(x, t)|^2 dx dt \leq L, \tag{2.3}$$

and for almost every  $t \in R^+$  we have

$$\int_{R^3} p(x, t) \Delta \phi(x) dx = - \int_{R^3} D_i u_i(x, t) D_j u_i(x, t) \phi(x) dx \tag{2.4}$$

if  $\phi \in C^\infty(R^3, R)$ ,  $\phi$  is bounded,  $|x| |\Delta \phi(x)|$  is bounded, and  $\Delta \phi \in C_0^\infty(R^3, R)$ .

*Part 2.* We assume that the conditions

$$\begin{aligned} \phi \in D^\infty(R^3 \times R^+, R); \quad \phi(x, t) \geq 0 \text{ for all } (x, t); \\ \text{and } \phi, D\phi, \Delta \phi + D_t \phi \text{ are bounded} \end{aligned} \tag{2.5}$$

imply that (2.6) holds. Note that (2.2), (2.3), Lemma 3.2, and Lemma 3.6 can be used to show that the integrals in (2.6) exist.

$$-2^{-1} (\int |u|^2 (D_t \phi + \Delta \phi)) + \int |Du|^2 \phi \leq \int u_i (2^{-1} |u|^2 + p) D_i \phi. \tag{2.6}$$

*Hypothesis II* is the following: In addition to Hypothesis I, we assume

$$- \int u_i (D_i \phi + \Delta \phi) = \int u_j u_i D_j \phi + \int p D_i \phi \tag{2.7}$$

for every  $i \in \{1, 2, 3\}$  and  $\phi \in C_0^\infty(R^3 \times R^+, R)$ .

Hypothesis I is a weak form of the classical Navier-Stokes equations

$$D_t u_i = -u_j D_j u_i - D_i p + \Delta u_i + f_i, \quad \text{div}(u) = 0 \tag{2.8}$$

where the external force  $f$  satisfies  $\text{div}(f) = 0$  and  $f(x, t) \cdot u(x, t) \leq 0$ . Hypothesis II is a weak form of (2.8) with  $f = 0$ . We will prove

**2.1. Theorem.** *If Hypothesis I holds then there exist a function  $u' : R^3 \times R^+ \rightarrow R^3$  and a set  $S \subset R^3 \times R^+$  satisfying the following: The functions  $u$  and  $u'$  are equal almost everywhere, the two dimensional Hausdorff measure of  $S$  is finite,  $S \cap \{(x, t) : t \geq \varepsilon\}$  is compact for every  $\varepsilon > 0$ , and  $|u'|$  is bounded on every compact set  $K \subset R^3 \times R^+$  which satisfies  $K \cap S = \emptyset$ .*

The proof of this theorem includes a priori estimates on the size of  $|u'|$ . It is possible to show that the Hausdorff dimension of  $S$  is at most  $7/4$ . We also have

**2.2. Theorem.** *If Hypothesis II holds and  $S$  is as in Theorem 2.1 then there exists a function  $u'' : R^3 \times R^+ \rightarrow R^3$  such that  $u$  and  $u''$  are equal almost everywhere, and  $u''$  is continuous on  $(R^3 \times R^+) - S$ .*

**2.3. Theorem.** *If  $v : R^3 \rightarrow R^3$  is a square integrable function satisfying  $\text{div}(v) = 0$  then there exists  $u$  satisfying Hypothesis II and*

$$\begin{aligned}
 & - \int_{R^3} v_i(x)\phi(x, 0)dx - \int_{R^3 \times R^+} u_i(D_i\phi + \Delta\phi) \\
 & = \int_{R^3 \times R^+} u_j u_i D_j \phi + \int_{R^3 \times R^+} p D_i \phi
 \end{aligned} \tag{2.9}$$

if  $\phi : R^3 \times R \rightarrow R$  is smooth with compact support and  $i \in \{1, 2, 3\}$ .

Here (2.9) states that  $v$  is the initial condition for the solution  $u$ .

This type of partial regularity is similar to that obtained by Almgren for solutions to generalized variational problems [1]. The study of the relationship between Hausdorff measure and the geometry of turbulence was started by Mandelbrot [6].

The next three sections contain the proof of Theorems 2.1 and 2.2. The proof of Theorem 2.3 is outlined in Section 6.

### 3. Preliminary Estimates

Throughout this section we assume that Part I of Hypothesis I holds.

**3.1. Lemma.** *If  $f : R^3 \rightarrow R$ ,  $f \in L^2$ , and  $Df \in L^2$ , then*

- (1)  $\int |f|^6 \leq C(\int |Df|^2)^3$
- (2)  $\int |f|^3 \leq C\varepsilon^{-3}(\int |f|^2)^3 + C\varepsilon(\int |Df|^2)$  whenever  $0 < \varepsilon < \infty$ .

*Proof.* Part (1) is the first inequality in line 9, p. 127 of [9]. We use Hölder's inequality, part (1), and Young's inequality

$$ab \leq (1/4)(\delta^{-1}a)^4 + (3/4)(\delta b)^{4/3} \quad \text{for } a, b \geq 0 \quad \text{and } \delta = \varepsilon^{3/4}$$

to estimate

$$\begin{aligned}
 \int |f|^3 &= \int |f|^{3/2} |f|^{3/2} \\
 &\leq (\int (|f|^{3/2})^{4/3})^{3/4} (\int (|f|^{3/2})^4)^{1/4} \\
 &= (\int |f|^2)^{3/4} (\int |f|^6)^{1/4} \\
 &\leq C(\int |f|^2)^{3/4} (\int |Df|^2)^{3/4} \\
 &\leq C\varepsilon^{-3}(\int |f|^2)^3 + C\varepsilon(\int |Df|^2).
 \end{aligned}$$

**3.2. Lemma.** *If  $0 < T < \infty$  then  $\int_0^T \int_{\mathbb{R}^3} |u(x, t)|^3 dx dt \leq CL^{3/2} T^{1/4}$ .*

*Proof.* Using Lemma 3.1 with  $\varepsilon = L^{1/2} T^{1/4}$ , (2.2), and (2.3), we obtain

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^3} |u(x, t)|^3 dx dt \\ & \leq C\varepsilon^{-3} \left( \int_0^T \left( \int_{\mathbb{R}^3} |u(x, t)|^2 dx \right)^3 dt \right) + C\varepsilon \left( \int_0^T \int_{\mathbb{R}^3} |Du(x, t)|^2 dx dt \right) \\ & \leq CL^{3/2} T^{1/4}. \end{aligned}$$

**3.3. Definition.** We fix  $f_0 \in C_0^\infty(\mathbb{R}^3, \mathbb{R})$  and  $g_0 \in C_0^\infty(\mathbb{R}, \mathbb{R})$  such that  $\text{spt}(g_0) \subset [-1, 1]$ ,  $f_0 \geq 0$ ,  $g_0 \geq 0$ ,  $f_0(x) = f_0(-x)$ ,  $g_0(t) = g_0(-t)$ , and  $\int f_0 = \int g_0 = 1$ . For  $n = 1, 2, 3, \dots$  we set  $f_n(x) = n^3 f_0(nx)$  and  $g_n(t) = n g_0(nt)$ . We let  $A$  consist of all  $t \in \mathbb{R}^+$  such that the function  $p_t(x) = p(x, t)$  is locally integrable, (2.2) and (2.4) hold, the function  $d_t(x) = Du(x, t)$  is square integrable, the divergence of the function  $u_t(x) = u(x, t)$  is zero, and  $\lim_n (|u|^2 * g_n)(x, t) = |u(x, t)|^2$  for almost every  $x \in \mathbb{R}^3$ . Part 1 of Hypothesis I, Fubini's theorem, and [10, Theorem 1.25, p. 13] imply that  $A$  is almost all of  $\mathbb{R}^+$ .

**3.4. Lemma.** *If  $t \in A$ ,  $\alpha \in C_0^\infty(\mathbb{R}^3, \mathbb{R})$ ,  $\beta = 1 - \alpha$ ,  $\beta(x) = 0$  for all  $x$  in a neighborhood of 0,  $\alpha'(x) = -(4\pi|x|)^{-1}\alpha(x)$ , and  $\beta'(x) = -(4\pi|x|)^{-1}\beta(x)$ , then*

$$p(x, t) = -(D_i u_j D_j u_i * \alpha')(x, t) - (u_j u_i * D_{ij} \beta')(x, t)$$

*holds for almost every  $x \in \mathbb{R}^3$ .*

*Proof.* Define  $k: \mathbb{R}^3 - \{0\} \rightarrow \mathbb{R}$  by  $k(x) = -(4\pi|x|)^{-1}$ . Recalling Definition 3.3, we have  $\Delta(k * f_n) = f_n$  (see [9, p. 126]). Hence 3.3 and Part 1 of Hypothesis I yield the following for all  $x \in \mathbb{R}^3$ :

$$\begin{aligned} (p * f_n)(x, t) &= (p * \Delta(k * f_n))(x, t) \\ &= ((-D_i u_j D_j u_i) * (k * f_n))(x, t) \\ &= ((-D_i u_j D_j u_i) * (\alpha' * f_n))(x, t) + ((-D_i u_j D_j u_i) * (\beta' * f_n))(x, t) \\ &= ((-D_i u_j D_j u_i) * (\alpha' * f_n))(x, t) + ((-u_j u_i) * (D_{ij} \beta' * f_n))(x, t). \end{aligned}$$

Since  $\alpha' \in L^1$ ,  $\alpha' * f_n$  converges to  $\alpha'$  in the  $L^1$  norm (see [10, Theorem 1.18, p. 10]). Hence the assumption  $t \in A$  and [10, Theorem 1.3, p. 3] imply

$$\lim_n \int_{\mathbb{R}^3} |((-D_i u_j D_j u_i) * (\alpha' * f_n) - \alpha')(x, t)| dx = 0.$$

Hence [3, (11.26)] implies that there exists a subsequence  $n_k$  of the positive integers such that

$$\lim_k ((-D_i u_j D_j u_i) * (\alpha' * f_{n_k}))(x, t) = ((-D_i u_j D_j u_i) * \alpha')(x, t)$$

holds for almost every  $x \in \mathbb{R}^3$ . Since  $t \in A$  and  $D_{ij} \beta' * f_n$  converges uniformly to  $D_{ij} \beta'$ , we have

$$\lim_k ((-u_j u_i) * (D_{ij} \beta' * f_{n_k}))(x, t) = ((-u_j u_i) * D_{ij} \beta')(x, t)$$

for all  $x \in R^3$ . Finally, [10, Theorem 1.25, p. 13] yields

$$\lim_k (p * f_{h_n})(x, t) = p(x, t) \quad \text{for almost every } x \in R^3.$$

These statements yield the conclusion of the lemma.

**3.5. Lemma.** *If  $t \in A$  and  $0 < r < \infty$  then*

$$\begin{aligned} & \left( \int_{R^3} |p(x, t)|^2 dx \right)^{1/2} \\ & \leq Cr^{1/2} \left( \int_{R^3} |Du(x, t)|^2 dx \right) + Cr^{-3/2} \left( \int_{R^3} |u(x, t)|^2 dx \right). \end{aligned}$$

*Proof.* Given  $r$ , we fix  $\alpha$  and  $\beta$  as in Lemma 3.4 such that  $\beta(x) = 0$  for  $x \in B(0, r)$ ,  $\text{spt}(\alpha) \subset B(0, 2r)$ ,  $0 \leq \alpha(x) \leq 1$ ,  $|D\alpha(x)| \leq Cr^{-1}$ , and  $|D_{ij}\alpha(x)| \leq Cr^{-2}$ . Then we have  $\|\alpha'\|_2 \leq Cr^{1/2}$  and  $\|D_{ij}\beta'\|_2 \leq Cr^{-3/2}$ . Hence Lemma 3.4 and [10, Theorem 1.3, p. 3] yield the conclusion.

**3.6. Lemma.** *If  $0 < T < \infty$  then  $\int_0^T \int_{R^3} |u(x, t)| |p(x, t)| dx dt \leq CL^{3/2} T^{1/4}$ .*

*Proof.* Using the Schwarz inequality, Lemma 3.5 with  $r = T^{1/2}$ , (2.2), and (2.3), we estimate

$$\begin{aligned} & \int_0^T \int_{R^3} |u(x, t)| |p(x, t)| dx dt \\ & \leq \int_0^T \left( \int_{R^3} |u(x, t)|^2 dx \right)^{1/2} \left( \int_{R^3} |p(x, t)|^2 dx \right)^{1/2} dt \\ & \leq CL^{1/2} \left( \int_0^T T^{1/4} \left( \int_{R^3} |Du(x, t)|^2 dx \right) + T^{-3/4} \left( \int_{R^3} |u(x, t)|^2 dx \right) dt \right) \\ & \leq CL^{3/2} T^{1/4}. \end{aligned}$$

**3.7. Lemma.** *If Hypothesis I holds,  $s \in A$ ,  $B = R^3 \times [0, s]$ , and  $\phi$  satisfies (2.5) then*

$$\begin{aligned} & 2^{-1} \int_{R^3} |u(x, s)|^2 \phi(x, s) dx - 2^{-1} \int_B |u|^2 (D_t \phi + \Delta \phi) + \int_B |Du|^2 \phi \\ & \leq \int_B u_t (2^{-1} |u|^2 + p) D_i \phi. \end{aligned}$$

*Proof.* Let  $h_n: R^3 \times R \rightarrow R$  be the function that satisfies  $D_t h_n(x, t) = -g_n(t - s)$  (see Definition 3.3) and  $h_n(x, t) = 1$  for  $t < s - n^{-1}$ . Then  $h_n(x, t) = 0$  for  $t > s + n^{-1}$ . We obtain the conclusion by substituting  $\phi h_n$  for  $\phi$  in (2.6), taking the limit inferior over  $n$ , using Fatou's lemma, and observing that

$$\lim_n \int_{R^+} |u(x, t)|^2 \phi(x, t) g_n(t - s) dt = |u(x, s)|^2 \phi(x, s)$$

holds for almost every  $x \in R^3$  [this is a consequence of  $s \in A$  and the relation  $g_n(t) = g_n(-t)$ ].

**3.8. Lemma.** *If  $f:R^3 \rightarrow R$ ,  $f \in L^2$ , and  $Df \in L^2$  then for every  $a \in R^3$  and  $0 < r < \infty$  we have*

$$\left( \int_{B(a,r)} |f|^4 \right)^{1/2} \leq Cr^{1/2} \left( \int_{B(a,2r)} |Df|^2 \right) + Cr^{-3/2} \left( \int_{B(a,2r)} |f|^2 \right).$$

*Proof.* Let  $g \in C_0^\infty(R^3, R)$  satisfy  $\text{spt}(g) \in B(a, 2r)$ ,  $g(x) = 1$  if  $x \in B(a, r)$ ,  $0 \leq g(x) \leq 1$ , and  $|Dg(x)| \leq Cr^{-1}$ . We apply the Schwarz inequality, Lemma 3.1 (1), Young's inequality [4, p. 11], and the estimate  $|D(fg)| \leq |Df| |g| + |f| |Dg|$  to write

$$\begin{aligned} \int_{B(a,r)} |f|^4 &\leq \int |fg|^4 \\ &= \int |fg|^3 |fg| \\ &\leq (\int |fg|^6)^{1/2} (\int |fg|^2)^{1/2} \\ &\leq C (\int |D(fg)|^2)^{3/2} (\int |fg|^2)^{1/2} \\ &\leq Cr (\int |D(fg)|^2)^2 + Cr^{-3} (\int |fg|^2)^2 \\ &\leq Cr \left( \int_{B(a,2r)} |Df|^2 \right)^2 + Cr^{-3} \left( \int_{B(a,2r)} |f|^2 \right)^2. \end{aligned}$$

#### 4. The Basic Estimate

In this section we assume that Hypothesis I (Parts 1 and 2) holds. The section is devoted to proving the following:

**4.1. Theorem.** *There exist absolute constants  $\varepsilon$  and  $K$  satisfying the following: If  $a \in R^3$ ,  $b \in A$  (see Definition 3.3),  $\gamma > 0$ ,  $b - \gamma^2 > 0$ , and*

$$\int_{b-\gamma^2}^b \int_{R^3} |u(x,t)|(2^{-1}|u(x,t)|^2 + |p(x,t)|(|x-a|+\gamma)^{-4}) dx dt \leq \varepsilon \gamma^{-2} \quad (4.1)$$

then

$$\int_{B(a,\tau\gamma)} |u(x,b)|^2 dx \leq K\tau^3\gamma \quad \text{for } 0 < \tau \leq 1/2. \quad (4.2)$$

We fix  $a \in R^3$ ,  $b \in A$ , and  $\gamma > 0$  with  $b - \gamma^2 > 0$ . For integers  $k$  we set

$$E_k = \{(x,t) : |x-a| < \gamma 2^{-k}, b - \min(\gamma^2 2^{-2k}, \gamma^2) < t < b\}. \quad (4.3)$$

**4.2. Lemma.** *There exist  $\phi_n \in D^\infty(R^3 \times R^+, R)$  for  $n = 1, 2, 3, \dots$  such that  $\phi_n(x,t) \geq 0$ , the functions  $\phi_n$ ,  $D\phi_n$ , and  $\Delta\phi_n + D_t\phi_n$  are bounded,*

$$\phi_n(x,t) = 0 \quad \text{if } t \leq b - \gamma^2, \quad (4.4)$$

$$\sup(\phi_n + \gamma 2^{-n} |D\phi_n|, E_n) \leq C\gamma^{-3} 2^{3n}, \quad (4.5)$$

$$\sup(\phi_n + \gamma 2^{-k} |D\phi_n|, E_k - E_{k+1}) \leq C\gamma^{-3} 2^{3k} \quad \text{if } 0 \leq k < n, \quad (4.6)$$

$$\sup(\phi_n + \gamma 2^{-k} |D\phi_n|, E_k - E_{k+1}) \leq C\gamma^{-3} 2^{4k} \quad \text{if } k < 0, \quad (4.7)$$

$$\sup(|D_t\phi_n + \Delta\phi_n|, E_0) \leq C\gamma^{-5}, \quad (4.8)$$

$$\sup(|D_t\phi_n + \Delta\phi_n|, E_k - E_{k+1}) \leq C\gamma^{-5} 2^{4k} \quad \text{if } k < 0, \quad (4.9)$$

and

$$(|x - a| + \gamma 2^{-n})^{-4} \leq C\gamma^{-1} 2^n \phi_n(x, t) \quad \text{if} \quad b - \gamma^2 2^{-2n} \leq t \leq b. \tag{4.10}$$

*Proof.* We fix  $n$ . Let  $h_n: R^3 \rightarrow R^+$  be defined by  $h(x) = \gamma 2^{-n} (|x| + \gamma 2^{-n})^{-4}$ . We define  $F: R^3 \times \{t: t < 0\} \rightarrow R^+$  by

$$F(x, t) = (2\sqrt{\pi})^{-3} (-t)^{-3/2} \exp(|x|^2/(4t)).$$

The function  $F$  is the fundamental solution to the heat equation with time reversed. We define  $\psi_n: R^3 \times (-\infty, b + \gamma^2 2^{-2n}) \rightarrow R^+$  by

$$\psi_n(x, t) = (F * h_n)(x - a, t - (b + \gamma^2 2^{-2n})).$$

We have  $D_t \psi_n + \Delta \psi_n = 0$ , and the properties (4.5), (4.6), (4.7), (4.10) are satisfied if  $\phi_n$  is replaced by  $\psi_n$ . Let  $g: R \rightarrow [0, 1]$  be a  $C^\infty$  function such that  $g(t) = 0$  if  $t \leq b - \gamma^2$ ,  $g(t) = 1$  if  $b - \gamma^2/2 \leq t \leq b$ ,  $g(t) = 0$  if  $b + \gamma^2 2^{-2n-1} \leq t$ , and  $|D_t g(t)| \leq C\gamma^{-2}$  if  $b - \gamma^2 \leq t \leq b - \gamma^2/2$ . Define  $\phi_n \in D^\infty(R^3 \times R^+, R)$  by  $\phi_n(x, t) = g(t)\psi_n(x, t)$  if  $t < b + \gamma^2 2^{-2n}$ , and  $\phi_n(x, t) = 0$  if  $t > b + \gamma^2 2^{-2n-1}$ . Then (4.4) is clear, and (4.5), (4.6), (4.7), (4.10) follow from the corresponding estimates on  $\psi_n$ . We have  $(D_t \phi_n + \Delta \phi_n)(x, t) = D_t g(t)\psi_n(x, t)$  if  $t \leq b$ . In particular, we have  $(D_t \phi_n + \Delta \phi_n)(x, t) = 0$  if  $b - \gamma^2/2 \leq t \leq b$ . Now (4.8) and (4.9) follow from the estimates on  $\psi_n$ .

**4.3. Definition.** For  $k = 1, 2, 3, \dots$  we fix  $C^\infty$  functions  $r_k$  on a neighborhood of  $R^3 \times [b - \gamma^2, b]$  such that  $r_k(x, t) \in [0, 1]$ ,  $r_k(x, t) = 1$  if  $(x, t) \in R^3 \times (b - \gamma^2, b)$  and  $(x, t) \notin E_{k+1}$ ,  $r_k(x, t) = 0$  if  $(x, t) \in E_{k+2}$ , and  $\sup(|Dr_k|, E_{k+1} - E_{k+2}) \leq C\gamma^{-1} 2^k$ . For  $n = 1, 2, 3, \dots$  and  $\delta > 0$  the inequalities (4.11), (4.12), (4.13) will be known as Property  $P(n, \delta)$ :

$$\int_{b - \gamma^2}^b \int_{R^3} |u(x, t)|^2 (|x - a| + \gamma)^{-4} dx dt \leq \delta \gamma^{-1} \tag{4.11}$$

$$\int_{b - \gamma^2}^b \left| \int_{R^3} u_t(x, t) (2^{-1}|u(x, t)|^2 + p(x, t)) D_t \phi_n(x, t) dx \right| dt \leq \delta \gamma^{-2} \tag{4.12}$$

$$\begin{aligned} & \int_{b - \gamma^2}^b \left| \int_{R^3} u_t(x, t) (2^{-1}|u(x, t)|^2 + p(x, t)) D_t (r_n \phi_q)(x, t) dx \right| dt \\ & \leq \delta \gamma^{-2} \quad \text{if} \quad q > n. \end{aligned} \tag{4.13}$$

**4.4. Lemma.** *There exists an absolute constant  $M$  with the following property: If  $P(n, \delta)$  holds (see Definition 4.3) then  $P(n + 1, \delta + M\delta^{3/2} 2^{-n})$  holds.*

*Proof.* Suppose that  $P(n, \delta)$  holds for some  $n$  and  $\delta$ . Let  $s \in A$  (see Definition 3.3) with  $b - \gamma^2 2^{-2n} \leq s \leq b$  and set  $B_s = R^3 \times [0, s]$ . Using Lemma 3.7, Lemma 4.2, and  $P(n, \delta)$  [Parts (4.12) and (4.11)] we obtain

$$\begin{aligned} & 2^{-1} \int_{R^3} |u(x, s)|^2 \phi_n(x, s) dx + \int_{B_s} |Du|^2 \phi_n \\ & \leq 2^{-1} \int_{B_s} |u|^2 (D_t \phi_n + \Delta \phi_n) + \int_{B_s} u_t (2^{-1}|u|^2 + p) D_t \phi_n \\ & \leq C\gamma^{-1} \int_{b - \gamma^2}^s \int_{R^3} |u(x, t)|^2 (|x - a| + \gamma)^{-4} dx dt + \delta \gamma^{-2} \\ & \leq C\delta \gamma^{-2}. \end{aligned} \tag{4.14}$$

Now (4.10) and (4.14) yield

$$\begin{aligned}
 & \int_{\mathbb{R}^3} |u(x, s)|^2 (|x - a| + \gamma 2^{-n})^{-4} dx \\
 & \leq C \gamma^{-1} 2^n \int_{\mathbb{R}^3} |u(x, s)|^2 \phi_n(x, s) dx \\
 & \leq C \delta \gamma^{-3} 2^n \quad \text{if } s \in A \quad \text{and } b - \gamma^2 2^{-2n} \leq s \leq b.
 \end{aligned} \tag{4.15}$$

Using (4.3), (4.10), and (4.14) (with  $s = b$ ) we obtain

$$\int_{E_n} |Du|^2 \leq C \gamma^3 2^{-3n} \int_{B_b} |Du|^2 \phi_n \leq C \delta \gamma 2^{-3n}. \tag{4.16}$$

For  $q = n + 1, n + 2, n + 3, \dots$  we define  $h_q : \mathbb{R}^3 \times (b - \gamma^2, b) \rightarrow \mathbb{R}$  by (see Definition 4.3)

$$h_{n+1} = (1 - r_n) \phi_{n+1}, h_q = (r_{n+1} - r_n) \phi_q \quad \text{if } q > n + 1. \tag{4.17}$$

From Definition 4.3 and Lemma 4.2 we obtain

$$\begin{aligned}
 & h_q(x, t) = 0 \quad \text{if} \\
 & (x, t) \notin E_{n+1}, \|h_q\|_\infty \leq C \gamma^{-3} 2^{3n}, \|Dh_q\|_\infty \leq C \gamma^{-4} 2^{4n}.
 \end{aligned} \tag{4.18}$$

Let  $s \in A$  such that  $b - \gamma^2 2^{-2n-2} < s < b$ . Using (4.18), the Schwarz inequality, (4.15), Lemma 3.8, and (4.15) again, we obtain

$$\begin{aligned}
 & \left| \int_{\mathbb{R}^3} u_i(x, s) (2^{-1} |u(x, s)|^2) D_i h_q(x, s) dx \right| \\
 & \leq C \gamma^{-4} 2^{4n} \left( \int_{B(a, \gamma 2^{-n-1})} |u(x, s)|^2 dx \right)^{1/2} \left( \int_{B(a, \gamma 2^{-n-1})} |u(x, s)|^4 dx \right)^{1/2} \\
 & \leq C \delta^{1/2} \gamma^{-7/2} 2^{5n/2} \left( \int_{B(a, \gamma 2^{-n-1})} |u(x, s)|^4 dx \right)^{1/2} \\
 & \leq C \delta^{1/2} \gamma^{-3} 2^{2n} \left( \int_{B(a, \gamma 2^{-n})} |Du(x, s)|^2 dx \right) \\
 & \quad + C \delta^{1/2} \gamma^{-5} 2^{4n} \left( \int_{B(a, \gamma 2^{-n})} |u(x, s)|^2 dx \right) \\
 & \leq C \delta^{1/2} \gamma^{-3} 2^{2n} \left( \int_{B(a, \gamma 2^{-n})} |Du(x, s)|^2 dx \right) + C \delta^{3/2} \gamma^{-4} 2^n.
 \end{aligned} \tag{4.19}$$

Now we integrate (4.19) over  $s$  (recall Definition 3.3) and apply (4.16) and (4.3) to obtain

$$\begin{aligned}
 & \int_{b - \gamma^2 2^{-2n-2}}^b \left| \int_{\mathbb{R}^3} u_i(x, s) (2^{-1} |u(x, s)|^2) D_i h_q(x, s) dx \right| ds \\
 & \leq C \delta^{3/2} \gamma^{-2} 2^{-n}.
 \end{aligned} \tag{4.20}$$



We choose  $\alpha$  and  $\beta$  as in Lemma 3.4 such that  $0 \leq \alpha(x) \leq 1$ ,  $\alpha(x) = 0$  for  $|x| \geq \gamma 2^{-n-1}$ ,  $\beta(x) = 0$  for  $|x| \leq \gamma 2^{-n-2}$ ,  $|D\beta(x)| \leq C\gamma^{-1}2^n$ ,  $|D_{ij}\beta(x)| \leq C\gamma^{-2}2^{2n}$ , and  $|D_{ijk}\beta(x)| \leq C\gamma^{-3}2^{3n}$ . Then we have (see Lemma 3.4)

$$\|\alpha'\|_2 \leq C\gamma^{1/2}2^{-n/2}, |D_{ijk}\beta'(x)| \leq C(|x| + \gamma 2^{-n})^{-4} \quad \text{if } x \in R^3. \quad (4.21)$$

Let  $s \in A$  such that  $b - \gamma^2 2^{-2n-2} < s < b$ . We set  $g_s(x) = (D_i u_j D_j u_i)(x, s)$  for  $|x - a| < \gamma 2^{-n}$ , and  $g_s(x) = 0$  for  $|x - a| \geq \gamma 2^{-n}$ . Then the property  $\text{spt}(\alpha') \subset B(0, \gamma 2^{-n-1})$ , [10, Theorem 1.3, p. 3], and (4.21) yield

$$\begin{aligned} & \left( \int_{B(a, \gamma 2^{-n-1})} |((D_i u_j D_j u_i) * \alpha')(x, s)|^2 dx \right)^{1/2} \\ &= \left( \int_{B(a, \gamma 2^{-n-1})} |(g_s * \alpha')(x)|^2 dx \right)^{1/2} \\ &\leq \|g_s * \alpha'\|_2 \leq \|g_s\|_1 \|\alpha'\|_2 \leq C \left( \int_{B(a, \gamma 2^{-n})} |Du(x, s)|^2 dx \right) (\gamma^{1/2} 2^{-n/2}). \end{aligned} \quad (4.22)$$

Using (4.18), the Schwarz inequality, (4.15), and (4.22) we obtain

$$\begin{aligned} & \left| \int_{R^3} u_k(x, s) ((D_i u_j D_j u_i) * \alpha')(x, s) D_k h_q(x, s) dx \right| \\ &\leq C\gamma^{-4} 2^{4n} \left( \int_{B(a, \gamma 2^{-n-1})} |u(x, s)| |((D_i u_j D_j u_i) * \alpha')(x, s)| dx \right) \\ &\leq C\delta^{1/2} \gamma^{-3} 2^{2n} \int_{B(a, \gamma 2^{-n})} |Du(x, s)|^2 dx. \end{aligned} \quad (4.23)$$

If  $|x - a| \leq \gamma 2^{-n-1}$  then (4.21) and (4.15) yield

$$\begin{aligned} & |((u_j u_i) * D_{ijk}\beta')(x, s)| \leq C \int_{R^3} |u(y, s)|^2 (|x - y| + \gamma 2^{-n})^{-4} dy \\ &\leq C \int_{R^3} |u(y, s)|^2 (|y - a| + \gamma 2^{-n})^{-4} dy \leq C\delta \gamma^{-3} 2^n. \end{aligned} \quad (4.24)$$

Hence Definition 3.3, integration by parts, (4.24), (4.18), the Schwarz inequality, and (4.15) yield

$$\begin{aligned} & \left| \int_{R^3} u_k(x, s) ((u_j u_i) * D_{ij}\beta')(x, s) D_k h_q(x, s) dx \right| \\ &= \left| \int_{R^3} u_k(x, s) ((u_j u_i) * D_{ijk}\beta')(x, s) h_q(x, s) dx \right| \\ &\leq C(\delta \gamma^{-3} 2^n) (\gamma^{-3} 2^{3n}) \int_{B(a, \gamma 2^{-n-1})} |u(x, s)| dx \\ &\leq C\delta \gamma^{-6} 2^{4n} \left( \int_{B(a, \gamma 2^{-n-1})} |u(x, s)|^2 dx \right)^{1/2} (\text{measure}(B(a, \gamma 2^{-n-1})))^{1/2} \\ &\leq C\delta^{3/2} \gamma^{-4} 2^n. \end{aligned} \quad (4.25)$$

Now (4.23), (4.25), and Lemma 3.4 yield

$$\begin{aligned} & \left| \int_{R^3} u_k(x, s) p(x, s) D_k h_q(x, s) dx \right| \\ &\leq C\delta^{1/2} \gamma^{-3} 2^{2n} \left( \int_{B(a, \gamma 2^{-n})} |Du(x, s)|^2 dx \right) + C\delta^{3/2} \gamma^{-4} 2^n. \end{aligned} \quad (4.26)$$

Integration of (4.26) with respect to  $s$ , Definition 3.3, (4.16), (4.3), and (4.20) yield

$$\begin{aligned} & \int_{b-\gamma^2}^b \int_{R^3} u_i(x, s) (2^{-1}|u(x, s)|^2 + p(x, s)) D_i h_q(x, s) dx \Big| ds \\ & \leq M \delta^{3/2} \gamma^{-2} 2^{-n}, \end{aligned} \quad (4.27)$$

where  $M$  is an absolute constant.

Setting  $q = n + 1$  in (4.13) and (4.27), and using (4.17), (4.18), and (4.3), we obtain

$$\begin{aligned} & \int_{b-\gamma^2}^b \int_{R^3} u_i(x, s) (2^{-1}|u(x, s)|^2 + p(x, s)) D_i \phi_{n+1}(x, s) dx \Big| ds \\ & \leq (\delta + M \delta^{3/2} 2^{-n}) \gamma^{-2}. \end{aligned} \quad (4.28)$$

Using (4.13) and (4.27) with  $q > n + 1$ , and using (4.17), (4.18), and (4.3) once again, we obtain

$$\begin{aligned} & \int_{b-\gamma^2}^b \int_{R^3} u_i(x, s) (2^{-1}|u(x, s)|^2 + p(x, s)) D_i (r_{n+1} \phi_q)(x, s) dx \Big| ds \\ & \leq (\delta + M \delta^{3/2} 2^{-n}) \gamma^{-2} \quad \text{if } q > n + 1. \end{aligned} \quad (4.29)$$

Now (4.28), (4.29), and property  $P(n, \delta)$  imply that  $P(n + 1, \delta + M \delta^{3/2} 2^{-n})$  holds. Lemma 4.4 has been proved.

Now we can prove Theorem 4.1. Choose an absolute constant  $\delta_0$  such that  $M(2\delta_0)^{3/2} \leq \delta_0$  (see Lemma 4.4). We have

$$\begin{aligned} & (\delta_0(2 - 2^{-n+1})) + M(\delta_0(2 - 2^{-n+1}))^{3/2} 2^{-n} \\ & < 2\delta_0 - 2^{-n+1} \delta_0 + M(2\delta_0)^{3/2} 2^{-n} \leq \delta_0(2 - 2^{-(n+1)+1}). \end{aligned}$$

Hence Lemma 4.4 and the definition of  $P(n, \delta)$  yield that  $P(n, \delta_0(2 - 2^{-n+1}))$  implies  $P(n + 1, \delta_0(2 - 2^{-(n+1)+1}))$ . Hence induction yields that  $P(1, \delta_0)$  implies  $P(n, \delta_0(2 - 2^{-n+1}))$  for all  $n$ . Now the definition of  $P(n, \delta)$  yields

$$P(1, \delta_0) \text{ implies } P(n, 2\delta_0) \text{ for all } n. \quad (4.30)$$

There is an absolute constant  $\eta$  satisfying

$$\eta \int_{b-\gamma^2}^b \int_{R^3} (|x - a| + \gamma)^{-4} dx dt \leq \gamma.$$

Young's inequality (see [4, p. 11]) yields

$$|u|^2 \leq (2/3)((\delta_0 \eta \gamma^{-2})^{-1/3} |u|^2)^{3/2} + (1/3)((\delta_0 \eta \gamma^{-2})^{1/3})^3.$$

Hence we have

$$\begin{aligned} & \int_{b-\gamma^2}^b \int_{R^3} |u(x, t)|^2 (|x - a| + \gamma)^{-4} dx dt \\ & \leq C \gamma \left( \int_{b-\gamma^2}^b \int_{R^3} |u(x, t)|^3 (|x - a| + \gamma)^{-4} dx dt \right) + (1/3) \delta_0 \gamma^{-1}. \end{aligned} \quad (4.31)$$

Now the estimates  $|D_i \phi_1(x, t)| \leq C(|x - a| + \gamma)^{-4}$  and  $|D_i(r_1 \phi_q)(x, t)| \leq C(|x - a| + \gamma)^{-4}$  for  $b - \gamma^2 \leq t \leq b$  and  $q > 1$  (see Lemma 4.2 and Definition 4.3), and (4.31) yield the existence of an absolute constant  $\varepsilon$  such that (4.1) implies  $P(1, \delta_0)$ . Hence (4.30) yields

$$\text{Inequality (4.1) implies } P(n, 2\delta_0) \text{ for all } n. \quad (4.32)$$

The assumption  $b \in A$ , (4.32), and the argument that yielded (4.15) can be used to show that (4.1) implies

$$\int_{R^3} |u(x, b)|^2 (|x - a| + \gamma 2^{-n})^{-4} dx \leq C(2\delta_0)\gamma^{-3} 2^n .$$

Hence we have that (4.1) implies

$$\int_{B(a, \gamma 2^{-n})} |u(x, b)|^2 dx \leq C\delta_0 \gamma 2^{-3n} . \tag{4.33}$$

For  $0 < \tau \leq 1/2$  we choose  $n$  such that  $2^{-n} \geq \tau > 2^{-n-1}$ . Then (4.33) yields

$$\begin{aligned} \int_{B(a, \tau\gamma)} |u(x, b)|^2 dx &\leq \int_{B(a, \gamma 2^{-n})} |u(x, b)|^2 dx \\ &\leq C\delta_0 \gamma 2^{-3n} \leq C\delta_0 \gamma (2\tau)^3 = K\tau^3 \gamma \end{aligned}$$

where  $K$  is an absolute constant. Theorem 4.1 has been proved.

### 5. The Connection with Hausdorff Measure

Throughout this section we assume that Hypothesis I holds.

*5.1. Definition.* We define  $V: R^3 \times R^+ \rightarrow R$  by  $V = |u|(2^{-1}|u|^2 + |p|)$ . For every integer  $n$  we define  $Q_n: R^3 \times R \rightarrow R$  by  $Q_n(x, t) = (|x| + 2^{-n})^{-4}$  if  $-2^{-2n} \leq t \leq 2^{-2n}$ , and  $Q_n(x, t) = 0$  otherwise. For  $t \geq 2^{-2n}$  we set

$$V_n(x, t) = \int_0^\infty \int_{R^3} V(y, s) Q_n(x - y, t - s) dy ds .$$

We define  $B(n, p_1, p_2, p_3, p_4)$  to be the set of all  $(x, t) \in R^3 \times R$  satisfying  $p_i 2^{-n} \leq x_i \leq (p_i + 1) 2^{-n}$  for  $i \in \{1, 2, 3\}$ , and  $p_4 2^{-2n} \leq t \leq (p_4 + 1) 2^{-2n}$ . We set  $B(n) = \{B(n, p_1, p_2, p_3, p_4) : p_i \text{ is an integer for all } i, \text{ and } p_4 \geq 1\}$ .

From Lemma 3.2 and Lemma 3.6 we obtain

$$\int_0^T \int_{R^3} V(x, t) dx dt \leq CL^{3/2} T^{1/4} \quad \text{if } 0 < T < \infty . \tag{5.1}$$

If  $2^{-2n} \leq a < b$ , [10, Theorem 1.3, p. 3] yields

$$\int_a^b \int_{R^3} V_n(x, t) dx dt \leq C 2^{-n} \int_{a-2^{-2n}}^{b+2^{-2n}} \int_{R^3} V(x, t) dx dt . \tag{5.2}$$

**5.2. Lemma.** *There exists an absolute constant  $\theta$  such that the conditions  $B \in B(n)$  and*

$\int_B V_n \leq \theta 2^{-3n}$  *imply that  $|u|$  is essentially bounded on a neighborhood of  $B$ .*

*Proof.* Let  $B = B(n, p_1, p_2, p_3, p_4)$  and  $\gamma = 2^{-n-2}$ . We set

$$\begin{aligned} U &= \{(x, t) \in R^3 \times R : (p_i - 1) 2^{-n} < x_i < (p_i + 2) 2^{-n} \text{ for } i \in \{1, 2, 3\}, \\ &\quad \text{and } p_4 2^{-2n} - 2^{-2n-4} < t < (p_4 + 1) 2^{-2n} + 2^{-2n-4}\} , \end{aligned}$$

$$\begin{aligned} D &= \{(x, t) \in R^3 \times R : p_i 2^{-n} \leq x_i \leq (p_i + 1) 2^{-n} \text{ for } i \in \{1, 2, 3\}, \\ &\quad \text{and } p_4 2^{-2n} + 2^{-2n-2} \leq t \leq (p_4 + 1) 2^{-2n} - 2^{-2n-2}\} . \end{aligned}$$

Now let  $(a, b) \in U$ . For every  $(y, s) \in D$  we have (see Definition 5.1)

$$\int_{b-\gamma^2}^b \int_{\mathbb{R}^3} V(x, t) (|x-a|+\gamma)^{-4} dx dt \leq C V_n(y, s).$$

Averaging over  $D$  and using the fact  $D \subset B$ , we obtain

$$\begin{aligned} & \int_{b-\gamma^2}^b \int_{\mathbb{R}^3} V(x, t) (|x-a|+\gamma)^{-4} dx dt \\ & \leq C(\text{measure}(D))^{-1} \left( \int_D V_n \right) \leq C\gamma^{-5} \int_B V_n. \end{aligned}$$

Since  $2^{-3n} = 2^6 \gamma^3$ , there exists an absolute constant  $\theta$  such that the property  $\int_B V_n \leq \theta 2^{-3n}$  implies that (4.1) holds for  $(a, b) \in U$ . Then we can use Theorem 4.1, Definition 3.3, and [9, Corollary 1, p. 5] to conclude that  $|u(a, b)|^2 \leq K(4\pi/3)^{-1} \gamma^{-2}$  holds for almost every  $(a, b) \in U$ .

**5.3. Definition.** The 2 dimensional Hausdorff measure of a set  $S \subset \mathbb{R}^3 \times \mathbb{R}$  is denoted by  $\mathcal{H}^2(S)$ . For the definition of Hausdorff measure, see [2, p. 171] (where  $X = \mathbb{R}^3 \times \mathbb{R}$  and the metric on  $X$  is the usual metric on  $\mathbb{R}^4$ ).

**5.4. Lemma.** For each integer  $k$  there exists a compact set  $S_k$  contained in  $\mathbb{R}^3 \times [2^{-k}, 2^{-k+1}]$  such that

$$\mathcal{H}^2(S_k) \leq C \int_{2^{-k-1}}^{2^{-k+2}} \int_{\mathbb{R}^3} V(x, t) dx dt \tag{5.3}$$

and for every  $(x, t) \in (\mathbb{R}^3 \times [2^{-k}, 2^{-k+1}]) - S_k$  there exists a neighborhood  $U$  of  $(x, t)$  such that  $|u|$  is essentially bounded on  $U$ .

*Proof.* Let  $k$  be given. For each integer  $n$  satisfying  $n \geq k+1$  and  $n \geq 0$  we set (see Lemma 5.2)  $D(n) = \left\{ B \in \mathcal{B}(n) : B \subset \mathbb{R}^3 \times [2^{-k}, 2^{-k+1}] \text{ and } \int_B V_n > \theta 2^{-3n} \right\}$ . We then set

$$S_k = \cap \{ (\cup \{ B : B \in D(n) \}) : n \geq k+1 \text{ and } n \geq 0 \}.$$

For each  $n$ , (5.2) yields

$$\sum_{B \in D(n)} \int_B V_n \leq C 2^{-n} \int_{2^{-k-1}}^{2^{-k+2}} \int_{\mathbb{R}^3} V(x, t) dx dt.$$

Hence the number of elements in  $D(n)$  is at most

$$C 2^{2n} \int_{2^{-k-1}}^{2^{-k+2}} \int_{\mathbb{R}^3} V(x, t) dx dt.$$

Hence (5.1) implies that  $S_k$  is compact, and we also have (using  $n \geq 0$ )

$$\begin{aligned} \sum_{B \in D(n)} (\text{diameter}(B))^2 & \leq \sum_{B \in D(n)} C 2^{-2n} \\ & \leq C \int_{2^{-k-1}}^{2^{-k+2}} \int_{\mathbb{R}^3} V(x, t) dx dt. \end{aligned}$$

Since the diameter of the sets in  $D(n)$  can be made arbitrarily small by taking  $n$  sufficiently large, and  $S_k$  is contained in  $\cup\{B : B \in D(n)\}$  for sufficiently large  $n$ , [2, p. 171] yields (5.3).

Now take  $(x, t) \in (R^3 \times [2^{-k}, 2^{-k+1}]) - S_k$ . There exists  $n \geq \max(k+1, 0)$  such that  $(x, t) \notin B$  for every  $B \in D(n)$ . However, there exists  $B \in B(n)$  such that  $B \subset R^3 \times [2^{-k}, 2^{-k+1}]$  and  $(x, t) \in B$ . Hence Lemma 5.2 implies that  $|u|$  is essentially bounded on a neighborhood of  $B$ , and hence on a neighborhood of  $(x, t)$ .

Now we can prove Theorem 2.1. For any integer  $n$ , (5.2) and (5.1) yield

$$\int_{2^{-2n}}^{2^{-2n+2}} \int_{R^3} V_n(x, t) dx dt \leq C 2^{-n} \int_0^{2^{-2n+3}} \int_{R^3} V(x, t) dx dt \leq WL^{3/2} 2^{-3n/2} \tag{5.4}$$

where  $W$  is an absolute constant. Let  $m$  be the integer that satisfies  $WL^{3/2} \leq \theta 2^{-3m/2} < 2^{3/2} WL^{3/2}$  (see Lemma 5.2). If  $n, p_1, p_2, p_3$  are integers such that  $n \leq m$  then, setting  $B_i = B(n, p_1, p_2, p_3, i)$  for  $i \in \{1, 2, 3\}$ , we obtain that (5.4) yields

$$\int_{B_i} V_n \leq WL^{3/2} 2^{-3n/2} \leq \theta 2^{-3m/2} 2^{-3n/2} \leq \theta 2^{-3n} \text{ for } i = 1, 2, 3 .$$

Hence Lemma 5.2 yields that  $|u|$  is essentially bounded on  $B_1, B_2,$  and  $B_3$ . By varying  $n$  and  $p_j, j=1, 2, 3$ , we obtain that  $|u|$  is locally essentially bounded on the set  $\{(x, t) : x \in R^3 \text{ and } t \geq 2^{-2m}\}$ . Actually, the proof of Lemma 5.2 shows that  $|u|$  is essentially bounded on that set. We define  $S = \cup\{S_k : k \geq 2m+1\}$ . The above and Lemma 5.4 yield that  $|u|$  is locally essentially bounded outside of  $S$ . Finally, the countable subadditivity of  $\mathcal{H}^2$ , (5.3), (5.1), and the definition of  $m$  yield

$$\mathcal{H}^2(S) \leq \sum_{k \geq 2m+1} \mathcal{H}^2(S_k) \leq 3C \int_0^{2^{-2m+1}} \int_{R^3} V(x, t) dx dt \leq CL^2 .$$

Theorem 2.1 has been proved.

We can prove Theorem 2.2 as follows: First, use Hypothesis II to imitate the proof of [7, Lemma 1.1] and derive identity (1.8) of [7] for almost every  $x, t_1,$  and  $t_2$ . Then use Theorem 2.1 to adapt the proof in the last paragraph of [7, Section 2] to our case.

### 6. Outline of Proof of Theorem 2.3

Let  $v$  be given as in Theorem 2.3. From [5] we obtain that there exist  $0 < L < \infty$  and  $(u, n) \in C^\infty(R^3 \times R^+, R^3)$  for  $n=1, 2, 3, \dots$  such that (see Definition 3.3)

$$\operatorname{div}(u, n) = 0 , \tag{6.1}$$

$$\int_{R^3} |(u, n)(x, t)|^2 dx \leq L \text{ for all } t \in R^+ , \tag{6.2}$$

$$\int_0^\infty \int_{R^3} |D(u, n)(x, t)|^2 dx dt \leq L , \tag{6.3}$$

$$\begin{aligned} & - \int_{R^3} v_i(x) \phi_i(x, 0) dx - \int_{R^3 \times R^+} (u, n)_i (D_t \phi_i + \Delta \phi_i) \\ & = \int_{R^3 \times R^+} ((u, n)_j * f_n)(u, n)_i D_j \phi_i \end{aligned} \tag{6.4}$$

whenever  $\phi \in C_0^\infty(R^3 \times R, R^3)$  satisfies  $\operatorname{div}(\phi) = 0$ . We also obtain from [5] that there exists an increasing sequence  $n_1, n_2, n_3, \dots$  of positive integers and a Lebesgue

measurable function  $u: R^3 \times R^+ \rightarrow R^3$  such that (2.1), (2.2), and (2.3) are satisfied, and we have

$$\lim_k \int_{R^3} |u(x, t) - (u, n_k)(x, t)|^2 dx = 0 \quad (6.5)$$

for almost every  $t \in R^3$ , and

$$D(u, n_k) \text{ converges weakly in } L^2 \text{ to } Du. \quad (6.6)$$

If  $0 < T < \infty$  then the Lebesgue dominated convergence theorem, (2.2), (6.2), and (6.5) yield

$$\lim_k \int_0^T \left( \int_{R^3} |u(x, t) - (u, n_k)(x, t)|^2 dx \right)^3 dt = 0. \quad (6.7)$$

From Lemma 3.1, (6.2), (2.2), (6.3), and (2.3) we obtain

$$\begin{aligned} & \int_0^T \int_{R^3} |u(x, t) - (u, n_k)(x, t)|^3 dx dt \\ & \leq C\varepsilon^{-3} \int_0^T \left( \int_{R^3} |u(x, t) - (u, n_k)(x, t)|^2 dx \right)^3 dt + C\varepsilon L \end{aligned} \quad (6.8)$$

for every  $0 < \varepsilon < \infty$ . Combining (6.7) and (6.8) (with varying  $\varepsilon$ ) we obtain

$$\lim_k \int_0^T \int_{R^3} |u(x, t) - (u, n_k)(x, t)|^3 dx dt = 0. \quad (6.9)$$

Let  $\alpha, \alpha', \beta, \beta'$  be as in Lemma 3.4. Define  $(p, n): R^3 \times R^+ \rightarrow R$  and  $p: R^3 \times R^+ \rightarrow R$  by

$$\begin{aligned} (p, n)(x, t) = & -(D_i((u, n)_j * f_n) D_j(u, n)_i * \alpha')(x, t) \\ & - (((u, n)_j * f_n)(u, n)_i * D_{ij} \beta')(x, t) \end{aligned} \quad (6.10)$$

$$p(x, t) = -(D_i u_j D_j u_i * \alpha')(x, t) - (u_j u_i * D_{ij} \beta')(x, t)$$

for almost all  $(x, t)$ . The argument in Lemma 3.5, the Schwarz inequality, (2.2), (6.2), (2.3), (6.3), Young's inequality, and [10, Theorem 1.3, p. 3] yield

$$\begin{aligned} & \left( \int_{R^3} |(p, n_k)(x, t) - p(x, t)|^2 dx \right)^{1/2} \\ & \leq Cr^{1/2} \int_{R^3} |(D(u, n_k) * f_{n_k})(x, t)|^2 + |D(u, n_k)(x, t)|^2 + |Du(x, t)|^2 dx \\ & \quad + Cr^{-3/2} \int_{R^3} |((u, n_k) * f_{n_k})(x, t) - u(x, t)| |(u, n_k)(x, t)| dx \\ & \quad + Cr^{-3/2} \int_{R^3} |u(x, t)| |(u, n_k)(x, t) - u(x, t)| dx \\ & \leq Cr^{1/2} \int_{R^3} |D(u, n_k)(x, t)|^2 + |Du(x, t)|^2 dx \\ & \quad + Cr^{-3/2} L^{1/2} \left( \int_{R^3} |((u, n_k) * f_{n_k})(x, t) - u(x, t)|^2 dx \right)^{1/2} \\ & \quad + Cr^{-3/2} L^{1/2} \left( \int_{R^3} |(u, n_k)(x, t) - u(x, t)|^2 dx \right)^{1/2} \end{aligned} \quad (6.11)$$

for almost all  $t \in R^+$ . The Schwarz inequality, the argument in Lemma 3.5, (2.2), (2.3), (6.2), (6.3), (6.11), Young's inequality, and [10, Theorem 1.3, p. 3] yield

$$\begin{aligned}
& \int_0^T \int_{R^3} |(u, n_k)_i(x, t)(p, n_k)(x, t) - u_i(x, t)p(x, t)| dx dt \\
& \leq \int_0^T \int_{R^3} |(u, n_k)(x, t) - n(x, t)| |(p, n_k)(x, t)| dx dt \\
& \quad + \int_0^T \int_{R^3} |u(x, t)| |(p, n_k)(x, t) - p(x, t)| dx dt \\
& \leq \int_0^T \left( \int_{R^3} |(u, n_k)(x, t) - u(x, t)|^2 dx \right)^{1/2} \left( \int_{R^3} |(p, n_k)(x, t)|^2 dx \right)^{1/2} dt \\
& \quad + \int_0^T \left( \int_{R^3} |u(x, t)|^2 dx \right)^{1/2} \left( \int_{R^3} |(p, n_k)(x, t) - p(x, t)|^2 dx \right)^{1/2} dt \\
& \leq Cr^{1/2} \int_0^T \left( \int_{R^3} |(u, n_k)(x, t) - u(x, t)|^2 dx \right)^{1/2} \left( \int_{R^3} |D(u, n_k)(x, t)|^2 dx \right) dt \\
& \quad + Cr^{-3/2} \int_0^T \left( \int_{R^3} |(u, n_k)(x, t) - u(x, t)|^2 dx \right)^{1/2} \left( \int_{R^3} |(u, n_k)(x, t)|^2 dx \right) dt \\
& \quad + L^{1/2} \int_0^T \left( \int_{R^3} |(p, n_k)(x, t) - p(x, t)|^2 dx \right)^{1/2} dt \\
& \leq Cr^{1/2} L^{3/2} + Cr^{-3/2} L \int_0^T \left( \int_{R^3} |(u, n_k)(x, t) - u(x, t)|^2 dx \right)^{1/2} dt \\
& \quad + Cr^{1/2} L^{3/2} + Cr^{-3/2} L \int_0^T \left( \int_{R^3} |((u, n_k) * f_{n_k})(x, t) - u(x, t)|^2 dx \right)^{1/2} dt \\
& \quad + Cr^{-3/2} L \int_0^T \left( \int_{R^3} |(u, n_k)(x, t) - u(x, t)|^2 dx \right)^{1/2} dt \tag{6.12}
\end{aligned}$$

for  $0 < T < \infty$ . Now we make  $r$  small and use (6.12), (2.2), (2.3), (6.2), (6.3), (6.5), the fact

$$\lim_k \int_{R^3} |((u, n_k) * f_{n_k})(x, t) - u(x, t)|^2 dx = 0$$

for almost every  $t \in R^+$  [see (6.5)], and the Lebesgue dominated convergence theorem to conclude

$$\lim_k \int_0^T \int_{R^3} |(u, n_k)_i(x, t)(p, n_k)(x, t) - u_i(x, t)p(x, t)| dx dt = 0 \tag{6.13}$$

Let  $\phi$  satisfy (2.5). From (6.1), (6.2), (6.3), (6.4), (6.10), and the usual arguments we conclude

$$\begin{aligned}
& -2^{-1} (\int |(u, n)|^2 (D_i \phi + \Delta \phi)) + \int |D(u, n)|^2 \phi \\
& = 2^{-1} \int ((u, n)_i * f_n) |(u, n)|^2 D_i \phi + \int (u, n)_i (p, n) D_i \phi. \tag{6.14}
\end{aligned}$$

Now (2.2), (6.2), and (6.5) yield

$$\lim_k \int |(u, n_k)|^2 (D_i \phi + \Delta \phi) = \int |u|^2 (D_i \phi + \Delta \phi).$$

Properties (2.3), (6.3), and (6.6) yield (recall  $\phi \geq 0$ )

$$\liminf_k \int |D(u, n_k)|^2 \phi \geq \int |Du|^2 \phi.$$

From (6.9) and (6.13) we obtain

$$\lim_k 2^{-1} \int ((u, n_k)_i * f_{n_k})(u, n_k)|^2 D_i \phi = \int u_i (2^{-1} |u|^2) D_i \phi,$$

$$\lim_k \int (u, n_k)_i (p, n_k) D_i \phi = \int u_i p D_i \phi.$$

Hence (6.14) yields (2.6). Properties (2.7) and (2.9) are a more immediate consequence of (6.1), (6.4), (6.10), and the usual estimates.

## References

1. Almgren, F.J., Jr.: Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints. *Memoirs of the American Mathematical Society* 165. Providence, R. I.: American Mathematical Society 1976
2. Federer, H.: *Geometric measure theory*. Berlin-Heidelberg-New York: Springer 1969
3. Hewitt, E., Stromberg, K.: *Real and abstract analysis*. Berlin-Heidelberg-New York: Springer 1965
4. Ladyzhenskaya, O.A.: *The mathematical theory of viscous incompressible flow*. Revised English edition. New York: Gordon and Breach 1964
5. Leray, J.: *Acta Math.* **63**, 193—248 (1934)
6. Mandelbrot, B.: Intermittent turbulence and fractal dimension kurtosis and the spectral exponent  $5/3 + B$ . In: *Turbulence and Navier-Stokes equation; Lecture Notes in Mathematics* 565. Berlin-Heidelberg-New York: Springer 1976
7. Scheffer, V.: Partial regularity of solutions to the Navier-Stokes equations. *Pacific J. M.*, to appear
8. Scheffer, V.: Turbulence and Hausdorff dimension. In: *Turbulence and Navier-Stokes equation; Lecture Notes in Mathematics* 565. Berlin-Heidelberg-New York: Springer 1976
9. Stein, E.M.: *Singular integrals and differentiability properties of functions*. Princeton: Princeton Univ. Press 1970
10. Stein, E.M., Weiss, G.L.: *Introduction to fourier analysis on euclidean spaces*. Princeton: Princeton Univ. Press 1971

Communicated by J. Glimm

Received April 8, 1977