

Letter to the Editor

A Re-Examination of the Zeolite-Promoted, Clay-Mediated Peptide Synthesis

Joseph T. Warden*, John J. McCullough⁺, Richard M. Lemmon, and Melvin Calvin

Laboratory of Chemical Biodynamics, Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720

Received April 18, 1974

Summary. It has been found that, contrary to recent reports, peptides are not detectable products from alanine-ATP-Decalso-montmorillonite mixtures. The peptides appear only when pre-formed adenylates are reacted in the presence of montmorillonite.

Key words: Amino-acyl Adenylates - Prebiotic Peptide Formation.

The abiogenic synthesis of amino acids, nucleic acid bases, sugars, and nucleotides has been well documented in recent years (Calvin, 1969; Kenyon & Steinman, 1969; Orgel, 1973; Lemmon, 1970; Ponnampuruma, 1971). However, attempts to produce long chain polypeptides and polynucleotides under simulated aqueous, prebiotic conditions have generally been unsuccessful. Inspired by the suggestion of Bernal (1951) that aqueous prebiotic synthesis might be facilitated by adsorption of reactants on clays, Paecht-Horowitz et al. (1970) demonstrated that amino-acyl adenylates condense in the presence of the clay crystallite, montmorillonite, to yield polypeptides whose molecular weights were grouped around certain favored values. Although amino-acid adenylates undergo polymerization in basic aqueous media in the absence of montmorillonite, Paecht-Horowitz et al. (1967), Lewinsohn et al. (1967), it has also been shown that the clay-mediated polycondensation results in a substantially larger yield and an increase in polymer lengths (Paecht-Horowitz et al., 1970, 1973; Banda & Ponnampuruma, 1971).

*Present address: Department of Chemistry, Rensselaer Polytechnic Institute, Troy, N.Y.12181, USA.

+Present address: Department of Chemistry, McMaster University, Hamilton, Ontario, Canada.

A recent modification of this procedure by Paecht-Horowitz & Katchalsky (1973) utilizes a synthetic ion-exchange zeolite, Decalso F, as the "catalyst" for the formation of amino-acyl adenylate from ATP and amino acids. From their studies of a simple aqueous mixture of Decalso F, montmorillonite, ATP, and alanine (pH 7.0, 37°C), the Israeli workers reported the disappearance of free alanine over a period of ~70 hrs, with the concomitant formation of polymeric alanine. The polymer products were reported to be similar to those obtained, using pre-formed adenylate, from the montmorillonite-aided polymerization. However, as is reported below, we have been unable to substantiate the Paecht-Horowitz and Katchalsky report concerning the synthesis of either amino-acyl adenylates or peptides utilizing a zeolite catalyst. It is important for researchers in this area to realize that the peptide formation does not occur if free amino acids (in contrast to amino-acyl adenylates) are used in the reaction mixture.

Polymerization trials were conducted according to the prescription given recently by Paecht-Horowitz & Katchalsky (1973). Briefly, the reaction mixture typically contained 10^{-4} M DL-alanine (including 10 μ Ci DL-alanine-2- 14 C; autoradiography of this labeled compound revealed no detectable radioimpurity), 5×10^{-2} M ATP, 500 mg/l montmorillonite, and 100 g/l Decalso. The polymerization mixture was stirred for 70-100 hrs at 37°C and the pH was maintained at 7.2 by a pH stat (Radiometer Co., Copenhagen). When alanyl adenylate (labeled with 10 μ Ci alanine-2- 14 C) was utilized as the precursor for polymerization, the procedure of Paecht-Horowitz et al. (1970) was adopted. Synthesis of DL-alanyl-2- 14 C adenylate followed the method of Meister & Scott (1961).

After completion of the polycondensation trial, the clay suspension was subjected to a 2 hrs centrifugation (25,000 x g); the clay pellet was washed with distilled water, resuspended, and centrifuged, followed by ultrafiltration of the supernatant through a XM-50 filter (Amicon). Less than 0.5% of the alanine-containing material (radioactivity) was trapped by the filter and the clay. This procedure effectively removed small clay or zeolite particles. The resulting supernatant was lyophilized and resuspended into a minimal amount of distilled water (2-4 ml) and stored at -15°C until subsequent analysis for polymeric alanine.

The supernatants from the reaction mixtures were assayed for alanine and *low molecular weight* polypeptides by means of a Beckman amino acid analyzer. Generally, a sample of each reaction mixture was hydrolyzed (6 N HCl, 100°C, 6 hrs) and analyzed for free alanine (taken as the initial concentration).

Subsequently, the supernatant from the polymerization trial was analyzed for alanine and the value obtained compared with the initial concentration. Additionally, short-chain polymers of alanine (n approx. 2-5) could be detected and quantitatively determined by the amino acid analyzer. The supernatants, before and after the polymerization trials, were routinely assayed for radioactivity by liquid scintillation counting. This procedure assured us that all amino acid was accounted for after the polymerization attempts.

Additional characterization of the polymerization samples was obtained by thin-layer chromatography on Macherey-Nagel, Polygram CEL 300 UV cellulose-coated plastic strips. Buffer systems employed were either phenol-water (5:1 w/w) or n-butanol-acetic acid water (5:2:3 by vol). The chromatograms were sprayed with ninhydrin solution and were additionally analyzed for radioactive components by X-ray film autoradiography.

Aliquots (1 ml) of the supernatants from the reaction mixtures were layered on a Sephadex G-15 column (2.5 x 90 cm) and eluted with 0.05 M HCl. The eluted fractions (5 ml) were monitored for alanine and alanine peptides either by liquid scintillation counting or by ultraviolet absorption at 216 nm.

DL-Alanine (A grade) and adenosine-5'-triphosphate, disodium salt (A grade) were products of Calbiochem. DL-Alanine-2-¹⁴C was obtained from Mallinckrodt, Labeled Compounds Division. Decalso (Permutit Company) was repetitively ground and passed through a sieve of 100 μ . (We have been unable to find any supplier of the Decalso "F" that was used by the Israeli group.) Final particle diameters were on the average less than 20 μ . Montmorillonite # 25 (John C. Lane Tract, Upton, Wyoming) was procured from Wards Natural Science Establishment and prepared according to the method of Posner & Quirk (1964). In addition, we differentially centrifuged the montmorillonite and our polymerization was conducted with samples having average platelet diameters < 2000 \AA .

Although the report by Paecht-Horowitz & Katchalsky (1973) indicated essentially complete alanine polymerization, we have been unable after nine trials to demonstrate any detectable polymerization when the Decalso was used as the template for adenylate synthesis. This assertion is supported by the analytical determinations of the free alanine, both before and after polymerization trials, that are presented in Table 1. We estimate that our analytical precision for these determinations was within 5%, and that 1% of

Table 1

Trial	Ala (initial) (μM)	Ala (final) (μM)	Decalso (g/l)	Montmorillonite (mg/l)
1	145	146	100	500
2	364	370	100	500
3	201	199	300	500
4	166	167	100	1000

the alanine converted into low molecular weight polymer should have been observable. Furthermore, TLC analysis of the supernatant revealed only alanine and AMP; no polymeric species was observed. Increasing the concentration of Decalso (fivefold) or montmorillonite two or threefold did not lead to detectable alanine polymerization. Likewise, lengthening the incubation time for the reaction mixture also had no effect on the final free alanine content; neither did substituting sodium tripolyphosphate for ATP.

Assuming that the polymerization might be thwarted because the intermediate adenylate was not formed, we attempted to monitor adenylate formation by using the hydroxamic acid- FeCl_3 color test described by Hestrin (1949). No evidence for alanyl adenylate was found, thus indicating that our Decalso did not promote adenylate formation.

To confirm our hypothesis that the ineffectiveness of Decalso in promoting adenylate production was the cause for failure in our polymerization experiments, we repeated the polymerization trials using alanine-2- ^{14}C adenylate and montmorillonite. Approximately 30% of the initial alanine (as determined by liquid scintillation counting and amino acid analysis) was converted into polymer in the presence of montmorillonite. Thin-layer chromatograms yielded a characteristic streaking typical for polymers with $n > 2$. In the absence of the clay only $\sim 10\%$ polymer was obtained. The elution pattern from Sephadex G-15 of the montmorillonite-catalyzed reaction products is illustrated in Fig. 1. These results support the assertion by Paecht-Horowitz et al. (1970) that clay-mediated polymerization provides both a significant increase in yield and degree of polymerization.

It is hard to understand the ineffectiveness of Decalso as a template for production of the amino-acyl adenylate in the light of the earlier reports by Paecht-Horowitz & Katchalsky. Despite our attempts to vary the treat-




Fig.1. Elution pattern from Sephadex G-15 of a montmorillonite mediated polymerization, utilizing pre-formed alanyl adenylate

ment of the zeolite, in hopes of finding the key to activating the catalytic activity, we have not been able to demonstrate Decalso-mediated adenylate synthesis. It may be that the Decalso utilized by the Israeli group differs chemically or physically from that available for our investigations. This suggestion is supported by a recent disclosure that the Polymer group in Israel has also experienced difficulty with the Decalso-catalyzed production of amino-acyl adenylates (personal communication from Dr. Paecht-Horowitz); this communication also suggested that a lower cation-exchange capacity of the zeolite they have been using recently is responsible for the diminished effectiveness of this material. However, our experiments indicate that increasing Decalso concentration fivefold does not stimulate adenylate formation. Although the Decalso-mediated procedure would be of great utility for prebiotic research, the experiments presented here indicate that further investigation of the properties of the synthetic zeolite is warranted.

Acknowledgements. We are thankful to Ms. Joan Caroon for the preparation of the purified montmorillonite. This investigation was supported by the U.S. Atomic Energy Commission.

References

- Banda, P.W., Ponnampereuma, C. (1971). *Space Life Sci.* 3, 54
- Bernal, J.D. (1951). *The physical basis of life.* London: Routledge and Kegan Paul
- Calvin, M. (1969). *Chemical evolution.* Oxford: Clarendon Press

- Hestrin, S. (1949). *J. Biol. Chem.* 180, 249
- Kenyon, D.H., Steinman, G. (1969). *Biochemical predestination*. New York: McGraw-Hill
- Lemmon, R.M. (1970). *Chem. Rev.* 70, 95
- Lewinsohn, R., Paecht-Horowitz, M., Katchalsky, A. (1967). *Biochim. Biophys. Acta* 140, 24
- Meister, A., Scott, S.J. (1961). *Biochem. Preparations* 8, 11
- Orgel, L.E. (1973). *The origins of life*. New York: Wiley
- Paecht-Horowitz, M., Berger, J., Katchalsky, A. (1970). *Nature* 228, 636
- Paecht-Horowitz, M., Katchalsky, A. (1967). *Biochim. Biophys. Acta* 140, 14
- Paecht-Horowitz, M., Katchalsky, A. (1973). *J. Mol. Evol.* 2, 91
- Ponnamperuma, C. (1971). *Quart. Rev. Biophys.* 4, 77
- Posner, A.M., Quirk, J.P. (1964). *Proc. Roy. Soc. A* 278, 35

Dr. Richard M. Lemmon
Laboratory of Chemical Biodynamics
University of California
Berkeley, California 94720, USA