

Scaling Limits for Interacting Diffusions^{*}

S. R. S. Varadhan

Courant Institute of Mathematical Sciences, New York University, 251 Mercer Street,
New York, NY 10012, USA

Received February 26, 1990; in revised form June 22, 1990

Abstract. We consider a large number of particles diffusing on a circle interacting through a drift resulting from the gradient of a pair potential whose support is of the order of the interparticle distance. We derive a nonlinear bulk diffusion equation for the density of the particle distribution on the circle. The diffusion coefficient is determined as a function of density in terms of standard thermodynamical objects.

1. Introduction

In this article we study the hydrodynamic limit for interacting Brownian motions on the one-dimensional circle. The interaction is between pairs of particles and is repulsive in nature. The scaling is such that the range of the interaction is of the same order as interparticle distance and therefore each particle interacts with only a finite number of nearby particles at any given time.

We obtain a nonlinear bulk diffusion equation and the diffusion coefficient is naturally expressed in terms of the thermodynamic functions of our one-dimensional system. The main limitations are the finiteness of volume that is forced because our basic space is the circle, the repulsive nature of the interactions that is assumed and that we are in one space dimension.

In this context the fluctuations around equilibrium have been studied earlier by H. Spohn [5] and the self diffusion in equilibrium by Guo and Papanicolaou [1]. Their results have been derived for infinite volume and arbitrary space dimension. Of course these results deal essentially with equilibrium problems where a lot of control through estimates is available.

Our method is similar to the one used in Guo, Papanicolaou and Varadhan [2] and uses estimates based on entropy and its rate of change. Section 2 describes

^{*} This research was supported by a grant from the National Science Foundation DMS-89-01682

the results precisely and Sects. 3-9 contain detailed proofs. Section 10 contains some results concerning the Gibbs measures of one dimensional systems. These results are all well known to the experts although it is difficult to find precise references to the results in the form we need. We have therefore included a quick exposition with sketches of proofs.

Along with S. Olla in [3] we have modified the methods of this article to extend these results to the case of interacting Ornstein-Uhlenbeck processes and they will appear in the next article.

2. Summary

We denote by S the circle of unit circumference and consider a system of N interacting Brownian motions with S as a state space satisfying the following system of stochastic differential equations:

$$dx_i(t) = -N \sum_{j:j \neq i} V'(N(x_i(t) - x_j(t)))dt + d\beta_i(t) , \quad i = 1, 2, \dots, N . \quad (2.1)$$

Here, β_1, \dots, β_N are N independent Brownian motions and $V(\cdot)$ is an even function on R satisfying the following assumptions:

- (i) $V \geq 0, V(0) > 0$ and V has compact support.
- (ii) V is once continuously differentiable.
- (iii) $V(\cdot)$ is repulsive in the sense that

$$\psi(z) = -zV'(z) \geq 0 . \quad (2.2)$$

Then the process $[x_1(t), \dots, x_N(t)]$ is a Markov process of diffusion type on S^N , the N -fold copy of S , with an infinitesimal generator given by

$$L_N = \frac{1}{2} \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} - N \sum_{j \neq i} V'(N(x_i - x_j)) \frac{\partial}{\partial x_j} . \quad (2.3)$$

It is easily verified that

$$L_N = \frac{1}{2} \sum_{i=1}^N \left(e^{\sum_{j \neq i} V(N(x_j - x_k))} \frac{\partial}{\partial x_i} e^{-\sum_{j \neq i} V(N(x_j - x_k))} \frac{\partial}{\partial x_j} \right) , \quad (2.4)$$

so that L_N is formally symmetric with respect to the measure

$$d\mu_N(x) = \frac{1}{Z_N} \exp \left[- \sum_{i,j} V(N(x_i - x_j)) \right] dx_1 \dots dx_N . \quad (2.5)$$

It is straightforward to verify that the Markov process $x(t)$ with L_N as generator is in fact reversible with respect to the invariant measure μ_N given by (2.5). The constant Z_N is for normalization and is so defined to make μ_N into a probability measure.

We start $x(0)$ with an initial distribution with density $f_N^0(x_1, \dots, x_N)$ with respect to the invariant density μ_N . The density at time t with respect to μ_N is denoted by $f_N^t(x_1, \dots, x_N)$ and is given as a solution of the forward equation

$$\frac{\partial f_N^t}{\partial t} = L_N f_N^t \quad \text{with} \quad f_N^t|_{t=0} = f_N^0 . \quad (2.6)$$

The empirical distribution of the process at time t is defined by

$$\xi_N(t, A) = \frac{1}{N} \sum_{i=1}^N \chi_A(x_i(t)) \quad \text{for } A \subset S, \tag{2.7}$$

and $\xi_N(t)$ is viewed as a random measure on S . If we denote by $M_1(S)$ the space of probability measures on S , one can view $\xi_N(\cdot)$ as a stochastic process with values in $M_1(S)$. In view of the continuity of the trajectories, we can pick a time $T < \infty$ and consider the space $C[[0, T], M_1(S)]$ of measure valued continuous functions on $[0, T]$, and our basic markov process with initial density f_N^0 will induce a measure Q_N on $C[[0, T], M_1(S)]$. Our main result is to show that under suitable assumptions on f_N^0 , the measures Q_N as $N \rightarrow \infty$ will concentrate on a single measure valued trajectory which is the solution of a certain nonlinear diffusion equation.

In order to describe this nonlinear diffusion equation we have to introduce some thermodynamic functions of one dimensional systems with pair interaction given by $V(\cdot)$. The partition function in a finite region $[0, l]$ with activity λ is given by

$$\hat{Z}(l, \lambda) = \sum_{n=0}^{\infty} \frac{e^{n\lambda}}{n!} \int_0^l \dots \int_0^l e^{-\sum_{i,j} V(x_i - x_j)} dx_1 \dots dx_n. \tag{2.8}$$

It is known [4] that the free energy defined by

$$F(\lambda) = \lim_{l \rightarrow \infty} \frac{\log \hat{Z}(l, \lambda)}{l} \tag{2.9}$$

exists and is a convex function of λ for all λ ,

$$\rho(\lambda) = \frac{dF}{d\lambda}, \tag{2.10}$$

is the ‘‘density’’ corresponding to the activity λ and is a continuous strictly monotone function of λ . This function of course can be inverted to yield $\lambda = \lambda(\rho)$ as a function of ρ . The free energy expressed as a function of ρ , i.e.

$$P(\rho) = F(\lambda(\rho)) \tag{2.11}$$

is called the pressure and is again a continuous strictly monotone function of the density ρ .

Our main result can now be stated. Let us assume that the initial densities $\{f_N^0\}$ have the following behavior as $N \rightarrow \infty$:

$$\int f_N^0 \log f_N^0 d\mu_N \leq CN \quad \text{for some } C < \infty \text{ and all } N; \tag{2.12}$$

$$\int_{A_{N,J,\varepsilon}} f_N^0 d\mu_N \rightarrow 0 \quad \text{as } N \rightarrow \infty \tag{2.13}$$

for every smooth test function J on S . Here $A_{N,J,\varepsilon}$ is the set of configurations (x_1, \dots, x_N) such that

$$A_{N,J,\varepsilon} = \left\{ x : \left| \frac{1}{N} \sum J(x_i) - \int J(\theta) \rho_0(\theta) d\theta \right| > \varepsilon \right\},$$

and $\rho_0(\theta)$ is some fixed density on S such that $\rho_0(\theta) \geq 0$ and $\int \rho_0(\theta) d\theta = 1$. In

order for (2.12) and (2.13) to be compatible $\rho_0(\theta)$ will have to satisfy some minimal regularity conditions. In any case (2.13) asserts that

$$\frac{\delta_{x_1} + \dots + \delta_{x_N}}{N} \Rightarrow \rho_0(\theta)d\theta$$

in the sense of weak convergence, in probability with respect to the initial distribution $\int_N^0 d\mu_N$.

Under these assumptions we establish that Q_N converges weakly to a limit which is the degenerate distribution on a single trajectory in $M_1(S)$ described by $\rho(\theta, t)d\theta$, where $\rho(\theta, t)$ is defined as the unique solution (satisfying mild regularity conditions) of

$$\frac{\partial \rho(\theta, t)}{\partial t} = \frac{1}{2} [P(\rho(\theta, t))]_{\theta\theta} \tag{2.14}$$

with

$$\rho(\theta, t)|_{t=0} = \rho_0(\theta) . \tag{2.15}$$

3. Outline of Proof

The proof of our result is based on the following ideas. For the sequence $\{Q_N\}$ of probability measures on the space $\Omega_T = C[[0, T], M_1(S)]$ of measure valued trajectories we first establish compactness under weak convergence.

Theorem 3.1. *The sequence $\{Q_N\}$ is tight on the space Ω_T .*

In order to establish that the limit Q_N exists and is concentrated on a single trajectory we establish properties satisfied by the support of any limit point Q , enough properties so that one is left with a single trajectory satisfying all of them.

Let us consider a smooth function $u(\theta)$ on S and the corresponding functional

$$\xi_N(t) = \frac{1}{N} \sum u(x_i(t)) .$$

By Itô's formula we see that

$$\begin{aligned} d\xi_N(t) &= \frac{1}{2N} \sum_{i=1}^N u''(x_i(t)) dt \\ &\quad - \sum_{i,j=1}^N V'(N(x_i(t)) - x_j(t)) u'(x_i(t)) dt \\ &\quad + \frac{1}{N} \sum_{i=1}^N u'(x_i(t)) d\beta_i(t) . \end{aligned} \tag{3.1}$$

By the law of large numbers, as $N \rightarrow \infty$, the third term goes to zero. As for the second term

$$\begin{aligned}
 & \sum_{i,j=1}^N V'(N(x_i(t) - x_j(t)))u'(x_i(t)) \\
 &= \frac{1}{2} \sum V'(N(x_i(t) - x_j(t)))(u'(x_i(t)) - u'(x_j(t))) \\
 & \quad \text{(by skew symmetry of } V') \\
 &\simeq \frac{1}{2} \sum V'(N(x_i(t) - x_j(t)))u''(x_i(t))(x_i(t) - x_j(t)) \\
 & \quad \text{by (mean value theorem)} \\
 &\simeq -\frac{1}{2N} \sum \psi(N(x_i(t) - x_j(t)))u'(x_i(t)) .
 \end{aligned}$$

If we are at some point (θ, t) in space time and the local density of particles is $\rho(\theta, t)$ then the system should be in local equilibrium there in a Gibbs state with density ρ and it is not a difficult calculation to show that in an exact Gibbs state of density ρ

$$\lim_{l \rightarrow \infty} \frac{1}{l} \sum_{x_i, x_j \in [0, l]} \psi(x_i - x_j) = P(\rho) - \rho . \tag{3.2}$$

One therefore hopes to establish that if at a typical time t , $\frac{1}{N}[\delta_{x_1(t)} + \dots + \delta_{x_N(t)}]$ is close to $\rho(t, \theta)d\theta$, then $\frac{1}{2N} \sum \psi(N(x_i(t) - x_j(t)))u''(x_i(t)) + \frac{1}{2N} \sum u''(x_i(t))$ should be close to $\frac{1}{2} \int P(\rho(t, \theta))u''(\theta)d\theta$.

In this manner we establish

Theorem 3.2. *Any limit point Q of Q_N satisfies*

- (i) $Q[\alpha(t, \cdot) : \alpha(t, d\theta) = \rho(t, \theta)d\theta \text{ for a.e. } t] = 1$.
- (ii) $E^Q[\rho : \rho(0, \theta) = \rho_0(\theta) \text{ a.e. } \theta] = 1$.
- (iii) $Q[P : \int u(\theta)\rho(t, \theta)d\theta - \int u(\theta)\rho_0(\theta)d\theta = \frac{1}{2} \int_0^t \int u''(\theta)P(\rho(t, \theta))d\theta] = 1$.

Given Theorem 3.2, Q lives on the set of weak solutions of

$$\begin{cases} \frac{\partial \rho}{\partial t} = \frac{1}{2}(P(\rho))_{\theta\theta} & \text{with} \\ \rho(t, \theta)|_{t=0} = \rho_0(\theta) . \end{cases} \tag{3.3}$$

One can establish the uniqueness of weak solutions of (3.3) under the assumptions

$$\int_0^T \int_S \rho^3(t, \theta)dt d\theta < \infty \tag{3.4}$$

and

$$\int_0^T \int_s \frac{1}{\rho(t, \theta)} \left[\frac{\partial}{\partial \theta} P(\rho(t, \theta)) \right]^2 d\theta < \infty . \tag{3.5}$$

In order then to complete the proof we only need to establish

Theorem 3.3. *Any limit point Q satisfies*

$$E^Q \left[\int_0^T \int_s \rho^3(t, \theta) dt d\theta \right] < \infty$$

and

$$E^Q \left[\int_0^T \int_s \frac{1}{\rho(t, \theta)} \left[\frac{d}{d\theta} P(\rho(t, \theta)) \right]^2 dt d\theta \right] < \infty .$$

Finally a word about the strategy to be used in establishing Theorems 3.1., 3.2 and 3.3. Since $N \rightarrow \infty$ we need to develop estimates that hold uniformly in N . We will need properties of the solutions f'_N , and $\tilde{f}_N = \frac{1}{T} \int_0^T f_N^s ds$, where f'_N solves

$$\frac{\partial f'_N}{\partial t} = L_N f'_N .$$

Our assumptions is that $\int f_N^0 \log f_N^0 d\mu_N \leq CN$. One knows that

$$\int f'_N \log f'_N d\mu_N = H_N(t)$$

is nonincreasing in t and

$$\frac{d}{dt} H_N(t) = -I_N(t) = -\frac{1}{2} \int \frac{|\nabla f'_N|^2}{f'_N} d\mu_N .$$

The functionals

$$H_N(t) = \int f \log f d\mu_N$$

and

$$I_N(t) = \frac{1}{2} \int \frac{|\nabla f|^2}{f} d\mu_N$$

are lower semicontinuous, non-negative and convex. Therefore one gets easily

$$\begin{aligned} H_N(\tilde{f}_N) &= H_N \left(\frac{1}{T} \int_0^T f_N^s ds \right) \\ &\leq \frac{1}{T} \int_0^T H(f_N^s) ds \\ &\leq H(f_N^0) \leq CN . \end{aligned}$$

Similarly,

$$\begin{aligned}
I_N(\tilde{f}_N) &\leq \frac{1}{T} \int_0^T I_N(f_N^s) ds \\
&= \frac{1}{T} [H_N(f_N^0) - H_N(f_N^T)] \\
&\leq \frac{1}{T} H_N(f_N^0) \\
&\leq \frac{C}{T} N .
\end{aligned}$$

In some sense our procedures are based solely on the two estimates derived above.

4. Some Estimates Based on Entropy

Given a density f_N relative to the invariant measure μ_N with entropy $H_N(f_N) = \int f_N \log f_N d\mu_N$, we shall obtain in this section some preliminary estimates based on a bound of the type

$$H_N(f_N) \leq AN . \quad (4.1)$$

Lemma 4.1. *There exists a constant C such that for every N and every f_N satisfying (4.1) we have*

$$E^{f_N} \left\{ \frac{1}{N} \sum_{i,j} V(N(x_i - x_j)) \right\} \leq A + C ,$$

where A is the same constant as in (4.1).

Proof.

$$\begin{aligned}
&E^{f_N} \left\{ \frac{1}{N} \sum_{i,j} V(N(x_i - x_j)) \right\} \\
&\leq \frac{1}{N} \log E^{\mu_N} \left\{ \exp \left[\sum V(N(x_i - x_j)) \right] \right\} + A \\
&\leq -\frac{1}{N} \log Z_N + A = -\frac{1}{N} \log \int e^{-\sum V(N(x_i - x_j))} dx_1 \dots dx_N + A \\
&\leq \frac{1}{N} \int \left(\sum V(N(x_i - x_j)) \right) dx_1 \dots dx_N + A \\
&\leq C + A .
\end{aligned}$$

From now on for the rest of the section we want to derive some consequences of an estimate of the form

$$E^{f_N} \left\{ \frac{1}{N} \sum V(N(x_i - x_j)) \right\} \leq B \quad \text{for all } N . \quad (4.2)$$

Let us divide the circle S into a large number l of small arcs length $\frac{1}{l}$. Let us denote them by S_1, S_2, \dots, S_l . If we are given a configuration of N points x_1, \dots, x_N we can form the frequency counts

$$g_i = \sum_{j=1}^N \chi_{S_i}(x_j)$$

of the number of points among x_1, \dots, x_N that fall in a given arc S_i for $1 \leq i \leq l$. We wish to estimate $\sum_{i=1}^l g_i^2$ in terms of $\sum_{i,j} V(N(x_i - x_j))$. Let us subdivide each arc S_i into $2k$ arcs of length $\frac{1}{2lk}$ and we assume that the size $\frac{1}{2lk}$ is of the order $\frac{\delta}{N}$, where δ is small, independent of N , and is so small that $V(x) \geq \eta$ for $|x| \leq \delta$ for some positive constant $\eta > 0$. For $1 \leq i \leq 2lk$, if we denote by h_i the frequency counts for the smaller arcs, then

$$\sum_1^l g_i^2 \leq 2k \sum_1^{2lk} h_i^2 .$$

If we translate the basic arcs S_i by half their lengths and denote them by $S_{i+1/2}$ with their frequency counts $g_{i+1/2}$, then we have just as well

$$\sum_1^l g_{i+1/2}^2 \leq 2k \sum_1^{2lk} h_i^2 .$$

If from among x_1, \dots, x_N , we have a pair x_r, x_s with $|x_r - x_s| \leq \frac{1}{2l}$, then they must belong either to the same S_i or $S_{i+1/2}$ for some i . Therefore

$$\begin{aligned} \sum_{i,j} \chi_{|x_i - x_j| \leq 1/2} (|x_i - x_j|) &\leq \sum g_i^2 + \sum g_{i+1/2}^2 \\ &\leq (4k) \sum_1^{2lk} h_i^2 \\ &\leq \frac{4k}{\eta} \sum V(N(x_i - x_j)) . \end{aligned} \tag{4.3}$$

If $W(x)$ is a function that is supported on $\left[-\frac{1}{2l}, \frac{1}{2l}\right]$ and is bounded there by a constant $\|W\|$ then

$$\sum_{i,j} W(x_i - x_j) \leq \|W\| \cdot \frac{4k}{\eta} \sum V(N(x_i - x_j)) , \tag{4.4}$$

provided only that $\frac{N}{2lk} = \delta$ is sufficiently small.

Lemma 4.2. *If W is a fixed compactly supported function then we can estimate*

$$E^{f_N} \frac{\lambda}{N^2} \sum W(\lambda(x_i - x_j)) \leq \|W\| \cdot B \cdot C ,$$

where $\|W\|$ is a bound for W , B is any constant satisfying (4.2) and C is a universal constant depending only on V and the size of the support of $W(\cdot)$. This estimate holds uniformly for all λ in the range $\varepsilon_0 \leq \lambda \leq N$.

Proof. We need basically l to be $c\lambda$ for fixed constant c in (4.4) and that forces on us a choice of $k = CN/\lambda$ for some other constant C . Now Lemma 4.2 is an immediate consequence of (4.2) and (4.4).

5. Some Estimates Based on the Dirichlet Form

In this section we want to explore some of the consequences of assuming that the density f_N satisfies

$$E^{f_N} \left[\frac{1}{N} \sum V(N(x_i - x_j)) \right] \leq B \tag{5.1}$$

and

$$\frac{1}{2} \int |\nabla f_N|^2 \frac{1}{f_N} d\mu_N = I_N(f_N) \leq DN . \tag{5.2}$$

The basic estimate uses integration by parts for any “suitable” test function $u = u(x_1, \dots, x_N)$

$$\begin{aligned} |E^{f_N}[L_N u]| &= \left| \int (L_N u) f_N d\mu_N \right| \\ &= \left| \frac{1}{2} \int \langle \nabla u, \nabla f_N \rangle d\mu_N \right| \\ &\leq \left(\frac{1}{2} \int |\nabla u|^2 f_N d\mu_N \right)^{1/2} \left(\frac{1}{2} \int |\nabla f_N|^2 \frac{1}{f_N} d\mu_N \right)^{1/2} \\ &\leq \left(\frac{1}{2} \int |\nabla u|^2 f_N d\mu_N \right)^{1/2} (DN)^{1/2} . \end{aligned} \tag{5.3}$$

Of course such an estimate is only as good as the test functions u that we can find to use in the estimate. The following class of test functions will be particularly useful:

$$u_N(x_1, \dots, x_N) = \sum_{i,j} \mathcal{G}_{N,\varepsilon,\lambda}(x_i - x_j) , \tag{5.4}$$

where $\mathcal{G}_{N,\varepsilon,\lambda}(x)$ is a function on R that has sufficiently small support that it can be viewed as a function on S . For each fixed λ and ε we will have a choice of u_N made for all sufficiently large N . The function $\mathcal{G}_{N,\varepsilon,\lambda}(x)$ is given by

$$\mathcal{G}_{N,\varepsilon,\lambda}(x) = \int_{-\infty}^x G_{N,\varepsilon,\lambda}(y) dy \tag{5.5}$$

and

$$G_{N,\varepsilon,\lambda}(x) = \int_{-\infty}^x g_{N,\varepsilon,\lambda}(y) dy , \tag{5.6}$$

where $g_{N,\varepsilon,\lambda}(\cdot)$ is a function R with support in a small interval around the origin,

satisfying in addition

$$\int_{-\infty}^{\infty} g_{N,\varepsilon,\lambda}(y)dy = 0 \tag{5.7}$$

and

$$\int_{-\infty}^{\infty} yg_{N,\varepsilon,\lambda}(y)dy = 0 , \tag{5.8}$$

so that G and \mathcal{G} have the same support as g . The function $g_{N,\varepsilon,\lambda}(y)$ will be of the form

$$g_{N,\varepsilon,\lambda}(y) = (N\varepsilon)g(N\varepsilon y) - \lambda g(\lambda y) , \tag{5.9}$$

where g is a nonnegative, smooth, symmetric function with compact support with total integral 1,

$$\int g(y)dy = 1 . \tag{5.10}$$

Clearly if λ is big enough and $\varepsilon > 0$ is arbitrary then for sufficiently large N , $u_N(x_1, \dots, x_N)$ is well defined.

In order to use the integration by parts estimate (5.3) we have to compute $\mathcal{L}_N u_N$ and $|\nabla u_N|^2$,

$$\begin{aligned} \mathcal{L}_N u_N &= N\varepsilon \sum g(N\varepsilon(x_i - x_j)) - \lambda \sum g(\lambda(x_i - x_j)) \\ &\quad - 2N \sum_i \left(\sum_j [G(N\varepsilon(x_i x_j)) - G(\lambda(x_i - x_j))] \right) \\ &\quad \cdot \left(\sum_k V'(N(x_i - x_k)) \right) , \end{aligned} \tag{5.11}$$

$$|\nabla u_N|^2 = 4 \sum_i \left| \sum_j [G(N\varepsilon(x_i - x_j)) - G(\lambda(x_i - x_j))] \right|^2 . \tag{5.12}$$

One can use the fact that V' is an odd function to rewrite (5.11) in the form

$$\begin{aligned} \mathcal{L}_N u_N &= N\varepsilon \sum g(N\varepsilon(x_i - x_j)) - \lambda \sum g(\lambda(x_i - x_j)) \\ &\quad - N \sum_{i,j,k} [G(N\varepsilon(x_i - x_j)) - G(N\varepsilon(x_k - x_j))] \\ &\quad - G(\lambda(x_i - x_j)) + G(\lambda(x_k - x_j))] V'(N(x_i - x_k)) . \end{aligned} \tag{5.12}$$

Let us take the special case of $\varepsilon = 1$ and $\lambda = 1$. We assume that the support of g is small enough that $\lambda = 1$ works. Otherwise we can take any value λ_0 for λ .

We can write inequality (5.3) in the form

$$\left| E^{f_N} \frac{1}{N^2} \mathcal{L}_N \mathcal{U}_N \right| \leq \frac{D^{1/2}}{N^{3/2}} \left[E^{f_N} \left\{ \frac{1}{2} |\nabla u_N|^2 \right\} \right]^{1/2} , \tag{5.13}$$

with $\varepsilon = 1$, the first term on the right-hand side of (5.11) is

$$E^{f_N} \left\{ \frac{1}{N} \sum g(N(x_i - x_j)) \right\} ,$$

which is uniformly bounded because of (5.1) and Lemma 4.2. The second term is

$$E^{f_N} \left\{ \frac{1}{N^2} \sum g(x_i - x_j) \right\} ,$$

which is bounded by a bound for g . The term

$$\begin{aligned} & \frac{1}{N} \sum_{i,j,k} [G(x_i - x_j) - G(x_k - x_j)] V'(N(x_i - x_k)) \\ & \leq \frac{C}{N} \sum_{i,j,k} |x_i - x_k| |V'(N(x_i - x_k))| \end{aligned}$$

by using a bound on the derivative of G . We can use the function $\psi(x) = -xV'(x)$ and write this term as

$$\frac{C}{N^2} \sum_{i,j,k} \psi(N(x_i - x_k)) \leq \frac{C}{N} \sum_{i,k} \psi(N(x_i - x_k)) ,$$

which is bounded by (5.1) and Lemma 4.2. It is easy to estimate

$$|\nabla u_N|^2 \leq CN^3$$

by using a bound for G . Since in the inequality (5.3) every term has been bounded except for a single term we obtain

$$E^{f_N} \left\{ \frac{1}{N} \sum_{i,j,k} [G(N(x_k - x_j)) - G(N(x_i - x_j))] V'(N(x_i - x_k)) \right\} \leq C , \quad (5.14)$$

where C is a bound depending only on B and D . The function $F(x - y, y - z) = V'(x - z)[G(z - y) - G(x - y)]$ is nonnegative and one can check easily that it is possible to pick g in such a manner that

$$F(x - y, y - z) \geq \psi(x - z)\psi(x - y) . \quad (5.15)$$

From (5.14) and (5.15) we obtain

$$\begin{aligned} E^{f_N} \left\{ \left(\frac{1}{N} \sum_{i,j} \psi(N(x_i - x_j)) \right)^2 \right\} & \leq E^{f_N} \left\{ \frac{1}{N} \sum_i \left(\sum_j \psi(N(x_i - x_j)) \right)^2 \right\} \\ & = E^{f_N} \left\{ \frac{1}{N} \sum_{i,j,k} \psi(N(x_i - x_j)) \psi(N(x_i - x_k)) \right\} \\ & \leq E^{f_N} \left\{ \frac{1}{N} \sum_{i,j,k} F(N(x_i - x_j), N(x_j - x_k)) \right\} \\ & \leq C . \end{aligned}$$

In other words (5.1) alone is sufficient to provide a bound for

$$E^{f_N} \left\{ \frac{1}{N} \sum \psi(N(x_i - x_j)) \right\}$$

while (5.1) and (5.2) provide a bound for

$$E^{f_N} \left\{ \frac{1}{N} \sum \psi(N(x_i - x_j)) \right\}^2 .$$

We have therefore proved Lemma 5.1.

Lemma 5.1. *There is a constant C depending only on B and D such that*

$$E^{f_N} \left\{ \frac{1}{N} \sum \psi(N(x_i - x_j)) \right\}^2 \leq C ,$$

$$E^{f_N} \left\{ \frac{1}{N} \sum_i \left[\sum_j \psi(N(x_i - x_j)) \right]^2 \right\} \leq C .$$

6. Compactness

In this section we will establish Theorem 3.1. Since the state space $M_1(S)$ is compact we need only establish estimates of modulus of continuity to appeal to Prohorov's theorem. It is therefore enough to prove

Lemma 6.1. *For any smooth function J on S, and $\epsilon > 0$,*

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} Q_N \left[\sup_{\substack{0 \leq s < t \leq T \\ |s-t| \leq \delta}} \left| \frac{1}{N} \sum J(x_i(s)) - \frac{1}{N} \sum J(x_i(t)) \right| \geq \epsilon \right] = 0 . \quad (6.1)$$

Proof. Let us take $s < t$. Then

$$\begin{aligned} & \frac{1}{N} \sum J(x_i(t)) - \frac{1}{N} \sum J(x_i(s)) \\ &= \frac{1}{2N} \sum_s^t J''(x_i(\sigma)) d\sigma + \frac{1}{N} \int_s^t \sum J'(x_i(\sigma)) d\beta_i(\sigma) \\ & \quad - \int_s^t \sum V'(N(x_i(\sigma)) - x_j(\sigma)) J'(x_i(\sigma)) d\sigma \\ &= \eta_1 + \eta_2 + \eta_3 . \end{aligned} \quad (6.2)$$

The first term is clearly bounded by

$$|\eta_1| \leq \frac{1}{2} C(t-s) , \quad (6.3)$$

where C is a bound for J'' .

The second term is a Martingale term and by Doob's inequality

$$\begin{aligned} & E \sup_{0 \leq t \leq T} \left| \frac{1}{N} \sum_0^t J'(x_i(\sigma)) d\beta_i(\sigma) \right|^2 \\ & \leq C \exp \left\{ E \left| \frac{1}{N} \sum_0^t J'(x_i(\sigma)) d\beta_i(\sigma) \right|^2 \right\} \end{aligned}$$

$$\begin{aligned}
 &= C \frac{1}{N^2} \int_0^T \sum |J'(x_i(\sigma))|^2 d\sigma \\
 &\leq \frac{C \cdot C_1^2 T}{N},
 \end{aligned}
 \tag{6.4}$$

where C_1 is a bound for J' .

The third term can be estimated by estimating the integrand

$$\begin{aligned}
 &\sum_{ij} V'(N(x_i(\sigma) - x_j(\sigma))) J'(x_i(\sigma)) \\
 &= \sum_{ij} V'(N(x_i(\sigma) - x_j(\sigma))) (J'(x_i(\sigma)) - J'(x_j(\sigma))) \\
 &\leq C \sum_{i,j} |V'(N(x_i(\sigma) - x_j(\sigma)))| |x_i(\sigma) - x_j(\sigma)|,
 \end{aligned}$$

where C is a bound for J'' . Recalling that $\psi(z) = -zV'(z)$ we can estimate

$$|\eta_3| \leq C \int_s^t \frac{1}{N} \sum |\psi(N(x_i(\sigma) - x_j(\sigma)))| d\sigma.$$

One can then complete the proof of Lemma 4.1 if we show the following estimate:

$$\sup_N E^{\mathcal{Q}_N} \int_0^T \left[\frac{1}{N} \sum \psi(N(x_i(\sigma) - x_j(\sigma))) \right]^2 d\sigma < \infty.$$

This follows from Lemma 5.1.

7. Local Gibbs States

In establishing the hydrodynamic scaling limit one of the problems we have to deal with is expressing average microscopic quantities like $\frac{1}{N} \sum \psi(N(x_i - y_j)) U(x_i)$ purely in terms of the macroscopic density function. This is of course possible only if the configuration x_1, \dots, x_n is suitably organized and we have to establish that during our stochastic dynamics most of the time the configurations are suitably organized with a very high probability.

To make this precise let us fix a value of l and consider a functional $H(\omega)$ which is a bounded and continuous functional of the configuration of points in the interval $(-l, l)$. We can view $H(\omega)$ as a functional of a point process on R , which depends however only on the configuration of points in $(-l, l)$. If we have a set (x_1, \dots, x_N) of N points in S , for any given x , we can look at the points $x_i: |x_i - x| < \frac{l}{4}$ and consider the points $N(x_i - x)$ on the line $(-\infty, \infty)$ obtained from the original (x_1, \dots, x_N) on S . The resulting point process on R , while it is somewhat arbitrary due to the cutoff at the edges namely around $\pm \frac{N}{4}$, for every l the configuration is natural and well defined on $[-l, l]$, provided N is large enough. We can evaluate our functional $H(\omega)$ at this configuration, which we denote by ω_N^x and obtain a value $H(\omega_N^x)$.

We are interested in the quantity

$$\xi_N(x_1, \dots, x_N) = \int_S H(\omega_N^x) U(x) dx . \tag{7.1}$$

Corresponding to the set x_1, \dots, x_N we have the empirical distribution

$$\alpha_N(dx) = \frac{1}{N} (\delta_{x_1} + \dots + \delta_{x_N}) .$$

We pick a mollifier $h(x)$ with compact support, i.e. a nonnegative function $h(x)$ with $\int h(x) dx = 1$, and assuming that the support of h is contained in $[-\frac{1}{2}, \frac{1}{2}]$, we can lift h as a function on S , and for $\lambda \geq 1$, $\lambda h(\lambda x)$ is a mollifier on S as well. We can then pick for $\lambda \geq 1$, a natural version of local density for the configuration (x_1, \dots, x_N) , by

$$\begin{aligned} \rho_\lambda(x) &= \frac{\lambda}{N} \sum h(\lambda(x_i - x)) \\ &= (h_\lambda * \alpha_N)(x) , \quad \text{where} \\ h_\lambda &= \lambda h(\lambda x) . \end{aligned} \tag{7.2}$$

For a one dimensional system, there are no phase transitions and we have a unique Gibbs state P_ρ viewed as a stationary point process with density ρ that corresponds to the given pair interaction $V(x)$. Since we expect the arrangement of points near x to be a random arrangement from P_ρ with $\rho = \rho_\lambda$, by (7.1) we expect to replace $\xi_N(x_1, \dots, x_N)$ by $\eta_{N,\lambda}(x_1, \dots, x_N)$, where

$$\eta_{N,\lambda}(x_1, \dots, x_N) = \int_S \hat{H}(\rho_\lambda(x)) U(x) dx \tag{7.3}$$

with

$$\hat{H}(\rho) = E^{P_\rho} [H(\omega)] . \tag{7.4}$$

The error between ξ_N and $\eta_{N,\lambda}$ is expected to be small provided λ is large and N is large and (x_1, \dots, x_N) is a typical configuration from a nice distribution $f_N d\mu_N$.

Let us denote by $\mathcal{A}_{N,B,D}$ the set of probability densities f_N on S^N that satisfy

$$F^{f_N} \left\{ \frac{1}{N} \sum V(N(x_i - x_j)) \right\} \leq B , \tag{7.5}$$

$$\frac{1}{2} \int \frac{|\nabla f_N|^2}{f_N} d\mu_N \leq DN . \tag{7.6}$$

Because of translation invariance of the expressions (7.5) and (7.6), if we replace f_N by

$$\hat{f}_N(x_1, \dots, x_N) = \int_S f_N(x_1 + a, \dots, x_N + a) da$$

due to linearity in the case of (7.5) and convexity in the case of (7.6), \hat{f}_N will satisfy the inequalities (7.5) and (7.6). In other words, if $f_N \in \mathcal{A}_{N,B,D}$ so does \hat{f}_N . Let us denote by $\mathcal{A}_{N,B,D}^0$, the translation invariant densities in $\mathcal{A}_{N,B,D}$. Our goal is to establish that for each fixed l and $H(\cdot)$

Theorem 7.1.

$$\lim_{\lambda \rightarrow \infty} \limsup_{N \rightarrow \infty} \sup_{f_N \in \mathcal{A}_{N,B,D}} E^{f_N} |\xi_N - \eta_{N,\lambda}| = 0 .$$

Since P_ρ depends continuously on ρ , $\hat{H}(\rho)$ is continuous in ρ and we can establish Theorem 7.1 in two steps.

Theorem 7.2.

$$\lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{f_N \in \mathcal{A}_{N,B,D}} E^{f_N} |\xi_N - \eta_{N,N\varepsilon}| = 0 ,$$

and

Theorem 7.3.

$$\lim_{\lambda \rightarrow \infty} \limsup_{N \rightarrow \infty} \sup_{f_N \in \mathcal{A}_{N,B,D}} E^{f_N} \int_S |\rho_\lambda(x) - \rho_{\varepsilon N}(x)| dx = 0 .$$

Remark. In view of the comment made earlier about translation invariance in Theorems 7.1, 7.2 and 7.3 we can in the proof replace $\mathcal{A}_{N,B,D}$ by $\mathcal{A}_{N,B,D}^0$.

We can write

$$\begin{aligned} & E^{f_N} \left| \int H(\omega_N^x) U(x) dx - \int \hat{H}(\rho_\lambda(x)) U(x) dx \right| \\ & \leq E^{f_N} \left| \int \left[\frac{N}{2a} \int_{|y-x| \leq a/N} H_0(\omega_N^x) dx \right] U(y) dy - \int \hat{H}(\rho_\lambda(x)) U(x) dx \right| \\ & \quad + \|H\| E^{f_N} \int \frac{N}{2a} \int_{|U-x| \leq a/N} |U(y) - u(x)| dy dx . \end{aligned}$$

To fixed a, H and any sequence $\lambda = \lambda_N$,

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \sup_{f_N \in \mathcal{A}_{N,B,D}} E^{f_N} \left| \int H(\omega_N^x) U(x) dx - \int \hat{f}(\rho_{\lambda_N}(x)) U(x) dx \right| \\ & \leq \|U\| \limsup_{N \rightarrow \infty} \sup_{f_N \in \mathcal{A}_{N,B,D}} E^{f_N} \left[\int dy \left| \frac{N}{2a} \int_{|y-x| \leq a/N} H(\omega_N^x) dx - \hat{H}(\rho_{\lambda_N}(y)) \right| \right] \end{aligned}$$

by assuming continuity of the function $U(x)$.

If we now replace f_N by \hat{f}_N then

$$\begin{aligned} & E^{\hat{f}_N} \left[\int dy \left| \frac{N}{2a} \int_{|y-x| \leq a/N} H(\omega_N^x) dx - \hat{H}(\rho_{\lambda_N}(y)) \right| \right] \\ & = E^{\hat{f}_N} \left[\int dy \left| \frac{N}{2a} \int_{|y-x| \leq a/N} H(\omega_N^x) dx - \hat{H}(\rho_{\lambda_N}(y)) \right| \right] \\ & = E^{\hat{f}_N} \left[\left| \frac{N}{2a} \int_{|x| \leq a/N} H(\omega_N^x) dx - \hat{H}(\rho_{\lambda_N}(0)) \right| \right] . \end{aligned}$$

The proof of Theorem 7.2 can therefore be reduced to the proof of the following theorem.

Theorem 7.4.

$$\overline{\lim}_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{f_N \in \mathcal{A}_{N,B,D}^0} E^{f_N} \left| \frac{N\varepsilon}{2} \int_{|x| \leq 1/N\varepsilon} H(\omega_N^x) dx - \hat{H}(\rho_{N\varepsilon}(0)) \right| = 0 .$$

The proof of Theorem 7.4 proceeds as follows.

The computation of $\int_{|x| \leq 1/N\varepsilon} H(\omega_N^x) dx$ and $\rho_{N\varepsilon}(0)$ requires only a knowledge of the location of the particles from among x_1, \dots, x_N that belong to some interval of the form $|x| \leq \frac{1}{N\varepsilon} + \frac{l_0}{N}$, where $[-l_0, l_0]$ is such that $H(\omega)$ depends only on the configuration there. So if we project $f_N d\mu_N$ onto the configurations in such an interval and expand the interval by a factor of N , we will get a set $\mathcal{B}_{N,B,D,\varepsilon}^0$ of point processes on the interval $|x| \leq \frac{1}{\varepsilon} + l_0$. Theorem 7.4 reduces to proving that

$$\lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\gamma \in \mathcal{B}_{N,B,D,\varepsilon}^0} E^\gamma \left| \frac{\varepsilon}{2} \int_{-1/\varepsilon}^{1/\varepsilon} H(\omega^x) dx - \hat{H}(\rho_\varepsilon(0)) \right| = 0 . \tag{7.7}$$

Let us note that because of translation invariance and blowup by a factor of N , the local density of particles is normalized at 1. Therefore the collection $\mathcal{B}_{N,B,D,\varepsilon}^0$ of point processes on $\left[-\left(\frac{1}{\varepsilon} + l_0\right), \left(\frac{1}{\varepsilon} + l_0\right)\right]$ is compact as $N \rightarrow \infty$ and we denote by $\mathcal{B}_{B,D,\varepsilon}$ the set of limit points. Verifying (7.7) reduces to verifying

$$\lim_{\varepsilon \rightarrow 0} \sup_{\nu \in \mathcal{B}_{B,D,\varepsilon}} E^\nu \left[\left| \frac{\varepsilon}{2} \int_{-1/\varepsilon}^{1/\varepsilon} H(\omega^x) dx - \hat{H}(\rho_\varepsilon(0)) \right| \right] = 0 . \tag{7.8}$$

In the appendix we have introduced the collection of Gibbs measure with boundary condition ω and particle number n on the interval $[-l, l]$. Let us denote the convex hull of these measures with n and ω varying by Γ . The subset with the average particle density less than or equal to 1 will be denoted by $\Gamma_l^{(1)}$. According to Theorem 10.3 of the appendix,

$$\lim_{l \rightarrow \infty} \sup_{\nu \in \Gamma_{l_0}^{(1)}} E^\nu \left[\left| \frac{1}{2l} \int_{-l}^l H(\omega^x) dx - \hat{H}(\rho_l(0)) \right| \right] = 0 ,$$

and we will complete the proof of (7.8) by establishing

Lemma 7.5. For any B, D and $\varepsilon > 0$

$$\mathcal{B}_{B,D,\varepsilon} \subset \Gamma_{1/\varepsilon + l_0}^{(1)} .$$

Proof. Since the density can only drop under the weak limit it is clear that it is sufficient to prove $\mathcal{B}_{B,D,\varepsilon} \subset \Gamma_{l_0 + 1/\varepsilon}$.

Let us assume that V is supported on some $[-c_0, c_0]$ and look at the set of limit points of the point process on a slightly larger interval of the form $\left[-\left(\frac{1}{\varepsilon} + l_0 + c_0\right), \frac{1}{\varepsilon} + l_0 + c_0\right]$ in the expanded scale. Let γ be any limit point. We want to characterize the restriction of ν as a point process on

$\left[-\left(\frac{1}{\varepsilon}+l_0\right),\left(\frac{1}{\varepsilon}+l_0\right)\right]$ which is a member of $\Gamma_{l_0+1/\varepsilon}$. Let us consider a diffusion on the configuration space of point processes on the interval $\left[-\left(\frac{1}{\varepsilon}+l_0+c_0\right),\frac{1}{\varepsilon}+l_0+c_0\right]$ under which the particles in $\left[-\left(\frac{1}{\varepsilon}+l_0\right),\left(\frac{1}{\varepsilon}+l_0\right)\right]$ diffuse with reflecting boundary conditions at the end point and those outside $\left[-\left(\frac{1}{\varepsilon}+l_0\right),\left(\frac{1}{\varepsilon}+l_0\right)\right]$ stay put. The generator is given by

$$L = \frac{1}{2} \sum \frac{\partial^2}{\partial x_i^2} - \sum_{i,j} V'(x_i-x_j) \frac{\partial}{\partial x_i} - \frac{1}{2} \sum_{i,\alpha} V'(x_i-y_\alpha) \frac{\partial}{\partial x_i},$$

where x_1, \dots, x_n are the particles inside and $\{y_\alpha\}, \dots$ are the particles outside. The outside configuration ω and the number n of particles inside do not change and for this diffusion L the family $\mu_{n,l'}^\omega$ with $l' = \left(\frac{1}{\varepsilon}+l_0\right)$ is precisely the set of extremal invariant measures. Therefore all we need to show is that ν is invariant for the diffusion L .

Let us take a function $u(x_1, \dots, x_n, b)$ of the form $u(x_1, \dots, x_n)\phi(b)$ with a smooth u , satisfying Neumann boundary conditions. Strictly speaking $u(x_1, \dots, x_n, b)$ is not a continuous function viewed as a function on configurations in $\left[-\left(\frac{1}{\varepsilon}+l+k_0\right),\frac{1}{\varepsilon}+l+k_0\right]$ because of the distinction between outside and inside variables. Its discontinuity points are on the set where a particle sits exactly on the boundary. These have probability zero because the homogeneity in space makes any local density at most one in the limit. So we can calculate

$$\begin{aligned} \int_{A_n} Ludv &= \lim_{N \rightarrow \infty} \int_{A_n} Ludv_N \\ &= \lim_{N \rightarrow \infty} \int_{A_{n,N}} \hat{L}_N \hat{u}_N f_N d\mu_N, \end{aligned}$$

where $f_N d\mu_N$ is the measure on S^N which produced the ν_N by restricting in the set $\mathcal{B}_{N,B,D,\varepsilon'}^0$ for some choice of ε' . A_n is the set with n points in $\left[-\left(\frac{1}{\varepsilon}+l_0\right),\left[\frac{1}{\varepsilon}+l_0\right]\right]$ and $A_{n,N}$ is the set with n points in $\left[-\frac{\left(\frac{1}{\varepsilon}+l_0\right)}{N},\frac{\left(\frac{1}{\varepsilon}+l_0\right)}{N}\right]$. An elementary calculation using integration by parts yields

$$\int_{A_{n,N}} \hat{L}_N \hat{u}_N f_N d\mu_N = \frac{1}{2N^2} \int_{A_{n,N}} \sum' \frac{\partial \hat{u}_n}{\partial x_i} \cdot \frac{\partial f_N}{\partial x_i} d\mu_N,$$

where \sum' is the summation over those variables x_1, \dots, x_n inside $\left[-\frac{\left(\frac{1}{\varepsilon}+l_0\right)}{N},\frac{\left(\frac{1}{\varepsilon}+l_0\right)}{N}\right]$. The factor $\frac{1}{N^2}$ comes from the stretching. $\frac{\partial \hat{u}_n}{\partial x_i}$ are of

order N and therefore we only have to prove

$$\sup_N \frac{1}{2} \int_{A_{n,N}} \sum' \left(\frac{\partial f_N}{\partial x_i} \right)^2 \cdot \frac{1}{f_N} d\mu_N < \infty .$$

We can calculate by translation invariance

$$\begin{aligned} \int \frac{1}{2} \sum' \left(\frac{\partial f_N}{\partial x_i} \right)^2 \frac{1}{f_N} d\mu_N &= \frac{2 \left(\frac{1}{\varepsilon} + l_0 \right)}{N} \int \frac{1}{2} \sum \left(\frac{\partial f_N}{\partial x_i} \right)^2 \frac{1}{f_N} d\mu_N \\ &\leq \frac{2 \left(\frac{1}{\varepsilon} + l_0 \right)}{N} \cdot ND \\ &= 2 \left(\frac{1}{\varepsilon} + l_0 \right) D , \end{aligned}$$

and we are done.

This proves Theorem 7.2.

We now turn to the proof of Theorem 7.3. We have to carry out some construction. Given any set (x_1, \dots, x_N) of N points in S , we have earlier associated the empirical distribution

$$\alpha_N(dx) = \frac{1}{N} [\delta_{x_1} + \dots + \delta_{x_N}]$$

and a mollified density

$$\rho_\lambda(x) = \frac{\lambda}{N} \sum h(\lambda(x - x_i)) .$$

For most configurations the density $\rho_{\varepsilon N}(x)$ can be highly oscillatory and $\int \hat{H}(\rho_{\varepsilon N}(x)) dx$ and $\int \hat{H}(\rho_\lambda(x)) dx$ for fixed ε and λ can be very far apart. To rule this out we plan to calculate the Young measure associated with the functions $\rho_{\varepsilon N}(\cdot)$ and show that for N large and ε small, these are nearly degenerate. Corresponding to a mollified density $\rho_\lambda(x)$ we can define a probability measure π on $S \times R^+$ by the relation

$$\int F(x, \rho) \pi(dx, d\rho) = \int_S F(x, \rho_\lambda(x)) dx .$$

π clearly enjoys the properties

$$\int \rho \pi(dx, d\rho) = \int \rho_\lambda(x) dx = 1 , \tag{7.9}$$

$$\int F(x) \pi(dx, d\rho) = \int F(x) dx , \tag{7.10}$$

or π projects to Lebesgue measure on S . If we denote by \mathcal{M} the space of probability measures on $S \times R^+$, then through π and ρ_λ we map x_1, \dots, x_N into \mathcal{M} . We already have a map \mathcal{L}_n that maps (x_1, \dots, x_n) into $M_1(S)$ by

$$\alpha_N(dx) = \frac{1}{N} [\delta_{x_1} + \dots + \delta_{x_N}] .$$

We now take both the maps and map (x_1, \dots, x_N) into the pair α_N, π , a point in $M_1(S) \times \mathcal{M}$. With the choice of $\lambda = N\varepsilon$, this produces an induced probability measure $\hat{Q}_{N,\varepsilon}$ on $M_1(S) \times \mathcal{M}$ for each f_N from $\mathcal{A}_{N,B,D}$ and choice of ε . The range of such $\hat{Q}_{N,\varepsilon}$ will be denoted by $\hat{\mathcal{A}}_{N,B,D,\varepsilon}$, their limit points as $N \rightarrow \infty$ by $\hat{\mathcal{A}}_{B,D,\varepsilon}$ and the limit points of these as $\varepsilon \rightarrow 0$ by $\hat{\mathcal{A}}_{B,D}$. Our goal is to prove Theorem 7.3 and the main step is

Theorem 7.6. *Let $\hat{Q} \in \hat{\mathcal{A}}_{B,D}$. Then for almost all (α, π) with respect to \hat{Q} ,*

$$\alpha(dx) = \rho(x)dx$$

and

$$\pi(dx, d\rho) = dx \delta_{\rho(x)}(d\rho)$$

for some function $\rho(x)$ in $L_1(S)$, i.e. the young measures are trivial.

We will reach Theorem 7.6 by a series of lemmas:

Lemma 7.7. *Let $\hat{Q} \in \hat{\mathcal{A}}_{B,D}$. Then \hat{Q} has the property*

$$\hat{Q}[\alpha : \alpha(dx) = \rho(x)dx \text{ for some } \rho] = 1 ,$$

and in fact

$$E^{\hat{Q}} \left[\int_S \rho^2(x) dx \right] \leq C \cdot B ,$$

where C is a universal constant.

Proof. From Lemma 4.2 we can conclude that

$$E^{\hat{Q}} \int_S \lambda W(\lambda(x-y)) \alpha(d\alpha) \alpha(dy) \leq \|W\| CB ,$$

where W is a nonnegative function of integral 1. Since the bound is uniform in λ we can let $\lambda \rightarrow \infty$ and establish Lemma 7.7 by Fatou's lemma.

Lemma 7.8. *For any Q_ε in $\hat{\mathcal{A}}_{B,D,\varepsilon}$,*

$$Q_\varepsilon[(\alpha, \pi) : \int \rho \pi(dx, d\rho) = 1] = 1 , \tag{7.11}$$

$$Q_\varepsilon[(\alpha, \pi) : \pi(dx, d\rho) = dx \pi_x(d\rho)] = 1 , \tag{7.12}$$

$$Q_\varepsilon[(\alpha, \pi) : \alpha(dx) = \rho(x)dx \text{ with } \rho(x) = \int \rho \pi_x(d\rho)] = 1 . \tag{7.13}$$

Proof.

$$\begin{aligned} E^{\hat{Q}_{N,\varepsilon}} \int \rho^2 \pi(dx, d\rho) &= E^{f_N} \int_S \rho_{\varepsilon N}^2(x) dx \\ &= E^{f_N} \int_S \left| \varepsilon \sum h(\varepsilon N(x_i - x)) \right|^2 dx \\ &= \frac{\varepsilon}{N} E^{f_N} \sum W(\varepsilon N(x_i - x_j)) \\ &\leq \|W\| BC \quad (\text{by Lemma 4.2}) , \end{aligned}$$

Now (7.11) is a consequence of the uniform integrability (7.9). Relation (7.12) follows from (7.10). (7.13) is just the routine fact that the mean of the Young measures defines the function which is the weak limit.

We now try to use the bound on the Dirichlet form and use inequality (5.3). The expression for $\mathcal{L}_N U_N$ given by (5.11) can be written as

$$\frac{1}{N^2} \mathcal{L}_N U_N = T_1 - T_2 + T_3 - T_4, \tag{7.14}$$

where

$$T_1 = \frac{\delta}{N} \sum g(N\delta(x_i - x_j)),$$

$$T_2 = \frac{\lambda}{N^2} \sum g(\lambda(x_i - x_j))$$

$$T_3 = \frac{1}{N} \sum_{i,j,k} [G(N\delta(x_k - x_j)) - G(N\delta(x_i - x_j))] V'(N(x_i - x_k)),$$

$$T_4 = \frac{1}{N} \sum_{i,j,k} [G(\lambda(x_k - x_j)) - G(\lambda(x_i - x_j))] V'(N(x_i - x_k)).$$

The inequality (5.3) takes the form

$$E^{f_N} [T_1 + T_3] \leq E^{f_N} [T_2 + T_4] + D^{1/2} (E^{f_N} T_5)^{1/2}, \tag{7.15}$$

where

$$T_5 = \frac{4}{N^3} \sum_i \left| \sum_j G(N\delta(x_i - x_j)) - G(\lambda(x_i - x_j)) \right|^2.$$

Because $|G(N\delta(x_i - x_j)) - G(\lambda(x_i - x_j))|$ is bounded by 1 and $G(N\delta x) - G(\lambda x)$ is supported on some interval of size λ^{-1} we can bound

$$|G(N\delta(x_i - x_j)) - G(\lambda(x_i - x_j))| \leq \chi(\lambda(x_i - x_j)),$$

where $\chi(x)$ is a bounded nonnegative continuous function with compact support. Therefore in (7.15), T_5 can be replaced by

$$T_6 = \frac{4}{N^2} \sum_i \sum_j \chi(\lambda(x_i - x_j)).$$

We next want to chase the inequality (7.15) through subsequences to get a limit point of $\hat{Q}_{N,\epsilon}$ in $\mathcal{A}_{B,D,\epsilon}$ and then again through limits finally to a limit \hat{Q} in $\mathcal{A}_{B,D}$.

We have

$$\begin{aligned} T_1 &= \frac{\delta}{N} \sum g(N\delta(x_i - x_j)) \\ &= \delta \int_S \sum g(N\delta(x_i - x_j)) h(N(x_i - x)) dx \\ &= \delta \int_S H(\omega_N^x) dx, \end{aligned}$$

where

$$H(\omega) = \sum g(\delta(x_i - x_j)) h(x_i).$$

If it were not for the fact that $H(\omega)$ is unbounded we could apply Theorem 7.2

and we would have

$$\lim_{N \rightarrow \infty} E^{fN} T_1 = E^Q \int dx [\int \hat{H}_\delta(\rho) \pi_x(d\rho)]$$

with

$$\begin{aligned} \hat{H}_\delta(\rho) &= E_\rho [\delta \sum g(\delta(x_i - x_j)) h(x_i)] \\ &= \int \delta g(\delta(x - y)) h(x) R_\rho^{(2)}(dx, dy) , \end{aligned}$$

where $R_\rho^{(2)}(dx, dy)$ is the two point correlation function relative to the Gibbs measure with density ρ . However since $H(\omega)$ is nonnegative and continuous Fatou's lemma can be used to conclude

$$E^Q \int_S dx [\int \hat{H}_\delta(\rho) \pi_x(d\rho)] \leq \lim_{N \rightarrow \infty} E^{fN} T_1 . \quad (7.16)$$

We next look at T_3 ,

$$\begin{aligned} T_3 &= \frac{1}{N} \sum_{i,j,k} [G(N\delta(x_k - x_j)) - G(N\delta(x_i - x_j))] V'(N(x_i - x_k)) \\ &= \int_S dx \sum [G(N\delta(x_k - x_j)) - G(N\delta(x_i - x_j))] V'(N(x_i - x_k)) h(N(x_i)) \\ &= \int_S H_1(\omega_N^x) dx , \end{aligned}$$

where

$$H_1(\omega) = \sum [G(\delta(x_k - x_j)) - G(\delta(x_i - x_j))] V'(x_i - x_k) h(x_i) .$$

One can check that because of the monotonicity of G and the fact that $xV'(x) \leq 0$ for all x , $H_1(\omega)$ is nonnegative. One can then obtain by exactly the same reasoning as above

$$E^Q \int_S dx [\int \hat{H}_{1,\delta}(\rho) \pi_x(d\rho)] \leq \lim_{N \rightarrow \infty} E^{fN} T_3 , \quad (7.17)$$

where

$$\hat{H}_{1,\delta}(\rho) = \int \int [G(\delta(z - y)) - G(\delta(x - y))] V'(x - y) h(x) R_\rho^{(3)}(dx, dy, dz) ,$$

where $R_\rho^{(3)}(dx, dy, dz)$ is the three point correlation.

We now turn to T_2 :

$$\begin{aligned} E^Q \int \int \lambda g(\lambda(x - y)) d\alpha(x) d\alpha(y) &= E^Q \int \int \lambda g(\lambda(x - y)) \rho(x) \rho(y) dx dy \\ &= \lim_{N \rightarrow \infty} E^{fN} T_2 . \end{aligned} \quad (7.18)$$

As for T_4

$$T_4 = \frac{1}{N} \sum_{i,j,k} [G(\lambda(x_k - x_j)) - G(\lambda(x_i - x_j))] V'(N(x_i - x_k)) .$$

We know that $V'(N(x_i - x_k)) = 0$ unless $|x_i - x_k| \leq C/N$. Therefore we can expand $G(\lambda(x_k - x_j)) - G(\lambda(x_i - x_j))$ by Taylor's formula with remainder and

$$|G(\lambda(x_k - x_j)) - G(\lambda(x_i - x_j))| \leq \lambda g(\lambda(x_k - x_j))(x_k - x_i) + \frac{C}{N} |x_k - x_i| ,$$

where the error term depends only on λ .

We can now write

$$T_4 = \frac{\lambda}{N^2} \sum g(\lambda(x_i - x_j)) \psi(N(x_i - x_k)) + \text{Error} ,$$

$$E^{fN}[|\text{Error}|] \leq E^{fN} \frac{C}{N^2} \sum_{i,j,k} \psi(N(x_i - x_j))$$

$$\leq \frac{C}{N} \text{ by Lemma 4.2.}$$

We now try to evaluate

$$\lim_{n \rightarrow \infty} E^{fN} \left[\frac{\lambda}{N^2} \sum g(\lambda(x_k - x_j)) \psi(N(x_i - x_k)) \right] .$$

The function $\lambda g(\lambda(x - y))$ on $S \times S$ can be approximated by functions of the form

$$\sum_{r=1}^l a_r(x) b_r(y)$$

uniformly on $S \times S$ and if we denote by

$$\lambda g(\lambda(x - y)) - \sum_{r=1}^l a_r(x) b_r(y) = c(x, y) ,$$

then

$$E^{fN} \left| \frac{1}{N^2} \sum c(x_k, x_j) \psi(N(x_i - x_k)) \right| \leq \frac{1}{N^2} \|C\| N E^{fN} \left[\sum \psi(N(x_i - x_j)) \right] \\ \leq \|C\| \text{ by Lemma 4.2.}$$

So the error is controlled uniformly in N . We can compute

$$\lim_{N \rightarrow \infty} E^{fN} \frac{1}{N^2} \sum a(x_k) b(x_j) \psi(N(x_i - x_k)) \\ = \lim_{N \rightarrow \infty} E^{fN} \left[\frac{1}{N} \sum b(x_j) \cdot \frac{1}{N} \sum a(x_k) \psi(N(x_i - x_k)) \right]$$

$$\left| \frac{1}{N} \sum a(x_k) \psi(N(x_i - x_k)) - \int_S a(y) \sum_{i,k} \psi(N(x_i - x_k)) h(N(x_k - y)) \right|$$

$$\leq \left| \frac{1}{N} \sum \psi(N(x_i - x_k)) \int [a(y) - (x_k)] N h(N(x_k - y)) dy \right|$$

$$\leq \frac{1}{N} \sum \psi(N(x_i - x_k)) \cdot \epsilon_N ,$$

where ε_N depends only on the modulus of continuity of a and goes to zero with N . Again by Lemma 4.2 we can control the error uniformly in N .

Now we can consider

$$\lim_{N \rightarrow \infty} E^{f_N} \left\{ \frac{1}{N} \sum b(x_i) \cdot \int_S a(y) \sum \psi(N(x_i - x_k)) h(N(x_k - y)) dy \right\},$$

and working as before we should end up with

$$\begin{aligned} & E^Q \left\{ \int_S b(x) \rho(x) dx \int_S a(y) \int_R \hat{H}_3(\rho) \pi_y(d\rho) \right\} \\ &= E^Q \left\{ \iint b(x) a(y) \rho(x) \int_R \hat{H}_3(\rho) \pi_y(d\rho) \right\}, \end{aligned}$$

where

$$\hat{H}_3(\rho) = \iint \psi(x - y) h(y) R_\rho^{(2)}(dx, dy).$$

However a serious problem here is that we are operating on the right side of the inequality and cannot afford the luxury of Fatou’s lemma. We have to have an actual identification of limit and therefore have to prove that the functionals

$$H_3(\omega) = \sum \psi(x_i - x_k) h(N(x_k - y))$$

are uniformly integrable and can be truncated to yield uniformly bounded versions with uniformly small errors. What we can use for this is an estimate of the form

$$E^{f_N} \int \left| \sum_{i,k} \psi(N(x_i - x_k)) h(N(x_k - y)) \right|^{4/3} dy \leq C.$$

a and b are uniformly bounded and cause no problems.

Lemma 7.9. *If $f_N \in \mathcal{A}_{N,B,D}$, then*

$$E^{f_N} \int \left| \sum_{i,k} \psi(N(x_i - x_k)) \hat{H}(N(x_k - y)) \right|^{4/3} dy \leq C,$$

where C depends only on B and D .

Proof. Let us denote $\psi(N(x_i - x_k))$ by ψ_{jk} , $h(N(x_k - y))$ by h_k and $\sum_i \psi_{ik}$ by T_k . Then

$$\begin{aligned} \left| \sum_{i,k} \psi_{ik} h_k \right| &= \left| \sum h_k T_k \right| \leq (\sum h_k)^{1/2} (\sum h_k T_k^2)^{1/2} \\ \left| \sum_{ik} \psi_{ik} h_k \right|^{4/3} &\leq (\sum h_k)^{2/3} (\sum h_k T_k^2)^{2/3} \\ E^{f_N} \int \left| \sum_{i,k} \psi_{ik} h_k \right|^{4/3} dy &\leq E^{f_N} \{ (\sum h_k)^{4/3} (\sum h_k T_k^2)^{2/3} \} \\ &\leq (E^{f_N} \int (\sum h_k)^2 dy)^{1/3} (E^{f_N} \int (\sum h_k T_k^2) dy)^{2/3}. \end{aligned}$$

But

$$E^{fN} \int |\sum h(N(x_k - y))|^2 dy = \frac{1}{N} E^{fN} \sum \chi(N(x_i - x_j))$$

is bounded. Moreover

$$E^{fN} \int \sum h(N(x_k - y)) T_k^2 dy = E^{fN} \frac{1}{N} \sum_i \left| \sum_j \psi(N(x_i - x_j)) \right|^2 ,$$

and this was shown to us uniformly bounded before in Lemma 5.1.

Turning to our original estimate now we have

$$\lim_{N \rightarrow \infty} E^{fN} T_4 = E^Q \left[\int \int \lambda g(\lambda(x - y)) \rho(x) dx dy \int \hat{H}_3(\rho) \pi_y(d\rho) \right] . \tag{7.19}$$

We now look at the final term

$$\lim_{N \rightarrow \infty} E^{fN} T_6 = 4E^Q \left[\int \int \chi(\lambda(x - y)) \rho(x) \rho(y) dx dy \right] .$$

Combining (7.15) with (7.16) through (7.19) we obtain for any measure Q in $\mathcal{A}_{b,D}$,

$$\begin{aligned} & E^Q \left[\int_S dx \left[\int \hat{H}_\delta(\rho) \pi_x(d\rho) \right] \right. \\ & \quad \left. + \int_S dx \left[\int \hat{H}_{1,\delta}(\rho) \pi_x(d\rho) \right] \right] \\ & \leq E^Q \left[\int \int \lambda g(\lambda(x - y)) \rho(x) \rho(y) dx dy \right. \\ & \quad \left. + \int \int \lambda g(\lambda(x - y)) \rho(x) dx dy \int \hat{H}_3(\rho) \pi_y(d\rho) \right] \\ & \quad + \sqrt{D} \left[4E^Q \left[\int \int \chi(\lambda(x - y)) \rho(x) \rho(y) dx dy \right] \right]^{1/2} . \tag{7.20} \end{aligned}$$

In the relation (7.20) the right-hand side is clearly bounded uniformly in terms of our constants B and D for each fixed λ . We first let $\delta \rightarrow 0$ in the left. From the factoring properties of the correlation functions $R_\rho^{(2)}(dx, dy)$ and $R_\rho^{(3)}(dx, dy, dz)$ we can conclude by an application of Fatou's lemma

$$\lim_{\delta \rightarrow 0} \int_S dz \int [\hat{H}_\delta(\rho) + \hat{H}_{1,\delta}(\rho)] \pi_x(d\rho) \geq \int_S dx \int \rho P(\rho) \pi_x(d\rho) ,$$

since by Theorem 10.4 $P(\rho) \sim \rho^2$ for large ρ , it follows that

Lemma 7.10. For $Q \in \mathcal{A}_{B,D}$,

$$\sup_{Q \in \mathcal{A}_{B,D}} E^Q \left[\int \rho^3(x) dx \right] \leq E^Q \left[\int_S dx \int \rho^3 \pi_x(d\rho) \right] < \infty .$$

We are interested in letting $\lambda \rightarrow \infty$ on the right-hand side. Let us look at each of the three terms on the right. Clearly $\int \int \lambda g(\lambda(x-y))\rho(x)\rho(y)dx dy \leq \int \rho^2(x)dx$, and since $E^Q \int \rho^2(x)dx < \infty$ we can use the dominated convergence theorem to conclude that

$$\lim_{\lambda \rightarrow \infty} E^Q \int \int \lambda g(\lambda(x-y))\rho(x)\rho(y)dx dy = E^Q \int \rho^2(x)dx . \quad (7.21)$$

The second term involves

$$\int \int \lambda g(\lambda(x-y))\rho(x)dx dy \int \hat{H}_3(\rho)\pi_y(d\rho)$$

and

$$\lim_{\lambda \rightarrow \infty} \int \int \lambda g(\lambda(x-y))\rho(x)dy \int \hat{H}_3(\rho)\pi_y(d\rho) = \int \rho(y) \int \hat{H}_3(\rho)\pi_y(d\rho) .$$

We can bound

$$\begin{aligned} & \int \int \lambda g(x-y)\rho(x)dx dy \int P(\rho)\pi_y(d\rho) \\ & \left[\int \left(\int \lambda g(\lambda(x-y))\rho(x)dx \right)^3 \right]^{1/3} \left[\int \left[\int P(\rho)\pi_y(d\rho) \right]^{3/2} dy \right]^{2/3} \\ & \leq \frac{1}{3} \int \lambda g(\lambda(x-y))\rho^3(x)dx dy + \frac{2}{3} \int [P(\rho)]^{3/2} \pi_y(d\rho) dy \\ & \leq A \text{ by Lemma 7.10.} \end{aligned}$$

The third term clearly goes to zero as $\lambda \rightarrow \infty$. Therefore for every $Q \in \mathcal{A}_{B,D}$,

$$E^Q \left[\int_S dx \int \rho P(\rho)\pi_x(d\rho) \right] \leq E^Q \left[\int_S dx \left[\int \rho \pi_x(d\rho) \int P(\rho)\pi_x(d\rho) \right] \right] . \quad (7.22)$$

Proof of Theorem 7.6. We can think of Eq. (7.22) as

$$E^Q \int_S dx \left\{ \int \rho P(\rho)\pi_x(d\rho) - \int \rho \pi_x(d\rho) \int P(\rho)\pi_x(d\rho) \right\} \leq 0 . \quad (7.23)$$

This means that on the average the covariance

$$\int \rho P(\rho)\pi_x(d\rho) - \int \rho \pi_x(d\rho) \int P(\rho)\pi_x(d\rho)$$

is nonnegative. However

$$\begin{aligned} & \int \rho P(\rho)\pi_x(d\rho) - \int \rho \pi_x(d\rho) \int P(\rho)\pi_x(d\rho) \\ & = \int \left(\rho - \int \rho \pi_x(d\rho) \right) \left(P(\rho) - P \left(\int \rho \pi_x(d\rho) \right) \right) \pi_x(d\rho) \\ & \geq 0 , \end{aligned}$$

because $P(\rho)$ is a nondecreasing function of ρ . The strictly increasing nature of $P(\rho)$ dictates that we have strict positivity unless $\pi_x(d\rho)$ is degenerate. This completes the proof of Theorem 7.6.

Having proved Theorem 7.6 the first application is an estimate on the regularity of densities in the support of any $Q \in \mathcal{A}_{B,D}$.

Theorem 7.11. For any $Q \in \mathcal{A}_{B,D}$,

$$E^Q \left[\int_S \{ [P(\rho(\theta))]_{\theta} \}^2 \frac{1}{\rho(\theta)} \right] \leq 2D .$$

Proof. Let us consider k functions $u_1(x), \dots, u_k(x)$ from $C^\infty(S)$ and l functions $F_1(x), \dots, F_l(x)$ also from $C^\infty(S)$. We consider as well the functions g_1, \dots, g_k from $C_0^\infty(\mathbb{R}^l)$. Let us denote by $\hat{x} = (x_1, \dots, x_N)$ a point of S^N . Then we define on S^N

$$U_n(\hat{x}) = \frac{1}{N} \sum_{j=1}^N u_r(x_j) ,$$

$$G_r(\hat{x}) = g_r \left(\frac{1}{N} \sum_{j=1}^N F_1(x_j), \dots, \frac{1}{N} \sum_{j=1}^N F_l(x_j) \right) .$$

We can perform integration by parts and write

$$\int \left\{ \sum_{r=1}^k [L_N U_r(\hat{x})][G_r(\hat{x})] \right\} f_N(\hat{x}) d\mu_N$$

$$= -\frac{1}{2} \int \left\{ \sum_{r=1}^k \langle \nabla U_r, \nabla G_r \rangle + \sum_{r=1}^k \left\langle U_r, \frac{\nabla f_N}{f_N} \right\rangle G_r \right\} f_N d\mu_N$$

$$= T_1 + T_2 . \tag{7.24}$$

Since $\left| \frac{\partial G_r}{\partial x_j} \right| \leq \frac{C}{N}$ it follows by an elementary calculation that $|T_1| \rightarrow 0$ as $N \rightarrow \infty$.

We can estimate T_2 by Schwartz's inequality to obtain

$$|T_2| \leq \frac{1}{2} \sqrt{2D} \left(\int \frac{1}{N} \sum_{j=1}^N \left(\sum_{r=1}^k u'_r(x_j) G_r(\hat{x}) \right)^2 \cdot f_N d\mu_N \right)^{1/2} .$$

The left-hand side of (7.24) can be calculated to be approximately

$$\frac{1}{2} \int \sum_{r=1}^k \left\{ \frac{1}{N} \sum_{j=1}^N u''_r(x_j) G_r(\hat{x}) \left(1 + \sum_{i=1}^N \psi(N(x_i - x_j)) \right) \right\} f_N d\mu_N .$$

For any $Q \in \mathcal{A}_{B,D}$, we can therefore obtain the inequality

$$\frac{1}{2} \left| E^Q \int \sum_{r=1}^k u''_r(x) G_r(\langle \rho, F_1 \rangle, \dots, \langle \rho, F_l \rangle) P(\rho(x)) dx \right|$$

$$\leq \frac{1}{2} \sqrt{2D} \left[E^Q \int \left| \sum u'_r(x) G_r(\langle \rho, F_1 \rangle, \dots, \langle \rho, F_l \rangle) \right|^2 \phi(x) dx \right]^{1/2} .$$

It is not difficult to pass from this estimate to

$$\left| E^Q \int \psi_{xx}(x, \omega) P(\rho(x, \omega)) dx \right| \leq \sqrt{2D} \left\{ E^Q \int \psi_x^2(x, \omega) \rho(x, \omega) dx \right\} . \tag{7.25}$$

Here ω is the generic point and Q is the measure on random densities $\rho(\cdot)$.

The estimate (7.25) will hold if the test function $\psi(x, \omega)$ is uniformly bounded along with ψ_x and ψ_{xx} . Since we can replace ψ_x by ψ we can assume

$$\left| E^\mathcal{Q} \int \psi_x(x, \omega) P(\rho(x, \omega)) dx \right| \leq \sqrt{2D} \left\{ E^\mathcal{Q} \int \psi^2(x, \omega) \rho(x, \omega) dx \right\}^{1/2} \quad (7.26)$$

for all bounded ψ with ψ_x bounded and having $\int \psi dx \equiv 0$ for almost all ω .

Since $E^\mathcal{Q} \int \rho^2(x, \omega) dx$ has a uniform bound in $\mathcal{A}_{B,D}$ we will have a bound of the form

$$\left| E^\mathcal{Q} \int \psi_x(x, \omega) P(\rho(x, \omega)) dx \right| \leq C \left(E^\mathcal{Q} \int \psi^4(x, \omega) dx \right)^{1/4}, \quad (7.27)$$

provided $\int \psi(x, \omega) dx \equiv 0$. This is enough to yield the estimate

$$E^\mathcal{Q} \int |[P(\rho(x, \omega))]_x|^{4/3} dx \leq C.$$

We can now rewrite (7.26) in the form

$$\left| E^\mathcal{Q} \int \psi(x, \omega) (P(\rho(x, \omega)))_x dx \right| \leq \sqrt{2D} \left\{ E^\mathcal{Q} \int \psi^2(x, \omega) \rho(x, \omega) dx \right\}^{1/2} \quad (7.28)$$

for all $\psi(x, \omega)$ such that $E^\mathcal{Q} \int \psi^4(x, \omega) dx < \infty$ and satisfy $\int \psi(x, \omega) dx \equiv 0$. Let $\psi(x, \omega)$ be any function with $E^\mathcal{Q} \int \psi^4(x, \omega) dx < \infty$. We write $\psi(x, \omega) = \psi_1(x, \omega) + h(\omega)$, where $\psi_1(x, \omega)$ has the property $\int \psi_1(x, \omega) dx \equiv 0$. Then because $(P(\rho(x, \omega)))_x$ is orthogonal to constants,

$$\begin{aligned} & \left| E^\mathcal{Q} \int \psi(x, \omega) P(\rho(x, \omega))_x dx \right| \\ &= \left| E^\mathcal{Q} \int \psi_1(x, \omega) (P(\rho(x, \omega)))_x dx \right| \\ &\leq \sqrt{2D} \left\{ E^\mathcal{Q} \int [\psi(x, \omega) - h(\omega)]^2 P(x, \omega) dx \right\}^{1/2} \\ &\leq \sqrt{2D} \left\{ E^\mathcal{Q} \int [\psi(x, \omega) - h(\omega)]^2 \right. \\ &\quad \left. \cdot [1 + \rho(x, \omega)] dx \right\}^{1/2}. \end{aligned} \quad (7.29)$$

Since $h(\omega) = 0 \int \psi(x, \omega) dx$ we can bound

$$E^\mathcal{Q} h^2(\omega) \leq E^\mathcal{Q} \int \psi^2(x, \omega) [1 + \rho(x, \omega)] dx.$$

Therefore

$$\left| E^\mathcal{Q} \int \psi(x, \omega) (P(\rho(x, \omega)))_x dx \right| \leq C \left[E^\mathcal{Q} \int \psi^2(x, \omega) [1 + \rho(x, \omega)] dx \right]^{1/2}$$

for all functions $\psi(x, \omega)$ with $E^\mathcal{Q} \int \psi^4(x, \omega) dx < \infty$. This is enough to conclude that

$$E^\mathcal{Q} \int \frac{\{P(\rho(x, \omega))_x\}^2}{1 + \rho(x, \omega)} dx < \infty. \quad (7.30)$$

We now go back to (7.29) and use for $\psi(x, \omega)$ the function $\frac{[P(\rho(x, \omega))]_x}{\varepsilon + \rho(x, \omega)}$. (This has to be justified.) Then $h(\omega) = 0$.

We now obtain

$$\begin{aligned} & E^Q \int \frac{[P(\rho(x, \omega))]_x^2}{\varepsilon + \rho(x, \omega)} dx \\ & \leq \sqrt{2D} \left(E^Q \int \frac{(P(\rho(x, \omega)))_x^2}{[\varepsilon + \rho(x, \omega)]^2} \rho(x, \omega) dx \right)^{1/2} \\ & \leq \sqrt{2D} \left(E^Q \int \frac{(P(\rho(x, \omega)))_x^2}{\varepsilon + \rho(x, \omega)} dx \right)^{1/2} . \end{aligned}$$

Because of (7.30) we can let $\varepsilon \rightarrow 0$ and obtain our theorem.

Now we have all the ingredients necessary to prove Theorem 7.3. If we calculate

$$\begin{aligned} & \lim_{\lambda \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{Q \in \mathcal{A}_{N, \varepsilon, D}} E^Q \left[\int |\rho_\lambda(x) - \rho_{N\varepsilon}(x)| dx \right] \\ & = \lim_{\lambda \rightarrow 0} \sup_{Q \in \mathcal{A}_{B, D}} E^Q \left[\left| \int \int h_\lambda(x-y) \rho(y) dy - \rho(x) \right| dx \right] \\ & = 0 \end{aligned}$$

because of the estimate in Theorem 7.11. Because $P(\rho)$ is strictly monotone and continuous in ρ any control on $\int \frac{(P(\rho))_\theta^2}{\rho} d\theta$ will provide control on the L_1 modulus of continuity of $\rho(\cdot)$.

8. Hydrodynamic Limit

Let Q_N be the family of probability measures induced on $C[[0, T]; M_1(S)]$ by looking at the trajectory $\frac{\delta_{x_1(\cdot)} + \dots + \delta_{x_n(\cdot)}}{\varepsilon}$ in the space of probability measures.

We have already established Theorem 3.1 asserting the compactness of Q_N . Let Q be any weak limit point. The results of Sect. 7 establish Theorems 3.2 and 3.3. All we need is the following lemma.

Lemma 8.1. *If we have the bounds*

$$\begin{aligned} & E^Q \int \rho^2(\theta, \omega) d\theta \leq C , \\ & E^Q \int \frac{(P(\rho))_\theta^2}{\rho} d\theta \leq C , \end{aligned}$$

then for some other constant C ,

$$E^Q \int \rho^3(\theta) d\theta \leq C .$$

Proof. Let us consider

$$\begin{aligned}
 S(\rho) &= [1 + P(\rho)]^{3/4}, \\
 (S(\rho))_\theta &= \frac{3}{4}[1 + P(\rho)]^{-1/4}[P(\rho)]_\theta, \\
 \int (S(\rho))_\theta^2 d\theta &\leq \frac{9}{16} \int \frac{(P(\rho))_\theta^2}{[1 + P(\rho)]^{1/2}} d\theta, \\
 &\leq C
 \end{aligned}$$

because of the bound $P(\rho) \geq c\rho^2$ for large ρ . Since $P(\rho) \geq C\rho^2$ we have $S(\rho) \geq C\rho^{3/2}$ for large ρ and we have $S(\rho) \leq C\rho^{3/2}$ with another constant. In other words we can control $\int S(\rho)d\theta$ and $\int (S(\rho))_\theta^2 d\theta$, and hence control $\int S^2(\rho)d\theta$.

Remark. Although the conclusion of the lemma is already implied by Lemma 7.10, it follows from the first two estimates assumed in the lemma.

9. Uniqueness

Let $\rho_0(\theta)$ be a function on $L_2(S)$. Assume $\rho(t, \theta)$ is a weak solution of

$$\frac{\partial \rho(t, \theta)}{\partial t} = \frac{1}{2} (P(\rho(t, \theta)))_{\theta\theta}$$

with

$$\rho(0, \theta) = \rho_0(\theta) .$$

Assume that the solution $\rho(t, \theta)$ has the property

$$\int_s^T \int_0^\infty \rho^3(t, \theta) d\theta dt < \infty \tag{9.1}$$

and

$$\int_s^T \int_0^\infty \frac{[P(\rho(t, \theta))]_\theta^2}{\rho(t, \theta)} dt d\theta < \infty . \tag{9.2}$$

Theorem 9.1. *A weak solution satisfying (9.1) and (9.2) is unique.*

Let if possible $u(t, \theta)$ and $v(t, \theta)$ be two solutions corresponding to the same initial data $\rho_0(\theta)$. Let us define the function

$$\omega(\theta, t) = \frac{1}{2} \int_0^t \{ [P(u(s, \theta))]_\theta - [P(v(s, \theta))]_\theta \} ds . \tag{9.3}$$

Using the fact that $\int_0^t \int_0^\infty \rho(s, \theta) ds d\theta < \infty$, it follows from (9.2) that

$$[P(u)]_\theta - [P(v)]_\theta \in L_1(S \times [0, t]) .$$

Hence $\omega(\theta, t)$ is well defined as a continuous function of t for almost all θ . An elementary computation yields

$$\frac{\partial \omega}{\partial \theta} = u - v \tag{9.4}$$

in the sense of distributions. Now

$$\begin{aligned} \frac{\partial}{\partial t} [\omega(\theta, t)]^2 &= 2\omega(\theta, t) \frac{\partial \omega}{\partial t} \\ &= \omega(\theta, t)[P(u) - P(v)]_\theta . \end{aligned}$$

We want to show that

$$\int \int_0^t |\omega(\theta, t)| \{ |[P(u)]_\theta| + |[P(v)]_\theta \} d\theta dt < \infty . \tag{9.5}$$

Since $\int \int \frac{|[P(u)]_\theta|^2}{u} d\theta dt$ and $\int \int \frac{|[P(v)]_\theta|^2}{v} d\theta dt$ are finite, it is sufficient to prove that

$$\int \int_0^t |\omega(\theta, t)|^2 u(\theta, t) d\theta dt$$

and

$$\int \int_0^t |\omega(\theta, t)|^2 v(\theta, t) d\theta dt$$

are finite. (9.1) and (9.4) are sufficient now to conclude (9.5). We therefore conclude that

$$\chi(\theta, t) = [\omega(\theta, t)]^2$$

is a continuously differentiable map of $[0, t]$ into $L_1(S)$. Moreover $\omega(\theta, t) \rightarrow 0$ a.e. θ as $t \rightarrow 0$,

$$\begin{aligned} \int \frac{\partial \chi}{\partial t} d\theta &= \int \omega(\theta, t)[P(u(\theta, t)) - P(v(\theta, t))]_\theta d\theta \\ &= - \int \left(\frac{\partial \omega}{\partial \theta} \right) (P(u) - P(v)) d\theta \\ &= - \int (u - v)(P(u) - P(v)) d\theta \leq 0 . \end{aligned}$$

Therefore $\int \chi(\theta, t) d\theta \equiv 0$. This proves uniqueness.

10. Appendix

In this section we will establish some results concerning finite and infinite volume Gibbs measures for our continuous system with finite range interaction. Although these results are not new and are part of the folklore in the field, there does not appear to be any source that we can refer to for a precise statement of the results in the form we need them. We will therefore provide a proof.

Our main assumption is that we have a repulsive pair potential, i.e. a function $V(x)$, which is nonnegative, even, continuously differentiable and has compact support. We denote by $[-c_0, c_0]$ the interval of support of the function $V(\cdot)$ and we assume further that $\psi(x) = -xV'(x) \geq 0$ and that $V(0) > 0$.

For any $l > 0$, we consider the exterior $[-l, l]^c$ of the interval $[-l, l]$ relative to the real line. A configuration ω is the realization of a point process and

represents a locally finite collection of points. Since our interaction has range c_0 we are only interested in the part of the realization that has to do with the border $[-l - c_0, -l] \cup [l, l + c_0]$ and therefore we can think of ω as an arbitrary but finite configuration of points in the border.

We shall think of $\omega = \{y_\alpha\}$, as a finite set of points located in $[-l - c_0, -l] \cup [l, l + c_0]$. The canonical partition function $Z(n, l, \omega)$ is defined as follows for $n \geq 0, l > 0$ and all ω ,

$$Z(n, l, \omega) = 1 \quad \text{if } n = 0 \quad \text{for all } l \text{ and } \omega, \tag{10.1}$$

$$Z(n, l, \omega) = \frac{1}{n!} \int_{-l}^l \dots \int_{-l}^l \exp \left[- \sum_{i \neq j=1}^n V(x_i - x_j) - 2 \sum_{i, \alpha} V(x_i - y_\alpha) \right] dx_1 \dots dx_n. \tag{10.2}$$

The canonical Gibbs measure $\mu_{n, l, \omega}$ is a point process on $[-l, l]$ with exactly n points or equivalently a measure on $[-l, l]^n$ with a density given by

$$f_{n, l, \omega}(x_1, \dots, x_n) = c_n \exp \left[- \sum_{i \neq j=1}^n V(x_i - x_j) - 2 \sum_{i, \alpha} V(x_i - y_\alpha) \right].$$

The normalizing constant c_n is of course given by

$$c_n^{-1} = n! Z(n, l, \omega).$$

When ω is the empty set we denote $Z(n, l, \omega)$ by $Z_{n, l}^0$ and $\mu_{n, l, \omega}$ by $\mu_{n, l}^0$.

The grand canonical partition function with activity λ is defined for every real λ and

$$\hat{Z}(\lambda, l, \omega) = \sum_{n=0}^{\infty} e^{\lambda n} Z(n, l, \omega),$$

and the grand canonical Gibbs measure is a point process $\hat{\mu}_{\lambda, l, \omega}$ on $[-l, l]$ which is thought of as a probability measure on

$$\bigcup_{n=0}^{\infty} [-l, l]^n.$$

It is given by

$$\hat{\mu}_{\lambda, l, \omega} = [\hat{Z}(\lambda, l, \omega)]^{-1} \sum e^{\lambda n} Z(n, l, \omega) \mu_{n, l, \omega}, \tag{10.3}$$

i.e. it is a convex combination of canonical Gibbs measures and the weights are proportional to $e^{\lambda n} Z(n, l, \omega)$.

Although we have only defined these Gibbs measures on intervals of the form $[-l, l]$ it is clear that they enjoy translation invariance of sorts and can be defined on any interval $[a, b]$ and depend only on the configuration in the appropriate border. Moreover, there is an internal consistency in the following two senses.

First, by definition, the canonical Gibbs measure is the conditional distribution of the grand canonical Gibbs measure when it is conditioned by the number of particles in the set.

Moreover if we take the grand canonical Gibbs distribution on $[-l, l]$ and condition it with respect to the configuration in $[a, b]^c$, where $-l \leq a < b \leq l$, then

the conditional distribution is the grand canonical Gibbs measure on $[a, b]$ with the same activity λ .

The infinite volume Gibbs measure μ_λ corresponding to activity λ , is defined as a stationary point process such that for every interval $[a, b]$ the conditional distribution of the configuration in $[a, b]$ given the configuration in $[a, b]^c$ is the grand canonical Gibbs measure $\hat{\mu}_{\lambda, l, \omega}$ with $l = \frac{1}{2}(b - a)$ and ω representing the exterior configuration. We shall state the main theorems we will use in our article and provide a sketch of the proof at the end, after establishing a few key estimates as lemmas.

Theorem 10.1. *The following thermodynamic functions exists:*

$$F(\lambda) = \lim_{l \rightarrow \infty} \frac{1}{2l} \log Z_{\lambda, l}^0 \quad \text{for } \lambda \text{ real ,}$$

$$\Psi(\rho) = \lim_{\substack{l \rightarrow \infty \\ n \rightarrow \infty \\ n/2k \rightarrow \rho}} \frac{1}{2l} \log Z_{n, k}^0 \quad \text{for } \rho \geq 0 .$$

The function $F(\lambda)$ is convex in λ and the function $\Psi(\rho)$ is concave in ρ . Moreover they are related by the equation

$$F(\lambda) = \sup_{\rho} [\lambda \rho + \Psi(\rho)] . \tag{10.4}$$

Theorem 10.2. *For each value of the activity parameter λ , there exists exactly one Gibbs measure μ_λ with activity λ . The point process μ_λ depends continuously on the parameter λ . As a point process its density is given by*

$$\rho(\lambda) = F'(\lambda) = \frac{dF(\lambda)}{d\lambda} .$$

In particular $F(\lambda)$ is once continuously differentiable. The function $\rho(\lambda)$ is continuous and nondecreasing. Its inverse is

$$\lambda = -\Psi'(\rho) = \frac{d\Psi(\rho)}{d\rho} .$$

We have in addition

$$\lim_{\lambda \rightarrow \infty} \rho(\lambda) = \infty , \quad \lim_{\lambda \rightarrow -\infty} \rho(\lambda) = 0 .$$

If I is a unit interval and $N(I)$ denotes the number of points in the interval I , then

$$E^{\mu_\lambda}[\{N(I)\}^r] < \infty \quad \text{for every } r \geq 1 .$$

Theorem 10.3. *Let $n \rightarrow \infty$ and $l \rightarrow \infty$ in such a manner that $\frac{n}{l} \rightarrow \rho$. Let $H(\omega)$ be a bounded continuous local function depending on the configuration in some fixed finite interval $[-l_0, l_0]$. Let us denote by $H_x(\omega)$ the same functional evaluated at the configuration in $[-l_0 + x, l_0 + x]$. We denote by $\hat{H}(\lambda)$ the expected value $E^{\mu_\lambda}[H(\omega)]$. Then for every $\varepsilon > 0$,*

$$\lim_{n, l} \mu_{n, l}^0 \left[\left| \frac{1}{2l} \int_{-(l-l_0)}^{(l-l_0)} H_x(\omega) dx - \hat{H}(\lambda) \right| \geq \varepsilon \right] = 0 , \tag{10.5}$$

where $\rho = F'(\lambda)$.

Moreover for every function $H(x)$ on $[-1, 1]$ which is nonnegative, has compact support with $\int \phi(x)dx = 1$,

$$\lim_{n,l} \mu_{n,l}^0 \left[\left| \frac{1}{l} \sum \phi\left(\frac{x_i}{l}\right) - \rho \right| \geq \varepsilon \right] = 0 . \tag{10.6}$$

Remark 1. It follows from (10.5) that if λ_1, λ_2 are two different activities such that $F'(\lambda_1) = F'(\lambda_2) = \rho$, then $\hat{H}(\lambda_1) = \hat{H}(\lambda_2)$ for all functions $H(\omega)$. This in turn implies $\mu_1 = \mu_{\lambda_1}$, and $\lambda_1 = \lambda_2$. Therefore $\rho(\lambda)$ is a strictly increasing function of λ and its inverse $-\Psi'(\rho)$ is a strictly increasing continuous function of ρ .

We define the function

$$P(\rho) = F(\lambda(\rho)) . \tag{10.7}$$

Theorem 10.4. *There exists constants $0 < c_1 < c_2 < \infty$ such that*

$$c_1 \rho^2 \leq P(\rho) \leq c_2 \rho^2 \text{ for } \rho \text{ large .}$$

Since $E^{\mu_\lambda}[\{N(I)\}'] < \infty$ for every μ_λ , we consider for every function $f(x_1, \dots, x_k)$ of k -variables which is smooth and has compact support on R^k , the variable

$$\xi_f = \sum f(x_1, \dots, x_k) ,$$

where the summation is over all k -tuples of k distinct points in the configuration. ξ_f has a finite expectation and

$$E^{\mu_\lambda} \xi_f = \int f(x_1, \dots, x_k) R_k^\lambda(dx_1, \dots, dx_k) .$$

The measure $R_k^\lambda(dx_1, \dots, dx_k)$ is the k -point correlation measure. It has translation invariance relative to $x_1, \dots, x_k \rightarrow x_1 + a, \dots, x_k + a$. The one point measure $R_k^\lambda(dx_1) = \rho(\lambda)dx_1$, where $\rho(\lambda)$ is the density.

We need the following identity between $P(\rho)$ and $R_2^\lambda(dx_1, dx_2)$.

Theorem 10.5. *If λ and ρ are related by $\lambda = \lambda(\rho)$, then*

$$P(\rho) = \rho + \int \psi(x_1 - x_2) \phi(x_2) R_1^\lambda(dx_1, dx_2) , \tag{10.8}$$

where $\phi(x)$ is any function on R that is smooth, has compact support, is nonnegative and has $\int \phi(x)dx = 1$.

Finally the ergodicity of μ^λ implies some asymptotic factorization of the correlation measures.

Theorem 10.6. *Let g_1, g_2 be functions on R that are smooth and have compact support. Let g_3 be a smooth function on R^2 having compact support. Then for any λ ,*

$$\lim_{k \rightarrow 0} k \int g_1(k(x_1 - x_2)) g_2(x_2) R_2^\lambda(dx_1, dx_2) = \rho^2 \left(\int g_1(x) dx \right) \left(\int g_2(x) dx \right) , \tag{10.9}$$

$$\begin{aligned} & \lim_{k \rightarrow 0} k \int g(k(x_1 - x_2)) g_2(x_2, x_3) R_3^\lambda(dx_1, dx_2, dx_3) \\ & = \rho \left(\int g_1(x) dx \right) \left(\int g_2(x_2, x_3) R_2^\lambda(dx_1, dx_2) \right) . \end{aligned} \tag{10.10}$$

We adopt the following conventions regarding constants. Constants will always be independent of n, l and ω . A, B, C will denote constants and will be

used at different times of denote different constants. We will use γ to denote constants that are strictly positive. The constant c_0 is the range of the interaction $V(\cdot)$. The quantity $\frac{n}{2l}$ will be denoted by ρ .

Lemma 10.7. *For every $l > c_0$ and ω we have*

$$Z_{n,l-c_0}^0 \leq Z(n, l, \omega) \leq Z_{n,l}^0 . \tag{10.11}$$

Proof. For $n=0$, there is nothing to prove. The upper bound is obtained by dropping the interaction terms involving the points in the border. For the lower bound we limit the integration to the situation where all the points x_1, \dots, x_n lie inside $[-(l-c_0), (l-c_0)]$.

Lemma 10.8. *There is a constant C such that*

$$\frac{d \log Z_{n,l}^0}{dl} \leq \frac{n}{l} + C \frac{n^2}{l^2} . \tag{10.12}$$

Proof. By differentiation we can write

$$\frac{d \log Z_{n,l}^0}{dl} = \frac{n}{l} + \frac{1}{l} \int \sum \psi(x_i - x_j) d\mu_{n,l}^0 .$$

[Change variables $x_i = ly_i$, differentiate and change back.] From formula (4.4) we have an estimate

$$\sum \psi(x_i - x_j) \leq C_1 \sum V(x_i - x_j)$$

for some C_1 . Since

$$d\mu_{n,l}^0 = c_n \exp[-\sum V(x_i - x_j)] d\theta ,$$

where $d\theta$ is volume on $[-l, l]^N$, it follows that

$$\int \sum V(x_i - x_j) d\mu_{n,l}^0 \leq \int \sum V(x_i - x_j) d\theta ,$$

where $d\theta$ is normalized volume. We can now calculate $\int \sum V(x_i - x_j) d\theta$ and obtain (10.12).

Lemma 10.9. *For every n and $l > 0$,*

$$z_{n+1,l}^0 \leq \frac{2l}{n+1} Z_{n,l}^0 . \tag{10.13}$$

Proof. If we ignore any interaction involving x_{n+1} and integrate out x_{n+1} we obtain (10.13).

Lemma 10.10. *Let $l = l_1 + l_2 + \frac{1}{2}c_0$ with $l_1 > 0, l_2 > 0$. Then*

$$Z_{n,l}^0 \geq \sum_{\substack{n_1 + n_2 = n \\ n_1 \geq 0, n_2 \geq 0}} Z_{n_1,l_1}^0 Z_{n_2,l_2}^0 .$$

In particular for any n_1, n_2 with $n_1 + n_2 = n$,

$$Z_{n,l}^0 \geq Z_{n_1,l_1}^0 Z_{n_2,l_2}^0 . \tag{10.14}$$

Proof. Let us divide the interval $[-l, l]$ into three parts I_1, I and I_2 with lengths $2l_1, c_0$ and $2l_2$ arranged in order. We limit the integration to the domain where I_1 and I_2 contain all the points (x_1, \dots, x_n) . Here n_1 and n_2 represent the number of particles in I_1 and I_2 respectively. The combinatorial factors adjust themselves and we get Lemma 10.10.

Lemma 10.11. *There exist constants A and B such that*

$$\mu_{n,l,\omega} \leq A e^{B\rho^2} \mu_{n,l}^0, \quad \text{provided } l \geq 2c_0 .$$

Proof. The density $f_{n,l,\omega}$ would be dominated by $f_{n,l}^0$ if it were not for the normalizing constant. We therefore need only estimate the ratio

$$Z_{n,l}^0 [Z(n, l, \omega)]^{-1} .$$

Lemmas 10.7 and 10.8 will establish a bound for this if we can keep l away from c_0 .

Lemma 10.12. *There are positive constants γ, c_1 and c_2 such that*

$$\frac{(2l)^n}{n!} \exp \left[-c_2 \frac{n^2}{l} \right] \leq Z_{n,l}^0 \leq \frac{(2l)^n}{n!} \exp \left[-\gamma \frac{n^2}{l} + c_1 n \right] . \quad (10.15)$$

Proof.

$$\frac{n! Z_{n,l}^0}{(2l)^n} = \frac{1}{(2l)^n} \int_{-l}^l \dots \int_{-l}^l \exp \left[-\sum_{i \neq j=1}^n V(x_i - x_j) \right] dx_1 \dots dx_n .$$

The lower bound follows from Jensen's inequality. On the other hand there are constants $\gamma > 0$ and c_1 such that

$$\sum_{i \neq j=1}^n V(x_i - x_j) \geq \gamma \frac{n^2}{l} - c_1 n$$

for every configuration (x_1, \dots, x_n) .

Lemma 10.13. *Let $I \subset [-l, l]$ be an arbitrary interval of length 1. Let E_k denote the event that the set I contains exactly k of the points from (x_1, \dots, x_n) . Then there are constants A, B, C and $\nu > 0$ such that*

$$\mu_{n,l}^0(E_k) \leq A \exp[-\nu k^2 + Bk\rho + c\rho^2] \quad (10.16)$$

for all n and $l > \frac{1}{2}$.

Proof. Let $i = [a, a + 1]$ for some $-l \leq a \leq l - 1$. Let $I_1 = [-l, a]$ and $I_2 = [a + 1, l]$. We let $l_1 = \frac{l+a}{2}$ and $l_2 = \frac{1}{2}(l - a - 1)$. Suppose that I_1, I_2 contain n_1, n_2 points respectively and $n_1 + n_2 + k = n$. If we ignore all interactions between points in different intervals and use the bound

$$\sum_{\substack{x_i, x_j \in I \\ i \neq j}} V(x_i - x_j) \geq \nu_1 k^2 - c_1 k ,$$

we obtain

$$\mu_{n,l}^0(E_k) \leq \frac{1}{k!} \exp[-\nu_1 k^2 + c_1 k] \cdot \frac{1}{Z_{n,l}^0} \sum_{n_1 + n_2 = n - k} Z_{n_1, l_1}^0 Z_{n_2, l_2}^0 . \quad (10.17)$$

From Lemma 10.10,

$$\begin{aligned} \sum_{n_1+n_2=n-k} Z_{n_1,l_1}^0 Z_{n_2,l_2}^0 &\leq Z_{n-k,l_1+l_2+c_0/2}^0 \\ &= Z_{n-k,l-(1/2)+(c_0/2)}^0 . \end{aligned}$$

We can now apply (10.14) to get

$$Z_{n,l+c_0+(kl/n)}^0 \geq Z_{n-k,l-(1/2)+(c_0/2)}^0 \cdot Z_{k,(1/2)+(kl/n)}^0 .$$

(10.17) then becomes

$$\mu_{n,l}^0(E_k) \leq \frac{1}{k!} \exp[-\nu_1 k^2 + c_1 k] \frac{Z_{n,l+c_0+(kl/n)}^0}{Z_{n,l}^0} \cdot \frac{1}{Z_{k,(1/2)+(kl/n)}^0} . \quad (10.18)$$

We can use bounds from Lemma 10.8 and the lower bound of Lemma 10.12 to complete the proof.

Lemma 10.14. *If N_I is the number of particles in the interval I then for some $\nu_1 > 0$ and $l > \frac{1}{2}$,*

$$E^{\mu_{n,l}^0} [\exp \{ \nu_1 N_I^2 \}] \leq A \exp [B \rho^2] ,$$

where A and B are some constants.

Proof. We pick $\nu_1 < \nu$ and sum

$$\sum \mu_{n,l}^0(E_k) e^{\nu_1 k^2} ,$$

using the bounds in Lemma 10.13.

Lemma 10.15. *For every m , there exists a constant c_m such that*

$$E^{\mu_{n,l}^0} [N_I^m] \leq C_m (1 + \rho)^m ,$$

for all n and l .

Proof. For any random variable X there is a constant c_m independent of X such that

$$E |X|^m \leq c_m (1 + \log E e^{X^2})^{m/2} .$$

If we now combine Lemma 10.11 with Lemmas 10.14 and 10.15 we get

Lemma 10.16. *For all $l > \frac{1}{2}$ and n ,*

$$\sup_{\omega} E^{\mu_{n,l}^{\omega}} [\exp \{ \nu N_I^2 \}] \leq A \exp [B \rho^2] , \quad (10.19)$$

$$\sup_{\omega} E^{\mu_{n,l}^{\omega}} [[N_I]^m] \leq C_m (1 + \rho)^m . \quad (10.20)$$

We now turn to the grand canonical Gibbs distribution and estimates on them.

Lemma 10.17. *Let $\hat{\mu}_{\lambda,l,\omega}$ be the grand canonical Gibbs measure. Then for any given $\lambda_0 > 0$ for all $\lambda \leq \lambda_0$, $l > \frac{1}{2}$ and ω we have the following bounds:*

$$E^{\hat{\mu}_{\lambda,l,\omega}} [\exp \{ \nu N_I^2 \}] \leq A \exp [B (1 + \lambda_0)^2] , \quad (10.21)$$

$$E^{\hat{\mu}_{\lambda,l,\omega}} [[N_I]^m] \leq C_m (1 + \lambda_0)^m . \quad (10.22)$$

Proof. According to Lemma 10.17 there exists $\alpha > 0$ such that

$$E^{\mu_{n,l}}[\exp[\alpha N_l^2]] \leq A \exp[B\rho^2] .$$

We can replace α by $\varepsilon\alpha$ and then B can be replaced by $\varepsilon\beta$ by Hölder's inequality.

We pick ε so small that εB is less than $\frac{\nu}{2}$, where ν is as in Lemma 10.13. We will continue to denote $\varepsilon\alpha$ by α . We now have

$$E^{\mu_{n,l,\omega}}[\exp[\alpha N_l^2]] \leq A \exp\left[\frac{\nu}{2}\rho^2\right] .$$

To estimate $E^{\hat{\mu}_{\lambda,l,\omega}}[\exp[N_l^2]]$ we must bound

$$\begin{aligned} \frac{A}{\hat{Z}(\lambda, l, \omega)} \sum e^{\lambda n} Z(n, l, \omega) \exp\left[\frac{\nu}{2}\rho^2\right] &= T , \\ T^{2l} &\leq \frac{a^{2l}}{\hat{Z}(\lambda, l, \omega)} \sum e^{\lambda n} Z(n, l, \omega) \exp\left[\frac{\nu}{2}(2l)\rho^2\right] . \end{aligned}$$

We use the trivial bound $\hat{Z}(\lambda, l, \omega) \geq 1$. Then

$$T^{2l} \leq A^{2l} \sum_{n=0}^{\infty} e^{\lambda n} \frac{(2l)^n}{n!} e^{-\nu(n^2/l) + cn + (r/4) \cdot (n^2/l)} .$$

One can estimate $\frac{(2l)^n}{n!}$ by e^{2l} and obtain (10.21) by direct estimation of the sum (more or less Gaussian sum). The inequality (10.22) follows from (10.21). The fact that the summation is over nonnegative n makes the estimates uniform for λ in $(-\infty, \lambda_0]$.

Now we turn our attention to a proof of our theorems.

Proof of Theorem 10.1. The existence of the free energy $F(\lambda)$ and the specific energy or entropy functional $\Psi(\rho)$ follows from standard subadditivity arguments. Lemma 10.10 provides the required subadditivity as well as the concavity. The formula for \hat{Z} in terms of Z can be used with a standard Laplace asymptotic formula to calculate the free energy in terms of the function $\Psi(\rho)$ i.e. to prove formula (10.4).

Proof of Theorem 10.2. For any real λ , we consider the family $\hat{\mu}_{\lambda,l}^0$ of grand canonical Gibbs measures as point processes on $[-l, l]$. We let $l \rightarrow \infty$. Lemma 10.17 allows us to take a subsequence and prove the existence of a Gibbs measure which may perhaps be nonstationary. A standard averaging argument over translations and another limit produces a stationary candidate.

For uniqueness let P_1, P_2 be two grand canonical Gibbs measures for the same activity λ . Let $[-a, a]$ be any arbitrary finite interval and we denote by $\|P_1 - P_2\|_a$ the variation norm on the σ field configurations on $[-a, a]$.

We have to show that $\|P_1 - P_2\|_a = 0$ for every a . Let us take $J_k = [x : (k-1)c_0 \leq |x| \leq kc_0]$ and define the conditional expectation operator,

$$(T_k F)(\omega) = \int F(\omega') \hat{\mu}_{\lambda, kc_0, \omega}(d\omega') ,$$

taking functions on the configuration space of J_k into those of J_{k+1} . Proving

uniqueness can easily be reduced to proving

$$||| T_k F ||| \leq \theta ||| F |||$$

for some $\theta < 1$ for all k and F . Here

$$\begin{aligned} ||| F ||| &= \frac{1}{2} \sup_{\omega_1, \omega_2} |F(\omega_1) - F(\omega_2)| \\ &= \inf_c \sup_{\omega} |F(\omega) - c| \end{aligned}$$

where the infimum is taken over constants. This however needs only the following lemmas.

Lemma 10.18. *Let E_k be the event that the interval J_k contains no points of the configuration. Then there is a $\delta > 0$ depending only on λ such that*

$$\hat{\mu}_{\lambda, kc_0, \omega}(E_k) \geq \delta$$

for all $k \geq 1$ and ω .

Proof. All we need is to estimate the ratio

$$\begin{aligned} \tau &= \frac{1}{\hat{Z}(\lambda, kc_0, \omega)} \left(\sum_{n=0}^{\infty} e^{\lambda n} Z(n, kc_0, \omega) \mu_{n, kc_0, \omega}(E_k) \right) \\ \tau &\geq \frac{1}{\hat{Z}(\lambda, kc_0, \omega)} \sum_{n=0}^{\infty} e^{\lambda n} Z_{n, (k-1)c_0}^0 \\ &\geq \frac{\sum_{n=0}^{\infty} e^{\lambda n} Z_{n, (k-1)c_0}^0}{\sum_{n=0}^{\infty} e^{\lambda n} Z_{n, kc_0}^0} \end{aligned}$$

We use the estimate

$$\frac{Z_{n, (k-1)c_0}^0}{Z_{n, kc_0}^0} \geq e^{-A\rho^2 - B}$$

from Lemma 10.8 and we write

$$\begin{aligned} \tau &\geq \frac{\sum_{n=0}^{\infty} e^{\lambda n} Z_{n, kc_0}^0 e^{-A(n/2)^2}}{\sum_{n=0}^{\infty} e^{\lambda n} Z_{n, kc_0}^0} \cdot e^{-B} \\ &\geq e^{-B} e^{-AR^2} \left(1 - \sum_{n \geq 2lR} e^{\lambda n} Z_{n, kc_0}^0 e^{-A(n/2)^2} \right) \end{aligned}$$

We have used $\sum_{n=0}^{\infty} e^{\lambda n} Z_{n, kc_0}^0 \geq 1$.

Now we use the estimates of Lemma 10.12 to complete the proof.

Continuous dependence is an easy consequence of uniform bounds and uniqueness.

Uniqueness of the Gibbs measure yields the differentiability of $F(\lambda)$ and the continuity of $\rho(\lambda)$. The bounds on the moments of $N(l)$ are contained in Lemma 10.17. The relation between $F'(\lambda)$ and $\Psi'(\rho)$ is a consequence of the Legendre transform (10.4). To see the limits of $\rho(\lambda)$ as $\lambda \rightarrow \pm \infty$, the behavior as $\lambda \rightarrow -\infty$ will follow from the easily established relation $\lim_{\lambda \rightarrow -\infty} F(\lambda) = 0$. The behavior for $\lambda \rightarrow \infty$ is obtained by proving $c_1 \lambda^2 \leq F(\lambda) \leq c_2 \lambda^2$ as $\lambda \rightarrow \infty$. These are deduced from Lemma 10.12.

Proof of Theorem 10.3. This theorem, known often as the theorem on the equivalence of ensembles, is an easy consequence of large deviation theory. We note first that the uniqueness of Gibbs measures implies that each μ_λ is ergodic, and in fact, for nice local functions $H(\omega)$,

$$\overline{\lim}_{l \rightarrow \infty} \frac{1}{2l} \log \mu_\lambda \left\{ \left| \frac{1}{2l} \int_{-l}^l H_x(\omega) dx - \hat{H}(\lambda) \right| \geq \varepsilon \right\} \leq -\eta(\varepsilon) < 0 . \quad (10.23)$$

If E_l is the event that the set $[-l-1, -l] \cup [l, l+1]$ has no points in it, then from Lemma 10.18 it follows easily that

$$\mu_\lambda[E_l] \geq \delta > 0 \quad \text{uniformly as } l \rightarrow \infty . \quad (10.24)$$

Combining (10.23) and (10.24)

$$\overline{\lim}_{l \rightarrow \infty} \frac{1}{2l} \log \hat{\mu}_{\lambda,l}^0 \left\{ \left| \frac{1}{2l} \int_{-l}^l H_x(\omega) dx - \gamma ZH(\lambda) \right| \geq \varepsilon \right\} \leq -\eta(\varepsilon) < 0 . \quad (10.25)$$

One has a similar result for the number of points $N(l)$ in the interval $[-l, l]$,

$$\overline{\lim}_{l \rightarrow \infty} \frac{1}{2l} \log \mu_{\lambda,l}^0 \left\{ \left| \frac{N(l)}{2l} - \rho(\lambda) \right| \geq \varepsilon \right\} \leq -\eta(\varepsilon) < 0 . \quad (10.26)$$

In particular

$$\overline{\lim}_{l \rightarrow \infty} \mu_{\lambda,l}^0 \left\{ \left| \frac{N(l)}{2l} - \rho(\lambda) \right| \leq \varepsilon \right\} = 1 . \quad (10.27)$$

If we look at the ratios

$$\frac{\mu_{\lambda,l}^0[N(l) = k + 1]}{\mu_{\lambda,l}^0[N(l) = k]} = \frac{Z_{k+1,l}^0}{Z_{k,l}^0}$$

in the range where $\left| \frac{k}{2l} - \rho(\lambda) \right| \leq \varepsilon$, then Lemma 10.9 provides an upper bound for the ratio and Lemmas 10.10 and 10.8 provide a lower bound for it. Since $\rho(\lambda) > 0$ we can make the bounds uniform over k in the range we are interested in. In turn this means

$$\mu_{\lambda,l}^0 \left[\left| \frac{N(l)}{2l} - \rho(\lambda) \right| \leq \varepsilon \right] \leq e^{c_l \varepsilon} \cdot 2l \varepsilon \cdot \inf_{\left| \frac{k}{2l} - \rho(\lambda) \right| \leq \varepsilon} \mu_{\lambda,l}^0[N(l) = k] . \quad (10.28)$$

It follows now from (10.28) that

$$\lim_{\substack{l \rightarrow \infty \\ k \rightarrow \infty \\ k/2l \rightarrow \rho(\lambda)}} \frac{1}{2l} \log \mu_{\lambda, l}^0 [N(l) = k] = 0 \quad (10.29)$$

If we combine (10.23) with (10.29) we obtain the first half of Theorem 10.3. The second half is similar. We start with the estimate from large deviations.

$$\overline{\lim}_{l \rightarrow \infty} \frac{1}{2l} \log \hat{\mu}_\lambda \left\{ \left| \sum \phi(lx_i) - \rho(\lambda) \right| \geq \varepsilon \right\} \leq -\eta(\varepsilon) < 0 \quad (10.30)$$

if ϕ is nonnegative, is supported on $[-1, 1]$, and has $\int \phi(x) dx = 1$. This covers Theorem 10.3 except when $n/2l \rightarrow 0$. But this case is trivial anyway since most intervals in $[-l, l]$ would have to be empty.

Proof of Theorem 10.4. It follows from the bounds in Lemma 10.2 that for $\lambda \rightarrow +\infty$ there exist constants $0 < c_1 < c_2 < \infty$ such that

$$c_1 \lambda^2 \leq F(\lambda) \leq c_2 \lambda^2 \quad (10.31)$$

Since $F(\lambda)$ is convex, this implies for some $0 < c_1 < c_2 < \infty$,

$$c_1 \lambda \leq \rho(\lambda) = F'(\lambda) \leq c_2 \lambda \quad (10.32)$$

Now because $P(\rho) = F(\lambda(\rho))$, (10.31) and (10.32) imply our theorem.

Proof of Theorem 10.5. We have the identity from Lemma 10.8,

$$\begin{aligned} \frac{d \log Z_{m, l}^0}{dl} &= \frac{n}{l} + \frac{1}{l} \int \sum \psi(x_i - x_j) d\mu_{nl}^0 \\ &= \frac{n}{l} + \frac{1}{l} \int \int_{-l}^l \sum \psi(x_i - x_j) \phi(x_j - x) dx d\mu_{nl}^0 \\ &\quad + \text{a small error} \end{aligned}$$

Clearly

$$\lim_{l \rightarrow \infty} \frac{dZ_{n, l}^0}{dl} = 2\rho + 2 \int \Psi(x_1 - x_2) \phi(x_2) R_2^\lambda(dx_1, dx_2) \quad (10.33)$$

Since $Z_{n, l}^0 \sim 2l \Psi(\rho) = 2l \Psi\left(\frac{n}{2l}\right)$,

$$\frac{dZ_{n, l}^0}{dl} = 2 \Psi\left(\frac{n}{2l}\right) - 2 \cdot l \cdot \frac{n}{2l^2} \Psi'\left(\frac{n}{2l}\right)$$

Therefore

$$\begin{aligned} \lim_{l \rightarrow \infty} \frac{dZ_{n, l}^0}{dl} &= 2 \Psi(\rho) - 2\rho \Psi'(\rho) \\ &= 2 \Psi(\rho) + 2\lambda \rho \\ &= 2F(\lambda) \end{aligned} \quad (10.34)$$

Combining (10.33) and (10.34)

$$P(\rho) = F(\lambda) = \rho + \int \psi(x_1 - x_2) \phi(x_2) R_2^\lambda(dx_1, dx_2) .$$

Proof of Theorem 10.6. Consider

$$\xi_{k,f} = \sum f_k(x_1, x_2) ,$$

where $f_k(x_1, x_2) = kg_1(k(x_1 - x_2))g_2(x_2)$,

$$\lim_{k \rightarrow \infty} \xi_{k,f} = \rho \sum g_2(x_2) \quad \text{a.e. by the ergodic theorem .}$$

If we take expectations with respect to μ_λ we obtain (10.9). Relation (10.10) is similar. Lemma 10.17 provides enough uniform integrability to justify interchanging limits and integration.

References

1. Guo, M.Z., Papanicolaou, G.C.: Self diffusion of interacting Brownian particles, Probabilistic Methods in Mathematical Physics, pp. 113–151. Ito, K, Ikeda N. (eds.). Tokyo: Kinokuniya 1987
2. Guo, M.Z., Papanicolaou, G.C., Varadhan, S.R.S.: Nonlinear diffusion limit for a system with nearest neighbor interaction. Commun. Math. Phys. **118**, 31–59 (1988)
3. Spohn, H.: Equilibrium fluctuations for interacting Brownian particles. Commun. Math. Phys. **103**, 1–33 (1986)
4. Olla, S., Varadhan, S.R.S.: Scaling limit for interacting Ornstein-Uhlenbeck Processes. Commun. Math. Phys. **135**, 355–378 (1991)
5. Ruelle, David: Statistical mechanics; rigorous results. New York: Benjamin 1969

Communicated by J.L. Lebowitz