

The roles of neuromelanin, binding of metal ions, and oxidative cytotoxicity in the pathogenesis of Parkinson's disease: a hypothesis

W. S. Enochs^{1,2}, T. Sarna³, L. Zecca⁴, P. A. Riley⁵, and H. M. Swartz^{1,6}

¹University of Illinois College of Medicine, Urbana, IL, and

²Department of Radiology, Massachusetts General Hospital and
Harvard Medical School, Boston, MA, U.S.A.

³Jagellonian University, Krakow, Poland

⁴Istituto Technologie Biomediche Avanzate, CNR, Milano, Italy

⁵University of London, London, United Kingdom

⁶Dartmouth Medical School, Hanover, NH, U.S.A

Accepted December 13, 1993

Summary. A characteristic feature of both Parkinson's disease (idiopathic paralysis agitans) and normal aging is loss of pigmented neurons in the substantia nigra. This has been found to correlate with the accumulation of neuromelanin and with oxidative stress in this brain region, but a clear association between these factors has not been established. Based on our recent demonstration that neuromelanin is a true melanin, containing bound metal ions in situ, we present a general model for its accumulation in vivo and the hypotheses (1) that it has a cytoprotective function in the sequestration of redox-active metal ions under normal conditions but (2) that it has a cytotoxic role in the pathogenesis of Parkinson's disease. Thus, neuromelanin accumulates normally through the autooxidation of catecholamines and serves tightly to bind redox-active metal ions, processes which would accelerate under conditions of intracellular or extracellular oxidative stress. Based on the known properties of melanin, however, neuromelanin also has the potential for exacerbating oxidative stress, eg by generating H₂O₂ when it is intact or by releasing redox-active metal ions if it loses its integrity; these reactions also would modulate the reactivity of the neuromelanin. By overwhelming intracellular antioxidative defense mechanisms, such a positive-feedback cycle could turn a condition of chronic or repeated oxidative stress in vulnerable neurons into an acute crisis, leading to cellular death. If the cumulative stress in duration and/or degree is severe enough, neuronal depletion could be sufficient to cause Parkinson's disease during life. One possible trigger for this cascade is suggested by the increased nigral iron contents in postmortem parkinsonian brains and the correlation of this disease with urban living where exposure to heavy metal ions is high:

the saturation of neuromelanin with redox-active metal ions. Parkinson's disease therefore may be a form of accelerated aging in the substantia nigra associated with environmental toxins in which neuromelanin has a central, active role.

Key words: Neuromelanin, melanins, metal ions, free radicals, oxygen toxicity, substantia nigra, aging, Parkinson's disease, idiopathic paralysis agitans.

Introduction

Parkinson's disease (PD, idiopathic paralysis agitans) is a neurological disorder of the extrapyramidal system characterized clinically by akinesia, rigidity, and tremor (Stern, 1990). Its most consistent pathological finding is extensive degeneration of the naturally pigmented dopaminergic neurons projecting from the pars compacta of the substantia nigra to the striatum (caudate nucleus and putamen) (Greenfield and Bosanquet, 1953; Pakkenberg and Brody, 1965), which causes a functional deficiency of dopamine in the latter area (Hornykiewicz, 1966; Bernheimer et al., 1973). In both PD and normal aging, the most highly pigmented neurons in this brain region are most susceptible to death (Pakkenberg and Brody, 1965; Graham, 1979; Mann and Yates, 1979, 1983; Hirsch et al., 1988). Although a parkinsonian syndrome similar to PD has been shown to result from acute exposure to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) (Langston et al., 1983; Singer et al., 1987; Adams and Odunze, 1991b), the etiologic mechanism of idiopathic PD remains unknown. Proposed causes of the latter have included viral infection (Elizan and Casals, 1983; Mattock et al., 1988), arteriosclerosis (Issidorides, 1971; Bernheimer et al., 1973), heredity (Poirier et al., 1991; Tanner, 1992), environmental and endogenous toxins (Langston et al., 1987; Fahn, 1989; Goldsmith et al., 1990; Poirier et al., 1991; Furtado and Mazurek, 1991; Tanner, 1989, 1992), and mechanical disruption of neurons by accumulating neuromelanin (Mann and Yates, 1977, 1983).

Since the discovery of parkinsonism induced in primates by exposure to MPTP, research into the etiology of idiopathic PD in humans has intensified, and strong circumstantial evidence has accrued suggesting that neuronal loss in the substantia nigra may result from lethal oxidative stress [for reviews, see Cohen (1983, 1986), Graham (1984), Hornykiewicz and Kish (1986), Fahn (1989), Halliwell (1989), Youdim et al. (1989), Martilla and Rinne (1989), Olanow (1990), Götz et al. (1990), Ben-Shachar and Youdim (1990), Adams and Odunze (1991a,b), Olanow (1992), Jenner (1992), Jenner et al. (1992), Hirsch (1992), Dexter et al. (1992), Ben-Shachar et al. (1992)]. This largely is based on: the production of hydrogen peroxide occurring with the normal and abnormal metabolism of dopamine (Cohen, 1983, 1986; Graham, 1984); increased amounts of redox-active metal ions, principally iron, observed in diseased substantia nigra and the known ability of such ions in vitro to reduce hydrogen peroxide to the potentially highly toxic hydroxyl radical (Halliwell, 1989; Adams and Odunze, 1991b; Dexter et al.,

1992; Riederer et al., 1992; Ben-Shachar et al., 1992); evidence for a disease-related reduction in antioxidative capacity in this brain region (Jenner, 1992; Jenner et al., 1992); and the presence of increased products of lipid peroxidation (Jenner, 1992; Jenner et al., 1992). While a role for neuromelanin in this process occasionally has been suggested (Lindquist et al., 1987; Youdim et al., 1989; Ben-Schachar and Youdim, 1990; Ben-Shachar et al., 1992; Hirsch, 1992; Jellinger et al., 1992), however, such a role has not been clearly defined nor has an attempt been made to relate the preceding observations to the accumulation of neuromelanin.

By combining previous and current work by others with recent results from our laboratories, we present a comprehensive hypothesis for the pathogenesis of PD which centers on an active role for neuromelanin. This is contrary to prior suggestions that it has a passive role, such as the mechanical disruption of neurons (Mann and Yates, 1977, 1983) or the binding and concentrating of environmental toxins similar to MPTP (Lindquist et al., 1987; D'Amato et al., 1987; Fahn, 1989). Our proposal provides a general model for the accumulation of neuromelanin in catecholaminergic neurons and suggests a cytoprotective function for it under normal conditions. However, it also offers a plausible mechanism for the depletion of pigmented neurons during normal aging and its acceleration in PD, postulating a specific etiologic trigger for the latter that is consistent with the cytoprotective function.

Nature of neuromelanin

Neuromelanin is a poorly-defined pigment [for reviews, see Marsden (1969), Van Woert and Ambani (1974), Barden (1975, 1981), Barden and Brizzee (1987)] that accumulates normally with age in the central catecholaminergic neurons of most species (Bazelon and Fenichel, 1967; Mann and Yates, 1974; Bogerts, 1981). It is most abundant in the substantia nigra and locus coeruleus (which are dopaminergic and noradrenergic nuclei, respectively, in the mesencephalon) (Bazelon and Fenichel, 1967; Bogerts, 1981) and increases in amount between species according to their phylogenetic proximity to man (Marsden, 1961). Histological studies have demonstrated that neuromelanin occurs as a set of membrane-limited granules within the neuronal perikaryon (Moses et al., 1966; Hirosawa, 1968), while extensive histochemical analyses provide strong evidence that: these granules are inactivated lysosomes; the pigment occurs in association with lipofuscin; and it has a high content of sulfur [for reviews, see Barden (1975, 1981), Barden and Brizzee (1987)]. Spectroscopic and degradative studies further indicate that a major precursor of neuromelanin is the catecholamine neurotransmitter specific to the pigmented neuron (e.g. dopamine in the substantia nigra and norepinephrine in the locus coeruleus) (Van Woert et al., 1967; Maeda and Wegmann, 1969; Nordgren et al., 1971; Das et al., 1978; Carstam et al., 1991; Zecca et al., 1992).

Based on results with the usual histochemical and physicochemical methods for detecting melanins, neuromelanin has been classified as a

melanin (Lillie and Yamada, 1960; Marsden, 1969; Van Woert and Ambani, 1974; Barden, 1975, 1981; Barden and Brizzee, 1987); however, these methods often give ambiguous results (Sarna and Swartz, 1978). By comparison, electron paramagnetic resonance (EPR, or equivalently, electron spin resonance, ESR) spectroscopy is a highly sensitive and specific technique for investigating melanins [for review, see Sealy et al. (1980)] but has been under-utilized in the study of neuromelanin. Although the latter pigment has been shown by EPR spectroscopy to contain a stable population of organic free radicals which increases upon exposure to visible light (Van Woert et al., 1967; Mikulski, 1970), properties characteristic of more typical melanins (Sarna and Swartz, 1978), these early studies were limited in scope. Since then, little more has been done despite the development of a fairly complete understanding of the paramagnetic nature of melanins in general (Sealy et al., 1980).

Using an established set of EPR criteria for the unambiguous identification and characterization of melanins (Sarna and Swartz, 1978; Enochs et al., 1993a), we have demonstrated recently that human neuromelanin purified from normal substantia nigra is a true melanin (Enochs et al., 1993b). However, compared to purified bovine eye melanin (a typical and fairly well-characterized eumelanin) and various synthetic melanins, it is an atypical melanin that is not well modeled by synthetic dopamine melanin. Some of the unusual features of neuromelanin can be explained by postulating two distinct sources for its free radicals, the dominant one possibly derived from a precursor containing sulfur. This is consistent with recent degradative studies of neuromelanin indicating that it may be a heterogeneous copolymer of dopamine and cysteine or glutathione (Carstam et al., 1991; Zecca et al., 1992).

We also have demonstrated by EPR spectroscopy the presence of significant amounts of paramagnetic metal ions in unprocessed human substantia nigra, part of which at least are bound to neuromelanin in situ (Enochs et al., 1993b). Although other physicochemical studies have shown that this brain region grossly contains large quantities of Fe^{3+} , Cu^{2+} , and Mn^{2+} (Warren et al., 1960; Cumings, 1968; Earle, 1968; Larsen et al., 1979; Sofic et al., 1988; Dexter et al., 1989, 1991; Riederer et al., 1989; Uitti et al., 1989; Hirsch et al., 1991), histochemical studies have suggested that the metal ions are localized to storage granules within glial cells and are not endogenous to neuromelanin granules in neurons (Barden, 1971; Hill and Switzer, 1984; Connor et al., 1990; Jellinger et al., 1990; Sofic et al., 1991). However, our data have been confirmed recently by other sophisticated physicochemical techniques (Swartz et al., 1992; Perl and Good, 1992; Jellinger et al., 1992), indicating that histochemical staining reactions for metal ions may not be reliable when the ions are bound to melanin.

Interaction of typical melanins with O₂ under physiological conditions and their potential toxicity in vivo

Our unambiguous demonstration that neuromelanin is a true (albeit atypical) melanin containing bound redox-active metal ions has significant pathophysiological implications when the reactivity of typical melanins toward molecular oxygen (O₂) and its reduced forms is considered [for review, see Sarna and Swartz (1993)]. In the presence of O₂, nonilluminated melanin at physiological pH is a low-level source in vitro of both superoxide anion (O₂^{•-}) and hydrogen peroxide (H₂O₂) (Felix et al., 1978; Sarna et al., 1980a; Sarna and Sealy, 1984). Superoxide anion is produced from the univalent reduction of O₂ by hydroquinonoid and semiquinonoid subunits in melanin (Table 1, Eqs. 1a,b); although most of this O₂^{•-} is reoxidized to O₂ by quinonoid subunits (Eq. 1b), part of it spontaneously dismutates (Eq. 2) or is reduced further by melanin to form H₂O₂ (Eqs. 3a,b) (Korytowski et al., 1985). This process is accelerated greatly by exposure to ultraviolet (UV) or even visible light (Korytowski et al., 1987). Because H₂O₂ is relatively inert toward pure melanin at physiologic pH, the majority of it appears in the bulk solution (Korytowski and Sarna, 1990).

When containing bound redox-active metal ions or especially in the presence of simple chelate complexes of these ions, nonilluminated melanin also is a source in vitro of the highly reactive hydroxyl radical (OH[•]), even at physiologic pH. This occurs through a Fenton-type (Halliwell, 1981) decomposition of endogenous or exogenous H₂O₂ (Eq. 5a) and, like the production of O₂^{•-} and H₂O₂, is accelerated by exposure to UV or visible light (Korytowski et al., 1987). Simple chelate complexes of redox-active

Table 1. Reactions between melanins, bound redox-active metal ions, and oxygen and its reduced forms

1a	$\text{Mel-Q}^{2-} + \text{O}_2 \rightleftharpoons \text{Mel-Q}^{\cdot-} + \text{O}_2^{\cdot-}$
b	$\text{Mel-Q}^{\cdot-} + \text{O}_2 \rightleftharpoons \text{Mel-Q} + \text{O}_2^{\cdot-}$
2	$\text{O}_2^{\cdot-} + \text{O}_2^{\cdot-} + 2\text{H}^+ \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$
3a	$\text{Mel-Q}^{2-} + \text{O}_2^{\cdot-} + 2\text{H}^+ \rightarrow \text{Mel-Q}^{\cdot-} + \text{H}_2\text{O}_2$
b	$\text{Mel-Q}^{\cdot-} + \text{O}_2^{\cdot-} + 2\text{H}^+ \rightarrow \text{Mel-Q} + \text{H}_2\text{O}_2$
4	$\text{Chel-M}^n + \text{H}_2\text{O}_2 \rightarrow \text{Chel-M}^{n+1} + \text{OH}^{\cdot} + \text{OH}^-$
5a	$\text{Mel-M}^n + \text{H}_2\text{O}_2 \rightarrow \text{Mel-M}^{n+1} + \text{OH}^{\cdot} + \text{OH}^-$
b	$\text{Mel}_{\text{red}}\text{-M}^{n+1} \rightarrow \text{Mel}_{\text{ox}}\text{-M}^n$
6	$\text{Mel} + \text{OH}^{\cdot} \rightarrow \text{Mel}_{\text{ox}}$

*Mel-Q*²⁻ Hydroquinonoid melanin subunits, *Mel-Q* Quinonoid melanin subunits, *Mel-Q*^{•-} Semiquinonoid melanin subunits, *O*₂ Molecular oxygen, *O*₂^{•-} Superoxide anion, *H*₂*O*₂ Hydrogen peroxide, *OH*[•] Hydroxyl radical, *OH*⁻ Hydroxide anion, *H*⁺ Proton, *Chel-M*^{*n*} Chelate-metal ion complex, *Mel-M*^{*n*} Melanin-metal ion complex, *Mel*_{red} Reduced melanin, *Mel*_{ox} Oxidized melanin

metal ions are known to cause an analogous breakdown of H_2O_2 (Eq. 4) (Halliwell, 1981), but this reaction is expected quickly to cease as the ions are oxidized to their higher valencies. By contrast, melanin is a redox polymer which can reduce the oxidized metal ions, leading to cycling of their valencies such that a steady-state flux of $\text{OH}\cdot$ is possible under conditions of constant H_2O_2 concentration or O_2 tension (Eqs. 5a,b) (Korytowski and Sarna, 1990). The vast majority of $\text{OH}\cdot$ is scavenged immediately at the site of its formation, thus causing peroxidative degradation of the melanin (Eq. 6) (Sarna et al., 1986; Korytowski and Sarna, 1990) while simultaneously limiting damage to other molecules that may be present. Specifically, melanin has been shown to accelerate the rate of production of $\text{OH}\cdot$ in the presence of H_2O_2 and Fe^{3+} or Cu^{2+} , which occurs because of reduction of Fe^{3+} to Fe^{2+} or Cu^{2+} to Cu^+ (Hintz and Kalyanaram, 1986; Korytowski et al., 1987; Pilas et al., 1988; Korytowski and Sarna, 1990).

Since melanin has a high affinity for metal ions (White, 1958; Bruenger et al., 1967; Potts and Au, 1976; Sarna et al., 1976, 1980b, 1981; Larsson and Tjalve, 1978; Froncisz et al., 1980; Lyden et al., 1984; Ben-Shachar et al., 1991) and pigmented tissues have been shown to contain impressive amounts of metal ions bound to it in situ, including a variety of redox-active species (Bowness et al., 1952; Stein, 1955; Horcicko et al., 1973; Szerkeres, 1975; Parkinson et al., 1979; Sarna et al., 1980; Okazaki et al., 1985; Enochs et al., 1993b), the observations above suggest that natural melanin may be a source in vivo of O_2^- , H_2O_2 , and $\text{OH}\cdot$, especially on exposure to UV light. Since these species potentially are highly cytotoxic [for reviews, see Halliwell and Gutteridge (1984)], this appears to contradict the generally accepted role of melanin in photoprotection (Pathak et al., 1976; Kollias et al., 1991). However, it is interesting that in skin and hair, which are most exposed to such illumination, the cells specialized for the synthesis of melanin (melanocytes) transfer their melanin to a cell line destined for terminal inactivation and sloughing (keratinocytes) (Jimbow et al., 1976). Furthermore, melanocytes in both the skin and retina are screened at least partly from UV light by the overlying stratum corneum and epidermis in the first instance and by the cornea, iris, and lens in the second (Boettner and Wolter, 1962; Pathak et al., 1976; Kollias et al., 1991).

Under normal conditions, the potential toxicity of melanin in vivo also may be limited by its occurrence within membrane-limited organelles (melanosomes) (Jimbow et al., 1976). Due to its low solubility in lipids (Rumyantseva et al., 1979) and presumably slow diffusion out of the melanosome, any intramelanosomal O_2^- would tend to be reoxidized by melanin or reduced further to H_2O_2 , while any $\text{OH}\cdot$ generated from the decomposition of intramelanosomal or extramelanosomal H_2O_2 would react immediately with melanin or other contents of the organelle (including its membrane). Thus, the melanosome normally would act as a low-level source mainly of H_2O_2 , which then could be scavenged readily by intracellular peroxidases and catalase [for review, see Halliwell (1981)].

Based on the paramagnetic characteristics of melanin and its high affinity for metal ions, at least two cytoprotective functions other than photoabsorption have been proposed for natural melanin. One is the intracellular scavenging of extraneous or light-induced radicals (Commoner et al., 1954; Mason et al., 1960; Cotzias et al., 1964; Lukiewicz, 1972; Barbeau, 1984a; Sarna et al., 1986), but this is difficult to reconcile with the presence of a relatively impermeable lipid membrane around the melanosome. Another is the sequestering of redox-active metal ions from the cytoplasm under normal conditions, as recently suggested for melanin in the retinal pigment epithelium (RPE) (Sarna, 1992). By the mechanisms above, however, intense or prolonged illumination (especially at short wavelengths) may cause sufficient structural deterioration of melanin that its capacity for binding metal ions is lost and sufficient secondary peroxidation of the melanosomal membrane that its integrity is disrupted. This could result in leakage of the melanosomal contents, including free metal ions, into the cytoplasm with grave consequences for the cell. A similar but more gradual process also could occur during normal aging of melanosomes and would be especially problematic for postmitotic cells, such as those in the RPE, for which no renewal of melanin occurs (Sarna, 1992).

A model for the accumulation of neuromelanin

Most existing data strongly support an autooxidative mechanism for the accumulation of neuromelanin in catecholaminergic neurons (Van Woert et al., 1967; Maeda and Wegmann, 1969; Nordgren et al., 1971; Das et al., 1978; Graham, 1979; Carstam et al., 1991; Zecca et al., 1992). While our recent results have confirmed that neuromelanin is a true melanin (Enochs et al., 1993b), previous histochemical analyses indicate that it may occur in association with lipofuscin (Van Woert and Ambani, 1974; Barden, 1975, 1981; Barden and Brizzee, 1987), and a recent study has shown that it has an intrinsic component of fatty acids (Zecca et al., 1992). As the classic "age pigment," lipofuscin generally is accepted as a nonbiodegradable product of lipid peroxidation that collects in the lysosomes of postmitotic cells during normal aging [for reviews, see Dolman and MacLeod (1981), Barden and Brizzee (1987)]. Therefore, it is reasonable to assume that the accumulation of neuromelanin in catecholaminergic brain nuclei occurs by analogy to lipofuscin elsewhere in the central nervous system.

Because no enzymatic mechanism for its formation has been found, neuromelanin most likely is a consequence of the occasional oxidation of catecholamines (e.g. dopamine in the substantia nigra and norepinephrine in the locus coeruleus). Catecholamines *in vitro* undergo autooxidation in the presence of O₂, though inefficiently at physiologic pH (Graham et al., 1978a,b; Sealy et al., 1984), and also enzymatic oxidation in the presence of H₂O₂ and peroxidase (Okun et al., 1971; Kalyanaraman et al., 1984). Since the normal catabolism of catecholamines by monoamine oxidase and their autooxidation by O₂ are known sources of H₂O₂ (Cohen, 1983, 1986; Graham, 1984), the intracellular milieu of catecholaminergic neurons would

predispose toward a nonphysiological oxidation of catecholamines in vivo to their corresponding semiquinones and quinones. However, while these species are relatively reactive electrophiles and thus potentially cytotoxic themselves (Graham et al., 1978b; Kalyanaraman et al., 1987), they normally would undergo prompt reduction to the original catecholamine by antioxidants such as glutathione and ascorbate (Heacock and Laidlaw, 1958; Heacock and Scott, 1959; Tse et al., 1976).

Despite the presence of antioxidative mechanisms in catecholaminergic neurons for maintaining their neurotransmitters in a reduced state, it is expected that this system is not perfect. Some of the semiquinones and quinones formed by the oxidation of catecholamines may escape reduction on occasion and randomly form dimers and oligomers with the original catecholamine (Graham et al., 1978b; Sealy et al., 1984) or become reductively coupled to the cysteinyl groups of GSH or proteins (Graham, 1978b; Rosengren et al., 1985; Fornstedt et al., 1986, 1989; Kalyanaraman et al., 1987; d'Ischia et al., 1987a,b; Ito et al., 1988) or possibly to other cellular constituents. These products probably could not be catabolized in general and would gradually collect in lysosomes, perhaps copolymerizing with the products of lipid peroxidation to form a type of melanized lipofuscin unique to catecholaminergic brain nuclei. Since neurons are post-mitotic cells, the accumulation of neuromelanin, like lipofuscin elsewhere, thus would be a measure of the cumulative exposure (in both duration and degree) of individual neurons to oxidative stress.

This model for the accumulation of neuromelanin and our observation of paramagnetic metal ions bound to it in situ (Enochs et al., 1993b) are consistent with previous suggestions that it is a waste product of dopamine metabolism (Graham, 1978a). Nevertheless, this origin still is compatible with a cytoprotective function for neuromelanin under normal conditions. Catecholaminergic neurons are exposed to unusually high levels of H_2O_2 because of the enzymatic catabolism and autooxidation of dopamine. The relative stability of H_2O_2 and its ability to diffuse long distances in cells have led to the proposal that the cytotoxicity of H_2O_2 ultimately results from its decomposition to $OH\cdot$, catalyzed in a Fenton-type reaction by free or loosely bound redox-active metal ions in the cytoplasm and causing site-specific peroxidation of lipids, proteins, and nucleic acids (Halliwell and Gutteridge, 1984). In analogy to melanosomes, neuromelanin granules may sequester such metal ions normally, thus reducing their cytoplasmic concentration and confining the products of their reaction with H_2O_2 to the granules. Through this process, neuromelanin granules also could complement established mechanisms for scavenging H_2O_2 , such as glutathione peroxidase and catalase (Halliwell, 1981), by acting as scavengers themselves of any excess H_2O_2 escaping enzymatic degradation.

Hypothetical pathogenesis for the loss of pigmented neurons during normal aging and in Parkinson's disease

Based on the model above for the accumulation of neuromelanin, two possibilities are suggested for the pathogenesis of neuronal loss in the sub-

stantia nigra during normal aging and in PD, which are consistent with observations that the most highly pigmented neurons in this brain region are most susceptible to death (Pakkenberg and Brody, 1965; Graham, 1979; Mann and Yates, 1979, 1983; Hirsch et al., 1988) and evidence that this may be due to oxidative stress (Cohen, 1983, 1986; Graham, 1984; Hornykiewicz and Kish, 1986; Fahn, 1989; Halliwell, 1989; Youdim et al., 1989; Martilla and Rinne, 1989; Olanow, 1990, 1992; Gotz et al., 1990; Ben-Shachar and Youdim, 1990; Adams and Odunze, 1991a,b; Jenner, 1992; Jenner et al., 1992; Hirsch, 1992; Dexter et al., 1992; Ben-Shachar et al., 1992). First, neuromelanin may form as an indicator of oxidative stress, reflecting an independent, yet-unidentified source of the stress which is endogenous or exogenous to catecholaminergic neurons and generates neuromelanin at a proportionate rate. Second, neuromelanin itself may contribute to oxidative stress as a source of H_2O_2 when it is intact or of redox-active metal ions if it loses its integrity. More likely still is an interplay between these two roles of indicator and contributor of oxidative stress in which neuromelanin could have a role in its own formation.

According to the model, neuromelanin and lipofuscin normally form early in life as the result of independent oxidative stressor(s) and imperfect antioxidative defense mechanisms, thus accumulating gradually with age. Regardless of whether neuromelanin has a cytoprotective function, the formation of both pigments is expected to accelerate in specific neurons experiencing chronic or repeated oxidative stress. Whereas both neuromelanin and lipofuscin granules also contain paramagnetic metal ions bound to the pigments in situ (Vistnes et al., 1983; Enochs et al., 1993b), however, only the former is susceptible to peroxidative degradation and leakage of its potentially toxic contents into the cytoplasm. This is analogous to melanosomes but would occur by exposure to extragranular H_2O_2 from the metabolism of dopamine rather than to intramelanosomal H_2O_2 generated by illumination. Since neurons are postmitotic and neuromelanin apparently does not undergo turnover (Lindquist et al., 1987), such degradation could occur gradually with normal aging of the oldest granules or acutely by a sudden oxidative insult. Over time, the contribution of neuromelanin to oxidative stress thus could result in a positive-feedback cycle leading to a condition of chronic oxidative stress with increased degradation of metal-laden old neuromelanin, eventually overwhelming antioxidative defense mechanisms, with an acute crisis ending in neuronal death.

Such a positive-feedback cycle of increasing oxidative stress not only provides a plausible explanation for the loss of pigmented neurons in the substantia nigra at advanced ages and in idiopathic PD but also is compatible with suggestions that the latter is a form of accelerated aging in this brain region (Barbeau, 1984b; Langston, 1989; Wolters and Calne, 1989). During normal aging, neuronal death occurring sporadically with age in relatively few vulnerable neurons, though progressing somewhat faster at advanced ages, would not cause a functional deficiency of dopaminergic neurons during the average human lifespan since symptoms do not appear until more than 80% of the neurons are lost (Bernheimer et al., 1973). However, chronic oxidative stress in the substantia nigra as a whole would

accelerate this process, and a single intense oxidative insult or repeated smaller insults could lead to the simultaneous degradation of many neuromelanin granules with the acute loss of a large number of neurons. Whether parkinsonism subsequently developed during life would depend on: the density of neurons in the substantia nigra at birth; the age of the individual (determining the number of remaining neurons, the average neuronal content of neuromelanin, and the efficacy of antioxidative defense mechanisms (Fahn, 1989) when the stress began or the insult occurred; and the duration and intensity of the stress or insult. Furthermore, an acute oxidative insult occurring remotely and causing a subcritical depletion of neurons (less than 80%) eventually could lead to parkinsonism because of chronic stress in the remaining neurons secondary to compensatory overactivity (Hornykiewicz, 1966; Bernheimer et al., 1973; Fornstedt et al., 1989), with a more rapid decline in their number than previously.

The proximate trigger(s) for the pathogenesis of idiopathic PD according to the hypothesis above is not specified. Certainly any substance or process would be a candidate if it raised the intraneuronal concentration of H_2O_2 or independently altered the structure or composition of the neuromelanin granule, making it more susceptible to degradation or otherwise decreasing its postulated cytoprotective ability. One strong possibility consistent with the latter is suggested by the increased content of iron observed in the postmortem substantia nigras of patients with PD (Dexter et al., 1992; Riederer et al., 1992; Ben-Shachar et al., 1992; Perl and Good, 1992; Jellinger et al., 1992) and the greater incidence of this disease in industrial areas where exposure to heavy metals is high (Tanner, 1989; Goldsmith et al., 1990). Elevated levels of redoxactive metal ions in the microenvironment of the neuromelanin granule could lead to saturation of neuromelanin with these ions and the sequelae above. Our hypothesis thus provides for a multifactorial etiology for PD but one in which neuromelanin has a central, active role in its pathogenesis.

Future directions

Additional information is required fully to evaluate our model for the accumulation of neuromelanin and our hypotheses that it has a cytoprotective function under normal conditions but a cytotoxic role in the pathogenesis of PD. This includes: a more precise determination of the nature of purified neuromelanin and its physicochemical properties, including any changes associated with aging or PD; the identification and quantitation of the types and amounts of metal ions actually contained within intact neuromelanin granules and any age- or disease-related differences in them; and a determination of the reactivities of genuine neuromelanin towards O_2 and its reduced forms, both with and without its normal complement of metal ions, and how these reactivities compare with those of more typical melanins. Elucidating the physicochemical structure of neuromelanin also would lead to a more complete understanding of the mechanism of its formation and would facilitate the development of a synthetic model pigment for use in

extensive investigations of its reactivity *in vitro* and, using cultured cells, its potential toxicity *in vivo*.

With such information and a validation of our model and hypotheses, methods for the diagnosis of PD at a subclinical stage might be further advanced, such as using magnetic resonance or PET imaging of the midbrain to detect subtle changes in the anatomical structure, chemical composition, or metabolism of the substantia nigra (Olanow, 1992; Eidelberg, 1992). Furthermore, novel rational approaches to medical therapy, prophylaxis, or prevention at this early stage might be devised to complement the successful uses of L-dopa and the monoamine oxidase B inhibitor deprenyl in symptomatic PD (Hornykiewicz, 1973; Parkinson Study Group, 1993). Although a recent study indicates that administration of alpha tocopherol does not slow the progression of established PD (Parkinson Study Group, 1993), the success of an approach using ascorbate (Fahn, 1989) or other antioxidants (Hall, 1992) or one using metal chelators such as deferoxamine (Shoulson, 1992) would be consistent with our proposals.

Acknowledgements

The authors gratefully acknowledge the support of this research by the Parkinson's Disease Foundation, the United Parkinson's Disease Foundation, the American Parkinson's Disease Foundation, and the Polish National Committee for Scientific Research (grant KBN-2225).

References

- Adams JD, Odunze IN (1991a) Biochemical mechanisms of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine toxicity: could oxidative stress be involved in the brain? *Biochem Pharmacol* 41: 1099–1105
- Adams JD, Odunze IN (1991b) Oxygen free radicals and Parkinson's disease. *Free Radic Biol Med* 10: 161–169
- Barbeau A (1984a) Manganese and extrapyramidal disorders. *Neurotoxicology* 5: 13–36
- Barbeau A (1984b) Etiology of Parkinson's disease: a research strategy. *Can J Neurol Sci* 11: 24–28
- Barden H (1971) The histochemical distribution and localization of copper, iron, neuromelanin and lysosomal enzyme activity in the brain of aging rhesus monkey and the dog. *J Neuropathol Exp Neurol* 30: 650–667
- Barden H (1975) The histochemical relationships and nature of neuromelanin. In: Brody H, Harman D, Ordy JM (eds) *Aging*, vol 1. Raven Press, New York, pp 79–117
- Barden H (1981) The biology and chemistry of neuromelanin. In: Sohal RS (ed) *Age pigments*. Elsevier/North-Holland Biomedical Press, Amsterdam, pp 155–180
- Barden H, Brizzee KR (1987) The histochemistry of lipofuscin and neuromelanin. In: Totaro EA, Glees P, Pisanti FA (eds) *Advances in the biosciences*, vol 64. Pergamon Press, Oxford, pp 339–392
- Bazelon M, Fenichel GM (1967) Studies on neuromelanin I. A melanin system in the human adult brainstem. *Neurology* 17: 512–519
- Ben-Shachar D, Youdim MBH (1990) Selectivity of melanized nigra-striatal dopamine neurons to degeneration in Parkinson's disease may depend on iron-melanin interaction. *J Neural Transm [Suppl]* 29: 251–258
- Ben-Shachar D, Riederer P, Youdim MBH (1991) Iron-melanin interaction and lipid peroxidation: implications for Parkinson's disease. *J Neurochem* 57: 1609–1614

- Ben-Shachar D, Eshel G, Riederer P, Youdim MBH (1992) Role of iron and iron chelation in dopaminergic-induced neurodegeneration: implications for Parkinson's disease. *Ann Neurol [Suppl]* 32: S105–S110
- Bernheimer H, Birkmayer W, Hornykiewicz O, Jellinger K, Seitelberger F (1973) Brain dopamine and the syndromes of Parkinson and Huntington: clinical, morphological and neurochemical correlations. *J Neurol Sci* 20: 415–455
- Boettner EA, Wolter JR (1962) Transmission in the ocular media. *Invest Ophthalmol Vision Res* 1: 776–781
- Bogerts B (1981) A brainstem atlas of catecholaminergic neurons in man, using melanin as a natural marker. *J Comp Neurol* 197: 63–80
- Bowness JM, Morton RA, Shakir MH, Stubbs AL (1952) Distribution of copper and zinc in mammalian eyes. Occurrence of metals in melanin fractions from eye tissues. *Biochem J* 51: 521–530
- Bruenger FW, Stover BJ, Atherton DR (1967) The incorporation of various metal ions into in vivo- and in vitro-produced melanin. *Radiat Res* 32: 1–12
- Carstam R, Brinck C, Hindemith-Augustsson A, Rorsman H, Rosengren E (1992) The neuromelanin of the human substantia nigra. *Biochem Biophys Acta* 1097: 152–160
- Cohen G (1983) The pathobiology of Parkinson's disease: biochemical aspects of dopamine neuron senescence. *J Neural Transm [Suppl]* 19: 89–103
- Cohen G (1986) Monoamine oxidase, hydrogen peroxide, and Parkinson's disease. *Adv Neurol* 45: 119–125
- Commoner B, Townsend J, Pake GE (1954) Free radicals in biological materials. *Nature* 74: 689–691
- Connor JR, Menzies SL, St. Martin SM, Musson EJ (1990) Cellular distribution of transferrin, ferritin, and iron in normal and aged human brains. *J Neurosci Res* 27: 595–611
- Cotzias GC, Papavasiliou PS, Van Woert MH, Sakamoto A (1964) Melanogenesis and extrapyramidal diseases. *Fed Proc* 23: 713–718
- Cummings JN (1968) Trace metals in the brain and in Wilson's disease. *J Clin Pathol* 21: 1–7
- D'Amato RJ, Alexander GM, Schwartzman RJ, Kitt CA, Price DL, Snyder SH (1987) Evidence for neuromelanin involvement in MPTP-induced neurotoxicity. *Nature* 327: 324–326
- Das KC, Abramson MB, Katzman R (1978) Neuronal pigments: spectroscopic characterization of human brain melanin. *J Neurochem* 30: 601–605
- Dexter DT, Wells FR, Lees AJ, Agid Y, Jenner P, Marsden CD (1989) Increased nigral iron content and alterations in other metal ions occurring in brain in Parkinson's disease. *J Neurochem* 52: 1830–1836
- Dexter DT, Carayon A, Javoy-Agid F, Agid Y, Wells FR, Daniel SE, Lees AJ, Jenner P, Marsden CD (1991) Alterations in the levels of iron, ferritin and other trace metals in Parkinson's disease and other neurodegenerative diseases affecting the basal ganglia. *Brain* 114: 1953–1975
- Dexter DT, Jenner P, Schapira AHV, Marsden CD (1992) Alterations in levels of iron, ferritin, and other trace metals in neurodegenerative diseases affecting the basal ganglia. *Ann Neurol [Suppl]* 32: S84–S100
- d'Ischia M, Napolitano A, Protà G (1987) Sulphydryl compounds in melanogenesis. Part I. Reaction of cysteine and glutathione with 5,6-dihydroxyindoles. *Tetrahedron* 43: 5351–5356
- d'Ischia M, Napolitano A, Protà G (1987) Sulphydryl compounds in melanogenesis. Part II. Reactions of cysteine and glutathione with dopachrome. *Tetrahedron* 43: 5357–5362
- Dolman CL, MacLeod PM (1981) Lipofuscin and its relation to aging. *Adv Cell Neurobiol* 2: 205–247
- Earle KM (1968) Studies on Parkinson's disease including X-ray fluorescent spectroscopy of formalin-fixed brain tissue. *J Neuropathol Exp Neurol* 27: 1–14

- Eidelberg D (1992) Positron emission tomography studies in parkinsonism. *Neurol Clin* 10(2): 421–433
- Elizan TS, Casals J (1983) The viral hypothesis in Parkinsonism. *J Neural Transm [Suppl]* 19: 75–88
- Enochs WS, Nilges MJ, Swartz HM (1993a) A standardized test for the identification and characterization of melanins using electron paramagnetic resonance (EPR) spectroscopy. *Pigment Cell Res* 6: 91–99
- Enochs WS, Nilges MJ, Swartz HM (1993b) Purified human neuromelanin, synthetic dopamine melanin as a potential model pigment, and the normal human substantia nigra: characterization by electron paramagnetic resonance (EPR) spectroscopy. *J Neurochem* 61: 68–79
- Fahn S (1989) The endogenous toxin hypothesis of the etiology of Parkinson's disease and a pilot trial of high-dosage antioxidants in an attempt to slow the progression of the illness. *Ann NY Acad Sci* 570: 186–196
- Fahn S (1992) A pilot trial of high-dose alpha-tocopherol and ascorbate in early Parkinson's disease. *Ann Neurol [Suppl]* 32: S128–S132
- Felix CC, Hyde JS, Sarna T, Sealy RC (1978) Melanin photoreactions in aerated media: electron spin resonance evidence for production of superoxide and hydrogen peroxide. *Biochem Biophys Res Commun* 84: 335–341
- Fornstedt B, Rosengren E, Carlsson A (1986) Occurrence and distribution of 5-S-cysteinyl derivatives of dopamine, DOPA, and DOPAC in the brains of eight mammalian species. *Neuropharmacology* 25: 451–454
- Fornstedt B, Brun A, Rosengren E, Carlsson A (1989) The apparent autoxidation rate of catechols in dopamine-rich regions of human brains increases with the degree of depigmentation of substantia nigra. *J Neural Transm [P-D Sect]* 1: 279–295
- Furtado JCS, Mazurek MF (1991) MPTP-induced neurotoxicity and the quest for a preventive therapy for Parkinson's disease. *Can J Neurol Sci* 18: 77–82
- Froncisz W, Sarna T, Hyde JS (1980) Cu^{2+} probe of metal ion binding sites in melanin using electron paramagnetic resonance spectroscopy I. Synthetic melanins. *Arch Biochem Biophys* 202: 289–303
- Goldsmith JR, Herishanu Y, Abarbanel JM, Weinbaum Z (1990) Clustering of Parkinson's disease points to environmental etiology. *Arch Environ Health* 45: 88–94
- Götz ME, Freyberger A, Riederer P (1990) Oxidative stress: a role in the pathogenesis of Parkinson's disease. *J Neural Transm [Suppl]* 29: 241–249
- Graham DG (1978a) Oxidative pathways for catecholamines in the genesis of neuromelanin and cytotoxic quinones. *Mol Pharmacol* 14: 633–643
- Graham DG, Tiffany SM, Bell WR, Gutknecht WF (1978b) Autoxidation versus covalent binding of quinones as the mechanism of toxicity of dopamine, 6-hydroxydopamine, and related compounds toward C1300 neuroblastoma cells in vitro. *Mol Pharmacol* 14: 644–653
- Graham DG (1979) On the origin and significance of neuromelanin. *Arch Pathol Lab Med* 103: 359–362
- Graham DG (1984) Catecholamine toxicity: a proposal for the molecular pathogenesis of manganese neurotoxicity and Parkinson's disease. *Neurotoxicology* 5: 83–96
- Greenfield JG, Bosanquet FD (1953) The brain-stem lesions in parkinsonism. *J Neurol Neurosurg Psychiatry* 16: 213–226
- Hall ED (1992) Novel inhibitors of iron-dependent lipid peroxidation for neurodegenerative disorders. *Ann Neurol [Suppl]* 32: S137–S142
- Halliwell B (1981) Free radicals, oxygen toxicity and aging. In: Sohal RS (ed) *Age pigments*. Elsevier/North-Holland Biomedical Press, Amsterdam, pp 1–62
- Halliwell B, Gutteridge JMC (1984) Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J* 219: 1–14
- Halliwell B (1989) Oxidants and the central nervous system: some fundamental questions. *Acta Neurol Scand* 126: 23–33

- Heacock RA, Laidlaw BD (1958) Reduction of adrenochrome with ascorbic acid. *Nature* 182: 526–527
- Heacock RA, Scott BD (1959) The reduction of adrenochrome with thiols. *Can J Biochem Physiol* 37: 1087–1088
- Hill JM, Switzer RC (1984) The regional distribution and cellular localization of iron in the rat brain. *Neuroscience* 11: 595–603
- Hintz P, Kalyanaraman B (1986) Metal ion-induced activation of molecular oxygen in pigmented polymers. *Biochem Biophys Acta* 883: 41–45
- Hirosawa K (1968) Electron microscopic studies on pigment granules in the substantia nigra and locus coeruleus of the Japanese monkey (*Macaca fuscata yuki*). *Z Zellforsch* 88: 187–203
- Hirsch EC, Graybiel AM, Agid YA (1988) Melanized dopaminergic neurons are differentially susceptible to degeneration in Parkinson's disease. *Nature* 334: 345–348
- Hirsch EC, Brandel JP, Galle P, Javoy-Agid F, Agid Y (1991) Iron and aluminum increase in the substantia nigra of patients with Parkinson's disease: an X-ray microanalysis. *J Neurochem* 56: 446–451
- Hirsch EC (1992) Why are nigral catecholaminergic neurons more vulnerable than other cells in Parkinson's disease? *Ann Neurol [Suppl]* 32: S88–S93
- Horcicko J, Borovansky J, Duchon J, Prochazkova B (1973) Distribution of zinc and copper in pigmented tissues. *Z Physiol Chem* 354: 203–204
- Hornykiewicz O (1966) Dopamine (3-hydroxytryptamine) and brain function. *Pharmacol Rev* 18: 925–964
- Hornykiewicz O (1973) Parkinson's disease: from brain homogenate to treatment. *Fed Proc* 32: 183–190
- Hornykiewicz O, Kish SJ (1986) Biochemical pathophysiology of Parkinson's disease. *Adv Neurol* 45: 19–34
- Issidorides MR (1971) Neuronal vascular relationships in the zona compact of normal and parkinsonian substantia nigra. *Brain Res* 25: 289–299
- Ito S, Imai Y, Jimbow K, Fujita K (1988) Incorporation of sulfhydryl compounds into melanins in vivo. *Biochem Biophys Acta* 964: 1–7
- Jellinger K, Paulus W, Grundke-Iqbal I, Riederer P, Youdim MBH (1990) Brain iron and ferritin in Parkinson's and Alzheimer's diseases. *J Neural Transm [P-D Sect]* 2: 327–340
- Jellinger K, Kienzl E, Rumpelmair G, Riederer P, Stachelberger H, Ben-Shachar D, Youdim MBH (1992) Iron-melanin complex in substantia nigra of parkinsonian brains: an X-ray microanalysis. *J Neurochem* 59: 1168–1171
- Jenner P (1992) What process causes nigral cell death in Parkinson's disease? *Neurol Clin* 10: 387–403
- Jenner P, Dexter DT, Sian J, Schapira AHV, Marsden CD (1992) Oxidative stress as a cause of nigral cell death in Parkinson's disease and incidental Lewy body disease. *Ann Neurol [Suppl]* 32: S82–87
- Jimbow K, Quevedo WC, Fitzpatrick TB, Szabo G (1976) Some aspects of melanin biology: 1950–1975. *J Invest Dermatol* 67: 72–89
- Kalyanaraman B, Felix CC, Sealy RC (1984) Peroxidatic oxidation of catecholamines: a kinetic electron spin resonance investigation using the spin stabilization approach. *J Biol Chem* 259: 7584–7589
- Kalyanaraman B, Premovic PI, Sealy RC (1987) Semiquinone anion radicals from addition of amino acids, peptides, and proteins to quinones from oxidation of catechols and catecholamines. *J Biol Chem* 262: 1080–1087
- Kollias N, Sayre RM, Chedekel LZ, Chedekel MR (1991) Photoprotection by melanin. *J Photochem Photobiol B Biol* 9: 135–160
- Korytowski W, Hintz P, Sealy RC, Kalyanaraman B (1985) Mechanism of dismutation of superoxide produced during autooxidation of melanin pigments. *Biochem Biophys Res* 131: 659–665

- Korytowski W, Pilas B, Sarna T, Kalyanarman B (1987) Photoinduced generation of hydrogen peroxide and hydroxyl radicals in melanins. *Photochem Photobiol* 45: 185–190
- Korytowski W, Sarna T (1990) Bleaching of melanin pigments. *J Biol Chem* 265: 12410–12416
- Langston JW, Ballard P, Tetrud JW, Irwin I (1983) Chronic parkinsonism in humans due to a product of meperidine-analog synthesis. *Science* 219: 979–980
- Langston JW, Irwin I, Ricaurte GA (1987) Neurotoxins, parkinsonism, and Parkinson's disease. *Pharmacol Ther* 32: 19–49
- Langston JW (1989) Current theories on the cause of Parkinson's disease. *J Neurol Neurosurg Psychiatry [Spec Suppl]*: 13–17
- Larsen NA, Pakkenberg H, Damsgaard E, Heydorn K (1979) Topographical distribution of arsenic, manganese, and selenium in the human brain. *J Neurol Sci* 42: 407–416
- Larsson B, Tjalve H (1978) Studies on the melanin-affinity of metal ions. *Acta Physiol Scand* 104: 479–484
- Lillie RD, Yamada H (1960) Histochemical studies on the neuromelanins. *Okajimas Folia Anatom Japon* 36: 155–163
- Lindquist NG, Larsson BS, Lyden-Sokolowski A (1987) Neuromelanin and its possible protective and destructive properties. *Pigment Cell Res* 1: 133–136
- Lukiewicz S (1972) The biological role of melanin I. New concepts and methodical approaches. *Folia Histochem Cytochem* 10: 93–108
- Lyden A, Larsson BS, Lindquist NG (1984) Melanin affinity of manganese. *Acta Pharmacol Toxicol* 55: 133–138
- Maeda T, Wegmann R (1969) Infrared spectrometry of locus coeruleus and substantia nigra pigments in human brain. *Brain Res* 14: 673–681
- Mann DMA, Yates PO (1974) Lipoprotein pigments – Their relationship to aging in the human nervous system II. The melanin content of pigmented nerve cells. *Brain* 97: 489–498
- Mann DMA, Yates PO, Barton CM (1977) Neuromelanin and RNA in cells of substantia nigra. *J Neuropathol Exp Neurol* 36: 379–383
- Mann DMA, Yates PO (1979) The effects of ageing on the pigmented nerve cells of the human locus caeruleus and substantia nigra. *Acta Neuropathol* 47: 93–97
- Mann DMA, Yates PO (1983) Possible role of neuromelanin in the pathogenesis of Parkinson's disease. *Mech Age Dev* 21: 193–203
- Marsden CD (1961) Pigmentation in the nucleus substantia nigrae of mammals. *J Anat* 95: 256–261
- Marsden CD (1969) Brain melanin. In: Wolman M (ed) *Pigments in pathology*. Academic Press, New York, pp 395–420
- Marttila RJ, Rinne UK (1989) Oxygen toxicity protecting enzymes in aging and Parkinson's disease. In: Calne DB (ed) *Parkinsonism and aging*. Raven Press, New York, pp 89–95
- Mason HS, Ingram DJE, Allen B (1960) The free radical property of melanins. *Arch Biochem Biophys* 86: 225–230
- Mattock C, Marcot M, Stern G (1988) Could Parkinson's disease follow intrauterine influenza? A speculative hypothesis. *J Neurol Neurosurg Psychiatry* 51: 753–756
- Mikulski T (1970) Electron spin resonance (ESR) and infrared spectra of the eye melanin and of the pigment in substantia nigra. *Acta Physiol Pol* 21: 175–180
- Moses HL, Ganote CE, Beaver DL, Schuffman SS (1966) Light and electron microscopic studies of pigment in human and rhesus monkey substantia nigra and locus coeruleus. *Anat Rec* 155: 167–184
- Nordgren L, Rorsman H, Rosengren AM, Rosengren E (1971) Dopa and dopamine in the pigment of substantia nigra. *Experientia* 27: 1178–1179

- Okazaki M, Kuwata K, Miki Y, Shiga S, Shiga T (1985) Electron spin relaxation of synthetic melanin and melanin-containing human tissue as studied by electron spin echo and electron spin resonance. *Arch Biochem Biophys* 242: 197–205
- Okun MR, Donnellan B, Lever WF (1971) Peroxidase-dependent oxidation of tyrosine or dopa to melanin in neurons. *Histochemie* 25: 289–296
- Olanow CW (1990) Oxidation reactions in Parkinson's disease. *Neurology [Suppl 3]* 40: 32–37
- Olanow CW (1992) Magnetic resonance imaging in parkinsonism. *Neurol Clin* 10: 405–420
- Olanow CW (1992) An introduction to the free radical hypothesis in Parkinson's disease. *Ann Neurol [Suppl]* 32: S2–S9
- Pakkenberg H, Brody H (1965) The number of nerve cells in the substantia nigra in paralysis agitans. *Acta Neuropathol* 5: 320–324
- Parkinson TF, Millikan LE, Anderson DF (1979) Manganese, copper and zinc concentrations in human skin lesions. *Int J Appl Radiat Isotopes* 30: 411–415
- Parkinson Study Group (1993) Effects of tocopherol and deprenyl on the progression of disability in early Parkinson's disease. *N Engl J Med* 328: 176–183
- Pathak MA, Jimbow K, Szabo G, Fitzpatrick TB (1976) Sunlight and melanin pigmentation. In: Smith KC (ed) *Photochemical and photobiological reviews*, vol 1. Plenum Press, New York, pp 211–239
- Perl DP, Good PF (1992) Comparative techniques for determining cellular iron distribution in brain tissues. *Ann Neurol [Suppl]* 32: S76–S81
- Pilas B, Sarna T, Kalyanaraman B, Swartz HM (1988) The effect of melanin on iron associated decomposition of hydrogen peroxide. *Free Radic Biol Med* 4: 285–293
- Poirier J, Kogan S, Gauthier S (1991) Environment, genetics and idiopathic Parkinson's disease. *Can J Neurol Sci* 18: 70–76
- Potts AM, Au PC (1976) The affinity of melanin for inorganic ions. *Exp Eye Res* 22: 487–491
- Riederer P, Sofic E, Rausch W-D, Schmidt B, Reynolds GP, Jellinger K, Youdim MBH (1989) Transition metals, ferritin, glutathione, and ascorbic acid in parkinsonian brains. *J Neurochem* 52: 515–520
- Riederer P, Dirr A, Goetz M, Sofic E, Jellinger K, Youdim MBH (1992) Distribution of iron in different brain regions and subcellular compartments in Parkinson's disease. *Ann Neurol [Suppl]* 32: S101–S104
- Rosengren E, Linder-Eliasson E, Carlsson A (1985) Detection of 5-S-cysteinyl-dopamine in human brain. *J Neural Transm* 63: 247–253
- Rumyantseva GY, Weiner LM, Molin YN, Budker VG (1979) Permeation of liposome membrane by superoxide radical. *FEBS Lett* 108: 477–480
- Sarna T, Hyde JS, Swartz HM (1976) Ion-exchange in melanin. An electron spin resonance study with lanthanide probes. *Science* 192: 1132–1134
- Sarna T, Swartz HM (1978) Identification and characterization of melanin in tissues and body fluids. *Folia Histochem Cytochem* 16: 275–286
- Sarna T, Duleba A, Korytowski W, Swartz H (1980a) Interaction of melanin with oxygen. *Arch Biochem Biophys* 200: 140–148
- Sarna T, Froncisz W, Hyde JS (1980b) Cu^{2+} probe of metal binding sites in melanin using electron paramagnetic resonance spectroscopy II. Natural melanin. *Arch Biochem Biophys* 202: 304–313
- Sarna T, Korytowski W, Pasenkiewicz-Gierula M, Gudowska E (1981) Ion-exchange studies in melanins. In: Seiji M (ed) *Pigment cell 1981: phenotypic expression in pigment cells*. University of Tokyo Press, Tokyo, pp 23–29
- Sarna T, Sealy RC (1984) Photoinduced oxygen consumption in melanin systems. Action spectra and quantum yields for eumelanin and synthetic melanin. *Photochem Photobiol* 39: 69–74

- Sarna T, Pilas B, Land EJ, Truscott TG (1986) Interaction of radicals from water radiolysis with melanin. *Biochim Biophys Acta* 883: 162–167
- Sarna T, Swartz HM (1993) Interactions of melanin with oxygen (and related species). In: Scott G (ed) *Atmospheric oxidation and antioxidants*, vol III. Elsevier, Amsterdam, pp 129–169
- Sarna T (1992) Properties and function of the ocular melanin: a photobiophysical view. *J Photochem Photobiol B* 12: 215–258
- Sealy RC, Felix CC, Hyde JS, Swartz HM (1980) Structure and reactivity of melanins: influence of free radicals and metal ions. In: Prior WA (ed) *Free radicals in biology*, vol 4. Academic Press, New York, pp 209–259
- Sealy RC, Puzyna W, Kalyanaraman B, Felix CC (1984) Identification by electron paramagnetic spectroscopy of free radicals produced during autooxidative melanogenesis. *Biochem Biophys Acta* 800: 269–276
- Shoulson I (1992) Neuroprotective clinical strategies for Parkinson's disease. *Ann Neurol [Suppl]* 32: S143–S145
- Singer TP, Castagnoli N, Ramsay RR, Trevor AJ (1987) Biochemical events in the development of parkinsonism induced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *J Neurochem* 49: 1–8
- Sofic E, Riederer P, Heinsen H, Beckmann H, Reynolds GP, Hebenstreit G, Youdim MBH (1988) Increased iron (III) and total iron content in postmortem substantia nigra of parkinsonian brain. *J Neural Transm* 74: 199–205
- Sofic E, Paulus W, Jellinger K, Riederer P, Youdim MBH (1991) Selective increase of iron in substantia nigra zona compacta of parkinsonian brains. *J Neurochem* 56: 978–982
- Stein WD (1955) Chemical composition of the melanin granule and its relation to the mitochondrion. *Nature* 175: 256–257
- Stern GM (ed) (1990) *Parkinson's disease*. Johns Hopkins University Press, Baltimore
- Swartz HM, Sarna T, Zecca L (1992) Modulation by neuromelanin of the availability and reactivity of metal ions. *Ann Neurol [Suppl]* 32: S69–S75
- Szekeres L (1975) Fine structure and X-ray microanalysis of melanosomes in pigmented nevi and melanomas. *Arch Derm Forsch* 252: 291–304
- Tanner CM (1989) The role of environmental toxins in the etiology of Parkinson's disease. *Trends Neurol Sci* 12: 49–54
- Tanner CM (1992) Epidemiology of Parkinson's disease. *Neurol Clin* 10(2): 317–329
- Tse DCS, McCreery RL, Adams RN (1976) Potential oxidative pathways of brain catecholamines. *J Med Chem* 19: 37–40
- Uitti RJ, Rajput AH, Rozdilsky B, Bickis M, Wollin T, Yuen WK (1989) Regional metal concentrations in Parkinson's disease, other chronic neurological diseases, and control brains. *Can J Neurol Sci* 16: 310–314
- Van Woert MH, Prasad KN, Borg DC (1967) Spectroscopic studies of substantia nigra pigment in human subjects. *J Neurochem* 14: 707–716
- Van Woert MH, Ambani LM (1974) Biochemistry of neuromelanin. *Adv Neurol* 5: 215–223
- Vistnes AI, Henriksen T, Nicolaissen B, Armstrong D (1983) Free radicals and aging. Electron spin resonance studies on neuronal lipopigments and cells grown in vitro. *Mech Age Dev* 22: 335–345
- Warren PJ, Earl CJ, Thompson RHS (1960) The distribution of copper in human brain. *Brain* 83: 709–717
- White LP (1958) Melanin: a naturally occurring cation exchange material. *Nature* 182: 1427–1428
- Wolters EC, Calne DB (1989) Is Parkinson's disease related to aging? In: Calne DB (ed) *Parkinsonism and aging*. Raven Press, New York, pp 125–132
- Youdim MBH, Ben-Shachar D, Riederer P (1989) Is Parkinson's disease a progressive siderosis of substantia nigra resulting in iron and melanin-induced neurodegeneration? *Acta Neurol Scand* 126: 47–54

Zecca L, Mecacci C, Seraglia R, Parati E (1992) The chemical characterization of melanin contained in substantia nigra of human brain. *Biochem Biophys Acta* 1138: 6–10

Authors' address: H. M. Swartz, M.D., Ph.D., 7252 Strassenburgh Hall, Rm 308, Hanover, NH 03755-3863, U.S.A.

Received June 14, 1993