

Inhibition by L-Phenylalanine of Tryptophan Transport by Synaptosomal Plasma Membrane Vesicles: Implications in the Pathogenesis of Phenylketonuria

E. HERRERO, M. C. ARAGON, C. GIMENEZ and F. VALDIVIESO

Departamento de Bioquímica y Biología Molecular, Centro de Biología Molecular, Facultad de Ciencias, Universidad Autónoma de Madrid, Madrid-34, Spain

Phenylalanine is accumulated in the genetically linked deficiency phenylketonuria. The effect of L-phenylalanine on the transport of tryptophan was studied using membrane vesicles from rat-brain synaptosomes. Phenylalanine at similar concentrations to those found in phenylketonuric patients competitively inhibits tryptophan uptake, with a K_i of the same order as the K_m for tryptophan. This inhibition could be responsible for the depletion of serotonin found in phenylketonuria.

Although the basic biochemical defect in phenylketonuria—the genetically linked deficiency of the hepatic phenylalanine hydroxylase system—is well characterized, the mechanisms by which increased phenylalanine causes brain dysfunction are still a subject of research. Several lines of experimental evidence suggest two principal causes for the brain dysfunction in phenylketonuria: defective myelination and impairment in the synthesis of neurotransmitter amines (for a review see Blau, 1979).

In patients with phenylketonuria an abnormal indole metabolism in general, and impairment in the serotonin metabolism in particular, as well as significantly depleted serotonin concentrations, have been found (Vorhees *et al.*, 1981; McKean, 1972). The serotonin depletion, which appears to be related to the high increase of phenylalanine levels, has been proposed as a current cause for the pathogenesis of brain dysfunction in phenylketonuria.

On the other hand, because the concentration of the aromatic amino acid tryptophan, precursor of serotonin in brain cells (McKean *et al.*, 1968; Tyfield and Holton, 1976) is relatively small when compared to the concentration of most other amino acids, it appears that the transport process across plasma membranes might be the rate-limiting step controlling its concentration (Grahame-Smith and Parfitt, 1970; Lajtha, 1974).

In our laboratory we have demonstrated that membrane vesicles derived from brain synaptosomes transport tryptophan by an Na^+ -gradient-dependent system analogous to those previously described for other amino acids in similar preparations (Kanner, 1978; Kanner and Sharon, 1980; Aragon *et al.*, 1981, 1982; Mayor *et al.*, 1981; Marvizon *et al.*, 1981).

The presence of high- and low-affinity transport systems for neurotransmitter amino acids and neurotransmitter precursors has been demonstrated in different preparations of brain (Logan and Snyder, 1972; Kuhar and Zarbin, 1978; Mandell and Knapp,

1977; Aragon *et al.*, 1981). High-affinity Na^+ -dependent systems for tryptophan uptake are thought to be implicated in the maintenance of appropriate tryptophan levels in the nerve cell to ensure serotonin synthesis (Fernston and Wurtman, 1971; Grahame-Smith, 1971), whereas the low affinity system would subservise general metabolic functions (Hedqvist and Stjarne, 1969; Johnston and Iversen, 1971).

The results indicate that phenylalanine at physiological concentrations does inhibit the uptake of tryptophan by membrane vesicles derived from rat brain synaptosomes. This appears to be the most likely cause for the serotonin depletion in phenylketonuria.

MATERIALS AND METHODS

L-[G- ^3H] Tryptophan (sp. radioactivity 3.1 Ci/mmol) was obtained from The Radiochemical Centre, Amersham, Bucks., UK. Ficoll 400 was obtained from Pharmacia. All other materials were of the highest purity available.

Adult male rats of the Wistar strain, weighing 150–200 g, were used. Membrane vesicles were isolated from rat brain essentially as described previously (Kanner, 1978; Aragon *et al.*, 1981).

L-Tryptophan uptake was determined by a filtration technique. Portions (20 μl) of the suspension of membrane vesicles (about 0.1 mg of protein), preloaded with 120 mmol/l KCl–22 mmol/l potassium phosphate buffer, pH 7.4 (KCl medium), were pre-incubated for 1 min at 25°C. Uptake was started by adding 100 μl of a solution containing L-[G- ^3H] tryptophan (5 $\mu\text{mol/l}$ final concentration) in 120 mmol/l NaCl 22 mmol/l sodium phosphate, pH 7.4 (NaCl medium) or in KCl medium. The experiment was terminated by diluting with 5 ml of ice-cold 0.8 mol/l NaCl, and immediately filtering through a moistened Millipore filter, RAWP 02500 (1.2 μm pore size), attached to a vacuum assembly. The filters were rinsed twice with the ice-cold medium

Dilution, filtration and washing procedures were performed within 15 s. The filters were dried at 60°C, placed in microvials and their radioactivity was measured in a liquid-scintillation counter (Beckman LS-350). All the experiments were corrected for a control obtained by diluting the membrane suspension before adding the radioactive substrate solution. All solutions used in the preparation of the membrane vesicles and in the uptake experiments were prepared with distilled and de-ionized water and had been filtered through Millipore filters (0.45 μm) to avoid possible bacterial contamination. The osmolarity of all solutions was kept constant during the experiments. The pH of external and internal medium was 7.4 throughout the experiments.

Membrane proteins were determined according to the method of Resch *et al.* (1971).

RESULTS

Aromatic amino acids interact in their influx into brain cells both *in vivo* (Guroff and Udenfriend, 1962; Blasberg and Lajtha, 1965; McKean *et al.*, 1968) and *in vitro* (Neame, 1961; Pratt, 1976; Barbosa *et al.*, 1970; Vahvelainen and Oja, 1975). Although common mechanisms for transporting aromatic amino acids have been proposed on the basis of this mutual interference, it is questionable whether they share a common transport system (Vahvelainen and Oja, 1975; Lähdesmäki and Hannus, 1977).

The effect of L-phenylalanine on the uptake of L-tryptophan by synaptosomal plasma membrane vesicles from rat brain, has been tested at concentrations similar to those found in the plasma of phenylketonuric patients (Figure 1). As shown in Figure 1A phenylalanine inhibits the tryptophan uptake, both in the presence of a Na^+ gradient, as well as under non-gradient conditions (i.e. KCl medium). Figure 1B shows the effect of phenylalanine on the specific Na^+ -dependent tryptophan uptake. Over the past few years, the existence of two processes of uptake, sodium dependent, in brain preparations, with high and low affinities for tryptophan have been postulated (Grahame-Smith and Parfitt, 1970; Mandell and Knapp, 1977; Laakso and Oja, 1979; Korpi, 1980). The kinetic data of the high-affinity system ($K_m = 35 \mu\text{mol/l}$) in the presence of a Na^+ gradient and 10 and 40 $\mu\text{mol/l}$ L-phenylalanine in the medium are shown in Figure 2. The analysis of these data by Lineweaver-Burk plots demonstrates that L-phenylalanine is a competitive inhibitor for the tryptophan uptake with a K_i of 41 $\mu\text{mol/l}$. Phenylalanine at concentrations of 2.5 mmol/l, also inhibits the low-affinity system ($K_m = 1.46 \text{ mmol/l}$) competitively with a K_i of 1.50 mmol/l (data not shown).

DISCUSSION

Because the membrane vesicles preparation allows the use of a well-defined ionic environment and energy sources and avoids metabolic and compartmentation interference, the nature and kinetic properties of the transport systems for aromatic amino acids in the brain can be studied more effectively. Results support the




Figure 1 Effect of phenylalanine on tryptophan uptake by membrane vesicles from rat brain. The vesicles were preloaded with KCl medium, and incubated as described in the Materials and Methods section in the presence of 5 $\mu\text{mol/l}$ L-[G- ^3H] tryptophan and 0 (○, ●), 10 (□, ■) or 500 (△, ▲) μmol L-phenylalanine in NaCl (○, □, △) or KCl medium (●, ■, ▲). The compositions of the media are described in the Methods section. The specific Na^+ -dependent tryptophan uptake (dashed lines) was obtained by subtracting the uptake in the NaCl medium (○, □, △) from that in the KCl medium (●, ■, ▲) at each time

postulate that phenylalanine and tryptophan have common transport systems (Blasberg and Lajtha, 1965; Sershen and Lajtha, 1979) since they inhibit to a considerable extent each others' influx competitively. The inhibition constants calculated from various different inhibitor concentrations (Figure 2), are of the same order of magnitude as the K_m value. These results are consistent with the idea that the influx of several amino acids will be reduced in children with phenylketonuria (Pratt, 1981). Interference in the transport of tryptophan by a large excess of phenylalanine, as demonstrated in the present report using a membrane vesicles preparation, might be of significance in phenylketonuria where phenylalanine accumulates in the tissues and body fluids of patients. These increased concentrations of phenylalanine, by decreasing the availability of the precursors tryptophan and tyrosine (Aragón *et al.*, 1982), might be the primary cause of serotonin and catecholamine depletion in phenylketonuria (Figure 3).

The authors thank Professor F. Mayor for helpful advice and encouragement. Mrs Mercedes López is gratefully acknowledged for her excellent technical assistance. This work was




Figure 2 Kinetic and double reciprocal plot of initial (30 s) uptake of tryptophan with various concentrations of tryptophan in the presence of 0 (Δ), 10 (\square) or 40 (\circ) $\mu\text{mol/l}$ phenylalanine. The membrane vesicles, preloaded with KCl

medium, were incubated for 30 s as described in the Materials and Methods section in the NaCl medium and radioactive tryptophan at the concentrations indicated. Lineweaver-Burk plots have been generated using the least-squares fitting method.




Figure 3 Phenylalanine interference on the transport of tryptophan and tyrosine in neuron-presynaptic membrane.

supported by grants from the Fondo de Investigaciones Sanitarias.

MS received 11.6.82

Accepted for publication 19.10.82

References

Aragon, M. C., Gimenez, C., Mayor, Jr. F., Marvizon, J. G. and Valdivieso, F. Tyrosine transport by membrane vesicles isolated from rat brain. *Biochim. Biophys. Acta* 646 (1981) 465–470
 Aragón, M. C., Gimenez, C. and Valdivieso, F. Inhibition by phenylalanine of tyrosine transport by synaptosomal plasma membrane vesicles: implications in the pathogenesis

of phenylketonuria. *J. Neurochem.* 39 (1982) 1185–1187
 Barbosa, E., Joanny, P. and Carriol, J. Accumulation active du tryptophane dans le cortex cérébral isolé du rat. *C. R. Soc. Biol. (Paris)* 164 (1970) 345–350
 Blasberg, R. G. and Lajtha, A. Substrate specificity on steady-state amino acid transport in mouse brain slices. *Arch. Biochem.* 112 (1965) 361–377
 Blau, K. Phenylalanine hydroxylase deficiency: Biochemical, physiological, and clinical aspects of phenylketonuria and related phenylalaninemias. In Youdim, M. B. H. (ed) *Aromatic Amino Acid Hydroxylases and Mental Disease*. Wiley, New York, NY (1979), pp. 77–139
 Fernston, J. W. and Wurtzman, R. J. Brain serotonin content: physiological dependence on plasma tryptophan levels. *Science* 173 (1974) 149–152
 Grahame-Smith, D. G. and Parfitt, A. G. Tryptophan transport across the synaptosomal membrane. *J. Neurochem.* 17 (1970) 1339–1352
 Grahame-Smith, D. G. Studies in vivo on the relationship between brain tryptophan, brain 5-HT synthesis, and hyperactivity in rats treated with a monoamine oxidative inhibitor and L-tryptophan. *J. Neurochem.* 18 (1971) 1053–1066
 Guroff, G. and Udenfriend, S. Studies on aromatic amino acid uptake by rat brain *in vivo*. *J. Biol. Chem.* 237 (1962) 803–806
 Hedqvist, P. and Stjarne, L. The relative role of recapture and of 'de novo' synthesis for the maintenance of neurotransmitter homeostasis in noradrenergic nerves. *Acta Physiol. Scand.* 76 (1969) 270–276
 Johnston, G. A. R. and Iversen, L. L. Glycine uptake in rat central nervous systems slices and homogenates: Evidence for different uptake systems in spinal cord and cerebral cortex. *J. Neurochem.* 18 (1971) 1951–1961
 Kanner, B. I. Active transport of α -aminobutyric acid by membrane vesicles isolated from rat brain. *Biochemistry* 17 (1978) 1207–1211
 Kanner, B. I. and Sharon, I. Active transport of L-proline by membrane vesicles isolated from rat brain. *Biochim. Biophys. Acta* 600 (1980) 185–194
 Korpi, E. R. Tryptophan and phenylalanine transport in rat cerebral cortex slices as influenced by sodium ions. *Neurochem. Res.* 5 (1980) 415–431

- Kuhar, M. J. and Zarbin, M. A. Synaptosomal transport: A chloride dependence for choline, GABA, glycine, and several other compounds. *J. Neurochem.* 31 (1978) 251–256
- Laakso, M. L. and Oja, S. S. Transport of tryptophan and tyrosine in rat brain slices in the presence of lithium. *Neurochem. Res.* 4 (1979) 411–423
- Lädhesmäki, P. and Hannus, M.-L. Effect of aromatic acids on the influx of aromatic amino acids in rat brain slices. *Exp. Brain. Res.* 30 (1977) 539–548
- Lajtha, A. Amino acid transport in the brain *in vivo* and *in vitro*. In Wolstenholme, G. E. W. and Fitzsimons, D. W. (eds.) *Aromatic Amino Acids in the Brain*, Elsevier, Amsterdam, 1974, pp. 25–41
- Logan, W. J. and Snyder, S. H. High affinity uptake systems for glycine, glutamic and aspartic acids in synaptosomes of rat central nervous tissues. *Brain. Res.* 42 (1972) 413–431
- Mandell, A. J. and Knapp, S. Regulation of serotonin biosynthesis in brain: role of the high affinity uptake of tryptophan into serotonergic neurons. *Fed. Proc., Fed. Am. Soc. Exp. Biol.* 36 (1977) 2142–2148
- Marvizón, J. G., Mayor, Jr. F., Aragon, M. C., Gimenez, C., and Valdivieso, F. L-Aspartate transport into plasma membrane vesicles derived from rat brain synaptosomes. *J. Neurochem.* 37 (1981) 1401–1406
- Mayor, Jr. F., Marvizón, J. G., Aragon, M. C., Gimenez, C. and Valdivieso, F. Glycine transport into plasma membrane vesicles derived from rat brain synaptosomes. *Biochem. J.* 198 (1981) 535–541
- McKean, C. H., Boggs, D. E. and Peterson, N. A. The influence of high phenylalanine and tyrosine on the concentration of essential amino acids in brain. *J. Neurochem.* 15 (1968) 235–241
- McKean, C. H. The effects of high phenylalanine concentration on serotonin and catecholamine metabolism in the human brain. *Brain Res.* 47 (1972) 469–476
- Neame, K. D. Phenylalanine as inhibitor of transport of amino-acids in brain. *Nature (London)* 192 (1961) 173–174
- Pratt, O. E. The transport of metabolizable substances into the living brain. In Levi, G., Battistin, L. and Lajtha, A. (eds.) *Transport Phenomena in the Nervous System: Physiological and Pathological Aspects*. Advances in Experimental Medicine and Biology. Plenum Press, New York, 1976, pp 55–75
- Pratt, O. E. The needs of the brain for amino acids and how they are transported across the blood–brain barrier. In Belton, N. R. and Toothill, C. (eds.) *Transport and Inherited Disease*. MTP Press, Lancaster, 1981, pp. 87–122
- Resch, K., Imm, W., Ferber, E., Wallach, D. F. M. and Fisher, H. Quantitative determination of soluble and membrane proteins through their native fluorescence. *Naturwissenschaften* 58 (1971) 220
- Sershen, H. and Lajtha, A. Inhibition pattern of analogs indicates the presence of ten or more transport systems for amino acids in brain cells. *J. Neurochem.* 32 (1979) 719–726
- Tyfield, L. A. and Holton, J. B. The effect of high concentrations of histidine on the level of other amino acids in plasma and brain of the mature rat. *J. Neurochem.* 26 (1976) 101–105
- Vahvelainen, M.-L. and Oja, S. S. Kinetic analysis of phenylalanine-induced inhibition in the saturable influx of tyrosine, tryptophan, leucine and histidine into brain cortex slices from adult and 7-day-old rats. *J. Neurochem.* 24 (1975) 885–892
- Vorhees, C. V., Butcher, R. E. and Berry, H. K. Progress in experimental phenylketonuria: a critical review. *Neurosci. Biobehav. Rev.* 5 (1981) 177–190