

## Extractability of nickel and its concentration in cultivated plants in Ni rich ultramafic soils of New Caledonia

Laurent L'Huillier and Sylvie Edighoffer

*Agropedology Laboratory, ORSTOM, B.P. A5, 98 848 Noumea cedex, New-Caledonia\**

Received 31 January 1996. Accepted in revised form 3 September 1996

*Key words:* bioavailability, crop plants, DTPA, maize, Ni uptake, toxicity

### Abstract

The presence of higher-than-normal quantities of nickel is one of the most general features of ultramafic soils and is often suspected as the reason for their infertility. This study on the bioavailability of Ni in ultramafic soils derived from peridotites in New Caledonia showed important variations depending on the position of the soil in the landscape. In piedmont and non hydromorphic colluvio-alluvial soils, Ni was poorly absorbed by cultivated plants. In contrast, crops species grown in the plain soils, especially those found in the colluvio-alluvial and plain soils subject to temporary reducing conditions, possessed very high and even toxic Ni concentrations. Extraction of Ni by DTPA 5 mM was an effective method of estimating Ni bioavailability in these soils. The regression equation developed with only DTPA-extractable Ni explained 88% of the variability in tomato Ni concentration. Extractable Ni might originate from the association of Ni with primary alterable minerals, organic matter and goethite.

### Introduction

Soils derived from ultramafic rocks (ultramafic soils) have been the focus of intensive studies, especially concerning the possible causes of their infertility. Proctor and Woodell (1975) pointed out that identification of the main factor conditioning plant growth in each case was difficult, owing to the great variability of these soils.

Nickel is known to have toxic effects on biological systems, especially on plants (Foy et al., 1978), and its abundance in ultramafic soils is often cited in the literature as an important factor (Robertson, 1992; Soane and Saunder, 1959). The total Ni concentration of ultramafic soils is often greater than 3000  $\mu\text{g g}^{-1}$  (Baker et al., 1992; Brooks, 1987) whereas the majority of soils contain less than 500  $\mu\text{g g}^{-1}$  Ni with a mean between 20 and 40  $\mu\text{g g}^{-1}$  (Uren, 1992).

Many plants found on ultramafic soils accumulate and tolerate high Ni concentrations (Jaffré et al., 1976; Reeves, 1992; Vergnano Gambi et al., 1982) due to physiological adaptations (Gabbrielli et al., 1990; Lee et al., 1978), which suggests that Ni might be relative-

ly bioavailable in these soils. Crops plants grown on these soils could then exhibit a reduction of growth and yield, because they are in general very sensitive to Ni (Davis and Beckett, 1978; Frank et al., 1982; Soane and Saunder, 1959).

Ultramafic soils of New Caledonia possess amongst the highest concentrations of Ni in the world (between 0.5 and 2%). Moreover, the extensive area occupied by these soils (about 1/3 of the country, or 500,000 ha) and the development of agriculture in this region present an interesting model for testing Ni phytotoxicity.

The aim of this study is therefore: (1) to determine whether the problems encountered in crop development on these soils are related to excessive Ni uptake, (2) to test the efficiency of a common single chemical extractant in evaluating the bioavailability of Ni in these soils.

### Materials and methods

#### *Soils*

Sixty surface soils (0–20 cm) were collected in southern New Caledonia in the La Coulee and La Lembi

\* FAX No.: +687264326. E-mail: huillier@noumea.orstom.nc


Table 1. Chemical and physical properties of the ultramafic soils studied. Values are means  $\pm$  S.E. (*n* in parenthesis)

Soil	Particle size distribution (%) <sup>a</sup>			pH (H <sub>2</sub> O) <sup>b</sup>	pH (KCl) <sup>b</sup>	C (%) <sup>c</sup>	N (%) <sup>d</sup>	Exchangeable cations (cmol kg <sup>-1</sup> ) <sup>e</sup>				Total elements (%) <sup>f</sup>				
	Sand	Silt	Clay					Ca	Mg	K	Na	Si	Fe	Ni		
1. Piedmont (n=6)	28.9 $\pm$ 4.6	39.0 $\pm$ 7.2	31.1 $\pm$ 10.7	5.1 $\pm$ 0.1	5.8 $\pm$ 0.1	1.04 $\pm$ 0.29	0.06 $\pm$ 0.01	0.12 $\pm$ 0.05	0.10 $\pm$ 0.06	0.06 $\pm$ 0.03	0.06 $\pm$ 0.04	0.06 $\pm$ 0.04	1.8 $\pm$ 0.4	0.4 $\pm$ 0.2	26.7 $\pm$ 0.3	0.62 $\pm$ 0.06
2. Colluvio-alluvial soil (n=24)	50.6 $\pm$ 6.6	30.9 $\pm$ 3.8	16.8 $\pm$ 5.3	6.2 $\pm$ 0.1	6.3 $\pm$ 0.1	1.40 $\pm$ 0.11	0.11 $\pm$ 0.01	1.50 $\pm$ 0.37	0.18 $\pm$ 0.07	0.04 $\pm$ 0.02	1.36 $\pm$ 0.57	5.7 $\pm$ 0.8	0.8 $\pm$ 0.1	23.9 $\pm$ 0.5	0.51 $\pm$ 0.03	
3. Hydromorphic colluvio-alluvial soil (n=4)	26.1 $\pm$ 5.2	32.5 $\pm$ 6.4	40.1 $\pm$ 10.3	5.9 $\pm$ 0.2	5.5 $\pm$ 0.3	2.71 $\pm$ 0.44	0.20 $\pm$ 0.03	1.75 $\pm$ 0.35	9.62 $\pm$ 2.61	0.40 $\pm$ 0.30	0.35 $\pm$ 0.25	24.4 $\pm$ 7.0	5.9 $\pm$ 1.8	17.4 $\pm$ 2.8	0.91 $\pm$ 0.02	
4. Hydromorphic plain (n=4)	37.4 $\pm$ 9.2	35.4 $\pm$ 8.3	28.3 $\pm$ 6.7	6.0 $\pm$ 0.2	5.8 $\pm$ 0.2	1.42 $\pm$ 0.32	0.11 $\pm$ 0.03	1.05 $\pm$ 0.20	4.70 $\pm$ 1.55	0.20 $\pm$ 0.10	0.25 $\pm$ 0.15	13.8 $\pm$ 2.2	5.7 $\pm$ 0.2	16.8 $\pm$ 0.8	0.89 $\pm$ 0.09	
5. Plain (n=22)	50.4 $\pm$ 8.8	31.9 $\pm$ 7.2	15.5 $\pm$ 3.2	6.6 $\pm$ 0.1	6.4 $\pm$ 0.1	1.47 $\pm$ 0.13	0.11 $\pm$ 0.01	2.13 $\pm$ 0.25	1.07 $\pm$ 0.23	0.16 $\pm$ 0.04	0.68 $\pm$ 0.14	8.4 $\pm$ 0.9	3.5 $\pm$ 0.4	19.9 $\pm$ 0.8	0.85 $\pm$ 0.04	

<sup>a</sup>Sand: 50-2000  $\mu$ m; silt: 2-50  $\mu$ m; clay: <2  $\mu$ m.<sup>b</sup>pH in 1:2.5 soil:water or soil:KCl 1 M suspension.<sup>c</sup>Total organic carbon (Walkley-Black).<sup>d</sup>Total Kjeldahl nitrogen.<sup>e</sup>Exchangeable cations and CEC determined by the Tucker method at pH 7.0<sup>f</sup>HNO<sub>3</sub> 11 N then HClO<sub>4</sub> 6 N (4 h at boiling).

Table 2. Chemical and physical properties of the ultramafic soils used in the greenhouse experiment

Soil	Horizon (cm)	Particle size distribution (%)			pH		C (%)	N (%)	Exchangeable cations ( $\text{cmol kg}^{-1}$ )				CEC ( $\text{cmol kg}^{-1}$ )	Total elements (%)		
		Sand	Silt	Clay	(H <sub>2</sub> O)	(KCl)			Ca	Mg	K	Na		Si	Fe	Ni
Piedmont	0-20	30.8	45.0	24.1	5.1	5.7	1.83	0.09	0.15	0.21	0.13	0.13	2.5	1.0	25.8	0.76
Piedmont	40-60	35.7	47.3	17.0	4.7	6.1	0.42	0.03	0.02	0.01	0.03	0.03	1.9	0.7	25.9	0.70
Plain	0-20	29.1	49.6	21.3	6.6	6.2	2.10	0.15	1.21	4.48	0.45	0.22	12.1	5.7	16.9	0.85
Plain	40-60	20.9	61.4	17.6	6.6	6.2	1.45	0.11	0.56	4.53	0.16	0.13	11.0	6.4	17.8	0.96



- 1 : Piedmont.
- 2 : Colluvio-alluvial soil.
- 3 : Colluvio-alluvial soil with temporary reducing conditions.
- 4 : Plain soil with temporary reducing conditions.
- 5 : Plain (alluvial terrace).

Figure 1. Position in the landscape of the different type of ultramafic soils studied.

valleys (166°36' E-22°14'S) that are the major agricultural areas of the region. They are highly weathered oxisols (Acrorthox) derived from peridotite. Samples were classified in five different soil types (Figure 1) previously described on the basis of differences in their chemistry and physical properties (Becquer et al., 1995; Bourdon and Becquer, 1992). Soils were air dried and sieved through a 2 mm sieve. Their characteristics are given in Table 1.

DTPA, (diethylenetriaminepentaacetic acid) was used as the extractant method for studying the bioavailability of Ni (DTPA-Ni): 5 g of dry soil < 2 mm was mixed in a solution of 25 mL DTPA 5 mM + CaCl<sub>2</sub> 10 mM at a pH of 5.3 for a period of 1 hour (Becquer et al., 1995; after Lindsay and Norvell, 1978). After centrifuging and filtration of solids, Ni was obtained by atomic absorption spectrophotometry AAS (Varian AA300).

#### Maize experiment

A greenhouse experiment with maize (*Zea mays* L. cv. GH 5010) was conducted using a randomized block design with four types of oxisols (piedmont and non hydromorphic plain soils, sampled at depths of 0-20

cm and 40-60 cm) and ten repetitions of each. Soils characteristics are given in Table 2. They were sieved through a 6 mm sieve and homogenised before the filling of the pots (5.1 kg of plain soil per pot; 6.2 kg of piedmont soil per pot). They received an addition of nutrients previously determined to alleviate any deficiencies ( $\text{mg kg}^{-1}$  of soil): 140 N, 1000 P, 69 K, 37 Ca, 23 Mg, 30 S, 0.7 B, 1.1 Zn, 1.6 Cu, 0.2 Mo. After 2 days of germination at 27 °C on cotton moistened with distilled water, the most vigorous maize seedlings were transferred to the pots (one plant by pot). Plants were harvested after 28 days of growth. Water was continually supplied to the pots via the capillary action of fibreglass mesh which was fed by an external water source. The temperature in the greenhouse varied from 24 to 38 °C, the relative humidity from 90 to 50% (day/night), and the photoperiod was 13 hours.

#### Plant sampling and analysis

Plant material (except maize) was sampled in agricultural land from adult crop plants during their fruiting period. Mature non senescent leaves were sampled for chemical analysis. All the soils described above and the plant material were collected in association: the soil was taken just beside the plant.

At harvest, plants were rinsed with demineralised water, dried at 105 °C for 24 h and weighed. For analysis, samples of dried and ground plant material were ashed and digested with concentrated HCl. After filtration, the residue was dried, weighed and treated with hydrofluoric acid to eliminate silica. This element was determined by weight loss of the residue. Cation concentrations obtained after filtration were determined in diluted solutions containing 1% HCl, 0.2% H<sub>3</sub>BO<sub>3</sub> and 0.5% La<sub>2</sub>O<sub>3</sub> by flame emission spectroscopy (K) or AAS (Ca, Mg, Fe, Mn, Ni). Phosphorus was determined colorimetrically at 420 nm according to the vanadomolybdate method (Fiske and Subbarow,

1925). Nitrogen was determined using the Kjeldahl method.

#### Statistical analysis

Results obtained from the greenhouse experiment were examined by analysis of variance. The significant differences between means were analysed by Student t test at the 5% confidence limit. Regression equations were performed on all data using step by step regression available on Statview version 4.02, with significance defined at the 1% confidence level.

### Results

The geochemical nature of the different soils is relatively homogeneous (Tables 1 and 2). However, the colluvio-alluvial and plain soils possess the highest concentrations of Ni, and exceptionally high concentrations of Si and Mg. Levels of these elements are lower in the piedmont soils, but there is a higher concentration of Fe which corresponds to greater alteration.

#### Greenhouse experiment

The maize grown in the test soils exhibited significant differences in growth (Table 3). Maize showed a weaker root and shoot development in plain soils (0-20 cm and especially 40-60 cm: respectively 15% and 39% reduction in shoot dry weight) than in piedmont soils. No particular symptoms were observed during maize development.

Mineral analysis of maize shoots revealed significant differences for many elements between plants grown on the four soils (Table 3). However, variations in growth showed the stronger correlation with shoot Ni concentrations ( $r$  significant at 1%, after Figure 2).

Results of extraction of Ni by DTPA showed similar differences between soils (Table 5): plain soils released much more Ni than piedmont soils, but no significant differences were observed between horizons. Figure 3 shows that the Ni extracted by DTPA correlated relatively well with Ni concentrations in maize shoots.

#### Sampling from agricultural land

Sampling and analysis were carried out on the soils and the associated crops that are grown in the region. They were classified in terms of their position in the land-




Figure 2. Correlation between shoot dry weight and shoot Ni concentration, and between root dry weight and shoot Ni concentration of maize plants grown in the greenhouse experiment. \*\*Significant at 1%.




Figure 3. Correlation between Ni concentration in maize shoots and Ni extracted by DTPA. Data from the greenhouse experiment. \*\*Significant at 1%.

scape (Figure 1) according to Becquer et al. (1995), and Bourdon and Becquer (1992). Important differences were noted in relation to their position: plain and colluvio-alluvial soils with temporary reducing conditions in particular were distinguishable by the highest levels of leaf Ni concentrations in crop plants

Table 3. Dry matter yield and mineral composition of shoots of maize grown on ultramafic soils used in the greenhouse experiment. Values are means  $\pm$ SE ( $n=10$ )

Soil	Yield (mg $\text{plt}^{-1}$ )		Si	N	P	K	Ca	Mg	Fe	Mn	Ni
	Shoot	Root									
Piedmont 0-20 cm	35.1a $\pm 3.9$	2.0a $\pm 0.4$	1.5a $\pm 0.1$	2.8a $\pm 0.3$	0.22a $\pm 0.02$	1.5a $\pm 0.2$	0.54a $\pm 0.05$	0.46a $\pm 0.03$	121a $\pm 32$	115a $\pm 9$	4.8a $\pm 0.6$
Piedmont 40-60 cm	33.0a $\pm 3.0$	2.6a $\pm 0.6$	0.7b $\pm 0.1$	3.0a $\pm 0.4$	0.23a $\pm 0.01$	1.4a $\pm 0.3$	0.68a $\pm 0.08$	0.29b $\pm 0.02$	103a $\pm 26$	73b $\pm 9$	2.9b $\pm 0.3$
Plain 0-20 cm	29.8b $\pm 3.1$	1.4b $\pm 0.3$	2.0c $\pm 0.1$	3.1a $\pm 0.3$	0.38b $\pm 0.05$	2.5b $\pm 0.3$	0.24b $\pm 0.02$	0.64c $\pm 0.04$	134a $\pm 60$	56c $\pm 7$	6.6c $\pm 1.6$
Plain 40-60 cm	21.5c $\pm 2.3$	1.1b $\pm 0.3$	2.1c $\pm 0.1$	3.4a $\pm 0.2$	0.37b $\pm 0.03$	2.5b $\pm 0.3$	0.25b $\pm 0.03$	0.82d $\pm 0.07$	102a $\pm 50$	75b $\pm 6$	12.5d $\pm 1.7$

Different letters in a column indicate significant difference at the 0.05 confidence level.

Table 4. Ni concentrations in the leaves ( $\mu\text{g g}^{-1} \text{ dw}$ ) of some plants cultivated on the ultramafic soils studied. Values are means  $\pm$  SE ( $n$ )

	1. Piedmont	2. Colluvio- alluvial soil	3. Colluvio- alluvial soil with reducing conditions	4. Plain soil with reducing conditions	5. Plain soil
Banana tree				53.5 $\pm$ 11.5(2)	22.1 $\pm$ 11.0(7)
Carrot				59.1 $\pm$ 6.0(2)	41.0 $\pm$ 10.1(3)
Chinese cabbage					48.5 $\pm$ 10.5(2)
Courgette		11.6 $\pm$ 1.2(3)			55.5 $\pm$ 1.5(2)
Eggplant		19.4 $\pm$ 4.6(4)			94.0(1)
Lemon tree		44.0 $\pm$ 2.0(2)	82.1 $\pm$ 16.3(2)		
Mango tree	25.0 $\pm$ 8.6(4)	23.5 $\pm$ 6.5(2)			
Pawpaw tree		14.5 $\pm$ 1.5(2)			
Radish					85.0(1)
Tomato		13.9 $\pm$ 4.8(14)			47.5 $\pm$ 4.7(5)

Table 5. Concentrations in DTPA-extractable Ni (Ni-DTPA) in the ultramafic soils used in the greenhouse experiment or sampled from the field. Values are means  $\pm$  SE ( $n$ )

Greenhouse experiment		Sampling from the field	
Soil	Ni-DTPA ( $\mu\text{g g}^{-1} \text{ soil}$ )	Soil	Ni-DTPA ( $\mu\text{g g}^{-1} \text{ soil}$ )
Piedmont 0-20 cm	18.5 $\pm$ 1.6 ( $n=10$ )	Piedmont	6.9 $\pm$ 3.9 ( $n=6$ )
Piedmont 40-60 cm	1.3 $\pm$ 0.4 ( $n=10$ )	Colluvio-alluvial soil	9.1 $\pm$ 4.5 ( $n=24$ )
Plain 0-20 cm	84.2 $\pm$ 6.7 ( $n=10$ )	Colluvio-alluvial soil with reducing conditions	252.0 $\pm$ 100.0 ( $n=4$ )
Plain 40-60 cm	74.3 $\pm$ 4.1 ( $n=10$ )	Plain soil with reducing conditions	164.5 $\pm$ 40.7 ( $n=4$ )
		Plain	48.8 $\pm$ 18.8 ( $n=22$ )

grown on these types of soil, with Ni concentrations often exceeding  $50 \mu\text{g g}^{-1}$  dw (Table 4). Despite the high levels of fertilization applied by the farmers, crop yields on these soils were much lower than their production potential (about 1.5 to 3 fold lower). Even in the absence of precise information on yields, it appears that there were no important differences between the five types of soil. Only the lemon trees (for the same age and same species *Citrus aurantifolia*) grown in colluvio-alluvial soils with reducing conditions showed chlorotic leaves and significantly lower growth and production than in colluvio-alluvial soils. Except for lemon trees, no particular symptoms were observed in the field.

The same differences as for leaf Ni concentrations between the type of soils were found at the level of DTPA-extractable Ni (Table 5).

In addition, there is a significant correlation between Ni concentrations extracted by DTPA (Ni-DTPA) from the different soils and Ni concentrations in the crop plants examined (Figure 4a). The significance of this correlation is even greater if only tomato plants or eggplants are selected ( $r = 0.95$  and  $0.99$ ; Figures 4 b,c).

Regression equations for the prediction of Ni concentration in tomato leaves could be developed (Table 7). Only equations with all variables significant at 0.01 probability are presented. DTPA-extractable Ni alone accounted for 88% of the variability in tomato Ni concentration.  $R^2$  values were not significantly improved by the inclusion of other soil characteristics in the equation.

## Discussion

The flora of ultramafic soils has been the focus of many studies due to its originality (Brooks, 1987; Lee et al., 1977; Reeves, 1992). In contrast, there is still very little known on the effect of Ni within these soils on plant development (Baker et al., 1992; Robertson, 1992; Uren, 1992).

This study shows the existence of very important variations in Ni bioavailability in the ultramafic soils examined which relate to the position of the soil in the landscape and the soil horizon.

Results of the greenhouse experiment (Table 3) showed that Ni concentrations in maize shoots were all significantly different between the four soils (piedmont and plain soils, for the horizons 0-20 and 40-60 cm). This was also noticed in the growth on the test

plants. The high Ni concentrations recorded for maize grown in the plain soils - especially the horizon 40-60 cm - are probably connected with their poor growth, because they were similar or superior to the concentrations considered toxic in the literature (see Table 6). The strong correlation between the dry weight of roots and shoots and Ni concentrations in maize shoots - stronger than all other elements - reinforce further the hypothesis (Figure 2). However, based on the mineral composition of the maize shoots (Table 3), it appears that four factors might intervene: (1) a low Ca concentration in maize grown in the plain soils. But Ca concentrations of 0.24% are not considered as deficient for maize (Jones, 1967); (2) an excess of Mg, which is often reported for ultramafic soils (Proctor et al., 1981); (3) a strongly imbalanced Mg/Ca ratio; (4) an excess of Ni. It is also possible that combined excess concentrations of Ni and Mg led to a decrease in the growth of maize.

Cultivated plants grown in the field (Table 4) also showed very strong variations in foliar Ni concentration depending on the position of the soil in the landscape. Moreover, leaf Ni concentrations were often very high. In general, crop plants contain less than  $5 \mu\text{g g}^{-1}$  dw Ni in their shoots (Hutchinson and Whitby, 1974; Vanselow, 1966). Nearly all the crops sampled from the plain soils, and especially the colluvio-alluvial and plain formations subject to temporary reducing conditions, possessed Ni concentrations greater than  $40 \mu\text{g g}^{-1}$  dw (Table 4) which are normally considered toxic for most cultivated plants (Table 6). For instance, Ni concentrations found in eggplants and lemon trees cultivated in the plain and colluvio-alluvial soils subject to temporary reducing conditions were probably toxic (compare Tables 4 and 6). This could be connected with the chlorotic leaves and the low growth and production of lemon trees in colluvio-alluvial soils with reducing conditions.

Despite the lack of information in the literature about upper critical levels of Ni for crops, it is likely that Ni plays a significant role in growth of plants on ultramafic soils. However, Ni toxicity is probably limited to certain types of oxisols particular to specific areas in the landscape: plain soils, and especially colluvio-alluvial and plain soils subject to temporary reducing conditions (Figure 1).

The very strong variability in the bioavailability of Ni in relation to the position of the soil in the landscape reinforces the need to establish a method of chemically evaluating Ni bioavailability in these soils. We therefore tested the efficiency of DTPA that is com-




Figure 4. Correlations between DTPA-extractable Ni and shoot Ni concentrations in all crop plants (a), in tomato (b), or in eggplant (c) sampled from the field on ultramafic soils. \*\* Significant at 1%.

monly used (Haq et al., 1980; Lindsay and Norvell, 1978; Sauerbeck and Hein, 1991). For the soils used in the greenhouse experiment, the correlation between Ni concentrations in maize shoots and Ni-DTPA was significant at 1% ( $r=0.74$ ; Figure 3). However, this correlation is dangerous since figure 3 show two different data populations, corresponding to two types of soil very different: plain soils with high Ni-DTPA and piedmont soils with low Ni-DTPA. The difference in Ni uptake by maize between the surface (0-20 cm) and sub-surface (40-60 cm) horizons of plain soils was not

found with the DTPA extraction. The reasons for that are not clear. Nevertheless, a satisfactory correlation was obtained between Ni concentrations in all crop plants sampled in the field and Ni-DTPA ( $r=0.61$ ; Figure 4a). The correlations were particularly strong for tomato plants ( $r=0.94$ ) and eggplants ( $r=0.99$ ; Figures 4b,c).

A very satisfactory result was obtained from the regression equations, whereby Ni extraction by DTPA explained by itself 88% of the variation in Ni concentration in tomato leaves (Table 7). The multiple cor-

Table 6. Toxic Ni concentrations in different crop plants

Plant	Stage	Ni concentration ( $\mu\text{g g}^{-1}$ dw) <sup>a</sup>	Effects	Reference
Alfalfa	83 days	44 (S)	Reduced growth	Halstead et al. (1969)
Barley	control at 5 L	12 (S)	Upper critical level	Davis and Beckett (1978)
Bean	75 days	47 (L)	- 60% seeds weight	Piccini and Malavolta (1992)
Citrus	-	40 (L)	Toxic	Vanselow (1966)
Eggplant	4 months	24 (L)	-16% plant dw	Salim et al. (1988)
Maize	18 days	8.4 (S)	-6% shoots dw	Wallace (1989)
	75 cm at 9 <sup>th</sup> L	12 (S)	Upper critical level	L'Huillier (1994)
Oat	adult	42 (L)	Reduced growth	Hunter and Vergnano (1952)
Soya	18 days	13 (L)	-22% shoots dw	Wallace (1989)
Wheat	23 days	25 (L)	-10% leaves dw	Taylor (1988)

<sup>a</sup> Organ analysed: L: Leaves, S: Shoots.

Table 7. Regression equations for the prediction of Ni concentrations in Tomato developed with DTPA-extractable Ni and other soil characteristics as independent variables ( $n=19$ )

Regression step	$R^2$	Variable added	Final equation		
			Coefficient	Standard error	F
1	0.88	Ni-DTPA (Constant: $12.3 \pm 1.5$ )	0.375	0.035	113.2**

\*\* Significant at the 0.01 confidence level.

relation coefficient ( $R^2$ ) did not significantly improve when other soil characteristics were included into the regression equation. The data from Table 7 can be presented in equation form as follows:

Ni in Tomato ( $\mu\text{g g}^{-1}$  dw) =  $12.3 + 0.375$  Ni-DTPA ( $\mu\text{g g}^{-1}$  soil) ( $R^2 = 0.88$ ,  $n = 19$ )

DTPA is therefore a chemical extractant that by itself is a good indicator of Ni bioavailability and toxicity risks in ultramafic soils. Hughes and Noble (1991) are among the few who have tested chemical extractants in ultramafic soils. They showed that none of the ten single extractants they tested were capable of indicating Ni availability for the flora of ultramafic soils in the Eastern Transvaal. However, the correlations with the vegetation they studied probably lacked some precision and DTPA was not tested.

The regression equations for the prediction of Ni extractability by DTPA (Table 8) allowed us to highlight soil characteristics which may play a significant role in Ni extractability. The levels of CEC, pH(KCl), silica, iron and carbon in the soil explained 89% of the variation in Ni extractability by DTPA with significant probabilities at 1%. This suggests that Ni was more extractable - and probably more bioavailable - in soils with a high level of silica, iron and probably organic

matter (for CEC and carbon). The negative coefficient for pH(KCl) suggests that Ni was more extractable in low pH soils. CEC in these soils mainly originates from organic matter, so this one might be a significant source of bioavailable Ni, especially in the surface horizons of these soils as already shown (Becquer et al., 1995). Silicates might constitute an important source of bioavailable Ni as they can originate from easily weatherable primary minerals (rich in Ni and Si), which can be transported by alluviation and colluviation in colluvio-alluvial and plain soils. Nickel has also been shown to occur in association with iron oxides in mafic soils (Schwertmann and Latham, 1986; Singh and Gilkes, 1992; Uren, 1992) which may represent another significant source of Ni for plants, in particular goethite which is more abundant in plain oxisols (Schwertmann and Latham, 1986). These results are in accordance with Jenne (1968) who showed the important effect of hydrous Fe oxides on Ni availability and emphasised the crucial role of redox potential and pH in determining the availability of these hydrous oxides.

In conclusion, our results show that Ni bioavailability in ultramafic soils is very variable. The levels are low in non hydromorphic colluvio-alluvial and piedmont soils, as previously observed by Angelone et al.



Table 8. Regression equations for the prediction of DTPA-extractable Ni developed with soil characteristics as independent variables ( $n=60$ )

Regression step	R <sup>2</sup>	Variable added	Final equation		
			Coefficient	Standard error	F
1	0.76	CEC	4.8	1.2	16.0**
2	0.81	pH(KCl)	-36.2	10.5	11.8**
3	0.85	Si	31.1	5.6	30.6**
4	0.87	Fe	11.6	2.9	15.8**
5	0.89	C	2.3	0.9	5.6**

(Constant:  $-126.2 \pm 106.4$ )

\*\*Significant at the 0.01 confidence level.

(1993) on ultramafic soils. In contrast, Ni bioavailability is high in plain soils and especially in colluvial and plain soils subject to temporary reducing conditions. Crops cultivated on these soils probably suffer from an excess in Ni uptake. Extraction by DTPA gives a good estimation of Ni bioavailability in ultramafic soils.

The regulations concerning the maximum concentrations of Ni allowed in agricultural soils treated with sewage sludge (30 to 75 mg kg<sup>-1</sup> of dry soil in the EEC; McGrath, 1993) are not practically adaptable to ultramafic soils. It would probably be desirable to establish different limits based on the types of soil, or to take into account the bioavailability of Ni in the soil assessed by a chemical extractant.

Complementary studies are necessary to determine the applicability of DTPA for other soils and also to define the upper critical levels of Ni for many crop species. This would enable a greater understanding of the toxicity risks on ultramafic soils and soils which are in general rich in Ni.

## Acknowledgements

The authors would like to thank the Southern Province of New Caledonia for their financial aid.

## References

- Angelone M, Vaselli O, Bini C and Coradossi N 1993 Pedochemical evolution and trace elements availability to plants in ophiolitic soils. *Sci. Total Environ.* 129, 291–309.
- Baker A J M, Proctor J and Reeves R D 1992 The Vegetation of Ultramafic (Serpentine) Soils. Proceedings of The First International Conference on Serpentine Ecology. University of California, Davis, 19–22 June 1991. Intercept, Andover, UK. 509 p.
- Becquer T, Bourdon E and Pétard J 1995 Disponibilité du nickel le long d'une toposéquence de sols développés sur roches ultramafiques de Nouvelle-Calédonie. *C. R. Acad., Sci. Paris* 321, 585–592.
- Bourdon E and Becquer T 1992 Etude de l'organisation pédologique des sols ferrallitiques des massifs du Sud de la Grande Terre: zones de la Coulée et de la Lembi. Caractérisation physico-chimique des sols. Nouméa, ORSTOM, Agropédol. 16. 88 p.
- Brooks R R 1987 Serpentine and Its Vegetation. A Multidisciplinary Approach. Diocorides Press, Portland, OR.
- Davis R D and Beckett P H T 1978 Upper critical levels of toxic elements in plants. II. Critical levels of copper in young barley, wheat, rape, lettuce and ryegrass, and of nickel and zinc in young barley and ryegrass. *New Phytol.* 80, 23–32.
- Fiske C H and Subbarow Y 1925 The colorimetric determination of phosphorus. *J. Biol. Chem.* 66, 375–400.
- Foy C D, Chaney R L and White M C 1978 The physiology of metal toxicity in plants. *Annu. Rev. Plant Physiol.* 29, 511–566.
- Frank R, Stonefield K I, Suda P and Potter J W 1982 Impact of nickel contamination on the production of vegetables on an organic soil, Ontario, Canada. *Sci. Total Environ.* 26, 41–65.
- Gabbriellini R, Pandolfini T, Vergnano O and Palandri M R 1990 Comparison of two species with different nickel tolerance strategies. *Plant and Soil* 122, 271–277.
- Halstead R L, Finn B J and McLean A J 1969 Extractability of nickel added to the soils and its concentration in plants. *Can. J. Soil. Sci.* 49, 335–342.
- Haq A V, Bates T E and Soon Y K 1980 Comparison of extractants for plant-available Zn, Cd, Ni and Cu in contaminated soils. *Soil Sci. Soc. Am. J.* 44, 772–777.
- Hughes J C and Noble A D 1991 Extraction of chromium, nickel and iron and the availability of chromium and nickel to plants from some serpentinite-derived soils from the eastern Transvaal as revealed by various single and sequential extraction techniques. *Commun. Soil. Sci. Plant Anal.* 22, 1753–1766.
- Hunter J G and Vergnano O 1952 Nickel toxicity in plant. *Ann. Appl. Biol.* 39, 279–284.
- Hutchinson T C and Whitby L M 1974 Heavy metal pollution in the Sudbury mining and smelting region of Canada. I. Soil and vegetation contamination by nickel, copper and other metals. *Environ. Cons.* 1, 123–132.
- Jaffré T, Brooks R R, Lee J and Reeves R D 1976 *Sebertia acuminata*: a hyperaccumulator of nickel from New Caledonia. *Science* 193, 579–580.
- Jenne E A 1968 Controls of Mn, Fe, Co, Ni, Cu, and Zn concentrations in soils and water: the significant role of hydrous Mn and Fe

- oxides. *In* Trace Inorganics in Water. Ed. R F Gould. pp 337–387. Adv. Chem. Series 73, Am. Chem. Soc., Washington, DC, USA.
- Jones J B Jr 1967 Interpretation of plant analysis for several agronomic crops. *In* Soil Testing and Plant Analysis. pp 49–58. Soil Sci. Soc. Am., Special Publ. Series No. 2. SSSA, Madison, WI, USA.
- L'Huillier L 1994 Biodisponibilité du nickel dans les sols ferrallitiques ferritiques de Nouvelle-Calédonie. Effets toxiques de Ni sur le développement et la physiologie du maïs. Ph. D. Thesis, Montpellier II University, Montpellier, France.
- Lee J, Brooks R R, Reeves R D, Boswell C R and Jaffré T 1977 Plant-soil relationships in a New Caledonian serpentine flora. *Plant and Soil* 46, 675–680.
- Lee J, Reeves R D, Brooks R R and Jaffré T 1978 The relation between nickel and citric acid in some nickel-accumulating plants. *Phytochemistry* 17, 1033–1035.
- Lindsay W L and Norvell W A 1978 Development of a DTPA soil test for zinc, iron, manganese and copper. *Soil Sci. Soc. Am. J.* 42, 421–428.
- McGrath S P 1993 Soil quality in relation to agricultural uses. *In* Integrated Soil and Sediment Research: A Basis for Proper Protection. Eds. H J P Eijsackers and T Hamers. pp 187–200. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Piccini D F and Malavolta E 1992 Effect of nickel on two common bean cultivars. *J. Plant Nutr.* 15, 2343–2350.
- Proctor J and Woodell S J 1975 The ecology of serpentine soils. *Adv. Ecol. Res.* 9, 255–366.
- Proctor J, Johnston W R, Cottam D A and Wilson A B 1981 Field-capacity water extracts from serpentine soils. *Nature* 294, 245–246.
- Reeves R D 1992 The hyperaccumulation of nickel by serpentine plants. *In* The Vegetation of Ultramafic (Serpentine) Soils. Eds. A J M Baker, J Proctor and R D Reeves. pp 253–277. Intercept, Andover, UK.
- Robertson A I 1992 The relation of nickel toxicity to certain physiological aspects of serpentine ecology: some facts and a new hypothesis. *In* The Vegetation of Ultramafic (Serpentine) Soils. Eds. A J M Baker, J Proctor and R D Reeves. pp 331–336. Intercept, Andover, UK.
- Salim R, Haddad M and El-Khatib I 1988 Effect of nickel treatment on the growth of eggplant. *J. Environ. Sci. Health* A23, 369–379.
- Sauerbeck D R and Hein A 1991 The nickel uptake from different soils and its prediction by chemical extractions. *Water Air Soil Pollut.* 57–58, 861–871.
- Schwertmann U and Latham M 1986 Properties of iron oxides in some new Caledonian oxisols. *Geoderma* 39, 105–123.
- Singh B and Gilkes R J 1992 Properties and distribution of iron oxides and their association with minor elements in the soils of south-western Australia. *J. Soil Sci.* 43, 77–98.
- Soane B D and Saunderson D H 1959 Nickel and chromium toxicity of serpentine soils in southern Rhodesia. *Soil Sci.* 88, 322–330.
- Taylor G J 1988 Multiple metal stress in *Triticum aestivum*, differentiation between additive, multiplicative, antagonistic and synergistic effects. *Can. J. Bot.* 67, 2272–2276.
- Uren N C 1992 Forms, reactions, and availability of nickel in soils. *Adv. Agron.* 48, 141–203.
- Vanselow A P 1966 Nickel. *In* Diagnostic Criteria for Plants and Soils. Ed. H D Chapman. pp 302–309. University of California, Citrus Research Center and Agric. Experiment. Station, Riverside, USA.
- Vergnano Gambi O, Gabrielli R and Pancaro L 1982 Nickel, chromium and cobalt in plants from Italian serpentine areas. *Acta Oecol. Oecol. Plant* 3, 291–306.
- Wallace A 1989 Effect of liming on trace element interactions in plants. *Soil Sci.* 147, 416–421.

Section editor: L V Kochian