

## e9 Leptospiral Pneumonia

Sunil Karande, Nitin Satam, Madhuri Kulkarni, Renu Bharadwaj<sup>1</sup> and Sae Pol<sup>1</sup>

*Department of Pediatrics, Lokmanya Tilak Municipal Medical College and General Hospital, Sion, Mumbai and*  
*<sup>1</sup>Department of Microbiology, Byramjee Jeejeebhoy Medical College & Sassoon General Hospitals, Pune, India.*

**Abstract.** Severe leptospirosis rarely presents with primary pulmonary manifestations, without any associated jaundice or renal dysfunction. The authors report a nine-year-old boy who presented with complaints of abrupt onset of high fever; with myalgia, headache, and pain in right chest region, productive cough with hemoptysis and vomiting developing over the past 72 hours. Chest radiograph showed consolidation in the right upper lobe with air bronchogram. A history of contact with sewage water and presence of conjunctival suffusion in a child with pneumonia made us suspect leptospirosis. Following prompt initiation of parenteral penicillin therapy the child's complaints resolved over the next five days. Dri-Dot test to detect anti-*Leptospira* antibodies was positive. The diagnosis of leptospirosis was confirmed by a positive microagglutination test to *Leptospira interrogans* serovar Australis by a fourfold rise in antibody titer in paired sera collected during convalescence. Leptospirosis presenting with pulmonary hemorrhage has been associated with significant mortality but it can be successfully treated with early clinical suspicion of alveolar hemorrhage and prompt therapy.

[Indian J Pediatr 2005; 72 (1) : e9-e12] E-mail: karandesunil@yahoo.com

## e13 Cutaneous Tuberculosis Mimicking Sporotrichosis

Suvasini Sharma, Ramesh Choudhary, Monica Juneja, Chander Grover and Belum Siva Nagi Reddy

*Departments of Pediatrics and Dermatology and STD, Maulana Azad Medical College and Associated Lok Nayak Hospital, New Delhi, India.*

**Abstract.** The authors describe an unusual presentation of lupus vulgaris in a 5 year old boy. The lesions had a linear arrangement with proximal spread mimicking sporotrichosis. Histopathology of the lesions revealed tuberculoid granulomas. Tubercular etiology was confirmed by the demonstration of acid fast bacilli in the smears from the regional lymph node aspirate stained with Ziehl Neelsen stain, and growth of *Mycobacterium tuberculosis* in the aspirate culture. The patient showed marked improvement of his lesions on anti-tubercular treatment.

[Indian J Pediatr 2005; 72 (1) : e13-e15] E-mail id: suvasinisharma@rediffmail.com

## e16 Pontine Cavernous Hemangioma Presenting with Horizontal Gaze Palsy

Vimla Menon, Rohit Saxena and Zia Chaudhary

*Dr. Rajendra Prasad Centre for Ophthalmic Sciences, All India Institute of Medical Sciences, New Delhi, India*

**Abstract.** Pontine cavernous hemangioma presenting as horizontal gaze palsy is extremely rare. A 32 years old patient presented with left horizontal gaze palsy with left esotropia. A large pontine mass was present which was removed in toto using a sub-occipital craniotomy. Post-operatively, the gaze palsy showed recovery and the diplopia decreased. Follow-up MRI showed no residual mass. [Indian J Pediatr 2005; 72 (1) : e16-e18] E-mail: rohitsaxena80@hotmail.com

## e19 Undescended Thymus Presenting as Midline Neck Swelling

Shashiraj and Anju Aggarwal

*Department of Pediatrics, University College of Medical Sciences and Guru Tegh Bahadur Hospital, Delhi, India.*

**Abstract.** A seven month male child presented with midline neck swelling which was visible in the suprasternal notch when the child cried. Computed Tomography revealed that it was thymic tissue (solid). There was no thymic tissue in the normal position. Undescended thymus or ectopic thymus is a rare cause of neck mass. Solid type as seen in this case constitute 10% of ectopic thymic tissue, 90% being of cystic variety. Caution should be exercised in excision of such masses as they may be the only thymic tissue. [Indian J Pediatr 2005; 72 (1) : e19-e20] E-mail : a\_anju@vsnl.net

## e21 Waugh's Syndrome

P.L.N.G. Rao and Vijay Kumar

*Kasturba Medical College & Hospital, Manipal, Karnataka, India.*

**Abstract.** A case of Waugh's Syndrome, association of malrotation and intussusception, is being reported. It is suggested that this possibility must be kept in mind and looked for, when a case of intussusception is being treated either by surgery or by hydrostatic reduction. If not when the child develops intestinal problems like obstruction, secondary to malrotation, in the post operative period, it could be misdiagnosed as recurrent intussusception.

[Indian J Pediatr 2005; 72 (1) : e21-e22] E-mail: plngmh@yahoo.com