

On a new unified integral

MRIDULA GARG and SHWETA MITTAL

Department of Mathematics, University of Rajasthan, Jaipur 302 004, India
 E-mail: garghome@satyam.net.in; shwetamittal2003@yahoo.co.in

MS received 15 March 2003; revised 27 November 2003

Abstract. In the present paper we derive a unified new integral whose integrand contains products of Fox H -function and a general class of polynomials having general arguments. A large number of integrals involving various simpler functions follow as special cases of this integral.

Keywords. Fox H -function; general class of polynomials; hypergeometric function.

1. Introduction

The H -function introduced by Fox [1], will be represented and defined in the following manner:

$$\begin{aligned} H_{p,q}^{m,n}[x] &= H_{p,q}^{m,n} \left[x \left| \begin{matrix} (a_1, \alpha_1), \dots, (a_p, \alpha_p) \\ (b_1, \beta_1), \dots, (b_q, \beta_q) \end{matrix} \right. \right] \\ &= \frac{1}{2\pi i} \int_L \frac{\prod_{j=1}^m \Gamma(b_j - \beta_j \xi) \prod_{j=1}^n \Gamma(1 - a_j + \alpha_j \xi)}{\prod_{j=m+1}^q \Gamma(1 - b_j + \beta_j \xi) \prod_{j=n+1}^p \Gamma(a_j - \alpha_j \xi)} x^\xi d\xi. \end{aligned} \quad (1.1)$$

For the nature of contour L in (1.1), the convergence, existence conditions and other details of the H -function, one can refer to [3].

The general class of polynomials introduced by Srivastava [4] is defined in the following manner:

$$S_V^U[x] = \sum_{K=0}^{[V/U]} \frac{(-V)_{UK} A(V, K)}{K!} x^K, \quad V = 0, 1, 2, \dots, \quad (1.2)$$

where U is an arbitrary positive integer and coefficients $A(V, K)$, ($V, K \geq 0$) are arbitrary constants, real or complex.

2. Main result

$$\begin{aligned} &\int_0^\infty x^{\lambda-1} [x + a + (x^2 + 2ax)^{1/2}]^{-v} H_{p,q}^{m,n} [y \{x + a + (x^2 + 2ax)^{1/2}\}^{-\mu}] \\ &\quad \times S_V^U [z \{x + a + (x^2 + 2ax)^{1/2}\}^{-\alpha}] dx \end{aligned}$$

$$\begin{aligned}
&= 2a^{-v} \left(\frac{1}{2}a\right)^\lambda \Gamma(2\lambda) \sum_{K=0}^{\lfloor V/U \rfloor} (-V)_{UK} A(V, K) \frac{(z/a^\alpha)^K}{K!} H_{p+2, q+2}^{m, n+2} \\
&\quad \times \left[ya^{-\mu} \left| \begin{matrix} (-v - \alpha K, \mu), (1 + \lambda - v - \alpha K, \mu), (a_1, \alpha_1), \dots, (a_p, \alpha_p) \\ (b_1, \beta_1), \dots, (b_q, \beta_q), (1 - v - \alpha K, \mu), (-v - \alpha K - \lambda, \mu) \end{matrix} \right. \right],
\end{aligned} \tag{2.1}$$

where

- (i) $\mu > 0, \operatorname{Re}(\lambda, v, \alpha) > 0,$
(ii) $\operatorname{Re}(\lambda) - \operatorname{Re}(v) - \mu \min_{1 \leq j \leq m} \operatorname{Re} \left(\frac{b_j}{\beta_j} \right) < 0.$

Proof. To obtain the result (2.1), we first express Fox H -function involved in its left-hand side in terms of contour integral using eq. (1.1) and the general class of polynomials $S_V^U[x]$ in series form given by eq. (1.2). Interchanging the orders of integration and summation (which is permissible under the conditions stated with (2.1)) and evaluating the x -integral with the help of the result given below [2]:

$$\begin{aligned}
&\int_0^\infty x^{z-1} [x + a + (x^2 + 2ax)^{1/2}]^{-v} dx \\
&= 2va^{-v} \left(\frac{1}{2}a\right)^z [\Gamma(1 + v + z)]^{-1} \Gamma(2z) \Gamma(v - z), \quad 0 < \operatorname{Re}(z) < v,
\end{aligned}$$

we easily arrive at the desired result (2.1).

3. Special case

If in the integral (2.1) we reduce $S_V^U[x]$ to unity and Fox H -function to Gauss hypergeometric function [3], we arrive at the following result after a little simplification:

$$\begin{aligned}
&\int_0^\infty x^{\lambda-1} [x + a + (x^2 + 2ax)^{1/2}]^{-v} \\
&\quad \times {}_2F_1(a, b; c; y(x + a + (x^2 + 2ax)^{1/2})^{-1}) dx \\
&= 2^{1-\lambda} v \Gamma(2\lambda) a^{\lambda-v} \frac{\Gamma(v - \lambda)}{\Gamma(v + \lambda + 1)} \\
&\quad \times {}_4F_3(a, b, v - \lambda, v + 1; c, v, v + \lambda + 1; y/a),
\end{aligned} \tag{3.1}$$

where

$$0 < \operatorname{Re}(\lambda) < \operatorname{Re}(v), |y| < |a|.$$

The importance of the result given by (3.1) lies in the fact that it not only gives the value of the integral but also ‘augments’ the coefficients in the series in the integrand to give a ${}_4F_3$ series as the integrated series.

A number of other integrals involving functions that are special cases of Fox H -function [3] and/or the general class of polynomials [5] can also be obtained from (2.1) but we do not record them here.

Acknowledgements

The authors are thankful to the worthy referee for his very valuable suggestions. The first author is thankful to the University Grants Commission, New Delhi for providing necessary financial assistance to carry out the present work. The authors are thankful to K C Gupta, Jaipur for his useful suggestions.

References

- [1] Fox C, The G and H functions as symmetrical Fourier kernels, *Trans. Am. Math. Soc.* **98** (1961) 395–429
- [2] Oberhettinger F, Tables of Mellin transforms (Berlin, Heidelberg, New York: Springer-Verlag) (1974) p. 22
- [3] Srivastava H M, Gupta K C and Goyal S P, The H -functions of one and two variables with applications (New Delhi and Madras: South Asian Publ.) (1982) p. 11, 18–19
- [4] Srivastava H M, A contour integral involving Fox's H -function, *Indian J.Math.* **14** (1972) 1–6
- [5] Srivastava H M and Singh N P, The integration of certain products of the multivariable H -function with a general class of polynomials, *Rend. Circ. Mat. Palermo*, **32** (1983) 157–187