

THE FUNDAMENTAL SOLUTION OF THE SPACE-TIME FRACTIONAL ADVECTION-DISPERSION EQUATION

F. HUANG AND F. LIU*

ABSTRACT. A space-time fractional advection-dispersion equation (ADE) is a generalization of the classical ADE in which the first-order time derivative is replaced with Caputo derivative of order $\alpha \in (0, 1]$, and the second-order space derivative is replaced with a Riesz-Feller derivative of order $\beta \in (0, 2]$. We derive the solution of its Cauchy problem in terms of the Green functions and the representations of the Green function by applying its Fourier-Laplace transforms. The Green function also can be interpreted as a spatial probability density function (*pdf*) evolving in time. We do the same on another kind of space-time fractional advection-dispersion equation whose space and time derivatives both replacing with Caputo derivatives.

AMS Mathematics Subject Classification: 26A33, 49K20, 44A10.

Key words and phrases: Time-space fractional advection-dispersion equation, Fourier transform, Laplace transform, Mittag-Leffler function, Green function, Caputo derivative, Riesz-Feller derivative.

1. Introduction

An equation commonly used to describe solute transport in aquifers is the advection-dispersion equation(ADE) (Liu et al., 2002, 2003a, 2004):

$$\frac{\partial u}{\partial t} = -\nu \frac{\partial u}{\partial x} + \mathcal{D} \frac{\partial^2 u}{\partial x^2}, \quad (1)$$

where u is solute concentration, the positive constants ν , \mathcal{D} are represent the average fluid velocity and the dispersion coefficient, x is the spatial domain, t is time. The ADE is a deterministic equation describing a probability function for the location of particles in a continuum. The fundamental solutions of the ADE over time will be Gaussian densities with means and variances based on the values of the macroscopic transport coefficients ν and \mathcal{D} .

Received January 9, 2004. Revised October 14, 2004. *Corresponding author.

© 2005 Korean Society for Computational & Applied Mathematics and Korean SIGCAM .

The classical ADE with a local (or asymptotical constant) dispersion tensor is a very handy predictive equation, since solutions are easily gained. The fractional-order forms of the ADE are similarly useful. Some partial differential equations of space-time fractional order were successfully used for modelling relevant physical processes (Mainardi, 1997; Benson et al., 1998,2000; El-Sayed and Aly, 2002; Basu and Acharya, 2002). Numerous authors have shown the equivalence between the transport equations that used fractional-order derivatives and some heavy-tailed motions which extended the predictive capability of models built on the stochastic process of Brownian motion, which is the basis for the classical ADE. The motions can be heavy-tailed, implying extremely long-term correlation and fractional derivatives in time and/or space.

For example, Benson and his collaborator have derived the application of a fractional ADE(see Benson et al., 2000; Benson, 1998; Meerschaert et al., 1999). There are some other authors who considered the fractional ADE. A space-fractional ADE with Eulerian derivation was derived by Schumer et al.(2001), which is used to describe solute plume evolution with a large probability of particles moving significantly ahead of and behind the mean solute velocity.

For example, it can be used in groundwater hydrology to model the transport of passive tracers carried by fluid flow in a porous medium (see Meerschaert and Tadjeran, where a Riemann fractional derivation instead of the Eulerian derivation and a practical numerical method to solve it was developed). Integro-differential equation which interpolates the heat equation and the wave equation is considered by Fujita (1990). He gave the representation of the solution by the fundamental solution and showed some properties. The solution of the bi-fractional differential equation was developed by Kilbas et al. (2004).

The fundamental solution of these problems was established and its moments are calculated. Anh et al. (2001,2002,2003) considered spectral analysis of fractional kinetic equations with random data, renormalization and homogenization of fractional diffusion equations with random data, and harmonic analysis of fractional diffusion-wave equations, respectively. The time-fractional advection-dispersion equation was considered by Liu, Anh et al.(2003b) and its complete solution was obtained by using variable transformation, Mellin and Laplace transforms, and properties of H-functions. We further extend this work and derived its solutions in half-space and a bounded space domain (Huang and Liu, 2004).

This paper is a continuation of these papers. In this paper we intend to consider two kinds of space-time fractional advection-dispersion equations. For both of them, the time fractional derivative is defined in the Caputo sense, while the space fractional derivatives are Riesz-Feller derivatives and Caputo (or Riemann-Liouville) derivatives, respectively.

2. The time-space fractional ADE with Riesz-Feller space derivative

We first consider the equation

$${}_tD_*^\alpha u(x, t) = -\nu D_x u(x, t) + \mathcal{D}D_\theta^\beta u(x, t), \quad x \in \mathbb{R}, \quad t \in \mathbb{R}^+, \quad (2)$$

with the following initial and boundary conditions

$$u(x, 0) = g(x), \quad x \in \mathbb{R}, \quad u(\mp\infty, t) = 0, \quad t > 0, \quad (3)$$

where the real parameters β, θ are always restricted as follows

$$0 < \beta \leq 2, \quad (4)$$

$$|\theta| \leq \min\{\beta, 2 - \beta\}. \quad (5)$$

We denote ${}_tD_*^\alpha$ is the Caputo time-fractional derivative of order α (see Podlubny, 1999; or Appendix B),

D_θ^β is the Riesz-Feller space-fractional derivative of order β (see Appendix A), and $D_x = \frac{\partial}{\partial x}$,

$u(x, t)$ and $g(x)$ are both the (real) field variable, and sufficiently well-behaved functions.

We can consult the literature of Mainardi et al. (2001) for the properties and more details about the Riesz-Feller fractional derivative. By the Fourier-Laplace transform, we represent the solution of the Cauchy problem by the integral formula in terms of the Green functions. The representations of the Green functions also are obtained by a composition rule.

2.1 Some properties of the Green function

Applying Laplace and Fourier transforms to Eq. (2) with initial condition (3) with respect to variable t and x respectively, by taking into account the Laplace transform for the Caputo time fractional derivative (52), and the Fourier transform for the Riesz-Feller space fractional derivative (42), we obtain the following nonhomogeneous differential equation

$$s^\alpha \tilde{u}(x, s) - s^{\alpha-1} g(x) = -\nu D_x \tilde{u}(x, s) + \mathcal{D}D_\theta^\beta \tilde{u}(x, s), \quad (6)$$

$$s^\alpha \hat{\tilde{u}}(k, s) - s^{\alpha-1} \hat{g}(k) = i\nu k \hat{\tilde{u}}(k, s) - \mathcal{D}\psi_\theta^\beta(k) \hat{\tilde{u}}(k, s) \quad (7)$$

where $\psi_\theta^\beta(k) = |k|^\beta e^{i(\text{sign}k)\theta\pi/2}$, and $i^2 = -1$. So we derive

$$\hat{\tilde{u}}(k, s) = \frac{s^{\alpha-1}}{s^\alpha - (i\nu k - \mathcal{D}\psi_\theta^\beta(k))} \hat{g}(k). \quad (8)$$

To invert the Laplace transform in (8), we recall the Laplace transform pair,

$$E_\alpha(ct^\alpha) \xleftrightarrow{\mathcal{L}} \frac{s^{\alpha-1}}{s^\alpha - c}, \quad \Re(s) > |c|^{1/\alpha}, \quad (9)$$

with $c \in \mathcal{C}$, $0 < \alpha \leq 1$, where E_α denotes the entire transcendental function, known as the Mittag-Leffler function of order α , defined in the complex plane by the power series

$$E_\alpha(z) := \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + 1)}, \quad \alpha > 0, \quad z \in \mathcal{C}. \quad (10)$$

For detailed informations on the Mittag-Leffler-type functions and their Laplace transforms, the reader may consult *e.g.* Erdélyi, Magnus and Oberhettinger, pp.1953-1954; Djrbashian, 1966; Podlubny, 1999; Gorenflo and Mainardi, 1997.

Going back to the time domain, we have

$$\widehat{u}(k, t) = E_\alpha \left[(i\nu k - \mathcal{D}\psi_\theta^\beta(k)) t^\alpha \right] \widehat{g}(k). \quad (11)$$

Furthermore, we invert the Fourier transform in above equation to obtain (see Podlubny, 1999)

$$\begin{aligned} u(x, t) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ikx} E_\alpha \left[(i\nu k - \mathcal{D}\psi_\theta^\beta(k)) t^\alpha \right] \widehat{g}(k) dk \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ikx} E_\alpha \left[(i\nu k - \mathcal{D}\psi_\theta^\beta(k)) t^\alpha \right] \int_{-\infty}^{+\infty} e^{iky} g(y) dy dk \\ &= \int_{-\infty}^{+\infty} \left(\frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ik(x-y)} E_\alpha \left[(i\nu k - \mathcal{D}\psi_\theta^\beta(k)) t^\alpha \right] dk \right) g(y) dy \\ &= \int_{-\infty}^{+\infty} G_{\alpha, \beta}^\theta(x-y, t) g(y) dy, \end{aligned} \quad (12)$$

where

$$G_{\alpha, \beta}^\theta(x, t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ikx} E_\alpha \left[(i\nu k - \mathcal{D}\psi_\theta^\beta(k)) t^\alpha \right] dk.$$

It is the Green function (or fundamental solution), which is the formal solution of (2) corresponding to $g(x) = \delta(x)$ (the Dirac delta function). Therefore, we obtain

$$\widehat{G}_{\alpha, \beta}^\theta(k, t) = E_\alpha \left[(i\nu k - \mathcal{D}\psi_\theta^\beta(k)) t^\alpha \right], \quad (13)$$

$$\widehat{\widetilde{G}}_{\alpha, \beta}^\theta(k, s) = \frac{s^{\alpha-1}}{s^\alpha - (i\nu k - \mathcal{D}\psi_\theta^\beta(k))}. \quad (14)$$

In view of the conjugate property of the Mittag-Leffler function $\left(\overline{E_\alpha(z)} = E_\alpha(\overline{z}) \right)$, we have

$$\begin{aligned} \overline{\widehat{G}_{\alpha, \beta}^\theta(-k, t)} &= \overline{E_\alpha \left[(-i\nu k - \mathcal{D}\psi_\theta^\beta(-k)) t^\alpha \right]} \\ &= E_\alpha \left[\overline{(-i\nu k - \mathcal{D}\psi_\theta^\beta(-k)) t^\alpha} \right] \end{aligned}$$

$$\begin{aligned} &= E_\alpha \left[(i\nu k - \mathcal{D}\psi_\theta^\beta(k))t^\alpha \right] \\ &= \widehat{G}_{\alpha,\beta}^\theta(k, t). \end{aligned} \tag{15}$$

Furthermore, we easily recognize

$$\widehat{G}_{\alpha,\beta}^\theta(0, t) = E_\alpha(0) = 1, \quad t \geq 0. \tag{16}$$

Provided that $G_{\alpha,\beta}^\theta(x, t)$ does exist as inverse Fourier transform of (13), equations (15)-(16) ensure that $G_{\alpha,\beta}^\theta(x, t)$ is real and normalized, *i.e.*,

$$G_{\alpha,\beta}^\theta(x, t) \in \mathbb{R}, \quad \int_{-\infty}^{+\infty} G_{\alpha,\beta}^\theta(x, t) = 1.$$

It remains to prove that $G_{\alpha,\beta}^\theta(x, t)$ is positive which ensures that the Green function is the spatial probability density for different values of the relevant parameters α, β . It can be obtained by deriving its explicit representation as indicated below, which is also ensures that $G_{\alpha,\beta}^\theta(x, t)$ does exist as inverse Fourier transform .

2.2 The explicit representation of the Green function

The space fractional advection-dispersion ($\alpha = 1$).

Let us first consider $\alpha = 1, 0 < \beta \leq 2$ (space fractional advection-dispersion including standard advection-dispersion for $\beta = 2$ with $\theta = 0$). In this case, we reduce the Mittag-Leffer function in (13) to the exponential function. To derive the Green function in the space and time domain, we recover the characteristic function of a class of Lévy strictly stable densities $p_\beta^\theta(x)$ (Mainardi, Luchko, and Pagnini, 2001). Then using the notation introduced in Appendix, we write

$$\begin{aligned} \widehat{G}_{1,\beta}^\theta(k, t) &= e^{(i\nu k - \mathcal{D}\psi_\theta^\beta(k))t} = e^{i\nu t k} e^{-\mathcal{D}t\psi_\theta^\beta(k)} \\ &= \widehat{P}_1^1(k; -\nu t) \widehat{P}_\beta^\theta(k; \mathcal{D}t), \end{aligned} \tag{17}$$

where

$$\widehat{P}_\beta^\theta(k; c) = e^{-c\psi_\theta^\beta(k)}, \quad c \in \mathbb{R}. \tag{18}$$

By using the known scaling rule for the Fourier transform,

$$f(ax) \xleftrightarrow{\mathcal{F}} |a|^{-1} \widehat{f}(k/a), \quad a \in \mathbb{R}, \tag{19}$$

we have

$$P_\beta^\theta(x; c) = |c|^{-1/\beta} p_\beta^\theta\left(x/c^{1/\beta}\right), \tag{20}$$

which is non negative. We can consult Appendix A for the stable probability density $p_\beta^\theta(x)$. Inverting the Fourier transform of (17), we have

$$G_{1,\beta}^\theta(x, t) = \int_{-\infty}^{+\infty} P_1^1(x - y; -\nu t) P_\beta^\theta(y; \mathcal{D}t) dy, \tag{21}$$

which is also non negative.

For the standard ADE ($\beta = 2$ with $\theta = 0$), we know

$$p_1^1(x) = \delta(x+1), \quad p_2^0(x) = \frac{1}{2\sqrt{\pi}}e^{-x^2/4}.$$

Then the Green function of standard ADE is

$$G_{1,2}^0(x,t) = \frac{1}{2\nu\sqrt{\mathcal{D}\pi t^3}}e^{-(x+\nu t)^2/4\mathcal{D}t}. \quad (22)$$

Composition rule for the Green function with $0 < \alpha < 1$.

To express the general Green function of the space-time fractional ADE for $0 < \alpha < 1, 0 < \beta \leq 2$, we note that the Fourier-Laplace transform of the Green function (14) can be re-written in integral form as in Saichev and Zaslavsky(1997); Mainardi, Luchko and Pagnini(2001)

$$\begin{aligned} \widehat{\widetilde{G}}_{\alpha,\beta}^\theta(k,s) &= \frac{s^{\alpha-1}}{s^\alpha - (i\nu k - \mathcal{D}\psi_\theta^\beta(k))} \\ &= s^{\alpha-1} \int_0^{+\infty} \exp\left\{-u\left[s^\alpha - (i\nu k - \mathcal{D}\psi_\theta^\beta(k))\right]\right\} du \\ &= \int_0^{+\infty} \exp\left\{u\left[i\nu k - \mathcal{D}\psi_\theta^\beta(k)\right]\right\} \left(s^{\alpha-1}e^{-us^\alpha}\right) du \\ &= \int_0^{+\infty} \widehat{G}_{1,\beta}^\theta(k,u)\widetilde{G}_{2\alpha}(u,s)du, \end{aligned} \quad (23)$$

where

$$\widetilde{G}_\beta(x,s) = s^{\beta/2-1}e^{-|x|s^{\beta/2}}, \quad x \in \mathbb{R}, \quad \Re(s) > 0, \quad (24)$$

with solution

$$G_\beta(x,t) = t^{\beta/2}M_{\beta/2}\left(|x|/t^{\beta/2}\right), \quad x \in \mathbb{R}, \quad t \geq 0, \quad (25)$$

where $M_{\beta/2}$ denotes the so-called M function(of the Wright type) of order $\beta/2$, whose general properties can be found in Podlubny(1999); Gorenflo, Luchko and Mainardi(2000); Mainardi, Luchko and Pagnini(2001).

Going back to the space-time domain we obtain the relation

$$G_{\alpha,\beta}^\theta(x,t) = \int_0^{+\infty} G_{1,\beta}^\theta(x,u)G_{2\alpha}(u,t)du. \quad (26)$$

Then we can derive that the Green function is non negative for the non negative prosperities of $G_{1,\beta}^\theta$ and $G_{2\alpha}$.

3. The space-time fractional ADE with Caputo space derivative

Now we consider the space-time fractional AED

$${}_tD_*^\alpha u(x, t) = -\nu D_x u(x, t) + \mathcal{D}D^\beta u(x, t), \quad x \in \mathbb{R}, \quad t \in \mathbb{R}^+, \quad (27)$$

with the same initial and boundary conditions (3). Where D^β is the Riemann-Liouville or Caputo space fractional derivative of order β with the lower terminal $a = -\infty$, *i.e.*, $D^\beta = {}_{-\infty}D^\beta = {}_{-\infty}^c D^\beta$ (see (51)).

Properties and more details about the Riemann-Liouville and Caputo fractional derivative can be found in texts on Fractional Calculus (Podlubny, 1999; Gorenflo and Mainardi, 1997 or Appendix B). We use the same above technique to obtain the following results (Podlubny, 1999)

$$u(x, t) = \int_{-\infty}^{+\infty} G_{\alpha, \beta}(x - y, t)g(y)dy, \quad (28)$$

where $G_{\alpha, \beta}(x, t)$ is the Green function (or fundamental solution), which is the formal solution of (27) corresponding to $g(x) = \delta(x)$ (the Dirac delta function). The Green function is characterized by

$$\widehat{G}_{\alpha, \beta}(k, t) = E_\alpha \left[\left(\nu((ik)) + \mathcal{D}(-ik)^\beta \right) t^\alpha \right], \quad (29)$$

$$\widehat{\widehat{G}}_{\alpha, \beta}(k, s) = \frac{s^{\alpha-1}}{s^\alpha - \left(\nu((ik)) + \mathcal{D}(-ik)^\beta \right)}. \quad (30)$$

The Green function is also real and normalized, *i.e.*,

$$G_{\alpha, \beta}(x, t) \in \mathbb{R}, \quad \int_{-\infty}^{+\infty} G_{\alpha, \beta}(x, t) = 1.$$

To ensure that the Green function is the spatial probability densities for different values of the relevant parameters α, β , we must further show that the Green function is non negative. In fact, we can also obtain their explicit representation.

First, we consider the space fractional AED($\alpha = 1$). Then

$$\begin{aligned} \widehat{G}_{1, \beta}(k, t) &= e^{[\nu k + \mathcal{D}(-ik)^\beta]t} = e^{\nu tk} e^{(-ik)^\beta \mathcal{D}t} \\ &= \widehat{P}_1(k; -\nu t) \widehat{P}_\beta(k; \mathcal{D}t), \end{aligned} \quad (31)$$

where

$$\widehat{P}_\beta(k; c) = e^{(-ik)^\beta c}, \quad c \in \mathbb{R}. \quad (32)$$

By the same technique as in Huang and Liu(2004b), we obtain

$$P_\beta(x, c) = \begin{cases} (|c|)^{-1/\beta} p_\beta^{-\beta} \left(\frac{x}{(-c)^{1/\beta}} \right), & 0 < \beta \leq 1, \\ (|c|)^{-1/\beta} p_\beta^{2-\beta} \left(\frac{x}{c^{1/\beta}} \right), & 1 < \beta \leq 2, \end{cases} \quad (33)$$

which is non negative where $p_\beta^{-\beta}(x)$ and $p_\beta^{2-\beta}(x)$ are the special stable probability density functions for $p_\beta^\theta(x)$ (see Appendix A). So

$$G_{1,\beta}(x, t) = \int_{-\infty}^{+\infty} P_1(x - y; -\nu t) P_\beta(y; \mathcal{D}t) dy, \quad (34)$$

which is also non negative.

For the case $0 < \alpha < 1$, we can also obtain a composition rule for the Green function by using the same technique as above

$$\widehat{G}_{\alpha,\beta}(k, s) = \int_0^{+\infty} \widehat{G}_{1,\beta}(k, u) \widetilde{G}_{2\alpha}(u, s) du, \quad (35)$$

where \widetilde{G}_β is defined as (24)-(25). So

$$G_{\alpha,\beta}(x, t) = \int_0^{+\infty} G_{1,\beta}(x, u) G_{2\alpha}(u, t) du. \quad (36)$$

Acknowledgements

This research has been supported by the National Natural Science Foundation of China 10271098.

Appendix: Preliminaries

Here we present an introduction to the Riesz-Feller and Caputo fractional derivatives starting from their representation in the Fourier and Laplace transform domain, respectively (Feller, 1952; Mainardi, Luchko and Pagnini, 2001; Gorenflo and Mainardi, 1997; Podlubny, 1999).

Appendix A: The Fourier transform and the Riesz-Feller fractional derivative

Let

$$\widehat{f}(k) = \mathcal{F}\{f(x); k\} = \int_{-\infty}^{+\infty} e^{+ikx} f(x) dx, \quad k \in \mathbb{R}, \quad (37)$$

be the Fourier transform of a well-behaved function $f(x)$, and let

$$f(x) = \mathcal{F}^{-1}\{\widehat{f}(k); x\} = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ikx} \widehat{f}(k) dk, \quad x \in \mathbb{R}, \quad (38)$$

be the inverse Fourier transform.

The Fourier transform of a convolution of two functions states

$$\mathcal{F}\{h(x) * f(x); k\} = \mathcal{F}\left\{\int_{-\infty}^{+\infty} h(x-y)f(y)dy; k\right\} = \widehat{h}(k)\widehat{f}(k). \quad (39)$$

Let

$$0 < \beta \leq 2, \quad |\theta| \leq \begin{cases} \beta & \text{if } 0 < \beta \leq 1, \\ 2 - \beta & \text{if } 1 < \beta \leq 2, \end{cases} \quad (40)$$

and denote by $p_\beta^\theta(x)$ for $x \in R$ the stable probability density whose characteristic function(Fourier transform) is

$$\widehat{p}_\beta^\theta(k) = e^{-\psi_\theta^\beta(k)}, \quad (41)$$

where $\psi_\theta^\beta(k) = |k|^\beta e^{i(\text{sign}k)\theta\pi/2}$, (see Feller(1952) for the general theory of stable probability distributions). The explicit representations and properties of the Riesz-Feller also can be found in Feller(1952); Mainardi, Luchko and Pagnini(2001). The Riesz-Feller fractional derivative of order β and skewness θ defines as

$$\mathcal{F}\{D_\theta^\beta f(x); k\} = -\psi_\theta^\beta(k)\widehat{f}(k), \quad (42)$$

Thus, we recognize that the Riesz-Feller derivative is required to be the pseudo-differential operator whose symbol $-\psi_\theta^\beta(k)$ is the logarithm of the characteristic function of a general Lévy strictly stable probability density with index of stability β and asymmetry parameter θ (improperly called skewness) according to Feller's(1952) parameterizations, as revisited by Gorenflo and Mainardi(1997). In fact, for $0 < \beta < 2$ and $|\theta| \leq \min\{\beta, \beta - 2\}$, the Riesz-Feller derivative reads

$$D_\theta^\beta f(x) = \frac{\Gamma(1 + \beta)}{\pi} \left\{ \sin[(\beta + \theta)\pi/2] \int_{-\infty}^{+\infty} \frac{f(x + \xi) - f(x)}{\xi^{1+\beta}} d\xi + \sin[(\beta - \theta)\pi/2] \int_{\infty}^{+\infty} \frac{f(x - \xi) - f(x)}{\xi^{1+\beta}} d\xi \right\}. \quad (43)$$

and for $\beta = 2$, $D_0^2 = \frac{d^2}{dx^2}$.

Appendix B: The Laplace transform and the Caputo fractional derivative

The Laplace transform of a function $f(t), 0 < t < \infty$, is defined as follows (Mainardi, 1997; Podlubny, 1999):

$$\widetilde{f}(s) = \mathcal{L}\{f(t); s\} = \int_0^\infty e^{-st} f(t) dt, \quad (44)$$

and its inverse is given by the formula

$$f(t) = \mathcal{L}^{-1}\{\widetilde{f}(s); t\} = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{st} \widetilde{f}(s) ds, \quad \gamma = \Re(s) > s_0. \quad (45)$$

It follows that

$$\mathcal{L}\{t^\nu; s\} = \frac{\Gamma(\nu+1)}{s^{\nu+1}}, \quad \nu > -1. \quad (46)$$

The Laplace transform of the convolution of two functions is the product of their Laplace transforms,

$$\mathcal{L}\{h(t) * f(t)\} = \mathcal{L}\left\{\int_0^t h(t-\tau)f(\tau)d\tau; s\right\} = \tilde{h}(s)\tilde{f}(s). \quad (47)$$

Now if f is continuous, the fractional integral of order α of f is

$${}_t D_*^{-\alpha} f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} f(\tau) d\tau, \quad \alpha > 0 \quad (48)$$

which is a convolution integral. From (46)-(48), we have

$$\mathcal{L}\{{}_t D_*^{-\alpha} f(t)\} = \frac{1}{\Gamma(\alpha)} \mathcal{L}\{t^{\alpha-1}\} \mathcal{L}\{f(t)\} = s^{-\alpha} \tilde{f}(s), \quad \alpha > 0. \quad (49)$$

The fractional derivative of order α is defined in the Caputo sense

$${}_t D_*^\alpha f(t) = \begin{cases} \frac{d^n f(t)}{dt^n}, & \alpha = n \in N, \\ \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-\tau)^{n-\alpha-1} \frac{d^n f(\tau)}{d\tau^n} d\tau, & n-1 < \alpha < n. \end{cases} \quad (50)$$

It was proposed by Caputo first in his paper(1967). For the lower terminal $a = -\infty$, we have the relation

$$-{}_\infty D_t^\alpha f(t) = -{}_\infty^c D_t^\alpha f(t). \quad (51)$$

There are two fundamental formulas

$$\mathcal{L}\left\{\frac{d^\alpha}{dt^\alpha} f(t), p\right\} = s^\alpha \tilde{f}(s) - \sum_{k=0}^{n-1} s^{\alpha-1-k} f^{(k)}(0^+), \quad n-1 < \alpha \leq n, \quad n \in N, \quad (52)$$

$$\mathcal{F}\{-{}_\infty D_t^\alpha f(t), k\} = \mathcal{F}\{-{}_\infty^c D_t^\alpha f(t), k\} = (-ik)^\alpha \hat{f}(k). \quad (53)$$

REFERENCES

1. V.V. Anh and N.N. Leonenko, *Spectral analysis of fractional kinetic equations with random data*, J. Stat. Physics, **104**, N5/6 (2001), 1349-1387.
2. V.V. Anh and N.N. Leonenko, *Renormalization and homogenization of fractional diffusion equations with random data*, Probab. Theory Rel. Fields, **124** (2002), 381-408.
3. V.V. Anh and N.N. Leonenko, *Harmonic analysis of fractional diffusion-wave equations*, Applied Math. Comput., **48(3)** (2003), 239-252.
4. M. Basu and D.P. Acharya, *On quadratic fractional generalized solid bi-criterion*, J. Appl. Math. and Computing(old:KJCAM) **2**(2002), 131-144.
5. D.A. Benson, *The fractional advection dispersion equation: Development and application*, Ph.D. thesis, Univ. of Nev. Reno, 1998.

6. D.A. Benson, S.W. Wheatcraft and M.M. Meerschaert, *Application of a fractional advection-dispersion equation*, Water Resources Research, **36**(6) (2000), 1403-1412.
7. M. Caputo, *Linear model of dissipation whose Q is almost frequency independent-II*, Geophys. J. R. Astr. Soc., **13** (1967), 529-539.
8. M.M. Djrbashian, *Integral transforms and representations of functions in the complex plane*, Nauka, 1966(in russian).
9. A.M.A. El-Sayed and M.A.E. Aly, *Continuation theorem of fractional order evolutionary integral equations*, J. Appl. Math. and Computing(ol:KJCAM) **2**(2002), 525-534.
10. A. Erdélyi, W. Magnus, F. Oberhettinger, F.G. Tricomi, *Higher transcendental functions*, **3**, New York, McGraw-Hill, (1953-1954).
11. Y. Fujita, *Integrodifferential equation which interpolates the heat equation and the wave equation*, Osaka, J. Math. **27** (2004), 309-321.
12. A.A. Kilbas, T. Pierantozzi, J. Trujillo, *On the solution of fractional evolution equations*, J. Phys. A: Math. Gen. **37** (2004), 3271-3283.
13. F. Liu, I. Turner and V. Anh, *An unstructured mesh finite volume method for modelling saltwater intrusion into coastal aquifer*, J. Appl. Math. and Computing(ol:KJCAM)**9** (2002), 391-407.
14. F. Liu, I. W. Turner, V. Anh and N. Su, *A two-dimensional finite volume method for transient simulation of time-, scale- and density-dependent transport in heterogeneous aquifer systems*, J. Appl. Math. and Computing **11**(2003a), 215-241.
15. F. Liu, I. W. Turner, V. Anh and P. Zhuang, *Time fractional advection-dispersion equation*, J. Appl. Math. and Computing **13**(2003b), 233-245.
16. F. Liu, V.V. Anh and I. Turner, *Numerical solution of the space fractional Fokker-Plank Equation*, J. Comp. Appl. Math., **166** 2004, 209-319.
17. W. Feller, *On a generalization of Marcel Riesz's potentials and the semigroups generated by them*, Meddeanden lunds Universitets Matematiska Seminarium (Comm. Sé. M.athém. Université de Lund), Tome suppl. dédié à M.Riesz, Lund, (1952), 73-81.
18. R. Gorenflo and F. Mainardi, *Approximation of Lévy-Feller diffusion by random walk*, ZAA, **18** (1999), 231-246.
19. R. Gorenflo, Yu. Luchko and F. Mainardi, *Wright function as scale-invariant solutions of the diffusion-wave equation*, J. Comp. Appl. Math., **118** (2000), 175-191.
20. R. Gorenflo and F. Mainardi, *Fractional calculus: integral and differential equations of fractional order*, in A.Carpinteri and Mainardi(Editors), Fractals and Fractional Calculus in Continuum Mechanics, Wien and New York, Springer Verlag, (1997), 223-273.
21. F. Huang and F. Liu, *The time fractional diffusion equation and advection-dispersion equation*, Submitted to the Australian and New Zealand Industrial and Applied Mathematic Journal(ANZIAM), 2004, in press.
22. F. Mainardi, *Fraction calculus: some basic problems in continuum and statistical mechanics (A. Carpinteri, F. Mainardi, Eds.)*, Fractal and Fractional Calin Continuum Mechanics, Springer, Wien, (1997), 291-348.
23. F. Mainardi, Y. Luchko, G. Pagnini, *The fundamental solution of the space-time fractional diffusion equation*, Fractional Calculus and Applied Analysis, **4** (2001), 153-1925.
24. M.M. Meerschaert, D.A. Benson and B. Bäumer, *Multidimensional advection and fractional dispersion*, Phys. Rev. E. **59**(5), (1999), 5026-5028.
25. M.M. Meerschaert and C.Tadjeeran, *Finite difference approximations for fractional advection-dispersion equations*, unr.edu/homepage/mcubed/frade.pdf.
26. I. Podlubny, *Fractional differential equations*, Academic press, San Diego, 1999.
27. R.Schumer, D.A.Benson, M.M.Meerschaert, S.W.Wheatcraft, *Eulerian derivation of the fractional advection-dispersion equation*, Journal of Contaminant Hydrology **48** (2001), 69-88.

28. A.Saichev and G.Zaslavsky, *Fractional kinetic equations: solutions and applications*, Chaos **7** (1997), 753-764.

Fawang Liu received his MSc from Fuzhou University in 1982 and PhD from Trinity College, Dublin, in 1991. Since graduation, he has been working in computational and applied mathematics at Fuzhou University, Trinity College Dublin and University College Dublin, University of Queensland, Queensland University of Technology and Xiamen University. Now he is a Professor at Xiamen University. His research interest is numerical analysis and techniques for solving a wide variety of problems in applicable mathematics, including semiconductor device equations, microwave heating problems, gas-solid reactions, singular perturbation problem, saltwater intrusion into aquifer systems and fractional differential equations.

Department of Mathematical Sciences, Xiamen University, Xiamen 361005, China
School of Mathematical Sciences, Queensland University of Technology, Qld. 4001, Australia.

e-mail: fwliu@xmu.edu.cn

Fenghui Huang received her MSc and PhD from Xiamen University, Xiamen, China, in 2001 and 2004. Now She is a lecturer at South China University of Technology. Her research interest is numerical computation for PDE, especially, solving the variety of problems in the Computational Fluid Dynamics, such as incompressible fluid flow and turbulence. She also pay respect to some applied problems in fractional differential equations.

Department of Mathematical Sciences, Xiamen University, Xiamen 361005, China
School of Mathematical Sciences, South China University of Technology, Guangzhou 510640, China.

e-mail: huangfh@scut.edu.cn