

Antimutagenic Effect of Plant Flavonoids in the *Salmonella* Assay System

Jae Sue Choi¹, Kun Young Park², Suk Hee Moon², Sook Hee Rhee² and Han Suk Young³

¹Dept. of Food and Nutrition, National Fisheries University of Pusan, Pusan 608-737, ²Dept. of Food Science and Nutrition, Pusan National University, Pusan 608-735 and ³College of Pharmacy, Pusan National University, Pusan 608-735, Korea

(Received August 21, 1993)

The antimutagenic effects of 27 kinds of plant flavonoids on the mutagenicity of aflatoxin B₁(AFB₁) and N-methyl-N'-nitro-N-nitrosoguanidine(MNNG) in *Salmonella typhimurium* TA100 were investigated. In the mixed applications of AFB₁ (1 µg/plate) with the flavonoids (300 µg/plate) in the presence of a mammalian metabolic activation system (S9 mix), chrysin, apigenin, luteolin and its glucoside, kaempferol, fisetin, morin, naringenin, hesperetin, persicogenin, (+)-catechin and (-)-epicatechin showed the antimutagenic effect against AFB₁ with more than 70% inhibition rate. A little or no antimutagenicities except flavone against MNNG (0.5 µg/plate) were observed. For the antimutagenicity of the flavonoids on AFB₁, the flavonoid structure that contains the free 5-, 7-hydroxyl group seemed to be essential. However, saturation of the 2,3-double bond or elimination of the 4-keto group did not affect the activity.

Key words: Antimutagenicity, Flavonoid, *Salmonella* assay system

INTRODUCTION

Flavonoids, which are among the most ubiquitously distributed compounds in the plant kingdom (as well as some mosses, liverworts, fungi and ferns) have been shown to possess a variety of biochemical and pharmacological activities. In addition to their biological activities, several kinds of naturally occurring plant flavonoids have been implicated as genotoxic agents (Bjeldanes *et al.*, 1977; Brown, 1980; Nagao *et al.*, 1981). Since flavonoids occur in all higher plants they are, and always have been, a common constituent of diet. It has been estimated that the "average" daily diet contains about 1 gram of flavonoids (Kuhnau, 1976). Thus the significance of their biological activities with regard to the toxicities has attracted much attention by several workers.

In the course of search for antimutagenic principles from Chinese medicines, we demonstrated that the methanol extract obtained from the whole plants of *Orostachys japonicus* showed the antimutagenic activity (Park *et al.*, 1991a). And also we reported the active principles as flavonoids (Park *et al.*, 1991b). This

kind of result is contradictory to the previous results that flavonoids exhibit the mutagenicities. The results that flavonoids suppress/or reduce the mutagenicity of various chemicals are reported by several workers (Hwang *et al.*, 1983, Francis *et al.*, 1989; Heo, *et al.*, 1992).

Since as naturally occurring substances they are thought to be harmless, we screened the mutagenic and antimutagenic activities of 27 flavonoids isolated from various plants using a microsuspension technique of the *Salmonella*/microsome assay (Ames test). The antimutagenic activity was assayed against mutagens of aflatoxin B₁(AFB₁) and N-methyl-N'-nitro-N-nitrosoguanidine (MNNG). The present report gives data of chemical structures of flavonoids and their antimutagenicity.

MATERIALS AND METHODS

Chemicals

Most of the flavonoids tested were extracted from various plants, purified and characterized according to the previously described procedures (Choi *et al.*, 1986, 1987, 1990).

AFB₁ (Sigma Chemical Co., St. Louis, MO, USA) was dissolved in spectrophotometric grade dimethylsulfo-

Correspondence to: Jae Sue Choi, Dept. of Food and Nutrition, National Fisheries University of Pusan, Pusan 608-737, Korea

xide (DMSO). MNNG (Aldrich Chemical Co. Milwaukee, WI, USA) was dissolved in sterile distilled water. Flavonoid samples (flavone, chrysin, diosmin and hesperidin) were obtained from Sigma Chemical Co., and all of the flavonoids were dissolved in DMSO for the test.

Ames Mutagenicity/antimutagenicity Assay

Salmonella typhimurium strain TA100 histidine requiring mutants were used as bacterial strains. They were maintained as described by Maron and Ames (1983).

S9 mix to activate the mutagen was also prepared by the method of Maron and Ames (1983). Aroclor 1254 in corn oil (200 mg/ml) was injected to male Sprague-Dawley rats (body weight about 200 g) for inducing the liver enzymes. Five days later, liver was excised, homogenized with 0.15 M KCl (3 ml/g liver) using homogenizer (Potter-Elvehjem apparatus, USA) and the homogenate was centrifuged at 9,000 g for 10 min (4°C). Supernatant (S9 fraction) was kept frozen at -80°C until use. The S9 fraction (10%) was mixed with MgCl₂-KCl salts (2%), 1 M glucose 6-phosphate (0.5%), 1 M NADP (4%), 0.2 M phosphate buffer (pH 7.4), and sterilized water prior to use for the Ames assay.

Mutagenicity/antimutagenicity test was carried out by a modified plate incorporation test (liquid preincubation of the organisms with the test compounds) (Matsushima *et al.*, 1980). A 0.5 ml of S9 mix (or 0.5 ml of phosphate buffer for-MNNG) distributed in sterile capped-tubes in an ice bath. Then 0.1 ml of testers from an overnight culture ($1\sim 2\times 10^9$ cells/ml) and 0.1 ml of test compounds (50 µl of mutagen and 50 µl of 0.5~5% (catechins) or 300 µg/plate (all test flavonoids) were added. The tubes were gently vortexed and preincubated at 37°C for 30 min. The top agar (2 ml) kept at 45°C were added to each tube and vortexed for 3 seconds. The entire mixture was overlaid on the minimal agar plate. The plates were incubated at 37°C for 48 hrs and then the revertant bacterial colonies on each plate were counted. Toxicity tests for the different levels of flavonoid samples were also carried out (Maron and Ames, 1983), and the flavonoid concentrations employed for the antimutagenic test in this study did not show any toxicity to the tester strain.

RESULTS

Flavones

Flavone and five related compounds were tested for (anti) mutagenicities and their antimutagenic activities are given in Table I. These compounds except diosmin revealed strong antimutagenic effect against

Table I. Antimutagenic effect of flavones and their glycosides against aflatoxin B₁ (AFB₁, 1 µg/plate) and N-methyl-N'-nitro-N-nitrosoguanidine (MNNG, 0.5 µg/plate) in *Salmonella typhimurium* TA100^a

Compounds	Substitution	Inhibition%	
		AFB ₁	MNNG
Flavone	—	51%	79%
Chrysin	5,7-diOH	90%	-12%
Apigenin	5,7,4'-triOH	88%	6%
Luteolin (Lt)	5,7,3',4'-tetraOH	89%	7%
Cynaroside	Lt-7-O-Glu	80%	-14%
Diosmin	5,3'-diOH,4'-OMe,7-O-Rut	3%	31%

^a 300 µg/plate of flavonoid solutions was employed for the test. The results are the means of triplicate. Glu; glucoside, Rut; rutinoside

Table II. Antimutagenic effect of flavonols and their glycosides against aflatoxin B₁ (AFB₁, 1 µg/plate) and N-methyl-N'-nitro-N-nitrosoguanidine (MNNG, 0.5 µg/plate) in *Salmonella typhimurium* TA100^a


Compounds	Substitution	Inhibition%	
		AFB ₁	MNNG
Kaempferol (kp)	5,7,4'-triOH	75%	14%
Quercetin (Qc)	5,7,3',4'-tetraOH	55%	2%
Fisetin	7,3',4'-triOH	80%	31%
Morin	5,7,2',4'-tetraOH	82%	8%
Populnin	Kp-7-O-Rha	66%	8%
Hyperin	Qc-3-O-Gal	4%	31%
Isoquercitrin	Qc-3-O-Glu	13%	17%
Quercitrin	Qc-3-O-Rha	0%	13%
Avicularin	Qc-3-O-Ara(f)	26%	15%

^a 300 µg/plate of flavonoid solutions was employed for the test. The results are the means of triplicate. Rha; rhamnoside, Gal; galactoside, Ara(f); arabinofuranoside

AFB₁, however, a little or no antimutagenicities except flavone against MNNG were observed.

Flavonols

The antimutagenicities of four flavonols and their five glycosides were tested (Table II). All the flavonols tested showed antimutagenic activity against AFB₁, but little antitagenicity except fisetin against MNNG was observed. Only fisetin which does not have a hydroxyl group at position 5 was antimutagenic at concentrations tested against both AFB₁ and MNNG. Thus, the hydroxyl group at position 5 seems to be important for antimutagenicity in this case. Morin which is the 2'-hydroxy derivative of kaempferol, showed more enhanced antimutagenicity than kaempferol against AFB₁. Therefore, hydroxy groups in the B ring also affect the antimutagenicity. In plants, flavonols are often present as glycosides. Interestingly the quercetin glycosides (hyperin, quercitrin, isoquercitrin, and avicularin)



Scheme 1. Structure and numbering of flavonoids.

were not antimutagenic against AFB₁ or MNNG compared to those of flavonols.

Flavanones and Flavanonols

The antimutagenicities of three flavanones, five flavanone glycosides and two flavanonols were tested (Table III). Among them, naringenin, hesperetin, and persicogenin revealed strong antimutagenic activities against AFB₁. Methylation or glycosylation of the 7-hydroxyl group does affect the antimutagenic potential. However, aromadendrin and taxifolin showed weak antimutagenic effect against AFB₁. They are the 2,3-dihydro derivatives of kaempferol and quercetin, respectively.

(+)-Catechin and (-)-Epicatechin

The antimutagenicities of (+)-catechin and (-)-epicatechin structurally related to flavonoids were tested (Tables IV and V). (+)-Catechin and (-)-epicatechin showed strong antimutagenic activity against AFB₁ with dose-dependant manners. Little or no antimutagenicity was observed against MNNG.

Discussion

In this work, we tested the antimutagenicities of various naturally occurring flavonoid aglycones, flavo-

Table III. Antimutagenic effect of flavanones and their glycosides against aflatoxin B₁ (AFB₁, 1 µg/plate) and N-methyl-N'-nitro-N-nitrosoguanidine (MNNG, 0.5 µg/plate) in *Salmonella typhimurium* TA100^a

Compounds	Substitution	Inhibition%	
		AFB ₁	MNNG
Naringenin (Nr)	5,7,4'-triOH	85%	9%
Hesperetin (He)	5,7,3'-triOH,4'-OH	88%	20%
Persicogenin (Pe)	5,3'-diOH,7,4'-diOMe	71%	15%
Hesperetin 5-Glu	He-5-O-Glu	49%	8%
Hesperidin	He-7-O-Glu	-21%	24%
Prunin	Nr-7-O-Glu	-4%	13%
Naringin	Nr-7-O-Hes	10%	-25%
Persiconin	Pe-5-O-Glu	59%	2%
Aromadendrin	3,5,7,4'-tetraOH	27%	12%
Taxifolin	3,5,7,3',4'-pentaOH	59%	27%

^a 300 µg/plate of flavonoid solutions was employed for the test. The result are the mean of triplicate. Hes; hesperidoside.

Table IV. Effects of (+)-catechin and (-)-epicatechin on the inhibition of aflatoxin B₁ (AFB₁, 1 µg/plate) mutagenicity in *Salmonella typhimurium* TA100

Treatment	Revertants/plate	Inhibition rate (%)
AFB ₁	941 ± 44	
Spontaneous	106 ± 9	
(+)-catechin		
0.5%	598 ± 57	41
1.0%	494 ± 27	54
2.5%	286 ± 27	78
5.0%	195 ± 9	89
AFB ₁	787 ± 18	
Spontaneous	132 ± 7	
(-)-epicatechin		
0.5%	709 ± 20	12
1.0%	546 ± 6	37
2.5%	458 ± 21	50
5.0%	282 ± 42	77

noid glycosides and some synthetic flavonoids. Among the 27 flavonoids tested, flavones and some of their glycoside showed strong antimutagenic effect against AFB₁. Among them only flavone revealed strong antimutagenic activity to both AFB₁ and MNNG.

Although the mutagenicities of various flavonoids have been reported by many workers, Nagao *et al.* (1981) found that flavones having no hydroxyl group at position 3 were not mutagenic to TA100 or TA98 at concentrations of up to 200~300 µg/plate. Thus, absence of the hydroxyl group at position 3 seems to be essential for the antimutagenicity of flavones.

Naturally occurring flavonol aglycones, kaempferol, quercetin, fisetin and morin showed strong antimutagenic effect against AFB₁ in this study, but Pamukcu *et al.* (1980) reported that quercetin was carcinogenic.

Table V. Effects of (+)-catechin and (-)-epicatechin on the inhibition of N-methyl-N'-nitro-N-nitrosoguanidine (MNNG, 0.5 µg/plate) mutagenicity in *Salmonella typhimurium* TA100

Treatment	Revertants/plate	Inhibition rate (%)
MNNG	962 ± 27	
Spontaneous	142 ± 4	
(+)-catechin		
0.5%	938 ± 40	3
1.0%	920 ± 2	5
2.5%	933 ± 52	4
5.0%	888 ± 21	9
(-)-epicatechin		
MNNG	1032 ± 71	
Spontaneous	169 ± 20	
0.5%	1041 ± 168	—
1.0%	1114 ± 131	—
2.5%	1029 ± 44	0
5.0%	1105 ± 119	—

Quercetin was found to induce transformations of all cryopreserved Syrian golden hamsters *in vitro*, sister Chromatid exchange in Chinese hamster cells *in vitro*, and mutation in mouse lymphoma cells L5178Y (Amacher *et al.*, 1980; Sugimura, 1979; Umezawa *et al.*, 1977). Quercetin was also shown to induce intestinal and bladder carcinoma in rats. However, there is a report (Sugimura, 1979) that it was noncarcinogenic in mice, hamsters, and rats. Though the results on quercetin are contradictory, our result showed this compound was antimutagenic toward AFB₁. It seems that dose levels or different experimental systems might cause the differences. In the case of flavonol compounds, glycosylation at position 3 or 7 reduced the activity significantly.

Flavanones and flavanonols which are the 2,3-dihydro derivatives of flavones and flavonols, respectively, revealed strong antimutagenic activities toward AFB₁. Their antimutagenic activities are similar to those of flavones, but they are stronger than flavonols.

As flavonol compounds did, glycosylation at position 7 of flavanones was not antimutagenic. Methylation of the 7-hydroxyl group reduced the activity. However the magnitude was not significant than glycosylation of the 7-hydroxyl group. Glycosylation at position 5 of flavanones also reduced the activity significantly. Thus, the free hydroxyl group at position 5 or 7 seems to be important for the antimutagenicity.

Our results define several structural features essential for the antimutagenic activity of flavonoids in the strain TA100 against AFB₁ and MNNG. From the above data, it appears that the following generalization may be made regarding the structural features essential for the antimutagenicity of flavonoids. Those with a free hydroxyl group at position 5 and 7 were active unless

the 3 position of the C ring was glycosylated. Saturation of the 2,3-double bond or elimination of the 4-keto group does not affect the activity for AFB₁.

Since the precise mechanism of the antimutagenic activity of flavonoids in the bacterial strain system is uncertain, the possible antimutagenic mechanism of flavonoids was demonstrated as inhibitory action on DNA-adduct formation through interaction with microsomal activating enzymes (Francis *et al.*, 1989).

Although Macgregor and Jurd (1978) described that structural features which appear essential for mutagenic activity of the flavonoid in TA98 strain are a basic flavonoid ring structure with (1) a free hydroxyl group at the 3 position, (2) double bond at the 2,3 position, (3) a keto group at the 4 position, and (4) a structure which permits the proton of the 3-hydroxyl group to tautomerise to a 3-keto compound, the essential feature of the hypothesis proposed from our results is that it seems not necessary to have more structural requirement for antimutagenic activity compared to the mutagenicity.

ACKNOWLEDGEMENT

This work was supported by the research grant from KOSEF (90-0500-03).

REFERENCES CITED

- Amacher, D. E., Paillet, S. C., Turner, G. N., Ray, V. A. and Salsburg, D. S., Point mutations at the thymidine kinase locus in L5187Y mouse lymphoma cells. II. Test validation and interpretation. *Mutat. Res.*, 72, 447-449 (1980).
- Bjeldanes, L. F. and Chang, G. W., Mutagenic activities of quercetin and related compounds. *Science*, 197, 577-578 (1977).
- Brown, J. P., A review of genetic effects of naturally occurring flavonoids, anthraquinones and related compounds. *Mutat. Res.*, 75, 243-277 (1980).
- Choi, J. S., Young, H. S., Park, J. C., Choi, J. H. and Woo, W. S., Flavonoids from the leaves of *Rhododendron brachycarpum*. *Arch. Pharm. Res.*, 9, 233-236 (1986).
- Choi, J. S., Young, H. S., Park, J. C., Choi, J. H. and Woo, W. S., Further study on the constituents of *Rhododendron brachycarpum*. *Arch. Pharm. Res.*, 10, 169-172 (1987).
- Choi, J. S., Woo, W. S., Young, H. S. and Park, J. H., Phytochemical study on *Prunus davidiana*. *Arch. Pharm. Res.*, 13, 374-378 (1990).
- Francis, A. R., Shetty, T. K. and Bhattacharya, R. K., Modifying role of dietary factors on the mutagenicity of aflatoxin B₁: *In vitro* effect of plant flavonoids. *Mut. Res.*, 222, 393-401 (1989).

- Heo, M. Y., Yu, K. S., Kim, K. H., Kim, H. P. and Au, W. W., Anticlastogenic effect of flavonoids against mutagen-induced micronuclei in mice. *Mut. Res.*, 284, 243-249 (1992).
- Huang, M. T., Wood, A. W., Newmark, H. L., Sayer, J. M., Yagi, H., Jerina, D. M. and Conney, A. H., Inhibition of the mutagenicity of bay region diol epoxides of polycyclic aromatic hydrocarbons by plant flavonoids. *Carcinogenesis*, 4, 1631-1637 (1983).
- Kunau, J., The flavonoids. A class of semiessential food components: their role in human nutrition. *World Review of Nutrition and Dietetics*, 24, 117-191 (1976).
- Macgregor, J. T. and Jurd, L., Mutagenicity of plant flavonoids: structural requirements for mutagenic activity in *Salmonella typhimurium*. *Mutat. Res.*, 54, 297-309 (1978).
- Maron, D. M. and Ames, B. N., Revised methods for the *Salmonella* mutagenicity test. *Mutation Res.*, 113, 173-215 (1983).
- Matsushima, T., Sugimura, T., Nagao, M., Yahagi, T., Shirai, A. and Sawamura, M., Short term test systems for detecting carcinogens, In Norpoth, K. H. and Garner, R. C. (Eds), *Factors modulating mutagenicity in microbial test*. Springer, Berlin, 1980, pp. 273.
- Nagao, M., Morita, N., Yahagi, J., Shimizu, M., Kuroyanagi, M., Fukuoka, M., Yoshihira, K., Natori, S., Fujino, T. and Sugimura, T., Mutagenicities of 61 flavonoids and 11 related compounds. *Environ. Mut.*, 3, 401-419 (1981).
- Pamukcu, A. M., Yalciner, S., Hatcher, J. F. and Bryan, G. T., Quercetin, a rat intestinal and bladder carcinogen present in bracken fern (*Pteridium aquilinum*). *Cancer Res.*, 40, 3468-3472 (1980).
- Park, H. J., Moon, S. H., Park, K. Y., Choi, J. S., Chung, H. Y., Young, H. S. and Suh, S. S., Antimutagenic effect of *Orostachys japonicus*. *Yakhak Hoeji*, 35, 253-257 (1991a).
- Park, H. J., Young, H. S., Park, K. Y., Rhee, S. H., Chung, H. Y. and Choi, J. S., Flavonoids from the whole plants of *Orostachys japonicus*. *Arch. Pharm. Res.*, 14, 167-171 (1991b).
- Sugimura, T., Naturally occurring carcinogens-mutagens and modulators of carcinogenesis, In Miller, J. A., Miller, E. C., Sugimura, T., Takayama, S. and Hirono, I. (Eds), *Naturally occurring genotoxic carcinogens*. Baltimore, University Park Press, 1979, pp. 241.
- Umezawa, K., Matsushima, T., Sugimura, T., Hirakawa, T., Tanaka, M., Katoh, Y. and Takayama, S., *In vitro* transformation of hamster embryo cells by quercetin. *Toxicol. Lett.*, 1, 175-178 (1977).