

Antitumor and Antiinflammatory Constituents from *Celtis sinensis*

Dae Keun Kim, Jong Pil Lim, Jin Wook Kim, Hee Wook Park, and Jae Soon Eun

College of Pharmacy, Woosuk University, Samrye 565-701, Korea

(Received September 15, 2004)

Eight compounds were isolated from the methanolic extract of the twigs of *Celtis sinensis* through repeated silica gel and Sephadex LH-20 column chromatography. Their chemical structures were elucidated as two triterpenoids, germanicol and epifriedelanol, two amide compounds, *trans-N*-caffeoyltyramine and *cis-N*-coumaroyltyramine, two lignan glycoside, pinoresinol glycoside and pinoresinol rutinoside, and two steroids by spectroscopic analysis.

Key words: *Celtis sinensis*, Germanicol, Epifriedelanol, Amide compounds, Pinoresinol glycoside, Pinoresinol rutinoside, Antitumor, Antiinflammatory

INTRODUCTION

Seltis sinensis Persoon (Ulmaceae) is widely distributed in Korea and has been often used as folk medicine in lumbago, gastric disease, abdominal pain and urticaria eczema (But *et al.*, 1997). Previously, we reported the isolation of *trans-N-p*-coumaroyltyramine as an acetylcholinesterase inhibitor (Kim and Lee, 2003). In the course of our continuing work, two triterpenoids, germanicol and epifriedelanol (antitumor principle, Kundu *et al.*, 2000), two amide compounds, *trans-N*-caffeoyltyramine and *cis-N*-coumaroyltyramine, two lignan glycoside, pinoresinol glycoside and pinoresinol rutinoside (anti-inflammatory principle, Cho *et al.*, 2001), and two steroids were isolated. This paper describes the isolation and structural characterization of these compounds.

MATERIALS AND METHODS

General experimental procedures

¹H- and ¹³C-NMR spectra were determined on a JEOL JMN-EX 400 spectrometer in CDCl₃ and CD₃OD. IR spectra were obtained on a JASCO FT/IR 410 spectrometer and UV spectra were recorded on Shimadzu UV-1601 UV-Visible spectrophotometer. TLC work was carried out using plates coated with silica gel 60 F₂₅₄ (Merck Co.).

Silica gel column chromatography was performed on Merck silica gel 60 (230-400 mesh). Sephadex LH-20 was used for the column chromatography (Pharmacia, 25-100 μm). The column used for LPLC was Lobar-A (Merck Lichroprep Si 60, 240-10 mm). All solvents were routinely distilled prior to use. Other chemicals were commercial grade without purification.

Plant materials

The twigs of *C. sinensis* were collected and air-dried in June 2001 at Bongdong, Chonbuk, Korea. A voucher specimen was deposited in the herbarium of the college of pharmacy, Woosuk University (WSU-01-020).

Extraction and isolation

The shade dried plant material (500 g) was extracted (three times with MeOH at room temperature) and filtered. The filtrate was evaporated *in vacuo* to give a dark brownish residue. The resultant methanolic extract (95 g) was followed by successive solvent partitioning to give *n*-hexane (7 g), CHCl₃ (15 g), EtOAc (5 g), *n*-BuOH (30 g) and H₂O soluble fractions. *n*-Hexane soluble fraction was chromatographed on silica gel column (*n*-hexane-CHCl₃-EtOAc, 12:1:1) to give four fractions (H1-H4). Fraction H3 was purified by LPLC (Lobar A column, *n*-hexane-CHCl₃-EtOAc, 10:1:1) to give **1** (8 mg). Silica gel column chromatography of the CHCl₃ soluble fraction with CHCl₃-EtOAc-MeOH (10:1:1) gave six fractions (C1-C6). The major fraction C1 was chromatographed on silica gel column using CHCl₃-EtOAc (40:1) to yield five fractions

Correspondence to: Dae Keun Kim, College of Pharmacy, Woosuk University, Samrye 565-701, Korea
Tel: 82-63-290-1574, Fax: 82-63-290-1567
E-mail: dkkim@mail.woosuk.ac.kr

(C11-C15). Fraction C11 was purified by Sephadex LH-20 column with MeOH to give **2** (10 mg). Recrystallization of the fraction C15 with *n*-hexane-acetone (10:1) gave **3** (15 mg). Silica gel column chromatography of the EtOAc soluble fraction with CHCl₃-EtOAc-MeOH (10:1:1) gave five fractions (E1-E5). The major fraction E2 was rechromatographed on the Sephadex LH-20 column (MeOH) and purified by Lobar-A column chromatography (CHCl₃-MeOH, 7:1) to yield compound **4** (7 mg), and **5** (9 mg). Recrystallization of the fraction E4 with MeOH gave **6** (15 mg). 15 g of *n*-BuOH soluble fraction was chromatographed over Sephadex LH-20 column with MeOH to give four fractions (B1-B4). The major fraction B2 was chromatographed on silica gel column using CHCl₃-MeOH-H₂O (50:10:1) to yield four fractions (B21-B24). Fraction B22 was purified by Lobar-A column (CHCl₃-MeOH, 5:1) to give **7** (11 mg). Fraction B24 was purified by Lobar-A column (MeOH) to give **8** (11 mg).

Germanicol (**1**) (Koch *et al.*, 2005)

mp 173-174°C; ¹H-NMR (400 MHz, DMSO-*d*₆) δ: 4.85 (1H, s, H-19), 3.20 (1H, dd, *J*=11.2, 4.8 Hz, H-3α), 1.08, 1.02, 0.97, 1.88, 0.77, 0.73 (each 3H, s, CH₃), 0.94 (6H, s, 2×CH₃). ¹³C-NMR (100 MHz, DMSO-*d*₆) δ: 142.7 (C-18), 129.7 (C-19), 79.0 (C-3), 55.5 (C-5), 51.2 (C-9), 43.3 (C-14), 40.7 (C-8), 38.9 (C-4), 38.9 (C-13), 38.4 (C-1), 37.7 (C-16), 37.4 (C-22), 37.2 (C-10), 34.6 (C-7), 34.3 (C-17), 33.3 (C-21), 32.4 (C-20), 31.4 (C-29), 29.2 (C-30), 28.0 (C-23), 27.5 (C-15), 27.4 (C-2), 26.2 (C-12), 25.3 (C-28), 21.1 (C-11), 18.2 (C-6), 16.7 (C-26), 16.1 (C-25), 15.4 (C-24), 14.6 (C-27).

Epifriedelanol (**2**) (Matsunaga *et al.*, 1993)

mp. 277-278°C; ¹H-NMR (400 MHz, CDCl₃) δ: 3.74 (1H, m, H-3), 1.17, 1.00, 0.99, 0.98, 0.96, 0.86 (each 3H, s, CH₃), 0.95 (6H, s, 2×CH₃). ¹³C-NMR (100 MHz, CDCl₃) δ: 72.8 (C-3), 61.3 (C-10), 53.2 (C-8), 49.1 (C-4), 42.8 (C-18), 41.7 (C-6), 39.7 (C-13), 39.3 (C-22), 38.4 (C-14), 37.8 (C-9), 37.1 (C-5), 36.1 (C-16), 35.5 (C-11), 35.3 (C-19), 35.2 (C-29), 35.0 (C-2), 32.8 (C-21), 32.0 (C-15), 32.1 (C-28), 31.85 (C-30), 30.6 (C-12), 29.7 (C-17), 28.2 (C-20), 20.1 (C-26), 18.7 (C-27), 18.2 (C-25), 17.5 (C-1), 16.4 (C-24), 15.8 (C-7), 11.6 (C-23).

β-Sitosterol (**3**)

mp 277-278°C; ¹H-NMR (400 MHz, CD₃OD) δ: 3.74 (1H, m, H-3), 1.17, 1.00, 0.99, 0.98, 0.96, 0.86 (each 3H, s, CH₃), 0.95 (6H, s, 2×CH₃). ¹³C-NMR (100 MHz, CD₃OD) δ: 72.8 (C-3), 61.3 (C-10), 53.2 (C-8), 49.1 (C-4), 42.8 (C-18), 41.7 (C-6), 39.7 (C-13), 39.3 (C-22), 38.4 (C-14), 37.8 (C-9), 37.1 (C-5), 36.1 (C-16), 35.5 (C-11), 35.3 (C-19), 35.2 (C-29), 35.0 (C-2), 32.8 (C-21), 32.0 (C-15), 32.1 (C-28), 31.85 (C-30), 30.6 (C-12), 29.7 (C-17), 28.2

(C-20), 20.1 (C-26), 18.7 (C-27), 18.2 (C-25), 17.5 (C-1), 16.4 (C-24), 15.8 (C-7), 11.6 (C-23).

trans-*N*-Caffeoyltyramine (**4**)

mp 216-217°C; ¹H-NMR (400 MHz, CD₃OD) δ: 7.33 (1H, d, *J*=15.6 Hz, H-7), 7.02 (1H, d, *J*=1.5 Hz, H-2), 7.00 (2H, d, *J*=8.4 Hz, H-2', 6'), 6.95 (1H, dd, *J*=8.4, 1.5 Hz, H-6), 6.69 (1H, d, *J*=8.4 Hz, H-5), 6.62 (2H, d, *J*=8.4 Hz, H-3', 5'), 6.30 (1H, d, *J*=15.6 Hz, H-8), 3.38 (2H, t, *J*=7.5 Hz, H-8), 2.66 (2H, t, *J*=7.5 Hz, H-7), ¹³C-NMR (100 MHz, CD₃OD) δ: 167.2 (C-9), 155.6 (C-4'), 147.5 (C-4), 145.5 (C-3), 141.0 (C-7), 131.0 (C-1), 130.7 (C-2', 6'), 127.1 (C-1), 121.0 (C-6), 117.3 (C-8), 116.3 (C-5), 116.0 (C-3', 5'), 114.1 (C-2), 42.0 (C-8'), 34.0 (C-7').

cis-*N*-Coumaroyltyramine (**5**)

mp 234-235°C; ¹H-NMR (400 MHz, CD₃OD) δ: 7.26 (2H, d, *J*=8.8 Hz, H-2', 6'), 6.91 (2H, d, *J*=8.4 Hz, H-2, 6), 6.65 (2H, d, *J*=8.8 Hz, H-3', 5'), 6.62 (2H, d, *J*=8.4 Hz, H-3, 5), 6.51 (1H, d, *J*=12.8 Hz, H-8), 5.69 (1H, d, *J*=12.8 Hz, H-7), 3.29 (2H, t, *J*=7.5 Hz, H-8'), 2.59 (2H, t, *J*=7.5 Hz, H-7'), ¹³C-NMR (100 MHz, CD₃OD) δ: 170.4 (C-9), 159.4 (C-4), 156.9 (C-4'), 138.1 (C-7), 132.3 (C-2', 6'), 131.2 (C-1'), 130.7 (C-2, 6), 127.9 (C-1), 121.4 (C-8), 116.2 (C-3, 5), 116.0 (C-3, 5), 42.3 (C-8'), 35.5 (C-7').

β-Sitosterol-3-*O*-glucoside (**6**)

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 5.42 (1H, m, H-6), 4.95 (1H, d, *J*=5.0, H-1'), 3.80-3.00 (sugar H), 1.05 (3H, s, 19-CH₃), 1.03-0.85 (12H, 21, 29, 27, 26-CH₃), 0.71 (3H, s, 18-CH₃). ¹³C-NMR (100 MHz, DMSO-*d*₆) δ: 140.4 (C-5), 121.3 (C-6), 76.8 (C-3), 56.1 (C-14), 55.3 (C-17), 49.6 (C-9), 45.1 (C-24), 41.8 (C-13), 38.3 (C-12), 36.8 (C-1), 36.2 (C-10), 35.4 (C-20), 33.3 (C-22), 31.4 (C-7), 31.3 (C-8), 29.2 (C-2), 28.7 (C-25), 27.7 (C-16), 25.4 (C-23), 23.7 (C-15), 22.6 (C-28), 20.6 (C-11), 19.7 (C-26), 19.1 (C-19), 18.9 (C-27), 18.6 (C-21), 11.7 (C-29), 11.6 (C-18), 100.7 (C-1'), 76.8 (C-3'), 76.6 (C-5'), 73.4 (C-2'), 70.1 (C-4'), 61.0 (C-6').

Pinosresinol-4-*O*-glucoside (**7**)

¹H-NMR (400 MHz, CD₃OD) δ: 7.13 (1H, d, *J*=8.4 Hz, H-5), 7.01 (1H, d, *J*=1.6 Hz, H-2), 6.92 (1H, d, *J*=1.6 Hz, H-2'), 6.90 (1H, dd, *J*=8.4, 1.6 Hz, H-6), 6.80 (1H, dd, *J*=8.4, 1.6 Hz, H-6'), 6.76 (1H, d, *J*=8.4 Hz, H-5'), 4.87 (1H, d, *J*=7.8 Hz, H-1"), 4.74 (1H, d, *J*=4.0 Hz, H-7), 4.68 (1H, d, *J*=4.4 Hz, H-7'), 4.13 (2H, m, H-9β, 9'β), 3.85 (3H, s, OCH₃), 3.84 (3H, s, OCH₃), 3.82 (2H, m, H-9α, 9'α), 3.67-3.18 (sugar H), 3.11 (2H, m, H-8, 8'), ¹³C-NMR (100 MHz, CD₃OD) δ: 150.9 (C-4), 149.1 (C-4), 147.5 (C-3'), 147.3 (C-3), 137.4 (C-1), 133.7 (C-1'), 120.0 (C-6'), 119.8 (C-6), 118.0 (C-5), 116.1 (C-5'), 111.6 (C-2), 110.9 (C-2'), 102.8 (C-1"), 87.5 (C-7'), 87.1 (C-7), 78.2 (C-5"), 77.8 (C-3"), 74.9 (C-2"), 72.7 (C-9'), 72.7 (C-9), 71.3 (C-4"), 62.5 (C-6"),

56.7 (C-OCH₃), 56.4 (C-OCH₃), 55.5 (C-8'), 55.3 (C-8).

Pinoresinol-4-O-rutinoside (8)

UV λ_{\max} (MeOH) nm: 280, 235, IR ν_{\max} (KBr) cm⁻¹: 3350, 1630, 1520, FABMS: m/z 689.0447 (M+Na), ¹H-NMR (400 MHz, CD₃OD) δ : 7.05 (1H, d, $J=8.4$ Hz, H-5), 6.95 (1H, d, $J=1.6$ Hz, H-2), 6.87 (1H, d, $J=1.6$ Hz, H-2'), 6.88 (1H, dd, $J=8.4, 1.6$ Hz, H-6), 6.73 (1H, dd, $J=8.4, 1.6$ Hz, H-6'), 6.72 (1H, d, $J=8.4$ Hz, H-5'), 4.80 (1H, d, $J=7.8$ Hz, H-1''), 4.68 (1H, d, $J=4.0$ Hz, H-7), 4.66 (1H, s, H-1'''), 4.62 (1H, d, $J=4.4$ Hz, H-7'), 4.14 (3H, m, H-9 β , 9' β), 3.78 (3H, s, OCH₃), 3.76 (3H, s, OCH₃), 3.95-3.02 (sugar H), 3.02 (2H, m, H-8, 8'), 1.15 (3H, d, $J=6.4$ Hz, H-6'''). ¹³C-NMR (100 MHz, CD₃OD) δ : 150.7 (C-4), 148.9 (C-4), 147.0 (C-3'), 147.0 (C-3), 137.2 (C-1), 133.6 (C-1'), 119.9 (C-6'), 119.7 (C-6), 117.9 (C-5), 116.0 (C-5'), 111.5 (C-2), 110.9 (C-2'), 102.4 (C-1''), 101.9 (C-1'''), 87.2 (C-7'), 86.8 (C-7), 77.6 (C-5''), 76.7 (C-3''), 74.7 (C-2''), 73.9 (C-4''), 72.6 (C-9'), 72.5 (C-9), 72.2 (C-3'''), 72.0 (C-2'''), 71.3 (C-4''), 69.7 (C-5'''), 67.7 (C-6''), 56.7 (C-OCH₃), 56.6 (C-OCH₃), 55.2 (C-8'), 55.1 (C-8), 17.9 (C-6''').

RESULTS AND DISCUSSION

In the course of phytochemical study of the MeOH

extract from the twigs of *C. sinensis*, eight compounds were isolated by the chromatographic separation of each fraction. They were identified as two triterpenoids, two amide compounds, two lignan glycoside, and two steroids by spectroscopic analysis.

Compound **1** was obtained as an amorphous powder from MeOH. ¹H-NMR spectrum of **1** showed eight methyl singlet signals. The singlet peak at δ 4.85 (1H, s) was identified as H-19 proton signal attached on an olefinic double bond. In the ¹³C-NMR spectrum, 30 carbon signals were observed, which included two olefinic carbons at δ 142.7 and 129.7, and an oxygenated carbon at δ 79.0. As the result, compound **1** was identified as germanicol, which was reported from *Lactuca indica* and *Schaefferia cuneifolia* (Park *et al.*, 1995; Gonzalez *et al.*, 1989). The NMR spectra of **2** was similar to that of **1** but it did not show the olefinic double bond. The ¹H-NMR spectrum of **2** revealed the presence of a hydroxyl group at C-3 with δ 3.74 (1H, m). In the ¹³C-NMR spectrum, 30 carbon signals were observed, which included an oxygenated carbon at δ 72.8. From the comparison of all spectral data of compound **2**, they were in good agreement with epifriedelanol, which was reported in the literatures (Hong *et al.*, 1990; Ahmad *et al.*, 1994). Compounds **3** and **6** were identified as β -sitosterol (**3**) and β -sitosterol-3-O-glucoside (**6**) by




Fig. 1. Isolated compounds from *Celtis sinensis*

comparing physicochemical and spectral data with published literatures (Do *et al.*, 1988; Park *et al.*, 1994).

Compounds **4** and **5** have similar patterns in their NMR spectra. The ¹H-NMR spectrum of **4** showed two ortho-coupled doublets each of two protons with a *J* value of 8.4 Hz at δ 7.00 (2H, d, H-2', 6') and 6.62 (2H, d, H-3', 5'), indicating the presence of a 1,4-disubstituted aromatic ring. In addition, two coupled triplets corresponding to the methylene protons at δ 3.38 (2H, t, *J*=7.5 Hz, H-8) and 2.66 (2H, t, *J*=7.5 Hz, H-7) revealed the presence of tyramine moiety in **4**. The signals at δ 7.02 (1H, d, *J*=1.5 Hz, H-2), 6.95 (1H, dd, *J*=8.4, 1.5 Hz, H-6) and 6.69 (1H, d, *J*=8.4 Hz, H-5), and two *trans*-coupled vinylic protons at δ 7.33 (1H, d, *J*=15.6 Hz, H-7) and 6.30 (1H, d, *J*=15.6 Hz, H-8) revealed the presence of caffeoyl moiety in **4**. The ¹³C-NMR spectrum of **4** showed a carbonyl carbon, 10 aromatic carbons, two methylene carbons and two vinylic carbons. The structure of **4** was determined to be *trans*-*N*-caffeoyltyramine on the basis of the above evidences, together with a comparison of the above data with those published in the literature (Lajide *et al.*, 1996; Han *et al.*, 2002).

Compound **5** was obtained as an amorphous powder from MeOH. In the ¹³C-NMR spectrum of **5**, 13 carbon signals were observed, which included a carbonyl carbon, 8 aromatic carbons, two methylene carbons and two vinylic carbons. The ¹H-NMR spectrum of **5** showed two groups of ortho-coupled doublets each of two protons with a *J* value of 8.8 Hz at δ 7.26 (2H, d, H-2', 6') and 6.65 (2H, d, H-3', 5'), and a *J* value of 8.4 Hz at δ 6.91 (2H, d, H-2, 6) and 6.62 (2H, d, H-3, 5). In addition, two coupled triplets corresponding to the methylene protons at δ 3.29 (2H, t, *J*=7.5 Hz, H-8') and 2.59 (2H, t, *J*=7.5 Hz, H-7') and two *cis*-coupled vinylic protons at δ 6.51 (1H, d, *J*=12.8 Hz, H-8) and 5.69 (1H, d, *J*=12.8 Hz, H-7) were observed. On the basis of the above evidences, together with a comparison of the above data with those published in the literature, the structure of **5** was determined to be *cis*-*N*-coumaroyltyramine (Wu *et al.*, 1994).

The ¹H-NMR spectrum of **7** showed the typical pattern of two coupling groups of 1,3,4-trisubstituted benzene rings at δ 7.13 (1H, d, *J*=8.4 Hz, H-5), 7.01 (1H, d, *J*=1.6 Hz, H-2), 6.92 (1H, d, *J*=1.6 Hz, H-2'), 6.90 (1H, dd, *J*=8.4, 1.6 Hz, H-6), 6.80 (1H, dd, *J*=8.4, 1.6 Hz, H-6') and 6.76 (1H, d, *J*=8.4 Hz, H-5'), two methoxy groups at δ 3.85 and 3.84, and an anomeric proton at δ 4.87 (1H, d, *J*=7.8 Hz, H-1"). Eight proton signals at δ 4.74 (1H, d, *J*=4.0 Hz, H-7), 4.68 (1H, d, *J*=4.4 Hz, H-7'), 4.13 (2H, m, H-9 β , 9' β), 3.82 (2H, m, H-9 α , 9' α), and 3.11 (2H, m, H-8, 8') were assigned to a *cis*-diequatorial substituted 7,7-diaryldioxabicyclo[3,3,0]octane (Pérez *et al.*, 1995). On the basis of these data, the structure of **7** was elucidated as pinoresinol-4-*O*-glucopyranoside (Tsukamoto *et al.*,

1984; Chiba *et al.*, 1979).

Compound **8** showed a molecular ion peak at *m/z* 666 in the mass spectrum. The NMR spectra of **8** were similar to those of **7**, suggesting it has similar skeleton. The main difference was sugar moiety of aliphatic region. Another anomeric peak at δ _H4.66 (1H, s, H-1"), δ _C101.9 (C-1"), and five rhamnose sugar signal were observed in the ¹³C-NMR spectrum. In the HMBC spectrum, it was evident that the two methoxy groups are placed at each 3 and 3 position of the two aromatic rings. A correlation peak of the H-1 signal of glucose moiety at δ 4.80 (1H, d, *J*=7.8 Hz, H-1") and the C-4 signal of the phenyl group at δ _C150.7 suggested that the glucose moiety is connected to a 4-hydroxy group at the phenyl group. And a correlation peak in the HMBC spectrum of the H-1 signal for rhamnose at δ 4.66 (1H, s, H-1"), the C-6 signal for glucose at δ _C67.7 suggested that these two sugar moieties were connected by a 1→6 linkage between β -glucose and α -rhamnose, consistent with a rutinose structure (Agrawal, 1989). In conclusion, the structure of **8** was determined to be pinoresinol-4-*O*-[α -rhamnopyranosyl (1→6)-*O*- β -glucopyranoside].

Previously, epipinoresinol-4-*O*-[α -rhamnopyranosyl (1→6)-*O*- β -glucopyranoside] and pinoresinol-4-*O*-[α -rhamnopyranosyl(1→2)-*O*- β -glucopyranoside] reported in the literature (Lee *et al.*, 1999; Batirov *et al.*, 1985), but compound **8** has not been reported in any other plant.

ACKNOWLEDGEMENT

This research was supported by a grant (M103KD010014-04K0401-01410) from Plant Diversity Research Center of 21st Century Frontier Research Program funded by Ministry of Science and Technology of Korean government.

REFERENCES

- Agrawal, P. K., Carbon-13 NMR spectrum of flavonoids, Elsevier, New York, p. 341, (1989).
- Ahmad, V. U. and Rahman, A. U., Handbook of natural products data, vol. 2; Pentacyclic triterpenoids, Elsevier, New York, p. 566-567, (1994).
- Batirov, E., Kh., Matkarimov, A. D., Malikov, V. M., and Yagudaev, M. R., Versicoside-a novel lignan glycoside from *Haplophyll versicolor*. *Khim. Prir. Soedin.*, 5, 624-628 (1985).
- But, Paul P. H. Kimura, T. Guo, J. X., and Sung, C. K., *International collation of traditional and folk medicine: Part 2*. World scientific, Singapore, p. 22-23, (1997).
- Chiba, M., Okabe, K., Hisada, S., Shima, K., Takemoto, T., and Nishibe, S., Elucidation of the structure of a new lignan glucoside from *Olea europaea* by carbon-13 nuclear magnetic resonance spectroscopy. *Chem. Pharm. Bull.*, 27,

- 2868-2873 (1979).
- Cho, J. Y., Kim, A. R., and Park, M. H., Lignans from the rhizomes of *Coptis japonica* differentially act as anti-inflammatory principles. *Planta Med.*, 67, 312-316 (2001).
- Do, J. C., Son, K. H., and Kang, S. S., Studies on the constituents of the roots of *Rubus parvifolius* (L). Isolation of (-)-epicatechin. *Kor. J. Pharmacogn.*, 19, 170-173 (1988).
- Gonzalez, A. G., Mendoza, J. J., Ravelo, A. G., Luis J. G., and Dominguez, X. A., Δ^{18} oleanane triterpene from *Schaefferia cuneifolia*. *J. Nat. Prod.*, 52, 567-570 (1989).
- Han, S. H., Lee, H. H., Lee, I. S., Moon Y. M., and Woo, E. R., A new phenolic amide from *Lycium chinense* Miller. *Arch. Pharm. Res.*, 25, 433-437 (2002).
- Hong, N. D., Rho, Y. S., Kim, N. J., and Kim, J. S., Studies on the constituents of Ulmi Cortex. *Kor. J. Pharmacogn.*, 21 (3) 201-204 (1990).
- Kim, D. K. and Lee, K., Inhibitory effect of *trans-N-p*-coumaroyl tyramine from the twigs of *Celtis sinensis* on the acetylcholinesterase. *Arch. Pharm. Res.*, 26, 735-738 (2003).
- Koch, B. P., Harder, J., Lara, R. J., and Kattner, G., The effect of selective microbial degradation on the composition of mangrove derived pentacyclic triterpenols in surface sediments. *Organic Geochem.*, 36, 273-285 (2005).
- Kundu, J. K., Rouf, A. S., Hossain, M. N., Hasan, C. M., and Rashid, M. A., Antitumor activity of epifriedelanol from *Vitis trifolia*. *Fitoterapia*, 71, 577-579 (2000).
- Lajide, L., Escoubas, P., and Mizutani, J., Termite antifeedant activity in *Xylopi aethiopica*. *Phytochemistry*, 40, 1105-1112 (1995).
- Lee, S. J., Yun, Y. S., Lee, I. K., Ryoo, I. J., Yun, B. S., and Yoo, I. D., An antioxidant lignan and other constituents from the root bark of *Hibiscus syriacus*. *Planta Med.*, 65, 658-660 (1999).
- Matsunaga, S., Tanaka, R., Takaoka, Y., In Y., Ishida, T., Mawardi, R., and Ismail, H. B. M., 26-Nor-D:A-friedooleanane triterpenes from *Phyllanthus Watsonii*. *Phytochem.*, 32, 165-170 (1993).
- Park, H. J., Lee, M. S., Lee, E., Choi, M. Y., Cha, B. C., Jung, W. T., and Young, H. S., Serum cholesterol lowering effect of triterpene acetate obtained from *Lactuca indica*. *Kor. J. Pharmacogn.*, 26, 40-46 (1995).
- Park, S. W., Yook, C. S., and Lee, H. K., Chemical components from the fruits of *Crataegus pinnatifida* var. *psilosa*. *Kor. J. Pharmacogn.*, 25, 328-335 (1994).
- Pérez, C., Almonacid, L. N., Trujillo, J. M., González, A. G., Alonso, S. J., and Navarro, E., Lignans from *Apollonias barbujana*. *Phytochemistry*, 40, 1511-1513 (1995).
- Tsukamoto, H., Hisada, S., and Nishibe, S., Lignans from bark of *Fraxinus mandshurica* var. *japonica* and *F. japonica*. *Chem. Pharm. Bull.*, 32, 4482-4489 (1984).
- Wu, T. S., Ou, L. F., and Teng, C. M., Aristolochic acids, aristolactam alkaloids and amides from *Aristolochia kankauensis*. *Phytochemistry*, 36, 1063-1068 (1994).