

Influence of herbaceous cover, shelter and land cover structure on wild rabbit abundance in NW Portugal

José Carlos CARVALHO and Pedro GOMES

Carvalho J. C. and Gomes P. 2004. Influence of herbaceous cover, shelter and land cover structure on wild rabbit abundance in NW Portugal. *Acta Theriologica* 49: 63–74.

The European wild rabbit *Oryctolagus cuniculus* (Linnaeus, 1758) populations in the Iberian Peninsula has suffered a serious decline. Therefore, the knowledge about the factors that influence rabbit distribution and abundance is of major interest for conservation and management programmes. Rabbit relative abundance was evaluated by pellet counting in relation to herbaceous ground cover, shelter availability (tall scrub cover and gaps in rocks) and land cover structure in the Peneda-Gerês National Park (PGNP), NW Portugal. Rabbit abundance was higher at intermediate levels of herbaceous cover, but no significant statistical differences were detected. A strong association between the abundance index and tall scrub cover was verified, and when this cover was rare, an association with gaps in rocks was observed. At the land cover level, rabbit abundance was associated with high interspersed sites of rocks with matorral and high interspersed sites of tall scrubland. Evidence from this study highlighted the importance of shelter and the interspersed sites of open/shelter habitats for rabbits. This pattern could be interpreted as an anti-predator strategy. Therefore, management actions should favour the growing and establishing of tall scrub patches and the selective cutting in highly continuous areas of tall scrub vegetation.

Departamento de Biologia, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal, e-mail: pagomes@bio.uminho.pt

Key words: *Oryctolagus cuniculus*, shelter, abundance, food, interspersed

Introduction

The European wild rabbit *Oryctolagus cuniculus* (Linnaeus, 1758) is native to the Iberian Peninsula (Arthur 1989). This lagomorph is a key species in its environments, since it is an important prey for a large number of predators, some of them threatened, eg Iberian lynx *Lynx pardinus* and the Imperial eagle *Aquila adalberti* (Delibes and Hiraldo 1981, Iborra *et al.* 1990, Palomares 2001a), and also an important game species (Rogers *et al.* 1994). However, rabbit populations are seriously depleted in result of several factors, largely habitat destruction, over-hunting and diseases, such as, myxomatosis and more recently RVH (rabbit viral haemorrhagic) disease (Blanco and Villafuerte 1993). In some depleted populations, generalist predators may exert a regulating influence on rabbits retarding their recovery (Trout and Tittensor 1989, Pech *et al.* 1992, Banks *et al.* 1998, Banks 2000). The investigation of which habitat factors influence rabbit distribution and abundance are crucial for management and recovery plans for its populations.

Distribution and abundance of rabbits have been studied in various Mediterranean regions of the Iberian Peninsula (Rogers and Myers 1979, Soriguer 1983, Moreno *et al.* 1996, Palomares *et al.* 1996, Martins and Borralho 1998, Palomares *et al.* 2001), but little in areas with different biogeoclimatic conditions, like the North of Portugal. In Mediterranean regions rabbits were more abundant in areas suitable for warren building or which could provide an abundant supply of tall shrub vegetation well interspersed with areas dominated by herbaceous vegetation and/or agricultural lands (Rogers and Myers 1979, Soriguer 1983, Moreno *et al.* 1996, Palomares *et al.* 1996, Martins and Borralho 1998, Palomares *et al.* 2001). Although, the interspersion factor, between openfield patches (utilized for foraging) and cover (utilized for refuge against predators), has been implicitly recognized in these studies, no attempt has been made to measure this variable in relation to rabbit abundance. A study carried out on a large scale in Spain concluded that the distribution and abundance of rabbits is related to soil and climate (Blanco and Villafuerte 1993). However, Fa *et al.* (1999) argued that on a smaller scale habitat conditions, such as availability of food and shelter, might be determinant.

The aim of this paper is to clarify the influence of herbaceous cover, shelter availability and land cover structure (with emphasis on the interspersion between open/shelter habitats) on rabbit abundance in a transition region between Atlantic and Mediterranean vegetation, at a local scale, within the NW Portugal.

Material and methods

Study area

The study was carried out in the Peneda-Gerês National Park (PGNP), NW Portugal, near Montalegre (41°49'N, 07°47'W). The region (4606 ha) is located in a plateau, interrupted by mountains and small valleys, where the dominant lithological type is the granite, between 800 and 1400 m in altitude. Vegetation falls in the transition zone between Ibero-Atlantic acidiphilous oak woodlands and Supra-Mediterranean vegetation (Ozenda and Borel 2000). The area is dominated by matorral scrub composed of *Erica cinerea*, *Ulex* sp. and *Chamaespartium tridentatum*, within a mosaic of oakwoods (*Quercus pyrenaica* and *Quercus robur*) interspersed with pastures, agricultural fields and patches of tall scrubs of *Genista* sp. and *Cytisus* sp. Annual rainfall is about 2000 mm and annual average temperature is between 7.5 and 10°C.

A land cover map was developed from rectified orthophotomaps (freely available in <http://ortos.igeo.pt/ortofotos>), with a cell resolution of 16 m², by supervised classification for all pixels, in order to allow sampling procedures and land cover analysis (Fig. 1). Based on vegetation cover and land use, six land cover units were considered: deciduous forest, forest in regeneration, matorral, tall scrubland, pasture/agriculture fields, and rocks (a brief description of the land cover units is presented in Table 1). The map was extensively ground checked and corrected. Only minor corrections were required since the high resolution of the orthophotomaps allowed a precise classification.

Rabbit abundance

Pellet counts provide a simple and reliable means for estimating rabbit abundance (Palomares 2001b). We counted all the pellets observed along 86 belt transects of 400 × 4 m. The length of the transect was determined after initial trials. These trials revealed that rabbits were at very low densities, and for that reason, a longer transect was selected in order to maximize the detection of




Fig. 1. Land cover map of the study area (northeastern part of Peneda-Gerês National Park), obtained by supervised classification of aerial ortophotomaps (1995 flight) and field validation.

Table 1. Land cover units description.

| Land cover unit | Description | Occupied area (ha) | Percentage of the total area (%) |
|------------------------------|--|--------------------|----------------------------------|
| Deciduous forest | Patches of oakwood (<i>Quercus pyrenaica</i> and <i>Quercus robur</i>) with a mean tree height superior to 4 m. The understorey cover is mainly composed by <i>Ilex aquifolium</i> and <i>Crataegus monogyna</i> . | 578.7 | 12.6 |
| Regenerating forest | Patches of oakwood in regeneration after fire or cut, with a mean tree height inferior or equal to 4 m. | 309.9 | 6.7 |
| Matorral | Short scrubland dominated by <i>Erica cinerea</i> , <i>Ulex</i> sp. and <i>Chamaespartium tridentatum</i> ; mean scrub height inferior or equal to 1 m. | 2267.6 | 49.2 |
| Tall scrubland | Patches of tall scrubs of <i>Genista</i> sp. and <i>Cytisus</i> sp. resulting from forest degradation or abandoned fields. | 427.3 | 9.3 |
| Pastures/agricultural fields | Fields of grassland utilized for grazing bovine cattle interspersed with cereal fields of <i>Solanum tuberosum</i> , <i>Secale cereale</i> and <i>Zea mays</i> . | 688.9 | 15.0 |
| Rocks | Emerged rocks of granite dispersed mainly in the matorral. | 145.8 | 3.2 |
| Others | Mainly small villages and roads. | 187.5 | 4.0 |

rabbit pellets. The width of the transect was selected in order to maximize the observation of pellets by the researcher. The relative density of rabbits is expressed as the number of pellets/m².

Stratified sampling was followed by dividing the study area into 1 × 1 km units and randomly selecting one transect for each vegetal/landscape formation in each plot. The land cover map was used as a support for this procedure. In general, only those units covering more than 25% of the plot were sampled, although some minor units potentially used by rabbits were considered (rocks and patches of tall scrubland). A Global Positioning System (GPS; model Garmin 12XL, Garmin Corp., Olathe, Kansas, USA) was used to record the mean point position (ie the centre) of each transect. Only records with a GPS error lesser than 5 m were considered. When GPS error was greater than 5 m, we made subsequent records until the defined error limit was reached.

Herbaceous cover

Rabbits use a wide spectrum of foods, but mostly consume herbaceous plants (Soriguer 1988, Chapuis and Gaudin 1995). Seasonal and spatial variation in the diet of the wild rabbit in Portugal demonstrated also the importance of grass-forbs (Martins *et al.* 2002). These studies indicated that the genera *Lolium*, *Holcus*, *Bromus*, *Briza*, *Vulpia*, *Agrostis*, *Senecio*, *Echium* and *Leontodon* are usually consumed by rabbits. These plants are abundant in the herbaceous communities of the northeastern PGNP. By this reason we used the percentage of herbaceous ground cover as an index of food supply.

The sampling procedure consisted in dividing each belt transect into 10 (10 × 4 m) segments and estimating the percentage of herbaceous cover by visual observation following Higgins *et al.* (1996). These values were averaged for the 10 segments and then reclassified in 5 classes to reduce bias produced by visual estimation: 0–20%, 21–40%, 41–60%, 61–80%, 81–100%.

Shelter availability

Rabbits use tall shrub vegetation for refuge but could also use gaps between rocks (Rogers *et al.* 1994). Therefore, shelter availability was estimated by assigning to each transect, a class of percentage of ground covered by tall shrub vegetation, using the same procedure as for the estimation of herbaceous cover. Four classes of shelter availability were considered: 0–20%, 21–40%, 41–60%, > 60%. When shelter availability was = 20%, the presence or absence of other alternative refuges (gaps between rocks) was also recorded.

Land cover structure

The percentage cover by each land cover unit was measured on 3.14 ha circular plots (approximately the allometric home range of rabbits, Villafuerte 1994) around the mean point position (ie the centre) of each transect. Transects were located in sites where the immediately surrounding vegetation was similar to the area covered by the circular plot. Therefore, we assumed that the land cover data evaluated in the circle was representative of the surface crossed by the transect.

The interspersation between food patches and shelter patches was determined by the index of Patton (described in Litvaitis *et al.* 1996). This index measures the irregularity of a patch, based on the area-to-perimeter ratio of a circle, and has the following form:

$$I = P/2(\pi A)^{1/2}$$

where P is the perimeter of the edge between shelter patches (tall scrubland, rocks and forest) and foraging patches (pasture/agricultural fields and matorral) and A is an area of reference of 3.14 ha. All variables were estimated by GIS analysis with the software package MF-Works, version 2.6.3 (Keigan Systems, London, Ontario, Canada).

Statistical analysis

After testing the normality of the frequency distribution of the data on pellet-counts with the Kolmogorov-Smirnov test, the non-parametric analysis of variance (Kruskal-Wallis test) was chosen for detecting statistical differences between herbaceous cover and shelter availability; pairwise comparisons were made by Dunn test. Mann-Whitney *U*-test was also used to compare sites without any shelter alternative to shrubs and sites with alternative refuge (gaps between rocks). All these statistical procedures were performed in the Prism (version 3) software package (Motulsky 1999) with a significant level of 0.05.

The multidimensional characteristic of the land cover data requires a multivariate approach, in order to avoid the problem of the non-independence of variables. Therefore, a PCA (Principal Component Analysis) was performed from the land cover data. PCA creates independent factors which are a linear combination of the original variables. Prior to the PCA, land cover variables were arcsine transformed (Zar 1998). The PCA factors obtained were related with the rabbit relative abundance index, using the Spearman correlation coefficient, so that the distribution of the species could be described in relation to the land cover structure. Nevertheless, in order to search for humped shaped relationships between rabbit abundance and land cover data, non-parametric regression smoothing was used. These methods provide versatile techniques to explore general non-linear relationships between two variables (Hardle 1990). The LOWESS (LOcally WEighted Scatter plot Smoothing) was used to describe the relationship between the rabbit relative abundance (ie response variable) and the PCA factors (ie explanatory variables). The LOWESS technique is a k-nearest neighbour smoother adequate for handling data sets with outliers and a “noisy” relationship between variables, like the one here reported (for more details see Hardle 1990). PCA and LOWESS analysis were performed using the software ADE-4 (Thioulouse *et al.* 1997). The ADE-4 allows to select the number of neighbours, to use in the LOWESS procedure, in order to minimize the average error of the smoother curve in relation to the observed data.

Results

Herbaceous cover and shelter availability

A total of 86 transects was surveyed. Pellets were found in 40 transects (46.5%) with a mean relative density of 1.83 pellets/m² (SD = 3.71). Frequency distribution of data on pellet counts was not normally distributed (Kolmogorov-Smirnov test: $D = 0.3249, p < 0.0001$) justifying the choice for non-parametric statistical tests.

Rabbit abundance was highest at intermediate levels of herbaceous cover (the maximum abundance was found in the class of 41–60%), suggesting a possible humped shaped relationship between the two variables. A remarkable low value of rabbit abundance was observed in the highest class of herbaceous cover. Nevertheless, abundance of rabbits was highly variable (large SD values) and no significant statistical differences were found among the five classes considered (Table 2).

Results on pellet counts indicated that mean relative rabbit densities increased with increasing tall scrub cover. Statistical differences were detected among the 4 classes of percentage of ground covered by tall scrubs. Interclass comparisons with Dunn test showed significant differences between the two major classes (41–60% and > 60%) and the minor class (0–20%) of tall scrub cover (Table 2).

In places where tall shrub cover was = 20%, rabbits were more abundant when an alternative shelter existed (gaps between rocks). Mean (\pm SD) relative density of rabbits was 0.27 ± 0.93 pellets/m² in areas with no gaps between rocks ($n = 40$), and 1.23 ± 2.17 pellets/m² in areas with gaps between rocks ($n = 16$). These differences were statistical significantly (Mann-Whitney U -test: $U = 155.50, p < 0.05$).

Land cover structure

From the PCA on the land cover data, three factors were retained for subsequent analysis. The three PCs explain 66.6% of the total variance (Table 3). The first

Table 2. Comparison of mean relative densities of rabbits (pellets/m²) as a function of 5 classes of percentage of herbaceous cover and 4 classes of percentage of tall scrub cover.

| Variable | Class (%) | <i>n</i> | Mean \pm SD | Kruskall-Wallis test | | Dunn test – significant different pairs ($p < 0.05$) |
|--------------------------------|-----------|----------|-----------------|----------------------|----------|--|
| | | | | <i>H</i> | <i>p</i> | |
| Percentage of herbaceous cover | | | | | | |
| | 0–20 | 10 | 1.99 \pm 2.81 | 8.273 | ns | none |
| | 21–40 | 18 | 2.17 \pm 4.53 | | | |
| | 41–60 | 23 | 2.44 \pm 4.47 | | | |
| | 61–80 | 13 | 2.35 \pm 3.87 | | | |
| | 81–100 | 22 | 0.59 \pm 1.92 | | | |
| Percentage of tall scrub cover | | | | | | |
| | 0–20 | 56 | 0.64 \pm 1.81 | 25.158 | < 0.0001 | 0–20% vs 41–60% 0–20% vs > 60% |
| | 21–40 | 6 | 1.55 \pm 2.14 | | | |
| | 41–60 | 18 | 3.56 \pm 4.61 | | | |
| | > 60 | 6 | 8.12 \pm 6.80 | | | |

Table 3. Results of PCA from land cover data. The correlation between the original variables and the 3 first PC factors, along with eigenvalues and the cumulative percentage of variance explained, are presented.

| Variable | PC-1 | PC-2 | PC-3 |
|--|---------|---------|---------|
| Eigenvalue | 1.8798 | 1.5968 | 1.1867 |
| Cumulative % of variance | 26.9 | 49.7 | 66.6 |
| % cover by deciduous forest | -0.5837 | 0.1813 | 0.5653 |
| % cover by regenerating forest | 0.0160 | 0.0159 | 0.5115 |
| % cover by matorral | 0.5970 | -0.7393 | -0.1862 |
| % cover by tall scrubland | 0.2305 | 0.6667 | -0.4388 |
| % cover by pasture/agricultural fields | -0.6799 | 0.1217 | -0.4388 |
| % cover by rocks | 0.5996 | 0.1106 | 0.2686 |
| Interspersion | 0.5896 | -0.6793 | 0.2123 |




Fig. 2. Relationship between rabbit abundance (pellets/m²) and the first PCA factor. A LOWESS smoother curve is superposed. The vertical line in the middle of the graph separates negative and positive PC factor scores.

factor shows a gradient from sites characterized by a high interspersion between matorral and rocks (positive scores) towards sites dominated by continuous areas of deciduous forest and pastures/agricultural fields (negative scores). The second factor describes a gradient from high interspersion sites of tall scrubland (positive scores) to sites dominated by continuous areas of matorral (negative scores). The third factor generates a gradient from places dominated by tall scrubland and pastures/agricultural fields (negative scores) towards places dominated by deciduous forest and forest in regeneration (positive scores). The rabbit relative abundance index is significantly correlated with the first two factors ($r_S = 0.5575$, $p < 0.0001$ for PCA-1; $r_S = 0.4633$, $p < 0.0001$ for PCA-2), but was not correlated with the third factor ($r_S = -0.1810$, $p > 0.05$).

In Figs 2 and 3, two scatterplots of rabbit relative abundance index versus PCA – scores (factor one and two) are presented. Superposed are LOWESS smoother curves which allow a more easily interpretation of the relationship between variables. Hence, higher pellet densities were clearly associated with positive scores of the first and second factors.




Fig. 3. Relationship between rabbit abundance (pellets/m²) and the second PCA factor. A LOWESS smoother curve is superposed. The vertical line in the middle of the graph separates negative and positive PC factor scores.

Discussion

In Mediterranean regions, habitat selection by rabbits may be determined by a compromise between food availability and protection from predators (Moreno *et al.* 1996) provided both by bushes and warrens (Palomares and Delibes 1997). In the northeastern PGNP, rabbit abundance seems to have a humped shaped relationship with an index of food supply (percentage of herbaceous cover). This means that rabbits are less abundant in places where herbaceous cover reaches extreme values. Low herbaceous cover could be a limiting factor when rabbits are at high densities, since it may result in a food shortage. High herbaceous cover are associated with openfield, where shelter is not available, and rabbits are more vulnerable to predation (by humans and native predators) (Lombardi *et al.* 2003).

Despite the above considerations, a highly variable response of rabbits to herbaceous cover, in the northeastern PGNP, was observed (large SD values), so that no significant statistical differences were found. This could be a consequence of two factors. Firstly, rabbit abundance appears to be very low in comparison with previous studies. The mean and maximum pellet densities recorded in our study area were well below the values recorded for southern Spain and southern France (Rogers and Myers 1979, Rogers 1981, Palomares *et al.* 1996, Fa *et al.* 1999). Hence, rabbit populations in northeastern PGNP are probably far from the habitat carrying capacity limit in food supply terms. Secondly, the northeastern PGNP is situated in a transition area between Ibero-Atlantic acidiphilous oak woodlands vegetation and Supra-Mediterranean vegetation (Ozenda and Borel 2000) and seasonality of rainfall and herbaceous vegetation production are not so marked as in Mediterranean habitats. Therefore, the availability of herbaceous plants for rabbits to eat is probably more limited in Mediterranean regions. However, it should be noticed that only the percentage of ground cover by herbaceous plants was estimated and related with rabbit abundance in this study. This is a rough measure of food supply. Some aspects more meaningful, such as herbaceous biomass or volume and food quality, were not evaluated. Therefore, we argue that this subject needs more intensive research to clarify the influence of food supply on rabbit abundance in other areas outside the Mediterranean region.

Rabbit abundance was positively associated with shrub cover and, in the absence of this cover, a positive association with gaps in rocks was verified. The importance of shrub cover has also been observed in Mediterranean habitats but, in these regions, rabbits may also use open areas if warrens were available nearby (Palomares and Delibes 1997). Lombardi *et al.* (2003) found that in grassland rabbits lived in groups linked to aggregated burrows. However, in our case warrens were detected only in < 10% of the 86 transects, which could be a consequence of the soil characteristics. Two main types of soil situation were found: deep and very humid soils, corresponding to the valleys were oak wood forest or permanent pastures/ agricultural fields were present, and squeletic soils scattered with rocks of various sizes, dispersed throughout the matorral, in the highest part of the region. Thus,

contrary to Mediterranean regions, in northeastern PGNP the nature of the soil makes warren construction difficult not allowing them to explore open areas. Therefore, rabbits may react by staying closer to shrub cover and gaps between rocks.

The analysis at the land cover level showed that rabbit abundance was associated to sites of high interspersion between rocks and matorral (first PCA factor) and high interspersion sites of tall scrubland (second PCA factor). These results clearly highlighted the importance of shelter and the interspersion between open/shelter habitats for rabbits, as hypothesized in other studies for Mediterranean zones of Iberian Peninsula (Rogers and Myers 1979, Soriguer 1983, Moreno *et al.* 1996, Palomares *et al.* 1996, Martins and Borralho 1998, Palomares *et al.* 2001).

The abundance pattern of wild rabbit in the PGNP seems to be determined by an anti-predator strategy (Jaksic and Soriguer 1981, Villafuerte 1994, Villafuerte and Moreno 1997, Banks *et al.* 1999). This strategy consists in the utilization of closed areas during daylight (and gaps in rocks) for protection against diurnal birds of prey (and human hunting), but not from nocturnal or crepuscular carnivores (the red fox and the wild cat are the main mammal predators of rabbits in the PGNP; Carvalho 2001) whose prey tactics are favored by shrub vegetation (Villafuerte 1994, Moreno *et al.* 1996). Hence, during the night, rabbits tend to use edges between open/closed areas for hiding in bushes or gaps in rocks, foraging and/or run in openfield (Moreno *et al.* 1996, Villafuerte and Moreno 1997, Martins and Borralho 1998).

The results obtained in this study suggested that management actions should favor the growth and establishment of patches of natural shrubs (Palomares *et al.* 2001), similar in structure to tall scrubland of *Genista* sp. and *Cytisus* sp., scattered in areas of continuous matorral and pastures/agricultural fields. This strategy may enhance the availability of shelter conditions and the interspersion of open/closed areas. Continuous areas of tall scrubland, where interspersion is lower, can also be a problem for rabbits, since the risk of predation by nocturnal carnivores is higher (Moreno *et al.* 1996). Therefore, in these places selective vegetation cutting should be undertaken to the level where interspersion reaches an appropriate value. We suggest, an interspersion value, calculated according to the method here described, equal or superior to 1.

Acknowledgements: We are grateful to two anonymous referees for constructive criticisms and valuable comments.

References

- Arthur C. P. 1989. Origine et histoire du lapin. Bulletin mensuel, Office National de la chasse 135: 13–21.
- Banks P. B. 2000. Can foxes regulate rabbit populations? The Journal of Wildlife Management 64: 401–406.
- Banks P. B., Dickman C. R. and Newsome A. E. 1998. Ecological costs of feral predator control: foxes and rabbits. The Journal of Wildlife Management 62: 766–772.

- Banks P. B., Hume I. D. and Crowe O. 1999. Behavioural, morphological and dietary response of rabbits to predation risk from foxes. *Oikos* 85: 247–256.
- Blanco J. C. and Villafuerte R. 1993. Factores ecologicos que influyen sobre las poblaciones de conejos, incidencia de la enfermedad hemorragica. ICONA, Madrid: 1–123.
- Carvalho J. C. 2001. [Distribution and abundance of wild rabbit: influence of habitat quality and predators]. MSc thesis, Universidade do Minho, Braga, Portugal: 1–101. [In Portuguese]
- Chapuis J. L. and Gaudin J. C. 1995. Utilisation des ressources trophiques par le lapin de garenne (*Oryctolagus cuniculus*) en garrigue sèche aménagée. *Gibier Faune Sauvage* 12: 213–230.
- Delibes M. and Hiraldo F. 1981. The rabbit as a prey in the Iberian mediterranean ecosystem. [In: Proceedings of the World Lagomorph Conference (1979). K. Myers and C. D. MacInnes, eds]. University of Guelph, Ontario: 654–663.
- Fa J. E., Sharples C. M. and Bell D. J. 1999. Habitat correlates of European rabbit (*Oryctolagus cuniculus*) distribution after the spread of RVHD in Cadiz Province, Spain. *Journal of Zoology*, London 249: 83–96.
- Hardle W. 1990. Applied nonparametric regression. Cambridge University Press, Cambridge: 1–352.
- Higgins K. F., Oldemeyer J. L., Jenkins K. J., Clambey G. K. and Harlow R. F. 1996. Vegetation sampling and measurement. [In: Research and management techniques for wildlife and habitats. Fifth ed., rev. T. A. Bookhout, ed]. The Wildlife Society, Bethesda, Md: 567–591.
- Iborra O., Arthur C. P. and Bayle P. 1990. Importance trophique du lapin de garenne pour les grands rapaces provençaux. *Vie Millieu* 40 (2/3): 177–188.
- Jaksic F. and Soriguer R. C. 1981. Predation upon the European rabbit (*Oryctolagus cuniculus*) in mediterranean habitats of Chile and Spain: a comparative analysis. *Journal of Animal Ecology* 50: 269–285.
- Litvaitis J. A., Titus K. and Anderson E. M. 1996. Measuring vertebrate use of terrestrial habitats and foods. [In: Research and management techniques for wildlife and habitats. Fifth ed., rev. T. A. Bookhout, ed]. The Wildlife Society, Bethesda, Md: 254–274.
- Lombardi L., Fernandez N., Moreno S. and Villafuerte R. 2003. Habitat-related differences in rabbit (*Oryctolagus cuniculus*) abundance, distribution and activity. *Journal of Mammalogy* 84: 26–36.
- Martins H. and Borralho R. 1998. Evaluation of wild rabbit habitat selection (*Oryctolagus cuniculus* L. 1758) in a zone of the Center of Portugal by analysis of presence signs]. *Silva Lusitana* 6 (1): 73–88. [In Portuguese with English summary]
- Martins H., Milne J. A. and Rego F. 2002. Seasonal and spatial variation in the diet of the wild rabbit (*Oryctolagus cuniculus* L.) in Portugal. *Journal of Zoology*, London 258: 395–404.
- Moreno S., Villafuerte R. and Delibes M. 1996. Cover is safe during the day but dangerous at night: the use of vegetation by European wild rabbits. *Canadian Journal of Zoology* 74: 1656–1660.
- Motulsky H. J. 1999. Analysing data with Graphprism. Graphpad Software Inc., San Diego, CA: 1–379.
- Ozenda P. and Borel J. L. 2000. An ecological map of Europe: why and how? *Compte Rendu de l'Académie des Sciences, Paris, Sciences de la vie/Life Sciences* 323: 983–994.
- Palomares F. 2001a. Vegetation structure and prey abundance requirements of the Iberian lynx: implications for the design of reserves and corridors. *Journal of Applied Ecology* 38: 9–18.
- Palomares F. 2001b. Comparison of 3 methods to estimate rabbit abundance in a Mediterranean environment. *Wildlife Society Bulletin* 29: 578–585.
- Palomares F., Calzada J. and Revilla E. 1996. El manejo del hábitat y la abundancia de conejos: diferencias entre dos áreas potencialmente idénticas. *Revista Florestal* 9 (1): 201–210.
- Palomares F. and Delibes M. 1997. Predation upon European rabbits and their use of open and closed patches in Mediterranean habitats. *Oikos* 80: 407–410.
- Palomares F., Delibes M., Revilla E., Calzada J. and Fedriani J. M. 2001. Spatial ecology of Iberian lynx and abundance of European rabbits in Southwestern Spain. *Wildlife Monographs* 65: 1–36.
- Pech R. P., Sinclair A. R. E., Newsome A. E. and Catling P. C. 1992. Limits to predator regulation of rabbits in Australia: evidence from predator-removal experiments. *Oecologia* 89: 102–112.

- Rogers P. M., Arthur C. P. and Soriguer R. C. 1994. The rabbit in continental Europe. [In: The European rabbit: the history and biology of a successful colonizer. H. V. Thompson and C. M. King, eds]. Oxford University Press, Oxford: 22–63.
- Rogers P. M. and Myers K. 1979. Ecology of the European wild rabbit, *Oryctolagus cuniculus* (L.), in mediterranean habitats: I. Distribution in the landscape of the Coto Doñana, S. Spain. *Journal Applied Ecology* 16: 691–703.
- Rogers P. M. 1981. Ecology of the European wild rabbit, *Oryctolagus cuniculus* (L.), in mediterranean habitats: II. Distribution in the landscape of the Camargue, S. France. *Journal Applied Ecology* 18: 355–371.
- Soriguer R. C. 1983. El conejo: papel ecológico y estrategia de vida en los ecosistemas mediterráneos. XV Congreso de Fauna Cinegética y Silvestre. Trujillo, España: 517–542.
- Soriguer R. C. 1988. Alimentación del conejo (*Oryctolagus cuniculus* L. 1758) en Doñana. *Doñana Acta Vertebrata* 15 (1): 141–150.
- Thioulouse J., Chessel D., Dolédec S. and Olivier J. M. 1997. ADE-4: a multivariate analysis and graphical display software. *Statistics and Computing* 7: 75–83.
- Trout R. C. and Tittensor A. M. 1989. Can predators regulate wild rabbit *Oryctolagus cuniculus* population density in England and Wales? *Mammal Review* 19: 153–173.
- Villafuerte R. 1994. Riesgo de predación y estrategias defensivas del conejo, *Oryctolagus cuniculus*, en el Parque Nacional de Doñana. PhD thesis, Universidad de Córdoba, Córdoba: 1–229.
- Villafuerte R. and Moreno S. 1997. Predation risk, cover type and group size in European rabbits in Doñana (SW Spain). *Acta Theriologica* 42: 225–230.
- Zar J. H. 1998. *Biostatistical analysis*. 4th ed. Prentice Hall, New Jersey: 1–929.

Received 23 July 2002, accepted 12 November 2003.

Associate Editors were Bogumiła Jędrzejewska and Krzysztof Schmidt.