

Potential of natural bed soil in adsorption of heavy metals in industrial waste landfill

¹*M. Esmaili Bidhendi; ¹A. R. Karbassi; ¹A. Baghvand; ²M. Saeedi; ¹A. H. Pejman

¹Graduate Faculty of Environment, University of Tehran, Tehran, Iran

²Environmental Research Laboratory, School of Civil Engineering, Iran University of Science and Technology, Tehran, Iran

Received 28 April 2009; revised 24 June 2009; accepted 30 January 2010; available online 1 June 2010

ABSTRACT: Development of higher welfare could not be realized unless by energy consumption and other natural resources. Growth of industrial complexes has shown an unprecedented trend during recent years. Many of these towns have no treatment systems for the industrial wastes leachates. Besides, the chemical composition of wastes in such complexes varies considerably due to the different kinds of industries. It is endeavored in the present work to study the natural potential of soil to treat leachate of such industrial wastes. For this purpose, the Aliabad industrial complex in Tehran – Garmsar road was selected as the study area. The potential of adsorption of elements such as nickel, copper, cadmium, zinc, chromium, lead and manganese was investigated. The results indicated that the soil potential to adsorb heavy metals (except for manganese) was very high (95 %) in the adsorption of heavy metals (except for manganese). Further, chemical partitioning studies revealed that heavy metals are associated with various soil phases such as loosely bonded ions, sulfide and organics to various extents. Among the mentioned soil phases, one can deduce that major portion of metal contaminants is absorbed as loosely bonded ions. Organic bond and sulfide bond are in the 2nd and 3rd positions of metal contaminants adsorption, respectively. The results of the present study apparently showed that soil column had ample capacity to adsorb metal contaminants. Thus, determination of soil potential in adsorption of heavy metals during site selection is as important criteria.

Keywords: Heavy metal; Industry; Leachate; Waste landfill

INTRODUCTION

Industrial complexes are one of the major industrial centers, which play a remarkable role in progress and promotion of industry in developing countries. The concept beyond establishment of industrial complex is to control pollution emission through an integrated manner. Normally, this observation related to sanitary and environmental principles has led to increased amount of pollution in these complexes. Industrial activities are identified as a major cause for metal contamination of the ecosystem (Charlesworth *et al.*, 2003; Al-Khashman, 2004; Banat *et al.*, 2005; Karbassi *et al.*, 2008). Depending on type of industrial activities, huge amount of industrial disposal is produced. Due to lack of properly equipped plants and sanitary dumping sites operated within the required standards, the industrial wastes are released in an adhoc manner to the environment. Because of possessing highly

hazardous ingredient like different kinds of heavy metals, they are considered as the main sources of pollution. In many of the developing countries, solid wastes are dumped in open areas regardless of any eco-sanitary concern (Vastanthi *et al.*, 2008). Solid waste leachate which is the greatest that treat to groundwater possess various chemical and biological contaminants (Longe and Enekwechi, 2007; Mahvi, 2008; Suthar and Singh, 2008). Hence, regulatory measures related to the waste management and environmental conservancy is significant in industrial complexes. Indeed, due to the presence of different chemical compositions such as heavy metals, hydrocarbons and poisonous materials, as well as potential capability of producing leachate, industrial wastes and landfills without environmental safety are considered as one of the pollutant sources of ground water and surface water (Zheng *et al.*, 1991; Lee and Jones, 1993; Christensen *et al.*, 2001; Cheng *et al.*,

*Corresponding Author Email: m.esmaili@gmail.com
Tel.: +98912 1836 670; Fax: +9821 6640 7719



2004; Chen *et al.*, 2005; Nouri *et al.*, 2008; Malakootian *et al.*, 2009). The produced leachates is normally comprised of organic and inorganic compositions. In addition, with elapse of a time, the produced leachate permeates into ground systems leading to change of physical and chemical properties of groundwater (Vastani *et al.*, 2008). The leachate composition and its pollution intensity depend on many aspects such as landfill age, waste ingredients and hydraulic conditions of landfill (Qasim and Chiang, 1994; Okafor and Opuene, 2007). Produced leachate essentially contain elements such as cadmium, manganese, potassium, nitrogen, heavy metals (e.g. iron, copper, manganese, chromium, nickel, lead) and organic compositions like poly aromatic hydrocarbons, Acetone, Benzene, Toluene, Chloroforms (Freeze and Cherry, 1979; Lee *et al.*, 1986; Bagchi, 1989).

The density of mentioned materials in leachates and ground water depends on wastes composition (Fang, 1995). Leachates travel from unsaturated zones to ground waters surfaces. Some of the pollutant materials may be adsorbed by soil particles within their movement through the several of soil strata (Fang, 1995). When the leachate is mixed up with groundwater, it would be dispersed in the flow direction that eventually leads to spatial pollution (Vastani *et al.*, 2008). The rate and intensity of pollution in aquifers depend on rate of pollutants transfer from soil (Vastani *et al.*, 2008). Thus many factors in soil such as cation exchange capacity (CEC), clay minerals, organics and carbonates must be studied. Nevertheless, due to simplicity, low cost and efficiency and ignorance, solid wastes are often dumped in buried locations (Barrett and Lawlor, 1995). Bury and dump of solid wastes in dump places leads to many environmental problems due to emission of odor, air pollutants and effluents (Fatta *et al.*, 1999; Panjeshahi and Ataei, 2008). Also landfill is known to be sources of groundwater pollution (Allen, 2001). Gharaibeh and Masad (1989) have shown that dissolution of solid waste by rain produces a noticeable amount of polluted water in the form of leachate. Environmental problem of wastes are much more serious with industrial wastes that contain hazardous chemicals. This implies that contaminants may remain in their initial form for a long period (Fatta *et al.*, 1999). Potentially becoming hazardous to the exposed ecosystems. Issues related to heavy metal associated with industrial waste require considerable attention. For example, toxicological

impacts of various heavy metals, irrespective of their uses in the industry are known to be of an environmental concern (Aucott, 2006). For example, lead has been reported to cause various diseases such as brain damage, anorexia, mental deficiency and anemia also cadmium has been implicated in agonistic and antagonistic effects on hormones and enzymes leading to many of malformations such as renal damage (Lewis, 1991; Low *et al.*, 2000; Bulut and Baysal, 2006). Likewise, Pb and Cd are classified as carcinogens (EPA 1992b; Pekey, 2006). Nevertheless, because of owning exclusive characteristics, some of them such as Hg, Cd and Pb are often used in production of industrial products like different batteries, switches, electronic boards and so on (Aucott, 2006). Futta *et al.* (1997) showed in their study conducted on Athena Region that groundwater quality is highly influenced by leachate. High density of leachate in ground water of Athena region was also found. Heavy metals, investigated in this study, were Ni, Mn, Cr, Pb, Zn, Cu and Cd, each of which has been reported for various health problems non-biodegradable and with the possibility of accumulation in the food chain (Langston, 1990; Nameni *et al.*, 2008). This study attempts to investigate the impact of the heavy metal in the industrial waste disposal site of Aliabad industrial complex on the soils, by natural bed soil as an adsorption media to reduce such contamination.

Study area

Aliabad industrial complex is one of the first industrial complexes of Tehran Province located in Tehran – Garmsar road, 55 km away from Tehran City. The study location is situated at 35° 21'23" N and 51°55'59" E. The active industries in the complex involve metal, minerals, food, electricity, electronic, textile and chemicals. The geographical location of the investigated complex is shown in Fig. 1.

MATERIALS AND METHODS

A sample was extracted at 1m depth (intact) from the site by a hand auger. The sample was transferred into a polyethylene pipe and transported to the laboratory of Faculty of Environment, University of Tehran. Industrial leachates were collected in High Density Poly-ethylene (HDPE) bottles. A part of leachates was filtered for trace metal measurement by atomic absorption spectrophotometry (AAS). Then the






Fig. 1: Study area map




Fig. 2: Schematic view of the soil column



remaining leachates were poured on the top of soil sample and the outlet leachate was collected at the bottom of the soil column. The experiment was carried out for 35 continuously days. The schematic view of the soil column is depicted in Fig. 2. Two sets of soil samples were selected for chemical analysis. The first set includes natural soils from area of study and the second set encompasses the soil column that was used for purification of heavy metals in laboratory. At the end of 35th day, we repeated soil sampling at the same height as mentioned before. Then the obtained seven soil samples and eight collected leachate samples were prepared in accordance to Environmental Protection Agency (EPA) method number 3010 A (EPA, 1992a) and 7000 B (EPA, 2007). The concentrations of elements in samples have been analyzed using flame atomic absorption spectrophotometer.

RESULTS AND DISCUSSION

The results of soil analysis show that the project location has mainly coarse grains that are classified in GW-GM, GM, SM, SC-SM classes. It should be noted that fine texture of soil were classified in CL-ML and ML classes. The soil cohesion coefficient was determined to be 0.1 kg/cm² and the natural specific weight of the soil was about 1.85 g/cm³. Complementary data of soil column are presented in Table 1.

Table 2 shows the concentration of Ni, Cu, Cd, Zn, Pb, Cr and Mn in the leachates that are collected at 5, 10, 15, 20, 25, 30 and 35 days from bottom of the soil column.

For instance, in this study, it was observed that the initial concentration of Mn in the leachate was 0.75 mg/L, which was decreased to 0.39 mg/L after release of leachate over the soil sample. Other elements

also such as Cr, Pb, Zn, Cd, Cu and Ni were absorbed and dissipated by more than 95 %. The results of Table 2 show that major portions of contaminants were absorbed in the first 5 days of leachate release into the soil column. After accomplishment of first level of experiments, the contents of elements were measured in different sections of soil (Table 3). The comparison of elemental concentrations of intact soil with those of contaminated ones at various column depths was determined and provided the first line of proof that heavy metals adsorption by soil layers. Adsorption of heavy metals by soil layers at different depths should not be considered as purification of metal contaminants. In fact, type of association of elements with different phases of soil is very important in their fate. Typically, five important combinations naming “Loosely bonded ions”, “Sulfide bond”, “Organic-Metal bond”, “Resistant bond” and “Within lattice bond” are critical in understanding the contaminant dynamics of adsorption. The first three bonds are known as anthropogenic sources and the last two bonds are indicative of lithogenous sources. However, it is worthy to note that the contaminants in any of three anthropogenic bonds can be released due to partial variations in physical and chemical properties of soil. Such release may lead to surface and ground water pollution. The Sulfide bond that is indicative of redox conditions can actually occur where oxygen is depleted in the environment. The presence of elements such as Fe and Mn in this combination indicates formation of sulfides in the environment. It might be argued that Organo-Metallic bonds are more resistant to environmental changes than the other two discussed bonds. The results of mentioned combinations are presented in Table 4-6.

Table 1: Soil complementary data at Aliabad industrial complex

Depth (cm)	Layers classification	pH	Natural density (g/cm ³)	Dry density (g/cm ³)	Moisture (%)	Dissolved chloride (Cl)	Solvable sulphate (SO ₃)
0	GW-GM	8.64	1.83	1.78	5.1	0.05	0.39
20	GM	8.52	1.60	1.56	5.6	0.06	0.35
40	SM	8.2	1.61	1.59	5.7	0.04	0.38
60	SC-SM	8.2	1.62	1.55	6.2	0.06	0.33
80	CL-ML	8.05	1.65	1.61	6.3	0.08	0.27
100	ML	7.98	1.7	1.66	6.6	0.09	0.25

GW-GM: Well-graded gravel with silt and sand
 GM: Silty gravel with sand
 SM: Silty sand
 SC-SM: Silty clayey sand
 CL-ML: Silty clay with sand
 ML: Sandy silt with gravel



Table 2: Metal concentrations in the seven leachate samples collected from bottom of soil column and initial leachate

Sample	Elements (mg/L)						
	Ni	Cu	Cd	Zn	Pb	Cr	Mn
Initial leachate	9.5	3.5	2.3	4.5	4.5	135	0.75
Leachate at day 5	0.138	0.289	0.044	0.163	0.579	0.63	0.39
Leachate at day 10	0.071	0.228	0.013	0.152	0.551	0.64	0.39
Leachate at day 15	0.095	0.269	0.012	0.129	0.548	0.71	0.39
Leachate at day 20	0.016	0.291	0.011	0.104	0.679	0.64	0.39
Leachate at day 25	0.040	0.335	0.028	0.092	0.599	0.63	0.39
Leachate at day 30	0.064	0.643	0.039	0.105	0.651	0.58	0.40
Leachate at day 35	0.120	0.321	0.055	0.074	0.479	0.54	0.39
Max.	0.138	0.643	0.055	0.163	0.679	0.71	0.40
Min.	0.016	0.228	0.011	0.074	0.479	0.54	0.39
SD*	0.043	0.138	0.017	0.032	0.067	0.053	0.003

* SD: Standard Diviation

Table 3: Concentration of Heavy metals in various soil depths

Sample	(mg/kg)						
	Ni	Cu	Cd	Zn	Pb	Cr	Mn
Intact soil	44	24	2.8	60	56	58	716
Soil at 0 cm (surface)	397	203	9.6	263	186	64	820
Soil at 20 cm	76	75	7.3	114	67	81	754
Soil at 40 cm	76	81	9	117	73	74	781
Soil at 60 cm	70	75	7.9	111	73	73	775
Soil at 80 cm	59	76	7	117	72	73	782
Soil at 100 cm	59	56	7.8	109	75	76	819
Max.	397	203	9.6	263	186	81	820
Min.	59	56	7	109	67	64	754
SD	134.532	53.924	1.003	61.076	46.617	5.540	26.051

Table 4: Loosely bonded ions in soils

Sample	(mg/kg)						
	Ni	Cu	Cd	Zn	Pb	Cr	Mn
Intact soil	5.2	4.675	2.3	8.425	35.28	7.75	45.25
Soil at 0 cm (surface)	218.925	91.725	5.82	111.575	117.366	8.5	59
Soil at 20 cm	10.325	6.725	5.75	8.275	42.3	9.25	44
Soil at 40 cm	10.375	11.15	6.3	10.7	43.125	9.5	57.25
Soil at 60 cm	10.325	5.075	5.53	7.6	42.375	10	351.25
Soil at 80 cm	6.575	11.6	4.9	11.5	44.225	7.75	41.5
Soil at 100 cm	8.325	10.5	4.575	5.15	46.05	8	47.5
Max.	218.925	91.725	6.3	111.575	117.366	10	351.25
Min.	6.575	5.075	4.575	5.15	42.3	7.75	41.5
SD	85.639	33.869	0.635	42.082	30.141	0.889	123.247



Table 5: Sulfides ions in soils

Sample	(mg/kg)						
	Ni	Cu	Cd	Zn	Pb	Cr	Mn
Intact soil	0	0	0.086	0	6.49	0	0
Soil at 0 cm(surface)	67.1	22.35	1.62	22.1	21.576	9.6	0
Soil at 20 cm	0	0	0	1.6	7.775	0.25	0
Soil at 40 cm	0	4.8	1.476	3.6	13.35	1.75	0
Soil at 60 cm	0	0	1.29	0	13.2	2	253.25
Soil at 80 cm	0	1.65	1.15	4.775	15.625	0.25	0
Soil at 100 cm	0	0	0.825	0	16.275	0.5	0
Max.	67.1	22.35	1.62	22.1	21.576	9.6	253.25
Min.	0	0	0	0	7.775	0.25	0
SD	27.393	8.798	0.587	8.429	4.530	3.613	103.388

Table 6: Organo-metallic ions in soils

Sample	Chemical segregation analysis results – organic- metallic bond (mg/kg)						
	Ni	Cu	Cd	Zn	Pb	Cr	Mn
Intact soil	7.625	9.95	0.078	3.15	9	8	9.75
Soil at 0 cm(surface)	5.075	11.825	1.73	2.775	29.908	8.7	24
Soil at 20 cm	3.95	11.325	0.5	4.125	10.775	11	9.75
Soil at 40 cm	4.7	10.775	1.053	3.675	13.975	7.75	9.75
Soil at 60 cm	6.25	10.45	0.924	3.4	13.475	10.25	10
Soil at 80 cm	4.2	11.4	0.825	3.85	11.125	8	9.75
Soil at 100 cm	3.25	9.9	1.6	2.725	11.58	8.25	10
Max.	6.25	11.825	1.73	4.125	29.908	11	24
Min.	3.25	9.9	0.5	2.725	10.775	7.75	9.75
SD	1.035	0.705	0.472	0.573	7.349	1.324	5.778

Normally, major portion of absorbed elements is found in loose bond that can be released into the environment as result of changes in physical and chemical changes in soil properties. Also, it should be stated that manganese sulfides in soil column are not formed. Therefore, it might be concluded that the environment is in oxidation condition.

CONCLUSION

According to the sample analysis of present study, the industrial leachates was found to have high levels of heavy metal and could be considered as hazardous and toxic to the environment. The results of present study clearly show that soil (i.e. natural bed soil) of studied area has a good potential to absorb heavy metals. This was evident from the obtained results in this study which indicated a

reduction of all the metals by adsorption within the first 5 days. However, such adsorption is associated as loosely bonded ions that can easily be released into the environment by minor changes in physical and chemical conditions of soil. Hence, along with site selection of hazardous wastes' landfill, the cation exchange capacity of soil at various depths must be carefully determined. It is also concluded that removal of metals by soil profile does not mean an ultimate fate for contamination. In other words, the chemical bonds of metals with various soil phases are vital.

ACKNOWLEDGMENTS

The authors acknowledge financial and scientific support provided by the Faculty of Environment at the University of Tehran.



REFERENCES

- Al-Khashman, O. A., (2004). Heavy metal distribution in dust, street dust and soil from the work place in Karak Industrial Estate, Jordan Atmos. Environ., 38 (39), 6803-6812 (10 pages).
- Allen, A. R., (2001). Containment landfills: The myth of sustainability. J. Eng. Geol., 60 (1-4), 3-19 (17 pages).
- Aucott, M., (2006). The fate of heavy metals in landfills: A Review. Prepared for the industrial ecology, pollution prevention and the NY-NJ harbor. Project of the New York Academy of Sciences.
- Bagchi, A., (1989). Design, construction and monitoring of sanitary landfills. John Wiley and Sons, New York.
- Banat, K. M.; Howari, F. M.; Al-Hamad, A. A., (2005). Heavy metals in urban soils of central Jordan: Should we worry about their environmental risks? Environ. Res., 97 (3), 258-273 (16 pages).
- Barrett, A.; Lawlor, J., (1995). The economics of waste management in Ireland. Economic and Social Research Institute, Dublin.
- Bulut, Y.; Baysal, Z., (2006). Removal of Pb (II) from wastewater using wheat bran. J. Environ. Manage., 78 (2), 107- 113 (7 pages).
- Charlesworth, S.; Everett, M.; McCarthy, R.; Ordonez, A.; De Miguel, E., (2003). A comparative study of heavy metals concentration and distribution in deposited street dusts in a large and a small urban area: Birmingham and Coventry, west Midlands, UK. Environ. Int., 29 (5), 563-573 (11 pages).
- Chen, T. B.; Zheng, Y. M.; Lei, M.; Huang, Z. C.; Wu, H. T.; Chen, H.; Fan, K. K.; Yu, K.; Wu, X.; Tian, Q. Z., (2005). Assessment of heavy metal pollution in surface soils of urban parks in Beijing, China. Chemosphere, 60 (4), 542-551 (10 pages).
- Cheng, Z.; Zheng, Y.; Mortlock, R.; Van Geen, A., (2004). Rapid multi-element analysis of groundwater by high-resolution inductively coupled plasma mass spectrometry. Anal. Bioanal. Chem., 379 (3), 512-518 (7 pages).
- Christensen, T. H.; Kjeldsen, P.; Bjerg, P. L.; Jensen, D. L.; Christensen, J. B.; Baun, A.; Albrechtsen, H. J.; Heron, G., (2001). Biogeochemistry of landfill leachate plumes. Appl. Geochem., 16 (7-8), 659-718 (59 pages).
- EPA, (1992a). Integrated risk information system (IRIS). National centre for environmental assessment, Office of research and development, United State Environmental Protection Agency, Washington D.C.
- EPA, (1992b). Flame atomic absorption spectrophotometry. Environmental Protection Agency. Revision 2.
- EPA, (2007). Acid digestion of aqueous samples and extracts for total metals for analysis by FLAA or ICP Spectroscopy. Environmental Protection Agency. Method # 3010A, Revision 1.
- Fang, H. Y., (1995). Bacteria and tree root attack on liners, in: Sarby, R. W., (Ed.), Composition of leachate from waste disposal sites. Waste disposal by landfill: Green '93 - Proceedings of a symposium on Geotechnics Related to the European Environment, Bolton, UK, Balkema, Rotterdam, 215-221 (7 pages).
- Fatta, D.; Papadopoulos, A.; Loizidou, M., (1999). A Study on the landfill leachate and its impact on the ground water quality of the greater area. Environ. Geochem. Hlth., 21 (2), 175-190 (16 pages).
- Freeze, R. A.; Cherry, J. A., (1979). Ground Water. Prentice - Hall Inc., 604.
- Futta, D.; Yoscoc, C.; Haralambous, K. J.; Loizidou, M., (1997). An assessment of the effect of landfill leachate on groundwater quality. 6th. Int. landfill symposium. S. Margherita di Pule, Gagliari, Italy. 181-187 (7 pages).
- Gharaibeh, S. H.; Masad, A., (1989). Die problematite der ahablbeseitigung in Jordan. Fallstudie fur enstasicklungs lander. Wasser und Boden, 10, 620-622 (3 pages).
- Karbassi, A. R.; Monavari, S. M.; Bidhendi, G. R. N.; Nouri, J.; Nematpour, K., (2008). Metal pollution assessment of sediment and water in the Shur River. Environ. Monitor. Assess., 147 (1-3), 107-116 (10 pages).
- Langston, W., (1990). Toxic effects of metals and the incidence of metal pollution in marine coastal ecosystem, in: Furness, R. W.; Rainbow P. S., (Eds.), Heavy metals in the marine environment, Boca Raton. CRC Press Inc. 101-122 (22 pages).
- Lee, G. F.; Jones L. A., (1993). Groundwater pollution by municipal landfills: Leachate composition, detection and water quality significance, Proc. Sardinia '93 IV International Landfill Symposium, Sardinia, Italy, 1093-1103 (11 pages).
- Lee, G. F.; Jones, R. A.; Ray, C., (1986). Sanitary landfill leachate recycle. Biocycle., 27 (1), 36-38 (3 pages).
- Lewis, R. J., (1991). Hazardous chemicals desk reference, 2nd. Eds., Van Nostrand Reinhold, New York, USA, 1-71 (71 pages).
- Longe, E. O.; Enekwechi, L. O., (2007). Investigation on potential groundwater impacts and influence of local hydrology on natural attenuation of leachate at a municipal landfill. Int. J. Environ. Sci. Tech., 4 (1), 133-140 (8 pages).
- Low, K. S.; Lee, G. K.; Liew, S. C., (2000). Sorption of cadmium and lead from aqueous solutions by spent grain. Proc. Biochem., 36 (1), 59-64 (6 pages).
- Mahvi, A. H., (2008). Application of agricultural fibers in pollution removal from aqueous solution. Int. J. Environ. Sci. Tech., 5 (2), 275-285 (11 pages).
- Malakootian, M.; Nouri, J.; Hossaini, H., (2009). Removal of heavy metals from paint industries wastewater using Leca as an available adsorbent. Int. J. Environ. Sci. Tech., 6 (2), 183-190 (8 pages).
- Nameni, M.; Alavi Moghadam, M. R.; Arami, M., (2008). Adsorption of hexavalent chromium from aqueous solutions by wheat bran. Int. J. Environ. Sci. Tech., 5 (2), 161-168 (8 pages).
- Nouri, J.; Mahvi, A. H.; Jahed, G. R.; Babaei, A. A., (2008). Regional distribution pattern of groundwater heavy metals resulting from agricultural activities. Environ. Geol. 55 (6), 1337-1343 (7 pages).
- Okafor, E. Ch.; Opuene, K., (2007). Preliminary assessment of trace metals and polycyclic aromatic hydrocarbons in the sediments. Int. J. Environ. Sci. Tech., 4 (2), 233-240 (8 pages).
- Panjeshahi, M. H.; Ataei, A., (2008). Application of an environmentally optimum cooling water system design in water and energy conservation. Int. J. Environ. Sci. Tech., 5 (2), 251-262 (12 pages).



Heavy metals adsorption in industrial waste landfill

- Pekey, H., (2006). The distribution and sources of heavy metals in Izmit Bay surface sediments affected by a polluted stream. *Mar. Pollut. Bull.*, 52 (10), 1197-1208 (**12 pages**).
- Qasim, S. R.; Chiang, W., (1994). Sanitary landfill leachate. CRC Press.
- Suthar, S.; Singh, S., (2008). Vermicomposting of domestic waste by using two epigeic earthworms (*Perionyx excavatus* and *perionyx sansibaricus*). *Int. J. Environ. Sci. Tech.*, 5 (1), 99-106 (**8 pages**).
- Vasanthi, P.; Kaliappan, S.; Srinivasaraghavan, R., (2008). Impact of poor solid waste management on ground water. *Environ. Monit. Assess.*, 143 (1-3), 227-238 (**11 pages**).
- Zheng, C.; Bennett, G. D.; Andrews, C. B., (1991). Analysis of groundwater remedial alternatives of a superfund site. *Groundwater*, 29 (6), 838-848 (**11 pages**).

AUTHOR (S) BIOSKETCHES

Esmacili Bidhendi, M., M.Sc., Graduate Faculty of the Environment, University of Tehran, P. O. Box 14155-6135, Tehran, Iran.
Email: m.esmaieli@gmail.com

Karbassi, A. R., Ph.D., Assistant Professor, Graduate Faculty of the Environment, University of Tehran, P. O. Box 14155-6135, Tehran, Iran. Email: arkarbassi738@yahoo.com

Baghvand, A., Ph.D, Assistant Professor, Graduate Faculty of the Environment, University of Tehran, P. O. Box 14155-6135, Tehran, Iran.
Email: baghvand@ut.ac.ir

Saeedi, M., Ph.D, Assistant Professor, Environmental Research Laboratory, School of Civil Engineering, Iran University of Science and Technology, Tehran, Iran.
Email: msaeedi@iust.ac.ir

Pejman, A. H., M.Sc., Graduate Faculty of the Environment, University of Tehran, P. O. Box 14155-6135, Tehran, Iran.
Email: ahp9325@gmail.com

How to cite this article: (Harvard style)

Esmacili Bidhendi, M.; Karbassi, A. R.; Baghvand, A.; Saeedi, M.; Pejman, A. H., (2010). Potential of natural bed soil in adsorption of heavy metals in industrial waste landfill. Int. J. Environ. Sci. Tech., 7 (3), 545-552.

