

Singular foliations for M-theory compactification

Elena Mirela Babalic^{a,b} and Calin Iuliu Lazaroiu^c

^a*Department of Theoretical Physics, National Institute of Physics and Nuclear Engineering, Str. Reactorului no. 30, P.O.BOX MG-6, Postcode 077125, Bucharest-Magurele, Romania*

^b*Department of Physics, University of Craiova, 13 Al. I. Cuza Str., Craiova 200585, Romania*

^c*Center for Geometry and Physics, Institute for Basic Science (IBS), Pohang 790-784, Republic of Korea*

E-mail: mbabalic@theory.nipne.ro, calin@ibs.re.kr

ABSTRACT: We use the theory of singular foliations to study $\mathcal{N} = 1$ compactifications of eleven-dimensional supergravity on eight-manifolds M down to AdS_3 spaces, allowing for the possibility that the internal part ξ of the supersymmetry generator is chiral on some locus \mathcal{W} which does not coincide with M . We show that the complement $M \setminus \mathcal{W}$ must be a *dense* open subset of M and that M admits a singular foliation $\bar{\mathcal{F}}$ endowed with a longitudinal G_2 structure and defined by a closed one-form ω , whose geometry is determined by the supersymmetry conditions. The singular leaves are those leaves which meet \mathcal{W} . When ω is a Morse form, the chiral locus is a finite set of points, consisting of isolated zero-dimensional leaves and of conical singularities of seven-dimensional leaves. In that case, we describe the topology of $\bar{\mathcal{F}}$ using results from Novikov theory. We also show how this description fits in with previous formulas which were extracted by exploiting the $\text{Spin}(7)_\pm$ structures which exist on the complement of \mathcal{W} .

KEYWORDS: Differential and Algebraic Geometry, Flux compactifications, M-Theory

ARXIV EPRINT: [1411.3497](https://arxiv.org/abs/1411.3497)

Contents

1	Introduction	1
2	Basics	3
3	Parameterizing a Majorana spinor on M	5
3.1	Globally valid parameterization	5
3.2	The chirality decomposition of M	7
3.3	A topological no-go theorem	8
3.4	The singular distribution \mathcal{D}	9
3.5	Spinor parameterization and G_2 structure on the non-chiral locus	10
3.6	Spinor parameterization and $\text{Spin}(7)_\pm$ structures on the loci \mathcal{U}^\pm	11
3.7	Comparing spinors and G structures on the non-chiral locus	16
3.8	The singular foliation of M defined by \mathcal{D}	17
4	Relating the G_2 and $\text{Spin}(7)$ approaches on the non-chiral locus	18
4.1	The G_2 and $\text{Spin}(7)_\pm$ decompositions of $\Omega^4(\mathcal{U})$	19
4.2	The G_2 and $\text{Spin}(7)_\pm$ parameterizations of F	20
4.3	Relating the G_2 torsion classes to the Lee form and characteristic torsion of the $\text{Spin}(7)_\pm$ structures	22
4.4	Relation to previous work	23
5	Description of the singular foliation in the Morse case	24
5.1	Types of singular points	24
5.2	The regular and singular foliations defined by a Morse 1-form	25
5.3	Behavior of the singular leaves near singular points	27
5.4	Combinatorics of singular leaves	29
5.5	Homology classes of compact leaves	32
5.6	The Novikov decomposition of M	32
5.7	The foliation graph	34
5.8	The fundamental group of the leaf space	37
5.9	On the relation to compactifications of M-theory on 7-manifolds	38
5.10	A non-commutative description of the leaf space?	40
6	Conclusions and further directions	41
A	Proof of the topological no-go theorem	42
B	The case $\kappa = 0$	43
B.1	When M is compact	43
B.2	When M is non-compact	46
C	Comparison with the results of [1]	47
D	Generalized bundles and generalized distributions	52

E Some topological properties of singular foliations defined by a Morse one-form	55
E.1 Some topological invariants of M	55
E.2 Estimate for the number of splitting saddle points	55
E.3 Estimates for c and N_{\min}	56
E.4 Criteria for existence and number of homologically independent compact leaves	56
E.5 Generic forms	56
E.6 Exact forms	57
E.7 Behavior under exact perturbations	57

1 Introduction

$\mathcal{N} = 1$ flux compactifications of eleven-dimensional supergravity on eight-manifolds M down to AdS_3 spaces [1, 2] provide a vast extension of the better studied class of compactifications down to 3-dimensional Minkowski space [3–5], having the advantage that they are already consistent at the classical level [2]. They form a useful testing ground for various proposals aimed at providing unified descriptions of flux backgrounds [6] and may be relevant to recent attempts to gain a better understanding of F-theory [7]. When the internal part ξ of the supersymmetry generator is everywhere non-chiral, such backgrounds can be studied [8] using foliations endowed with longitudinal G_2 structures, an approach which permits a geometric description of the supersymmetry conditions while providing powerful tools for studying the topology of such backgrounds.

In this paper, we extend the results of [8] to the general case when the internal part ξ of the supersymmetry generator is allowed to become chiral on some locus $\mathcal{W} \subset M$. Assuming that $\mathcal{W} \neq M$, i.e. that ξ is not everywhere chiral, we show that, at the classical level, the Einstein equations imply that the chiral locus \mathcal{W} must be a set with empty interior, which therefore is negligible with respect to the Lebesgue measure of the internal space. As a consequence, the behavior of geometric data along this locus can be obtained from the non-chiral locus $\mathcal{U} \stackrel{\text{def.}}{=} M \setminus \mathcal{W}$ through a limiting process. The geometric information along the non-chiral locus \mathcal{U} is encoded [8] by a regular foliation \mathcal{F} which carries a longitudinal G_2 structure and whose geometry is determined by the supersymmetry conditions in terms of the supergravity four-form field strength. When $\emptyset \neq \mathcal{W} \subsetneq M$, we show that \mathcal{F} extends to a singular foliation $\bar{\mathcal{F}}$ of the whole manifold M by adding leaves which are allowed to have singularities at points belonging to \mathcal{W} . This singular foliation “integrates” a cosmooth¹ [11–14] singular distribution \mathcal{D} (a.k.a. generalized sub-bundle of TM), defined as the kernel distribution of a closed one-form ω which belongs to a cohomology class $\mathfrak{f} \in H^1(M, \mathbb{R})$ determined by the supergravity four-form field strength. The set of zeroes of ω coincides with the chiral locus \mathcal{W} . In the most general case, $\bar{\mathcal{F}}$ can be viewed as a

¹Note that \mathcal{D} is *not* a singular distribution in the sense of Stefan-Sussmann [9, 10] (it is cosmooth rather than smooth). See appendix D.

Haefliger structure [15] on M . The singular foliation $\bar{\mathcal{F}}$ carries a longitudinal G_2 structure, which is allowed to degenerate at the singular points of singular leaves. On the non-chiral locus \mathcal{U} , the problem can be studied using the approach of [8] or the approach advocated in [1], which makes use of two $\text{Spin}(7)_\pm$ structures. We show explicitly how one can translate between these two approaches and prove that the results of [8] agree with those of [1] along this locus.

While the topology of singular foliations defined by a closed one-form can be extremely complicated in general, the situation is better understood in the case when ω is a Morse one-form. The Morse case is generic in the sense that such 1-forms constitute an open and dense subset of the set of all closed one-forms belonging to the cohomology class \mathfrak{f} . In the Morse case, the singular foliation $\bar{\mathcal{F}}$ can be described using the *foliation graph* [16–18] associated to the corresponding decomposition of M (see [18–20] and [21]–[29]), which provides a combinatorial way to encode some important aspects of the foliation’s topology — up to neglecting the information contained in the so-called *minimal components* of the decomposition, components which should possess an as yet unexplored non-commutative geometric description. This provides a far-reaching extension of the picture found in [8] for the everywhere non-chiral case $\mathcal{U} = M$, a case which corresponds to the situation when the foliation graph is reduced to either a circle (when \mathcal{F} has compact leaves, being a fibration over S^1) or to a single so-called exceptional vertex (when \mathcal{F} has non-compact dense leaves, being a minimal foliation). In the minimal case of the backgrounds considered [8], the exceptional vertex corresponds to a noncommutative torus which encodes the noncommutative geometry [30, 31] of the leaf space.

The paper is organized as follows. Section 2 gives a brief review of the class of compactifications we consider, in order to fix notations and conventions. Section 3 discusses a geometric characterization of Majorana spinors ξ on M which is inspired by the rigorous approach developed in [32–34] for the method of bilinears [35], in the case when the spinor ξ is allowed to be chiral at some loci. It also gives the Kähler-Atiyah parameterizations of this spinor which correspond to the approach of [8] and to that of [1] and describes the relevant G -structures using both spinors and idempotents in the Kähler-Atiyah algebra of M . In the same section, we give the general description of the singular foliation $\bar{\mathcal{F}}$ as the Haefliger structure defined by the closed one-form ω . Section 4 describes the relation between the G_2 and $\text{Spin}(7)_\pm$ parameterizations of the fluxes as well as the relation between the torsion classes of the leafwise G_2 structure and the Lee form and characteristic torsion of the $\text{Spin}(7)_\pm$ structures defined on the non-chiral locus. The same section gives the comparison of the approach of [8] with that of [1] along that locus. Section 5 discusses the topology of the singular foliation $\bar{\mathcal{F}}$ in the Morse case while section 6 concludes. The appendices contain various technical details.

Notations and conventions. Throughout this paper, M denotes an oriented, connected and compact smooth manifold (which will mostly be of dimension eight), whose unital commutative \mathbb{R} -algebra of smooth real-valued functions we denote by $\Omega^0(M) = C^\infty(M, \mathbb{R})$. Given a subset A of M , we let \bar{A} denote the closure of A in M (taken with respect to the manifold topology of M). The *large topological frontier* (also called *topolog-*

ical boundary) of A is defined as $\text{Fr}(A) \stackrel{\text{def.}}{=} \bar{A} \setminus \text{Int}(A)$, where $\text{Int}(A)$ denotes the interior of A . The *small topological frontier* is $\text{fr}(A) \stackrel{\text{def.}}{=} \bar{A} \setminus A$. Notice that $\text{fr}(A) \subseteq \text{Fr}(A)$ and that $\text{fr}(A) = \text{Fr}(A)$ when A is open, in which case we speak simply of the *frontier* of A . All fiber bundles we consider are smooth.² We use freely the results and notations of [8, 32–34], with the same conventions as there. To simplify notation, we write the geometric product \diamond of [32–34] simply as juxtaposition while indicating the wedge product of differential forms through \wedge . If $\mathcal{D} \subset TM$ is a singular (a.k.a. generalized) distribution on M and \mathcal{U} is an open subset of M such that $\mathcal{D}|_{\mathcal{U}}$ is a regular Frobenius distribution (see appendix D), we let $\Omega_{\mathcal{U}}(\mathcal{D}) = \Gamma(\mathcal{U}, \wedge(\mathcal{D}|_{\mathcal{U}})^*)$ denote the $\mathcal{C}^\infty(\mathcal{U}, \mathbb{R})$ -module of $\mathcal{D}|_{\mathcal{U}}$ -longitudinal differential forms defined on \mathcal{U} . When $\dim M = 8$, then for any 4-form $\omega \in \Omega^4(M)$ we let $\omega^\pm \stackrel{\text{def.}}{=} \frac{1}{2}(\omega \pm *\omega)$ denote the selfdual and anti-selfdual parts of ω (namely, $*\omega^\pm = \pm\omega^\pm$). When M is eight-dimensional, we let $\Omega^{4\pm}(M)$ denote the spaces of selfdual and anti-selfdual four-forms, respectively. We use the “Det” convention for the wedge product and the corresponding “Perm” convention for the symmetric product. Hence given a local coframe e^a of M , we have:

$$\begin{aligned} e^{a_1} \wedge \dots \wedge e^{a_k} &\stackrel{\text{def.}}{=} \sum_{\sigma \in S_k} \epsilon(\sigma) e^{a_{\sigma(1)}} \otimes \dots \otimes e^{a_{\sigma(k)}}, \\ e^{a_1} \odot \dots \odot e^{a_k} &\stackrel{\text{def.}}{=} \sum_{\sigma \in S_k} e^{a_{\sigma(1)}} \otimes \dots \otimes e^{a_{\sigma(k)}}, \end{aligned} \tag{1.1}$$

without prefactors of $\frac{1}{k!}$ in the right hand side, where S_k is the symmetric group on k letters and $\epsilon(\sigma)$ denotes the signature of a permutation σ . This is the convention used, for example, in [36]. We use $\text{Sym}_0^2(T^*M)$ to denote the space of traceless symmetric covariant 2-tensors on M and $\text{Sym}_{\mathcal{U},0}^2(\mathcal{D}^*)$ to denote the space of traceless symmetric covariant 2-tensors defined on \mathcal{U} and which are longitudinal to the Frobenius distribution $\mathcal{D}|_{\mathcal{U}}$, when \mathcal{D} is as above. By definition, a $\text{Spin}(7)_+$ structure on M is a $\text{Spin}(7)$ structure with respect to the orientation chosen for M while a $\text{Spin}(7)_-$ structure is a $\text{Spin}(7)$ structure with respect to the opposite orientation.

2 Basics

We start with a brief review of the set-up, in order to fix notation. As in [1, 2], we consider 11-dimensional supergravity [37] on an eleven-dimensional connected and paracompact spin manifold \mathbf{M} with Lorentzian metric \mathbf{g} (of ‘mostly plus’ signature). Besides the metric, the classical action of the theory contains the three-form potential with four-form field strength $\mathbf{G} \in \Omega^4(\mathbf{M})$ and the gravitino Ψ , which is a Majorana spinor of spin 3/2. The bosonic part of the action takes the form:

$$S_{\text{bos}}[\mathbf{g}, \mathbf{C}] = \frac{1}{2\kappa_{11}^2} \int_{\mathbf{M}} R\nu - \frac{1}{4\kappa_{11}^2} \int_{\mathbf{M}} \left(\mathbf{G} \wedge \star \mathbf{G} + \frac{1}{3} \mathbf{C} \wedge \mathbf{G} \wedge \mathbf{G} \right),$$

where κ_{11} is the gravitational coupling constant in eleven dimensions, ν and R are the volume form and the scalar curvature of \mathbf{g} and $\mathbf{G} = d\mathbf{C}$. For supersymmetric bosonic

²The “generalized bundles” [11, 12] considered occasionally in this paper are *not* fiber bundles.

classical backgrounds, both the gravitino and its supersymmetry variation must vanish, which requires that there exist at least one solution η to the equation:

$$\delta_\eta \Psi \stackrel{\text{def.}}{=} \mathfrak{D}\eta = 0, \tag{2.1}$$

where \mathfrak{D} denotes the supercovariant connection. The eleven-dimensional supersymmetry generator η is a Majorana spinor (real pinor) of spin 1/2 on \mathbf{M} .

As in [1, 2], consider compactification down to an AdS₃ space of cosmological constant $\Lambda = -8\kappa^2$, where κ is a positive real parameter — this includes the Minkowski case as the limit $\kappa \rightarrow 0$. Thus $\mathbf{M} = N \times M$, where N is an oriented 3-manifold diffeomorphic to \mathbb{R}^3 and carrying the AdS₃ metric g_3 while M is an oriented, compact and connected Riemannian eight-manifold whose metric we denote by g . The metric on \mathbf{M} is a warped product:

$$ds^2 = e^{2\Delta} ds^2 \quad \text{where} \quad ds^2 = ds_3^2 + g_{mn} dx^m dx^n. \tag{2.2}$$

The warp factor Δ is a smooth real-valued function defined on M while ds_3^2 is the squared length element of the AdS₃ metric g_3 . For the field strength \mathbf{G} , we use the ansatz:

$$\mathbf{G} = \nu_3 \wedge \mathbf{f} + \mathbf{F}, \quad \text{with} \quad \mathbf{F} \stackrel{\text{def.}}{=} e^{3\Delta} F, \quad \mathbf{f} \stackrel{\text{def.}}{=} e^{3\Delta} f, \tag{2.3}$$

where $f \in \Omega^1(M)$, $F \in \Omega^4(M)$ and ν_3 is the volume form of (N, g_3) . For η , we use the ansatz:

$$\eta = e^{\frac{\Delta}{2}} (\zeta \otimes \xi),$$

where ξ is a Majorana spinor of spin 1/2 on the internal space (M, g) (a section of the rank 16 real vector bundle S of indefinite chirality real pinors) and ζ is a Majorana spinor on (N, g_3) .

Assuming that ζ is a Killing spinor on the AdS₃ space (N, g_3) , the supersymmetry condition (2.1) is equivalent with the following system for ξ :

$$\mathbb{D}\xi = 0, \quad Q\xi = 0, \tag{2.4}$$

where

$$\mathbb{D}_X = \nabla_X^S + \frac{1}{4}\gamma(X \lrcorner F) + \frac{1}{4}\gamma((X_\sharp \wedge f)\nu) + \kappa\gamma(X \lrcorner \nu), \quad X \in \Gamma(M, TM)$$

is a linear connection on S (here ∇^S is the connection induced on S by the Levi-Civita connection of (M, g) , while ν is the volume form of (M, g)) and

$$Q = \frac{1}{2}\gamma(d\Delta) - \frac{1}{6}\gamma(\iota_f \nu) - \frac{1}{12}\gamma(F) - \kappa\gamma(\nu)$$

is a globally-defined endomorphism of S . As in [1, 2], we do not require that ξ has definite chirality.

The set of solutions of (2.4) is a finite-dimensional \mathbb{R} -linear subspace $\mathcal{K}(\mathbb{D}, Q)$ of the infinite-dimensional vector space $\Gamma(M, S)$ of smooth sections of S . Up to rescalings by smooth nowhere-vanishing real-valued functions defined on M , the vector bundle S has

two admissible pairings \mathcal{B}_\pm (see [34, 38, 39]), both of which are symmetric but which are distinguished by their types $\epsilon_{\mathcal{B}_\pm} = \pm 1$. Without loss of generality, we choose to work with $\mathcal{B} \stackrel{\text{def.}}{=} \mathcal{B}_+$. We can in fact take \mathcal{B} to be a scalar product on S and denote the corresponding norm by $\| \cdot \|$ (see [32, 33] for details). Requiring that the background preserves exactly $\mathcal{N} = 1$ supersymmetry amounts to asking that $\dim \mathcal{K}(\mathbb{D}, Q) = 1$. It is not hard to check [32] that \mathcal{B} is \mathbb{D} -flat:

$$d\mathcal{B}(\xi', \xi'') = \mathcal{B}(\mathbb{D}\xi', \xi'') + \mathcal{B}(\xi', \mathbb{D}\xi''), \quad \forall \xi', \xi'' \in \Gamma(M, S). \quad (2.5)$$

Hence any solution of (2.4) which has unit \mathcal{B} -norm at a point will have unit \mathcal{B} -norm at every point of M and we can take the internal part ξ of the supersymmetry generator to be everywhere of norm one.

3 Parameterizing a Majorana spinor on M

3.1 Globally valid parameterization

Fixing a Majorana spinor $\xi \in \Gamma(M, S)$ which is everywhere of \mathcal{B} -norm one, consider the inhomogeneous differential form:

$$\check{E}_{\xi, \xi} = \frac{1}{16} \sum_{k=0}^8 \check{\mathbf{E}}_{\xi, \xi}^{(k)} \in \Omega(M), \quad (3.1)$$

whose rescaled rank components have the following expansions in any local orthonormal coframe $(e^a)_{a=1\dots 8}$ of M defined on some open subset U :

$$\check{\mathbf{E}}_{\xi, \xi}^{(k)} =_U \frac{1}{k!} \mathcal{B}(\xi, \gamma_{a_1 \dots a_k} \xi) e^{a_1 \dots a_k} \in \Omega^k(M).$$

The conditions:

$$\check{E}^2 = \check{E}, \quad \mathcal{S}(\check{E}) = 1, \quad \tau(\check{E}) = \check{E} \quad (3.2)$$

encode the fact that an inhomogeneous form $\check{E} \stackrel{\text{def.}}{=} \check{E}_{\xi, \xi}$ is of the type (3.1) for some Majorana spinor ξ which is everywhere of norm one. As a result of the last condition in (3.2), the non-zero components of \check{E} have ranks $k = 0, 1, 4, 5$ and we have $\mathcal{S}(\check{E}_{\xi, \xi}) = \check{\mathbf{E}}_{\xi, \xi}^{(0)} = \|\xi\|^2 = 1$, where \mathcal{S} is the canonical trace of the Kähler-Atiyah algebra. Hence:

$$\check{E} = \frac{1}{16} (1 + V + Y + Z + b\nu), \quad (3.3)$$

where we introduced the notations:

$$V \stackrel{\text{def.}}{=} \check{\mathbf{E}}^{(1)}, \quad Y \stackrel{\text{def.}}{=} \check{\mathbf{E}}^{(4)}, \quad Z \stackrel{\text{def.}}{=} \check{\mathbf{E}}^{(5)}, \quad b\nu \stackrel{\text{def.}}{=} \check{\mathbf{E}}^{(8)}. \quad (3.4)$$

Here, b is a smooth real valued function defined on M and ν is the volume form of (M, g) , which satisfies $\|\nu\| = 1$; notice the relation $\mathcal{S}(\nu \check{E}_{\xi, \xi}) = b$. On a small enough open subset $U \subset M$ supporting a local coframe (e^a) of M , one has the expansions:

$$\begin{aligned} V &= _U \mathcal{B}(\xi, \gamma_a \xi) e^a, & Y &= _U \frac{1}{4!} \mathcal{B}(\xi, \gamma_{a_1 \dots a_4} \xi) e^{a_1 \dots a_4}, \\ Z &= _U \frac{1}{5!} \mathcal{B}(\xi, \gamma_{a_1 \dots a_5} \xi) e^{a_1 \dots a_5}, & b &= _U \mathcal{B}(\xi, \gamma(\nu)\xi). \end{aligned} \quad (3.5)$$

One finds [32] that (3.2) is equivalent with the following relations which hold globally on M :

$$\begin{aligned} \|V\|^2 &= 1 - b^2 \geq 0, & \|Y^\pm\|^2 &= \frac{7}{2}(1 \pm b)^2, \\ \iota_V(*Z) &= 0, & \iota_V Z &= Y - b * Y, \\ (\iota_\alpha(*Z)) \wedge (\iota_\beta(*Z)) \wedge (*Z) &= -6\langle \alpha \wedge V, \beta \wedge V \rangle \iota_V \nu, & \forall \alpha, \beta \in \Omega^1(M). \end{aligned} \quad (3.6)$$

Notice that the first relation in the second row is equivalent with $V \wedge Z = 0$, which means that V and Z commute in the Kähler-Atiyah algebra of (M, g) .

Remark. Let (R) denote the second relation (namely $\iota_V Z = Y - b * Y$) on the second row of (3.6). Separating the selfdual and anti-selfdual parts shows that (R) is *equivalent* with the following two conditions:

$$(\iota_V Z)^\pm = (1 \mp b)Y^\pm. \quad (3.7)$$

Proposition. Relations (3.6) imply that the following normalization conditions hold globally on M :

$$\|Y\|^2 = 7(1 + b^2), \quad \|Z\|^2 = 7(1 - b^2). \quad (3.8)$$

Proof. The first equation in (3.8) follows from the last relations on the first row of (3.6) by noticing that $\|Y\|^2 = \|Y^+\|^2 + \|Y^-\|^2$ (since $\langle Y^+, Y^- \rangle = 0$). We have:

$$\|\iota_V Z\|^2 = \|*\iota_V Z\|^2 = \|V \wedge (*Z)\|^2 = \|V\|^2 \|*Z\|^2 = \|V\|^2 \|Z\|^2, \quad (3.9)$$

where in the middle equality we used the first equation on the second row of (3.6), which tells us that $*Z$ is orthogonal on V . The second equation in (3.8) now follows from (3.9) and from the identity:

$$\|\iota_V Z\|^2 = (1 - b)^2 \|Y^+\|^2 + (1 + b)^2 \|Y^-\|^2 = 7(1 - b^2) = 7\|V\|^2,$$

where we used (3.7) and both relations in the first row of (3.6). ■

The twisted selfdual and twisted anti-selfdual parts of \check{E} . The identity $\nu^2 = 1$ implies that the elements:

$$R^\pm \stackrel{\text{def.}}{=} \frac{1}{2}(1 \pm \nu)$$

are complementary idempotents in the Kähler-Atiyah algebra:

$$(R^\pm)^2 = R^\pm, \quad R^\pm R^\mp = 0, \quad R^+ + R^- = \text{id}_{\Omega(M)}. \quad (3.10)$$

The (anti)selfdual part of a four-form $\omega \in \Omega^4(M)$ can be expressed as:

$$\omega_\pm = R^\pm \omega.$$

Notice that this relation also gives the twisted (anti)selfdual parts [32] of an inhomogeneous form $\omega \in \Omega(M)$. The identities:

$$Y R^\pm = R^\pm Y = Y^\pm, \quad (1 + b\nu)R^\pm = (1 \pm b)R^\pm$$

allow us to compute the twisted selfdual part \check{E}^+ and twisted anti-selfdual part \check{E}^- of \check{E} :

$$\check{E}^\pm = \check{E}R^\pm = \frac{1}{16} [(1 \pm b + V + Z)R_\pm + Y^\pm] \in \Omega(M). \quad (3.11)$$

The following decomposition holds globally on M :

$$\check{E} = \check{E}^+ + \check{E}^-.$$

3.2 The chirality decomposition of M

Let $S^\pm \subset S$ be the rank eight subbundles of S consisting of positive and negative chirality spinors (the eigen-subbundles of $\gamma(\nu)$ corresponding to the eigenvalues $+1$ and -1). Since $\gamma(\nu)$ is \mathcal{B} -symmetric, S^+ and S^- give a \mathcal{B} -orthogonal decomposition $S = S^+ \oplus S^-$. Decomposing a normalized spinor as $\xi = \xi^+ + \xi^-$ with $\xi^\pm \stackrel{\text{def.}}{=} \frac{1}{2}(\text{id}_S \pm \gamma(\nu))\xi \in \Gamma(M, S^\pm)$, we have:

$$\|\xi\|^2 = \|\xi^+\|^2 + \|\xi^-\|^2 = 1$$

and:

$$b = \mathcal{B}(\xi, \gamma(\nu)\xi) = \|\xi^+\|^2 - \|\xi^-\|^2.$$

These two relations give:

$$\|\xi^\pm\|^2 = \frac{1}{2}(1 \pm b). \quad (3.12)$$

Notice that b equals ± 1 at a point $p \in M$ iff $\xi_p \in S_p^\pm$. Since $\|V\|^2 = 1 - b^2$, the one-form V vanishes at p iff ξ_p is chiral i.e. iff $\xi_p \in S_p^+ \cup S_p^-$. Consider the *non-chiral locus* (an open subset of M):

$$\begin{aligned} \mathcal{U} &\stackrel{\text{def.}}{=} \{p \in M | \xi \notin S_p^+ \cup S_p^-\} = \{p \in M | \xi_p^+ \neq 0 \text{ and } \xi_p^- \neq 0\} \\ &= \{p \in M | V_p \neq 0\} = \{p \in M | |b(p)| < 1\}, \end{aligned}$$

and its closed complement, the *chiral locus*:

$$\begin{aligned} \mathcal{W} &\stackrel{\text{def.}}{=} M \setminus \mathcal{U} = \{p \in M | \xi_p \in S_p^+ \cup S_p^-\} = \{p \in M | \xi_p^+ = 0 \text{ or } \xi_p^- = 0\} \\ &= \{p \in M | V_p = 0\} = \{p \in M | |b(p)| = 1\}. \end{aligned}$$

The chiral locus \mathcal{W} decomposes further as a disjoint union of two closed subsets, the *positive and negative chirality loci*:

$$\mathcal{W} = \mathcal{W}^+ \sqcup \mathcal{W}^-,$$

where:

$$\mathcal{W}^\pm \stackrel{\text{def.}}{=} \{p \in M | \xi_p \in S_p^\pm\} = \{p \in M | b(p) = \pm 1\} = \{p \in M | \xi_p^\mp = 0\}.$$

The extreme cases $\mathcal{W}^+ = M$ or $\mathcal{W}^- = M$, as well as $\mathcal{W}^+ = \mathcal{W}^- = \emptyset$ are allowed. However, the case $\mathcal{U} = \emptyset$ with both \mathcal{W}^+ and \mathcal{W}^- nonempty (then $M = \mathcal{W}^+ \sqcup \mathcal{W}^-$) is forbidden (recall that b is smooth and hence continuous while M is connected). Since ξ does not vanish on M , we have:

$$\mathcal{U}^\pm \stackrel{\text{def.}}{=} \mathcal{U} \cup \mathcal{W}^\pm = \{p \in M | \xi_p^\pm \neq 0\}.$$

Remark. Since $|b| \leq 1$ on M , the sets \mathcal{W}^\pm (when non-empty) consist of critical points of b , namely the absolute maxima and minima of b on M . Hence the differential of b vanishes at every point of \mathcal{W} . In general \mathcal{W}^\pm can be quite ‘wild’ (they can be very far from being immersed submanifolds of M).

3.3 A topological no-go theorem

Recall that M is compact. The following result clarifies the kind of topologies of the chiral loci which are of physical interest.

Theorem. Assume that the supersymmetry conditions, the Bianchi identity and equations of motion for G as well as the Einstein equations are satisfied. There exist only the following four possibilities:

1. The set \mathcal{W}^+ coincides with M and hence \mathcal{W}^- and \mathcal{U} are empty. In this case, ξ is a chiral spinor of positive chirality which is covariantly constant on M and we have $\kappa = f = F = 0$ while Δ is constant on M .
2. The set \mathcal{W}^- coincides with M and hence \mathcal{W}^+ and \mathcal{U} are empty. In this case, ξ is a chiral spinor of negative chirality which is covariantly constant on M and we have $\kappa = f = F = 0$ while Δ is constant on M .
3. The set \mathcal{U} coincides with M and hence \mathcal{W}^+ and \mathcal{W}^- are empty.
4. At least one of the sets \mathcal{W}^+ or \mathcal{W}^- is non-empty but both of these sets have empty interior. In this case, \mathcal{U} is dense in M and the union $\mathcal{W} = \mathcal{W}^+ \cup \mathcal{W}^-$ coincides with the topological frontier $\text{Fr}(\mathcal{U}) = \text{fr}(\mathcal{U}) = \bar{\mathcal{U}} \setminus \mathcal{U}$ of \mathcal{U} .

The proof of the theorem is given in appendix A.


Remarks.

- The theorem is a strengthening of an observation originally made in [2] in the case when ξ is nowhere-chiral.
- The theorem holds in classical supergravity only. One may be able to avoid its conclusions by considering quantum corrections.
- Cases 1 and 2 correspond to the classical limit of the compactifications studied in [3–5]. Case 3 was studied in [2, 8].

The study of Case 4 is the focus of the present paper. Due to the theorem, we shall from now on assume that we are in this case, i.e. that \mathcal{W} is non-empty and that it coincides with the frontier of \mathcal{U} ; in particular, we can assume that the closure of \mathcal{U} coincides with M :

$$M = \bar{\mathcal{U}} = \mathcal{U} \sqcup \mathcal{W}, \quad \mathcal{W} = \text{Fr}\mathcal{U}.$$

In figure 1, we sketch the chirality decomposition in two sub-cases of Case 4, which correspond to the assumptions that the one-form $\omega \stackrel{\text{def.}}{=} 4\kappa e^{3\Delta}V$ is of Morse and Bott-Morse type, respectively.



(a) Sketch of the chiral loci in the Morse sub-case of Case 4 of the Theorem. In this case, each of \mathcal{W}^+ and \mathcal{W}^- is a finite set of points, with the points of \mathcal{W}^+ indicated in red and those of \mathcal{W}^- indicated in blue.

(b) Sketch of \mathcal{W}^\pm in the Bott-Morse sub-case of Case 4 of the Theorem. The connected components of \mathcal{W} are submanifolds of various dimensions, shown respectively in red and blue for \mathcal{W}^+ and \mathcal{W}^- .

Figure 1. Sketch of chiral loci in two sub-cases of Case 4 of the Theorem, for the case of a two-dimensional manifold M . The non-chiral locus \mathcal{U} is the complement of \mathcal{W} in M and is indicated by white space, after performing appropriate cuts which allow one to map M to some region of the plane which is not indicated explicitly. The figures should be interpreted with care in our case $\dim M = 8$.

3.4 The singular distribution \mathcal{D}

The one-form V determines a singular (a.k.a. generalized) distribution \mathcal{D} (generalized sub-bundle of TM) which is defined through:

$$\mathcal{D}_p \stackrel{\text{def.}}{=} \ker V_p, \quad \forall p \in M.$$

This singular distribution is *cosmooth* (rather than smooth) in the sense of [11] (see appendix D). Notice that \mathcal{D} is smooth iff ξ is everywhere non-chiral — i.e. iff $\mathcal{W} = \emptyset$, which is the case studied in [8]; in that case, \mathcal{D} is a regular Frobenius distribution. Since in this paper we assume $\mathcal{W} \neq \emptyset$, it follows that \mathcal{D} is *not* a singular distribution in the sense of Stefan-Sussmann [9, 10]. The set of regular points of \mathcal{D} equals the non-chiral locus \mathcal{U} and we have:

$$\begin{aligned} \text{rk} \mathcal{D}_p &= 7 & \text{when } p \in \mathcal{U}, \\ \text{rk} \mathcal{D}_p &= 8 & \text{when } p \in \mathcal{W}. \end{aligned}$$

In particular, the restriction $\mathcal{D}|_{\mathcal{U}}$ is a regular Frobenius distribution on the non-chiral locus \mathcal{U} . As in [8], we endow $\mathcal{D}|_{\mathcal{U}}$ with the orientation induced by that of M using the unit norm vector field $n \stackrel{\text{def.}}{=} \hat{V}^\sharp = \frac{1}{\|V\|} V^\sharp$, which corresponds to the $\mathcal{D}|_{\mathcal{U}}$ -longitudinal volume form:

$$\nu_{\mathcal{T}} \stackrel{\text{def.}}{=} \iota_{\hat{V}} \nu|_{\mathcal{U}} = n \lrcorner \nu|_{\mathcal{U}} \in \Omega_{\mathcal{U}}^7(\mathcal{D}).$$

Let $*_{\perp} : \Omega_{\mathcal{U}}(\mathcal{D}) \rightarrow \Omega_{\mathcal{U}}(\mathcal{D})$ denote the corresponding Hodge operator along the Frobenius distribution $\mathcal{D}|_{\mathcal{U}}$:

$$*_{\perp} \omega = *(\hat{V} \wedge \omega) = -\iota_{\hat{V}}(*\omega) = \tau(\omega)\nu_{\mathcal{T}}, \quad \forall \omega \in \Omega_{\mathcal{U}}(\mathcal{D}). \tag{3.13}$$

3.5 Spinor parameterization and G_2 structure on the non-chiral locus

Proposition [8]. Relations (3.2) are equivalent on \mathcal{U} with the following conditions:

$$V^2|_{\mathcal{U}} = 1 - b^2, \quad Y|_{\mathcal{U}} = (1 + b\nu)|_{\mathcal{U}}\psi, \quad Z|_{\mathcal{U}} = V|_{\mathcal{U}}\psi, \quad (3.14)$$

where $\psi \in \Omega_{\mathcal{U}}^4(\mathcal{D})$ is the canonically normalized coassociative form of a G_2 structure on the Frobenius distribution $\mathcal{D}|_{\mathcal{U}}$ which is compatible with the metric $g|_{\mathcal{D}}$ induced by g and with the orientation of $\mathcal{D}|_{\mathcal{U}}$.

Let $\varphi \stackrel{\text{def.}}{=} *_1\psi \in \Omega_{\mathcal{U}}^3(\mathcal{D})$ be the associative form of the G_2 structure on $\mathcal{D}|_{\mathcal{U}}$ mentioned in the proposition. We have [8]:

$$\psi = \frac{1}{1 - b^2}VZ = \frac{1}{1 - b^2}(1 - b\nu)Y \in \Omega_{\mathcal{U}}^4(\mathcal{D}), \quad (3.15)$$

$$\varphi = \frac{1}{\|V\|} * Z = \frac{1}{\sqrt{1 - b^2}}Z\nu \in \Omega_{\mathcal{U}}^3(\mathcal{D}). \quad (3.16)$$

On the non-chiral locus, one can parameterize \check{E} as [8]:

$$\check{E}|_{\mathcal{U}} = \frac{1}{16}(1 + V + b\nu)(1 + \psi) = P|_{\mathcal{U}}\Pi, \quad (3.17)$$

where:

$$P \stackrel{\text{def.}}{=} \frac{1}{2}(1 + V + b\nu) \in \Omega(\mathcal{M}), \quad \Pi \stackrel{\text{def.}}{=} \frac{1}{8}(1 + \psi) \in \Omega_{\mathcal{U}}(\mathcal{D})$$

and where $P|_{\mathcal{U}}$ and Π are commuting idempotents in the Kähler-Atiyah algebra of \mathcal{U} . Notice the relations:

$$\varphi = *_1\psi = *(\hat{V} \wedge \psi), \quad *\varphi = -\hat{V} \wedge \psi, \quad *\psi = \hat{V} \wedge \varphi \quad (3.18)$$

and:

$$V\varphi = -\varphi V = V \wedge \varphi, \quad V\psi = \psi V = V \wedge \psi. \quad (3.19)$$

The selfdual and anti-selfdual parts of ψ . We have:

$$\psi^{\pm} = \frac{1}{2}(\psi \pm *\psi) = \frac{1}{2}(\psi \pm \hat{V} \wedge \varphi) \in \Omega(\mathcal{U}). \quad (3.20)$$

Lemma. The four-forms $\psi^{\pm} \in \Omega(\mathcal{U})$ satisfy the relations:

$$\hat{V}\psi^{\pm}\hat{V} = \psi^{\mp}, \quad (3.21)$$

$$\psi^+\psi^- = \psi^-\psi^+ = 0, \quad (3.22)$$

$$\psi^{\pm} = \frac{Y^{\pm}}{1 \pm b}|_{\mathcal{U}}, \quad (3.23)$$

$$\|\psi^+\|^2 = \|\psi^-\|^2 = \frac{7}{2}. \quad (3.24)$$

Proof. Using $\psi^\pm = R^\pm \psi$, relation (3.22) follows immediately from the fact that ν commutes with ψ . The last relation in (3.19) gives:

$$\hat{V}\psi\hat{V} = \psi \quad \text{on } \mathcal{U}. \quad (3.25)$$

Using the fact that \hat{V} and ν anti-commute in the Kähler-Atiyah algebra while ψ and ν commute (because ν is twisted central), relation (3.25) implies (3.21). Separating Y into its selfdual and anti-selfdual parts and using the fact that $\nu Y = Y\nu = *Y$, the last equality in (3.15) implies (3.23), which implies (3.24) when combined with the first relation in (3.8). ■

Proposition. The inhomogeneous differential forms:

$$\Pi^\pm \stackrel{\text{def.}}{=} R^\pm|_{\mathcal{U}}\Pi = \Pi R^\pm|_{\mathcal{U}} = \frac{1}{8} (R^\pm|_{\mathcal{U}} + \psi^\pm) = \frac{1}{16} (1 \pm \nu|_{\mathcal{U}} + 2\psi^\pm) \in \Omega(\mathcal{U})$$

satisfy $\Pi = \Pi^+ + \Pi^-$ and $\hat{V}\Pi^\pm\hat{V} = \Pi^\mp$ and are orthogonal idempotents in the Kähler-Atiyah algebra of \mathcal{U} :

$$(\Pi^\pm)^2 = \Pi^\pm, \quad \Pi^\pm\Pi^\mp = 0.$$

Furthermore, we have:

$$\check{E}^\pm|_{\mathcal{U}} = P|_{\mathcal{U}}\Pi^\pm. \quad (3.26)$$

Notice that Π^\pm are twisted (anti-)selfdual:

$$\Pi^\pm\nu = \pm\Pi^\pm.$$

Proof. Notice that ψ and R^\pm commute since ψ and ν commute. The conclusion now follows immediately using the properties of Π and R^\pm . ■

3.6 Spinor parameterization and $\text{Spin}(7)_\pm$ structures on the loci \mathcal{U}^\pm

Extending ψ^\pm to \mathcal{U}^\pm . Notice that $P \in \Omega(M)$ is globally defined on M while $\Pi \in \Omega(\mathcal{U})$ is only defined on the non-chiral locus.

Proposition. The four-form ψ^\pm has a continuous extension to the locus \mathcal{U}^\pm , which we denote through $\bar{\psi}^\pm \in \Omega^4(\mathcal{U}^\pm)$. Namely:

$$\bar{\psi}^\pm \stackrel{\text{def.}}{=} \frac{1}{1 \pm b} (Y^\pm|_{\mathcal{U}^\pm}) \in \Omega^4(\mathcal{U}^\pm).$$

Furthermore, the idempotents $\Pi^\pm \in \Omega(\mathcal{U})$ have continuous extensions to idempotents $\bar{\Pi}^\pm \in \Omega(\mathcal{U}^\pm)$, which are given by:

$$\bar{\Pi}^\pm \stackrel{\text{def.}}{=} \frac{1}{8} (R^\pm|_{\mathcal{U}^\pm} + \bar{\psi}^\pm) = \frac{1}{16} (1 + 2\psi^\pm \pm \nu) \in \Omega(\mathcal{U}^\pm) \quad (3.27)$$

and which are twisted (anti-)selfdual:

$$\bar{\Pi}^\pm R^\pm|_{\mathcal{U}^\pm} = \bar{\Pi}^\pm, \quad \bar{\Pi}^\pm R^\mp|_{\mathcal{U}^\pm} = 0.$$

Remarks.

1. Notice that (3.23) does not provide any information about the limit of ψ^\mp along \mathcal{W}^\pm , so ψ^\mp (and hence also Π^\mp) will not generally have an extension to \mathcal{U}^\pm . However, (3.24) tells us that ψ^\mp is bounded on M . In particular, we have:

$$\lim_{b \rightarrow \pm 1} (V\psi^\mp) = \lim_{b \rightarrow \pm 1} (\psi^\mp V) = 0. \tag{3.28}$$

2. On the locus \mathcal{W}^\pm we have:

$$b|_{\mathcal{W}^\pm} = \pm 1, \quad V|_{\mathcal{W}^\pm} = Z|_{\mathcal{W}^\pm} = Y^\mp|_{\mathcal{W}^\pm} = 0, \tag{3.29}$$

where the last relations follow from the last equation in (3.6) and from (3.23). The remaining conditions in (3.6) are automatically satisfied.

3. Notice the relation:

$$Y^\pm|_{\mathcal{W}^\pm} = 2\bar{\psi}^\pm|_{\mathcal{W}^\pm},$$

which follows from the fact that $b|_{\mathcal{W}^\pm} = \pm 1$.

Proof. Since $Y^\pm \in \Omega(M)$ is well-defined on M , the conclusion follows immediately from relation (3.23) and from the fact that $1 \pm b$ does not vanish on \mathcal{U}^\pm . The relations satisfied by $\bar{\Pi}^\pm$ on \mathcal{U}^\pm follow by continuity from the similar relations satisfied by Π^\pm on \mathcal{U} . ■

While Π^\mp does not generally have an extension to \mathcal{W}^\pm , the product $P\Pi^\mp$ has zero limit on \mathcal{W}^\pm :

Proposition. We have $P|_{\mathcal{W}^\pm} = R^\pm$ as well as:

$$\exists \lim_{b \rightarrow \pm 1} P\Pi^\mp = \check{E}^\mp|_{\mathcal{W}^\pm} = 0, \quad \check{E}^\pm|_{\mathcal{W}^\pm} = \bar{\Pi}^\pm|_{\mathcal{W}^\pm} = \frac{1}{8} (R^\pm + \bar{\psi}^\pm)|_{\mathcal{W}^\pm} = \frac{1}{16} (1 \pm \nu + 2\bar{\psi}^\pm)|_{\mathcal{W}^\pm}. \tag{3.30}$$

Proof. The relation $P|_{\mathcal{W}^\pm} = R^\pm$ is obvious. The other statements follow from (3.11) and (3.26) using (3.29). ■

The Spin(7) $_\pm$ structures on \mathcal{U}^\pm .

Lemma. Let $(e^a)_{a=1\dots 8}$ be a local coframe defined over an open subset $U \subset M$ and let $\eta \in \Gamma(U, S)$. Then:

$$\mathcal{B}(\gamma^a \eta, \gamma^b \eta) = g^{ab} \|\eta\|^2,$$

where $\gamma^a = \gamma(e^a)$ and $g^{ab} = \langle e^a, e^b \rangle$.

Proof. Using the property $(\gamma^a)^t = \gamma^a$ and the fact that $(\gamma^a \gamma^b)^t = \gamma^b \gamma^a$, compute:

$$\mathcal{B}(\gamma^a \eta, \gamma^b \eta) = \mathcal{B}(\eta, \gamma^a \gamma^b \eta) = \mathcal{B}(\eta, \gamma^b \gamma^a \eta) = \frac{1}{2} \mathcal{B}(\eta, \{\gamma^a, \gamma^b\} \eta) = g^{ab} \mathcal{B}(\eta, \eta) = g^{ab} \|\eta\|^2. \quad \blacksquare$$

When η is non-vanishing everywhere on U , the proposition implies that the spinors $\gamma^a \eta$ form a linearly-independent set of sections of S above U . Taking η to have chirality ± 1 and recalling that γ^a map S^\pm into S^\mp and that $\text{rk} S^+ = \text{rk} S^- = 8$, this gives:

Corollary. Let $(e^a)_{a=1\dots 8}$ be a local orthonormal coframe defined over an open subset $U \subset M$ and $\eta \in \Gamma(U, S^\pm)$ be a spinor of chirality ± 1 which is nowhere vanishing on U . Then $(\gamma^a \eta)_{a=1\dots 8}$ is a \mathcal{B} -orthogonal local frame of S^\mp above U . Every local section $\xi \in \Gamma(U, S^\mp)$ expands in this frame as:

$$\xi = \frac{1}{\|\eta\|^2} \sum_{a=1}^8 \mathcal{B}(\xi, \gamma_a \eta) \gamma^a \eta.$$

Proposition. Let U be an open subset of M which supports an orthonormal coframe e^a of (M, g) . Then:

1. If ξ^+ is everywhere non-vanishing on U , then ξ^- expands above U as $\xi^- = \sum_{a=1}^8 L_a^+ \gamma^a \xi^+ = \gamma(L^+) \xi^+$, where L_a^+ are the coefficients of the one-form $L^+ = L_a^+ dx^a = \frac{1}{1+b} V$.
2. If ξ^- is everywhere non-vanishing on U , then ξ^+ expands above U as $\xi^+ = \sum_{a=1}^8 L_a^- \gamma^a \xi^- = \gamma(L^-) \xi^-$, where L_a^- are the coefficients of the one-form $L^- = L_a^- dx^a = \frac{1}{1-b} V$.

Proof. Assume that ξ^+ (respectively ξ^-) vanishes nowhere on U . The corollary shows that ξ^\mp expands as $\xi^\mp = \sum_{a=1}^8 L_a^\pm \gamma^a \xi^\pm$ where:

$$L_a^\pm = \frac{1}{\|\xi^\pm\|^2} \mathcal{B}(\xi^\mp, \gamma_a \xi^\pm). \quad (3.31)$$

Recalling that S^+ and S^- are \mathcal{B} -orthonormal while γ^a are \mathcal{B} -symmetric, we find:

$$\mathcal{B}(\xi^+, \gamma_a \xi^-) = \mathcal{B}(\xi^-, \gamma_a \xi^+) = \frac{1}{2} \mathcal{B}(\xi, \gamma_a \xi) = \frac{1}{2} V_a.$$

Using this and (3.12), equation (3.31) becomes $L_a^\pm = \frac{1}{1\pm b} V_a$. ■

Remarks.

1. The “+” case of (3.31) was used in [1], where no explicit expression for L^+ (which is denoted by L in loc. cit.) was given.³
2. Notice that L^+ and L^- are not independent (they are proportional to each other) and that each of them contains the same information as V and b .

Recalling (3.12), consider the unit norm spinors (of chirality ± 1):

$$\eta^\pm = \sqrt{1 + \|L^\pm\|^2} \xi^\pm = \sqrt{\frac{2}{1\pm b}} \xi^\pm \in \Gamma(\mathcal{U}^\pm, S^\pm). \quad (3.32)$$

Using the fact that $\|\eta^\pm\| = 1$ while $\mathcal{B}(\eta^\pm, \gamma_{a_1\dots a_k} \eta^\pm)$ vanishes unless $k \equiv_4 0$, we find:

$$\check{E}_{\eta^\pm, \eta^\pm} = \frac{1}{16} (1 + \Phi^\pm \pm \nu) \in \Omega(\mathcal{U}^\pm), \quad (3.33)$$

where:

$$\Phi^\pm \stackrel{\text{def.}}{=} \frac{1}{4!} \mathcal{B}(\eta^\pm, \gamma_{a_1\dots a_4} \eta^\pm) e^{a_1\dots a_4} = \check{E}_{\eta^\pm, \eta^\pm}^{(4)} = \frac{2}{1\pm b} \check{E}_{\xi^\pm, \xi^\pm}^{(4)} \in \Omega^4(\mathcal{U}^\pm) \quad (3.34)$$

and where we noticed that $\mathcal{B}(\eta^\pm, \gamma(\nu) \eta^\pm) = \pm 1$.

³Notice that L^+ is not a quadratic function of ξ , since it involves the denominator $1 + b$ and thus it is not homogeneous under rescalings $\xi \rightarrow \lambda \xi$ with $\lambda \neq 0$.

Proposition. The four-form Φ^+ is selfdual while the four-form Φ^- is anti-selfdual. They satisfy the following relations on the locus \mathcal{U}^\pm :

$$\Phi^\pm = 2\bar{\psi}^\pm. \quad (3.35)$$

In particular, the inhomogeneous form (3.33) coincides with the extension (3.27) of Π^\pm to this locus:

$$\check{E}_{\eta^\pm, \eta^\pm} = \bar{\Pi}^\pm$$

and we have:

$$\|\Phi^\pm\|^2 = 14. \quad (3.36)$$

Moreover, the restriction of Φ^+ is the canonically-normalized calibration defining a Spin(7) structure on the open submanifold \mathcal{U} of M while the restriction of Φ^- is the canonically-normalized calibration defining a Spin(7) structure on the orientation reversal of \mathcal{U} .

Proof. Recalling that $\xi^\pm = \frac{1}{2}(1 \pm \gamma(\nu))\xi$, the identities $\check{E}_{\xi, \gamma(\nu)\xi} = \check{E}_{\xi, \xi}\nu$ and $\check{E}_{\gamma(\nu)\xi, \xi} = \nu\check{E}_{\xi, \xi}$ of [32] and the fact that ν is involutive and twisted central give:

$$\begin{aligned} \check{E}_{\xi^\pm, \xi^\pm} &= \frac{1}{4} (\check{E}_{\xi, \xi} \pm \nu\check{E}_{\xi, \xi} \pm \check{E}_{\xi, \xi}\nu + \nu\check{E}_{\xi, \xi}\nu) = \frac{1}{4} (\check{E}_{\xi, \xi} + \pi(\check{E}_{\xi, \xi})) (1 \pm \nu) \\ &= \frac{1}{2} \check{E}_{\xi, \xi}^{\text{ev}} (1 \pm \nu) = \frac{1}{2} (\check{E}_{\xi, \xi}^{\text{ev}} \pm * \tau(\check{E}_{\xi, \xi}^{\text{ev}})). \end{aligned}$$

Since the Hodge operator preserves $\Omega^4(M)$ and since the reversion τ of the Kähler-Atiyah algebra restricts to the identity on the space of four-forms, this implies:

$$\check{\mathbf{E}}_{\xi^\pm, \xi^\pm}^{(4)} = \frac{1}{2} (\check{\mathbf{E}}_{\xi, \xi}^{(4)} \pm * \check{\mathbf{E}}_{\xi, \xi}^{(4)}) = \frac{1}{2} (Y \pm *Y) = Y^\pm,$$

where the superscript \pm indicates the selfdual/anti-selfdual part. Substituting this into (3.34) gives relation (3.35). The statements of the proposition regarding the restrictions of Φ^\pm to the open submanifold \mathcal{U} follow from the fact that η_\pm is a Majorana-Weyl spinor of norm one and of chirality ± 1 ; it is well-known [40] that giving such a spinor on an eight-manifold \mathcal{U} induces Spin(7) structures on the underlying manifold or on its orientation reversal, whose normalized calibrations are given by (3.34). In particular, (3.36) holds on \mathcal{U} since there it amounts to the condition that Φ^\pm are canonically normalized. By continuity, this implies that (3.36) also holds on \mathcal{W}^\pm . ■

Remarks.

1. The proposition implies that the following relation holds on the non-chiral locus:

$$\check{E}_{\xi, \xi}|_{\mathcal{U}} = P|_{\mathcal{U}} (\check{E}_{\eta^+, \eta^+} + \check{E}_{\eta^-, \eta^-}).$$

This shows how the idempotent $\check{E}_{\xi, \xi}|_{\mathcal{U}}$ which characterizes the normalized Majorana spinor ξ on the locus \mathcal{U} relates to the two idempotents $\check{E}_{\eta^\pm, \eta^\pm}|_{\mathcal{U}} = \Pi^\pm$ which characterize the Majorana-Weyl spinors η^\pm and which encode the Spin(7) $_\pm$ structures through the Kähler-Atiyah algebra. While $\check{E}_{\eta^+, \eta^+}$ depends only on the positive chirality spinor η^+ and $\check{E}_{\eta^-, \eta^-}$ depends only on the negative chirality spinor η^- , the

idempotent P contains the quantities b and V , each of which involves both chirality components of the spinor ξ :

$$b = \|\xi^+\|^2 - \|\xi^-\|^2, \quad V = 2\mathcal{B}(\xi^+, \gamma_m \xi^-) e^m = (1 - b^2) \mathcal{B}(\eta^+, \gamma_m \eta^-) e^m.$$

The object P encodes in the Kähler-Atiyah algebra the $SO(7)$ structure which corresponds to the distribution \mathcal{D} on \mathcal{U} . Finally, notice that the idempotent Π encodes the G_2 structure along the distribution \mathcal{D} . Notice that P and Π commute, while P and Π_\pm do not commute.

2. Equation (3.35) implies that Φ^\pm coincides with $\pm Y^\pm$ on the locus \mathcal{W}^\pm since $b = \pm 1$ there. Notice that (3.36) agrees via (3.35) with the last equations in (3.6).

Spinor parameterization on the loci \mathcal{U}^\pm . On the locus \mathcal{U} , relations (3.14) and (3.35) give:

$$\begin{aligned} Z|_{\mathcal{U}} &= \frac{1}{2}V(\Phi^+ + \Phi^-), \\ Y|_{\mathcal{U}} &= \frac{1}{2}[(1+b)\Phi^+ + (1-b)\Phi^-]. \end{aligned} \tag{3.37}$$

In these relations, Φ^+ and Φ^- are not independent but related through:

$$\Phi^\mp = \hat{V}\Phi^\pm\hat{V}$$

as a consequence of (3.21). Hence on the non-chiral locus we can eliminate Φ^\mp in terms of Φ^\pm to obtain the following non-redundant parameterizations:

$$Z|_{\mathcal{U}} = \frac{1}{2}\sqrt{1-b^2}(\hat{V}\Phi^\pm + \Phi^\pm\hat{V}), \quad Y|_{\mathcal{U}} = \frac{1}{2}[(1 \pm b)\Phi^\pm + (1 \mp b)\hat{V}\Phi^\pm\hat{V}],$$

which give:

$$\begin{aligned} 16\check{E}|_{\mathcal{U}} &= P|_{\mathcal{U}}(\Pi_\pm + \hat{V}\Pi_\pm\hat{V}) \\ &= 1 + V + \frac{1}{2}[(1 \pm b)\Phi^\pm + (1 \mp b)\hat{V}\Phi^\pm\hat{V}] + \frac{1}{2}\sqrt{1-b^2}(\hat{V}\Phi^\pm + \Phi^\pm\hat{V}) + b\nu. \end{aligned}$$

This imply the following parameterizations on the loci \mathcal{U}^\pm :

$$16\check{E}|_{\mathcal{U}^\pm} = 1 + V + \frac{1}{2}\left[(1 \pm b)\Phi^\pm + \frac{1}{1 \pm b}V\Phi^\pm V\right] + \frac{1}{2}(V\Phi^\pm + \Phi^\pm V) + b\nu,$$

where it is understood that (see (3.28)):

$$\lim_{b \rightarrow \pm 1} V\Phi^\mp = \lim_{b \rightarrow \pm 1} \Phi^\mp V = 0$$

and hence (see (3.30)):

$$16\check{E}|_{\mathcal{W}^\pm} = \bar{\Pi}^\pm|_{\mathcal{W}^\pm} = \frac{1}{16}(1 + \Phi^\pm \pm \nu)|_{\mathcal{W}^\pm}.$$

Up to expressing V and b through L^\pm , this is the parameterization which corresponds to the approach of [1].

G structure	Spin(7) ₊	Spin(7) ₋	G ₂ (on $\mathcal{D} _{\mathcal{U}}$)	SO(7) ($\mathcal{D} _{\mathcal{U}}$)
spinor	η^+	η^-	$\eta_0 = \frac{1}{\sqrt{2}}(\eta^+ + \eta^-)$	—
idempotent	$\Pi^+ = \frac{1}{16}(1 + \Phi^+ + \nu)$	$\Pi^- = \frac{1}{16}(1 + \Phi^- - \nu)$	$\Pi = \Pi^+ + \Pi^- = \frac{1}{8}(1 + \psi)$	$P = \frac{1}{2}(1 + V + b\nu)$
forms	$\Phi^+ = 2\psi^+$	$\Phi^- = 2\psi^-$	φ and $\psi = *_{\perp}\varphi$	b and V
extends to	\mathcal{U}^+	\mathcal{U}^-	\mathcal{U}	\mathcal{U}

Table 1. Summary of various G structures and of their reflections in the Kähler-Atiyah algebra.

3.7 Comparing spinors and G structures on the non-chiral locus

Equation (3.20) gives:

$$\Phi^{\pm}|_{\mathcal{U}} = 2\psi^{\pm} = \psi \pm \hat{V} \wedge \varphi,$$

i.e.:

$$(\Phi^{\pm}|_{\mathcal{U}})_{\top} = \pm\varphi, \quad (\Phi^{\pm}|_{\mathcal{U}})_{\perp} = \psi. \tag{3.38}$$

The relation $\xi^{\mp} = \gamma(L^{\pm})\xi^{\pm}$ gives $\eta^{\mp} = \gamma(\hat{V})\eta^{\pm}$, which shows that the everywhere normalized spinor:

$$\eta_0 \stackrel{\text{def.}}{=} \frac{1}{\sqrt{2}}(\eta^+ + \eta^-) \in \Gamma(\mathcal{U}, S) \tag{3.39}$$

is a Majorana spinor along \mathcal{D} in the seven-dimensional sense, i.e. we have $D(\eta_0) = \eta_0$ where $D \stackrel{\text{def.}}{=} \gamma(\hat{V})$ is the real structure of S , when the latter is viewed as a complex spinor bundle over \mathcal{D} (see [8]). The identity $\check{E}_{\eta^{\pm}, \eta^{\mp}}^{(4)} = 0$ implies the following spinorial expression for ψ :

$$\psi = \check{E}_{\eta_0, \eta_0}^{(4)} = \frac{1}{4!} \mathcal{B}(\eta_0, \gamma_{a_1 \dots a_4} \eta_0) e^{a_1 \dots a_4}. \tag{3.40}$$

The relation $\xi^{\mp} = \gamma(L^{\pm})\xi^{\pm}$ gives $\eta^{\mp} = \gamma(\hat{V})\eta^{\pm}$, which implies:

$$\eta_0 = \frac{1}{\sqrt{2}}(\text{id}_S + \gamma(\hat{V}))\eta^+ = \frac{1}{\sqrt{2}}(\text{id}_S + \gamma(\hat{V}))\eta^-.$$

Notice that $\frac{1}{2}(\text{id}_S + \gamma(\hat{V}))$ is an idempotent endomorphism of S . As explained in [8], the spinor η_0 induces the G_2 structure of the distribution \mathcal{D} . The situation is summarized in table 1.

Remarks.

1. None of the G structures in table 1 extends to M . In fact, the structure group $\text{SO}(8)$ of the frame bundle of M does not globally reduce, in general, to any proper subgroup. As pointed out in [1], this is due to the fact that the action of $\text{Spin}(8)$ on the fibers $S_p \simeq \mathbb{R}^{16}$ of S (which is the action of $\text{Spin}(8)$ on the direct sum $\mathfrak{8}_s \oplus \mathfrak{8}_c$ of the positive and negative chirality spin 1/2 representations) is not transitive when restricted to the unit sphere $S^{15} \subset \mathbb{R}^{16}$. As shown in loc. cit., one can in some sense “cure” this problem by considering the manifold $\hat{M} \stackrel{\text{def.}}{=} M \times S^1$, using the fact that $\text{Spin}(9)$ acts transitively on S^{15} . However, such an approach does not immediately provide useful information on the geometry of M , in particular the geometry of the

singular foliation $\bar{\mathcal{F}}$ discussed in the next subsection is not immediately visible in that approach. It was also shown in loc. cit. that one can repackage the information contained in the $\text{Spin}(7)_{\pm}$ structures into a generalized $\text{Spin}(7)$ structure on \hat{M} in the sense of [41, 42]. In particular, it is easy to check that relations (4.8) of [1] are equivalent with some of the exterior differential constraints which can be obtained by expanding equation (3.5) of [8] into its rank components — exterior differential constraints which were discussed at length in [32] and in the appendix of [8]. As shown in detail in [8], those exterior differential constraints do not suffice to encode the full supersymmetry conditions for such backgrounds.

2. The fact that the structure group of TM does not globally reduce beyond $\text{SO}(8)$ in this class of examples illustrates some limits of the philosophy that flux compactifications can be described using reductions of structure group. That philosophy is based on the observation that a collection of (s)pinors defines a *local* reduction of structure group over any open subset of the compactification manifold M along which the stabilizer of the pointwise values of those spinors is fixed up to conjugacy in the corresponding Spin or Pin group. However, such a reduction does *not* generally hold globally on M , since the local reductions thus obtained can “jump” — in our class of examples, the jump occurs at the points of the chiral locus \mathcal{W} . The appropriate notion is instead that of *generalized* reduction of structure group, of which the class of compactifications considered here is an example. In this respect, we mention that the cosmooth generalized distribution \mathcal{D} can be viewed as providing a generalized reduction of structure group of M , which is an ordinary reduction from $\text{SO}(8)$ to $\text{SO}(7)$ only when restricted to its regular subset \mathcal{U} , on which $\mathcal{D}|_{\mathcal{U}}$ provides [8] an almost product structure. We also mention that the conditions imposed by supersymmetry can be formulated globally by using an extension of the language of Haefliger structures (see section 3.8), an approach which can in fact be used to give a fully general approach to flux compactifications. It is such concepts, rather than the classical concept of G structures [43], which provide the language appropriate for giving *globally valid* descriptions of the most general flux compactifications.

3.8 The singular foliation of M defined by \mathcal{D}

As in [8], one can show that the one-form:

$$\omega \stackrel{\text{def.}}{=} 4\kappa e^{3\Delta} V$$

satisfies the following relations which hold globally on M as a consequence of the supersymmetry conditions (2.4):

$$\begin{aligned} d\omega &= 0, \\ \omega &= \mathbf{f} - d\mathbf{b}, \quad \text{where } \mathbf{b} \stackrel{\text{def.}}{=} e^{3\Delta} b. \end{aligned} \tag{3.41}$$

As a result of the first equation, the generalized distribution $\mathcal{D} = \ker V = \ker \omega$ determines a singular foliation $\bar{\mathcal{F}}$ of M , which degenerates along the chiral locus \mathcal{W} , since that locus coincides with the set of zeroes of ω . The second equation implies that ω belongs to the cohomology class $\mathfrak{f} \in H^1(M, \mathbb{R})$ of \mathbf{f} .

Since \mathcal{D} is cosmooth rather than smooth, the notion of singular foliation which is appropriate in our case⁴ is that of Haefliger structure [15]. More precisely, $\bar{\mathcal{F}}$ can be described as the Haefliger structure defined as follows. Consider an open cover $(U_\alpha)_{\alpha \in I}$ of M such that each U_α is simply-connected and let $\omega_\alpha \stackrel{\text{def.}}{=} \omega|_{U_\alpha} \in \Omega^1(U_\alpha)$. We have $\omega_\alpha = d\mathbf{h}_\alpha$ for some $\mathbf{h}_\alpha \in \Omega^0(U_\alpha)$, where \mathbf{h}_α are determined up to shifts:

$$\mathbf{h}_\alpha \rightarrow \mathbf{h}'_\alpha + \mathbf{c}_\alpha, \quad \mathbf{c}_\alpha \in \mathbb{R}. \tag{3.42}$$

For any $\alpha, \beta \in I$ and any $p \in U_\alpha \cap U_\beta$, consider the orientation-preserving diffeomorphism $\phi_{\alpha\beta}(p) \in \text{Diff}_+(\mathbb{R})$ of the real line given by the translation:

$$\phi_{\alpha\beta}(p)(x) \stackrel{\text{def.}}{=} x + \mathbf{h}_\beta(p) - \mathbf{h}_\alpha(p) \quad \forall x \in \mathbb{R}.$$

Then $\phi_{\alpha\beta}(p)(\mathbf{h}_\alpha(p)) = \mathbf{h}_\beta(p)$. The germ $\hat{\phi}_{\alpha\beta}(p)$ of $\phi_{\alpha\beta}(p)$ at $\mathbf{h}_\alpha(p)$ is an element of the Haefliger groupoid Γ_1^∞ and it is easy to check that $\hat{\phi}_{\alpha\beta} : U_\alpha \cap U_\beta \rightarrow \Gamma_1^\infty$ is a Haefliger cocycle on M :

$$\hat{\phi}_{\beta\gamma}(p) \circ \hat{\phi}_{\alpha\beta}(p) = \hat{\phi}_{\alpha\gamma}(p) \quad \forall \alpha, \beta, \gamma \in I, \quad \forall p \in U_\alpha \cap U_\beta \cap U_\gamma.$$

Moreover, the shifts (3.42) correspond to transformations:

$$\hat{\phi}_{\alpha\beta} \rightarrow \hat{\phi}'_{\alpha\beta} = \hat{\mathbf{q}}_\beta \circ \hat{\phi}_{\alpha\beta} \circ \hat{\mathbf{q}}_\alpha^{-1},$$

where $\hat{\mathbf{q}}_\alpha : U_\alpha \rightarrow \Gamma_1^\infty$ are defined by declaring that $\hat{\mathbf{q}}_\alpha(p)$ is the germ at $p \in U_\alpha$ of the orientation-preserving diffeomorphism $\mathbf{t}_\alpha \in \text{Diff}_+(\mathbb{R})$ given by the following translation of the real line:

$$\mathbf{t}_\alpha(x) = x + \mathbf{c}_\alpha \quad \forall x \in \mathbb{R}.$$

It follows that the closed one-form ω determines a well-defined element of the non-Abelian cohomology $\in H^1(M, \Gamma_1^\infty)$, which is the Haefliger structure defined by ω . The singular foliation $\bar{\mathcal{F}}$ which “integrates” \mathcal{D} can be identified with this element.

The approach through Haefliger structures allows one to define rigorously the singular foliation $\bar{\mathcal{F}}$ in the most general case, i.e. without making any supplementary assumptions on the closed one-form ω . In general, such singular foliations can be extremely complicated and little is known about their topology and geometry. However, the description of $\bar{\mathcal{F}}$ simplifies when ω is a closed one-form of Morse or Bott-Morse type. In section 5, we discuss the Morse case, recalling some results which apply to $\bar{\mathcal{F}}$ in that situation.

4 Relating the G_2 and $\text{Spin}(7)$ approaches on the non-chiral locus

On the non-chiral locus \mathcal{U} , we have the regular foliation \mathcal{F} which is endowed with a longitudinal G_2 structure having associative and coassociative forms φ and ψ . We also have a $\text{Spin}(7)_+$ and a $\text{Spin}(7)_-$ structure, which are determined respectively by the calibrations

⁴Notice that this is not the notion of singular foliation considered in [44, 45], which is instead based on Stefan-Sussmann (i.e. smooth, rather than cosmooth) distributions.

$\Phi^\pm = 2\psi^\pm = \psi \pm \hat{V} \wedge \varphi$. Given this data, one can relate various quantities determined by (\mathcal{D}, φ) to quantities determined by Φ^\pm as we explain below. We stress that the results of this subsection are independent of the supersymmetry conditions (2.4) and hence they hold in the general situation described above. We mention that the relation between the type of G_2 structure induced on an oriented submanifold of a Spin(7) structure manifold and the intrinsic geometry of such submanifolds was studied in [46, 47].

4.1 The G_2 and Spin(7) $_{\pm}$ decompositions of $\Omega^4(\mathcal{U})$

The group G_2 has a natural fiberwise rank-preserving action on the graded vector bundle $\wedge(\mathcal{D}|_{\mathcal{U}})^*$, which is given at every $p \in \mathcal{U}$ by the local embedding of G_2 as the stabilizer $G_{2,p}$ in $\text{SO}(\mathcal{D}_p)$ of the 3-form $\varphi_p \in \wedge^3(\mathcal{D}_p^*)$. Since $\text{SO}(\mathcal{D}_p)$ embeds into $\text{SO}(T_p M)$ as the stabilizer of the 1-form $V_p \in T_p^* M$, this induces a rank-preserving action of $G_{2,p}$ on $\wedge T_p^* \mathcal{U}$ which can be described as follows. Decomposing any form $\omega \in \wedge T_p^* \mathcal{U}$ as $\omega = \omega_{\perp} + \hat{V} \wedge \omega_{\top}$, the action of an element of g of G_2 on ω is given by the simultaneous action of g on the components ω_{\perp} and ω_{\top} , both of which belong to $\wedge \mathcal{D}_p^*$. The corresponding representation of G_2 at p is equivalent with the direct sum of the representations in which the components ω_{\top} and ω_{\perp} transform at p . In particular, $F_{\perp,p}$ and $F_{\top,p}$ transform in a G_2 representation which is equivalent with the direct sum $\wedge^3 \mathcal{D}_p^* \oplus \wedge^4 \mathcal{D}_p^*$. The group Spin(7) is embedded inside $\text{SO}(T_p M)$ in two ways, namely as the stabilizers Spin(7) $_{\pm,p}$ of the selfdual 4-forms Φ_p^{\pm} . Then (3.35) shows that $G_{2,p}$ is the stabilizer of V_p in Spin(7) $_{\pm,p}$. The action of $G_{2,p}$ on $\wedge T_p^* M$ is obtained from that of Spin(7) $_{\pm,p}$ by restriction. Hence the irreducible components of the action of Spin(7) $_{\pm,p}$ on $\wedge^k(T_p^* M)$ decompose as direct sums of the irreducible components of the action of $G_{2,p}$ on the same space. We have the following decompositions into irreps. (see, for example, [48, 49]):

$$\begin{aligned}
 \wedge^4 T_p^* M &= \wedge_{1,\pm}^4 T_p^* M \oplus \wedge_{7,\pm}^4 T_p^* M \oplus \wedge_{27,\pm}^4 T_p^* M \oplus \wedge_{35,\pm}^4 T_p^* M & \text{for } \text{Spin}(7)_{\pm,p}, \\
 \wedge^4 T_p^* M &= \wedge_1^4 T_p^* M \oplus \wedge_7^4 T_p^* M \oplus \wedge_{27}^4 T_p^* M & \text{for } G_{2,p}, \\
 \wedge^3 T_p^* M &= \wedge_1^3 T_p^* M \oplus \wedge_7^3 T_p^* M \oplus \wedge_{27}^3 T_p^* M & \text{for } G_{2,p},
 \end{aligned}
 \tag{4.1}$$

where the numbers used as lower indices indicate the dimension of the corresponding irrep. The last two of these decompositions imply similar decompositions into irreps. of $G_{2,p}$ for the spaces of selfdual and anti-selfdual three- and four-forms:

$$(\wedge^4 T_p^* M)^{\pm} = \wedge_1^4 T_p^* M \oplus \wedge_7^4 T_p^* M \oplus \wedge_{27}^4 T_p^* M \quad \text{for } G_{2,p}.
 \tag{4.2}$$

Furthermore, we have:

$$\begin{aligned}
 (\wedge^4 T_p^* M)^{\pm} &= \wedge_{1,\pm}^4 T_p^* M \oplus \wedge_{7,\pm}^4 T_p^* M \oplus \wedge_{27,\pm}^4 T_p^* M & \text{for } \text{Spin}(7)_{\pm,p}, \\
 (\wedge^4 T_p^* M)^{\mp} &= \wedge_{35,\pm}^4 T_p^* M & \text{for } \text{Spin}(7)_{\pm,p},
 \end{aligned}
 \tag{4.3}$$

where the \pm superscripts indicate the subspaces of selfdual and anti-selfdual forms while the \pm subscripts indicate which of the Spin(7) $_p$ subgroups of $\text{SO}(T_p M)$ we consider. Comparing these two decompositions, one sees immediately that the irreps of Spin(7) $_{\pm,p}$ appearing

G_2 representation	1	7	27
$F_{\perp} \in \Omega_{\mathcal{U}}^4(\mathcal{D})$	$\text{tr}_g(\hat{h})$	$\alpha_1 \in \Omega_{\mathcal{U}}^1(\mathcal{D})$	$h^{(0)} \in \text{Sym}_{\mathcal{U},0}^2(\mathcal{D}^*)$
$F_{\top} \in \Omega_{\mathcal{U}}^3(\mathcal{D})$	$\text{tr}_g(\hat{\chi})$	$\alpha_2 \in \Omega_{\mathcal{U}}^1(\mathcal{D})$	$\chi^{(0)} \in \text{Sym}_{\mathcal{U},0}^2(\mathcal{D}^*)$

Table 2. The G_2 parameterization of F on the non-chiral locus.

in (4.3) decompose as follows under the G_2 action on $\wedge^4 T_p^* M$ which was discussed above:

$$\begin{aligned} \wedge_{\mathbf{k},\pm}^4 T_p^* M &= \wedge_k^4 T_p^* M, \quad \text{for } k = 1, 7, 27 \\ \wedge_{\mathbf{35},\pm}^4 T_p^* M &= \wedge_1^4 T_p^* M \oplus \wedge_7^4 T_p^* M \oplus \wedge_{27}^4 T_p^* M. \end{aligned} \quad (4.4)$$

Let $\omega^{(k)} \in \Omega_k(\mathcal{U})$ and $\omega_{\pm}^{[\mathbf{k}]} \in \Omega_{\mathbf{k}}(\mathcal{U})$ denote the (pointwise) projections of a form ω on the irreps of G_2 and $\text{Spin}(7)_{\pm}$ respectively.

4.2 The G_2 and $\text{Spin}(7)_{\pm}$ parameterizations of F

G_2 parameterization. Recall from [8] that $F|_{\mathcal{U}} = F_{\perp} + \hat{V} \wedge F_{\top}$ and $f|_{\mathcal{U}} = f_{\perp} + f_{\top} \hat{V}$, where $f_{\top} \in \Omega^0(\mathcal{U})$, $f_{\perp} \in \Omega_{\mathcal{U}}^1(\mathcal{D})$, $F_{\top} \in \Omega_{\mathcal{U}}^3(\mathcal{D})$ and $F_{\perp} \in \Omega_{\mathcal{U}}^4(\mathcal{D})$, with:

$$\begin{aligned} F_{\perp} &= F_{\perp}^{(7)} + F_{\perp}^{(S)} \quad \text{where} \quad F_{\perp}^{(7)} = \alpha_1 \wedge \varphi \in \Omega_{\mathcal{U}}^4(\mathcal{D}), \quad F_{\perp}^{(S)} = -\hat{h}_{kl} e^k \wedge \iota_{e^l} \psi \in \Omega_{\mathcal{U},S}^4(\mathcal{D}) \\ F_{\top} &= F_{\top}^{(7)} + F_{\top}^{(S)} \quad \text{where} \quad F_{\top}^{(7)} = -\iota_{\alpha_2} \psi \in \Omega_{\mathcal{U},7}^3(\mathcal{D}), \quad F_{\top}^{(S)} = \chi_{kl} e^k \wedge \iota_{e^l} \varphi \in \Omega_{\mathcal{U},S}^3(\mathcal{D}). \end{aligned} \quad (4.5)$$

Here $\alpha_1, \alpha_2 \in \Omega_{\mathcal{U}}^1(\mathcal{D})$, while $\hat{h} = \frac{1}{2} \hat{h}_{ij} e^i \odot e^j$ and $\chi = \frac{1}{2} \chi_{ij} e^i \odot e^j$ are sections of the bundle $\text{Sym}_{\mathcal{U}}^2(\mathcal{D}^*)$. We have $F_{\top}^{(S)} = F_{\top}^{(1)} + F_{\top}^{(27)}$ with $F_{\top}^{(1)} \in \Omega_{\mathcal{U}}^3(\mathcal{D})$, $F_{\top}^{(27)} \in \Omega_{\mathcal{U},27}^3(\mathcal{D})$ and a similar decomposition for $F_{\perp}^{(S)}$. The last relations correspond to the decompositions of χ and \hat{h} into their homothety parts $\text{tr}(\chi)g|_{\mathcal{D}}$, $\text{tr}(\hat{h})g|_{\mathcal{D}}$ and traceless parts:

$$\chi^{(0)} \stackrel{\text{def.}}{=} \chi - \frac{1}{7} \text{tr}(\chi)g|_{\mathcal{D}}, \quad h^{(0)} = \hat{h} - \frac{1}{7} \text{tr}(\hat{h})g|_{\mathcal{D}}.$$

Let $h, \hat{\chi} \in \text{Sym}_{\mathcal{U}}^2(\mathcal{D}^*)$ denote the symmetric tensors defined through:

$$h_{ij} \stackrel{\text{def.}}{=} \hat{h}_{ij} - \frac{1}{3} \text{tr}_g(\hat{h})g_{ij}, \quad \hat{\chi}_{ij} \stackrel{\text{def.}}{=} \chi_{ij} - \frac{1}{4} \text{tr}_g(\chi)g_{ij},$$

where:

$$\text{tr}_g(\chi) = -\frac{4}{3} \text{tr}_g(\hat{\chi}), \quad \text{tr}_g(\hat{h}) = -\frac{3}{4} \text{tr}_g(h).$$

The situation is summarized in table 2.

$\text{Spin}(7)_{\pm}$ parameterization. The discussion of the previous subsection gives the following decompositions of the selfdual and anti-selfdual parts of F :

$$F^{\pm} = F_{\pm}^{[1]} + F_{\pm}^{[7]} + F_{\pm}^{[27]} \in \Omega^{4\pm}(\mathcal{U}), \quad F^{\mp} = F_{\pm}^{[35]} \in \Omega^{4\mp}(\mathcal{U}).$$

Since the Hodge operator intertwines $\text{Spin}(7)_{\pm}$ representations, we have:

$$\begin{aligned} \left(F_{\pm}^{[\mathbf{k}]} \right)_{\perp} &= \pm *_{\perp} \left(F_{\pm}^{[\mathbf{k}]} \right)_{\top} \quad \text{for } k = 1, 7, 27, \\ \left(F_{\pm}^{[35]} \right)_{\perp} &= \mp *_{\perp} \left(F_{\pm}^{[35]} \right)_{\top}. \end{aligned}$$

One can parameterize $F_{\pm}^{[k]}$ through a zero-form $\mathcal{F}_{\pm}^{[1]} \in \Omega^0(\mathcal{U})$, a 2-form $\mathcal{F}_{\pm}^{[7]} \in \Omega^2(\mathcal{U})$, a \mathcal{D} -longitudinal traceless symmetric covariant tensor $\mathcal{F}_{\pm}^{[27]} \in \text{Sym}_{\mathcal{U},0}^2(\mathcal{D}^*)$ and a traceless symmetric covariant tensor $\mathcal{F}_{\pm}^{[35]} \in \text{Sym}_0^2(T^*\mathcal{U})$, which are defined by:

$$\begin{aligned} F_{\pm}^{[1]} &= \frac{1}{42} \mathcal{F}_{\pm}^{[1]} \Phi^{\pm}, \\ F_{\pm}^{[7]} &= \frac{1}{96} \Phi \Delta_1 \mathcal{F}_{\pm}^{[7]}, \\ F_{\pm}^{[27]} &= \frac{1}{24} \left(\mathcal{F}_{\pm}^{[27]} \right)_{ij} e^i \wedge \iota_{e^j} \Phi^{\mp}, \\ F_{\pm}^{[35]} &= \frac{1}{24} \left(\mathcal{F}_{\pm}^{[35]} \right)_{ab} e^a \wedge \iota_{e^b} \Phi^{\pm}. \end{aligned} \quad (4.6)$$

The quantities $\mathcal{F}^{[k]}$ with $k = 1, 7, 35$ can be recovered from F through the relation:

$$6(\iota_{e^a} F) \Delta_3 (\iota_{e^b} \Phi^{\pm}) = g_{ab} \mathcal{F}_{\pm}^{[1]} + \left(\mathcal{F}_{\pm}^{[7]} \right)_{ab} + \left(\mathcal{F}_{\pm}^{[35]} \right)_{ab}. \quad (4.7)$$

Define:

$$\begin{aligned} \beta_{1\pm} &\stackrel{\text{def.}}{=} \left(\mathcal{F}_{\pm}^{[7]} \right)_{\top} \in \Omega_{\mathcal{U}}^1(\mathcal{D}), \\ \beta_{2\pm} &\stackrel{\text{def.}}{=} n \lrcorner \mathcal{F}_{\pm}^{[35]} = \left(\mathcal{F}_{\pm}^{[35]} \right)_{1j} e^j \in \Omega_{\mathcal{U}}^1(\mathcal{D}), \\ \sigma_{\pm} &\stackrel{\text{def.}}{=} \frac{1}{2} \left(\mathcal{F}_{\pm}^{[35]} \right)_{ij} e^i \odot e^j \in \text{Sym}_{\mathcal{U}}^2(\mathcal{D}^*), \end{aligned} \quad (4.8)$$

where e_a is a local orthonormal frame such that $e_1 = n \stackrel{\text{def.}}{=} \hat{V}^{\#}$ and $j = 2, \dots, 8$. The fact that $F_{\pm}^{[7]}$ is (anti-)selfdual implies:

$$\left(\mathcal{F}_{\pm}^{[7]} \right)_{\perp} = \mp \iota_{\beta_{1\pm}} \varphi. \quad (4.9)$$

Choosing an orthonormal frame with $e_1 = n = \hat{V}^{\#}$ and recalling (3.38), relations (4.6) and (4.8) give the following parameterization of F , which refines the parameterization used in [1] by taking into account the decomposition into directions parallel and perpendicular to \hat{V} :

$$\begin{aligned} \left(F_{\pm}^{[1]} \right)_{\top} &= \pm \frac{1}{42} \mathcal{F}_{\pm}^{[1]} \varphi, & \left(F_{\pm}^{[1]} \right)_{\perp} &= \frac{1}{42} \mathcal{F}_{\pm}^{[1]} \psi, \\ \left(F_{\pm}^{[7]} \right)_{\top} &= \frac{1}{24} \iota_{\beta_{1\pm}} \psi, & \left(F_{\pm}^{[7]} \right)_{\perp} &= \mp \frac{1}{24} \beta_{1\pm} \wedge \varphi, \\ \left(F_{\pm}^{[27]} \right)_{\top} &= \mp \frac{1}{24} \left(\mathcal{F}_{\pm}^{[27]} \right)_{ij} e^i \wedge \iota_{e^j} \varphi, & \left(F_{\pm}^{[27]} \right)_{\perp} &= \frac{1}{24} \left(\mathcal{F}_{\pm}^{[27]} \right)_{ij} e^i \wedge \iota_{e^j} \psi, \\ \left(F_{\pm}^{[35]} \right)_{\top} &= \pm \frac{1}{24} \left[\iota_{\beta_{2\pm}} \psi - \frac{4}{7} (\text{tr} \sigma_{\pm}) \varphi + \left(\sigma_{\pm}^{(0)} \right)_{ij} e^i \wedge \iota_{e^j} \varphi \right], \\ \left(F_{\pm}^{[35]} \right)_{\perp} &= \frac{1}{24} \left[\beta_{2\pm} \wedge \varphi + \frac{4}{7} (\text{tr} \sigma_{\pm}) \psi + \left(\sigma_{\pm}^{(0)} \right)_{ij} e^i \wedge \iota_{e^j} \psi \right]. \end{aligned} \quad (4.10)$$

To arrive at the above, we used the relations:

$$\varphi \Delta_1 \left(\mathcal{F}_{\pm}^{[7]} \right)_{\perp} = \mp 3 \iota_{\beta_{1\pm}} \psi, \quad \psi \Delta_1 \left(\mathcal{F}_{\pm}^{[7]} \right)_{\perp} = \mp 3 \beta_{1\pm} \wedge \varphi,$$

which follow from (4.9) and the identities given in the appendix of [50].

Spin(7) $_{\pm}$ representation	1	7	27	35
component	$F_{\pm}^{[1]} \in \Omega^{4\mp}(\mathcal{U})$	$F_{\pm}^{[7]} \in \Omega^{4\mp}(\mathcal{U})$	$F_{\pm}^{[27]} \in \Omega^{4\mp}(\mathcal{U})$	$F_{\pm}^{[35]} \in \Omega^{4\pm}(\mathcal{U})$
\mathcal{U} -tensors	$\mathcal{F}_{\pm}^{[1]} \in \Omega^0(\mathcal{U})$	$\mathcal{F}_{\pm}^{[7]} \in \Omega^2(\mathcal{U})$	$\mathcal{F}_{\pm}^{[27]} \in \text{Sym}_{\mathcal{U},0}^2(\mathcal{D}^*)$	$\mathcal{F}_{\pm}^{[35]} \in \text{Sym}_0^2(T^*\mathcal{U})$
\mathcal{D} -tensors	$\mathcal{F}_{\pm}^{[1]} \in \Omega^0(\mathcal{U})$	$\beta_{1\pm} \in \Omega_{\mathcal{U}}^1(\mathcal{D})$	$\mathcal{F}_{\pm}^{[27]} \in \text{Sym}_{\mathcal{U},0}^2(\mathcal{D}^*)$	$\beta_{2\pm} \in \Omega_{\mathcal{U}}^1(\mathcal{D})$ $\sigma \in \text{Sym}_{\mathcal{U}}^2(\mathcal{D}^*)$

Table 3. The Spin(7) $_{\pm}$ parameterization of F on the non-chiral locus and its \mathcal{D} -refined version.

Relating the G_2 and Spin(7) $_{\pm}$ parameterizations of F . Relation (4.4) implies:

$$\begin{aligned}
 \left(F_{\pm}^{[k]}\right)_{\mp} &= \frac{1}{2} \left(F_{\mp}^{(k)} \pm *_\perp F_{\pm}^{(k)}\right), & \left(F_{\pm}^{[k]}\right)_{\pm} &= \frac{1}{2} \left(F_{\pm}^{(k)} \pm *_\perp F_{\mp}^{(k)}\right) \quad \text{for } k=1, 7, 27, \\
 \left(F_{\pm}^{[35]}\right)_{\mp} &= \frac{1}{2} \left(F_{\mp} \mp *_\perp F_{\pm}\right), & \left(F_{\pm}^{[35]}\right)_{\pm} &= \frac{1}{2} \left(F_{\pm} \mp *_\perp F_{\mp}\right).
 \end{aligned} \tag{4.11}$$

Comparing (4.10) with (4.11) and using the G_2 parameterization of F_{\mp} and F_{\pm} given in (4.5), one can express the quantities in the last row of table 3 in terms of α_1, α_2 and $\hat{h}, \hat{\chi}$:

$$\begin{aligned}
 \mathcal{F}_{\pm}^{[1]} &= -12 \text{tr}_g \left(\hat{h} \pm \hat{\chi}\right), \\
 \sigma_{\pm} &= -12 \left(\hat{h} \mp \hat{\chi}\right), \\
 \mathcal{F}_{\pm}^{[27]} &= -12 \left(\hat{h}^{(0)} \pm \hat{\chi}^{(0)}\right), \\
 \beta_{1\pm} &= -12(\alpha_2 \pm \alpha_1), \\
 \beta_{2\pm} &= +12(\alpha_1 \mp \alpha_2).
 \end{aligned} \tag{4.12}$$

These simple relations provide the connection between the G_2 parameterization (4.5) and the refined Spin(7) $_{\pm}$ parameterizations (4.10), thus allowing one to relate the G_2 and Spin(7) $_{\pm}$ decompositions of F .

4.3 Relating the G_2 torsion classes to the Lee form and characteristic torsion of the Spin(7) $_{\pm}$ structures

Recall that the *Lee form* of the Spin(7) $_{\pm}$ structure determined by Φ^{\pm} on \mathcal{U} is the one-form defined through:

$$\theta_{\pm} \stackrel{\text{def.}}{=} \pm \frac{1}{7} * (\Phi^{\pm} \wedge \delta \Phi^{\pm}) = -\frac{1}{7} * [\Phi^{\pm} \wedge (*d\Phi^{\pm})] \in \Omega^1(\mathcal{U}) \implies \Phi^{\pm} \wedge \delta \Phi^{\pm} = \mp 7 * \theta_{\pm}, \tag{4.13}$$

where we use the conventions of [51] and the fact that $*\Phi^{\pm} = \pm \Phi^{\pm}$. Also recall from loc. cit. that there exists a unique g -compatible connection ∇^c with skew-symmetric torsion such that $\nabla^c \Phi^{\pm} = 0$. This connection is called the *characteristic connection* of the Spin(7) $_{\pm}$ structure. Its torsion form (obtained by lowering the upper index of the torsion tensor of ∇^c) is given by:

$$T_{\pm} = -\delta \Phi^{\pm} \mp \frac{7}{6} * (\theta_{\pm} \wedge \Phi^{\pm}) = -\delta \Phi^{\pm} - \frac{7}{6} \iota_{\theta_{\pm}} \Phi^{\pm} = \pm * \left(d\Phi^{\pm} - \frac{7}{6} \theta_{\pm} \wedge \Phi^{\pm} \right) \in \Omega^3(\mathcal{U}) \tag{4.14}$$

and is called the *characteristic torsion* of the $\text{Spin}(7)_\pm$ structure. The normalization relation $\|\Phi^\pm\|^2 = 14$, i.e. $\Phi^\pm \wedge \Phi^\pm = \pm 14\nu$ implies $\Phi^\pm \wedge \iota_{\theta_\pm} \Phi^\pm = \pm 7 * \theta_\pm$. Thus $\Phi^\pm \wedge T_\pm = \mp \frac{7}{6} * \theta_\pm$, where we used (4.13) and (4.14). It follows that the Lee form is determined by the characteristic torsion through the equation:

$$\theta_\pm = \pm \frac{6}{7} * (\Phi^\pm \wedge T_\pm). \quad (4.15)$$

Relation (4.14) shows that the exterior derivative of Φ^\pm takes the form:

$$d\Phi^\pm = \frac{7}{6} \theta_\pm \wedge \Phi^\pm \mp * T_\pm = \pm [* (\Phi^\pm \wedge T_\pm)] \wedge \Phi^\pm \mp * T_\pm. \quad (4.16)$$

Recall the relation (see [8]):

$$D_n \psi = -3\vartheta \wedge \varphi,$$

where $\vartheta \in \Omega^1(\mathcal{D})$. Together with (3.38) and with the formula for the exterior derivative of longitudinal forms (see appendix C. of [8]), this gives:

$$\begin{aligned} (d\Phi^\pm)_\top &= \pm (H_\sharp \mp 3\vartheta - 3\tau_1) \wedge \varphi - \left(\frac{4}{7} \text{tr} A \pm \tau_0 \right) \psi - A_{jk}^{(0)} e^j \wedge \iota_{e^k} \psi \mp *_\perp \tau_3, \\ (d\Phi^\pm)_\perp &= 4\tau_1 \wedge \psi + *_\perp \tau_2, \end{aligned}$$

which implies:

$$\begin{aligned} (*d\Phi^\pm)_\top &= -\tau_2 - 4\iota_{\tau_1} \varphi, \\ (*d\Phi^\pm)_\perp &= \mp \iota_{(H_\sharp \mp 3\vartheta - 3\tau_1)} \psi - \left(\frac{4}{7} \text{tr} A \pm \tau_0 \right) \varphi + A_{jk}^{(0)} e^j \wedge \iota_{e^k} \varphi \mp \tau_3. \end{aligned} \quad (4.17)$$

Using this relation and (3.38), we can compute θ_\pm from (4.13) and then determine T_\pm from equation (4.14). We find:

$$\begin{aligned} (\theta_\pm)_\top &= -\frac{4}{7} \text{tr} A \mp \tau_0, & (\theta_\pm)_\perp &= -\frac{4}{7} (H_\sharp \mp 3\vartheta - 6\tau_1), \\ (T_\pm)_\top &= -\frac{2}{3} \iota_{(\pm H_\sharp - 3\vartheta)} \varphi \mp \tau_2, & (T_\pm)_\perp &= -\frac{1}{6} \left(\frac{4}{7} \text{tr} A \pm \tau_0 \right) \varphi - \frac{1}{3} \iota_{(H_\sharp \mp 3\vartheta + 3\tau_1)} \psi \pm A_{jk}^{(0)} e^j \wedge \iota_{e^k} \varphi - \tau_3. \end{aligned} \quad (4.18)$$

To arrive at the last two relations, we used the identities:

$$\iota_{\tau_2} \varphi = \iota_{\tau_3} \psi = \langle \tau_3, \varphi \rangle = 0,$$

which follow from relations (B.13) and (B.14) given in appendix B of [8] upon using the fact that $\tau_3 \in \Omega_{\mathcal{U}, 27}^3(\mathcal{D})$.

4.4 Relation to previous work

The problem of determining the fluxes f, F in terms of the geometry along the locus \mathcal{U}^+ was considered in reference [1], where the quantities denoted here by L^+, Φ^+ were denoted simply by L, Φ . Using the results of the previous subsections, one can show that the relations given in Theorem 3 of [8] are equivalent, on the non-chiral locus \mathcal{U} , with equations (3.16), (3.17) and (3.18) of [1]. This solves the problem of comparing the approach of loc. cit. with that of [2, 8]. The major steps of the comparison with loc. cit. are given in appendix C.

5 Description of the singular foliation in the Morse case

In this section, we consider the case when the closed one-form $\omega \in \Omega^1(M)$ is Morse. This case is generic in the sense that Morse one-forms form a dense open subset of the set of all closed one-forms belonging to the fixed cohomology class \mathfrak{f} — hence a form which satisfies equations (3.41) can be replaced by a Morse form by infinitesimally perturbing b . Singular foliations defined by Morse one-forms were studied in [21]–[29] and [52]–[57]. Let $\Pi_{\mathfrak{f}} = \text{im}(\text{per}_{\mathfrak{f}}) \subset \mathbb{R}$ be the period group of the cohomology class \mathfrak{f} and $\rho(\mathfrak{f}) = \text{rk}\Pi_{\mathfrak{f}}$ be its irrationality rank. The general results summarized in the following subsection hold for any smooth, compact and connected manifold of dimension d which is strictly bigger than two, under the assumption that the set of zeroes of ω (which in Novikov theory [58] is called the set of *singular points*):

$$\text{Sing}(\omega) \stackrel{\text{def.}}{=} \{p \in M | \omega_p = 0\}$$

is non-empty. Notice that $\text{Sing}(\omega)$ is a finite set since M is compact and since the zeroes of a Morse 1-form are isolated. The complement:

$$M^* \stackrel{\text{def.}}{=} M \setminus \text{Sing}(\omega)$$

is a non-compact open submanifold of M . Below, we shall use the notations \mathcal{F}_{ω} for the regular foliation induced by ω on M^* and $\bar{\mathcal{F}}_{\omega}$ for the singular foliation induced on M . In our application we have $n = 8$ and:

$$\text{Sing}(\omega) = \mathcal{W}, \quad M^* = \mathcal{U}, \quad \mathcal{F}_{\omega} = \mathcal{F}, \quad \bar{\mathcal{F}}_{\omega} = \bar{\mathcal{F}}.$$

5.1 Types of singular points

Let $\text{ind}_p(\omega)$ denote the Morse index of a point $p \in \text{Sing}(\omega)$, i.e. the Morse index at p of a Morse function $h_p \in C^{\infty}(U_p, \mathbb{R})$ such that dh_p equals $\omega|_{U_p}$, where U_p is some vicinity of p . This index does not depend on the choice of U_p and h_p . Let:

$$\begin{aligned} \text{Sing}_k(\omega) &\stackrel{\text{def.}}{=} \{p \in \text{Sing}(\omega) | \text{ind}_p(\omega) = k\}, & k = 1, \dots, d \\ \Sigma_k(\omega) &\stackrel{\text{def.}}{=} \{p \in \text{Sing}(\omega) | \text{ind}_p(\omega) = k \text{ or } \text{ind}_p(\omega) = d - k\}, & k = 1, \dots, \left\lfloor \frac{d}{2} \right\rfloor. \end{aligned}$$

Thus $\Sigma_k(\omega) = \text{Sing}_k(\omega) \cup \text{Sing}_{d-k}(\omega)$ for $k < \frac{d}{2}$ and $\Sigma_{d_0}(\omega) = \text{Sing}_{d_0}(\omega)$ when $d = 2d_0$ is even. In a small enough vicinity of $p \in \text{Sing}_k(\omega)$ (which we can assume to equal U_p by shrinking the latter if necessary), the Morse lemma applied to h_p implies that there exists a local coordinate system (x_1, \dots, x_d) such that:

$$h_p = - \sum_{j=1}^k x_j^2 + \sum_{j=k+1}^d x_j^2.$$

Definition. The elements of $\Sigma_0(\omega)$ are called *centers* while all other singularities of ω are called *saddle points*. The elements of $\Sigma_1(\omega)$ are called *strong saddle points*, while all other saddle points are called *weak*.

Remark. Strong saddle points are sometimes called “conical points”. That terminology can lead to confusion, since all singular points which are not centers are conical singularities of the singular leaf to which they belong (see below), in the sense that the singular leaf can be modeled by a cone (with one or two sheets) in a vicinity of such a singular point. In other references, a “conical point” means any singularity which is not a center, i.e. what we call a saddle point.

5.2 The regular and singular foliations defined by a Morse 1-form

The regular foliation \mathcal{F}_ω . The Morse 1-form ω defines a regular foliation \mathcal{F}_ω of the open submanifold M^* , namely the foliation which, by the Frobenius theorem, integrates the regular Frobenius distribution $\ker(\omega)|_{M^*}$. Following [27], we say that a singular point $p \in \text{Sing}(\omega)$ *adjoins* a leaf L^5 of \mathcal{F}_ω if the union $\{p\} \cup L$ is connected; notice that a center cannot adjoin any leaf of \mathcal{F}_ω . Let:

$$s(L) \stackrel{\text{def.}}{=} \{p \in \text{Sing}(\omega) | p \text{ adjoins } L\} \subset \text{Sing}(\omega).$$

The set $s(L)$ is contained in the intersection of $\text{Sing}(\omega)$ with the small topological frontier $\text{fr}(L)$ of L :

$$s(L) \subseteq \text{fr}(L) \cap \text{Sing}(\omega). \tag{5.1}$$

Notice that this inclusion can be strict; a beautifully drawn example illustrating this in the two-dimensional case can be found in [29] (see figure 2(c) of loc. cit.). We have $s(L) \cap \Sigma_0(\omega) = \emptyset$ and hence $s(L) = \bigsqcup_{k=1}^{\lfloor \frac{d}{2} \rfloor} s_k(L)$, where:

$$s_k(L) \stackrel{\text{def.}}{=} s(L) \cap \Sigma_k(L).$$

Classification of the leaves of \mathcal{F}_ω .

- *Compactifiable and non-compactifiable leaves.* We say that a leaf L of \mathcal{F}_ω is *compactifiable* if the set $L \cup \text{Sing}(\omega)$ is compact, which amounts to the condition that the small topological frontier $\text{fr}(L) \stackrel{\text{def.}}{=} \bar{L} \setminus L$ of L in M is a (possibly void) subset of $\text{Sing}(\omega)$ and hence a finite set. With this definition, compact leaves of \mathcal{F}_ω are compactifiable, but not all compactifiable leaves are compact. A *non-compactifiable leaf* of \mathcal{F}_ω is a leaf which is not compactifiable; obviously such a leaf is also non-compact. The closure of a non-compactifiable leaf is a set with non-empty interior [52, 55], so the small frontier of such a leaf is an infinite set.
- *Ordinary and special leaves.* The leaf L is called *ordinary* if $s(L)$ is empty and *special* if $s(L)$ is non-empty. An ordinary leaf is either compact or non-compactifiable. Any non-compact but compactifiable leaf is a special leaf, but there also exist non-compactifiable special leaves (see table 4).

⁵This should not be confused with the quantity L^\pm discussed in subsection 4.4 (or with the quantity denoted by L in [1]).

type of L	compactifiable		non-compactifiable
	compact	non-compact	
ordinary	Y	—	Y
special	—	Y	Y
$\text{Card}(\text{fr}L)$	finite		infinite

Table 4. Classification of the leaves of \mathcal{F}_ω , where the allowed combinations are indicated by the letter “Y”. A compactifiable leaf is ordinary iff it is compact and it is special iff it is non-compact. A non-compactifiable leaf may be either ordinary or special. Non-compactifiable leaves coincide [52, 55] with those leaves whose small frontier is an infinite set, while compactifiable leaves are those leaves whose small frontier is finite.

The foliation \mathcal{F}_ω has only a finite number of special leaves, because the local form of leaves near the points of $\text{Sing}(\omega)$ (see below) shows that at most two special leaves can contain each such point in their closures (recall that we assume $d \geq 3$). We shall see later that each non-compactifiable leaf (whether special or not) covers densely some open and connected subset of M^* . Notice that every singular point which is not a center adjoins some special leaf. Hence:

$$\Sigma_k(\omega) = \cup_{L=\text{special leaf of } \mathcal{F}_\omega} s_k(L), \quad \forall k = 1 \dots \left\lfloor \frac{d}{2} \right\rfloor. \tag{5.2}$$

The singular foliation $\bar{\mathcal{F}}_\omega$. One can describe [18, 58] the singular foliation $\bar{\mathcal{F}}_\omega$ of M defined by ω as the partition of M induced by the equivalence relation \sim defined as follows. We put $p \sim q$ if there exists a smooth curve $\gamma : [0, 1] \rightarrow M$ such that:

$$\gamma(0) = p, \quad \gamma(1) = q \quad \text{and} \quad \omega(\dot{\gamma}(t)) = 0 \quad \forall t \in [0, 1].$$

The leaves of $\bar{\mathcal{F}}_\omega$ are the equivalence classes of this relation; they are connected subsets of M (which need not be topological manifolds when endowed with the induced topology). Any such leaf is either of the form $\{p\}$ where $p \in \Sigma_0(\omega)$ is a center or is a topological subspace of M of Lebesgue covering dimension equal to $n - 1$.

Remark. We stress that $\bar{\mathcal{F}}_\omega$ is not generally a foliation of M in the ordinary sense of foliation theory but (as explained in the previous section) it should be viewed as a Haefliger structure. It is not even a \mathcal{C}^0 -foliation, i.e. a foliation in the category of topological manifolds (locally Euclidean Hausdorff topological spaces), because singular leaves of $\bar{\mathcal{F}}_\omega$ which pass through strong saddle points can be locally disconnected by removing those points and hence are not topological manifolds.

Regular and singular leaves of $\bar{\mathcal{F}}_\omega$. A leaf \mathcal{L} of $\bar{\mathcal{F}}_\omega$ is called *singular* if it intersects $\text{Sing}(\omega)$ and *regular* otherwise. The regular leaves of $\bar{\mathcal{F}}_\omega$ coincide with the ordinary leaves of \mathcal{F}_ω ; notice that every center singularity is a singular leaf. On the other hand, each singular leaf which is not a center is a disjoint union of a finite number of special leaves

of \mathcal{F}_ω and of some subset $s(\mathcal{L}) \stackrel{\text{def.}}{=} \mathcal{L} \cap \text{Sing}(\omega)$ of $\text{Sing}(\omega)$, which we shall call the *set of singular points of \mathcal{L}* . We have:

$$\mathcal{L} \setminus s(\mathcal{L}) = L_1 \sqcup \dots \sqcup L_r \sqcup L'_1 \sqcup \dots \sqcup L'_t,$$

where L_1, \dots, L_r are compactifiable special leaves while L'_1, \dots, L'_t are non-compactifiable special leaves of \mathcal{F}_ω . We also have $s(\mathcal{L}) = s^c(\mathcal{L}) \cup s^{\text{nc}}(\mathcal{L})$ (generally a non-disjoint union), with:

$$s^c(\mathcal{L}) \stackrel{\text{def.}}{=} \cup_{i=1}^r s(L_i), \quad s^{\text{nc}}(\mathcal{L}) \stackrel{\text{def.}}{=} \cup_{j=1}^t (L'_j).$$

The singular leaf \mathcal{L} decomposes as:

$$\mathcal{L} = \mathcal{L}^c \sqcup \mathcal{L}^{\text{nc}}, \tag{5.3}$$

where the *compact part* and *non-compact part* of \mathcal{L} are defined through:

$$\begin{aligned} \mathcal{L}^c &\stackrel{\text{def.}}{=} \bar{L}_1 \cup \dots \cup \bar{L}_r = L_1 \sqcup \dots \sqcup L_r \sqcup s^c(\mathcal{L}) \\ \mathcal{L}^{\text{nc}} &\stackrel{\text{def.}}{=} \mathcal{L} \setminus \mathcal{L}^c = (L'_1 \sqcup \dots \sqcup L'_t) \sqcup (s(\mathcal{L}) \setminus s^c(\mathcal{L})). \end{aligned} \tag{5.4}$$

The set $s^c(\mathcal{L})$ consists of those singular points of \mathcal{L} which lie on the compact part \mathcal{L}^c . Notice that both the compact and non-compact parts of \mathcal{L} can be void and that a non-compactifiable special leaf component L'_j of \mathcal{L} can adjoin points from $s^c(\mathcal{L})$ as well as from $s(\mathcal{L}) \setminus s^c(\mathcal{L})$ simultaneously; furthermore, \mathcal{L}^{nc} may meet itself at certain points of $s(\mathcal{L}) \setminus s^c(\mathcal{L})$.⁶ When \mathcal{L} is a center leaf $\{p\}$, we define $\mathcal{L}^c \stackrel{\text{def.}}{=} s(\mathcal{L}) = \{p\}$ and $\mathcal{L}^{\text{nc}} \stackrel{\text{def.}}{=} \emptyset$. Notice that any non-empty subset A of \mathcal{L} determines \mathcal{L} as the saturation of A with respect to the equivalence relation \sim . If S_ω denotes the union of $\text{Sing}(\omega)$ with all special leaves of \mathcal{F}_ω , then the singular leaves of $\bar{\mathcal{F}}_\omega$ (including the centers) coincide with the connected components of S_ω . Notice that $\text{fr}(L_i) = s(L_i) = \bar{L}_i \cap \text{Sing}(\omega)$ for each compactifiable leaf component L_i , $i = 1 \dots r$. The compact sets \bar{L}_i meet themselves or each other only in strong saddle points. In particular, we have:

$$\bar{L}_{i_1} \cap \bar{L}_{i_2} = s(L_{i_1}) \cap s(L_{i_2}) = s_1(L_{i_1}) \cap s_1(L_{i_2}) \subset \Sigma_1(\omega) \quad \text{for } 1 \leq i_1 < i_2 \leq r.$$

The following definition generalizes the notion of generic Morse function:

Definition. The Morse form ω is called *generic* if every singular leaf of $\bar{\mathcal{F}}_\omega$ contains exactly one singular point $p \in \text{Sing}(\omega)$.

5.3 Behavior of the singular leaves near singular points

In a small enough vicinity of $p \in \text{Sing}_k(\omega)$, the singular leaf \mathcal{L}_p passing through p is modeled by the locus $Q_k \subset \mathbb{R}^n$ given by the equation $h_p = 0$, where p corresponds to the origin of \mathbb{R}^n . One distinguishes the cases (see tables 5 and 6):

- $k \in \{0, n\}$, i.e. p is a center. Then $\mathcal{L}_p = \{p\}$ and the nearby leaves of \mathcal{F}_p are diffeomorphic to S^{n-1} .

⁶We thank I. Gelbukh for drawing our attention to these points.






Name	Morse index	Local form of \mathcal{L}_p	Local form of regular leaves
Center	0 or n	$\bullet = \{p\}$	
Weak saddle	between 2 and $n - 2$		
Strong saddle	1 or $n - 1$		

Table 5. Types of singular points p . The first and third figure on the right depict the case $d = 3$ for centers and strong saddles, while the second figure attempts to depict the case $d > 3$ for a weak saddle (notice that weak saddles do not exist unless $d > 3$). In that case, the topology of the leaves does not change locally when they “pass through” the weak saddle point. \mathcal{L}_p denotes the singular leaf of $\bar{\mathcal{F}}_\omega$ which passes through p .

- $2 \leq k \leq n - 2$, i.e. p is a weak saddle point. Then Q_k is diffeomorphic to a cone over $S^{k-1} \times S^{n-k-1}$ and $\mathbb{R}^n \setminus Q_k$ has two connected components while $Q_k \setminus \{p\}$ is connected. Removing p does not *locally* disconnect \mathcal{L}_p .
- $k \in \{1, n - 1\}$, i.e. p is a strong saddle point. Then Q_k is diffeomorphic to a cone over $\{-1, 1\} \times S^{n-2}$ and $\mathbb{R}^n \setminus Q_k$ has three connected components while $Q_k \setminus \{0\}$ has two components. Removing p *locally* disconnects \mathcal{L}_p . A strong saddle point $p \in \Sigma_1(\omega)$ is called *splitting* [27] (or *blocking* [54]) if it adjoins two different (special) leaves of the regular foliation \mathcal{F}_ω and it is called *non-splitting* (or a *transformation point* [27]) if it adjoins a single (special) leaf of \mathcal{F}_ω (see table 6). If a singular leaf \mathcal{L} contains only one splitting point, then removing it disconnects \mathcal{L} . If a singular leaf \mathcal{L} contains more than one splitting point, then removing it may not disconnect \mathcal{L} (an example of such behavior is given in [27, figure 7(b)]).

We have a decomposition $\Sigma_1(\omega) = \Sigma_1^{\text{sp}}(\omega) \sqcup \Sigma_1^{\text{nsp}}(\omega)$ of the set of strong saddle points, where:

$$\Sigma_1^{\text{sp}}(\omega) \stackrel{\text{def.}}{=} \{p \in \Sigma_1(\omega) | p \text{ is a splitting singularity}\}$$

$$\Sigma_1^{\text{nsp}}(\omega) \stackrel{\text{def.}}{=} \{p \in \Sigma_1(\omega) | p \text{ is a non - splitting singularity}\}.$$

Taking into account the local behavior of leaves near the various types of singular points, we find that the decomposition (5.2) is disjoint for $k \neq 1$:

$$\Sigma_k(\omega) = \sqcup_{L=\text{special leaf of } \mathcal{F}_\omega} s_k(L), \quad \forall k = 2 \dots \left\lfloor \frac{d}{2} \right\rfloor.$$



Singularity type	Example of global shape for \mathcal{L}_p
Splitting	
Non-splitting	

Table 6. Types of strong saddle points. The figures illustrate the two types through two simple examples in the case $d = 3$. The figure in the first row uses different colors to indicate two different special compactifiable leaves of \mathcal{F}_ω which are subsets of the same singular leaf of $\bar{\mathcal{F}}_\omega$, each of them adjoining the same splitting singular point. The figure in the second row shows a single special compactifiable leaf of \mathcal{F}_ω which adjoins a single non-splitting singular point.

while the decomposition for $k = 1$ may fail to be disjoint:

$$\Sigma_1(\omega) = \cup_{L=\text{special leaf of } \mathcal{F}_\omega} s_1(L). \tag{5.5}$$

More precisely:

$$\begin{aligned} \Sigma_1^{\text{nsp}}(\omega) &= \sqcup_{L=\text{special leaf of } \mathcal{F}_\omega} s_1^{\text{nsp}}(L) \\ \Sigma_1^{\text{sp}}(\omega) &= \cup_{L=\text{special leaf of } \mathcal{F}_\omega} s_1^{\text{sp}}(L), \end{aligned}$$

where we defined:

$$s_1^{\text{nsp}}(L) \stackrel{\text{def.}}{=} s_1(L) \cap \Sigma_1^{\text{nsp}}(\omega), \quad s_1^{\text{sp}}(L) \stackrel{\text{def.}}{=} s_1(L) \cap \Sigma_1^{\text{sp}}(\omega)$$

and where the second union may be non-disjoint. This is because two distinct special leaves of \mathcal{F}_ω can meet each other only at a strong saddle point which is a splitting singularity.

5.4 Combinatorics of singular leaves

Definition. A singular leaf of $\bar{\mathcal{F}}_\omega$ which is not a center is called a *strong singular leaf* if it contains at least one strong saddle point and a *weak singular leaf* otherwise.

A weak singular leaf is obtained by adjoining weak saddle points to a single special leaf of \mathcal{F}_ω . Such singular leaves are mutually disjoint and their singular points determine a partition of the set $\Sigma_{>1}(\omega) \stackrel{\text{def.}}{=} \cup_{k=2}^{\lfloor \frac{d}{2} \rfloor} \Sigma_k(\omega)$. The situation is more complicated for strong singular leaves, as we now describe.

At each $p \in \Sigma_1(\omega)$, consider the strong singular leaf \mathcal{L} passing through p . The intersection of $\mathcal{L} \setminus \{p\}$ with a sufficiently small neighborhood of p is a disconnected manifold diffeomorphic to a union of two cones without apex, whose rays near p determine a connected cone $C_p \subset T_p M$ inside the tangent space to M at p (see the last row of table 5). The set $\dot{C}_p \stackrel{\text{def.}}{=} C_p \setminus \{0_p\}$ (where 0_p is the zero vector of $T_p M$) has two connected components, thus $\pi_0(\dot{C}_p)$ is a two-element set. Hence the finite set:

$$\hat{\Sigma}_1(\omega) \stackrel{\text{def.}}{=} \sqcup_{p \in \Sigma_1(M)} \pi_0(\dot{C}_p)$$

is a double cover of $\Sigma_1(\omega)$ through the projection σ that takes $\pi_0(\dot{C}_p)$ to $\{p\}$. Consider the complete unoriented graph having as vertices the elements of $\hat{\Sigma}_1(\omega)$. This graph has a dimer covering given by the collection of edges:

$$\hat{\mathcal{E}} = \left\{ \pi_0(\dot{C}_p) \mid p \in \Sigma_1(\omega) \right\},$$

which connect vertically the vertices lying above the same point of $\Sigma_1(\omega)$ (see figure 2). If L is a special leaf of \mathcal{F}_ω and $p \in \Sigma_1(\omega)$ adjoins L , then the connected components of the intersection of L with a sufficiently small vicinity of p are locally approximated at p by one or two of the connected components of \dot{C}_p . The second case occurs iff p is a non-splitting strong saddle point (see table 6). Hence L determines a subset $\hat{s}_1(L)$ of $\hat{\Sigma}_1(\omega)$ such that $\sigma(\hat{s}_1(L)) = s_1(L)$ and such that the fiber of $\hat{s}_1(L)$ above a point $p \in s_1(L)$ has one element if p is a splitting singularity and two elements if p is non-splitting. If L' is a different special leaf of \mathcal{F}_ω , then the sets $\hat{s}_1(L')$ and $\hat{s}_1(L)$ are disjoint, even though their projections $s_1(L)$ and $s_1(L')$ through σ may intersect in $\Sigma_1(\omega)$. Hence the special leaves of \mathcal{F}_ω define a partition of $\hat{\Sigma}_1(\omega)$:

$$\hat{\Sigma}_1(\omega) = \sqcup_{L=\text{special leaf of } \mathcal{F}_\omega} \hat{s}_1(L),$$

which projects through σ to the generally non-disjoint decomposition (5.5). Viewing $\hat{\mathcal{E}}$ as a disconnected graph on the vertex set $\hat{\Sigma}_1(\omega)$, we let \mathcal{E} denote the (generally disconnected) graph obtained from $\hat{\mathcal{E}}$ upon identifying all vertices belonging to $\hat{s}_1(L)$ for each special leaf L of \mathcal{F}_ω , i.e. by collapsing $\hat{s}_1(L)$ to a point for each special leaf⁷ L . Let $p : \hat{\mathcal{E}} \rightarrow \mathcal{E}$ denote the corresponding projection. The graph \mathcal{E} has one vertex for each special leaf of \mathcal{F}_ω which adjoins some strong saddle point and an edge for each strong saddle point. Notice that this edge is a loop when the strong saddle point is a non-splitting singularity, since a non-splitting singularity adjoins a single special leaf. A strong singular leaf of \mathcal{F}_ω can be written as:

$$\mathcal{L} = \left(\sqcup_{\alpha=1}^{r+t} L''_\alpha \right) \sqcup s(\mathcal{L}), \tag{5.6}$$

where L''_α are special leaves of \mathcal{F}_ω (compactifiable or not). Its set of strong saddle singular points $s_1(\mathcal{L}) = \cup_{\alpha=1}^{r+t} s_1(L''_\alpha)$ is the projection through σ of the set $\hat{s}_1(\mathcal{L}) \stackrel{\text{def.}}{=} \sqcup_{\alpha=1}^{r+t} \hat{s}_1(L''_\alpha)$. Let $\hat{\mathcal{E}}_\mathcal{L}$ be the (generally disconnected) subgraph of $\hat{\mathcal{E}}$ consisting of those edges of $\hat{\mathcal{E}}$ which meet $\hat{s}_1(\mathcal{L})$. Then $s_1(\mathcal{L})$ is obtained from $\hat{\mathcal{E}}_\mathcal{L}$ by contracting each edge to a single point. If all special leaves L of \mathcal{F}_ω are known, then $\hat{\mathcal{E}}_\mathcal{L}$ uniquely determines the strong singular leaf \mathcal{L} . Indeed, $\hat{\mathcal{E}}_\mathcal{L}$ contains the information about how the special leaves which form \mathcal{L} meet themselves and each other at the strong saddle points. Since \mathcal{L} is connected and maximal with this property, the graph $\mathcal{E}_\mathcal{L}$ obtained from $\hat{\mathcal{E}}_\mathcal{L}$ by identifying to a single point the vertices of each of the subsets $\hat{s}_1(L''_\alpha)$ is a connected component of \mathcal{E} . It follows that the strong singular leaves of \mathcal{F}_ω are in one to one correspondence with the connected components of the graph \mathcal{E} — namely, their subgraphs $\hat{\mathcal{E}}_\mathcal{L}$ are the preimages through p of those components.

⁷If $s_1(L)$ is empty, this operation does nothing.




Figure 2. Example of the graphs $\hat{\mathcal{E}}$ and \mathcal{E} for a Morse form foliation $\bar{\mathcal{F}}_\omega$ with two compact strong singular leaves. The regular foliation \mathcal{F}_ω of M^* has four special leaves, each of which is compactifiable; they are depicted using four different colors. At the bottom of the picture, we depict $\Sigma_1(\omega)$ as well as the schematic shape of the special leaves in the case $d = 3$. The strong singular leaves of $\bar{\mathcal{F}}_\omega$ correspond to the left and right parts of the figure at the bottom; each of them is a union of two special leaves of \mathcal{F}_ω and of singular points. Each special leaf corresponds to a vertex of \mathcal{E} .

In our application, the set $\text{Sing}(\omega) = \mathcal{W} = \mathcal{W}^+ \sqcup \mathcal{W}^-$ consists of positive and negative chirality points of ξ , which are the points where b attains the values $b = \pm 1$. Relation (3.41) implies that \mathbf{f} satisfies:

$$\oint_\gamma \mathbf{f} = 0$$

for any smooth closed curve $\gamma \in \mathcal{L} \setminus \mathcal{W}$ and hence \mathbf{f} restricts to a trivial class in singular cohomology along each leaf \mathcal{L} of $\bar{\mathcal{F}}$:

$$\iota^*(\mathbf{f}) = 0 \in H^1(\mathcal{L}, \mathbb{R}),$$

where $\iota : \mathcal{L} \hookrightarrow M$ is the inclusion map while $H^1(\mathcal{L}, \mathbb{R})$ is the first singular cohomology group (which coincides with the first de Rham cohomology group when \mathcal{L} is non-singular). The pull-back of \mathbf{f} to $\mathcal{L} \setminus \mathcal{W}$ is given by:

$$\mathbf{f}|_{\mathcal{L} \setminus \mathcal{W}} = \mathbf{f}_\perp = d_\perp \mathbf{b}.$$

Notice that f_\perp and b have well-defined limits (equal to f_p and $b(p) \in \{-1, 1\}$) at each singular point $p \in \mathcal{L} \cap \mathcal{W}$ of a singular leaf \mathcal{L} . If $p_1, p_2 \in \mathcal{L} \cap \mathcal{W}$ are two singular points lying on the same singular leaf \mathcal{L} and $\gamma : (0, 1) \rightarrow \mathcal{L} \setminus \mathcal{W}$ is a smooth path which has limits at 0, 1 given by p_1 and p_2 , then the integral $\int_\gamma \mathbf{f}$ is well-defined and given by:

$$\int_\gamma \mathbf{f} = e^{3\Delta(p_2)} b(p_2) - e^{3\Delta(p_1)} b(p_1),$$

where $b(p_i) \in \{-1, 1\}$.

5.5 Homology classes of compact leaves

Let H_ω be the (necessarily free) subgroup of $H_{n-1}(M, \mathbb{Z})$ generated by the compact leaves of \mathcal{F}_ω and let $c(\omega) \stackrel{\text{def.}}{=} \text{rk}H_\omega$ denote the number of homologically independent compact leaves. It was shown in [21] that H_ω admits a basis consisting of homology classes $[L_i]$ ($i = 1, \dots, c(\omega)$) of compact leaves⁸ and that the homology class of any compact leaf L of \mathcal{F}_ω expands in this basis as:

$$[L] = \sum_{i=1}^{c(\omega)} n_i [L_i] \quad \text{where } n_i \in \{-1, 1\}.$$

Furthermore [21, 23], there exists a system of \mathbb{Z} -linearly independent one-cycles $\gamma_i \in H_1(M, \mathbb{Z})$ ($i = 1, \dots, c(\omega)$) such that $(\gamma_i, [L_j]) = \delta_{ij}$ and such that γ_i provide a direct sum decomposition:

$$H_1(M, \mathbb{Z}) = \langle \gamma_1, \dots, \gamma_{c(\omega)} \rangle \oplus \iota_*(H_1(\Delta)),$$

where $\iota : \Delta \hookrightarrow M$ is the inclusion map. Let $\mathcal{H}_\omega \stackrel{\text{def.}}{=} H_\omega \cap (\ker \text{per}_\omega)^\perp$. Then [25] the subgroup \mathcal{H}_ω is a direct summand in H_ω while H_ω is a direct summand in $H_{n-1}(M, \mathbb{Z})$. Furthermore, only the following values are allowed for $\text{rk}\mathcal{H}_\omega$:

$$\text{rk}\mathcal{H}_\omega \in \{0, \dots, \rho(\omega) - 2\} \cup \{\rho(\omega)\}.$$

5.6 The Novikov decomposition of M

What we shall call the ‘‘Novikov decomposition’’ is a generalization of the Morse decomposition [59–61], which was introduced in [19, 20] (see also [18, 57]) and used extensively in [21]–[29]; the name is motivated by analogy with ‘‘Morse decomposition’’, due to the role which this decomposition plays in the modern study of the topology of closed one-forms [58]. Define C^{\max} to be the union of all compact leaves and C^{\min} to be the union of all non-compactifiable leaves of \mathcal{F}_ω ; it is clear that these two subsets of M are disjoint. Then it was shown in [52, 55] that both C^{\max} and C^{\min} are open subsets of M which have a common topological small frontier F^9 given by the (disjoint) union $F_0 \cup \text{Sing}(\omega)$, where F_0 is the union of all those leaves of \mathcal{F}_ω which are compactifiable but non-compact:

$$\text{fr}C^{\max} = \text{fr}C^{\min} = F \stackrel{\text{def.}}{=} F_0 \sqcup \text{Sing}(\omega).$$

Each of the open sets C^{\max} and C^{\min} has a finite number of connected components, which are called the *maximal* and *minimal* components of the set $M \setminus F = C^{\max} \sqcup C^{\min}$. We let:

- $N_{\max}(\omega) \stackrel{\text{def.}}{=} |\pi_0(C^{\max})|$ denote the number of maximal components
- $N_{\min}(\omega) \stackrel{\text{def.}}{=} |\pi_0(C^{\min})|$ denote the number of minimal components.

⁸Such a basis is provided by the homology classes of the compact leaves corresponding to the edges of any spanning tree of the foliation graph defined below.

⁹This should not be confused with the internal part of the flux which is denoted by the same letter.

Indexing these by C_j^{\max} and C_a^{\min} (where $j = 1, \dots, N_{\max}(\omega)$ and $a = 1, \dots, N_{\min}(\omega)$), we have:

$$C^{\max} = \sqcup_{j=1}^{N_{\max}(\omega)} C_j^{\max}, \quad C^{\min} = \sqcup_{a=1}^{N_{\min}(\omega)} C_a^{\min} \tag{5.7}$$

and hence (since (5.7) are *finite* and *disjoint* unions) we also have:

$$\begin{aligned} \overline{C^{\max}} &= \cup_{j=1}^{N_{\max}(\omega)} \overline{C_j^{\max}}, & \overline{C^{\min}} &= \cup_{a=1}^{N_{\min}(\omega)} \overline{C_a^{\min}}, \\ F = \text{fr} C^{\max} &= \cup_{j=1}^{N_{\max}(\omega)} \text{fr} C_j^{\max} = \text{fr} C^{\min} = \cup_{a=1}^{N_{\min}(\omega)} \text{fr} C_a^{\min}. \end{aligned} \tag{5.8}$$

Notice that the unions appearing in these equalities need not be disjoint anymore, in particular the small frontiers of two distinct maximal components can intersect each other and similarly for two distinct minimal components. Let:¹⁰

$$\Delta \stackrel{\text{def.}}{=} M \setminus C^{\max} = \overline{C^{\min}} = C^{\min} \sqcup F$$

be the union of all non-compact leaves and singularities. This subset has a finite number (which we denote by $v(\omega)$) of connected components Δ_s :

$$\Delta = \sqcup_{s=1}^{v(\omega)} \Delta_s. \tag{5.10}$$

The connected components of F (which are again in finite number) are finite unions of singular points and of non-compact but compactifiable leaves of \mathcal{F}_ω which coincide with the ‘compact parts’ of the singular leaves of $\overline{\mathcal{F}}_\omega$ (see (5.3)).

One can show [18, 53] that each maximal component C_j^{\max} is diffeomorphic to the open unit cylinder over any of the (compact) leaves L_j of the restricted foliation $\mathcal{F}_\omega|_{C_j^{\max}}$, through a diffeomorphism which maps this restricted foliation to the foliation of the cylinder given by its sections $L_j \times \{t\}$:

$$C_j^{\max} \simeq L_j \times (0, 1). \tag{5.11}$$

In particular, we have:

$$\rho(\omega|_{C_j^{\max}}) = 0.$$

Being connected, each non-compactifiable leaf L of \mathcal{F}_ω is contained in exactly one minimal component. It was shown in [55] (see also appendix of [52]) that L is *dense* in that minimal component. Furthermore, one has [18, 52]:

$$\rho(\omega|_{C_a^{\min}}) \geq 2, \quad a = 1, \dots, N_{\min}(\omega).$$

In particular, any minimal component C_a^{\min} must satisfy $b_1(C_a^{\min}) \geq 2$.

Definition. The foliation \mathcal{F}_ω is called *compactifiable* if each of its leaves is compactifiable, i.e. if it has no minimal components.

¹⁰ Δ should not be confused with the warp factor.

5.7 The foliation graph

Since each maximal component C_j^{\max} is a cylinder, its frontier consists of either one or two connected components. When the frontier of C_j^{\max} is connected, there exists exactly one connected component Δ_{s_j} of Δ such that $\text{fr}C_j^{\max} \subset \Delta_{s_j}$. When the frontier of C_j^{\max} has two connected components, there exist distinct indices s'_j and s''_j such that these components are subsets of $\Delta_{s'_j}$ and $\Delta_{s''_j}$, respectively. These observations allow one to define a graph as follows [18, 57]:

Definition. The *foliation graph* Γ_ω of ω is the unoriented graph whose vertices are the connected components Δ_s of Δ and whose edges are the maximal components C_j^{\max} . An edge C_j^{\max} is incident to a vertex Δ_s iff a connected component of $\text{fr}C_j^{\max}$ is contained in Δ_s ; it is a loop at Δ_s iff $\text{fr}C_j^{\max}$ is connected and contained in Δ_s . A vertex Δ_s of Γ_ω is called *exceptional* (or of *type II*) if it contains at least one minimal component; otherwise, it is called *regular* (or of *type I*).

The terminology *type I*, *type II* for vertices is used in [27]. Since M is connected, it follows that Γ_ω is a connected graph. Notice that Γ_ω can have loops and multiple edges as well as terminal vertices. Let $\text{deg}\Delta_s$ denote the degree (valency) of Δ_s as a vertex of the foliation graph. A regular vertex Δ_s can be of two types:

- A center singularity $\Delta_s = \{p\}$ (with $p \in \Sigma_0(\omega)$), when $\text{deg}\Delta_s = 1$. In this case, Δ_s is a terminal vertex of Γ_ω .
- A compact singular leaf when $\text{deg}\Delta_s \geq 2$.

Every exceptional vertex is a union of minimal components, singular points and compactifiable non-compact leaves of \mathcal{F}_ω . For any vertex Δ_s of the foliation graph, we have [27]:

$$|\Delta_s \cap \Sigma_1^{\text{sp}}(\omega)| \geq \text{deg}\Delta_s + 2m_{\Delta_s} - 2,$$

where m_{Δ_s} is the number of minimal components contained in Δ_s . In particular, a regular vertex with $\text{deg}\Delta_s > 2$ is a compact singular leaf which contains at least one splitting strong saddle singularity. The number of edges $e(\Gamma_\omega)$ equals $N_{\max}(\omega)$ while the number of vertices $v(\Gamma_\omega)$ equals $v(\omega)$. Furthermore, it was shown in [28] that the cycle rank $b_1(\Gamma_\omega)$ equals $c(\omega)$. Thus:

$$e(\Gamma_\omega) = N_{\max}(\omega), \quad v(\Gamma_\omega) = v(\omega) \leq |\text{Sing}(\omega)|, \quad b_1(\Gamma_\omega) = c(\omega).$$

The graph Euler identity $e(\Gamma_\omega) = v(\Gamma_\omega) + b_1(\Gamma_\omega) - 1$ implies:

$$N_{\max}(\omega) = c(\omega) + v(\omega) - 1 \leq c(\omega) + |\text{Sing}(\omega)| - 1,$$

where we noticed that $v(\omega) \leq |\text{Sing}(\omega)|$ since each Δ_s contains at least one singular point. An example of foliation graph is depicted in figure 3.





Figure 3. An example of foliation graph. Regular (a.k.a type I) vertices are represented by black dots, while exceptional (a.k.a. type II) vertices are represented by green blobs. All terminal vertices are regular vertices and correspond to center singularities. Notice that the graph can have multiple edges as well as loops.

Constraints on the foliation graph from the irrationality rank of ω . When the chiral locus \mathcal{W} is empty (i.e. when ω is nowhere-vanishing) we have $\text{Sing}(\omega) = \emptyset$ and $\bar{\mathcal{F}}_\omega = \mathcal{F}_\omega$ is a regular foliation. Even though this doesn't fit our assumption $\text{Sing}\omega \neq \emptyset$, one can define a (degenerate) foliation graph also in this situation (which was considered in [8]). In this case, knowledge of the irrationality rank of \mathfrak{f} determines the topology of the foliation \mathcal{F}_ω for any $\omega \in \mathfrak{f}$. Namely, one has only two possibilities (see figure 4):

- $\rho(\mathfrak{f}) = 1$, i.e. \mathfrak{f} is projectively rational. Then there exists exactly one maximal component (which coincides with M) and no minimal component. The foliation “graph” consists of one loop and has no vertices; M is a fibration over S^1 as a consequence of Tischler’s theorem [62].
- $\rho(\mathfrak{f}) > 1$, i.e. \mathfrak{f} is projectively irrational. There exists exactly one minimal component (which coincides with M) and no maximal component, i.e. \mathcal{F}_ω is a minimal foliation. Then the foliation graph consists of a single exceptional vertex and every leaf of \mathcal{F}_ω is dense in M . As explained in [8], the noncommutative geometry of the leaf space is described by the C^* algebra $C(M/\mathcal{F}_\omega)$ of the foliation, which is a non-commutative torus of dimensions $\rho(\omega)$. Notice that this refined topological information is not reflected by the foliation graph.

The situation is much more complicated when $\text{Sing}(\omega)$ is non-empty, in that knowledge of $\rho(\omega)$ does not suffice to specify the topology of the foliation. In this case, knowledge of $\rho(\mathfrak{f})$ allows one to say only the following:

- When $\rho(\mathfrak{f}) = 1$, then the foliation \mathcal{F}_ω is compactifiable for any $\omega \in \mathfrak{f}$ [18] and the inequality (5.12) below requires $c(\omega) \geq 1$. Hence the foliation graph Γ_ω has only



(a) Foliation graph when $\mathcal{W} = \emptyset$ and $\rho(\omega) = 1$. (b) Foliation graph when $\mathcal{W} = \emptyset$ and $\rho(\omega) > 1$.

Figure 4. Degenerate foliation graphs in the everywhere non-chiral case.

regular vertices and must have at least one cycle. Except for this, nothing else can be said about \mathcal{F}_ω only by knowing that $\rho(\mathfrak{f}) = 1$. Indeed, it was shown in [28] that any compactifiable Morse form foliation $\mathcal{F}_{\omega'}$ with $c(\omega') \geq 1$ can be realized as the foliation defined by a Morse form ω belonging to a projectively rational cohomology class. It was also shown in loc. cit. that such a foliation can in fact be realized by a Morse form of any irrationality rank lying between 1 and $c(\omega')$, inclusively.

- When $\rho(\mathfrak{f}) > 1$, then \mathcal{F}_ω may be either compactifiable or non-compactifiable, hence the foliation graph may or may not have exceptional vertices; when \mathcal{F}_ω is compactifiable, then Γ_ω has no exceptional vertices and has a number of cycles at least equal to $\rho(\omega)$. Criteria for compactifiability of \mathcal{F}_ω can be found in [18, 21, 25] and are given below.

Theorem [18, 21, 25]. The following statements are equivalent:

- (a) \mathcal{F}_ω is compactifiable.
- (b) The period morphism $\text{per}_\mathfrak{f} : \pi_1(M) \rightarrow \mathbb{R}$ factorizes through a group morphism $\pi_1(M) \rightarrow K$, where K is a free group.
- (c) $H_\omega^\perp \subset \ker \omega$.
- (d) $\text{rk} \mathcal{H}_\omega = \rho(\omega)$.

The first criterion above is Proposition 2 in [18, section 8.2]. Since $\mathcal{H}_\omega \subset H_\omega$, we have $\text{rk} \mathcal{H}_\omega \leq \text{rk} H_\omega = c(\omega)$ and the theorem shows that compactifiability of \mathcal{F}_ω requires:

$$\rho(\omega) \leq c(\omega). \quad (5.12)$$

Remark. By its construction, the foliation graph discards topological information about the restriction of the foliation to the minimal components of the Novikov decomposition, which are represented in the graph by exceptional vertices. As in the case $\text{Sing}(\omega) = \emptyset$, the C^* algebra of the foliation should provide more refined information about the topology of $\bar{\mathcal{F}}_\omega$ than the foliation graph. To our knowledge, this C^* algebra has not been computed for foliations given by a Morse 1-form.

The oriented foliation graph. For each maximal component C_j^{max} , the diffeomorphism (5.11) can be chosen¹¹ such that the sign of the integral $\int_{\gamma_j} \omega$ is positive along any

¹¹The sign of $\int_{\gamma_j} \omega$ does not depend on the choice of γ since ω vanishes on the leaves of \mathcal{F}_ω . If the sign is negative, then it can be made positive by composing the diffeomorphism (5.11) with $\text{id}_{L_j} \times R$, where $R \in \text{Diff}_-((0, 1))$ is any orientation-reversing diffeomorphism of the interval $(0, 1)$.

smooth curve $\gamma_j : (0, 1) \rightarrow C_j^{\max}$ which projects to the interval $(0, 1)$. Identifying the corresponding edge e_j with this interval, this gives a canonical orientation \vec{e}_j of e_j which corresponds to “moving along e_j in the direction of increasing value of h_j ”, where h_j is any locally-defined smooth function on an open subset of C_j^{\max} whose exterior derivative equals ω . It follows that the foliation graph Γ_ω admits a canonical orientation, which makes it into the *oriented foliation graph* $\vec{\Gamma}_\omega$.

Weights on the oriented foliation graph. Using the canonical orientation, the integrals:

$$w_j \stackrel{\text{def.}}{=} \int_{\gamma_j} \omega \tag{5.13}$$

(whose value does not depend on the choice of γ_j as above) provide canonical positive weights on $\vec{\Gamma}_\omega$ [18, 57]. These weights can be used [23] to describe the set of Morse 1-forms ω which have the property that $\vec{\mathcal{F}}_\omega = \vec{\mathcal{F}}$ for a fixed singular foliation $\vec{\mathcal{F}}$.

Expression for the weights in terms of \mathbf{b} and \mathbf{f} . In our application, the vector field $n = \hat{V}^\# \in \Gamma(TU)$ is orthogonal to the leaves of \mathcal{F} and satisfies:

$$n \lrcorner \omega = 4\kappa e^{3\Delta} \|V\| = n \lrcorner \mathbf{f} - \partial_n \mathbf{b} \geq 0 \tag{5.14}$$

as a consequence of (3.41). Equality with zero in the right hand side occurs only at the points of $\mathcal{W} = \text{Sing}(\omega)$. It follows that the orientation of the edges of the foliation graph is in the direction of n and that we can take γ_j to be any integral curve ℓ_j of the vector field $n|_{C_j^{\max}}$. Relation (5.14) gives:

$$w_j = \mathbf{b}_j(\gamma_j(1)) - \mathbf{b}_j(\gamma_j(0)) + \int_{\gamma_j} \mathbf{f}.$$

When \mathcal{F}_ω is compactifiable, this relation implies that the sum of weights along all edges of a cycle of the *oriented* foliation graph $\vec{\Gamma}_\omega$ equals the period of \mathbf{f} along the corresponding homology 1-cycle $\alpha \in H_1(M)$ of M :

$$\sum_{\vec{e}_j \text{ in a cycle of } \vec{\Gamma}_\omega} w_j = \int_\alpha \mathbf{f}.$$

5.8 The fundamental group of the leaf space

Even though the quotient topology of the leaf space $M/\vec{\mathcal{F}}_\omega$ can be very poor, one can use the classifying space \mathcal{G} of the holonomy pseudogroup of the regular foliation \mathcal{F}_ω [63] to define the fundamental group of the leaf space through [54]:

$$\pi_1(M/\vec{\mathcal{F}}_\omega) \stackrel{\text{def.}}{=} \pi_1(B\mathcal{G}).$$

Notice that $B\mathcal{G}$ is an Eilenberg-MacLane space of type $K(\pi, 1)$ [63], (i.e. all its homotopy groups vanish except for the fundamental group) since \mathcal{F}_ω is defined by a closed one-form and hence the holonomy groups of its leaves are trivial. One finds [54]:

$$\pi_1(M/\vec{\mathcal{F}}_\omega) = \pi_1(M)/\mathcal{L}_\omega,$$

where \mathcal{L}_ω is the smallest normal subgroup of $\pi_1(M)$ which contains the fundamental group of each leaf of \mathcal{F}_ω . Notice that $M \setminus \text{Sing}(\omega)$ is connected (since M is) and that the inclusion induces an isomorphism $\pi_1(M \setminus \text{Sing}(\omega)) \simeq \pi_1(M)$, since we assume $\dim M \geq 3$ and hence $\text{Sing}\omega$ has codimension at least 3 in M . In particular, the period map of ω can be identified with that of $\omega|_{M \setminus \text{Sing}(\omega)}$. Since ω vanishes along the leaves of \mathcal{F}_ω , this map factors through the projection $\pi_1(M) \rightarrow \pi_1(M/\bar{\mathcal{F}}_\omega)$, inducing a map $\text{per}_0(\omega) : \pi_1(M/\bar{\mathcal{F}}_\omega) \rightarrow \mathbb{R}$.

A minimal component C_a^{\min} is called *weakly complete* [54] if any curve $\gamma \subset C_a^{\min}$ contained in C_a^{\min} and for which $\int_\gamma \omega$ vanishes has its two endpoints on the same leaf of \mathcal{F}_ω ; various equivalent characterizations of weakly complete minimal components can be found in loc. cit. Let:

- $N'_{\min}(\omega)$ denote the number of minimal components which are not weakly complete
- $N''_{\min}(\omega)$ denote the number of minimal components which are weakly complete
- $C_{a_1}^{\min}, \dots, C_{a_k}^{\min}$ (where $1 \leq a_1 < \dots < a_{N''_{\min}(\omega)} \leq N_{\min}(\omega)$) denote those minimal components of the Novikov decomposition which are weakly complete
- $\omega_j \stackrel{\text{def.}}{=} \omega|_{C_{a_j}^{\min}}$ denote the restriction of ω to the weakly complete minimal component $C_{a_j}^{\min}$
- $\Pi_j(\omega) \stackrel{\text{def.}}{=} \Pi(\omega_j)$ denote the period group of ω_j . Then $\Pi_j(\omega)$ is a free Abelian group of rank $\text{rk}\Pi_j(\omega) = \rho(\omega_j) \geq 2$ [54].

With these notations, it was shown in [54] that $\pi_1(M/\bar{\mathcal{F}}_\omega)$ is isomorphic with a free product of free Abelian groups:

$$\pi_1(M/\bar{\mathcal{F}}_\omega) \simeq F_\omega * \Pi_1(\omega) * \dots * \Pi_{N''_{\min}(\omega)}(\omega),$$

where $*$ denotes the free product of groups. Furthermore [22, 54], the free group F_ω factors as:

$$F_\omega \simeq \pi_1(\Gamma_\omega) * \mathbb{Z}^{*K(\omega)},$$

where $\pi_1(\Gamma_\omega) \simeq \mathbb{Z}^{*c(\omega)}$ is the fundamental group of the foliation graph and $K(\omega)$ is a non-negative integer which satisfies $K(\omega) \geq N'_{\min}(\omega)$ and $K(\omega) + c(\omega) + N''_{\min}(\omega) \leq b'_1(M)$. Here, $b'_1(M)$ denotes the first noncommutative Betti number of M [52], whose definition is recalled in appendix E (which also summarizes some further information on the topology of $\bar{\mathcal{F}}$).

5.9 On the relation to compactifications of M-theory on 7-manifolds

One way in which one may attempt to think about our class of compactifications is via a two-step reduction of eleven-dimensional supergravity, as follows:

1. First, reduce eleven-dimensional supergravity along a leaf of the foliation down to a supergravity theory in four dimensions; this would of course be a *gauged* supergravity theory since the restrictions of F and f to a leaf are generally non-trivial.
2. Further reduce the resulting four-dimensional theory down to three dimensions, along the “one-dimensional space” orthogonal to the leaf.

This way of thinking, which corresponds to an attempt at generalizing the well-known, but much simpler case of “generalized Scherk-Schwarz compactifications with a twist” (see, for example, [64]), turns out to be rather naive, for the following reasons:

- In the general case when \mathcal{W} is nonempty and differs from M , there is no such thing as a “typical leaf” of the regular foliation \mathcal{F} of $\mathcal{U} = M \setminus \mathcal{W}$, in the sense that the leaves of this foliation are not all diffeomorphic with each other. As explained above, what happens instead is that the leaves of the restriction of \mathcal{F} to each of the maximal or minimal components of the Novikov decomposition of M are diffeomorphic with each other, which means that for each component of the Novikov decomposition one generally has a distinct diffeomorphism class of leaves. As such, it is unclear which of these seven-manifolds one is supposed to reduce on in step 1 above. Furthermore, the extended foliation $\bar{\mathcal{F}}$ also contains singular leaves, and it is not immediately clear (from a Physics perspective) how to correctly reduce eleven-dimensional supergravity, in the presence of fluxes, on such singular seven-manifolds. One should also note that the leaves of the restriction of \mathcal{F} to a minimal component of the Novikov decomposition are non-compact, so the reduction along such leaves cannot be understood as a Kaluza-Klein reduction in the ordinary sense.
- In general, there is no nice “one-dimensional space” transverse to the leaves. As explained above, the best candidate for such a space is a non-commutative space whose “commutative parts” can be described by the foliation graph, but where some unknown non-commutative pieces have to be pasted in at the exceptional vertices. It is of course already unclear how to correctly reduce a four-dimensional supergravity theory on a graph, let alone on a non-commutative space.

As pointed out in [8, subsection 4.4], many of the issues mentioned above already appear in the much simpler case when ξ is everywhere non-chiral. In that situation, the foliation graph is either a circle (and the Novikov decomposition is reduced to a single maximal component, all leaves being compact and mutually diffeomorphic, being the fibers of a fibration over the circle) or a non-commutative torus of dimension given by the projective irrationality rank of ω (in which case the Novikov decomposition is reduced to a single minimal component, all leaves being non-compact, mutually diffeomorphic and dense in M). Only the first of these two cases has a chance at a meaningful interpretation as a “generalized Scherk-Schwarz compactification with a twist”, where the twist is provided by the Ehresmann connection discussed in [8, appendix E], whose parallel transport generates the defining diffeomorphism ϕ_{a_f} which presents M as a mapping torus in that case (see [8, subsection 4.2]). A proper analysis of that case (which is the simplest of this class of compactifications) is already considerably more subtle than might seem at first sight, for the following reason. As shown in [8, subsection 2.6], the restriction of ξ to a leaf L of \mathcal{F} induces the spinor η_0 of equation (3.39) (see also [8, eq. (2.21)]) which, as shown in loc. cit., is the normalized Majorana spinor (in the seven-dimensional sense) along the seven-manifold L which induces its G_2 structure and which should be used to perform the compactification of eleven-dimensional supergravity on L — a reduction which

would constitute the first step outlined above. Notice, however, that what one needs in our case is not the standard $\mathcal{N} = 1$ compactification of eleven-dimensional supergravity on a 7-manifold with G_2 structure which is usually considered in the literature following [65], since the latter is a compactification down to four-dimensional Minkowski space — while what would be needed in our case would be a compactification down to a space which is related to $\text{AdS}_3 \times S^1$. Also recall from [8, subsection 2.6] that η_0 is a Majorana (a.k.a. real) spinor on L (in the seven-dimensional sense) with respect to a real structure which is dependent of the precise leaf L under consideration and not only of its diffeomorphism class. In particular, the G_2 structure depends on the leaf L (it varies from leaf to leaf) in the complicated manner described by Theorems 1 and 2 of [8] and it is not invariant under the parallel transport of the Ehresmann connection mentioned above, so proper analysis of the second step of the reduction is considerably more involved than what one might expect based on analogy with previous work on Scherk-Schwarz-like constructions.

A conceptually better (and more uniform) way to think of the “relation to seven-dimensional compactifications” (beyond the results of [8] and of this paper, which can be viewed as already providing such a relation since they express very explicitly the geometry of M in terms of the seven-dimensional geometry of the leaves of the foliation) is to consider the “partial decompactification limit” in which the leaf space is “large”. The correct way to formulate this mathematically employs the theory of adiabatic limits of foliations (see, for example, [66]), which, in its most general form, concerns their behavior when the leaf space (understood, in general, as a non-commutative space) is “large” in an appropriate spectral sense. This relates to extending the ordinary adiabatic argument (which lies behind a proper Kaluza-Klein formulation of the idea of “two-step reduction”) to the case of foliations. Though this subject is well-outside the scope of the present paper, we mention that such a way of formulating the problem leads to non-trivial mathematical questions given the fact that the adiabatic limit of foliations is poorly understood for the case of foliations which are not Riemannian, such as those which are of interest in our case (see Remark 3 after Theorem 2 of reference [8]). The adiabatic limit for the general situation when one has to deal with a singular foliation $\bar{\mathcal{F}}$ does not seem to have been investigated in the Mathematics literature.

5.10 A non-commutative description of the leaf space?

Recall from [8] that the leaf space of \mathcal{F} admits a very explicit description as a non-commutative torus in the everywhere non-chiral case (the case $\mathcal{U} = M$, when the foliation graph is reduced either to a circle or to a single exceptional vertex). This leads to the speculation [54] that the topological information which is lost when constructing the exceptional vertices of the foliation graph in the general case could be encoded by some sort of non-commutative geometry, as expected from the fact that such vertices are constructed by collapsing at least one minimal component of the Novikov decomposition to a single point; since the minimal components are foliated by dense leaves, the C^* -algebra of their leaf space must be non-commutative. Unfortunately, it is non-trivial to make this expectation precise, because one also has to take into account the effect of the singular leaves of $\bar{\mathcal{F}}$, so progress on this question would first require giving a proper definition/construction

of the C^* -algebras of singular foliations in the sense of Haefliger, a task which, to our knowledge, has not yet been carried out in the mathematics literature. One may hope that some modification of the construction of [44, 45] (which applies to singular foliations in the sense of Stefan-Sussmann) would lead to a solution of this problem for the case of Haefliger structures, a case which is logically orthogonal to that considered in loc. cit.

6 Conclusions and further directions

We studied $\mathcal{N} = 1$ compactifications of eleven-dimensional supergravity down to AdS_3 in the case when the internal part ξ of the supersymmetry generator is not required to be everywhere non-chiral, but under the assumption that ξ is not chiral everywhere. We showed that, in such cases, the Einstein equations require that the locus \mathcal{W} where ξ becomes chiral must be a set with empty interior and therefore of measure zero. The regular foliation of [8] is replaced in such cases by a singular foliation $\bar{\mathcal{F}}$ (equivalently, by a Haefliger structure on M) which “integrates” a cosmooth singular distribution (generalized bundle) \mathcal{D} on M . The singular leaves of $\bar{\mathcal{F}}$ are precisely those leaves which meet the chiral locus \mathcal{W} , thus acquiring singularities on that locus.

We discussed the topology of such singular foliations in the generic case when ω is a Morse one-form, showing that it is governed by the foliation graph of [16–18]. On the non-chiral locus, we compared the foliation approach of [8] with the $\text{Spin}(7)_\pm$ structure approach of [1], giving explicit formulas for translating between the two methods and showing that they agree. It would be interesting to study what supplementary constraints — if any — may be imposed on the topology of $\bar{\mathcal{F}}$ (and on its foliation graph) by the supersymmetry conditions; this would require, in particular, a generalization of the work of [18, 57].

The singular foliation $\bar{\mathcal{F}}$ is defined by a closed one-form ω whose zero set coincides with the chiral locus. Along the leaves of $\bar{\mathcal{F}}$ and outside the intersection of the latter with \mathcal{W} , the torsion classes are determined by the fluxes [8]. For the singular leaves in the Morse case, this leads to a more complicated version of the problems which were studied in [67, 68] for metrics with G_2 holonomy (the case of torsion-free G_2 structures).

The backgrounds discussed in this paper display a rich interplay between spin geometry, the theory of G-structures, the theory of foliations and the topology of closed one-forms [58]. This suggests numerous problems that could be approached using the methods and results of reference [8] and of this paper — not least of which concerns the generalization to the case of singular foliations of the non-commutative geometric description of the leaf space. In this regard, we note that a complete solution of this problem requires extending the construction of the C^* algebra of regular foliations to the case of singular foliations in the sense of Haefliger — a generalization which would be different from (and, in fact, “orthogonal” to) that performed in [44, 45] for the case of singular foliations in the sense of Stefan-Sussmann. This problem is unsolved already for the case of singular foliations defined by a Morse one-form (the difficulty being in how to deal with the singular leaves). It would be interesting to study quantum corrections to this class of backgrounds, with a view towards clarifying their effect on the geometry of $\bar{\mathcal{F}}$. As mentioned in the introduction,

the class of backgrounds discussed here appears to be connected with the proposals of [6] and [7], connections which deserve to be explored in detail.

One of the reasons why the class of backgrounds studied in this paper may be of wider interest is because, as pointed out in [1], the structure group of M does *not* globally reduce to a proper subgroup of $SO(8)$. This is the origin of the phenomena discussed in this paper, which illustrate the limitations of the theory of classical G-structures as well as of the theory of regular foliations. In its classical form [43], the former does not provide a sufficiently wide conceptual framework for a fully general *global* description of all flux compactifications.

Acknowledgments

The work of C.I.L. is supported by the research grant IBS-R003-G1 while E.M.B. acknowledges support from the strategic grant POSDRU/159/1.5/S/133255, Project ID 133255 (2014), co-financed by the European Social Fund within the Sectorial Operational Program Human Resources Development 2007–2013. This work was also financed by the CNCS-UEFISCDI grants PN-II-ID-PCE 121/2011 and 50/2011 and also by PN 09 37 01 02. The authors thank M. Grana, D. Tsimpis and especially I. Gelbukh for correspondence and suggestions.

A Proof of the topological no-go theorem

Lemma. If $\kappa = 0$, then F and f must vanish and Δ must be constant on M . Furthermore, both ξ^+ and ξ^- must be covariantly constant on M (and hence ξ is also covariantly constant) and b must be constant on M .

Proof. The scalar part of the Einstein equations takes the form [2]:

$$e^{-9\Delta}\square e^{9\Delta} + 72\kappa^2 = \frac{3}{2}\|F\|^2 + 3\|f\|^2.$$

Integrating this by parts on M when $\kappa = 0$, implies¹² that F and f must vanish while Δ must be constant on M . In this case $Q = 0$ and $\mathbb{D} = \nabla^S$ so the supersymmetry conditions (2.4) reduce to the condition that ξ is covariantly constant on M . Then (2.5) implies that each of ξ^+ and ξ^- are covariantly constant and hence b is constant on M while V , Y and Z are covariantly constant since ∇^S is a Clifford connection in the sense of [69]. Notice that both ξ^+ and ξ^- can still be non-vanishing so we can still have $|b| < 1$, in which case V is also non-vanishing and we still have a global reduction of structure group to G_2 on M . ■

Proof of the Theorem. The argument is based on the results of [1]. Let us assume that $\text{Int}\mathcal{W}$ is non-empty. Then at least one of the subsets \mathcal{W}^+ and \mathcal{W}^- has non-empty interior and we can suppose, without loss of generality, that $\text{Int}\mathcal{W}^+ \neq \emptyset$. Let U be an open non-void subset of \mathcal{W}^+ . By the definition of \mathcal{W}^+ , we must have $\xi = \xi^+$ and thus $b = +1$ and $V = 0$

¹²This was first noticed in [2].

at any point of \mathcal{W}^+ and hence of U . Since the one-form L of [1] (which we denote by L_+) is given in terms of V by expression (3.31), it follows that L_+ vanishes at every point of U . The second of equations (3.16) of [1] (notice that we *can* use the differential equations of [1] on the subset U of \mathcal{W}^+ since U is open) shows that the following relation holds on U :

$$e^{-12\Delta} * d * \left(e^{12\Delta} \frac{L_+}{1 + L_+^2} \right) - 4\kappa \frac{1 - L_+^2}{1 + L_+^2} = 0$$

and since $L_+|_U = 0$ this gives $\kappa = 0$. The Lemma now implies that b is constant on M and since the set \mathcal{W}^+ where b equals $+1$ is non-void by assumption, it follows that $b = +1$ on M i.e. that we must have $\mathcal{W}^+ = M$, which is Case 1 in the Theorem. Had we assumed that $\text{Int}\mathcal{W}^-$ were non-empty, we would have concluded in the same way that $\mathcal{W}^- = M$, which is Case 2 in the theorem.

The argument above shows that either Case 1 or Case 2 of the Theorem hold or that both \mathcal{W}^+ and \mathcal{W}^- must have empty interior. If at least one of them is a non-empty set, then we are in Case 4 of the Theorem. If both of them are empty sets, then \mathcal{U} coincides with M by the definition of \mathcal{U} , \mathcal{W}^+ and \mathcal{W}^- and we are in Case 3. In Case 4, the fact that \mathcal{W}^\pm have empty interiors and the fact that they are both closed and disjoint implies immediately that they are both contained in the closure of U and hence so is their union \mathcal{W} . Since M equals $\mathcal{U} \cup \mathcal{W}$, this implies that the closure of \mathcal{U} equals M i.e. that \mathcal{U} is dense in M . By the definition of \mathcal{U} and \mathcal{W} we have $\mathcal{W} = M \setminus \mathcal{U}$ and, since M is the closure of \mathcal{U} , this means that \mathcal{W} is the frontier of \mathcal{U} . ■

B The case $\kappa = 0$

For completeness, we briefly discuss the case $\kappa = 0$ (which corresponds to compactifications down to Minkowski space $\mathbb{R}^{1,2}$).

B.1 When M is compact

In this case, the lemma of appendix A implies that F and f vanish while ξ (thus also V) are covariantly constant on M and hence b (and thus the norm of V) are constant on M ; furthermore, Δ is constant on M . The operator \mathbb{D} of (2.4) reduces to ∇^S while Q reduces to zero. Since ∇^S is a Clifford connection on S while $\nabla\nu = 0$, it follows that ∇^S commutes with $\gamma(\nu)$. Therefore, the supersymmetry equation $\nabla^S\xi = 0$ implies $\nabla^S\xi^\pm = 0$. When ξ is non-chiral, this shows that the space of solutions $\dim\mathcal{K}(\nabla^S, 0)$ must be at least two-dimensional provided that it is non-trivial. In fact, existence of a non-trivial and non-chiral solution of the supersymmetry equations (2.4) is equivalent with existence of two non-trivial chiral solutions of opposite chirality. Due to this fact, the non-chiral case corresponds to $\mathcal{N} = 2$ (rather than $\mathcal{N} = 1$) supersymmetry of the effective 3-dimensional theory. Notice that this phenomenon is specific to the Minkowski case $\kappa = 0$, since the operator \mathbb{D} does not commute with $\gamma(\nu)$ when κ is non-vanishing. We thus distinguish the cases:

1. ξ has definite chirality at some point (and hence at every point) of M , which amounts to $|b| = 1$. Then the metric g of M has holonomy contained either in $\text{Spin}(7)_+$ (when

$\xi = \xi_+$, i.e. $b = +1$ and $\mathcal{W} = \mathcal{W}^+ = M$, $\mathcal{U} = \mathcal{W}^- = 0$, which is Case 1 of the topological no-go theorem of subsection 3.3) or in $\text{Spin}(7)_-$ (when $\xi = \xi^-$, i.e. $b = -1$ and $\mathcal{W} = \mathcal{W}^- = M$, $\mathcal{U} = \mathcal{W}^+ = 0$, which is Case 2 of the topological no-go theorem), while V vanishes identically on M . As a consequence, the distribution $\mathcal{D} = \ker V$ has corank zero and coincides with TM ; the foliation \mathcal{F} becomes a *codimension zero* foliation consisting of a single leaf equal to M . These cases correspond to the classical limit of the well-known compactifications of [3]. The holonomy equals $\text{Spin}(7)_\pm$ iff. M is simply-connected (which is allowed in this case). Notice that this class of Minkowski compactifications *cannot* be viewed as the $\kappa \rightarrow 0$ limit of the compactifications considered in [8] (which correspond to Case 3 of the topological no-go theorem) or in this paper (which correspond to Case 4 of the topological no-go theorem). In particular, one cannot take the limit $\kappa \rightarrow 0$ of the formulas given in Theorems 1, 2, 3 of [8] (which only apply to Case 3 or to the regular foliation \mathcal{F} of the non-chiral set \mathcal{U} of Case 4), since one encounters division by zero.

Remark. One can also consider for example the case when $\mathcal{K}(\nabla^S, 0) \subset \Gamma(M, S^+)$ has dimension 2, which corresponds to the classical limit of the compactifications considered in [70]. In this case, M has holonomy contained in $\text{SU}(4) \subset \text{Spin}(7)_+$, i.e. it is a Calabi-Yau fourfold. One can turn on fluxes by considering the leading quantum correction to the Bianchi identity for \mathbf{G} in such a way as to preserve $\mathcal{N} = 2$ supersymmetry [70] (in which case F must be a primitive (2, 2) form) or $\mathcal{N} = 1$ supersymmetry (in which case F satisfies a weaker constraint, see [71, subsection 3.1]).

2. ξ is nowhere chiral on M , which amounts to $|b| < 1$ and thus $\|V\| \neq 0$. Then $\mathcal{D} = \ker V$ is the (corank one) kernel distribution of a non-trivial covariantly constant one-form. The metric g of M has holonomy contained in G_2 , the G_2 group at every point $p \in M$ being contained in the subgroup of $\text{SO}(T_p M, g_p) \simeq \text{SO}(8)$ consisting of those proper rotations which preserve the one-form $V_p \in T_p^* M$ (rotations which form the group $\text{SO}(\mathcal{D}_p, g_p|_{\mathcal{D}_p}) \simeq \text{SO}(7)$). The holonomy group at p coincides with the intersection of the $\text{Spin}(7)_+$ and $\text{Spin}(7)_-$ subgroups given by the stabilizers of ξ_p^\pm . The regular foliation \mathcal{F} has codimension one, the restriction of the metric to each leaf having holonomy contained in G_2 . As explained above, such compactifications lead to an effective action having $\mathcal{N} = 2$ supersymmetry in 3 dimensions. They can be viewed as the limit $\kappa \rightarrow 0$ of Case 3 of the topological no-go theorem (a case which was studied in [8]), the supersymmetry enhancement arising at the value $\kappa = 0$. In fact, when $\|V\| \neq 0$, one can immediately take the limit $\kappa \rightarrow 0$ in the formulas of Theorems 1, 2 and 3 of loc. cit., since κ appears at most linearly in those expressions (and hence the limit is manifest). Using the fact that $b = 0$ while Δ and $\|V\|$ are constant in the limit of interest here, one immediately checks, for example, that Theorem 1 of [8] reduces to a tautology (since $F = 0$) while Theorem 2 gives $H = A = 0$, which is equivalent with the fact that V is covariantly constant ($\nabla V = 0$) as well as $\vartheta = 0$ (in the notations of loc. cit.), which is equivalent with $D_n \varphi = D_n \psi = 0$ (which is a consequence of the fact that \mathcal{D} is the kernel distribution of a covariantly constant

one-form) and $\tau_0 = \tau_1 = \tau_2 = \tau_3 = 0$, which shows that the G_2 structure has trivial torsion classes, thus corresponding, as expected, to a metric whose restriction to the leaves of \mathcal{F} has G_2 holonomy. Since $H = A = 0$, it follows by Reinhart’s criterion that \mathcal{F} is a Riemannian foliation (the metric g is bundle-like for \mathcal{F}). Notice that such $\mathcal{N} = 2$ Minkowski compactifications are different from the classical limit of the Calabi-Yau fourfold compactifications considered in [70], which instead correspond to the case when the two-dimensional space of solutions $\mathcal{K}(\nabla^S, 0)$ consists of chiral spinors of the same chirality and hence arise as a particular case of 1. above. In the case discussed here $\mathcal{K}(\nabla^S, 0) \subset \Gamma(M, S)$ is spanned by two chiral spinors of *opposite* chirality. As in [70], one could turn on fluxes in such Minkowski compactifications (while preserving $\mathcal{N} = 2$ supersymmetry) by considering the quantum correction to the Bianchi identity of \mathbf{G} which is induced by 5-brane anomaly cancellation, leading to a class of “first order corrected compactifications” which, in our opinion, deserve further study.

For reader’s convenience, we reproduce below some results of [8, Theorem 3] which are relevant for this discussion, where we use arrows to indicate the limit $\kappa = 0$ with constant Δ and constant b (as required by the lemma of appendix A). Solving the supersymmetry conditions, i.e. finding at least one non-trivial solution ξ for (2.4) which is everywhere non-chiral (and which can be taken to be everywhere of norm one) was shown in [8] to give the following constraints:

- On the fluxes $f \in \Omega^1(M)$ and $F \in \Omega^4(M)$:

$$\begin{aligned} f &= 4\kappa V + e^{-3\Delta} d(e^{3\Delta} b) \rightarrow 0, \\ F_{\perp} &= \alpha_1 \wedge \varphi - \hat{h}_{ij} e^i \wedge \iota_{e^j} \psi, \\ F_{\top} &= -\iota_{\alpha_2} \psi + \chi_{ij} e^i \wedge \iota_{e^j} \varphi, \end{aligned}$$

with:

$$\begin{aligned} \alpha_1 &= \frac{1}{2\|V\|} (db)_{\perp} \rightarrow 0, & \alpha_2 &= -\frac{b}{2\|V\|} (db)_{\perp} + \frac{3\|V\|}{2} (d\Delta)_{\perp} \rightarrow 0, \quad (\text{B.1}) \\ \text{tr}_g(\hat{\chi}) &= \kappa - \frac{1}{2\|V\|} (db)_{\top} \rightarrow 0, & \text{tr}_g(\hat{h}) &= 2\kappa b - \frac{3\|V\|}{2} (d\Delta)_{\top} + \frac{b}{2\|V\|} (db)_{\top} \rightarrow 0, \end{aligned}$$

where:

$$\hat{h}_{ij} = h_{ij}^{(0)} + \frac{1}{7} \text{tr}_g(\hat{h}) g_{ij}, \quad \chi_{ij} = \chi_{ij}^{(0)} + \frac{1}{7} \text{tr}_g(\chi) g_{ij}, \quad \text{tr}_g(\chi) = -\frac{4}{3} \text{tr}_g(\hat{\chi}).$$

In the above limit, the fluxes vanish by the no-go theorem and one finds $\chi = \hat{h} = 0$.

- On the quantities H , $\text{tr}A$ and ϑ of the foliation \mathcal{F} :

$$\begin{aligned} H_{\sharp} &= 3(d\Delta)_{\perp} - \frac{b}{\|V\|^2} (db)_{\perp} \rightarrow 0, \\ \text{tr}A &= -\frac{8\kappa b}{\|V\|} + 12(d\Delta)_{\top} - \frac{b(db)_{\top}}{\|V\|^2} \rightarrow 0, \\ \vartheta &= \frac{b}{2} (d\Delta)_{\perp} - \frac{1+b^2}{6\|V\|^2} (db)_{\perp} \rightarrow 0. \quad (\text{B.2}) \end{aligned}$$

- On the torsion classes (notice that τ_3 is not constrained) of the leafwise G_2 structure:

$$\begin{aligned}
 \tau_0 &= \frac{4}{7\|V\|} \left[2\kappa(3+b^2) - \frac{3b}{2}\|V\|(d\Delta)_\top + \frac{1+b^2}{2\|V\|}(db)_\top \right] \rightarrow 0, \\
 \tau_1 &= -\frac{3}{2}(d\Delta)_\perp \rightarrow 0, \quad \tau_2 = 0, \\
 \tau_3 &= \frac{1}{\|V\|} \left(\chi_{ij}^{(0)} - h_{ij}^{(0)} \right) e^i \wedge \iota_{e^j} \varphi \rightarrow 0.
 \end{aligned}
 \tag{B.3}$$

B.2 When M is non-compact

Even though our interest is specifically in the case when M is compact, it may be instructive to consider for the moment also the non-compact case (this is the only place in this paper where we shall do so). Let us assume that M is non-compact but connected and paracompact. In this case, the lemma of appendix A fails to hold (and hence non-vanishing fluxes are allowed) but the first part of the proof of the topological no-go theorem (which is independent of the lemma) still applies, showing that the condition $\kappa \neq 0$ still requires that the closed sets \mathcal{W}^+ and \mathcal{W}^- have empty interior. When $\kappa = 0$, however, any of these sets may acquire interior points. In that case, one has a background given by a warped product of $\mathbb{R}^{1,2}$ with the non-compact Riemannian manifold (M, g) and we have $\mathbf{f} = d\mathbf{b}$ on M . The geometry can be described as follows upon using the chirality decomposition $M = \mathcal{U} \sqcup \mathcal{W}^+ \sqcup \mathcal{W}^-$ (we must of course assume that the warp factor Δ is smooth on M also in the Minkowski limit, in order to have a meaningful physical interpretation in supergravity):

- The open submanifold \mathcal{U} of M can support non-vanishing fluxes $F|_{\mathcal{U}}$ and $f|_{\mathcal{U}}$ and carries a regular codimension one foliation \mathcal{F} (the foliation which integrates the kernel distribution of the one-form $V|_{\mathcal{U}}$) endowed with a longitudinal G_2 structure, whose geometry is determined by the case $\kappa = 0$,¹³ of Theorems 2, 3 of reference [8]. This is the non-compact version of the solutions discussed at point 2 of the previous subsection. Notice that both b and Δ may be non-constant in the non-compact case and hence the limit $\kappa \rightarrow 0$ (which is again trivial to take) is slightly different from that given in the previous subsection.
- Up to a conformal transformation, the restriction $g|_{\text{Int}\mathcal{W}^+}$ has holonomy contained in $\text{Spin}(7)_+$, the type of the solution along $\text{Int}\mathcal{W}^+$ being, locally, the classical limit (the limit when the effect of the tadpole term induced by 5-brane anomaly cancellation in M-theory is neglected) of the non-compact version of the solution considered in [3]; in particular, the restriction $F|_{\mathcal{W}^+}$ is self-dual while the restriction of f to \mathcal{W}^+ is completely determined by the warp factor, as in loc. cit. These conclusions follow either from the computations of [3] (computations which are local in nature and hence

¹³On the locus \mathcal{U} , one can set $\kappa = 0$ directly in all expressions given in [8, Theorems 1, 2, 3] (in particular, in the expressions reproduced above), since $\|V\|$ does not vanish anywhere on \mathcal{U} and since those expressions depend at most linearly on κ . Also notice that the one-form $e^{3\Delta}V$ is closed and that, when κ is nonzero, it has the same kernel distribution as the one-form $\omega = 4\kappa e^{3\Delta}V$. Since ω vanishes when $\kappa = 0$, it must of course be replaced with $e^{3\Delta}V$ when considering the limit $\kappa \rightarrow 0$ of the distribution \mathcal{D} .

apply on $\text{Int}\mathcal{W}^+$) or more directly by setting $\kappa = 0, b = +1, V = L = 0$ in the results of appendix C, which gives $F|_{\mathcal{W}^+} = F^{[27]}|_{\mathcal{W}^+}$ and $\theta_+ = -6d\Delta, T_+ = *(\Phi^+ \wedge d\Delta)$ on this locus (see the last remark in that appendix).

- The restriction of $F|_{\mathcal{W}^-}$ is anti-selfdual while $g|_{\text{Int}\mathcal{W}^-}$ is conformally of holonomy contained in $\text{Spin}(7)_-$, the type of the solution along $\text{Int}\mathcal{W}^-$ being, locally, the classical limit of the non-compact version of the solutions considered in [3], up to a change of orientation of $\text{Int}\mathcal{W}^-$.

Notice that the one-form V vanishes along \mathcal{W}^+ and \mathcal{W}^- but that it does not vanish anywhere on \mathcal{U} . In general, the closures \mathcal{W}^\pm of the open submanifolds $\text{Int}\mathcal{W}^\pm$ need not themselves be manifolds, since the frontiers $\text{Fr}(\mathcal{W}^\pm) = \text{fr}(\mathcal{W}^\pm)$ could be quite “wild”, i.e. quite far from being immersed submanifolds of M . Globally, the geometry of M can be described by saying that M admits¹⁴ a metric-compatible “cosmooth generalized G-structure of type $(G_2, \text{Spin}(7)_+, \text{Spin}(7)_-)$, supported on $(\mathcal{U}, \text{Int}\mathcal{W}^+, \text{Int}\mathcal{W}^-)$ ”, where the $\text{Spin}(7)_\pm$ components are conformally parallel. As in subsection 3.8, one can pack this information into a Haefliger structure, which amounts, geometrically, to adding “singular leaves” to the foliation \mathcal{F} , thus completing it to a singular foliation $\bar{\mathcal{F}}$. Namely, $\bar{\mathcal{F}}$ will contain supplementary leaves of codimension one which meet the frontier $\text{Fr}(\mathcal{W}) = \text{fr}(\mathcal{W})$ (being singular there) as well as supplementary leaves of codimension zero (dimension eight) which are given by the connected components of the open sets $\text{Int}\mathcal{W}^\pm$. The latter are open submanifolds of M whose induced metric has $\text{Spin}(7)_\pm$ holonomy. When the form ω is Morse, the sets \mathcal{W}^+ and \mathcal{W}^- are finite (and hence have empty interior) and the second kind of supplementary leaves do not appear; in this case, one has a codimension one Morse form foliation of the non-compact manifold M , which can again be described using a foliation graph. The geometric description given above could be used, in principle, to attempt a mathematical classification of all non-compact backgrounds given by warped products $\text{AdS}_3 \times_\Delta M$, but such a study lies well outside the scope of the present paper.

Remark. Note that F and f need not be “small” on the locus \mathcal{U} in this class of non-compact Minkowski reductions. The *small flux* approximation (with M non-compact) along the locus $\text{Int}\mathcal{W}^+$ was studied in [1].

C Comparison with the results of [1]

Recall that the positive chirality component ξ^+ of ξ is non-vanishing along the locus \mathcal{U}^+ and hence defines a $\text{Spin}(7)_+$ structure on the open submanifold \mathcal{U} of M . The locus \mathcal{U}^+ was studied in [1] using this $\text{Spin}(7)_+$ structure. In this appendix, we show that the results of [1] agree with those of [8] along the non-chiral locus \mathcal{U} when taking into account the relation between L and V given in subsection 3.6 and the relation between the G_2 and $\text{Spin}(7)_+$ parameterizations of the fluxes given in subsection 4.2. Note that reference [1] uses the notation $\Phi \stackrel{\text{def.}}{=} \Phi^+$ and $L \stackrel{\text{def.}}{=} L^+$. Accordingly, in this appendix we work only

¹⁴The concept of “generalized G-structure” requires some abstract mathematical development, which will be taken up in a different publication.

with the $\text{Spin}(7)_+$ structure and we drop the “+” superscripts and subscripts indicating this structure. Only the major steps of some computations (many of which were performed using code based on the package `Ricci` [72] for `Mathematica`[®]) are given below.

Equations for L (V). Using the relation $L = \frac{1}{1+b}V$, equations [1, (3.16)] take the following form when written in an arbitrary local frame of \mathcal{U} :

$$d(e^{3\Delta}V) = 0, \quad e^{-12\Delta} * d * (e^{12\Delta}V) - 8\kappa b = 0. \quad (\text{C.1})$$

These coincide with the equations discussed in the Remarks after Theorem 3 of [8].

Equations for fluxes in terms of L (V). The first two and last of relations (4.6) take the following coefficient form in the $\text{Spin}(7)_+$ case, being equivalent with equations [1, (C.2)]:

$$\begin{aligned} F_{a_1 a_2 a_3 a_4}^{[1]} &= \frac{1}{42} \Phi_{a_1 a_2 a_3 a_4} \mathcal{F}^{[1]}, \\ F_{a_1 a_2 a_3 a_4}^{[7]} &= \frac{1}{24} \Phi_{[a_1 a_2 a_3}^a \mathcal{F}_{a_4]a}^{[7]}, \\ F_{a_1 a_2 a_3 a_4}^{[35]} &= \frac{1}{6} \Phi_{[a_1 a_2 a_3}^a \mathcal{F}_{a_4]a}^{[35]}. \end{aligned} \quad (\text{C.2})$$

Furthermore, the $\text{Spin}(7)_+$ case of relation (4.7) has the following coefficient form, which is equivalent with [1, (C.1)]:

$$F_{a_1 a_2 a_3 a_4} \Phi^{a_1 a_2 a_3}{}_{a_5} = g_{a_4 a_5} \mathcal{F}^{[1]} + \mathcal{F}_{a_4 a_5}^{[7]} + \mathcal{F}_{a_4 a_5}^{[35]}. \quad (\text{C.3})$$

Reference [1] uses the notations:

$$\begin{aligned} (P^7)_{rs}^{pq} &\stackrel{\text{def.}}{=} \frac{1}{4} \left(\delta_{[r}^p \delta_{s]}^q - \frac{1}{2} \Phi_{rs}{}^{pq} \right), \\ (L \otimes \mathcal{F}^{[7]})_{a_1 a_2 a_3}^{48} &= 6 \left(L_{[a_1} \mathcal{F}_{a_2 a_3]}^{[7]} + \frac{1}{7} \Phi_{a_2 a_2 a_3}{}^b L^a \mathcal{F}_{ab}^{[7]} \right) \iff (L \otimes \mathcal{F}^{[7]})^{48} = 2L \wedge \mathcal{F}^{[7]} - \frac{1}{7} \iota_L \mathcal{F}^{[7]} \Phi, \\ (L \otimes F^{[27]})_{a_1 a_2 a_3}^{48} &\stackrel{\text{def.}}{=} L^a F_{a a_1 a_2 a_3}^{[27]} \quad \text{i.e.} \quad L \otimes F^{[27]} \stackrel{\text{def.}}{=} \iota_L F^{[27]}. \end{aligned} \quad (\text{C.4})$$

Using the relation $L = \frac{1}{1+b}V$ and the identity $\|V\|^2 = 1 - b^2$, one computes, for example:

$$\|L\|^2 = \frac{1-b}{1+b}, \quad 1 + \|L\|^2 = \frac{2}{1+b}, \quad 1 - \|L\|^2 = \frac{2b}{1+b}, \quad \frac{1 - \|L\|^2}{1 + \|L\|^2} = b, \quad \frac{L}{1 + \|L\|^2} = \frac{1}{2}V.$$

Due to such identities, equations [1, (3.17)] take the form:

$$\begin{aligned}
 f &= e^{-3\Delta} d(e^{3\Delta} b) + 4\kappa V, & (C.5) \\
 \frac{1}{12} \mathcal{F}^{[1]} &= \frac{\|V\|}{2(1+b)} e^{-3\Delta} [d(e^{3\Delta}(1+b))]_{\top} - \kappa(1+2b), \\
 \frac{1}{96} \mathcal{F}_{pq}^{[7]} &= -\frac{1}{2(1+b)} e^{-3\Delta} (P^7)_{pq}{}^{rs} V_r \partial_s (e^{3\Delta}(1+b)), \\
 \frac{1}{24} \mathcal{F}_{pq}^{[35]} &= -\frac{\|V\|}{1+b} \nabla_{(p} \hat{V}_{q)} + \frac{1+b^2}{2(1+b)\|V\|} \hat{V}_{(p} \nabla_{q)} b + \frac{3(1-b)\|V\|}{2(1+b)} \hat{V}_{(p} \nabla_{q)} \Delta + \mathbb{T}_{pq} - \\
 &\quad - \frac{1}{14(1+b)} \left[\frac{3(1-b)\|V\|}{1+b} (db)_{\top} + 9(1-b)\|V\|(d\Delta)_{\top} + 8(1-b)(1+2b)\kappa \right] \hat{V}_p \hat{V}_q - \\
 &\quad - \frac{1}{14(1+b)} \left[\frac{(1-b)\|V\|}{2(1+b)} (db)_{\top} + \frac{3}{2} (15-2b)\|V\|(d\Delta)_{\top} - (1+15b-2b^2)\kappa \right] g_{pq},
 \end{aligned}$$

where the quantity \mathbb{T}_{ab} (which appears in the last equation of [1, (3.17)]) can be expressed as:

$$\mathbb{T}_{ab} \stackrel{\text{def.}}{=} -\frac{1}{4} \Phi_{(a}{}^{cde} (L \otimes F^{[27]})_{b)cd} L_e = \frac{1}{4} \Phi_{(a}{}^{cde} L^f F_{b)fd}^{[27]} L_e = \frac{1-b}{2(1+b)} (\iota_{e(a} \Phi) \Delta_3 [(\iota_{e^b} F^{[27]})_{\parallel}]]. \quad (C.6)$$

In an orthonormal local frame with $e_1 = n$, we have:

$$\mathbb{T}_{11} = \mathbb{T}_{1j} = \mathbb{T}_{j1} = 0, \quad \mathbb{T}_{ij} = \frac{1-b}{2(1+b)} (\iota_{e(i} \varphi) \Delta_2^{\perp} (\iota_{e^j} F_{\top}^{[27]})) = -\frac{1-b}{24(1+b)} \mathcal{F}_{ij}^{[27]}.$$

The first equation in (C.5) coincides with a relation given in Theorem 3 of [8]. The second equation in (C.5) can be written as:

$$\mathcal{F}^{[1]} = 12 \left[\frac{3\|V\|}{2} (d\Delta)_{\top} + \frac{\|V\|}{2(1+b)} (db)_{\top} - \kappa(1+2b) \right], \quad (C.7)$$

while the third relation in (C.5) separates as follows into parts parallel and perpendicular to n :

$$\begin{aligned}
 \mathcal{F}_{\top}^{[7]} &= -6\|V\| \left[3(d\Delta)_{\perp} + \frac{(db)_{\perp}}{(1+b)} \right], \\
 \mathcal{F}_{\perp}^{[7]} &= 6\|V\| \left[3\iota_{(d\Delta)_{\perp}} \varphi + \frac{1}{1+b} \iota_{(db)_{\perp}} \varphi \right]. \quad (C.8)
 \end{aligned}$$

In an orthonormal frame as above, we find that the last equation in (C.5) amounts to:

$$\begin{aligned}
 \mathcal{F}_{11}^{[35]} &= 12 \left[-\frac{3}{2} \|V\| (d\Delta)_{\top} - \kappa(1-2b) + \frac{1+b}{2\|V\|} (db)_{\top} \right], \\
 \mathcal{F}_{1i}^{[35]} e^i &= 12 \left[\frac{1+b}{2\|V\|} (db)_{\perp} - \frac{3}{2} \|V\| (d\Delta)_{\perp} \right], \\
 \frac{1}{2} \mathcal{F}_{ij}^{[35]} e^i \odot e^j &= \frac{12}{7} \left[\frac{3}{2} \|V\| (d\Delta)_{\top} - \frac{1+b}{2\|V\|} (db)_{\top} + \kappa(1-2b) \right] g - 12 (h^{(0)} - \chi^{(0)}). \quad (C.9)
 \end{aligned}$$

Substituting the expressions for α_1, α_2 and $\hat{h}, \hat{\chi}$ given in Theorem 3 of [8], it is now easy to check that relations (C.7)–(C.9) are equivalent with:

$$\begin{aligned}
 \mathcal{F}^{[1]} &= -12\text{tr}(\hat{h} + \hat{\chi}), \\
 \mathcal{F}_{\perp}^{[7]} &= -12(\alpha_1 + \alpha_2), \\
 \mathcal{F}_{\perp}^{[7]} &= 12\iota_{(\alpha_1 + \alpha_2)}\varphi, \\
 \mathcal{F}_{11}^{[35]} &= 12\text{tr}(\hat{h} - \hat{\chi}), \\
 \mathcal{F}_{1i}^{[35]}e^i &= 12(\alpha_1 - \alpha_2), \\
 \frac{1}{2}\mathcal{F}_{ij}^{[35]}e^i \odot e^j &= -12(\hat{h} - \hat{\chi}),
 \end{aligned} \tag{C.10}$$

which in turn are equivalent with (4.12) when $\mathcal{F}^{[k]}$ are expressed in the $\text{Spin}(7)_+$ parameterization using (4.8) and (4.9).

Remark. To arrive at equations (C.9), one uses the relations:

$$\hat{V}_{(1;1)} = 0, \quad \hat{V}_{(1;j)} = \frac{1}{2}H_j, \quad \hat{V}_{(i;j)} = -A_{ij}, \tag{C.11}$$

which can be derived by using the local expressions given in appendix C of [8]. Notice that the tensor $\frac{1}{2}V_{(a,b)}e^a \odot e^b = \frac{1}{2}V_{a;b}e^a \odot e^b = \hat{V}_{(a;b)}e^a \otimes e^b$ is the Hessian¹⁵ $\text{Hess}(\hat{V})$ of \hat{V} , where we remind the reader that we use conventions (1.1), which were also used in [8].

Equations for the $\text{Spin}(7)_+$ structure in terms of V and of the fluxes. Reference [1] uses a one-form $\omega^1 \in \Omega^1(M)$ and a three-form $\omega^2 \in \Omega^3(M)$ which are given by [1, eq. (3.18)]:

$$\begin{aligned}
 \omega_m^1 &= \frac{\kappa}{2}L_m + \frac{3}{4}\partial_m\Delta + \frac{1}{168}\left(L_m\mathcal{F}^{[1]} - L^i\mathcal{F}_{im}^{[7]}\right) \Leftrightarrow \omega^1 = \frac{\kappa}{2}L + \frac{3}{4}d\Delta + \frac{1}{168}\left(\mathcal{F}^{[1]}L - \iota_L\mathcal{F}^{[7]}\right), \\
 \omega_{mpq}^2 &= \frac{1}{192}\left(L \otimes \mathcal{F}^{[7]}\right)_{mpq}^{48} + \frac{1}{4}\left(L \otimes F^{[27]}\right)_{mpq}^{48} \Leftrightarrow \omega^2 = \frac{1}{192}\left(2L \wedge \mathcal{F}^{[7]} - \frac{6}{7}\iota_{\iota_L\mathcal{F}^{[7]}}\Phi\right) + \frac{1}{4}\iota_L F^{[27]}.
 \end{aligned} \tag{C.12}$$

These forms satisfy the equation (cf. [1, eq. (3.15)]):

$$\partial_{[m}\Phi_{pqrs]} = -8\Phi_{[mpqr}\omega_{s]}^1 - \frac{4}{15}\varepsilon_{mpqrs}{}^{ijk}\omega_{ijk}^2 \iff d\Phi = -8\Phi \wedge \omega^1 + 8*\omega^2, \tag{C.13}$$

where to arrive at the coordinate-free relation we used the expression:

$$(*\omega^2)_{mpqrs} = -\frac{1}{5!}\varepsilon_{mpqrsabc}(\omega^2)^{abc}.$$

Defining $\theta' \in \Omega^1(M)$ and $T' \in \Omega^3(M)$ through:

$$\omega^1 \stackrel{\text{def.}}{=} -\frac{7}{48}\theta', \quad \omega^2 \stackrel{\text{def.}}{=} -\frac{1}{8}T', \tag{C.14}$$

¹⁵We define the Hessian of an arbitrary one-form $\omega \in \Omega^1(M)$ to be the symmetric part of the tensor $H(\omega) \stackrel{\text{def.}}{=} \nabla\omega \in \Gamma(M, T^*M \otimes T^*M) = \Omega^1(M) \otimes \Omega^1(M)$. Thus $H(\omega)(X, Y) = (\nabla_X\omega)(Y) = X(\omega(Y)) - \omega(\nabla_X Y)$ and $H(\omega)_{ab} = \omega_{b;a} = e_a(\omega_b) - \Omega_{ab}^c\omega_c$ in any (generally non-holonomic) local frame e_a of M , with the connection coefficients Ω_{ab}^c defined through $\nabla_{e_a}e_b = \Omega_{ab}^c e_c$. We have $\text{Hess}(\omega)_{ab} = \omega_{(a;b)}$. When $f \in \mathcal{C}^\infty(M, \mathbb{R})$, the tensor $\text{Hess}(df)$ coincides with the usual Hessian of f .

equations (C.13) take the form:

$$d\Phi = \frac{7}{6}\theta' \wedge \Phi - *T'. \tag{C.15}$$

Relation (4.16) tells us that the Lee-form θ and the characteristic torsion form T of the $\text{Spin}(7)_+$ structure form the particular solution of this inhomogeneous equation which also satisfies condition (4.15). It follows that (θ', T') must differ from (θ, T) through a solution (θ^0, T^0) of the homogeneous equation associated with (C.15), i.e. we must have:

$$\theta' = \theta + \theta^0, \quad T' = T + T^0 \quad \text{with} \quad T^0 = -\frac{7}{6}\iota_{\theta^0}\Phi, \tag{C.16}$$

where $\theta^0 \in \Omega^1(M)$. Using (3.35), we find:

$$T^0_{\perp} = -\frac{7}{6}(\theta^0_{\perp}\varphi + \iota_{\theta^0_{\perp}}\psi), \quad T^0_{\top} = \frac{7}{6}\iota_{\theta^0_{\top}}\varphi \tag{C.17}$$

and hence:

$$\begin{aligned} \omega^1_{\top} &= -\frac{7}{48}(\theta_{\top} + \theta^0_{\top}), & \omega^2_{\top} &= -\frac{1}{8}\left(T_{\perp} + \frac{7}{6}\iota_{\theta^0_{\perp}}\varphi\right) \\ \omega^1_{\perp} &= -\frac{7}{48}(\theta_{\perp} + \theta^0_{\perp}), & \omega^2_{\perp} &= -\frac{1}{8}\left[T_{\top} - \frac{7}{6}(\iota_{\theta^0_{\top}}\psi + \theta^0_{\top}\varphi)\right]. \end{aligned} \tag{C.18}$$

Using the refined $\text{Spin}(7)_+$ parameterization given in table 3 and relations (4.12), equations (C.12) can be seen to be equivalent with:

$$\begin{aligned} \omega^1_{\top} &= \frac{3}{4}(d\Delta)_{\top} + \frac{\kappa\|V\|}{2(1+b)} - \frac{\|V\|}{14(1+b)}\text{tr}_g(\hat{h} + \hat{\chi}), & \omega^2_{\top} &= \frac{\|V\|}{14(1+b)}\iota_{(\alpha_1+\alpha_2)}\varphi, \\ \omega^1_{\perp} &= \frac{3}{4}(d\Delta)_{\perp} + \frac{\|V\|}{14(1+b)}(\alpha_1+\alpha_2), & \omega^2_{\perp} &= \frac{3\|V\|}{56(1+b)}\iota_{(\alpha_1+\alpha_2)}\psi + \frac{\|V\|}{8(1+b)}(h^{(0)}_{ij} + \chi^{(0)}_{ij})e^i \wedge \iota_{e^j}\varphi. \end{aligned} \tag{C.19}$$

Combining (C.18) and (4.18), we find that equations (C.19) agree with the relations given for the torsion classes of the G_2 structure in Theorems 2 and 3 of [8] provided that:

$$\theta^0 = -\frac{1}{7}\theta. \tag{C.20}$$

Conclusion. Combining the results of the paragraph above, we conclude that equations [1, (3.16), (3.17), (3.18)] are *equivalent* on the non-chiral locus with the results of Theorems 2 and 3 of [8]. Furthermore, the results of section 4 and of this appendix provide a complete dictionary which allows one to translate between the language of [8] and that of [1] along the non-chiral locus.

Remark. When M is non-compact and $\kappa = 0$, setting $V = L = 0$ and $b = +1$ in the relations above allows us to determine the nature of the solution along the locus $\text{Int}\mathcal{W}^+$.¹⁶ Doing so in (C.5) and (C.12) and using (C.2) gives $f = 3d\Delta$ and $F^{[1]} = F^{[7]} = F^{[35]} = 0$ (thus $F|_{\mathcal{W}^+} = F^{[27]}|_{\mathcal{W}^+}$ is self-dual) and $\omega^1 = \frac{3}{4}d\Delta$, $\omega^2 = 0$. Relations (C.16) and (C.20) give $\theta' = \frac{6}{7}\theta$, so (C.14) implies $\theta = -6d\Delta$. Relation (C.13) gives $d\Phi = -6\Phi \wedge d\Delta$, hence (4.14) implies $T = *(\Phi \wedge d\Delta)$ on $\text{Int}\mathcal{W}^+$. It follows that the conformally transformed metric $e^{3\Delta}g|_{\text{Int}\mathcal{W}^+}$ has holonomy contained in $\text{Spin}(7)_+$ (the transformation rules of T and θ under a conformal transformation can be found, for example, in [51, Proposition 4.1]).

¹⁶Some of the relations obtained extend to \mathcal{W}^+ by continuity.

D Generalized bundles and generalized distributions

Let M be a connected and paracompact Hausdorff manifold. Recall that a *generalized subbundle* F of a vector bundle E on M is simply a choice of a subspace of each fiber of that bundle. A (*local*) *section* of F is a (local) section s of E such that $s(p) \in F_p$ for any point p lying in the domain of definition of s ; such a section is called *smooth* when it is smooth as a section of the bundle E . The set of smooth sections of E over any open subset U of M forms a module over $\mathcal{C}^\infty(U, \mathbb{R})$ which we denote by $\mathcal{C}^\infty(U, F)$. The modules $\mathcal{C}^\infty(U, F)$ need not be finitely generated; furthermore the module $\mathcal{C}^\infty(M, F)$ of global smooth sections of F need not be projective or finitely generated.¹⁷ We say that F is *algebraically locally finitely generated* if every point of M has an open neighborhood U such that $\mathcal{C}^\infty(U, F)$ is finitely generated as a $\mathcal{C}^\infty(U, \mathbb{R})$ -module. A generalized subbundle of E is called *regular* if it is an ordinary smooth subbundle of E . Some references for the theory of generalized subbundles are [11, 12].

The *rank* of a generalized sub-bundle F is the map $\text{rk}F : M \rightarrow \mathbb{N}$ which associates to each point of M the dimension of the fiber of F at that point. The *corank* of F is the function $\text{corank}F \stackrel{\text{def.}}{=} \dim M - \text{rk}F : M \rightarrow \mathbb{N}$. A point $p \in M$ is called a *regular point* for F if the rank function is locally constant at p . The *regular set* of F is the open subset of M consisting of all regular points, while its closed complement is the *singular set* of F ; this is the set of points where the rank of the fiber of F ‘jumps’. Notice that F is regular iff all points of M are regular for F , i.e. (since M is connected) iff the rank function of F is constant on M .

F is called *smooth* if its fiber at any point p of M is generated as a vector space by the values at p of some finite collection of smooth local sections of E (equivalently, if any point of F_p is the value at p of a smooth local section of E). It is called *cosmooth* if, for all $p \in M$, the fiber F_p can be presented as the intersection of the kernels of the values at p of the elements of a finite collection of smooth local sections of the bundle E^* dual to E ; this amounts to the condition that F is the polar of a smooth generalized subbundle of G of E^* , i.e. that each of its fibers F_p coincides with the subspace of E_p where all linear functionals from $G_p \subset E_p^* = \text{Hom}_{\mathbb{R}}(E_p, \mathbb{R})$ vanish. It is easy to see that the rank of a smooth generalized bundle is a lower semicontinuous function, while the rank of a cosmooth generalized subbundle is upper semicontinuous. As a consequence, the set of regular points of a generalized subbundle F is open and dense in M (hence the singular set is nowhere dense) when F is either smooth or cosmooth. Also notice that F is both smooth and cosmooth iff its rank function is constant on M i.e. iff F is regular.

It was shown in [11] that a generalized subbundle F of E is smooth iff there exists a finite collection $s_1 \dots s_N$ of smooth *global* sections of E such that F_p is the linear span of $s_1(p), \dots, s_N(p)$ for all $p \in M$; furthermore, the number N of sections needed to generate all fibers of F is bounded from above by $(1 + \dim M)\text{rk}E$. Hence any generalized subbundle of E is *pointwise* globally finitely-generated in this manner.¹⁸

¹⁷When F is an ordinary subbundle of E , the module of global sections is finitely generated and projective since we assume M to be connected, Hausdorff and paracompact.

¹⁸This, of course, does *not* imply that it is globally or locally *algebraically* finitely generated. See [11] for a counter-example.

A generalized subbundle of TM is called a *singular (or generalized) distribution* on M while a generalized subbundle of T^*M is called a *singular (or generalized) codistribution* on M . Notice that a regular generalized (co)distribution is the same as a Frobenius (co)distribution (a subbundle of the (co)tangent bundle).

Remark. Given a smooth generalized codistribution which is algebraically locally finitely generated, its polar need not be algebraically locally finitely generated. To see this, consider the following:

Example. Let $M = \mathbb{R}$ and take the smooth generalized codistribution generated by the one-form $V = f(x)dx$, where $f \in C^\infty(\mathbb{R}, \mathbb{R})$ is a smooth function which is everywhere non-vanishing outside the interval $[0, 1]$ and vanishing on $[0, 1]$. The dual \mathcal{D} of this codistribution has rank one on the interval $[0, 1]$ and rank zero on its complement. For $p = 0 \in [0, 1]$ and I any open interval containing p , the space $C^\infty(I, \mathcal{D}) \subset C^\infty(I, \mathbb{R})$ consists of all functions $h \in C^\infty(I, \mathbb{R})$ whose open support $\text{supp}(h) \stackrel{\text{def.}}{=} \{x \in \mathbb{R} | h(x) \neq 0\}$ is contained in the open interval $I_+ \stackrel{\text{def.}}{=} I \cap (0, +\infty)$. Such functions form an ideal of $C^\infty(I, \mathbb{R})$ which is not finitely generated.

A generalized distribution $\mathcal{D} \subset TM$ with polar generalized codistribution $\mathcal{D}^\circ \subset T^*M$ is called:

- Cartan integrable at a point $p \in M$ if there exists an immersed submanifold N of M , passing through p , such that $T_p N = \mathcal{D}_p$
- Cartan integrable, if it is Cartan integrable at every point of M
- Pfaff integrable, if the $C^\infty(M, \mathbb{R})$ -module of global smooth sections $C^\infty(M, \mathcal{D}^\circ) \subset \Omega^1(M)$ is globally generated by a finite number of exact forms (in particular, this requires that \mathcal{D}° is globally algebraically finitely generated). It is easy to see that Pfaff integrability implies that $C^\infty(M, \mathcal{D}^\circ)$ is a differential ideal of the (graded-commutative) differential graded ring $(\Omega(M), d, \wedge)$. This in turn implies (but generally is not equivalent with) Pfaff's condition, which states that any finite set $\omega_1, \dots, \omega_N$ of generators of $C^\infty(M, \mathcal{D}^\circ)$ over $C^\infty(M, \mathbb{R})$ has the property that $d\omega \wedge \omega_1 \wedge \dots \wedge \omega_N = 0$ for all $\omega \in C^\infty(M, \mathcal{D}^\circ)$.

Cartan integrability and Pfaff integrability are logically independent conditions when \mathcal{D} is not regular, i.e. there exist Pfaff integrable generalized distributions which are not Cartan integrable and Cartan integrable generalized distributions which are not Pfaff integrable. Furthermore, Pfaff's condition is no longer equivalent with Pfaff integrability, unlike the case when \mathcal{D} is regular. Conditions for Cartan integrability of cosmooth generalized distributions were given in [73].

Almost all cosmooth generalized distributions arising in practice fail to be globally Cartan integrable. Due to this fact, one usually adopts the following definition. A *leaf* of a cosmooth distribution \mathcal{D} is a maximal connected subset \mathcal{L} of M with the property that any two points p, q of \mathcal{L} can be connected by a smooth curve $\gamma : [0, 1] \rightarrow M$ ($\gamma(0) = p, \gamma(1) = q$) such that the tangent vector of γ at each $t \in (0, 1)$ lies inside the subspace $\mathcal{D}_{\gamma(t)}$. With this

definition, the leaves can be singular (i.e. they need not be immersed submanifolds of M) and Cartan integrability at a point insures existence of a leaf through that point which is locally an immersed submanifold of dimension equal to $\dim \mathcal{D}_p$. When \mathcal{D} fails to be Cartan integrable at p , the leaf through p is singular at p .

Remark. Our terminology agrees with that of [12] but differs from the notion used by other authors. For example:

- A *Stefan-Sussmann distribution* (i.e. a singular distribution in the sense of [9] and [10]) is what we call a *smooth* singular distribution. For such singular distributions Stefan and Sussmann proved a generalization of the Frobenius integrability theorem (see [13] and [14] for textbook treatments).
- What the authors of [44, 45] call singular distribution is what we call an algebraically locally finitely generated smooth distribution. For such singular distributions, the Stefan-Sussmann integrability theorem states (similarly to the Frobenius theorem) that \mathcal{D} is integrable iff it is locally involutive with respect to the Poisson bracket.¹⁹
- The integrability conditions for a non-regular cosmooth distribution (equivalently, for a non-regular smooth codistribution) are much more complicated [73] than those given by Stefan and Sussmann for smooth distributions.

The cosmooth singular distribution defined by V . Consider the codistribution $\mathcal{V} \subset T^*M$ on M which is generated at every point by V , i.e. $\mathcal{V}_p = \mathbb{R}V_p \subset T_p^*M$. This distribution is smooth (since V is) as well as globally algebraically finitely generated by the single smooth section V of T^*M . Let $\mathcal{D} \subset TM$ be the polar of this codistribution. Thus \mathcal{D} is the generalized subbundle of TM defined by associating to a point p of M the kernel of the one-form V_p (which coincides with the orthogonal complement in T_pM of the dual vector $n_p = V_p^\sharp$ at that point). It follows that \mathcal{D} is *cosmooth* (as the polar of a smooth codistribution) but that it need not be algebraically locally finitely generated (see the example above). Notice that \mathcal{D} is smooth iff it is a regular Frobenius distribution, which happens only when V is everywhere non-vanishing, i.e. when the Majorana spinor ξ is everywhere non-chiral. The fiber $\mathcal{D}_p = \ker V_p \subset T_pM$ of \mathcal{D} at a point $p \in M$ has rank seven when $V_p \neq 0$ and rank eight when $V_p = 0$. Since \mathcal{D} is cosmooth, its rank function $\text{rk}\mathcal{D} = 8 - \text{rk}\mathcal{V} : M \rightarrow \mathbb{N}$ is upper semicontinuous; its value at p equals 7 when $V_p \neq 0$ and equals 8 otherwise. Assuming that we are in Case 4 of the topological no-go theorem of subsection 3.3, it follows that $\text{corank}\mathcal{D}$ equals 1 on the non-chiral locus \mathcal{U} and vanishes on the chiral locus \mathcal{W} . The set of regular points of \mathcal{D} coincides with \mathcal{U} .

¹⁹For singular smooth distributions which are not algebraically finitely generated the integrability condition is more complicated — see [9, 10, 13, 14].

E Some topological properties of singular foliations defined by a Morse one-form

E.1 Some topological invariants of M

Let $b'_1(M)$ denote the *first noncommutative Betti number* [52] of M , i.e. the maximum rank of a quotient group of $\pi_1(M)$ which is a free group.²⁰ Let $\mathfrak{H}(M)$ denote the largest rank of a subgroup of $H^1(M, \mathbb{Z})$ on which the cup product vanishes identically. It was shown in [22] that $b'_1(M) \leq \mathfrak{H}(M)$. Moreover, $\mathfrak{H}(M)$ has the following properties which are useful in computations [19, 74]:

1. $\mathfrak{H}(M_1 \times M_2) = \max(\mathfrak{H}(M_1), \mathfrak{H}(M_2))$.
2. $\mathfrak{H}(M_1 \# M_2) = \mathfrak{H}(M_1) + \mathfrak{H}(M_2)$ for $\dim M_i \geq 2$, where $\#$ denotes the connected sum.
3. Let $r = \text{rk}(\ker \cup)$, where \cup is the cup product on $H^1(M, \mathbb{Z})$. Then:

$$\frac{b_1(M) + b_2(M)r}{b_2(M) + 1} \leq \mathfrak{H}(M) \leq \frac{b_1(M)b_2(M) + r}{b_2(M) + 1}.$$

Since $r \leq b_1(M)$, this gives $\mathfrak{H}(M) \leq b_1(M)$.

4. One has $\mathfrak{H}(T^n) = 1$ and $\mathfrak{H}(M_g^2) = g$ where T^n is the n -torus and M_g^2 an orientable closed surface of genus g .

Combining the inequalities above gives:

$$b'_1(M) \leq \mathfrak{H}(M) \leq b_1(M).$$

Notice that $H_{n-1}(M, \mathbb{Z})$ is torsion free since it is isomorphic to $H^1(M, \mathbb{Z}) \simeq \text{Hom}(\pi_1(M, \mathbb{Z}), \mathbb{Z})$ by Poincaré duality — since M is a manifold, both groups are finitely generated and thus free Abelian. If $A \subset H_{n-1}(M, \mathbb{Z})$ is any subgroup, we let $A^\perp \subset H_1^{\text{tf}}(M, \mathbb{Z})$ denote the polar of A with respect to the intersection pairing $(,) : H_1^{\text{tf}}(M, \mathbb{Z}) \times H_{n-1}(M, \mathbb{Z}) \rightarrow \mathbb{Z}$ (which is a perfect pairing).

E.2 Estimate for the number of splitting saddle points

Define:

$$D(\omega) \stackrel{\text{def.}}{=} 1 + \frac{|\Sigma_1^{\text{sp}}(\omega)| - |\Sigma_0(\omega)|}{2} \in \frac{1}{2}\mathbb{Z},$$

where the numbers appearing in the right hand side were defined in section 5. It was shown in [27] that $D(\omega) \geq 0$, equality being attained iff ω is exact. When ω is not exact, one further has $D(\omega) \geq 1$, i.e. $D(\omega)$ can never take the value $\frac{1}{2}$. All greater integer and half-integer values can be realized for some Morse form ω belonging to any given nontrivial cohomology class $\mathfrak{f} \in H^1(M, \mathbb{R}) \setminus \{0\}$.

²⁰Such quotient groups are allowed to be non-Abelian.

E.3 Estimates for c and N_{\min}

It was shown in [22] that:

$$c(\omega) + N_{\min}(\omega) \leq b'_1(M) \tag{E.1}$$

and that every value of $c(\omega)$ between zero and $b'_1(M)$ is attained by some ω which is generic and which has compactifiable foliation \mathcal{F}_ω (i.e. which has $N_{\min}(\omega) = 0$). This inequality implies the non-exact estimate $c(\omega) + N_{\min}(\omega) \leq \mathfrak{H}(M)$ of [28]. The latter reference also gives the following estimate which is independent from (E.1):

$$c(\omega) + 2N_{\min}(\omega) \leq b_1(M). \tag{E.2}$$

Finally, the following inequality holds [27]:

$$c(\omega) + N_{\min}(\omega) \leq D(\omega). \tag{E.3}$$

This implies an older estimate of [74]. Notice that $D(\omega)$ can be smaller, equal to or larger than $b'_1(\omega)$ so (E.3) is independent of (E.1) unless one has more information about the form ω .

E.4 Criteria for existence and number of homologically independent compact leaves

Theorem [25]. The following statements are equivalent:

- (a) \mathcal{F}_ω has at least one compact leaf L
- (b) There exists a smooth non-constant function $h \in C^\infty(M, \mathbb{R})$ (which need not be Morse!) such that $\omega \sim dh$
- (c) There exists a closed one-form α (which need not be Morse!) such that $\alpha \wedge \omega = 0$, α has integer periods (i.e. $[\alpha] \in H^1(M, \mathbb{Z})$) and α is not identically zero. Moreover, L can be chosen with $[L] \neq 0$ in $H_{n-1}(M, \mathbb{Z})$ iff α can be chosen with $[\alpha] \neq 0$ in $H^1(M, \mathbb{R})$.

Theorem [25]. The following statements are equivalent:

- (a) \mathcal{F}_ω has c homologically independent compact leaves
- (b) There exist c cohomologically independent (over \mathbb{R}) closed one-forms α_i with *integer* periods, each of which satisfies $\alpha_i \wedge \omega = 0$.

E.5 Generic forms

Recall that the Morse form ω is called generic if each singular leaf of \mathcal{F}_ω contains exactly one singular point. Some special properties of such Morse forms are summarized in the following:

Proposition [27]. Let ω be a generic Morse one-form. Then:

1. $D(\omega)$ is an integer and satisfies $D(\omega) \leq b'_1(M)$. Furthermore, any value between 0 and $b'_1(M)$ can be realized on M by some generic Morse 1-form ω .
2. All regular (a.k.a. type I) vertices of Γ_ω have degree at most 3 while each exceptional (a.k.a. type II) vertex contains exactly one minimal component.
3. If each of the minimal components of ω is weakly complete, then equality holds in (E.3).

E.6 Exact forms

Let the Morse one-form ω be exact, thus $\rho(\omega) = 0$. In this particular case, we have $\omega = dh$ for some globally-defined Morse function $h \in C^\infty(M, \mathbb{R})$. Since M is compact and connected, h attains its maximum and minimum on M and the image $h(M) \subset \mathbb{R}$ is a closed interval $[a_1, a_N]$, where $a_1 < \dots < a_N$ are the critical values of h . We have $\text{Sing}(\omega) = \cup_{j=1}^N S_j$, where $S_j \stackrel{\text{def.}}{=} \text{Sing}(\omega) \cap h^{-1}(a_j)$ is the set of those critical points of h having critical value a_j . The leaves of the singular foliation $\bar{\mathcal{F}}_\omega$ are the connected components of the level sets $h^{-1}(\{x\})$, where $x \in [a_1, a_N]$. The singular leaves are those connected components of $h^{-1}(a_j)$ which contain at least one point of S_j . Hence the foliation \mathcal{F}_ω is compactifiable and its foliation graph projects onto the chain graph which has a_j as its vertices. The singular points belonging to S_1 and S_N are centers, while the remaining critical points are saddle points. The geometry of such foliations is a classical subject in Morse theory [59–61]. In this case, the form ω is generic iff h is generic in the sense of Morse theory, i.e. iff $|S_j| = 1$ for all $j = 1, \dots, N$. In this case, M can be constructed by successively attaching handles starting from the ball $h^{-1}([0, a_1])$.

E.7 Behavior under exact perturbations

Fix $f \in H^1(M)$ and let $\Omega(f) \stackrel{\text{def.}}{=} \{\omega \in \Omega(M) | d\omega = 0 \text{ and } \omega \in f\}$ be endowed with the C^∞ topology. Define:

- $\Omega_{\mathcal{M}}(f) \stackrel{\text{def.}}{=} \{\omega \in \Omega(f) | \omega \text{ is Morse}\}$
- $\Omega_{\mathcal{K}}(f) \stackrel{\text{def.}}{=} \{\omega \in \Omega_{\mathcal{M}}(f) | \mathcal{F}_\omega \text{ has at least one compact leaf}\}$
- $\Omega_{\text{cf}}(f) \stackrel{\text{def.}}{=} \{\omega \in \Omega_{\mathcal{M}}(f) | \mathcal{F}_\omega \text{ is compactifiable}\}$
- $\Omega_{\text{gen}}(f) \stackrel{\text{def.}}{=} \{\omega \in \Omega_{\mathcal{M}}(f) | \mathcal{F}_\omega \text{ is generic}\}$

Theorem [26]. We have:

1. $\Omega_{\mathcal{M}}(f)$ is open and dense in $\Omega(f)$ while $\Omega_{\text{gen}}(f)$ is dense (but not necessarily open) in $\Omega(f)$ (and hence also in $\Omega_{\mathcal{M}}(f)$).
2. $\Omega_{\mathcal{K}}(f)$ and $\Omega_{\text{cf}}(f)$ are open in $\Omega(f)$
3. $\Omega_{\text{cf}}(f) \cap \Omega_{\text{gen}}(f)$ is open in $\Omega(f)$
4. The restriction of the function c (which counts the number of homologically independent compact leaves) to $\Omega_{\mathcal{K}}(f)$ is lower semicontinuous.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] D. Tsimpis, *M-theory on eight-manifolds revisited: $N = 1$ supersymmetry and generalized $spin(7)$ structures*, *JHEP* **04** (2006) 027 [[hep-th/0511047](#)] [[INSPIRE](#)].
- [2] D. Martelli and J. Sparks, *G structures, fluxes and calibrations in M-theory*, *Phys. Rev. D* **68** (2003) 085014 [[hep-th/0306225](#)] [[INSPIRE](#)].
- [3] K. Becker, *A note on compactifications on $spin(7)$ -holonomy manifolds*, *JHEP* **05** (2001) 003 [[hep-th/0011114](#)] [[INSPIRE](#)].
- [4] M. Becker, D. Constantin, S.J. Gates Jr., W.D. Linch III, W. Merrell and J. Phillips, *M theory on $spin(7)$ manifolds, fluxes and 3-D, $N = 1$ supergravity*, *Nucl. Phys. B* **683** (2004) 67 [[hep-th/0312040](#)] [[INSPIRE](#)].
- [5] D. Constantin, *Flux compactification of M-theory on compact manifolds with $spin(7)$ holonomy*, *Fortsch. Phys.* **53** (2005) 1272 [[hep-th/0507104](#)] [[INSPIRE](#)].
- [6] M. Graña, C.S. Shahbazi and M. Zambon, *$Spin(7)$ -manifolds in compactifications to four dimensions*, *JHEP* **11** (2014) 046 [[arXiv:1405.3698](#)] [[INSPIRE](#)].
- [7] F. Bonetti, T.W. Grimm and T.G. Pugh, *Non-Supersymmetric F-theory Compactifications on $Spin(7)$ Manifolds*, *JHEP* **01** (2014) 112 [[arXiv:1307.5858](#)] [[INSPIRE](#)].
- [8] E.M. Babalic and C.I. Lazaroiu, *Foliated eight-manifolds for M-theory compactification*, *JHEP* **01** (2015) 140 [[arXiv:1411.3148](#)] [[INSPIRE](#)].
- [9] P. Stefan, *Accessible sets, orbits, and foliations with singularities*, *Proc. London Math. Soc.* **29** (1974) 699.
- [10] H.J. Sussmann, *Orbits of families of vector fields and integrability of distributions*, *Trans. Am. Math. Soc.* **180** (1973) 171.
- [11] L.D. Drager, J.M. Lee, E.Park and K. Richardson, *Smooth distributions are finitely generated*, *Ann. Global Anal. Geom.* **41** (2012) 357. [[arXiv:1012.5641](#)].
- [12] F. Bullo and A.D. Lewis, *Geometric Control of Mechanical Systems*, *Texts Appl. Math.* **49** (2004) 1.
- [13] P.W. Michor, *Topics in Differential Geometry*, *Grad. Stud. Math.* **93** (2008) 1.
- [14] J.E. Marsden and T.S. Ratiu, *Introduction to Mechanics and Symmetry*, *Texts Appl. Math.* **17** (1999) 1, internet supplement.
- [15] A. Haefliger, *Homotopy and integrability*, in *Manifolds*, *Lect. Notes Math.* **197** (1971) 133.
- [16] I.A. Mel'nikova, *Compact foliations of Morse forms*, Ph.D. Thesis, Moscow State University, Moscow, Russia (1995).
- [17] I.A. Mel'nikova, *Singular points of a Morsian form and foliations*, *Moscow Univ. Math. Bull.* **51** (1996) 33.
- [18] M. Farber, G. Katz and J. Levine, *Morse theory of harmonic forms*, *Topology* **37** (1998) 469 [[dg-ga/9711007](#)].

- [19] I.A. Mel'nikova, *A test for non-compactness of the foliation of a Morse form*, *Russ. Math. Surv.* **50** (1995) 444.
- [20] I.A. Mel'nikova, *An indicator of the noncompactness of a foliation on M_g^2* , *Math. Notes* **53** (1993) 356.
- [21] I. Gelbukh, *Presence of minimal components in a Morse form foliation*, *Diff. Geom. Appl.* **22** (2005) 189.
- [22] I. Gelbukh, *Number of minimal components and homologically independent compact leaves for a Morse form foliation*, *Studia Sci. Math. Hungar.* **46** (2009) 547.
- [23] I. Gelbukh, *Ranks of collinear Morse forms*, *J. Geom. Phys.* **61** (2011) 425.
- [24] I. Gelbukh, *On collinear closed one-forms*, *Bull. Aust. Math. Soc.* **84** (2011) 322.
- [25] I. Gelbukh, *On compact leaves of a Morse form foliation*, *Publ. Math. Debrecen* **78** (2011) 37.
- [26] I. Gelbukh, *Close cohomologous Morse forms with compact leaves*, *Czech. Math. J.* **63** (2013) 515.
- [27] I. Gelbukh, *The number of split points of a Morse form and the structure of its foliation*, *Math. Slovaca* **63** (2013) 331.
- [28] I. Gelbukh, *On the structure of a Morse form foliation*, *Czech. Math. J.* **59** (2009) 207.
- [29] I. Gelbukh, *The number of minimal components and homologically independent compact leaves of a weakly generic Morse form on a closed surface*, *Rocky Mountains J. Math.* **43** (2013) 1537.
- [30] A. Connes, *A survey of foliations and operator algebras*, in *Operator algebras and applications, Part 1, Proc. Sympos. Pure Math.* **38** (1982) 521.
- [31] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego, CA, U.S.A (1994).
- [32] C.I. Lazaroiu, E.M. Babalic and I.A. Coman, *Geometric algebra techniques in flux compactifications (I)*, [arXiv:1212.6766](#) [INSPIRE].
- [33] C.I. Lazaroiu and E.M. Babalic, *Geometric algebra techniques in flux compactifications (II)*, *JHEP* **06** (2013) 054 [[arXiv:1212.6918](#)] [INSPIRE].
- [34] C.I. Lazaroiu, E.M. Babalic and I.A. Coman, *The geometric algebra of Fierz identities in arbitrary dimensions and signatures*, *JHEP* **09** (2013) 156 [[arXiv:1304.4403](#)] [INSPIRE].
- [35] K. Tod, *All Metrics Admitting Supercovariantly Constant Spinors*, *Phys. Lett. B* **121** (1983) 241 [INSPIRE].
- [36] M. Spivak, *A comprehensive introduction to differential geometry*, Publish or Perish, (1999).
- [37] E. Cremmer, B. Julia and J. Scherk, *Supergravity Theory in Eleven-Dimensions*, *Phys. Lett. B* **76** (1978) 409 [INSPIRE].
- [38] D.V. Alekseevsky and V. Cortes, *Classification of N -(super)-extended Poincare algebras and bilinear invariants of the spinor representation of $\text{Spin}(p, q)$* , *Commun. Math. Phys.* **183** (1997) 477. [[math/9511215](#)].
- [39] D.V. Alekseevsky, V. Cortes, C. Devchand and A.V. Proyen, *Polyvector Super-Poincaré Algebras*, *Commun. Math. Phys.* **253** (2005) 385.
- [40] D. Joyce, *Compact manifolds with special holonomy*, Oxford university press, Oxford, U.K. (2000).

- [41] F. Witt, *Generalised G_2 manifolds*, *Commun. Math. Phys.* **265** (2006) 275 [[math/0411642](#)] [[INSPIRE](#)].
- [42] F. Witt, *Special metric structures and closed forms*, Ph.D. Thesis, University of Oxford, Oxford, U.K. (2004) [math/0502443](#) [[INSPIRE](#)].
- [43] S.S. Chern, *The geometry of G structures*, *Bull. Am. Math. Soc.* **72** (1966) 167.
- [44] I. Androulidakis and G. Skandalis, *The holonomy groupoid of a singular foliation*, *J. Reine Angew. Math.* **626** (2009) 1 [[math/0612370](#)].
- [45] I. Androulidakis and M. Zambon, *Holonomy transformations for singular foliations*, *Adv. Math.* **256** (2014) 348 [[arXiv:1205.6008](#)].
- [46] A. Gray, *Vector cross products on manifolds*, *Trans. Am. Math. Soc.* **141** (1969) 465 [*Erratum ibid.* **148** (1970) 625].
- [47] M. Cabrera, *Orientable Hypersurfaces of Riemannian Manifolds with Spin(7)-Structure*, *Acta Math. Hung.* **76** (1997) 235.
- [48] M. Fernandez, *A classification of Riemannian manifolds with structure group Spin(7)*, *Ann. Mat. Pura Appl.* **143** (1986) 101.
- [49] S. Karigiannis, *Deformations of G_2 and Spin(7) Structures on Manifolds*, *Can. J. Math.* **57** (2005) 1012 [[math/0301218](#)].
- [50] S. Karigiannis, *Flows of G_2 -structures, I*, *Q. J. Math.* **60** (2009) 487 [[math.DG/0702077](#)].
- [51] S. Ivanov, *Connections with torsion, parallel spinors and geometry of Spin(7)-manifolds*, *Math. Res. Lett.* **11** (2004) 171. [[math/0111216](#)] [[INSPIRE](#)].
- [52] P. Arnoux and G. Levitt, *Sur l'unique ergodicité des 1-formes fermées singulières*, *Invent. Math.* **84** (1986) 141.
- [53] G. Levitt, *1-formes fermées singulières et groupe fondamental*, *Invent. Math.* **88** (1987) 635.
- [54] G. Levitt, *Groupe fondamental de l'espace des feuilles dans les feuilletages sans holonomie*, *J. Diff. Geom.* **31** (1990) 711.
- [55] H. Imanishi, *On codimension one foliations defined by closed one forms with singularities*, *J. Math. Kyoto Univ.* **19** (1979) 285.
- [56] H. Imanishi, *Structure of codimension 1 foliations without holonomy on manifolds with abelian fundamental group*, *J. Math. Kyoto Univ.* **19** (1979) 481.
- [57] K. Honda, *A note on Morse theory of harmonic one forms*, *Topology* **38** (1999) 223.
- [58] M. Farber, *Topology of Closed One-Forms*, *Math. Surv. Mon.* **108** (2004) 1.
- [59] J. Milnor, *Lectures on the h-cobordism theorem*, *Mathematics Notes*, Princeton University Press, (1965).
- [60] M. Audin and M. Damian, *Morse Theory and Floer Homology*, [Springer](#), (2014).
- [61] L. Nicolaescu, *An invitation to Morse theory*, second edition, [Springer](#), (2011).
- [62] D. Tischler, *On fibering certain foliated manifolds over S^1* , *Topology* **9** (1970) 153.
- [63] A. Haefliger, *Groupoïdes d'holonomie de classifiants*, in *Structures Transverses des Feuilletages*, *Asterisque* **116** (1984) 70.
- [64] H. Looyestijn, E. Plaushinn and S. Vandoren, *New potentials from Scherk-Schwarz reductions*, *JHEP* **12** (2010) 016 [[arXiv:1008.4286](#)] [[INSPIRE](#)].

- [65] P. Kaste, R. Minasian and A. Tomasiello, *Supersymmetric M-theory compactifications with fluxes on seven-manifolds and G structures*, *JHEP* **07** (2003) 004 [[hep-th/0303127](#)] [[INSPIRE](#)].
- [66] Y.A. Kordyukov, *Adiabatic limits and spectral geometry of foliations*, *Math. Ann.* **313** (1999) 763. [[dg-ga/9506005](#)].
- [67] S. Karigiannis, *Desingularization of G2 manifolds with isolated conical singularities*, *Geom. Topol.* **13** (2009) 1583. [[arXiv:0807.3346](#)].
- [68] S. Karigiannis and J.D. Lotay, *Deformation theory of G2 conifolds* [arXiv:1212.6457](#).
- [69] N. Berline, E. Getzler and M. Vergne, *Heat Kernels and Dirac Operators*, *Grundlehren Math. Wiss.* **298** (1992) 1, Springer, New York, U.S.A..
- [70] K. Becker and M. Becker, *M theory on eight manifolds*, *Nucl. Phys. B* **477** (1996) 155 [[hep-th/9605053](#)] [[INSPIRE](#)].
- [71] D. Prins and D. Tsimpis, *Type IIA supergravity and M -theory on manifolds with SU(4) structure*, *Phys. Rev. D* **89** (2014) 064030 [[arXiv:1312.1692](#)] [[INSPIRE](#)].
- [72] J.M. Lee, D. Lear, J. Roth, J. Coskey and L. Nave, *Ricci — A Mathematica package for doing tensor calculations in differential geometry*, <http://www.math.washington.edu/~lee/Ricci/>.
- [73] M. Freeman, *Fully integrable Pfaffian systems*, *Ann. Math.* **119** (1984) 465.
- [74] I.A. Mel'nikova, *Non-compact leaves of foliations of Morse forms*, *Math. Notes* **63** (1998) 760.