

CP and discrete flavour symmetries

Martin Holthausen,^a Manfred Lindner^a and Michael A. Schmidt^b

^aMax-Planck Institut für Kernphysik,
Saupfercheckweg 1, 69117 Heidelberg, Germany

^bARC Centre of Excellence for Particle Physics at the Terascale,
School of Physics, The University of Melbourne,
Victoria 3010, Australia

E-mail: martin.holthausen@mpi-hd.mpg.de, lindner@mpi-hd.mpg.de,
michael.schmidt@unimelb.edu.au

ABSTRACT: We give a consistent definition of generalised CP transformations in the context of discrete flavour symmetries. Non-trivial consistency conditions imply that every generalised CP transformation can be interpreted as a representation of an automorphism of the discrete group. This allows us to give consistent generalised CP transformations of popular flavour groups. We are able to clear up issues concerning recent claims about geometrical CP violation in models based on T' , clarify the origin of "calculable phases" in $\Delta(27)$ and explain why apparently CP violating scalar potentials of A_4 result in a CP conserving ground state.

KEYWORDS: CP violation, Discrete and Finite Symmetries, Beyond Standard Model, Neutrino Physics

ARXIV EPRINT: [1211.6953](https://arxiv.org/abs/1211.6953)

Contents

1	Introduction	1
2	Generalised CP and the outer automorphism group	2
3	Physical implications of a generalised CP symmetry	7
4	Application to questions in the literature	8
4.1	$Z_3 \cong \text{SG}(3, 1)$	8
4.2	$A_4 \cong (Z_2 \times Z_2) \rtimes Z_3 \cong \text{SG}(12, 3)$	9
4.3	$T' \cong \text{SG}(24, 3)$	13
4.4	$\Delta(27) \cong (Z_3 \times Z_3) \rtimes Z_3 \cong \text{SG}(27, 3)$	18
4.5	$Z_9 \rtimes Z_3 \cong \text{SG}(27, 4)$	21
4.6	$Q_8 \times A_4 \cong \text{SG}(96, 204)$	22
4.7	$S_4 \cong (Z_2 \times Z_2) \rtimes S_3 \cong \text{SG}(24, 12)$	25
4.8	$T_7 \cong Z_7 \rtimes Z_3 \cong \text{SG}(21, 1)$	25
4.9	$\Delta(108) \cong \text{SG}(108, 22)$ (or $\Delta(216) \cong \text{SG}(216, 95)$)	26
5	Conclusions and outlook	27

1 Introduction

After the discovery of a sizeable value of θ_{13} by the reactor experiments DoubleChooz [1], DayaBay [2] and RENO [3] the door has been pushed wide open to measure the last undetermined parameters of the Standard Model, namely the CP phases of the lepton sector. Of special interest is the Dirac CP-phase δ_{CP} as it can be experimentally determined in neutrino oscillation experiments in the foreseeable future.¹

In the lepton sector, there is the proud/infamous tradition to explain the structure of mixing angles through the introduction of non-abelian discrete symmetries. The relative lack of success with regard to the reactor angle θ_{13} has not deterred the field from using the same set of ideas to try and predict the missing CP phase δ_{CP} using discrete symmetries. For example, a geometrical origin of the CP phase has been discussed for the group $\Delta(27)$ [4–9] and there have been attempts to explain CP violation as a result of complex Clebsch-Gordan coefficients in models based on the group T' [10, 11]. However, sometimes definitions of CP have been used that are incompatible with the discrete flavour symmetry, leading to inconsistencies, as will be discussed in detail later. In order to relate CP violation to the complex Clebsch-Gordan coefficients, a CP symmetry has to be imposed on the Lagrangian, which is then broken spontaneously [12, 13].²

¹To discern Majorana phases from possible future signals of neutrinoless double beta decay experiments will always be model dependent and thus seem less promising.

² Recently, a general group-theoretic condition for spontaneous CP violation has been given in ref. [14].

We here give a consistent general definition of CP transformations in the context of non-abelian discrete flavour groups. We will show that in many cases it is not possible to define CP in the naive way, $\phi \rightarrow \phi^*$, but rather a non-trivial transformation in flavour space is needed. Indeed there is a one-to-one correspondence between generalised CP transformations [15–17] and the outer automorphism group of the flavour group. It should not be surprising that outer automorphisms play a role in the definition of CP as complex conjugation is an outer automorphism of the field of complex numbers and the definition of CP transformations as automorphisms in the context of gauge theories has been discussed long ago by Grimus and Rebelo [18]. Generalised CP transformations in the context of discrete symmetries have been used before in ref. [19–28].

While the outer automorphism group of continuous groups is either trivial or a Z_2 (with the sole exception of $SO(8)$, whose outer automorphism group is S_3), the outer automorphism group of discrete groups can be very rich. For example the well-known flavour group $\Delta(27)$ has an automorphism group of order 432.

As a result of our investigation of generalised CP transformations, we present consistent definitions of CP for all groups of order smaller than 31 that contain three dimensional representations. Highlights are the case of T' , where we show that there is one consistent definition of CP, which we apply to the models discussed in ref. [10, 11]. We show that this CP is spontaneously broken by the VEVs of the doublets and it is additionally explicitly broken by the phases of Yukawa couplings and therefore the results obtained have to be considered as unphysical and basis dependent. For the group $\Delta(27)$ we are able to explain the so-called calculable phases as a result of an accidental generalised CP symmetry that had so far been overlooked in the literature.³

The outline of the paper is as follows. In section 2, we define a generalised CP transformation and discuss its connection with the outer automorphism group. The implications of a generalised CP transformation for the physical phases are discussed in section 3. In section 4, we apply our general considerations to specific examples. In order to uniquely specify each group, we denote it by $SG(O, N)$ with O being its order and N , the number in the GAP [31] SmallGroups catalogue [32]. In particular, we will discuss all groups of order less than 31 with a three-dimensional representation. Finally, we conclude in section 5.

For the convenience of the reader, we will briefly define all relevant group theoretical notions in the text or in a footnote. More detailed knowledge can be gained from standard group theory text books. See [33–35] for an overview of discrete groups, which have been used in the context of flavour symmetries.

2 Generalised CP and the outer automorphism group

In order to simplify the discussion, we will focus on finite discrete groups only. We do not consider the transformation under the Lorentz group or any continuous symmetry group and therefore restrict ourselves to scalar multiplets unless stated otherwise. An extension to

³Accidental CP symmetries have also been observed in scalar potentials in models based on dihedral groups D_n and its double cover Q_n [29, 30].

higher spin representations of the Lorentz group and continuous groups is straightforward. Let us consider a scalar multiplet

$$\phi = \left(\varphi_R, \varphi_P, \varphi_P^*, \varphi_C, \varphi_C^* \right)^T \tag{2.1}$$

that contains fields in real(R), pseudo-real(P) and complex(C) representations of the discrete group G . Note that ϕ always contains the field and its complex conjugate. The discrete group G acts on ϕ as

$$\phi \xrightarrow{G} \rho(g)\phi, \quad g \in G. \tag{2.2}$$

where ρ is a representation $\rho : G \rightarrow GL(N, \mathbb{C})$, which is generally reducible. In fact $\rho(G) \subset U(N)$, since we are only considering unitary representations. The representation ρ decomposes in a block diagonal form

$$\rho = \begin{pmatrix} \rho_R & & & & \\ & \rho_P & & & \\ & & \rho_P^* & & \\ & & & \rho_C & \\ & & & & \rho_C^* \end{pmatrix}. \tag{2.3}$$

A generalised CP transformation has to leave $|\partial\phi|^2$ invariant and thus is of the form

$$\phi \xrightarrow{CP} U\phi^* \tag{2.4}$$

with U being a unitary matrix, which is not necessarily block-diagonal, because it generically interchanges representations. Even different real representations can be connected by such a CP transformation, as we will discuss later.

If the representation is real, i.e. $\rho = \rho^*$, there is always the trivial CP transformation $\phi \rightarrow \phi^*$, which acts trivially on the group. In the following, we will take ρ to be complex and *faithful*, i.e. ρ is injective. If ρ were not faithful then the theory would only be invariant under the smaller symmetry group isomorphic to $G/\ker \rho$ and the restricted representation would be faithful.

Note that eq. (2.4) in combination with eq. (2.1) implies the existence of a matrix W with $W^2 = \mathbb{1}$, which exchanges the complex conjugate components of ϕ ,

$$\phi^* = W\phi, \quad \text{which implies } \rho(g) = W\rho(g)^*W^{-1}. \tag{2.5}$$

See section 4.1 and especially eq. (4.3) for a concrete example. Comparing first performing a group transformation and then performing a CP transformation with the inverse order of operations and demanding that the resulting transformation is contained in the symmetry group G of the theory, as shown in figure 1, one finds the requirement that

$$U\rho(g)^*U^{-1} \in \text{Im}\rho \equiv \rho(G), \tag{2.6}$$

i.e. the CP transformation maps group elements $\rho(g)$ onto group elements $\rho(g')$. We will refer to this condition as consistency condition and denote models satisfying this condition




Figure 1: CP definition.

consistent. If the condition (2.6) is not fulfilled, the group G is not the full symmetry group of the Lagrangian and one would have to consider the larger group, which closes under CP transformations (2.6). We do not consider this case further and will assume that the group G is the full symmetry group of the Lagrangian. Hence, a generalised CP transformation preserves the group multiplication, i.e. $U\rho(g_1g_2)^*U^{-1} = U\rho(g_1)^*U^{-1}U\rho(g_2)^*U^{-1}$ and $U\mathbb{1}^*U^{-1} = \mathbb{1}$, and therefore is a homomorphism.⁴ Furthermore the CP transformation is bijective, since U is unitary and therefore invertible. Hence, CP is an automorphism⁵ of the group, as is depicted in figure 2.

Indeed, *the possible matrices U of eq. (2.6) form a representation of the automorphism group⁶ $\text{Aut}(G)$ of G , which we are showing in the following.*

U represents the automorphism $u : G \rightarrow G$ given by

$$u : g \in G \rightarrow \rho(g) \rightarrow U\rho(g)^*U^{-1} = \rho(g') \rightarrow g' = \rho^{-1}(U\rho(g)^*U^{-1}) \in G \quad (2.7)$$

or

$$U\rho(g)^*U^{-1} = \rho(u(g)) . \quad (2.8)$$

It is straightforward to show that this mapping $u : G \rightarrow G$ is an automorphism, indeed.

Vice versa, if $u : G \rightarrow G$ is an automorphism, we can explicitly construct a matrix U in the following way. We first extend G to a group G' containing G as a normal subgroup and $u(g) = g'gg'^{-1} \forall g \in G$ with $g' \in G'$. This can be achieved as follows. Taking the order of u ⁷ to be $\text{ord}(u) = n$, we define the homomorphism

$$\theta : Z_n = (\{0, \dots, n - 1\}, +) \rightarrow \text{Aut}(G) : 1 \rightarrow \theta_1 \equiv u , \quad (2.9)$$

which has a trivial kernel. This homomorphism thus defines the semi-direct product group $G' = G \rtimes_{\theta} Z_n$ with the group multiplication

$$(g_1, z_1) \star (g_2, z_2) = (g_1\theta_{z_1}(g_2), z_1 + z_2) . \quad (2.10)$$

⁴A (group) homomorphism $\mu : G \rightarrow H$ is a mapping preserving the group structure, i.e. $\mu(g_1g_2) = \mu(g_1)\mu(g_2) \forall g_{1,2} \in G$, $\mu(g^{-1}) = \mu(g)^{-1}$, and $\mu(E_G) = E_H$, where $E_{G,H}$ denotes the identity elements of G and H , respectively.

⁵An automorphism μ of a group G is a bijective homomorphism $\mu : G \rightarrow G$.

⁶The automorphism group $\text{Aut}(G)$ is the set of all automorphisms of G with composition as group multiplication.

⁷The order of a group element u of G is given by the smallest $n \in \mathbb{N}$ with $u^n = \text{id}_G$.




Figure 2: The matrix U that appears in the definition of CP defines an automorphism $u : G \rightarrow G$ of the group G .

Keeping track of the multiplication rules, we find

$$(E, 1) \star (g, z) \star (E, 1)^{-1} = (u(g), z) , \tag{2.11}$$

where E is the identity element of G . The outer⁸ automorphism u of G becomes an inner automorphism of G' and we can obtain a matrix representation of u (or equivalently $(E, 1)$) by the standard techniques for finding matrix representations of groups, for example by using the computer algebra system **GAP** [31]. In order to relate the matrix representation of u in the semidirect product group G' with the matrix U in the CP transformation of a representation ρ of G , we have to consider a representation ρ' of G' whose restriction to the subgroup G is the representation ρ ,⁹ i.e. $\rho'|_G = \rho$. In terms of the representation ρ' , eq. (2.11) translates to

$$\rho'((E, 1))\rho'((g, z))\rho'((E, 1)^{-1}) = \rho'((u(g), z)) , \tag{2.12}$$

which can be written in terms of a CP transformation

$$U\rho'((g, z))^*U^{-1} = \rho'((u(g), z)) \tag{2.13}$$

with $U = \rho'((E, 1))W$ using the matrix W introduced in eq. (2.5). Finally, we have to consider the restriction of ρ' to the subgroup G , i.e. $\rho'|_G = \rho$ and therefore

$$U\rho(g)^*U^{-1} = \rho(u(g)) . \tag{2.14}$$

with $\rho(g) = \rho'((g, 0))$. Hence there is a one-to-one correspondence between set of matrices U and the automorphism group of G . If ρ does not contain all representations that are connected via the outer automorphism u , there is no matrix U that fulfils the consistency condition (2.14). To implement this CP transformation, the vector ϕ therefore has to be enlarged by the missing representations. For example in case of the group $Q_8 \times A_4$, which we discuss in section 4.6, different real representations are interchanged by the matrix U and the CP transformation can only be implemented if all representations connected by the corresponding automorphism are present in the theory. The same is true for the group A_5

⁸An *inner automorphism* μ_h of a group G is an automorphism, which is represented by conjugation with an element $h \in G$, i.e. $\mu_h \equiv \text{conj}(h) : g \rightarrow hgh^{-1}$. If an automorphism can not be represented by conjugation with a group element, it is called an *outer automorphism*.

⁹There is always a representation ρ' , whose restriction to G , $\rho'|_G$, contains ρ . If $\rho'|_G \neq \rho$, one has to extend ρ to $\rho'|_G$, because the CP transformation connects the representation ρ to another representation.

where the two different real 3-dimensional representations are interchanged by the outer automorphism of the group.

The automorphisms form a group with composition as group multiplication, i.e. $u' = \tilde{u} \circ u$ is again an automorphism represented by

$$U' \rho(g) * U'^{-1} = \rho(u'(g)) \tag{2.15}$$

with

$$\rho(u'(g)) = \rho(\tilde{u}(u(g))) = \tilde{U}W\rho(u(g))W\tilde{U}^{-1} = \tilde{U}WU\rho(g)*U^{-1}W\tilde{U}^{-1} \tag{2.16}$$

and thus

$$U' = \tilde{U}WU. \tag{2.17}$$

The trivial automorphism $\text{id}(g) = g \ \forall g \in G$ is represented by $U = W$ and the inverse automorphism u^{-1} is represented by $WU^{-1}W$. We thus have a homomorphism from the automorphism group to the group of matrices U defined in eq. (2.4) with the conjunction $\star: (A, B) \rightarrow A \star B \equiv AWB$. With respect to this conjunction the matrices U form a representation of the automorphism group. ■

For any solution U of eq. (2.6) the matrix $\rho(g)U$ is also a solution for any $g \in G$, which corresponds to performing a CP transformation followed by a group transformation described by $\rho(g)$. The group transformation corresponds to an inner homomorphism, which does not pose any new restrictions.¹⁰ It is therefore sufficient to consider automorphisms with inner automorphisms modded out. *Hence the group of generalised CP transformations is given by the outer automorphism group, which is defined by*

$$\text{Out}(G) \equiv \text{Aut}(G)/\text{Inn}(G) , \tag{2.18}$$

where $\text{Inn}(G)$ denotes the *inner automorphism group*,¹¹ the set of all inner automorphisms. Moreover, since the invariance under a CP transformation leads to the invariance under the subgroup generated by CP, the physically distinct classes of CP are given by the subgroups of the outer automorphism group.

As we will be using the character table in the discussion of the different groups (see e.g. table 1), we will briefly comment on how automorphisms act on the character table. As automorphisms are mappings from the group into itself and there is a unique character table for each group up to reordering of rows and columns, automorphisms are symmetries of the character table and can not change the character table besides exchanging rows and columns. While the action of the automorphisms on conjugacy classes should be self-explanatory, the action on representations requires further comment: Note that eq. (2.8)

¹⁰There are interesting phenomenological consequences for inner automorphisms, if the CP symmetry is left unbroken in one sector of the theory like the neutrino sector as discussed in [28]. However, we are more interested in the consistent definition in the unbroken theory and therefore do not further consider inner automorphisms.

¹¹For every group G there is a natural group homomorphism $G \rightarrow \text{Aut}(G)$ whose image is $\text{Inn}(G)$ and whose kernel is the *centre of G* , $Z(G)$, i.e. the subset of G which commutes with all elements of G . In short $\text{Inn}(G) \cong G/Z(G)$. Thus, if G has trivial centre it can be embedded into its own automorphism group.

may be read as a similarity transformation between the representations ρ^* and $\rho \circ u$. By composition an automorphism therefore interchanges representations. This is the reason why we have chosen to define CP on the reducible representation shown in eq. (2.1).

Inner automorphisms act via conjugation on the group. Hence, they map elements of the individual conjugacy class onto elements of the same conjugacy class. Neither do they exchange representations and therefore they do not change the character table. Outer automorphisms on the other hand map elements from one conjugacy class to another as well as one representation to another and thus exchange rows and columns.

Finally, note that it is always possible to use the freedom of a multiplication with an arbitrary phase factor. Hence if $\phi \rightarrow U\phi^*$ is a generalised CP transformation, so is

$$\phi \rightarrow e^{i\alpha}U\phi. \tag{2.19}$$

This does not lead to any additional constraints but only changes the phase factors of the different couplings.

We follow [18] and call a basis where U may be represented by the identity matrix times a phase, $\phi \rightarrow e^{i\alpha}\phi^*$, a *CP basis*. Note that under a change of basis $\phi' = V\phi$ we have

$$\phi' \rightarrow (VUV^T)\phi' \tag{2.20}$$

and it is thus not always possible to perform a basis change to a CP basis where VUV^T is diagonal [36], but it is possible to go to a basis where U takes the form [37]

$$VUV^T = \begin{pmatrix} O_1 & & & \\ & \ddots & & \\ & & O_l & \\ & & & \mathbb{1}_m \end{pmatrix} \tag{2.21}$$

with $2l + m = \dim(U)$ and O_i being orthogonal 2×2 matrices.

3 Physical implications of a generalised CP symmetry

The existence of a generalised CP symmetry implies that there is no direct CP violation and CP violation can only be generated via spontaneous symmetry breaking. This has been studied in terms of weak basis invariants [38–40]. A necessary and sufficient set of weak basis invariants, which measure the CP violation in the lepton sector and vanish in the CP conserving case has been proposed in [40]. In the following, we will explicitly demonstrate that the weak basis invariant for Dirac CP violation vanishes for our generalised CP symmetry and refer the reader to [40] for the remaining weak basis invariants. Let us consider a left-handed lepton doublet $L = (\nu, e)^T$ with the following mass terms

$$\mathcal{L}_{\text{mass}} = -e^T M_e e^c - \frac{1}{2} \nu^T M_\nu \nu + \text{h.c.} . \tag{3.1}$$

It was shown in ref. [38–40] that Dirac-type CP violation ($\sin \delta_{CP} \neq 0$) is equivalent to

$$0 \neq \text{tr} [H_\nu, H_e]^3 \quad \text{with} \quad H_\nu = (M_\nu^\dagger M_\nu)^* \quad \text{and} \quad H_e = (M_e M_e^\dagger)^T . \tag{3.2}$$

	E	T	T^2
$\mathbf{1}_1$	1	1	1
$\mathbf{1}_2$	1	ω	ω^2
$\mathbf{1}_3$	1	ω^2	ω

Table 1: Character table of Z_3 with $\omega = e^{2\pi i/3}$. The outer automorphism $u : T \rightarrow T^2$ is indicated in blue.

If L transforms under a generalised CP transformation as

$$L \xrightarrow{\text{CP}} UL^C \equiv U (i\sigma_2 L^*) \tag{3.3}$$

where L^C denotes charge conjugation with respect to the Lorentz group and U is unitary, the weak basis invariants $H_{\nu,e}$ have to fulfil

$$H_\nu = U^T H_\nu^T U^* \quad H_e = U^T H_e^T U^* \tag{3.4}$$

and therefore (note $[A, B]^T = -[A^T, B^T]$)

$$\text{tr} [H_\nu, H_e]^3 = \text{tr} U^T [H_\nu^T, H_e^T]^3 U^* = -\text{tr} [H_\nu, H_e]^3 = 0 \tag{3.5}$$

and there is thus no Dirac-type CP violation.

4 Application to questions in the literature

In the following, we will apply our general discussion to specific groups. We will concentrate on the most popular groups, which have been used in the literature.

4.1 $Z_3 \cong \text{SG}(3, 1)$

Let us start the discussion of examples by taking the cyclic group with three elements $Z_3 \cong \langle T | T^3 = E \rangle$, which is the smallest group with complex representations. There is one non-trivial automorphism $u : T \rightarrow T^2$, which is outer and since all group elements commute, there is only the trivial inner automorphism, $\text{conj}(T) = \text{conj}(T^2) = \text{id}$. The structure of automorphism group is thus:

$$\begin{aligned} \text{Z}(Z_3) &\cong Z_3 & \text{Aut}(Z_3) &\cong Z_2 \\ \text{Inn}(Z_3) &\cong Z_1 & \text{Out}(Z_3) &\cong Z_2 \end{aligned} \tag{4.1}$$

Looking at the character table in table 1, we see that the outer automorphism $u : T \rightarrow T^2$ indicated in blue acts on the character table by interchanging the conjugacy classes represented by T and $u(T) = T^2$ and the representations $\mathbf{1}_2 \leftrightarrow \mathbf{1}_2 \circ u = \mathbf{1}_3$, i.e. the rows and columns of the character table, such that the table stays invariant, as an outer automorphism should do.

Let us consider a theory that contains the complex representation $\varphi \sim \underline{\mathbf{1}}_2$. The vector $\phi = (\varphi, \varphi^*)^T$ is acted upon by the group generator T as

$$\rho(T) = \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix} \tag{4.2}$$

and we have $\rho(T)^* = \rho(T^2) = \rho(u(T)) \in \text{Im}\rho$ and therefore $U = \mathbb{1}_2$ is a representation of the outer automorphism $u : T \rightarrow T^2$. The generalised CP transformation (2.4) is therefore just the usual $\varphi \rightarrow \varphi^*$.

The matrix W relating the representation $\underline{\mathbf{1}}_2 \oplus \underline{\mathbf{1}}_3$ with its complex conjugate is given by

$$W = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \tag{4.3}$$

and $U = W$ represents the trivial automorphism, or $\varphi \rightarrow \varphi$.

While here it is trivial to find a matrix U representing the outer automorphism u , it is instructive to demonstrate the general method of constructing the semi-direct product group $G' = Z_3 \rtimes CP$ introduced in section 2 explicitly. It is given by $Z_3 \times Z_2$, where Z_3 is generated by T and Z_2 by the automorphism u . Hence, its elements are

$$\{(E, \text{id}), (T, \text{id}), (T^2, \text{id}), (E, u), (T, u), (T^2, u)\} \tag{4.4}$$

and the multiplication is defined by

$$(g_1, u_1) \star (g_2, u_2) \equiv (g_1 u_1(g_2), u_1 \circ u_2), \tag{4.5}$$

which defines a non-abelian group of order 6 and it is isomorphic to S_3 being the only non-abelian group of order 6. It has two generators: (T, id) and (E, u) . The outer automorphism $u : T \rightarrow T^2$ of Z_3 corresponds to the inner automorphism $\text{conj}((E, u)) : Z_3 \times Z_2 \ni g \rightarrow (E, u) \star g \star (E, u)^{-1}$ of $Z_3 \times Z_2 \cong S_3$. The group S_3 has three representations: $\underline{\mathbf{1}}_1, \underline{\mathbf{2}}$ and $\underline{\mathbf{3}}$; only the 2-dimensional representation is faithful and the generators are given by

$$\rho_{\underline{\mathbf{2}}}((T, \text{id})) = \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}, \quad \text{and} \quad \rho_{\underline{\mathbf{2}}}((E, u)) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{4.6}$$

In terms of the subgroup $Z_3 = \langle T \rangle$, it decomposes in the direct sum of the representations $\underline{\mathbf{1}}_2$ and $\underline{\mathbf{1}}_3$ of Z_3 with the group generator $\rho(T) = \rho_{\underline{\mathbf{2}}}((T, \text{id}))$. The automorphism u is represented by the matrix $U' = \rho_{\underline{\mathbf{2}}}((E, u))$ and $\rho(g) \rightarrow \rho(u(g)) = U' \rho(g) U'^{-1}$ and therefore the non-trivial CP transformation belonging to the automorphism u is given by $\rho(g) \rightarrow \rho(u(g)) = U \rho(g)^* U^{-1}$ with $U = U' W = \mathbb{1}_2$, as we have found above. Clearly the trivial automorphism corresponds to (E, id) and is represented by $U' = \mathbb{1}_2$ or $U = W$.

4.2 $A_4 \cong (Z_2 \times Z_2) \rtimes Z_3 \cong \text{SG}(\mathbf{12}, \mathbf{3})$

There is a complete classification of automorphism groups for the alternating groups A_n , which is shown in table 2a. Most of them have a very similar structure. We will discuss

	$Z(S_n)$	$\text{Aut}(S_n)$	$\text{Inn}(S_n)$	$\text{Out}(S_n)$
$n \geq 4, n \neq 6$	Z_1	S_n	A_n	Z_2
$n = 1, 2$	Z_n	Z_1	Z_1	Z_1
$n = 3$	Z_3	Z_2	Z_1	Z_2
$n = 6$	Z_1	$S_6 \rtimes Z_2$	A_6	$Z_2 \times Z_2$

	E	T	T^2	S
$\underline{\mathbf{1}}_1$	1	1	1	1
$\underline{\mathbf{1}}_2$	1	ω	ω^2	1
$\underline{\mathbf{1}}_3$	1	ω^2	ω	1
$\underline{\mathbf{3}}$	3	0	0	-1

 (a) Structure of the automorphism group of A_n

 (b) Character table of A_4 .

Table 2: Relevant group structure of the alternating groups A_n .

the specific case of $A_4 = \langle S, T | S^2 = T^3 = (ST)^3 = E \rangle$ ¹² in detail. It is very important for model building and serves as our first non-trivial example. As it can be seen in table 2a, only the identity element commutes with all other elements and the natural homomorphism $n : A_4 \rightarrow \text{Aut}(A_4)$ defined by $n(g) = \text{conj}(g)$ is therefore injective. There is one non-trivial outer automorphism $u : (S, T) \rightarrow (S, T^2)$. Here and in the following, we only give the action of automorphisms on the generators of the group, which uniquely defines an automorphism. The character table of A_4 is given in table 2b and it is easy to verify that the automorphism u represents a symmetry of the character table, again interchanging the representations $\underline{\mathbf{1}}_2$ and $\underline{\mathbf{1}}_3$. Let us first discuss the case where we have only one real scalar field in the real representation $\phi \sim \underline{\mathbf{3}}_1$ using the Ma-Rajasekaran [41] basis:

$$\rho_{\underline{\mathbf{3}}_1}(S) = S_3 \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \rho_{\underline{\mathbf{3}}_1}(T) = T_3 \equiv \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}. \quad (4.7)$$

In this basis both group generators are real ($\rho(g)^* = \rho(g) \in \text{Im}\rho$) and one might be tempted to take $U = \mathbb{1}_3$ as this fulfils eq. (2.6). However, the map derived from $U = \mathbb{1}_3$ via eq. (2.8) is not equal to $u : (S, T) \rightarrow (S, T^2)$, but the trivial automorphism id_{A_4} , which is obviously not outer and therefore does not lead to additional constraints on the couplings.¹³ One also encounters this problem as soon as one considers contractions such as

$$(\phi\phi)_{\underline{\mathbf{1}}_2} = \frac{1}{\sqrt{3}} (\phi_1\phi_1 + \omega^2\phi_2\phi_2 + \omega\phi_3\phi_3) \quad (4.8)$$

which transform under this "CP" $\phi \rightarrow U\phi^* = \phi$ as

$$(\phi\phi)_{\underline{\mathbf{1}}_2} \rightarrow (\phi\phi)_{\underline{\mathbf{1}}_2} \sim \underline{\mathbf{1}}_2 \quad (4.9)$$

which is in conflict with the expectation that CP should involve complex conjugation such that

$$(\phi\phi)_{\underline{\mathbf{1}}_2} \rightarrow [(\phi\phi)_{\underline{\mathbf{1}}_2}]^* \sim \underline{\mathbf{1}}_3. \quad (4.10)$$

¹² A_4 has been introduced as flavour symmetry in the lepton sector in [41].

¹³Obviously it still acts non-trivially on the space-time symmetry group as well as possibly the gauge group.

Just imagine that the theory contains a real scalar triplet $\chi \sim \mathbf{3}$ and a singlet $\xi \sim \mathbf{1}_3$. If one defines CP as $\chi \rightarrow \chi$ and $\xi \rightarrow \xi^*$ then the invariant $(\chi\chi)_{\mathbf{1}_2} \xi$ under CP is mapped to $(\chi\chi)_{\mathbf{1}_2} \xi^*$, which is not invariant under the group and it is forbidden by the combination of A_4 and this definition of CP. Looking at this definition of CP, i.e. $\chi \rightarrow \chi^*$ and $\xi \rightarrow \xi^*$, we can easily check that it does not fulfil the consistency condition in eq. (2.6) and therefore the true symmetry group of the Lagrangian is not A_4 , but the group generated by A_4 and this CP transformation. However, it has been (implicitly) used in a number of works [42–44]¹⁴ without properly taking into account the enlarged symmetry group with its additional restrictions on the Lagrangian.

If we instead use the non-trivial solution of eq. (2.6), which has been discussed in [19–27]

$$U = U_3 \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \tag{4.11}$$

that corresponds to the outer automorphism $u : (S, T) \rightarrow (S, T^2)$ we immediately see that

$$(\phi\phi)_{\mathbf{1}_2} \rightarrow [(\phi\phi)_{\mathbf{1}_2}]^* \sim \mathbf{1}_3. \tag{4.12}$$

Note that this is the only non-trivial definition of CP (up to inner automorphisms) in any theory that involves the complex representations, since the outer automorphism group is Z_2 .

Using eq. (2.6), we can immediately see that the solution $U = \mathbb{1}_3$ for $\rho \sim \mathbf{3}$ leads to the trivial automorphism id_{A_4} (up to inner automorphism), when it is extended to the other representations. Let us consider the vector $\phi = (\xi, \xi^*, \chi)^T$ with $\xi \sim \mathbf{1}_3$ and $\chi \sim \mathbf{3}_1$ which transforms as

$$\rho(S) = \text{diag}(1, 1, S_3) \quad \rho(T) = \text{diag}(\omega, \omega^2, T_3) \tag{4.13}$$

and clearly fulfils $\rho(S)^* = \rho(S) \in \text{Im}\rho$ and $\rho(T)^* \notin \text{Im}\rho$. We are therefore forced to use $U = \text{diag}(1, 1, U_3)$, which gives $U\rho(T)^*U^{-1} = \rho(T^2) \in \text{Im}\rho$ and $U\rho(S)^*U^{-1} = \rho(S) \in \text{Im}\rho$ and represents the outer automorphism $u : (S, T) \rightarrow (S, T^2)$. The only consistent (meaning satisfying condition (2.6)) non-trivial CP transformation in this theory is thus $\xi \rightarrow \xi^*$ and $\chi \rightarrow U_3\chi^* = U_3\chi$. Adding the generator U to A_4 results in S_4 because A_4 can be embedded in $\text{Aut}(G)$.

Summarising our discussion, there is only one non-trivial CP transformation (up to inner automorphisms) acting on the reducible representation $\phi \sim \mathbf{1}_1 \oplus \mathbf{1}_2 \oplus \mathbf{1}_3 \oplus \mathbf{3}$, which takes the form $\phi \rightarrow U\phi^*$ with

$$U = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & U_3 \end{pmatrix}. \tag{4.14}$$

¹⁴The discussion of CP in ref. [42] has been corrected in ref. [45].

The trivial CP transformation corresponding to the trivial automorphism id_{A_4} is determined by $\phi \rightarrow U\phi^*$ with

$$U = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \mathbb{1}_3 \end{pmatrix}, \quad (4.15)$$

which is equivalent to the transformation $\phi \rightarrow \phi$ as can be easily checked. There are no other CP transformations (up to inner automorphisms).

Since this case is of some relevance to model building, let us dwell on it a bit more and repeat the discussion for the basis

$$S = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix} \quad (4.16)$$

first used by Altarelli and Feruglio [46]. Here the group elements are complex but the Clebsch-Gordon coefficients are real. The unique result of eq. (2.6) is $U = \mathbb{1}_3$ up to inner automorphisms. This basis is therefore a CP basis, as defined in eq. (2.20). Note that in this basis

$$(\phi\phi)_{\mathbf{1}_2} = (\phi_2\phi_2 + \phi_1\phi_3 + \phi_3\phi_1), \quad (\phi\phi)_{\mathbf{1}_3} = (\phi_3\phi_3 + \phi_1\phi_2 + \phi_2\phi_1) \quad (4.17)$$

and thus

$$(\phi\phi)_{\mathbf{1}_2} \rightarrow [(\phi\phi)_{\mathbf{1}_2}]^* \sim \mathbf{1}_3. \quad (4.18)$$

as it should be.

Let us look at a physical situation where a certain confusion about the definition of CP can be alleviated by our definition.¹⁵ If one considers the potential for one electroweak Higgs doublet transforming as $\mathbf{3}_1$ denoted by $\chi = (\chi_1, \chi_2, \chi_3)^T$ in the basis (4.7), there is one potentially complex coupling in the potential [41, 42, 44]

$$\lambda_5 (\chi^\dagger\chi)_{\mathbf{3}_1} (\chi^\dagger\chi)_{\mathbf{3}_1} + \text{h.c.} = \lambda_5 \left[(\chi_1^\dagger\chi_2)^2 + (\chi_2^\dagger\chi_3)^2 + (\chi_3^\dagger\chi_1)^2 \right] + \text{h.c.} \quad (4.19)$$

It can be easily checked that the generalised CP transformation $\chi \rightarrow U_3\chi^*$ acts as

$$I \equiv \left[(\chi_1^\dagger\chi_2)^2 + (\chi_2^\dagger\chi_3)^2 + (\chi_3^\dagger\chi_1)^2 \right] \rightarrow \left[(\chi_1^\dagger\chi_2)^2 + (\chi_2^\dagger\chi_3)^2 + (\chi_3^\dagger\chi_1)^2 \right] = I \quad (4.20)$$

and thus does not give a restriction on the phase of λ_5 . Note that the naive CP transformation $\chi \rightarrow \chi^*$ transforms the group invariant I into I^* and therefore restricts λ_5 to be real as was e.g. done in ref. [43]. However, we have seen that this naive CP transformation cannot be consistently implemented on the Lagrangian level if there are complex representations, unless it is either the trivial generalised CP transformation, id_{A_4} , or the symmetry group A_4 is extended such that it is closed under this naive CP transformation. Therefore it is

¹⁵For a related discussion, see [47–49].

inappropriate to call the phase of λ_5 a CP phase. This also explains an observation made in ref. [42], where it was shown that even for $\arg \lambda_5 \neq 0$ the VEV configuration

$$\langle \chi \rangle = V(1, 1, 1), \quad \langle \chi \rangle = V(1, 0, 0) \quad V \in \mathbb{R}, \quad (4.21)$$

which of course respects both, the trivial as well as the non-trivial, generalised CP transformations, can be obtained without fine-tuning. This would have been somewhat surprising, as usually symmetry conserving solutions cannot be obtained from explicitly symmetry breaking potentials. However, the phase of λ_5 does not break the consistent definition of generalised CP, i.e. does not violate condition (2.6), as does the VEV configuration (4.21), therefore everything is consistent.

4.3 $T' \cong \text{SG}(24, 3)$

The group $T' = \langle S, T | S^4 = T^3 = (ST)^3 = E \rangle \cong \text{SL}(2, 3)$,¹⁶ is also an important group in the context of CP violation [10, 11]. It has two elements $Z(T') = \{E, S^2\} \cong Z_2$ that commute with all group elements and therefore $\text{Inn}(T') \cong T'/Z(T') \cong A_4$. There is one non-trivial outer automorphism (up to inner automorphisms) $u : (S, T) \rightarrow (S^3, T^2)$. Therefore the automorphism structure can be summarised as:

$$\begin{aligned} Z(T') &\cong Z_2 & \text{Aut}(T') &\cong S_4 \\ \text{Inn}(T') &\cong A_4 & \text{Out}(T') &\cong Z_2 \end{aligned} \quad (4.22)$$

A non-trivial CP transformation therefore has to be a representation of u in the sense of eq. (2.6). Let us now see how it is represented for the various representations of T' .

There is a faithful pseudo-real representation

$$\mathbf{2}_1 : S = A_1, \quad T = \omega A_2 \quad (4.23)$$

with $\sigma_2^\dagger S \sigma_2 = S^*$ and $\sigma_2^\dagger T \sigma_2 = T^*$ and the two faithful complex representations

$$\mathbf{2}_2 : S = A_1 \quad T = \omega^2 A_2; \quad \mathbf{2}_3 : S = A_1, \quad T = A_2 \quad (4.24)$$

with $\sigma_2^\dagger S_2 \sigma_2 = S_2^*$ and $\sigma_2^\dagger T_2 \sigma_2 = T_2^*$ where

$$A_1 = \frac{-1}{\sqrt{3}} \begin{pmatrix} i & \tilde{\omega}\sqrt{2} \\ -\tilde{\omega}^{-1}\sqrt{2} & -i \end{pmatrix}, \quad A_2 = \begin{pmatrix} \omega & 0 \\ 0 & 1 \end{pmatrix} \quad (4.25)$$

with $\tilde{\omega} = e^{2\pi i/24}$. For all two-dimensional representations, we find the matrix

$$U = U_2 \equiv \text{diag}(\tilde{\omega}^{-5}, \tilde{\omega}^5) \quad (4.26)$$

which represents the automorphism u via $U\rho(g)^*U^{-1} = \rho(u(g))$. For the three-dimensional representation

$$\rho(S) = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega^2 & -1 & 2\omega \\ 2\omega & 2\omega^2 & -1 \end{pmatrix}, \quad \rho(T) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \quad (4.27)$$

¹⁶ T' has been first discussed in a particle physics context in [50].

the matrix U of eq. (2.4) is given by $U = \rho(T)$ with again $U\rho(T)^*U^{-1} = \rho(T^2)$, $U\rho(S)^*U = \rho(S^3)$, for the one dimensional representations we take $U = \rho(T)$ as for the three-dimensional representations.

In summary, we have thus found the one unique non-trivial outer automorphism (up to inner automorphisms) of T' and thus the unique CP transformation

$$\mathbf{1}_i \rightarrow \omega^{i-1} \mathbf{1}_i^* \quad \mathbf{2}_i \rightarrow \text{diag}(\tilde{\omega}^{-5}, \tilde{\omega}^5) \mathbf{2}_i^* \quad \mathbf{3} \rightarrow \text{diag}(1, \omega, \omega^2) \mathbf{3}^*. \quad (4.28)$$

Let us now use this insight to investigate a claim that there is geometrical CP violation in grand unified models based on T' [10, 11]. We consider the model discussed in [10] and introduce $(T_1, T_2) \sim \mathbf{2}_1$ which transforms as $\mathbf{10}$ of $SU(5)$ and includes the first two generations of up-type quarks and the flavons $\phi \sim \mathbf{3}$ and $\phi' \sim \mathbf{3}$. Auxiliary $Z_{12} \times Z_{12}$ symmetries are introduced such that the one-two sector of the mass matrix is described by¹⁷

$$\begin{aligned} -\mathcal{L}_{TT} &= y_c TT\phi^2 + y_u TT\phi'^3 + \text{h.c.} \quad (4.29) \\ &\equiv y_c \frac{3}{2} \frac{2-i}{2} (TT)\mathbf{3}(\phi^2)\mathbf{3} + y_u \frac{1}{3} [(TT)\mathbf{3}\phi']\mathbf{1}_3(\phi'^2)\mathbf{1}_2 + \text{h.c.} \\ &= y_c \frac{3}{2} \frac{2-i}{2} \{ (1-i)T_1T_2(\phi_1^2 - \phi_2\phi_3) + iT_1^2(\phi_2^2 - \phi_1\phi_3) + T_2^2(\phi_3^2 - \phi_1\phi_2) \} + \\ &+ y_u \frac{1}{3} \left\{ (2\phi'_1\phi'_3 + \phi'^2_2) (iT_1^2\phi'_1 + (1-i)T_1T_2\phi'_2 + T_2^2\phi'_3) \right\} + \text{h.c.}, \end{aligned}$$

where we have omitted (Higgs-) fields that do not transform under the flavour symmetry and a suppression by some high-energy scale of a sufficient power to make y_i dimensionless is understood.

It is assumed that the VEVs

$$\langle \phi' \rangle = (1, 1, 1)V', \quad \langle \phi \rangle = (0, 0, 1)V \quad V, V' \in \mathbb{R} \quad (4.30)$$

are real, which may be justified by a CP transformation. There is only one CP transformation¹⁸ left invariant, namely the one corresponding to the outer automorphism $u' = \text{conj}(T^2) \circ u$ represented on the three dimensional representation by the identity matrix

$$\mathbf{1}_i \rightarrow \mathbf{1}_i^* \quad \mathbf{2}_i \rightarrow \text{diag}(\omega\tilde{\omega}^{-5}, \omega^{-1}\tilde{\omega}^5) \mathbf{2}_i^* \quad \mathbf{3} \rightarrow \mathbf{3}^*. \quad (4.31)$$

and therefore $\langle \phi' \rangle \rightarrow \langle \phi' \rangle^*$ and $\langle \phi \rangle \rightarrow \langle \phi \rangle^*$.

This results in the following 1-2 block of the up-type quark mass matrix M_u :

$$y_u \begin{pmatrix} i & \frac{1-i}{2} \\ \frac{1-i}{2} & 1 \end{pmatrix} V'^3 + y_c \begin{pmatrix} 0 & 0 \\ 0 & 1 - \frac{i}{2} \end{pmatrix} V^2. \quad (4.32)$$

At this point the parameters $y_{u,c}$ and VEVs are chosen real and it is claimed that the phases emerging from the complex Clebsch-Gordon coefficients explain CP violation. Therefore it is natural to ask whether this choice of parameters can be justified by a symmetry. The

¹⁷We use the Clebsch-Gordan coefficients given in appendix A of [51] for the Kronecker products.

¹⁸Note that this also determines the global phase of U .

only candidate symmetry is a generalised CP symmetry of type (2.6), which we explicitly state in eq. (4.31). As we have shown how the various fields have to transform under the generalised CP symmetry we can now easily determine how the invariants of eq. (4.29) transform:¹⁹

$$CP[TT\phi^2] = -\frac{4+3i}{5}(TT\phi^2)^* \quad CP[TT\phi'^3] = -i(TT\phi'^3)^*. \quad (4.33)$$

Therefore invariance under CP requires $\arg(y_c) = -\frac{1}{2}\arg(-4-3i) = -\frac{1}{2}\arctan\frac{3}{4}$ and $\arg y_u = \frac{\pi}{4}$ and the generalised CP (4.31) is explicitly broken by real couplings y_u, y_c , which was assumed in ref. [10]. Note that also the relative phase between the two couplings does not agree with 'geometrical' CP violation. This also shows that the results obtained in ref. [10] are completely basis dependent and therefore unphysical.

Although the VEVs (4.30) are invariant under the generalised CP transformation (4.31), in the full model [10] there are additional scalar fields e.g. $\psi \sim \mathbf{2}_2$ with $\langle\psi\rangle \sim (1, 0)$ which are not invariant under the generalised CP transformation (4.31). Hence, if the phases of the couplings are changed in accordance with the consistent CP transformation (4.31), CP will be broken spontaneously. Obviously, all predictions depend on the VEV alignment. In ref. [10], no dynamical mechanism was given to generate the VEV configuration.

Different invariants were used in the other grand unified T' model [11] claiming a geometric origin of CP violation. In the following, we argue that the CP phases in this model do not have a geometric origin as well. The argument is done in two steps: 1) We choose a CP transformation, which is not broken by the VEVs. 2) CP is explicitly broken by two different couplings in the superpotential.

1) As the CP transformation defined in eq. (4.31) is not broken by real VEVs of the singlet and triplet flavons, it is enough to consider the four doublets $\psi^{(l)} = (\psi_1^{(l)}, \psi_2^{(l)})^T$ and $\tilde{\psi}^{(l)} = (\tilde{\psi}_1^{(l)}, \tilde{\psi}_2^{(l)})^T$, which obtain the VEVs

$$\langle\psi'\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \psi'_0, \quad \langle\tilde{\psi}'\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \tilde{\psi}'_0, \quad \langle\psi''\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \psi''_0, \quad \langle\tilde{\psi}''\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \tilde{\psi}''_0. \quad (4.34)$$

We modify the CP transformation (4.31) by a phase redefinition of the doublet fields

$$\begin{aligned} \psi' &\rightarrow \text{diag}(1, \omega^{-2}\tilde{\omega}^{10})\psi'^* & \psi'' &\rightarrow \text{diag}(\omega^2\tilde{\omega}^{-10}, 1)\psi''^* \\ \tilde{\psi}' &\rightarrow \text{diag}(1, \omega^{-2}\tilde{\omega}^{10})\tilde{\psi}'^* & \tilde{\psi}'' &\rightarrow \text{diag}(\omega^2\tilde{\omega}^{-10}, 1)\tilde{\psi}''^* \end{aligned} \quad (4.35)$$

such that the VEVs do not break this CP transformation.

¹⁹Note that inner automorphisms correspond to group transformations and therefore only outer automorphism can give non-trivial constraints when acting on group invariants. Here there is only one non-trivial outer automorphism(up to inner automorphisms).

2) Using this CP transformation, we consider two couplings in the superpotential²⁰

$$\begin{aligned}
 \mathcal{W}_{Y_u} &\supset y_{22} T T \tilde{\psi}''^2 \tilde{\zeta}' + y_{21} T T \tilde{\phi} \tilde{\psi}''^2 \tilde{\zeta}' & (4.36) \\
 &\equiv y_{22} ((T \tilde{\psi}'')_{\mathbf{3}} (T \tilde{\psi}'')_{\mathbf{3}})_{\mathbf{1}_3} \tilde{\zeta}' + y_{21} ((T \tilde{\phi})_{\mathbf{2}_2} \tilde{\zeta}')_{\mathbf{2}_3} (\tilde{\psi}' (T \tilde{\psi}')_{\mathbf{3}})_{\mathbf{2}_2} \\
 &= \frac{y_{22}}{\sqrt{3}} \tilde{\zeta}' \left\{ T_2^2 \tilde{\psi}_2''^2 + (1+i) T_1 \tilde{\psi}_1'' (T_1 \tilde{\psi}_2'' + T_2 \tilde{\psi}_1'') \right\} \\
 &+ \frac{y_{21}}{3\sqrt{2}} \tilde{\zeta}' \left\{ T_1^2 \left[(1-i) \tilde{\phi}_1 \tilde{\psi}_2''^2 - (1+i) \tilde{\phi}_2 \tilde{\psi}_2''^2 \right] + T_2^2 \left[(1+i) \tilde{\phi}_3 \tilde{\psi}_2''^2 - 2 \tilde{\phi}_2 \tilde{\psi}_1' \tilde{\psi}_2'' \right] \right. \\
 &\quad \left. + 2 T_1 T_2 \left[(1-i) \tilde{\phi}_1 \tilde{\psi}_1' \tilde{\psi}_2'' - i \tilde{\phi}_3 \tilde{\psi}_1''^2 \right] \right\}
 \end{aligned}$$

contributing to the 1-2 sector of the up-type quark mass matrix, where $T = (T_1, T_2)^T \sim \mathbf{2}_1$, $\tilde{\zeta}' \sim \mathbf{1}_2$ and $\tilde{\phi} = (\tilde{\phi}_1, \tilde{\phi}_2, \tilde{\phi}_3)^T \sim \mathbf{3}$. Similarly to the argument for [10], we can now easily determine how the invariants transform under the generalised CP transformation

$$CP[TT \tilde{\psi}''^2 \tilde{\zeta}'] = -i(TT \tilde{\psi}''^2 \tilde{\zeta}')^* \quad CP[TT \tilde{\phi} \tilde{\psi}''^2 \tilde{\zeta}'] = -(TT \tilde{\phi} \tilde{\psi}''^2 \tilde{\zeta}')^* . \quad (4.37)$$

Hence, CP is explicitly broken by choosing y_{22} and y_{21} real. As there is a relative phase difference between the two operators, it is not possible to redefine the CP transformation of T , such that there is no explicit CP violation.

The authors additionally propose a way to obtain the VEV alignment using driving fields using the method introduced in ref. [52]. Let us analyse the flavon potential in more detail using the generalised CP transformation (4.31), i.e. without the modification in eq. (4.35). It seems plausible to get real VEVs for the triplets and singlet fields, as these are eigenstates of CP. We will therefore concentrate on the doublets $\psi^{(l)}$ and $\tilde{\psi}^{(l)}$. The generalised CP transformation (4.31) fixes the phase (modulo π) of all couplings and in particular²¹

$$\begin{aligned}
 \mathcal{W}_f &\supset D_\psi (\lambda (\psi'')^2 + \kappa \phi \zeta') & (4.38) \\
 &= \frac{1}{\sqrt{3}} [D_{\psi_1} (\lambda \psi_2''^2 + \kappa \phi_3 \zeta') + D_{\psi_2} (i \lambda \psi_1''^2 + \kappa \phi_2 \zeta') + D_{\psi_3} ((1-i) \lambda \psi_1'' \psi_2'' + \kappa \phi_1 \zeta')]
 \end{aligned}$$

where $D_\psi \sim \mathbf{3}$ is a driving field, $\phi \sim \mathbf{3}$ and $\zeta' \sim \mathbf{1}_2$. CP invariance with respect to eq. (4.31) requires κ to be real and $\arg(\lambda) = \pi/4$. Assuming the VEV alignment (4.34), the F-term equation

$$0 \stackrel{!}{=} \frac{\partial \mathcal{W}_f}{\partial D_{\psi_1}} = \frac{1}{\sqrt{3}} (\lambda \psi_2''^2 + \kappa \phi_3 \zeta') \quad (4.39)$$

leads to a complex VEV for ψ'' with $\arg(\psi_0'') = 7\pi/8 + \mathbb{Z}\pi$ for $\lambda\kappa < 0$, which conserve CP, and $3\pi/8 + \mathbb{Z}\pi$ for $\lambda\kappa > 0$, which preserves $\psi_0'' \rightarrow -CP[\psi_0'']$, a different CP

²⁰A CP transformation relates the (holomorphic) superpotential with the anti-holomorphic superpotential. Similarly to eq. (4.29), we omit Higgs fields that do not transform under the flavour symmetry and a suppression by some high-energy scale Λ of a sufficient power to make y_i dimensionless is understood. We use the Clebsch-Gordan coefficients given in table 7 of ref. [11] itself. Note that the second operator is only defined up to a sign in ref. [11]. However, this does not affect the discussion.

²¹We dropped the fields ϵ_i , which are singlets of T' and are required to adjust the charges of the shaping symmetries.

transformation, which can also be extended to a symmetry of the full theory by changing the CP transformation of the doublets to $CP' : \mathbf{2}_i \rightarrow -CP[\mathbf{2}_i]$. Hence, it is not possible to break both CP transformations by the VEV of ψ'' alone. However, the VEV of a second doublet can break the remaining CP transformation, but the phases of the VEVs depend on a discrete choice. The other doublet VEVs are related to the VEV of ψ'' via $\arg(\tilde{\psi}''_0) = \arg(\psi''_0) + \mathbb{Z}\pi/2$ and $\arg(\psi'_0), \arg(\tilde{\psi}'_0) = -\arg(\psi''_0) + \mathbb{Z}\pi/2$, where the shift $\mathbb{Z}\pi/2$ depends on the sign of the respective couplings. Hence, there is a discrete set of phases of the VEVs. In analogy to the VEV of ψ'' , each VEV breaks one of the two CP transformations. Concluding, as soon as there are two VEVs, which break different CP transformations, it is possible to have CP violation.

It might be instructive to look at the potential for one doublet field $\psi \sim \mathbf{2}_2$ and study the VEV configurations that can be obtained in order to see if it is possible to obtain a phase prediction from a spontaneous breaking of the generalised CP. On renormalizable level there is only one coupling that depends on phases

$$\lambda \frac{\tilde{\omega}^2}{\sqrt{3}} (\psi_1(\psi_1^3 - (2 - 2i)\psi_2^3)) + \text{h.c.}, \tag{4.40}$$

where the phases have been adjusted such, that CP forces λ to be real. We will focus on VEVs of the form $\langle \psi \rangle = (Ve^{i\alpha}, 0)^T$ with $V > 0$ that conserve the Z_3 subgroup generated by T . For $\lambda < 0$ we find the minima $\{1, i, -1, -i\}(e^{i\pi 11/24}, 0)^T$, which conserve $\psi \rightarrow \{1, -1, 1, -1\}CP[\psi]$, and for $\lambda > 0$ we find the minima $\{1, i, -1, -i\}(e^{i\pi 5/24}, 0)^T$ which conserve $\psi \rightarrow -i\{1, -1, 1, -1\}CP[\psi]$. The additional solutions are due to fact that the phase dependent part of the potential has an accidental Z_4 symmetry $\psi \rightarrow i\psi$, which will most likely be broken in a full theory such that one would expect only the CP conserving solutions to survive. The required real VEVs cannot be obtained in this simple setup. Note that if the VEV of ψ conserves CP, the phases of the VEVs and of the couplings conspire that there is no CP violation, as shown in section 3.

Let us briefly summarise our view on geometrical CP violation in T' . To be able to talk about CP violation one has to apply the consistent CP symmetry of eq. (4.31) on the Lagrangian level. This will then fix the phase (modulo π) of most couplings. The phases of invariants, which are CP self-conjugate are not fixed by CP. In supersymmetric theories, the phases of all couplings in the superpotential are fixed (modulo π), because CP relates the superpotential with the anti-holomorphic superpotential. One could thus imagine a setup along the lines of [10, 11] where this has been implemented and therefore the only source of CP violation are the VEVs of the doublet scalar fields, which break CP spontaneously. However, the phases of the VEVs are only determined up to a finite discrete choice.

For usual spontaneous breaking of CP one would expect the phases of the fields to depend on potential parameters and therefore not be determined by the group symmetry structure. The only way to get 'calculable phases', i.e. phases that do not depend on potential parameters, seems to be if this CP breaking vacua is connected to an additional (accidental) CP symmetry of the potential as is the case for $\Delta(27)$ (see section 4.4). For T' , however, there cannot be such an additional generalised CP besides the CP transfor-

	E	$BABA$	ABA	A	BAB	AB	A^2	B^2	B	BA^2BAB	AB^2ABA
$\underline{1}_1$	1	1	1	1	1	1	1	1	1	1	1
$\underline{1}_2$	1	ω	ω^2	1	ω	ω^2	1	ω	ω^2	1	1
$\underline{1}_3$	1	ω^2	ω	1	ω^2	ω	1	ω^2	ω	1	1
$\underline{1}_4$	1	ω	ω	ω^2	ω^2	ω^2	ω	1	1	1	1
$\underline{1}_5$	1	ω^2	1	ω^2	1	ω	ω	ω	ω^2	1	1
$\underline{1}_6$	1	1	ω^2	ω^2	ω	1	ω	ω^2	ω	1	1
$\underline{1}_7$	1	ω^2	ω^2	ω	ω	ω	ω^2	1	1	1	1
$\underline{1}_8$	1	1	ω	ω	ω^2	1	ω^2	ω	ω^2	1	1
$\underline{1}_9$	1	ω	1	ω	1	ω^2	ω^2	ω^2	ω	1	1
$\underline{3}$	3	3ω	$3\omega^2$
$\underline{3}^*$	3	$3\omega^2$	3ω

Table 3: Character table of $\Delta(27)$. The first line indicates representatives of the different conjugacy classes. Zeroes in the character table are denoted by a dot . and ω is the third root of unity $\omega = e^{2\pi i/3}$. The arrows illustrate the generators of the outer automorphism group u_1 (blue) and u_2 (red).

mations which are connected to the unique non-trivial CP transformation by some group transformation, since the outer automorphism group is Z_2 .

4.4 $\Delta(27) \cong (Z_3 \times Z_3) \rtimes Z_3 \cong \text{SG}(27, 3)$

The group $\Delta(27) = \langle A, B | A^3 = B^3 = (AB)^3 = E \rangle$ ²² is another interesting group from the standpoint of CP violation. Its automorphism structure is quite involved. The centre of the group is isomorphic to Z_3 and generated by the group element $X = A^2BAB^2$ with $\text{conj}(X) = \text{id}$ and the inner automorphism group has the structure $Z_3 \times Z_3$. The outer automorphism group is generated by

$$u_1 : (A, B) \rightarrow (ABA^2, B^2AB), \quad u_2 : (A, B) \rightarrow (ABAB, B^2). \quad (4.41)$$

It is isomorphic to $\text{GL}(2, 3)$, i.e. the general linear group of 2×2 matrices over the field Z_3 . The multitude of outer automorphisms can be traced back to the various symmetries of the character table shown in table 3 that are due to the fact that there are so many one-dimensional representations. Together with the inner automorphisms these generators generate the full automorphism group, which is of order 432. In summary the automorphism structure presents itself as:

$$\begin{aligned} \text{Z}(\Delta(27)) &\cong Z_3 & \text{Aut}(\Delta(27)) &\cong (((Z_3 \times Z_3) \rtimes Q_8) \rtimes Z_3) \rtimes Z_2 \\ \text{Inn}(\Delta(27)) &\cong Z_3 \times Z_3 & \text{Out}(\Delta(27)) &\cong \text{GL}(2, 3). \end{aligned} \quad (4.42)$$

²² $\Delta(27)$ has been first used in the lepton sector in [53].

The outer automorphism u_1 acts on the representations as

$$\underline{\mathbf{1}}_2 \leftrightarrow \underline{\mathbf{1}}_4, \quad \underline{\mathbf{1}}_3 \leftrightarrow \underline{\mathbf{1}}_7, \quad \underline{\mathbf{1}}_6 \leftrightarrow \underline{\mathbf{1}}_8, \quad \underline{\mathbf{3}} \leftrightarrow \underline{\mathbf{3}}^* \quad (4.43)$$

where e.g. $\underline{\mathbf{1}}_2 \rightarrow \underline{\mathbf{1}}_4$ is to be read as $\rho_{\underline{\mathbf{1}}_4} = \rho_{\underline{\mathbf{1}}_2} \circ u_1$ etc., and the outer automorphism u_2 acts as

$$\underline{\mathbf{1}}_2 \rightarrow \underline{\mathbf{1}}_9 \rightarrow \underline{\mathbf{1}}_8 \rightarrow \underline{\mathbf{1}}_3 \rightarrow \underline{\mathbf{1}}_5 \rightarrow \underline{\mathbf{1}}_6 \rightarrow \underline{\mathbf{1}}_2 \quad (4.44)$$

From this it is trivial to determine the representations of the automorphisms for the one-dimensional representations. Let us therefore focus on the three dimensional representation $\underline{\mathbf{3}}$ generated by

$$\rho(A) = T_3, \quad \rho(B) = \text{diag}(1, \omega, \omega^2). \quad (4.45)$$

The two generators of the outer automorphism group act on $\phi \sim (\underline{\mathbf{3}}, \underline{\mathbf{3}}^*)$ as

$$U(u_1) = \begin{pmatrix} \tilde{U} & 0 \\ 0 & \tilde{U}^* \end{pmatrix} \quad \text{with} \quad \tilde{U} = \frac{1}{\sqrt{3}} \begin{pmatrix} \omega^2 & \omega & 1 \\ \omega & \omega^2 & 1 \\ 1 & 1 & 1 \end{pmatrix} \quad (4.46)$$

and

$$U(u_2) = \begin{pmatrix} 0 & \tilde{U} \\ \tilde{U}^* & 0 \end{pmatrix} \quad \text{with} \quad \tilde{U} = \begin{pmatrix} \omega^2 & 0 & 0 \\ 0 & 0 & \omega \\ 0 & \omega^2 & 0 \end{pmatrix}. \quad (4.47)$$

All automorphisms can be generated from the generators u_i by composition and the representation matrices $U(\text{aut})$ may be obtained with the help of eq. (2.17). We have therefore found a complete classification of possible CP transformations that may be implemented in a model based on $\Delta(27)$. There are 48 outer automorphisms generated by u_1 and u_2 that may in principle give physically distinct CP transformations with distinct physical implications, however as a model that is invariant under CP will also be invariant under CP^n it is sufficient to consider which subgroups of the automorphism groups is realised.

It is instructive to look at some of these subgroups in detail. Let us for example consider the CP transformation $\phi \rightarrow \phi^*$ or $U(h_1) = \mathbb{1}_3$ that corresponds to the outer automorphism $h_1 : (A, B) \rightarrow (A, B^2)$, which can be expressed in terms of the generators as $h_1 = u_1 \circ u_2^2 \circ u_1^{-1} \circ u_2 \circ u_1^{-1} \circ u_2^{-1} \circ u_1^{-1} \circ \text{conj}(A)^{-1} \circ u_1^{-1}$. This outer automorphism squares to one and therefore generates a Z_2 subgroup of the automorphism group. Contrary to the situation we have encountered before, where the outer automorphism group was a Z_2 , this is not the only solution. As a further example we may consider the Z_2 subgroup generated by $u_1 \circ u_2^2 \circ u_1^{-1} \circ u_2 \circ u_1^{-1} \circ u_2^{-2}$ with $h_2 : (A, B) \rightarrow (ABA, B)$ which according to eq. (2.17) is represented by

$$U(h_2) = \begin{pmatrix} \omega & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (4.48)$$

We will use this matrix later on. Let us now use this machinery to tackle a physical question, namely the so-called geometrical CP violation. 'Geometrical' CP-violation [4-8]

denotes the following: If one considers a triplet of Higgs doublets $H = (H_1, H_2, H_3) \sim \underline{\mathbf{3}}$ the only phase dependent term in the scalar potential is given by

$$I \equiv \sum_{i \neq j \neq k} (H_i^\dagger H_j)(H_i^\dagger H_k). \tag{4.49}$$

Let us now investigate how the term transforms under the two generators u_1 and u_2 of the outer automorphism group. We find

$$CP_{u_1}[I] = -\frac{1}{3}I^* + \frac{2}{3}I + \sum_i \frac{1}{3}(H_i^\dagger H_i)^2 + \sum_{i \neq j} (H_i^\dagger H_i)(H_j^\dagger H_j), \quad CP_{u_2}[I] = \omega^2 I \tag{4.50}$$

and we thus find the invariant combinations

$$CP_{u_1}[I - I^*] = I - I^* \quad CP_{u_2}[I] = I \tag{4.51}$$

Clearly invariance under u_1 requires further non-trivial relations among the other couplings in the scalar potential which do not depend on phases and thus do not concern us here.

Let us investigate the case where the theory is invariant under h_1 which corresponds to the 'usual' CP transformation $\phi \rightarrow \phi^*$ and forces the coupling λ_4 multiplying I to be real. For $\lambda_4 < 0$ one finds the global minimum

$$\langle H \rangle = \frac{v}{\sqrt{3}}(1, \omega, \omega^2) \tag{4.52}$$

and for $\lambda_4 > 0$ one finds

$$\langle H \rangle = \frac{v}{\sqrt{3}}(\omega^2, 1, 1). \tag{4.53}$$

Both VEV configurations correspond to generalised CP transformations $H \rightarrow UH^*$. For $\lambda_4 < 0$ it is for example given by $U = \rho(B^2)$ which is clearly part of $\Delta(27)$ and therefore up to an inner automorphism corresponds to h_1 . The phases of the VEVs thus do not imply spontaneous CP violation. For $\lambda_4 > 0$ the VEV configuration leaves the CP transformation corresponding to the outer automorphism h_2 given in eq. (4.48) invariant. However, there is something that is much harder to understand about this VEV configuration: the generalised CP symmetry corresponding to this configuration is not a symmetry of the Lagrangian. It would be a symmetry if the phase of λ_4 would be the same as ω , as $CP_{h_2}[I] = \omega I^*$. So here we are confronted with the puzzling situation where a VEV configuration is more symmetric than the original Lagrangian. This is also denoted as calculable phases.

This conundrum can be solved if there is a generalised CP trafo that is left invariant by the VEV and is compatible with λ_4 being real. Since we have a complete classification of all generalised CP transformations we can answer this question and indeed we find the CP transformation

$$\begin{pmatrix} H \\ H^* \end{pmatrix} = U \begin{pmatrix} H^* \\ H \end{pmatrix} \quad \text{with} \quad U = \begin{pmatrix} 0 & \tilde{U} \\ \tilde{U}^* & 0 \end{pmatrix}, \quad \tilde{U} = \begin{pmatrix} 0 & 0 & \omega^2 \\ 0 & 1 & 0 \\ \omega & 0 & 0 \end{pmatrix} \tag{4.54}$$

which represents the outer automorphism $u : (A, B) \rightarrow (AB^2AB, AB^2A^2)$ via eq. (2.8), where $u = u_2^3 \circ \text{conj}(A)$ and that gives

$$CP_u[\langle H \rangle] = \langle H \rangle \quad \text{for} \quad \langle H \rangle = \frac{v}{\sqrt{3}}(\omega^2, 1, 1), \quad CP_u[I] = I \quad (4.55)$$

Note that this CP transformation acts as $H \rightarrow \tilde{U}H$, which is not something you would naively expect, but it is an outer automorphism and therefore it is justified to call it a CP transformation. Furthermore, this becomes apparent when one looks at how the outer automorphism u acts on representations. It interchanges the one-dimensional representations

$$\underline{\mathbf{1}}_2 \leftrightarrow \underline{\mathbf{1}}_3, \quad \underline{\mathbf{1}}_5 \leftrightarrow \underline{\mathbf{1}}_9, \quad \underline{\mathbf{1}}_6 \leftrightarrow \underline{\mathbf{1}}_8, \quad (4.56)$$

making the "CP-character" of the transformation more apparent. An alternative independent explanation of geometric CP violation has been given in ref. [49].

4.5 $Z_9 \times Z_3 \cong \text{SG}(27, 4)$

Similarly to $\Delta(27)$, the group $Z_9 \times Z_3 = \text{SG}(27, 4) = \langle A, B | A^9 = B^3 = BAB^2A^2 = E \rangle$ ²³ has a more complicated automorphism group structure. The group is the semi-direct product of Z_9 generated by A (with $A^9 = E$) with Z_3 generated by B (with $B^3 = E$) defined by $BAB^{-1} = A^7$. The centre of the group is isomorphic to Z_3 and generated by A^3 . Hence, the inner automorphism group has the structure $Z_3 \times Z_3$. The outer automorphism group is generated by

$$\begin{aligned} u_1 : (A, B) &\rightarrow (AB, B^2A^6B^2A^3) \\ u_2 : (A, B) &\rightarrow (AB^4AB^4A^6, B^2A^6B^2A^6) \end{aligned} \quad (4.57)$$

and the structure of the automorphism group may be summarised as

$$\begin{aligned} \text{Z}(G) &\cong Z_3 & \text{Aut}(G) &\cong ((Z_3 \times Z_3) \times Z_3) \times Z_2 \\ \text{Inn}(G) &\cong Z_3 \times Z_3 & \text{Out}(G) &\cong S_3. \end{aligned} \quad (4.58)$$

There is a faithful three dimensional representation given by

$$\rho(A) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \quad \rho(B) = \begin{pmatrix} \omega^2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega \end{pmatrix}. \quad (4.59)$$

The generators of the outer automorphisms can be obtained in the same way as before and act on $(\underline{\mathbf{3}}, \underline{\mathbf{3}}^*)$ as

$$U(u_1) = \begin{pmatrix} 0 & \tilde{U} \\ \tilde{U}^* & 0 \end{pmatrix} \quad \text{with} \quad \tilde{U} = \text{diag}(1, 1, \omega^2) \quad (4.60)$$

and

$$U(u_2) = \begin{pmatrix} \tilde{U} & 0 \\ 0 & \tilde{U}^* \end{pmatrix} \quad \text{with} \quad \tilde{U} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}. \quad (4.61)$$

²³The possibility of having $Z_9 \times Z_3$ as a flavour group in the lepton sector has been first mentioned in ref. [54].

4.6 $Q_8 \times A_4 \cong \text{SG}(96, 204)$

Let us also consider our favourite group, $Q_8 \times A_4$ [55, 56] generated by S, T, X with

$$S^2 = T^3 = X^4 = (ST)^3 = SX SX^3 = T^2 X (T^2 X^3)^2 = (STX^3 T^2)^2 = E \quad (4.62)$$

the smallest group that may realise the VEV alignment. Its centre is given by $Z(Q_8 \times A_4) = \{E, X^2\} \cong Z_2$ and its outer automorphism group is generated by

$$\begin{aligned} h_4 : (S, T, X) &\rightarrow (S, T^2, SX), & h_5 : (S, T, X) &\rightarrow (S, T^2, X^3), \\ h_6 : (S, T, X) &\rightarrow (ST^2 STX^3, T, T^2 XT). \end{aligned} \quad (4.63)$$

These generators act on the character table and representations in the way indicated in table 5. Together with the inner automorphisms, the automorphism group is of order 576 and its structure may be summarised as:

$$\begin{aligned} Z(Q_8 \times A_4) &\cong Z_2 & \text{Aut}(Q_8 \times A_4) &\cong ((A_4 \times A_4) \rtimes Z_2) \rtimes Z_2 & (4.64) \\ \text{Inn}(Q_8 \times A_4) &\cong Z_2^4 \rtimes Z_3 & \text{Out}(Q_8 \times A_4) &\cong D_{12}. \end{aligned}$$

Let us discuss how the generators of the automorphism group may be represented upon the vector

$$\phi = \begin{pmatrix} \varphi_C \\ \varphi_C^* \end{pmatrix} \quad (4.65)$$

with $\varphi_C \sim \underline{4}_2$ upon which the group generators act as

$$\rho(S) = \begin{pmatrix} S_4 & 0 \\ 0 & S_4 \end{pmatrix}, \quad \rho(T) = \begin{pmatrix} \omega^2 T_4 & 0 \\ 0 & \omega T_4 \end{pmatrix}, \quad \text{and} \quad \rho(X) = \begin{pmatrix} X_4 & 0 \\ 0 & X_4 \end{pmatrix} \quad (4.66)$$

with

$$S_4 \equiv \sigma_3 \otimes \sigma_1, \quad T_4 \equiv \text{diag}(T_3, 1), \quad X_4 \equiv -i\sigma_2 \otimes \sigma_3 \quad (4.67)$$

and $\rho(S, X)^* = \rho(S, X)$ but $\rho(T)^* \notin \text{Im}\rho$. One solution to eq. (2.6) is the analogue of the A_4 case, $U = \text{diag}(U_4, U_4)$ with $U_4 = \text{diag}(\tilde{U}_3 \equiv T_3 U_3 T_3^{-1}, 1)$. The matrix U_3 has been defined in eq. (4.11). This generator acts on the generators of the group as

$$U\rho(S)^*U^{-1} = \rho(S), \quad U\rho(T)^*U^{-1} = \rho(T^2), \quad U\rho(X)^*U^{-1} = \rho(SX) \quad (4.68)$$

and therefore represents the automorphism h_4 . Before discussing other solutions to eq. (2.6), let us demonstrate how this outer automorphism can be represented for the other representations. For the representation $\underline{4}_1$ we find $U = U_4$. For the one-dimensional representations we have $U = 1$.

Clearly the relation (4.68) cannot be fulfilled by $\underline{3}_1$ as $\rho(X) = \mathbb{1}_3$

$$1 = U\rho(X)U^{-1} = \rho(SX) = S_3 \quad (4.69)$$

	S	T	X	FS	Z(G)
$\underline{\mathbf{1}}_1$	1	1	1	1	1
$\underline{\mathbf{1}}_2$	1	ω	1	0	1
$\underline{\mathbf{1}}_3$	1	ω^2	1	0	1
$\underline{\mathbf{4}}_1$	S_4	T_4	X_4	1	-1
$\underline{\mathbf{4}}_2$	S_4	$\omega^2 T_4$	X_4	0	-1
$\underline{\mathbf{4}}_3$	S_4	ωT_4	X_4	0	-1

	S	T	X	FS	Z(G)
$\underline{\mathbf{3}}_1$	S_3	T_3	$\mathbb{1}_3$	1	1
$\underline{\mathbf{3}}_2$	$T_3 S_3 T_3^2$	T_3	S_3	1	1
$\underline{\mathbf{3}}_3$	$T_3 S_3 T_3^2$	T_3	$T_3^2 S_3 T_3$	1	1
$\underline{\mathbf{3}}_4$	$\mathbb{1}_3$	T_3	$T_3 S_3 T_3^2$	1	1
$\underline{\mathbf{3}}_5$	$T_3^2 S_3 T_3$	T_3	$T_3^2 S_3 T_3$	1	1

Table 4: Representations of $Q_8 \times A_4$ in the chosen basis. The one-dimensional representations and the first three-dimensional one are the unfaithful A_4 representations (therefore $X = \mathbb{1}$), which the leptons are assigned to in refs. [55, 56]. The representation $\underline{\mathbf{4}}_1$ is used to break A_4 in the neutrino sector. Note that this representation is double valued, i.e. $Z(G) = X^2 = -\mathbb{1}$. FS is the Frobenius-Schur indicator $\frac{1}{|G|} \sum_{g \in G} \chi(g^2)$ that takes the values 1 for real, 0 for complex or -1 for pseudo-real representations, respectively. The matrices S_3 and T_3 have been defined in eq. (4.7).

for any U . The representation $\underline{\mathbf{3}}_1$ is rather part of a larger representation that also includes $\underline{\mathbf{3}}_5$:²⁴

$$S = \text{diag}(S_3, T_3^2 S_3 T_3), \quad T = \text{diag}(T_3, T_3^2), \quad X = \text{diag}(\mathbb{1}_3, T_3^2 S_3 T_3), \quad U = \begin{pmatrix} 0 & T_3 \\ T_3^2 & 0 \end{pmatrix}. \quad (4.70)$$

The real representations $\underline{\mathbf{3}}_{2,3,4}$ can be extended to representations of the CP-extended group by $U = \tilde{U}_3$. We have therefore seen that a CP transformation as defined in (2.6) can only be realised if both $\underline{\mathbf{3}}_1$ and $\underline{\mathbf{3}}_5$ are present in the Lagrangian, i.e. the condition of CP conservation requires non-trivial relations among real representations of the group, something one would not immediately suspect. To summarise a consistent definition of CP acts as

$$\underline{\mathbf{4}}_i \rightarrow U_4 \underline{\mathbf{4}}_i^* \quad \underline{\mathbf{3}}_i \rightarrow \tilde{U}_3 \underline{\mathbf{3}}_{f(i)}^* \quad \underline{\mathbf{1}}_i \rightarrow \underline{\mathbf{1}}_i^* \quad (4.71)$$

with $f : \{1, 2, 3, 4, 5\} \rightarrow \{5, 2, 3, 4, 1\}$.

The natural question is now if it is possible to have outer automorphisms of the group that act as CP in the sense that they interchange the complex representations $\underline{\mathbf{1}}_{2,3}$ and $\underline{\mathbf{4}}_{2,3}$ but transform the real representations only within themselves. This question can be answered using the explicit form of the generators of eq. (4.63).

An outer automorphism swaps conjugacy classes and representations in such a way as to leave the character table 5 invariant. For illustration look at the automorphism h_4 (4.68). It acts on the conjugacy classes as

$$G \cdot T \leftrightarrow G \cdot T^2, \quad G \cdot XT \leftrightarrow G \cdot SXT^2, \quad G \cdot X \leftrightarrow G \cdot SX \quad (4.72)$$

²⁴For $\underline{\mathbf{3}}_5$ we have $\rho(S) = \rho(X)$ and therefore eq. (4.68) would imply $\rho(S) = \rho(X) = \mathbb{1}_3$.

	E	T	SYX	SY	X^2	T^2	XT	S	SX	X	SXT^2
$\underline{1}_1$	1	1	1	1	1	1	1	1	1	1	1
$\underline{1}_2$	1	ω	1	1	1	ω^2	ω	1	1	1	ω^2
$\underline{1}_3$	1	ω^2	1	1	1	ω	ω^2	1	1	1	ω
$\underline{3}_1$	3	.	-1	-1	3	.	.	-1	-1	3	.
$\underline{3}_2$	3	.	3	-1	3	.	.	-1	-1	-1	.
$\underline{3}_3$	3	.	-1	3	3	.	.	-1	-1	-1	.
$\underline{3}_4$	3	.	-1	-1	3	.	.	3	-1	-1	.
$\underline{3}_5$	3	.	-1	-1	3	.	.	-1	3	-1	.
$\underline{4}_1$	4	1	.	.	-4	1	-1	.	.	.	-1
$\underline{4}_2$	4	ω^2	.	.	-4	ω	$-\omega^2$.	.	.	$-\omega$
$\underline{4}_3$	4	ω	.	.	-4	ω^2	$-\omega$.	.	.	$-\omega^2$

Table 5: Character table of $Q_8 \times A_4$. The first line indicates representatives of the different conjugacy classes. Zeroes in the character table are denoted by a dot . and ω is the third root of unity $\omega = e^{2\pi i/3}$ and $Y = T^2XT$. The arrows illustrate the generators of the outer automorphism group h_4 (blue), h_5 (red), h_6 (green).

where $G \cdot T \equiv \{gTg^{-1} : g \in G\}$, leaving all other conjugacy classes invariant. To obtain a symmetry of the character table one therefore needs to interchange the representations

$$\underline{1}_2 \leftrightarrow \underline{1}_3, \quad \underline{4}_2 \leftrightarrow \underline{4}_3, \quad \underline{3}_1 \leftrightarrow \underline{3}_5. \tag{4.73}$$

If we want to have a symmetry of the character table without interchanging any real representations that still acts as CP, we therefore have to have an automorphism that realises

$$G \cdot T \leftrightarrow G \cdot T^2, \quad G \cdot XT \leftrightarrow G \cdot SXT^2 \tag{4.74}$$

while keeping all other conjugacy classes invariant. No such automorphism exists, as can be inferred from eq. (4.63).²⁵

However, if we relax the condition to the point where we only demand that the representation $\underline{3}_1$ transforms into itself we have to search for outer automorphisms that realise

$$G \cdot T \leftrightarrow G \cdot T^2, \quad G \cdot XT \leftrightarrow G \cdot SXT^2 \quad G \cdot X \leftrightarrow G \cdot X. \tag{4.75}$$

Indeed there is a automorphism that realises this: $h_5 : (S, T, X) \rightarrow (S, T^2, X^3)$. An explicit matrix representation for representation $\underline{4}_1$ is given by

$$U_4(h_5) = \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ -1 & -1 & -1 & -1 \end{pmatrix} \tag{4.76}$$

²⁵It is convenient to use the computer algebra system GAP [31].

and for the representation $\underline{\mathbf{3}}_1$ we find $U = U_3$.

Having found a consistent CP transformation for a theory that contains only the representations $\underline{\mathbf{3}}_1$, $\underline{\mathbf{4}}_1$ and $\underline{\mathbf{1}}_i$ we can now ask ourselves the question that lead us to this study of generalised CP transformations. Namely if we take the flavon content of ref. [55] and promote all the fields to electroweak (EW) doublets $\chi \sim \underline{\mathbf{3}}_1$ and $\phi \sim \underline{\mathbf{4}}_1$, there is a purely imaginary coupling²⁶

$$\lambda \left(\chi^\dagger \chi \right)_{\underline{\mathbf{3}}_1, \mathbf{S}} \cdot \left(\phi^\dagger \phi \right)_{\underline{\mathbf{3}}_1} + \text{h.c.} \tag{4.77}$$

which breaks the accidental symmetry needed for vacuum alignment [55]. To forbid this imaginary coupling one might think that a CP symmetry can be invoked. However, the consistent CP transformation corresponding to h_5 (which is unique up to inner automorphisms) under which the EW doublets ϕ and χ transform as

$$\phi_i \rightarrow U_4 \phi^*, \quad \chi \rightarrow U_3 \chi^* \tag{4.78}$$

leaves the operator in eq. (4.77) invariant, even though it is purely imaginary.

For completeness we also give a representation of h_6

$$U_4(h_6) = \frac{1}{2} \begin{pmatrix} 1 & -1 & -1 & -1 \\ -1 & 1 & -1 & -1 \\ -1 & -1 & 1 & -1 \\ 1 & 1 & 1 & -1 \end{pmatrix} \tag{4.79}$$

from which all the other representation matrices can be derived using the Clebsch-Gordon coefficients.

4.7 $S_4 \cong (\mathbf{Z}_2 \times \mathbf{Z}_2) \rtimes S_3 \cong \mathbf{SG}(24, 12)$

There is a complete classification of automorphism groups for the symmetric group S_n , which we summarise in table 6. Since $\text{Out}(S_n) = \mathbf{Z}_1$ for $n \neq 2, 6$, there is no non-trivial generalised CP transformation, especially S_3 and S_4 , which have been introduced in [57, 58] and used in models explaining the leptonic mixing structure, do not allow for a non-trivial generalised CP transformation. The recently discussed generalised CP transformation in ref. [28] is an inner automorphism of S_4 , similarly the generalised CP in the framework of S_3 discussed in [59]. Obviously, it is always possible to apply a group transformation at the same time as a CP transformation. We therefore do not discuss S_3 or S_4 in more detail.

4.8 $T_7 \cong \mathbf{Z}_7 \rtimes \mathbf{Z}_3 \cong \mathbf{SG}(21, 1)$

The group $T_7 \cong \mathbf{Z}_7 \rtimes \mathbf{Z}_3 \cong \mathbf{SG}(21, 1) = \langle A, B | A^7 = B^3 = BAB^{-1}A^5 = E \rangle$ has been first used in particle physics in ref. [60]. In the basis used in [61], the generators A and B are given by

$$\rho(A) = \text{diag}(\eta, \eta^2, \eta^4) \quad \rho(B) = T_3 \tag{4.80}$$

²⁶ $(\chi^\dagger \chi)_{\underline{\mathbf{3}}_1, \mathbf{S}}$ is real and $(\phi^\dagger \phi)_{\underline{\mathbf{3}}_1}$ is purely imaginary, therefore the coupling has to be purely imaginary.

	$Z(S_n)$	$\text{Aut}(S_n)$	$\text{Inn}(S_n)$	$\text{Out}(S_n)$
$n \neq 2, 6$	Z_1	S_n	S_n	Z_1
$n = 2$	Z_2	Z_1	Z_1	Z_1
$n = 6$	Z_1	$S_6 \rtimes Z_2$	S_6	Z_2

Table 6: Group structure of the symmetric group S_n

for $\mathbf{3}_1$ with $\eta = e^{2\pi i/7}$. T_7 has a trivial centre and therefore the inner automorphism group $\text{Inn}(T_7)$ is isomorphic to T_7 itself. However, since $\rho(A)^* = \rho(A^6) \in \text{Im}\rho$ and $\rho(B)^* = \rho(B) \in \text{Im}\rho$, the outer automorphism group is non-trivial. Its generator $u : (A, B) \rightarrow (A^6, B)$ is thus represented by the identity matrix on the three dimensional representation and this basis is thus a CP basis. Concluding the structure of the automorphism group is described by

$$\begin{aligned} Z(T_7) &\cong Z_1 & \text{Aut}(T_7) &\cong \text{SG}(42, 2) & (4.81) \\ \text{Inn}(T_7) &\cong T_7 & \text{Out}(T_7) &\cong Z_2 . \end{aligned}$$

The outer automorphism exchanges the three-dimensional representations, while leaving the one-dimensional ones fixed, i.e.

$$\mathbf{1}_2 \rightarrow \mathbf{1}_2, \quad \mathbf{1}_3 \rightarrow \mathbf{1}_3 \quad \text{and} \quad \mathbf{3} \leftrightarrow \mathbf{3}^* . \quad (4.82)$$

4.9 $\Delta(108) \cong \text{SG}(108, 22)$ (or $\Delta(216) \cong \text{SG}(216, 95)$)

Recently [9], CP violation has been discussed in the context of $\Delta(108) = \Delta(3 \times 6^2)$,²⁷ which may be represented by a faithful three-dimensional representation as

$$\rho(\mathcal{S}) = S_3, \quad \rho(\mathcal{T}) = T_3 \quad \rho(\mathcal{T}') = \text{diag}(1, \omega, \omega^2) . \quad (4.83)$$

The model possesses an accidental $\mu - \tau$ exchange symmetry, which is generated by U_3 .²⁸ Including this generator $U = U_3$, the group becomes $\Delta(6 \times 6^2)$. A generalised CP transformation was defined on the faithful representation ℓ_R as

$$\ell_R \rightarrow iU_3 \ell_R^* , \quad (4.84)$$

where we have suppressed the Lorentz structure. This is equivalent to the automorphism $u : (\mathcal{S}, \mathcal{T}, \mathcal{T}') \rightarrow (\mathcal{S}, \mathcal{T}^2, \mathcal{T}')$, which is outer in $\Delta(3 \times 6^2)$ and inner in $\Delta(6 \times 6^2)$. In ref. [9] this has been consistently applied to all non-faithful representations which they consider.

Let us comment on the origin of maximal CP violation in their model, which seems to be in conflict with our general statement that there can be no CP violation. It is related to the breaking of the flavour symmetry in their model. One of the scalar fields breaking the flavour symmetry is the scalar ϕ transforming as

$$\rho(\mathcal{S}) = S_3, \quad \rho(\mathcal{T}) = T_3 \quad \rho(\mathcal{T}') = \mathbb{1}_3 , \quad (4.85)$$

²⁷ $\Delta(108)$ has been first used in the lepton sector in ref. [62]. There is a comprehensive study of $\Delta(3n^2)$ [63] and $\Delta(6n^2)$ [64] groups in the context of flavour symmetries.

²⁸The matrices S_3 , T_3 and U_3 have been defined in eq. (4.7) and eq. (4.11).

and thus transforms only under the subgroup $\langle \mathcal{S}, \mathcal{T} \rangle \cong A_4$ with the CP transformation $\phi \rightarrow U_3 \phi^*$. CP conservation would therefore require $v_2 = v_3^*$. However, they have to assume a large hierarchy in the VEVs of ϕ in order to accommodate the hierarchy in the charged lepton sector, which is given by $m_e : m_\mu : m_\tau = v_1 : v_2 : v_3$. Hence, the requirement $|v_2|/|v_3| = m_\mu/m_\tau \ll 1$ is the necessary ingredient for maximal CP violation in the model.

5 Conclusions and outlook

We have given consistency conditions for the definition of CP in theories with discrete flavour symmetries that have sometimes been overlooked in the literature. We have shown that every generalised CP transformation furnishes a representation of an outer automorphism and that generalised CP invariance implies vanishing CP phases. We have applied these ideas to popular flavour groups with three-dimensional representations and group order smaller than 31.²⁹ In particular, we have shown that there is one unique non-trivial CP transformation (up to group transformations) for the group T' , which we applied to the models discussed in ref. [10, 11]. We show that this CP is spontaneously broken by the VEVs of the doublets. The claimed geometric CP-violation in ref. [10] can only be viewed as an arbitrary basis-dependent explicit breaking of CP. In the case of $\Delta(27)$ we have shown that the so-called geometric phases may be viewed as the result of an accidental generalised CP transformation of the scalar potential. Finally, we showed in the case of A_4 that the phase of $(\chi^\dagger \chi)_{\mathbf{3}_1} (\chi^\dagger \chi)_{\mathbf{3}_1}$ in the potential of a single triplet does not break CP, which has also been independently shown in ref. [47–49]. This clarifies the recent observation that CP conserving solutions result from seemingly explicitly CP-breaking potentials [42].

The (outer) automorphism structure of small groups is very rich and it stands to wonder if not more physics might be hidden in there. We may speculate about this possibility in the following. S_4 is the smallest group that can really generate TBM (with all the caveats involved) and it is isomorphic to the automorphism group of A_4 . Maybe the accidental symmetry that makes A_4 to S_4 on the level of mass matrices is connected to this fact. This would open an interesting avenue for model building: interesting mixing patterns can be obtained from $\Delta(6n^2)$ but it is quite unappealing to start from such large groups, it might be nicer to start from smaller groups and obtain the accidental symmetry from the larger automorphism group in the same way as in A_4 models. As an example how complicated structures can arise from simpler ones, look at the automorphism group of $\Delta(27)$, which is of order 432. The smallest group whose automorphism group contains $\Delta(96)$ is given by $(Z_4 \times Z_4) \rtimes Z_2 \cong \text{SG}(32, 34)$. Further investigation of these ideas is left for future work.

Note added. While this work was being finalised, a related work [28] addressing CP in the context of discrete flavour symmetries appeared on the arxiv. We both give a general definition and discussion of generalised CP symmetries. Our work differs from [28] as

²⁹For completeness, we mention the group $A_4 \times Z_2 \cong \text{SG}(24, 13)$, which we did not discuss in detail. It has been mentioned in the survey of ref. [54]. Its automorphism group structure is directly inherited from A_4 with the addition that it has a non-trivial centre $Z(Z_2)$.

follows. They consider the physical implications for the lepton mixing parameters of a remnant CP symmetry in the neutrino sector. In particular, they discuss the groups S_4 as well as A_4 in more detail. We, on the other hand, emphasise the relation of generalised CP transformations to the automorphism group and especially the outer automorphism group. We perform a systematic study of all generalised CP transformations for all groups with a three-dimensional irreducible representation of order less than 31. In particular, we discuss the "calculable phases" in models based on $\Delta(27)$ and interpret them in terms of an accidental generalised CP transformation as well as comment on the recent claims of geometric CP violation in the context of T' models.

Acknowledgments

We would like to thank I. de Medeiros Varzielas for comments on the manuscript. Furthermore, we thank C. Hagedorn, A. Meroni, S. Petcov and especially M. Spinrath for discussions on CP violation in T' models. We thank the referee for his detailed comments. M.S. would like to acknowledge MPI für Kernphysik, where a part of this work has been done, for hospitality of its staff and the generous support. M.H. acknowledges support by the International Max Planck Research School for Precision Tests of Fundamental Symmetries. This work was supported in part by the Australian Research Council.

References

- [1] DOUBLE-CHOOZ collaboration, Y. Abe et al., *Indication for the disappearance of reactor electron antineutrinos in the Double CHOOZ experiment*, *Phys. Rev. Lett.* **108** (2012) 131801 [[arXiv:1112.6353](#)] [[INSPIRE](#)].
- [2] DAYA-BAY collaboration, F. An et al., *Observation of electron-antineutrino disappearance at Daya Bay*, *Phys. Rev. Lett.* **108** (2012) 171803 [[arXiv:1203.1669](#)] [[INSPIRE](#)].
- [3] RENO collaboration, J. Ahn et al., *Observation of Reactor Electron Antineutrino Disappearance in the RENO Experiment*, *Phys. Rev. Lett.* **108** (2012) 191802 [[arXiv:1204.0626](#)] [[INSPIRE](#)].
- [4] G. Branco, J. Gerard and W. Grimus, *GEOMETRICAL T VIOLATION*, *Phys. Lett.* **B 136** (1984) 383 [[INSPIRE](#)].
- [5] I. de Medeiros Varzielas and D. Emmanuel-Costa, *Geometrical CP-violation*, *Phys. Rev.* **D 84** (2011) 117901 [[arXiv:1106.5477](#)] [[INSPIRE](#)].
- [6] I. de Medeiros Varzielas, D. Emmanuel-Costa and P. Leser, *Geometrical CP-violation from Non-Renormalisable Scalar Potentials*, *Phys. Lett.* **B 716** (2012) 193 [[arXiv:1204.3633](#)] [[INSPIRE](#)].
- [7] I. de Medeiros Varzielas, *Geometrical CP-violation in multi-Higgs models*, *JHEP* **08** (2012) 055 [[arXiv:1205.3780](#)] [[INSPIRE](#)].
- [8] G. Bhattacharyya, I. de Medeiros Varzielas and P. Leser, *A common origin of fermion mixing and geometrical CP-violation and its test through Higgs physics at the LHC*, *Phys. Rev. Lett.* **109** (2012) 241603 [[arXiv:1210.0545](#)] [[INSPIRE](#)].

- [9] P. Ferreira, W. Grimus, L. Lavoura and P. Ludl, *Maximal CP-violation in Lepton Mixing from a Model with Delta(27) flavour Symmetry*, *JHEP* **09** (2012) 128 [[arXiv:1206.7072](#)] [[INSPIRE](#)].
- [10] M.-C. Chen and K. Mahanthappa, *Group Theoretical Origin of CP-violation*, *Phys. Lett. B* **681** (2009) 444 [[arXiv:0904.1721](#)] [[INSPIRE](#)].
- [11] A. Meroni, S. Petcov and M. Spinrath, *A SUSY SU(5) \times T' Unified Model of Flavour with large θ_{13}* , *Phys. Rev. D* **86** (2012) 113003 [[arXiv:1205.5241](#)] [[INSPIRE](#)].
- [12] T. Lee, *A Theory of Spontaneous T Violation*, *Phys. Rev. D* **8** (1973) 1226 [[INSPIRE](#)].
- [13] G.C. Branco, *Spontaneous CP-violation in Theories with More Than Four Quarks*, *Phys. Rev. Lett.* **44** (1980) 504 [[INSPIRE](#)].
- [14] H.E. Haber and Z. Surujon, *A Group-theoretic Condition for Spontaneous CP-violation*, *Phys. Rev. D* **86** (2012) 075007 [[arXiv:1201.1730](#)] [[INSPIRE](#)].
- [15] G. Ecker, W. Grimus and W. Konetschny, *Quark mass matrices in left-right symmetric gauge theories*, *Nucl. Phys. B* **191** (1981) 465 [[INSPIRE](#)].
- [16] G. Ecker, W. Grimus and H. Neufeld, *Spontaneous CP-violation in left-right symmetric gauge theories*, *Nucl. Phys. B* **247** (1984) 70 [[INSPIRE](#)].
- [17] H. Neufeld, W. Grimus and G. Ecker, *Generalized CP invariance, neutral flavor conservation and the structure of the mixing matrix*, *Int. J. Mod. Phys. A* **3** (1988) 603 [[INSPIRE](#)].
- [18] W. Grimus and M. Rebelo, *Automorphisms in gauge theories and the definition of CP and P*, *Phys. Rept.* **281** (1997) 239 [[hep-ph/9506272](#)] [[INSPIRE](#)].
- [19] P. Harrison and W. Scott, *$\mu - \tau$ reflection symmetry in lepton mixing and neutrino oscillations*, *Phys. Lett. B* **547** (2002) 219 [[hep-ph/0210197](#)] [[INSPIRE](#)].
- [20] P. Harrison and W. Scott, *Symmetries and generalizations of tri - bimaximal neutrino mixing*, *Phys. Lett. B* **535** (2002) 163 [[hep-ph/0203209](#)] [[INSPIRE](#)].
- [21] P. Harrison and W. Scott, *The Simplest neutrino mass matrix*, *Phys. Lett. B* **594** (2004) 324 [[hep-ph/0403278](#)] [[INSPIRE](#)].
- [22] W. Grimus and L. Lavoura, *A Nonstandard CP transformation leading to maximal atmospheric neutrino mixing*, *Phys. Lett. B* **579** (2004) 113 [[hep-ph/0305309](#)] [[INSPIRE](#)].
- [23] Y. Farzan and A.Y. Smirnov, *Leptonic CP-violation: Zero, maximal or between the two extremes*, *JHEP* **01** (2007) 059 [[hep-ph/0610337](#)] [[INSPIRE](#)].
- [24] A.S. Joshipura, B.P. Kodrani and K.M. Patel, *Fermion Masses and Mixings in a mu-tau symmetric SO(10)*, *Phys. Rev. D* **79** (2009) 115017 [[arXiv:0903.2161](#)] [[INSPIRE](#)].
- [25] W. Grimus and L. Lavoura, *mu-tau Interchange symmetry and lepton mixing*, [arXiv:1207.1678](#) [[INSPIRE](#)].
- [26] R. Mohapatra and C. Nishi, *S_4 Flavored CP Symmetry for Neutrinos*, *Phys. Rev. D* **86** (2012) 073007 [[arXiv:1208.2875](#)] [[INSPIRE](#)].
- [27] R. Krishnan, P. Harrison and W. Scott, *Simplest Neutrino Mixing from S_4 Symmetry*, [arXiv:1211.2000](#) [[INSPIRE](#)].
- [28] F. Feruglio, C. Hagedorn and R. Ziegler, *Lepton Mixing Parameters from Discrete and CP Symmetries*, [arXiv:1211.5560](#) [[INSPIRE](#)].

- [29] K.S. Babu and J. Kubo, *Dihedral families of quarks, leptons and Higgses*, *Phys. Rev. D* **71** (2005) 056006 [[hep-ph/0411226](#)] [[INSPIRE](#)].
- [30] K. Babu, K. Kawashima and J. Kubo, *Variations on the Supersymmetric Q_6 Model of Flavor*, *Phys. Rev. D* **83** (2011) 095008 [[arXiv:1103.1664](#)] [[INSPIRE](#)].
- [31] GAP, *GAP – Groups, Algorithms, and Programming, Version 4.5.5*, The GAP Group (2012), <http://www.gap-system.org>.
- [32] H.U.Besche, B.Eick, and E.O'Brien, *SmallGroups - library of all 'small' groups, GAP package, Version included in GAP 4.5.5*, The GAP Group (2002), <http://www.gap-system.org/Packages/sgl.html>.
- [33] W. Fairbairn and T. Fulton, *Some comments on finite subgroups of $SU(3)$* , *J. Math. Phys.* **23** (1982) 1747 [[INSPIRE](#)].
- [34] W. Grimus and P.O. Ludl, *Finite flavour groups of fermions*, *J. Phys. A* **45** (2012) 233001 [[arXiv:1110.6376](#)] [[INSPIRE](#)].
- [35] H. Ishimori et al., *An introduction to non-Abelian discrete symmetries for particle physicists*, *Lect. Notes Phys.* **858** (2012) 1.
- [36] G. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1 [[arXiv:1106.0034](#)] [[INSPIRE](#)].
- [37] G. Ecker, W. Grimus and H. Neufeld, *A standard form for generalized CP transformations*, *J. Phys. A* **20** (1987) L807 [[INSPIRE](#)].
- [38] M. Gronau, A. Kfir and R. Loewy, *Basis independent tests of CP-violation in fermion mass matrices*, *Phys. Rev. Lett.* **56** (1986) 1538 [[INSPIRE](#)].
- [39] J. Bernabeu, G. Branco and M. Gronau, *CP restrictions on quark mass matrices*, *Phys. Lett. B* **169** (1986) 243 [[INSPIRE](#)].
- [40] G. Branco, L. Lavoura and M. Rebelo, *Majorana neutrinos and CP-violation in the leptonic sector*, *Phys. Lett. B* **180** (1986) 264 [[INSPIRE](#)].
- [41] E. Ma and G. Rajasekaran, *Softly broken A_4 symmetry for nearly degenerate neutrino masses*, *Phys. Rev. D* **64** (2001) 113012 [[hep-ph/0106291](#)] [[INSPIRE](#)].
- [42] R. de Adelhart Toorop, F. Bazzocchi, L. Merlo and A. Paris, *Constraining Flavour Symmetries At The EW Scale I: The A_4 Higgs Potential*, *JHEP* **03** (2011) 035 [Erratum *ibid.* **1301** (2013) 098] [[arXiv:1012.1791](#)] [[INSPIRE](#)].
- [43] P. Ferreira and L. Lavoura, *Seesaw Neutrino Masses from an A_4 Model with Two Equal Vacuum expectation values*, [arXiv:1111.5859](#) [[INSPIRE](#)].
- [44] A. Machado, J. Montero and V. Pleitez, *Three-Higgs-doublet model with A_4 symmetry*, *Phys. Lett. B* **697** (2011) 318 [[arXiv:1011.5855](#)] [[INSPIRE](#)].
- [45] R. de Adelhart Toorop, F. Bazzocchi, L. Merlo and A. Paris, *Constraining Flavour Symmetries At The EW Scale I: The A_4 Higgs Potential*, *JHEP* **03** (2011) 035 [Erratum *ibid.* **1301** (2013) 098] [[arXiv:1012.1791](#)] [[INSPIRE](#)].
- [46] G. Altarelli and F. Feruglio, *Tri-bimaximal neutrino mixing, A_4 and the modular symmetry*, *Nucl. Phys. B* **741** (2006) 215 [[hep-ph/0512103](#)] [[INSPIRE](#)].
- [47] R. de Adelhart Toorop, *A Flavour of family symmetries in a family of flavour models*, PhD. thesis, Nikhef (2012).

- [48] I. Ivanov and E. Vdovin, *Classification of finite reparametrization symmetry groups in the three-Higgs-doublet model*, *Eur. Phys. J. C* **73** (2013) 2309 [[arXiv:1210.6553](#)] [[INSPIRE](#)].
- [49] A. Degee, I.P. Ivanov and V. Keus, *Geometric minimization of highly symmetric potentials*, *JHEP* **02** (2013) 125 [[arXiv:1211.4989](#)] [[INSPIRE](#)].
- [50] P. Frampton and T. Kephart, *Simple nonAbelian finite flavor groups and fermion masses*, *Int. J. Mod. Phys. A* **10** (1995) 4689 [[hep-ph/9409330](#)] [[INSPIRE](#)].
- [51] F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, *Tri-bimaximal Neutrino Mixing and Quark Masses from a Discrete Flavour Symmetry*, *Nucl. Phys. B* **775** (2007) 120 [Erratum *ibid.* **836** (2010) 127-128] [[hep-ph/0702194](#)] [[INSPIRE](#)].
- [52] S. Antusch, S.F. King, C. Luhn and M. Spinrath, *Right Unitarity Triangles and Tri-Bimaximal Mixing from Discrete Symmetries and Unification*, *Nucl. Phys. B* **850** (2011) 477 [[arXiv:1103.5930](#)] [[INSPIRE](#)].
- [53] I. de Medeiros Varzielas, S. King and G. Ross, *Neutrino tri-bi-maximal mixing from a non-Abelian discrete family symmetry*, *Phys. Lett. B* **648** (2007) 201 [[hep-ph/0607045](#)] [[INSPIRE](#)].
- [54] K.M. Parattu and A. Wingerter, *Tribimaximal Mixing From Small Groups*, *Phys. Rev. D* **84** (2011) 013011 [[arXiv:1012.2842](#)] [[INSPIRE](#)].
- [55] M. Holthausen and M.A. Schmidt, *Natural Vacuum Alignment from Group Theory: The Minimal Case*, *JHEP* **01** (2012) 126 [[arXiv:1111.1730](#)] [[INSPIRE](#)].
- [56] M. Holthausen, M. Lindner and M.A. Schmidt, *Lepton flavor at the electroweak scale: A complete A_4 model*, *Phys. Rev. D* **87** (2013) 033006 [[arXiv:1211.5143](#)] [[INSPIRE](#)].
- [57] S. Pakvasa and H. Sugawara, *Discrete Symmetry and Cabibbo Angle*, *Phys. Lett. B* **73** (1978) 61 [[INSPIRE](#)].
- [58] S. Pakvasa and H. Sugawara, *Mass of the t Quark in $SU(2) \times U(1)$* , *Phys. Lett. B* **82** (1979) 105 [[INSPIRE](#)].
- [59] S.L. Adler, *Model for particle masses, flavor mixing and CP-violation, based on spontaneously broken discrete chiral symmetry as the origin of families*, *Phys. Rev. D* **59** (1999) 015012 [Erratum *ibid.* **D 59** (1999) 099902] [[hep-ph/9806518](#)] [[INSPIRE](#)].
- [60] C. Luhn, S. Nasri and P. Ramond, *Tri-bimaximal neutrino mixing and the family symmetry semidirect product of $Z(7)$ and $Z(3)$* , *Phys. Lett. B* **652** (2007) 27 [[arXiv:0706.2341](#)] [[INSPIRE](#)].
- [61] C. Hagedorn, M.A. Schmidt and A.Y. Smirnov, *Lepton Mixing and Cancellation of the Dirac Mass Hierarchy in $SO(10)$ GUTs with Flavor Symmetries $T(7)$ and $Sigma(81)$* , *Phys. Rev. D* **79** (2009) 036002 [[arXiv:0811.2955](#)] [[INSPIRE](#)].
- [62] I. de Medeiros Varzielas, S. King and G. Ross, *Tri-bimaximal neutrino mixing from discrete subgroups of $SU(3)$ and $SO(3)$ family symmetry*, *Phys. Lett. B* **644** (2007) 153 [[hep-ph/0512313](#)] [[INSPIRE](#)].
- [63] C. Luhn, S. Nasri and P. Ramond, *The Flavor group $\Delta(3n^2)$* , *J. Math. Phys.* **48** (2007) 073501 [[hep-th/0701188](#)] [[INSPIRE](#)].
- [64] J. Escobar and C. Luhn, *The Flavor Group $\Delta(6n^2)$* , *J. Math. Phys.* **50** (2009) 013524 [[arXiv:0809.0639](#)] [[INSPIRE](#)].